

JEE MAIN

RAY OPTICS FORMULAE

Now that's how you REVISE

- Mohit Goenka, IIT Kharagpur

List of Content on Eduniti YouTube Channel:

1. PYQs Video Solution Topic Wise:
 - (a) JEE Main 2018/2020/2021 Feb & March
2. Rank Booster Problems for JEE Main
3. Part Test Series for JEE Main
4. JEE Advanced Problem Solving Series
5. Short Concept Videos
6. Tips and Tricks Videos
7. JEE Advanced PYQs

.....and many more to come




EDUNITI




Eduniti for Physics


RAY OPTICS



1. REFLECTION



2. VECTOR FORM OF REFLECTED RAY




$$\hat{r} = \hat{i} - 2(\hat{i} \cdot \hat{n}) \cdot \hat{n}$$

\hat{i} , \hat{r} and \hat{n} are unit vectors along incident ray, reflected ray and normal.




3. MINIMUM MIRROR LENGTH TO SEE ONE'S IMAGE



* Assuming eye level at A


4. ANGLE OF DEVIATION (δ)




$$\delta = \pi - 2i$$

Here it is
ACW rotation


$$\begin{aligned} S_{NET} &= S_1 + S_2 \\ &= 2\pi - 2\theta \text{ CW} \end{aligned}$$



5. MIRROR ROTATION



6. IMAGE VELOCITY



$$\vec{v}_o/M = -\vec{v}_I/M$$

$$\Rightarrow \vec{v}_o - \vec{v}_M = -(\vec{v}_I - \vec{v}_M)$$


$$\Rightarrow \boxed{\vec{v}_I = 2\vec{v}_M - \vec{v}_o}$$

* These are all velocities
Normal to mirror surface.



SPHERICAL MIRROR

F. MIRROR IMPORTANT TERMS




r : radius of aperture

$2r$: Aperture size

C : center of curvature

R : Radius of curvature



$$FP = R - \frac{R}{2} \sec \theta$$

NOTE:

If rays were PARAXIAL
(close to axis)

then θ is very small.

$\Rightarrow F$ is FOCUS

and

$$FP = \frac{R}{2}$$



8. MIRROR FORMULAE

$$\frac{1}{f} = \frac{1}{V} + \frac{1}{U} \quad \text{or} \quad V = \frac{Uf}{U-f}$$

- Put U, f with sign
- f is -VE for CONCAVE
and +VE for CONVEX
MIRROR

9. MAGNIFICATION


→ TransVERSE, $m = \frac{h_i}{h_o} = -\frac{V}{U} = -\frac{f}{U-f}$

- * Put terms with sign
- * Erect image $\Rightarrow m$ is +VE
- * INVERTED $\Rightarrow m$ is -VE

→ LONGITUDINAL

$$m = \frac{\text{Image Length along P.A.}}{\text{Object Length along PA}}$$

Ex:



10. Six standard cases of image formation in CONCAVE MIRROR.


→ Must remember them all

→ Magnified image

Object between
f and C
(Real image)
 m is -VE

Object between Pole
and focus
(Virtual image)
 m is +VE.

11. Diameter of image of SUN (Distant object)




$$d = \theta f'$$

↪ θ in radians



12. IMAGE VELOCITY



$$V_{I/M} = - \left(\frac{v}{u} \right)^2 V_{0/M}$$


If $V_M = 0$ then, $V_I = - \left(\frac{f}{u-f} \right)^2 V_0$

→ put u, f, V_0 with sign



REFRACTION

13. SNELL'S LAW



$$(a.) n_1 \sin i = n_2 \sin r$$


(b) As per this fig.

$$n_2 > n_1$$

$$(c) \frac{n_2}{n_1} = \frac{V_1}{V_2} = \frac{\lambda_1}{\lambda_2}$$

V_1 and V_2 are speed
of light in medium
1 and 2.

#




$$n_1 \sin \theta_1 = n_2 \sin \theta_2 = n_3 \sin \theta_3 \\ = \dots = n_6 \sin \theta_6$$

$$\therefore n \sin \theta = \text{constant}$$



14. IMAGE FORMATION DUE TO PLANE SURFACE

(a)



$$d' = d \frac{n_2}{n_1}$$

(1.) d' is Apparent depth


(2.) $d' < d$ If $n_1 > n_2$

$d' > d$ If $n_1 < n_2$

(3.) Paraxial rays

or observer is
looking from above.

(b)




$$d' = n \left(\frac{d_1}{n_1} + \frac{d_2}{n_2} + \frac{d_3}{n_3} \right)$$




15. SHIFTING DUE TO SLAB

(a)



(b)




$$\Delta s = t \left(1 - \frac{n_{\text{sur}}\sigma}{n_{\text{slab}}} \right)$$

$$\Delta S = t_1 \left(1 - \frac{n}{n_1} \right) + t_2 \left(1 - \frac{n}{n_2} \right) + t_3 \left(1 - \frac{n}{n_3} \right)$$


- (1.) valid for paraxial
- (2.) valid only if medium around slab is same



16. LATERAL DISPLACEMENT



$$\Delta s = \frac{t \sin(i - r)}{\cos r}$$

17. SPHERICAL REFRACTION
IMAGE FORMULAE

$$\frac{n_2}{v} - \frac{n_1}{u} = \frac{n_2 - n_1}{R}$$


- (1.) Put u, v, R with sign
 (2.) n_2 is medium where rays are going



18. THIN LENSES: LENS FORMULAE

$$\frac{1}{V} - \frac{1}{U} = \frac{1}{f} \quad \text{or} \quad V = \frac{Uf}{U+f}$$

- (1.) f is +VE for converging lens
and -VE for diverging lens
- (2.) Valid for paraxial rays



$$\frac{1}{f} = \left(\frac{n_l}{n_s} - 1 \right) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

19. MAGNIFICATION

$$m = \frac{h_i}{h_o} = \frac{V}{U} = \frac{f}{U+f}$$

→ Transverse mag.


- (1.) Valid only if lens is surrounded by only one medium.
- (2.) Put R_1, R_2 with sign

LENS MAKER'S FORMULAE

20. OPTICAL POWER

$$P = \frac{1}{f_l}$$


- UNIT: DIOPTER
- put f_l with sign
- put f_e in meters

21. Combination of thin lenses, f_{eq} 

$$\frac{1}{f_{eq}} = \frac{1}{f_1} + \frac{1}{f_2} + \frac{1}{f_3}$$

- (1.) Put f_1, f_2, f_3 with sign
- (2.) Lenses must be in contact
- (3.) f_1, f_2, f_3 are individual focal lengths w.r.t surrounding medium.



22. COMBINATION OF LENS and MIRRORS (f_{eq})

$$\frac{1}{f_{eq}} = \frac{1}{f_m} - \frac{2}{f_l}$$

(1.) Put f_m, f_l with sign.


(2.) If f_{eq} is +VE \Rightarrow equivalent convex mirror

If f_{eq} is -VE \Rightarrow concave mirror

If f_{eq} is ∞ \Rightarrow Plane Mirror



23. DISPLACEMENT METHOD TO MEASURE FOCAL LENGTH OF CONVEX LENS



$$f = \frac{D^2 - x^2}{4D}$$

D : Distance between screen and object

x : Distance between two positions of lens.


$$h_0 = \sqrt{h_1 h_2}$$

↳ h_1 : image height when lens is at posⁿ 1

↳ h_2 : when at posⁿ 2



24. CRITICAL ANGLE AND TIR



- (1.) Here $n_1 > n_2$
- (2.) i_c : critical angle for which $\sigma = 90^\circ$
- (3.) If $\theta > i_c \Rightarrow \text{TIR}$

$$\sin i_c = \frac{n_2}{n_1} \quad \text{or}$$

$$i_c = \sin^{-1}\left(\frac{n_2}{n_1}\right)$$

NOTE: Question type

- (1.) OPTICAL FIBRE
- (2.) slab or prism
- (3.) Light source in liquid, find area of Patch at top surface.



25. PRISM

(1.) A: ANGLE OF PRISM, $A = \gamma_1 + \gamma_2$

(2.) S : ANGLE OF Deviation
 $S = i + e - A$

(3.) For S to be minimum, S_{\min}
 $i = e \Rightarrow \gamma_1 = \gamma_2 = \gamma$


$$\therefore \frac{n_p}{n_s} = \frac{\sin\left(\frac{S_m + A}{2}\right)}{\sin\left(\frac{A}{2}\right)}$$

n_p : R.I of PRISM

n_s : RI of Surrounding

(4.) If A is very small (THIN PRISM), then

$$S = A \left(\frac{n_p}{n_s} - 1 \right)$$



26. DISPERSIVE POWER (ω)



$$\delta_R = A(n_R - 1), \quad \delta_Y = A(n_Y - 1), \quad \delta_V = A(n_V - 1)$$

n_R, n_Y, n_V are R.I. of medium for Red, yellow and violet color


(Cauchy's eqn, $n = A + \frac{B}{\lambda^2} + \frac{C}{\lambda^4} + \dots$)

$$(a.) \theta \text{ (Dispersion Angle)} = \delta_V - \delta_R = A(n_V - n_R)$$

$$(b.) \omega \text{ (Dispersive Power)} = \frac{\delta_V - \delta_R}{\delta_Y} = \frac{n_V - n_R}{n_Y - 1}$$



27. CONDITION for DEVIATION WITHOUT DISPERSION



$$\theta_{NET} = 0$$


$$\Rightarrow \theta_1 = \theta_2$$

or

$$A_1(n_{V_1} - n_{R_1}) = A_2(n_{V_2} - n_{R_2})$$

Achromatic Prism
Combination

28. CONDITION FOR DISPERSION WITHOUT DEVIATION



Here final emergent yellow
is parallel to incident
white light.

$$so, \delta_{NET} = 0 \Rightarrow \delta_1 = \delta_2$$

$$\Rightarrow A_1(n_{V_1} - 1) = A_2(n_{V_2} - 1)$$



29. SIMPLE MICROSCOPE ($m = \theta_c / \theta_0$), $\theta_0 = h_0 / D$

|

Image at D Image at ∞

$$m = 1 + \frac{D}{f}$$

$$m = \frac{D}{f}$$

30. COMPOUND MICROSCOPE

$$(f_e > f_o)$$

$$m = m_o \times m_e$$

Image at D (strained condⁿ) Image at ∞
 (Normal adjustment)

(a) $m = \frac{V_o}{U_o} \left(1 + \frac{D}{f_e} \right)$

* $m = \frac{L}{f_o} \left(1 + \frac{D}{f_e} \right)$

(b) Tube Length = $V_o + U_e$ $m = \frac{V_o}{U_o} \frac{D}{f_e}$

$V_o + f_e$

only magnitudes



30. TELESCOPE ($f_o > f_e$)

Image at D

$$(a.) m = \frac{f_o}{f_e} \left(1 + \frac{f_e}{D} \right)$$

$$(b.) \text{Tube length} = f_o + f_e$$

Image at ∞

$$m = f_o / f_e$$

$$\text{Tube Length} = f_o + f_e$$

NOTE: put without sign.

