

श्री
Balaji

Narendra Avasthi

PROBLEMS IN
**PHYSICAL
CHEMISTRY**

for
JEE
Main & Advanced

Free
20 YEARS
PREVIOUS PAPERS
(WITH SOLUTIONS)
WITH THIS BOOK
WORTH ₹300/-



Purchase
**Genuine
Books only**

A genuine copy of this book is covered by 3D holographic film printed with the logo of Shri Balaji Publications. As you tilt the book, the logo disappears at a certain angle.

15th
edition

श्री
Balaji

Problems in **PHYSICAL CHEMISTRY**

for **JEE**
Main & Advanced

by:

Narendra Avasthi

Director,
Vibrant Academy India (P) Ltd.
KOTA



SHRI BALAJI PUBLICATIONS

(EDUCATIONAL PUBLISHERS & DISTRIBUTORS)

AN ISO 9001-2008 CERTIFIED ORGANIZATION

Muzaffarnagar (U.P.) - 251001

For More Material Join: @JEEAdvanced_2024




Contents



1.	Stoichiometry	1-72
2.	Atomic Structure	73-129
3.	Gaseous State	130-193
4.	Thermodynamics	194-257
5.	Chemical Equilibrium	258-309
6.	Ionic Equilibrium	310-363
7.	Chemical Kinetics and Nuclear Chemistry	364-426
8.	Electrochemistry	427-478
9.	Dilute Solution	479-520
10.	Solid State	521-561
11.	Surface Chemistry	562-600



1



STOICHIOMETRY

Laws of Chemical Combination

Chemical reactions take place according to certain laws. These laws are called the Laws of Chemical Combination. "These are no longer useful in chemical calculations now but gives an idea of earlier methods of analysing and relating compounds by mass."

❖ **Law of Conservation of Mass [Lavoisier (1774)]**

During any physical or chemical change, the sum of masses of all substances present in reactions vessel remain conserved.

❖ **Law of Constant Composition or Definite Proportions [Proust (1799)]**

In a given chemical compound, the elements are always combined in the same proportions by mass.

❖ **Law of Multiple Proportions [Dalton (1803)]**

Whenever two elements form more than one compound, the different masses of one element that combine with the same mass of the other element are in the ratio of small whole numbers.

❖ **Law of Reciprocal Proportions [Richter (1792)]**

When two elements combine separately with a fixed mass of a third elements then the ratio of their masses in which they do so is either same or some whole number multiple of the ratio in which they combine with each other.

❖ **Gay-Lussac's Law of combining volumes**

According to Gay-Lussac's law of combining volume, when gases react together, they always do so in volumes which bear a simple ratio to one another and to the volumes of the products, if these are also gases, provided all measurements of volumes are done under similar conditions of temperature and pressure.

❖ **Avogadro's Law**

The volume of a gas (at fixed pressure and temperature) is proportional to the number of moles (or molecules of gas present). Mathematically we can write

$$V \propto n$$

Or Equal volumes of all gases under similar conditions of temperature and pressure contain equal number of molecules.

Dalton's Atomic Theory

- ❖ Matter consists of tiny particles called atoms.
- ❖ Atoms are indestructible. In chemical reactions, the atoms rearrange but they do not themselves break apart.
- ❖ In any sample of a pure element, all the atoms are identical in mass and other properties.
- ❖ The atoms of different elements differ in mass and other properties.
- ❖ When atoms of different elements combine to form compounds, new and more complex particles are formed. However, in a given compound the constituent atoms are always present in the same fixed numerical ratio.

Modern Atomic Theory

- ❖ **Atom is no longer considered to be indivisible :** It has been found that an atom has a complex structure and is composed of sub-atomic particles such as electrons, protons and neutrons.
- ❖ **Atoms of the same element may not be similar in all respects :**
Ex. Isotopes ($^{23}_{11}\text{Na}$, $^{24}_{11}\text{Na}$)
- ❖ **Atoms of different elements may be similar in one or more respects :**
Ex. Isobars. ($^{40}_{20}\text{Ca}$, $^{40}_{18}\text{Ar}$)
- ❖ **Atom is the smallest unit which takes part in chemical reactions :** Although atom is composed of sub-atomic particles, yet it is the smallest particle which takes part in chemical reactions.
- ❖ **The ratio in which atoms unite may be fixed and integral but may not be simple :** For example, in sugar molecule ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$), the ratio of C, H and O atoms is 12 : 22 : 11 which is not simple.
- ❖ **Atoms of one element can be changed into atoms of other element :**
Ex. Artificial Nuclear Reactions
- ❖ **The mass of atom can be changed into energy :** According to Einstein's equation $E = mc^2$ (E Energy, m mass, c the speed of light, i.e., $3 \times 10^{10} \text{ cm sec}^{-1}$), mass and energy are inter-convertible.

Mole Concept

- (a) **Definition of one mole :** One mole is a collection of that many entities as there are number of atoms exactly in 12 gm of C-12 isotope.
- (b) $1 \text{ u} = 1 \text{ amu} = \frac{1}{12}$ of mass of 1 atom of $\text{C}^{12} = \frac{1 \text{ g}}{N_A} = 1.66 \times 10^{-24} \text{ g}$
- (c) **For Elements :**
- ❖ 1 g atom = 1 mole of atoms = N_A atoms.
 - ❖ g atomic mass (GAM) = mass of N_A atoms in g.
 - ❖ Mole of atoms = $\frac{\text{Mass (g)}}{\text{GAM or molar mass}}$

(d) For molecules :

- ❖ 1 g molecule 1 mole of molecule N_A molecule.
- ❖ g molecular mass (GMM) mass of N_A molecule in g.
- ❖ Moles of molecules $\frac{\text{Mass (g)}}{\text{GMM or molar mass}}$

(e) For ionic compounds :

- ❖ 1 g formula unit 1 mole of formula unit N_A formula unit.
- ❖ g formula mass (GFM) mass of N_A formula unit in g.
- ❖ Mole of formula unit $\frac{\text{Mass (g)}}{\text{GFM or molar mass}}$

- (f)** 1 mole of a substance
- Contains 6.022×10^{23} particles
 - Mass as much as molecular mass/atomic mass/ionic mass in grams
 - If it is a gas, one mole occupies a volume of 22.4 L at 1 atm & 273 K

(g) Average or mean atomic mass : Average atomic mass of element

$$A_{\text{Avg.}} = \frac{A_1 x_1 + A_2 x_2 + \dots}{x_1 + x_2 + \dots}$$

Here A_1, A_2 are isotopic mass of element and x_1, x_2 are natural abundance of isotopes.

(h) Average or mean molar mass : The average molar mass of the different substances present in the container

$$M_{\text{Avg.}} = \frac{M_1 n_1 + M_2 n_2 + \dots}{n_1 + n_2 + \dots}$$

Here M_1, M_2 are molar mass of substances and n_1, n_2 are mole of substances present in the container.

Empirical Formula, Molecular Formula**(a) Empirical formula :** Formula depicting constituent atom in their simplest ratio.

Molecular formula : Formula depicting actual number of atoms in one molecule of the compound.

(b) Relation between molecular formula and empirical formula :

$$n = \frac{\text{Molecular mass}}{\text{Empirical Formula mass}}$$

(c) Densities :

- ❖ Density $\frac{\text{Mass}}{\text{Volume}}$, Unit: g/cc
- ❖ Relative density $\frac{\text{Density of any substance}}{\text{Density of reference substance}}$
- ❖ Specific gravity $\frac{\text{Density of any substance}}{\text{Density of water at } 4^\circ \text{C}}$
- ❖ Vapour density : Ratio of density of vapours to the density of hydrogen at similar pressure and temperature.


$$\text{Vapour density} = \frac{\text{Density of vapours at some temperature and pressure}}{\text{Density of H}_2 \text{ gas at same temperature and pressure}}$$

$$\text{Vapour density} = \frac{\text{Molecular mass}}{2}$$


Stoichiometry

Stoichiometry pronounced (“stoy – key – om – e – tree”) is the calculations of the quantities of reactants and products involved in a chemical reaction. Following methods can be used for solving problems.

(a) Mole Method (Balancing is required)



(b) Principle of Atom Conservation (P.O.A.C.) method (Balancing is not required)



Concept of Limiting Reagent

- (a) Limiting Reagent :** It is very important concept in chemical calculation. It refers to reactant which is present in minimum stoichiometry quantity for a chemical reaction. It is reactant consumed fully in a chemical reaction. So all calculations related to various products or in sequence of reactions are made on the basis of limiting reagent.
- (b) Calculation of Limiting Reagent :** Divide given moles of each reactant by their stoichiometric coefficient, the one with least ratio is limiting reagent.

Percentage Yield

$$\text{The percentage yield of product} = \frac{\text{Actual yield}}{\text{theoretical maximum yield}} \times 100$$

Concentration Terms

(a) For solutions (i.e., homogeneous mixtures) :

❖ If the mixture is not homogeneous, then none of them is applicable.

(i) % by mass $\frac{w}{W} \times \frac{\text{Mass of solute}}{\text{Mass of solution}} \times 100$

[X% by mass means 100 gm solution contains X gm solute ; (100 - X) gm solvent]

(ii) % $\frac{w}{V} \times \frac{\text{Mass of solute}}{\text{Volume of solution}} \times 100$

[X% $\frac{w}{V}$ means 100 mL solution contains X gm solute]

$$(iii) \quad \% \frac{v}{V} = \frac{\text{Volume of solute}}{\text{Volume of solution}} \times 100$$

❖ For gases % by volume is same as mole %

$$(iv) \quad \text{Mole \%} = \frac{\text{Moles of solute}}{\text{Total moles}} \times 100$$

$$(v) \quad \text{Mole fraction (X)} = \frac{\text{Moles of solute}}{\text{Total moles}}$$

$$(vi) \quad \text{Molarity (M)} = \frac{\text{Moles of solute}}{\text{Volume of solution (in litre)}}$$

$$(vii) \quad \text{Molality (m)} = \frac{\text{Moles of solute}}{\text{Mass of solvent (in kg)}}$$

$$(viii) \quad \text{Parts per million (ppm)} = \frac{\text{Mass of solute}}{\text{Mass of solvent}} \times 10^6 \sim \frac{\text{Mass of solute}}{\text{Mass of solution}} \times 10^6$$

$$(ix) \quad \text{Formality (F)} = \frac{\text{No. of formula unit}}{\text{Volume of solution (in litre)}}$$

(b) (i) On adding solvent in a solution (dilution) : Number of mole of solute remains constant


$$M_f V_f = M_i V_i$$

(ii) Mixing of two solutions of same solute

$$M_f V_f = M_1 V_1 + M_2 V_2 + \dots$$

(c) Volume strength of H_2O_2 :

H_2O_2 (aq) solution labelled as 'x V' 'volume H_2O_2 (for e.g., '20 V' H_2O_2), it means x volume of O_2 (in litre) at 1 atm and 273K that can be obtained from 1 litre of such a sample when it decomposes according to



Eudiometry

(For reactions involving gaseous reactants and products)

- ❖ Eudiometry or gas analysis involves the calculations based on gaseous reactions or the reactions in which at least two components are gaseous, in which the amounts of gases are represented by their volumes, measured at the same pressure and temperature.
- ❖ Gay-Lussac's law of volume combination holds good.
- ❖ Problem may be solved directly in terms of volume, in place of mole.

The volume of gases produced is often given by mentioning certain solvent which absorb certain gases.

Solvent	gas (es) absorbed
KOH (aq.)	$\text{CO}_2, \text{SO}_2, \text{Cl}_2$
Ammoniacal Cu_2Cl_2	CO
Turpentine oil	O_3
Alkaline pyrogallol	O_2
Water	NH_3, HCl
Anhydrous $\text{CuSO}_4/\text{CaCl}_2$	H_2O

Redox Processes

(a) Oxidation Number : It is the charge (real or imaginary) which an atom appears to have when it is in combined state. It may be a whole no. or fractional. An element may have different values of oxidation number depending . It depends on nature of compound in which it is present. There are some operational rules to determine oxidation number.

(b) Definition of Oxidation and Reduction :

- ❖ **Oxidation :** Addition of oxygen , removal of hydrogen , addition of electronegative element, removal of electropositive element , loss of electrons , increase in oxidation number (de-electronation).
- ❖ **Reduction :** Removal of oxygen, addition of hydrogen, removal of electronegative element, addition of electropositive element, gain of electrons, decrease in oxid. no. (electronation).
- ❖ **Redox Reaction :** A reaction in which oxidation & reduction occur simultaneously.

(c) Oxidising and Reducing Agents :

- ❖ **Oxidising Agents (Oxidants, Oxidisers) :** They oxidise others, themselves are reduced & gain electrons. e.g., $\text{O}_2, \text{O}_3, \text{HNO}_3, \text{MnO}_2, \text{H}_2\text{O}_2$, halogens, $\text{KMnO}_4, \text{K}_2\text{Cr}_2\text{O}_7, \text{KIO}_3, \text{Cl}(\text{SO}_4)_3, \text{FeCl}_3, \text{NaOCl}$, hydrogen ions. (Atoms present in their higher oxidation state.)

Compounds in which reacting atom is present in its highest oxidation state may act as oxidant only.

- ❖ **Reducing Agents (Reductants or Reducers) :** They reduce others, themselves get oxidised & lose electrons. Also called reductants or reducers. H_2 molecular form is weak reducing agent but Nascent hydrogen is strong reducing agent e.g., C, CO, $\text{H}_2\text{S}, \text{SO}_2, \text{SnCl}_2$, Sodium thio Sulphate ($\text{Na}_2\text{S}_2\text{O}_3$), Al, Na, $\text{CaH}_2, \text{NaBH}_4, \text{LiAlH}_4$ etc. Compounds in which reacting atom is present in its lowest oxidation state may act as reductant only.

Both Oxidising & Reducing Agents : Compounds having reactive atom in its intermediate (between highest and lowest oxidation states) oxidation state may act as both oxidant or reductant. e.g., $\text{SO}_2, \text{H}_2\text{O}_2, \text{O}_3, \text{NO}_2$, etc.

(d) Balancing of redox reactions :

- ❖ Ion - electron method
- ❖ Oxidation number method

[Concept involved is that, in any chemical reaction electrons (e^-) cannot be produced so no. of e^- s in L.H.S. & R.H.S. should be the same]

Equivalent Concept

- (a) **Law of Chemical equivalence** : It states that in any chemical reaction the equivalents of all the reactants and products must be same.



Equivalents of 'A' Equivalents of 'B' Equivalents of 'C'

- (b) **Terms used in equivalent concept** :

- ❖ Equivalent mass of A $\frac{\text{Molar mass of A}}{\text{Valency factor or } n \text{ factor}}$
- ❖ Equivalents of 'A' $\frac{\text{Mass of 'A' (in g)}}{\text{Equivalent mass of 'A'}}$
- ❖ Numbers of equivalents of 'A' no. of moles of 'A' n -factor
- ❖ **Normality (N)** : For solutions, concentration term **normality (N)** is used, which can be defined as "the number of equivalent of solute present in one litre (1000 mL) solution".

$$N = \frac{\text{Number of g - equivalents of solute}}{\text{Volume of solution (in L)}}$$

$$N = \frac{W(g)}{E} \times \frac{1000}{V \text{ (in mL)}}$$

$$N = \text{Molarity} \times \text{Valence factor}$$

$$\text{milli-equivalents of solute} = \frac{N}{1000} \times V \text{ (in mL)} = \frac{M}{1000} \times V \text{ (in mL)} \times n \text{ factor}$$

- (c) **Valence factor (n-factor) calculation** : n -factor here we mean a conversion factor by which we divide molar mass of substance to get equivalent mass and it depends on nature of substance which vary from one condition to another condition. We can divide n -factor calculations in two category.

- ❖ **In case of non-redox reaction.**

n -factor = moles of charge displaced per mole of species

- ❖ **In case of redox reaction.**

n -factor of oxidizing or reducing agent = mole of electrons gain or lost per mol of species.

- (d) **Volumetric analysis (Titration)** : Titration is a procedure for determining the concentration of a solution by allowing a carefully measured volume to react with a standard solution of another substance, whose concentration is known.

- ❖ **Primary standard** : A substance available in a pure form or state of known purity which is used in standardizing a solution.
- ❖ **Standardization** : The process by which the concentration of a solution is accurately ascertained.
- ❖ **Standard solution** : A solution whose concentration has been accurately determined.
- ❖ **Titrant** : The reagent (a standard solution) which is added from a burette standard term to react with the analyte.
- ❖ **Titrate** : This mainly involves titrations based chemistry. It can be divided into two major category.

(I) Non-redox reactions

(II) Redox reactions

(e) Type of reactions :**(i) Non-redox reactions :** This involve following kind of titrations :

1. Acid-Base titrations
2. Double indicator acid-base titration
3. Precipitation titration
4. Back titration

(ii) Redox reactions : This involve following kind of titrations :


1. Iodimetry titrations
2. Iodometry titrations
3. Back titration

(f) Titrations :**(I) Non-redox titrations**

(1) Acid-Base titration : To find out strength or concentration of unknown acid or base, it is titrated against base or acid of known strength. At the equivalence point, equivalents of titrate and titrant solution are equal. So by using law of chemical equivalence strength of unknown solution is determined.

At equivalence point, milliequivalents of acid = milliequivalents of base.

(2) Double indicator acid-base titration : In the acid-base titration the equivalence point is known with the help of indicator which changes its colour at the end point. In the titration of polyacidic base or polybasic acid there are more than one end point. Some times one indicator is not able to give colour change at every end point. So to find out each end point we have to use more than one indicator. For example, in the titration of Na_2CO_3 against HCl there are two end points.



When we use phenolphthalein in the above titration it changes its colour at first end point when NaHCO_3 is formed and with it we can not determine second end point. Similarly with methyl orange it changes its colour at second end point only and we can not determine first end point. It is because all indicator changes colour on the basis of pH of titration mixture. So in titration of NaHCO_3 , KHCO_3 against acid, phenolphthalein can not be used.

Titration (against acid)	Indicator	pH Range	n-factor
------------------------------------	------------------	-----------------	-----------------

Na_2CO_3	Phenolphthalein	8.3 – 10	1
Na_2CO_3	Methyl orange	3.1 – 4.4	2

Note : When we carry out dilution of solution, milliequivalent, equivalent, milli mole or mole of substance does not change because they represent amount of substance, however molar concentration may change.

(3) Precipitation titration : In precipitation titrations involving ionic reactions, we determine the strength of unknown solution of salt by titrating it against a reagent with which it can form precipitate. For example NaCl strength can be determine by titrating it against AgNO_3 solution with which it form white ppt. of AgCl .

At equivalence point,

Milliequivalent of NaCl milliequivalent of AgNO_3 used milliequivalent of AgCl formed

(II) Redox Titrations

At equivalence point,

milliequivalent of Oxidizing agent used milliequivalent of reducing agent reacted.

S.No.	Reagent	Half Reaction	n-factor of reagent
1.	FAS (Mohr's salt) [$\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$]	$\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + e^-$	1
2.	MnO_4^- (Permanganate ion) (in acidic medium)	$\text{MnO}_4^- + 8\text{H}^+ + 5e^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$	5
3.	MnO_4^- (in basic medium)	$\text{Mn}^{7+} + e^- \rightarrow \text{Mn}^{6+}$	1
4.	MnO_4^- (in mild basic or neutral medium)	$\text{Mn}^{7+} + e^- \rightarrow \text{Mn}^{4+}$	3
5.	$\text{Cr}_2\text{O}_7^{2-}$ (dichromate ion)	$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6e^- \rightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	6
6.	$\text{C}_2\text{O}_4^{2-}$ (Oxalate ion)	$\text{C}_2\text{O}_4^{2-} \rightarrow 2\text{CO}_2 + 2e^-$	2
7.	As_2O_3	$\text{As}_2\text{O}_3 + 5\text{H}_2\text{O} + 2\text{AsO}_4^{3-} + 10\text{H}^+ + 4e^-$	4
8.	CaOCl_2 (Bleaching powder)	$\text{CaOCl}_2 + \text{H}_2\text{O} + 2\text{KI} \rightarrow \text{Ca(OH)}_2 + \text{I}_2 + 2\text{KCl}$	2
9.	MnO_2	$\text{MnO}_2 + 4\text{HCl (Conc.)} \rightarrow \text{MnCl}_2 + \text{Cl}_2 + 2\text{H}_2\text{O}$	2
10.	IO_3^-	$\text{IO}_3^- + 5\text{I}^- + 6\text{H}^+ \rightarrow 3\text{I}_2 + 3\text{H}_2\text{O}$	5
11.	H_2O_2 (act as oxidizing agent)	$\text{H}_2\text{O}_2 + 2e^- \rightarrow 2\text{H}_2\text{O}$	2
12.	H_2O_2 (act as reducing agent)	$\text{H}_2\text{O}_2 \rightarrow \text{O}_2 + 2e^-$	2
13.	H_2O_2 (disproportion)	$\text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{O} + \frac{1}{2}\text{O}_2$	1
14.	Cl_2 (disproportion)	$3\text{Cl}_2 + 6\text{OH}^- \text{ (strong)} \rightarrow \text{ClO}_3^- + 5\text{Cl}^- + 3\text{H}_2\text{O}$	5/3
15.	H_2S (in acidic medium)	$\text{H}_2\text{S} + \text{I}_2 \rightarrow \text{S} + 2\text{HI}$	2
16.	Sn^{2+} (in acidic medium)	$\text{Sn}^{2+} + \text{I}_2 \rightarrow \text{Sn}^{4+} + 2\text{I}^-$	2
17.	N_2H_4	$\text{N}_2\text{H}_4 \rightarrow \text{N}_2 + 4\text{H}^+ + 4e^-$	4
18.	SO_3^{2-} (in acidic medium)	$\text{SO}_3^{2-} + \text{H}_2\text{O} \rightarrow \text{SO}_4^{2-} + 2e^- + 2\text{H}^+$	2
19.	$\text{Na}_2\text{S}_2\text{O}_3$ (Sodium thiosulphate or Hypo)	$2\text{Na}_2\text{S}_2\text{O}_3 + \text{I}_2 \rightarrow \text{Na}_2\text{S}_4\text{O}_6 + 2\text{NaI}$	1
20.	I_2	$\text{I}_2 + 2e^- \rightarrow 2\text{I}^-$	2

(1) Iodimetric Titration or Iodimetry : In such titrations iodine solution is used as an oxidant and iodine is directly titrated against a reducing agent. This type of titrations are used for the determination of strength of reducing agents like sulphides, arsenides, thiosulphates etc., by titrating them against a standard solution of iodine.

This type of titration involves free iodine, here iodine solution is treated with known sodium thiosulphate solution.



Equivalents of I_2 Equivalents of $Na_2S_2O_3$ used

(2) Iodometric Titration or Iodometry: It is an indirect method of estimation of iodine. In this titration an oxidizing agent is used to liberate iodine from solution containing iodide ion (e.g., KI) and the liberated iodine is treated with a standard solution of a reducing agent added from a burette. Here a neutral or an acidic solution of an oxidizing agent is used and the amount of liberated I_2 is equal to the equivalents of this oxidizing agent.

These titrations are used to determine the concentration of $K_2Cr_2O_7$, $KMnO_4$, $CuSO_4$, Ferric ions, H_2O_2 etc.

These titrations are carried out in following two steps :

❖ **Step-1 :** Oxidizing agent (X) KI (excess) I_2 reduced state of oxidant

Equivalents of (X) Equivalents of I_2

❖ **Step-2 :** Liberated I_2 $2Na_2S_2O_3$ $2NaI$ $Na_2S_4O_6$

Equivalents of I_2 Equivalents of $Na_2S_2O_3$ used

(g) Back titration : Back titration is used in volumetric analysis to find out excess amount of reagent added by titrating it with suitable reagent. It is also used to find out percentage purity of sample.

For example in an acid-base titration suppose we have added excess base $[B(OH)_q]$ in acid (H_nA) solution.

To find excess base it is back titrated with another acid (H_mB) of known concentration.

Equivalents of base equivalents of (H_nA H_mB)

q moles of base taken n moles of H_nA reacted m moles of H_mB reacted

Hardness of Water

(a) Definition of hard water : Hard water is having soluble salts of calcium and magnesium ions.

(b) Degree of hardness : Degree of hardness defined as number of parts by mass of $CaCO_3$ (or its equivalent quantities of other substance) present in million parts of mass of water.

$$\text{Hardness of water} = \frac{\text{Mass of } CaCO_3}{\text{Mass of water}} \times 10^6 \text{ ppm.}$$



Level 1

- Calculate number of neutrons present in 12×10^{25} atoms of oxygen (${}_8\text{O}^{17}$) :
(Given: $N_A = 6 \times 10^{23}$)
(a) 1800 (b) 1600 (c) $1800 N_A$ (d) $3200 N_A$
- If mass of one atom is 3.32×10^{-23} g, then calculate number of nucleons (neutrons and protons) present in 2 atoms of the element :
(a) 40 (b) 20 (c) 10 (d) $40 N_A$
- Calculate number of electrons present in 9.5 g of PO_4^{3-} :
(a) 6 (b) $5 N_A$ (c) $0.1 N_A$ (d) $4.7 N_A$
- What is the number of moles of O-atom in 126 amu of HNO_3 ?
(a) 2 (b) $\frac{2}{N_A}$ (c) 6 (d) $\frac{6}{N_A}$
- What is the charge of 96 amu of S^{2-} ?
(a) 2 C (b) 3.2×10^{-19} C (c) 9.6×10^{-19} C (d) 6 C
- A sample of sodium has a mass of 46 g. What is the mass of the same number of calcium atoms as sodium atoms present in given sample ?
(a) 46 g (b) 20 g (c) 40 g (d) 80 g
- The total number of neutrons present in 54 mL H_2O (l) are:
(a) $3 N_A$ (b) $30 N_A$ (c) $24 N_A$ (d) none of these
- Total number of electrons present in 48 g Mg^{2+} are:
(a) $24 N_A$ (b) $2 N_A$ (c) $20 N_A$ (d) none of these
- The number of neutrons in 5 g of D_2O (D is ${}_1^2\text{H}$) are:
(a) $0.25 N_A$ (b) $2.5 N_A$ (c) $1.1 N_A$ (d) none of these
- Cisplatin, an anticancer drug, has the molecular formula $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$. What is the mass (in gram) of one molecule ? (Atomic masses : Pt = 195, H = 1.0, N = 14, Cl = 35.5)
(a) 4.98×10^{-21} (b) 4.98×10^{-22} (c) 6.55×10^{-21} (d) 3.85×10^{-22}
- Aspirin has the formula $\text{C}_9\text{H}_8\text{O}_4$. How many atoms of oxygen are there in a tablet weighing 360 mg?
(a) 1.204×10^{23} (b) 1.08×10^{22} (c) 1.204×10^{24} (d) 4.81×10^{21}
- 20 g of an ideal gas contains only atoms of S and O occupies 5.6 L at 1 atm and 273 K. What is the molecular mass of gas ?
(a) 64 (b) 80 (c) 96 (d) None of these
- A sample of ammonium phosphate, $(\text{NH}_4)_3\text{PO}_4$, contains 6 moles of hydrogen atoms. The number of moles of oxygen atoms in the sample is:
(a) 1 (b) 2 (c) 4 (d) 6
- Total number of moles of oxygen atoms in 3 litre O_3 (g) at 27°C and 8.21 atm are:
(a) 3 (b) 1 (c) 1 (d) none of these
- 3.011×10^{22} atoms of an element weighs 1.15 gm. The atomic mass of the element is:
(a) 10 amu (b) 2.3 amu (c) 35.5 amu (d) 23 amu

16. One atom of an element x weighs 6.643×10^{-23} g. Number of moles of atoms in its 20 kg is:
(a) 4 (b) 40 (c) 100 (d) 500
17. Mass of one atom of the element A is 3.9854×10^{-23} g. How many atoms are contained in 1 g of the element A ?
(a) 2.509×10^{23} (b) 6.022×10^{23} (c) 12.044×10^{23} (d) None of these
18. Which of the following contains the largest mass of hydrogen atoms?
(a) 5.0 moles $C_2H_2O_4$ (b) 1.1 moles $C_3H_8O_3$
(c) 1.5 moles $C_6H_8O_6$ (d) 4.0 moles $C_2H_4O_2$
19. Which has minimum number of oxygen atoms ?
(a) 10 mL H_2O (l) (b) 0.1 mole of V_2O_5 (s)
(c) 12 gm O_3 (g) (d) 12.044×10^{22} molecules of CO_2
20. Rearrange the following (I to IV) in the order of increasing masses:
(I) 0.5 mole of O_3 (II) 0.5 gm atom of oxygen
(III) 3.011×10^{23} molecules of O_2 (IV) 5.6 litre of CO_2 at 1 atm and 273 K
(a) $II < IV < III < I$ (b) $II < I < IV < III$ (c) $IV < II < III < I$ (d) $I < II < III < IV$
21. If the volume of a drop of water is 0.0018 mL then the number of water molecules present in two drop of water at room temperature is:
(a) 12.046×10^{19} (b) 1.084×10^{18} (c) 4.84×10^{17} (d) 6.023×10^{23}
22. It is known that atom contain protons, neutrons and electrons. If the mass of neutron is assumed to half of its original value whereas that of proton is assumed to be twice of its original value then the atomic mass of $^{14}_6C$ will be :
(a) same (b) 14.28% less (c) 14.28% more (d) 28.56% less
23. Common salt obtained from sea-water contains 8.775% NaCl by mass. The number of formula units of NaCl present in 25 g of this salt is :
(a) 3.367×10^{23} formula units (b) 2.258×10^{22} formula units
(c) 3.176×10^{23} formula units (d) 4.73×10^{25} formula units
24. The number of hydrogen atoms present in 25.6 g of sucrose ($C_{12}H_{22}O_{11}$) which has a molar mass of 342.3 g is :
(a) 22×10^{23} (b) 9.91×10^{23} (c) 11×10^{23} (d) 44×10^{23}
25. Caffeine has a molecular mass of 194. If it contains 28.9% by mass of nitrogen, number of atoms of nitrogen in one molecule of caffeine is :
(a) 4 (b) 6 (c) 2 (d) 3
26. The density of water is 1g/mL. What is the volume occupied by 1 molecule of water ?
(a) 1.44×10^{23} mL (b) 1mL (c) 18 mL (d) 2.88×10^{23} mL
27. A 25.0 mm $\times$ 40.0 mm piece of gold foil is 0.25 mm thick. The density of gold is 19.32 g/cm^3 . How many gold atoms are in the sheet ? (Atomic weight : Au = 197.0)
(a) 7.7×10^{23} (b) 1.5×10^{23} (c) 4.3×10^{21} (d) 1.47×10^{22}
28. If average molecular mass of air is 29, then assuming N_2 and O_2 gases are there, which options are correct regarding composition of air ?
(i) 75% by mass of N_2 (ii) 75% by moles N_2 (iii) 72.41% by mass of N_2
(a) only (i) is correct (b) only (ii) is correct
(c) both (ii) and (iii) are correct (d) both (i) and (ii) are correct

29. Density of dry air containing only N_2 and O_2 is 1.15 g/L at 740 mm of Hg and 300 K. What is % composition of N_2 by mass in the air ?
(a) 78% (b) 85.5% (c) 70.02% (d) 62.75%
30. A gaseous mixture of H_2 and CO_2 gases contains 66 mass % of CO_2 . The vapour density of the mixture is:
(a) 6.1 (b) 5.4 (c) 2.7 (d) 10.8
31. The vapour density of a mixture containing NO_2 and N_2O_4 is 27.6. The mole fraction of N_2O_4 in the mixture is:
(a) 0.1 (b) 0.2 (c) 0.5 (d) 0.8
32. Density of an ideal gas at 2 atm and 600 K is 2 g/L.
Calculate relative density of this gas with respect to Ne(g) under similar conditions :
(Given : $R = 1/12$ atm L/mol.K)
(a) 2.5 (b) 2 (c) 3 (d) 5
33. Average atomic mass of magnesium is 24.31 amu. This magnesium is composed of 79 mole % of ^{24}Mg and remaining 21 mole % of ^{25}Mg and ^{26}Mg . Calculate mole % of ^{26}Mg .
(a) 10 (b) 11 (c) 15 (d) 16
34. Indium (atomic mass = 114.82) has two naturally occurring isotopes, the predominant one form has isotopic mass 114.9041 and abundance of 95.72%. Which of the following isotopic mass is the most likely for the other isotope ?
(a) 112.94 (b) 115.90 (c) 113.90 (d) 114.90
35. Calculate density of a gaseous mixture which consist of 3.01×10^{24} molecules of N_2 and 32 g of O_2 gas at 3 atm pressure and 860 K temperature (Given : $R = 1/12$ atm L/mole.K)
(a) 0.6 g/L (b) 1.2 g/L (c) 0.3 g/L (d) 12 g/L
36. A mixture of O_2 and gas "Y" (mol. mass 80) in the mole ratio $a : b$ has a mean molecular mass 40. What would be mean molecular mass, if the gases are mixed in the ratio $b : a$ under identical conditions ? (Assume that gases are non-reacting):
(a) 40 (b) 48 (c) 62 (d) 72
37. If water sample are taken from sea, rivers or lake, they will be found to contain hydrogen and oxygen in the approximate ratio of 1 : 8. This indicates the law of:
(a) multiple proportion (b) definite proportion
(c) reciprocal proportions (d) none of these
38. Hydrogen and oxygen combine to form H_2O_2 and H_2O containing 5.93% and 11.2% hydrogen respectively. The data illustrates :
(a) law of conservation of mass (b) law of constant proportion
(c) law of reciprocal proportion (d) law of multiple proportion
39. Which one of the following combinations illustrate law of reciprocal proportions ?
(a) N_2O_3, N_2O_4, N_2O_5 (b) $NaCl, NaBr, NaI$
(c) CS_2, CO_2, SO_2 (d) PH_3, P_2O_3, P_2O_5
40. Carbon and oxygen combine to form two oxides, carbon monoxide and carbon dioxide in which the ratio of the masses of carbon and oxygen is respectively 12 : 16 and 12 : 32. These figures illustrate the :
(a) law of multiple proportions (b) law of reciprocal proportions
(c) law of conservation of mass (d) law of constant proportions

41. A sample of calcium carbonate (CaCO_3) has the following percentage composition : Ca = 40%, C = 12%, O = 48%. If the law of constant proportions is true, then the mass of calcium in 4 g of a sample of calcium carbonate obtained from another source will be :
(a) 0.016 g (b) 0.16 g (c) 1.6 g (d) 16 g
42. The law of multiple proportion is illustrated by the two compounds :
(a) Sodium chloride and sodium bromide (b) Ordinary water and heavy water
(c) Caustic soda and caustic potash (d) Sulphur dioxide and sulphur trioxide
43. All the substances listed below are fertilizers that contribute nitrogen to the soil. Which of these is the richest source of nitrogen on a mass percentage basis ?
(a) Urea, $(\text{NH}_2)_2\text{CO}$ (b) Ammonium nitrate, NH_4NO_3
(c) Nitric oxide, NO (d) Ammonia, NH_3
44. One mole of element X has 0.444 times the mass of one mole of element Y. One atom of element X has 2.96 times the mass of one atom of ^{12}C . What is the atomic mass of Y ?
(a) 80 (b) 15.77 (c) 46.67 (d) 40.0
45. A given sample of pure compound contains 9.81 gm of Zn, 1.8×10^{23} atoms of chromium and 0.60 mole of oxygen atoms. What is the simplest formula ?
(a) ZnCr_2O_7 (b) ZnCr_2O_4 (c) ZnCrO_4 (d) ZnCrO_6
46. The formula of an acid is HXO_2 . The mass of 0.0242 moles of the acid is 1.657 g. What is the atomic mass of X ?
(a) 35.5 (b) 28.1 (c) 128 (d) 19.0
47. What is the empirical formula of vanadium oxide, if 2.74 g of the metal oxide contains 1.53 g of metal ?
(a) V_2O_3 (b) VO (c) V_2O_5 (d) V_2O_7
48. Determine the empirical formula of Kelvar, used in making bullet proof vests, is 70.6% C, 4.2% H, 11.8% N and 13.4% O :
(a) $\text{C}_7\text{H}_5\text{NO}_2$ (b) $\text{C}_7\text{H}_5\text{N}_2\text{O}$ (c) $\text{C}_7\text{H}_9\text{NO}$ (d) $\text{C}_7\text{H}_5\text{NO}$
49. The hydrated salt $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$ undergoes 63% loss in mass on heating and becomes anhydrous. The value of x is:
(a) 10 (b) 12 (c) 8 (d) 18
50. A 6.85 g sample of the hydrated $\text{Sr}(\text{OH})_2 \cdot x\text{H}_2\text{O}$ is dried in an oven to give 3.13 g of anhydrous $\text{Sr}(\text{OH})_2$. What is the value of x ? (Atomic masses : Sr = 87.60, O = 16.0, H = 1.0)
(a) 8 (b) 12 (c) 10 (d) 6
51. What percentage of oxygen is present in the compound $\text{CaCO}_3 \cdot 3\text{Ca}_3(\text{PO}_4)_2$?
(a) 23.3% (b) 45.36% (c) 41.94% (d) 17.08%
52. Dieldrin, an insecticide, contains C, H, Cl and O. Combustion of 29.72 mg of Dieldrin gave 41.21 mg CO_2 and 5.63 mg of H_2O . In a separate analysis 25.31 mg of Dieldrin was converted into 57.13 mg AgCl. What is the empirical formula of Dieldrin ?
(a) $\text{C}_6\text{H}_4\text{Cl}_3\text{O}$ (b) $\text{C}_8\text{H}_8\text{ClO}$ (c) $\text{C}_{12}\text{H}_8\text{Cl}_6\text{O}$ (d) $\text{C}_6\text{H}_4\text{Cl}_3\text{O}_2$
53. A gaseous compound is composed of 85.7% by mass carbon and 14.3% by mass hydrogen. Its density is 2.28 g/litre at 300 K and 1.0 atm pressure. Determine the molecular formula of the compound.
(a) C_2H_2 (b) C_2H_4 (c) C_4H_8 (d) C_4H_{10}

54. Complete combustion of 0.86 g of compound X gives 2.64 g of CO_2 and 1.26 g of H_2O . The lowest molecular mass X can have :
(a) 43 g (b) 86 g (c) 129 g (d) 172 g
55. The sulphate of a metal M contains 9.87% of M . This sulphate is isomorphous with $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$. The atomic mass of M is :
(a) 40.3 (b) 36.3 (c) 24.3 (d) 11.3
56. In an organic compound of molar mass 108 gmol^{-1} C, H and N atoms are present in 9 : 1 : 3.5 by mass. Molecular formula can be :
(a) $\text{C}_6\text{H}_8\text{N}_2$ (b) $\text{C}_7\text{H}_{10}\text{N}$ (c) $\text{C}_5\text{H}_6\text{N}_3$ (d) $\text{C}_4\text{H}_{18}\text{N}_3$
57. On analysis, a certain compound was found to contain 254 g of iodine (at. mass 127) and 80 g oxygen (at. mass 16). What is the formula of the compound ?
(a) IO (b) I_2O (c) I_5O_3 (d) I_2O_5
58. An element A is tetravalent and another element B is divalent. The formula of the compound formed from these elements will be :
(a) A_2B (b) AB (c) AB_2 (d) A_2B_3
59. A compound used in making nylon, contains 43.8% oxygen. There are four oxygen atoms per molecule. What is the molecular mass of compound?
(a) 36 (b) 116 (c) 292 (d) 146
60. Suppose two elements X and Y combine to form two compounds XY_2 and X_2Y_3 . 0.05 mole of XY_2 weighs 5 g while 3.011×10^{23} molecules of X_2Y_3 weigh 85 g. The atomic masses of X and Y are respectively :
(a) 20, 30 (b) 30, 40 (c) 40, 30 (d) 80, 60
61. 44 g of a sample on complete combustion gives 88 g CO_2 and 36 g of H_2O . The molecular formula of the compound may be:
(a) C_4H_6 (b) $\text{C}_2\text{H}_6\text{O}$ (c) $\text{C}_2\text{H}_4\text{O}$ (d) $\text{C}_3\text{H}_6\text{O}$
62. 40 milligram diatomic volatile substance (X_2) is converted to vapour that displaced 4.92 mL of air at 1 atm and 300 K. Atomic mass of element X is nearly:
(a) 400 (b) 240 (c) 200 (d) 100
63. Two elements X (at. mass 75) and Y (at. mass 16) combine to give a compound having 75.8% of X . The formula of the compound is:
(a) XY (b) X_2Y (c) X_2Y_2 (d) X_2Y_3
64. A sample of phosphorus that weighs 12.4 g exerts a pressure 8 atm in a 0.821 litre closed vessel at 527°C . The molecular formula of the phosphorus vapour is:
(a) P_2 (b) P_4 (c) P_6 (d) P_8
65. Manganese forms non-stoichiometric oxides having the general formula MnO_x . The value of x for the compound that analyzed 64% by mass Mn:
(a) 1.16 (b) 1.83 (c) 2 (d) 1.93
66. 1.44 gram of titanium (At. mass 48) reacted with excess of O_2 and produce x gram of non-stoichiometric compound $\text{Ti}_{1.44}\text{O}$. The value of x is:
(a) 2 (b) 1.77 (c) 1.44 (d) none of these
67. Which statement is false for the balanced equation given below ?
$$\text{CS}_2 + 3\text{O}_2 \rightarrow 2\text{SO}_2 + \text{CO}_2$$

- (a) One mole of CS_2 will produce one mole of CO_2
 (b) The reaction of 16 g of oxygen produces 7.33 g of CO_2
 (c) The reaction of one mole of O_2 will produce $2/3$ mole of SO_2
 (d) Six molecules of oxygen requires three molecules of CS_2
68. Which of the following setups is correct to calculate the mass (in g) of KClO_3 produced from the reaction of 0.150 moles of Cl_2 ?
- $$3\text{Cl}_2 + 6\text{KOH} \rightarrow 5\text{KCl} + \text{KClO}_3 + 3\text{H}_2\text{O}$$
- (a) 0.150 moles Cl_2 1 mole $\text{KClO}_3/3$ moles Cl_2 122.5 g/1 mole KClO_3
 (b) 0.150 moles Cl_2 1 mole $\text{KClO}_3/3$ moles Cl_2 1 mole $\text{KClO}_3/122.5$ g
 (c) 0.150 moles Cl_2 3 moles $\text{Cl}_2/1$ mole KClO_3 122.5 g/1 mole KClO_3
 (d) 0.150 moles Cl_2 3 moles $\text{Cl}_2/1$ mole KClO_3 1 mole $\text{KClO}_3/122.5$ g
69. 2.0 g of a sample contains mixture of SiO_2 and Fe_2O_3 . On very strong heating, it leaves a residue weighing 1.96 g. The reaction responsible for loss of mass is given below.
- $$\text{Fe}_2\text{O}_3(s) \rightarrow \text{Fe}_3\text{O}_4(s) + \text{O}_2(g), \text{ (unbalance equation)}$$
- What is the percentage by mass of SiO_2 in original sample ?
 (a) 10% (b) 20% (c) 40% (d) 60%
70. What volume of air at 1 atm and 273 K containing 21% of oxygen by volume is required to completely burn sulphur (S_8) present in 200 g of sample, which contains 20% inert material which does not burn. Sulphur burns according to the reaction
- $$\frac{1}{8} \text{S}_8(s) + \text{O}_2(g) \rightarrow \text{SO}_2(g)$$
- (a) 23.52 litre (b) 320 litre (c) 112 litre (d) 533.33 litre
71. For the reaction, $2\text{Fe}(\text{NO}_3)_3 + 3\text{Na}_2\text{CO}_3 \rightarrow \text{Fe}_2(\text{CO}_3)_3 + 6\text{NaNO}_3$
 initially 2.5 mole of $\text{Fe}(\text{NO}_3)_3$ and 3.6 mole of Na_2CO_3 are taken. If 6.3 mole of NaNO_3 is obtained then % yield of given reaction is:
 (a) 50 (b) 84 (c) 87.5 (d) 100
72. How many moles of P_4 can be produced by reaction of 0.10 moles $\text{Ca}_5(\text{PO}_4)_3\text{F}$, 0.36 moles SiO_2 and 0.90 moles C according to the following reaction ?
- $$4 \text{Ca}_5(\text{PO}_4)_3\text{F} + 18 \text{SiO}_2 + 30 \text{C} \rightarrow 3\text{P}_4 + 2\text{CaF}_2 + 18\text{CaSiO}_3 + 30 \text{CO}$$
- (a) 0.060 (b) 0.030 (c) 0.045 (d) 0.075
73. Some older emergency oxygen masks contains potassium superoxide, KO_2 which reacts with CO_2 and water present in exhaled air to produce oxygen according to the given equation. If a person exhales 0.667 g of CO_2 per minute, how many grams of KO_2 are consumed in 5.0 minutes?
- $$4\text{KO}_2 + 2\text{H}_2\text{O} + 4\text{CO}_2 \rightarrow 4\text{KHCO}_3 + 3\text{O}_2$$
- (a) 10.7 (b) 0.0757 (c) 1.07 (d) 5.38
74. The mass of N_2F_4 produced by the reaction of 2.0 g of NH_3 and 80 g of F_2 is 3.56 g. What is the per cent yield?
- $$2\text{NH}_3 + 5\text{F}_2 \rightarrow \text{N}_2\text{F}_4 + 6\text{HF}$$
- (a) 79.0 (b) 71.2 (c) 84.6 (d) None of these

- 75.** Calculate the mass of lime (CaO) obtained by heating 200 kg of 95% pure lime stone (CaCO_3):
 (a) 104.4 kg (b) 105.4 kg (c) 212.8 kg (d) 106.4 kg
- 76.** Phosphoric acid (H_3PO_4) prepared in a two step process.
 (1) $\text{P}_4 + 5\text{O}_2 \rightarrow \text{P}_4\text{O}_{10}$ (2) $\text{P}_4\text{O}_{10} + 6\text{H}_2\text{O} \rightarrow 4\text{H}_3\text{PO}_4$
 We allow 62 g of phosphorus to react with excess oxygen which form P_4O_{10} in 85% yield. In the step (2) reaction 90% yield of H_3PO_4 is obtained. Mass of H_3PO_4 produced is:
 (a) 37.485 g (b) 149.949 g (c) 125.47 g (d) 564.48 g
- 77.** 9 moles of "D" and 14 moles of E are allowed to react in a closed vessel according to given reactions. Calculate number of moles of B formed in the end of reaction, if 4 moles of G are present in reaction vessel. (Percentage yield of reaction is mentioned in the reaction)
 Step-1 $3\text{D} + 4\text{E} \xrightarrow{80\%} 5\text{C} + \text{A}$
 Step-2 $3\text{C} + 5\text{G} \xrightarrow{50\%} 6\text{B} + \text{F}$
 (a) 2.4 (b) 30 (c) 4.8 (d) 1
- 78.** The chief ore of Zn is the sulphide, ZnS. The ore is concentrated by froth floatation process and then heated in air to convert ZnS to ZnO.
 $2\text{ZnS} + 3\text{O}_2 \xrightarrow{80\%} 2\text{ZnO} + 2\text{SO}_2$
 $\text{ZnO} + \text{H}_2\text{SO}_4 \xrightarrow{100\%} \text{ZnSO}_4 + \text{H}_2\text{O}$
 $2\text{ZnSO}_4 + 2\text{H}_2\text{O} \xrightarrow{80\%} 2\text{Zn} + 2\text{H}_2\text{SO}_4 + \text{O}_2$
 The number of moles of ZnS required for producing 2 moles of Zn will be :
 (a) 3.125 (b) 2 (c) 2.125 (d) 4
- 79.** 0.8 mole of a mixture of CO and CO_2 requires exactly 40 gram of NaOH in solution for complete conversion of all the CO_2 into Na_2CO_3 . How many moles more of NaOH would it require for conversion into Na_2CO_3 , if the mixture (0.8 mole) is completely oxidised to CO_2 ?
 (a) 0.2 (b) 0.6 (c) 1 (d) 1.5
- 80.** Silver oxide (Ag_2O) decomposes at temperature 300°C yielding metallic silver and oxygen gas. A 1.60 g sample of impure silver oxide yields 0.104 g of oxygen gas. What is the per cent by mass of the silver oxide in the sample?
 (a) 5.9 (b) 47.125 (c) 94.25 (d) 88.2
- 81.** 342 g of 20% by mass of $\text{Ba}(\text{OH})_2$ solution (sp. gr. 0.57) is reacted with 1200 mL of 2 M HNO_3 . If the final density of solution is same as pure water then molarity of the ion in resulting solution which decides the nature of the above solution is:
 (a) 0.25 (b) 0.5 M (c) 0.888 M (d) None of these
- 82.** 100 mL of H_2SO_4 solution having molarity 1 M and density 1.5 g/mL is mixed with 400 mL of water. Calculate final molarity of H_2SO_4 solution, if final density is 1.25 g/mL :
 (a) 4.4 M (b) 0.145 M (c) 0.52 M (d) 0.227 M
- 83.** What volume of HCl solution of density 1.2 g/cm^3 and containing 36.5% by mass HCl, must be allowed to react with zinc (Zn) in order to liberate 4.0 g of hydrogen?
 (a) 333.33 mL (b) 500 mL (c) 614.66 mL (d) None of these
- 84.** An ideal gaseous mixture of ethane (C_2H_6) and ethene (C_2H_4) occupies 28 litre at 1 atm and 273 K. The mixture reacts completely with 128 g O_2 to produce CO_2 and H_2O . Mole fraction at C_2H_6 in the mixture is:
 (a) 0.6 (b) 0.4 (c) 0.5 (d) 0.8

85. Wood's metal contains 50.0% bismuth, 25.0% lead, 12.5% tin and 12.5% cadmium by mass. What is the mole fraction of tin ?
(Atomic mass : Bi 209, Pb 207, Sn 119, Cd 112)
(a) 0.202 (b) 0.158 (c) 0.176 (d) 0.221
86. The density of a 56.0% by mass aqueous solution of 1-propanol ($\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$) is 0.8975 g/cm^3 . What is the mole fraction of the 1-propanol ?
(a) 0.292 (b) 0.227 (c) 0.241 (d) 0.276
87. What is the molality of SO_4^{2-} ion in aqueous solution that contain 34.2 ppm of $\text{Al}_2(\text{SO}_4)_3$?
(Assume complete dissociation and density of solution 1 g/mL)
(a) $3 \times 10^{-4} \text{ M}$ (b) $2 \times 10^{-4} \text{ M}$ (c) 10^{-4} M (d) None of these
88. The relation between molarity (M) and molality (m) is given by :
(density of solution (g/mL), M_1 molecular mass of solute)
(a) $m = \frac{1000 M}{1000 - M_1}$ (b) $m = \frac{1000 M}{1000 + MM_1}$
(c) $m = \frac{1000 MM}{1000 + MM_1}$ (d) $m = \frac{1000 M}{1000 + MM_1}$
89. Molarity and molality of a solution of an liquid (mol. mass 50) in aqueous solution is 9 and 10 respectively. What is the density of solution?
(a) 1 g/cc (b) 0.95 g/cc (c) 1.05 g/cc (d) 1.35 g/cc
90. An aqueous solution of ethanol has density 1.025 g/mL and it is 2 M . What is the molality of this solution ?
(a) 1.79 (b) 2.143 (c) 1.951 (d) None of these
91. 0.2 mole of HCl and 0.2 mole of barium chloride were dissolved in water to produce a 500 mL solution. The molarity of the Cl^- ions is :
(a) 0.06 M (b) 0.09 M (c) 1.2 M (d) 0.80 M
92. Calculate the mass of anhydrous HCl in 10 mL of concentrated HCl (density = 1.2 g/mL) solution having 37% HCl by mass.
(a) 4.44 g (b) 4.44 mg (c) $4.44 \times 10^3 \text{ mg}$ (d) 0.444 g
93. Calculate the molality of 1 L solution of 80% H_2SO_4 (w/V), given that the density of the solution is 1.80 g mL^{-1} .
(a) 8.16 (b) 8.6 (c) 1.02 (d) 10.8
94. Fluoxymesterone, $\text{C}_{20}\text{H}_{29}\text{FO}_3$, is an anabolic steroid. A 500 mL solution is prepared by dissolving 10.0 mg of the steroid in water. 1.0 mL portion of this solution is diluted to a final volume of 1.00 L . What is the resulting molarity ?
(a) 1.19×10^{-10} (b) 1.19×10^{-7} (c) 5.95×10^{-8} (d) 2.38×10^{-11}
95. The 25 mL of a 0.15 M solution of lead nitrate, $\text{Pb}(\text{NO}_3)_2$ reacts with all of the aluminium sulphate, $\text{Al}_2(\text{SO}_4)_3$, present in 20 mL of a solution. What is the molar concentration of the $\text{Al}_2(\text{SO}_4)_3$?
$$3\text{Pb}(\text{NO}_3)_2(aq) + \text{Al}_2(\text{SO}_4)_3(aq) \rightarrow 3\text{PbSO}_4(s) + 2\text{Al}(\text{NO}_3)_3(aq)$$

(a) $6.25 \times 10^{-2} \text{ M}$ (b) $2.421 \times 10^{-2} \text{ M}$ (c) 0.1875 M (d) None of these
96. Concentrated HNO_3 is 63% HNO_3 by mass and has a density of 1.4 g/mL . How many millilitres of this solution are required to prepare 250 mL of a 1.20 M HNO_3 solution?
(a) 18.0 (b) 21.42 (c) 20.0 (d) 14.21

97. 50 mL of 20.8% (w/V) BaCl_2 (aq) and 100 mL of 9.8% (w/V) H_2SO_4 (aq) solutions are mixed. Molarity of Cl^- ions in the resulting solution is: (At. mass of Ba = 137)
(a) 0.333 M (b) 0.666 M (c) 0.1 M (d) 1.33 M
98. 100 mL of 10% NaOH (w/V) is added to 100 mL of 10% HCl (w/V). The nature of resultant solution is:
(a) alkaline (b) strongly alkaline (c) acidic (d) neutral
99. How many millilitres of 0.1 M H_2SO_4 must be added to 50 mL of 0.1 M NaOH to give a solution that has a concentration of 0.05 M in H_2SO_4 ?
(a) 400 mL (b) 200 mL (c) 100 mL (d) None of these
100. 1 M HCl and 2 M HCl are mixed in volume ratio of 4 : 1. What is the final molarity of HCl solution?
(a) 1.5 (b) 1 (c) 1.2 (d) 1.8
101. Three solutions X, Y, Z of HCl are mixed to produce 100 mL of 0.1 M solution. The molarities of X, Y and Z are 0.07 M, 0.12 M and 0.15 M respectively. What respective volumes of X, Y and Z should be mixed ?
(a) 50 mL, 25 mL, 25 mL (b) 20 mL, 60 mL, 20 mL
(c) 40 mL, 30 mL, 30 mL (d) 55 mL, 20 mL, 25 mL
102. A bottle of an aqueous H_2O_2 solution is labelled as '28.375 V H_2O_2 ' and the density of the solution (in g/mL) is 1.25. Choose the correct option:
(a) Molality of H_2O_2 solution is 2 (b) Molarity of H_2O_2 solution is 5
(c) Molality of H_2O_2 solution is 2.15 (d) None of these
103. The impure 6 g of NaCl is dissolved in water and then treated with excess of silver nitrate solution. The mass of precipitate of silver chloride is found to be 14 g. The % purity of NaCl solution would be:
(a) 95% (b) 85% (c) 75% (d) 65%
104. $\text{Al}_2(\text{SO}_4)_3$ solution of 1 molal concentration is present in 1 litre solution of density 2.684 g/cc. How many moles of BaSO_4 would be precipitated on adding excess of BaCl_2 in it ?
(a) 2 moles (b) 3 moles (c) 6 moles (d) 12 moles
105. A certain public water supply contains 0.10 ppb (part per billion) of chloroform (CHCl_3). How many molecules of CHCl_3 would be obtained in 0.478 mL drop of this water ?
(assumed $d = 1 \text{ g/mL}$)
(a) $4 \times 10^{13} N_A$ (b) $10^3 N_A$ (c) $4 \times 10^{10} N_A$ (d) None of these
106. Decreasing order (first having highest and then others following it) of mass of pure NaOH in each of the aqueous solution:
(i) 50 g of 40% (w/W) NaOH
(ii) 50 mL of 50% (w/V) NaOH [$d_{\text{soln.}} = 1.2 \text{ g/mL}$]
(iii) 50 g of 15 M NaOH [$d_{\text{soln.}} = 1 \text{ g/mL}$]
(a) i, ii, iii (b) iii, ii, i (c) ii, iii, i (d) ii, i, iii
107. What is the molar mass of diacidic organic Lewis base (B), if 12 g of its chloroplatinate salt ($\text{B H}_2\text{PtCl}_6$) on ignition produced 5 g residue of Pt?
(a) 52 (b) 58 (c) 88 (d) None of these

- 108.** On strong heating, one gram of the silver salt of an organic dibasic acid yields 0.5934 g of silver. If the mass percentage of carbon in it 8 times the mass percentage of hydrogen and one-half the mass percentage of oxygen, determine the molecular formula of the acid.
(a) $C_4H_6O_4$ (b) $C_4H_6O_6$ (c) $C_2H_6O_2$ (d) $C_5H_{10}O_5$
- 109.** 0.607 g of a silver salt of tribasic organic acid was quantitatively reduced to 0.37 g of pure Ag. What is the molecular mass of the acid?
(a) 207 (b) 210 (c) 531 (d) 324
- 110.** A sample of peanut oil weighing 2 g is added to 25 mL of 0.40 M KOH. After saponification is complete, 8.5 mL of 0.28 M H_2SO_4 is needed to neutralize excess of KOH. The saponification number of peanut oil is :
(saponification number is defined as the milligrams of KOH consumed by 1 g of oil)
(a) 146.72 (b) 223.44 (c) 98.9 (d) None of these
- 111.** 20 mL of a mixture of CO and H_2 were mixed with excess of O_2 and exploded and cooled. There was a volume contraction of 18 mL. All volume measurements corresponds to room temperature ($27^\circ C$) and one atmospheric pressure. Determine the volume ratio $V_1 : V_2$ of CO and H_2 in the original mixture.
(a) 1 : 2 (b) 3 : 2 (c) 2 : 3 (d) 4 : 1
- 112.** In the reaction,
 $2Al(s) + 6HCl(aq) \rightarrow 2Al^{3+}(aq) + 6Cl^{-}(aq) + 3H_2(g)$
(a) 6 L $HCl(aq)$ is consumed for every 3 L $H_2(g)$ produced
(b) 33.6 L $H_2(g)$ is produced regardless of temperature and pressure for every mole Al that reacts
(c) 67.2 L $H_2(g)$ at 1 atm and 273 K is produced for every mole Al that reacts
(d) 11.2 L $H_2(g)$ at 1 atm and 273 K is produced for every mole $HCl(aq)$ consumed
- 113.** The percentage by volume of C_3H_8 in a gaseous mixture of C_3H_8 , CH_4 and CO is 20. When 100 mL of the mixture is burnt in excess of O_2 , the volume of CO_2 produced is:
(a) 90 mL (b) 160 mL (c) 140 mL (d) none of these
- 114.** 40 mL gaseous mixture of CO, CH_4 and Ne was exploded with 15 mL of oxygen. After treatment with KOH the volume reduced by 9 mL and again on treatment with alkaline pyrogallol, the volume further reduced by 1.5 mL. Percentage of CH_4 in the original mixture is:
(a) 22.5 (b) 77.5 (c) 7.5 (d) 15
- 115.** A gaseous mixture of propane and butane of volume 3 litre on complete combustion produces 11.0 litre CO_2 under standard conditions of temperature and pressure. The ratio of volume of butane to propane is:
(a) 1 : 2 (b) 2 : 1 (c) 3 : 2 (d) 3 : 1
- 116.** Phosphorous has the oxidation state of +1 in:
(a) Orthophosphoric acid (b) Phosphorous acid
(c) Hypophosphorous acid (d) Metaphosphoric acid
- 117.** The oxidation state(s) of Cl in $CaOCl_2$ (bleaching powder) is/are:
(a) +1 only (b) -1 only (c) +1 and 1 (d) none of these
- 118.** The oxidation number of sulphur in S_8 , S_2F_2 , H_2S and H_2SO_4 respectively are:
(a) 0, 1, 2 and 6 (b) 2, 0, 2 and 6
(c) 0, 1, 2 and 4 (d) 2, 0, 2 and 6

- 119.** Fe shows an oxidation state of +1 in:
 (a) $\text{Fe}(\text{CO})_5$ (b) $[\text{Fe}(\text{H}_2\text{O})_5\text{NO}]\text{SO}_4$
 (c) $\text{Fe}_4[\text{Fe}(\text{CN})_6]_3$ (d) FeCl_4
- 120.** When SO_2 is passed into an acidified potassium dichromate solution, the oxidation numbers of sulphur and chromium in the final products respectively are:
 (a) +6, +6 (b) +6, +3 (c) 0, +3 (d) +2, +3
- 121.** The oxidation number of nitrogen atoms in NH_4NO_3 are:
 (a) +3, +3 (b) +3, 3 (c) 3, 5 (d) 5, 3
- 122.** The oxidation states of S-atoms in Caro's and Marshall's acids are:
 (a) 6, 6 (b) 6, 4 (c) 6, 6 (d) 4, 6
- 123.** In which of the following the oxidation number of oxygen has been arranged in increasing order :
 (a) OF_2 KO_2 BaO_2 O_3 (b) BaO_2 KO_2 O_3 OF_2
 (c) BaO_2 O_3 OF_2 KO_2 (d) KO_2 OF_2 O_3 BaO_2
- 124.** The oxidation numbers of oxygen in KO_3 , Na_2O_2 respectively are :
 (a) 3, 2 (b) 1, 0 (c) 0, 1 (d) -0.33, -1
- 125.** The oxidation number of phosphorus in $\text{Ba}(\text{H}_2\text{PO}_2)_2$ is :
 (a) -1 (b) +1 (c) +2 (d) +3
- 126.** If it is known that in $\text{Fe}_{0.96}\text{O}$, Fe is present in +2 and +3 oxidation state, what is the mole fraction of Fe^{2+} in the compound ?
 (a) 12/25 (b) 25/12 (c) 1/12 (d) 11/12
- 127.** Which of the following sequence of compounds is according to the decreasing order of the oxidation state of nitrogen ?
 (a) HNO_3 , NO , NH_4Cl , N_2 (b) HNO_3 , NO , N_2 , NH_4Cl
 (c) HNO_3 , NH_4Cl , NO , N_2 (d) NO , HNO_3 , NH_4Cl , N_2
- 128.** 2 mole of N_2H_4 loses 16 mole of electron is being converted to a new compound X. Assuming that all of the N appears in the new compound. What is the oxidation state of 'N' in X ?
 (a) 1 (b) 2 (c) 2 (d) 4
- 129.** When $\text{K}_2\text{Cr}_2\text{O}_7$ is converted to K_2CrO_4 , the change in the oxidation state of chromium is :
 (a) 0 (b) 6 (c) 4 (d) 3
- 130.** When a manganous salt is fused with a mixture of KNO_3 and solid NaOH , the oxidation number of Mn changes from +2 to :
 (a) +4 (b) +3 (c) +6 (d) +7
- 131.** In $\text{Fe}(\text{II})$ - MnO_4 titration, HNO_3 , is not used because:
 (a) it oxidises Mn^{2+} (b) it reduces MnO_4
 (c) it oxidises Fe^{2+} (d) it reduces Fe^{3+} formed
- 132.** Which species are oxidized and reduced in the reaction ?
 FeC_2O_4 KMnO_4 Fe^{3+} CO_2 Mn^{2+}
 (a) Oxidised : Fe, C ; Reduced : Mn (b) Oxidised : Fe ; Reduced : Mn
 (c) Reduced : Fe, Mn ; Oxidised : C (d) Reduced : C ; Oxidised : Mn, Fe

- 133.** In which of the following reactions, H_2O_2 is acting as a reducing agent ?
 (a) $\text{SO}_2 + \text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{SO}_4$ (b) $2\text{KI} + \text{H}_2\text{O}_2 + 2\text{KOH} \rightarrow \text{I}_2$
 (c) $\text{PbS} + 4\text{H}_2\text{O}_2 \rightarrow \text{PbSO}_4 + 4\text{H}_2\text{O}$ (d) $\text{Ag}_2\text{O} + \text{H}_2\text{O}_2 \rightarrow 2\text{Ag} + \text{H}_2\text{O} + \text{O}_2$
- 134.** Following reaction describes the rusting of iron
 $4\text{Fe} + 3\text{O}_2 \rightarrow 4\text{Fe}^{3+} + 6\text{O}^{2-}$
 Which one of the following statement is incorrect ?
 (a) This is an example of a redox reaction (b) Metallic iron is reduced to Fe^{3+}
 (c) Fe^{3+} is an oxidising agent (d) Metallic iron is a reducing agent
- 135.** Which reaction does not represent auto redox or disproportionation ?
 (a) $\text{Cl}_2 + \text{OH}^- \rightarrow \text{Cl}^- + \text{ClO}_3^- + \text{H}_2\text{O}$ (b) $2\text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{O} + \text{O}_2$
 (c) $2\text{Cu} \rightarrow \text{Cu}^{2+} + \text{Cu}$ (d) $(\text{NH}_4)_2\text{Cr}_2\text{O}_7 \rightarrow \text{N}_2 + \text{Cr}_2\text{O}_3 + 4\text{H}_2\text{O}$
- 136.** Which of the following is redox reaction ?
 (a) H_2SO_4 reacts with NaOH
 (b) In atmosphere, O_3 is formed from O_2 by lightning
 (c) Evaporation of H_2O
 (d) Oxides of nitrogen are formed from nitrogen & oxygen by lightning
- 137.** Which of the following is a redox-reaction ?
 (a) $2\text{Na}[\text{Ag}(\text{CN})_2] + \text{Zn} \rightarrow \text{Na}_2[\text{Zn}(\text{CN})_4] + 2\text{Ag}$
 (b) $\text{BaO}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + \text{H}_2\text{O}_2$
 (c) $\text{N}_2\text{O}_5 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3$
 (d) $\text{AgNO}_3 + \text{KI} \rightarrow \text{AgI} + \text{KNO}_3$
- 138.** For the redox reaction, $\text{MnO}_4^- + \text{C}_2\text{O}_4^{2-} + \text{H}^+ \rightarrow \text{Mn}^{2+} + \text{CO}_2 + \text{H}_2\text{O}$
 the correct coefficients of the reactants for the balanced reaction are respectively MnO_4^- , $\text{C}_2\text{O}_4^{2-}$, H^+ :
 (a) 2, 5, 16 (b) 16, 3, 12 (c) 15, 16, 12 (d) 2, 16, 5
- 139.** In a chemical reaction, $\text{K}_2\text{Cr}_2\text{O}_7 + x\text{H}_2\text{SO}_4 + y\text{SO}_2 \rightarrow \text{K}_2\text{SO}_4 + \text{Cr}_2(\text{SO}_4)_3 + z\text{H}_2\text{O}$; the value of x , y and z respectively are :
 (a) $x = 1, y = 3, z = 1$ (b) $x = 4, y = 1, z = 4$
 (c) $x = 3, y = 2, z = 1$ (d) $x = 2, y = 2, z = 1$
- 140.** Balance the following equation and choose the quantity which is the sum of the coefficients of the products :
 $\dots \text{CS}_2 + \dots \text{Cl}_2 \rightarrow \text{CCl}_4 + \dots \text{S}_2\text{Cl}_2$
 (a) 5 (b) 3 (c) 6 (d) 2
- 141.** Balance the following equation and choose the quantity which is the sum of the coefficients of reactants and products:
 $\dots \text{PtCl}_4 + \dots \text{XeF}_2 \rightarrow \text{PtF}_6 + \dots \text{ClF} + \dots \text{Xe}$
 (a) 16 (b) 13 (c) 18 (d) 12
- 142.** If 0.1 mole H_3PO_x is completely neutralised by 5.6 g KOH then select the true statement.
 (a) $x = 3$ and given acid is dibasic
 (b) $x = 4$ and given acid has no P-H linkage
 (c) $x = 2$ and given acid does not form acid salt
 (d) all of these

- 143.** When potassium permanganate is titrated against ferrous ammonium sulphate in acidic medium, the equivalent mass of potassium permanganate is:
- (a) $\frac{\text{molecular mass}}{3}$ (b) $\frac{\text{molecular mass}}{5}$
 (c) $\frac{\text{molecular mass}}{2}$ (d) $\frac{\text{molecular mass}}{10}$
- 144.** Equivalent mass of FeS_2 in the half reaction, $\text{FeS}_2 \rightarrow \text{Fe}_2\text{O}_3 + \text{SO}_2$ is :
- (a) $M/10$ (b) $M/11$ (c) $M/6$ (d) $M/1$
- 145.** The equivalent mass of HCl in the given reaction is:
- $$\text{K}_2\text{Cr}_2\text{O}_7 + 14\text{HCl} \rightarrow 2\text{KCl} + 2\text{CrCl}_3 + 3\text{Cl}_2 + \text{H}_2\text{O}$$
- (a) 16.25 (b) 36.5 (c) 73 (d) 85.1
- 146.** Equivalent mass of H_3PO_2 when it disproportionate into PH_3 and H_3PO_3 is:
- (a) M (b) $M/2$ (c) $M/4$ (d) $3M/4$
- 147.** In the following reaction, $\text{As}_2\text{S}_3 + \text{H}^+ + \text{NO}_3^- \rightarrow \text{NO} + \text{H}_2\text{O} + \text{AsO}_4^{3-} + \text{SO}_4^{2-}$
 the equivalent mass of As_2S_3 is related to its molecular mass by :
- (a) $M/2$ (b) $M/4$ (c) $M/24$ (d) $M/28$
- 148.** Sulphur forms the chlorides S_2Cl_2 and SCl_2 . The equivalent mass of sulphur in SCl_2 is :
- (a) 8 g/mol (b) 16 g/mol (c) 64.8 g/mol (d) 32 g/mol
- 149.** The equivalent mass of an element is 4. Its chloride has a vapour density 59.25. Then, the valency of the elements is :
- (a) 4 (b) 3 (c) 2 (d) 1
- 150.** 6×10^{-3} mole $\text{K}_2\text{Cr}_2\text{O}_7$ reacts completely with 9×10^{-3} mole X^{n+} to give XO_3 and Cr^{3+} . The value of n is :
- (a) 1 (b) 2 (c) 3 (d) None of these
- 151.** What mass of $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ (mol. mass = 126) should be dissolved in water to prepare 250 mL of centinormal solution which act as a reducing agent ?
- (a) 0.63 g (b) 0.1575 g (c) 0.126 g (d) 0.875 g
- 152.** The equivalent mass of the salt, $\text{KHC}_2\text{O}_4 \cdot \text{H}_2\text{C}_2\text{O}_4 \cdot 4\text{H}_2\text{O}$ when it act as reducing agent is :
- (a) $\frac{\text{Mol. mass}}{1}$ (b) $\frac{\text{Mol. mass}}{2}$ (c) $\frac{\text{Mol. mass}}{3}$ (d) $\frac{\text{Mol. mass}}{4}$
- 153.** The equivalent mass of divalent metal is W . The molecular mass of its chloride is :
- (a) $W + 35.5$ (b) $W + 71$ (c) $2W + 71$ (d) $2W + 35.5$
- 154.** When BrO_3^- ion reacts with Br^- in acid medium, Br_2 is liberated. The equivalent mass of Br_2 in this reaction is:
- (a) $\frac{5M}{8}$ (b) $\frac{5M}{3}$ (c) $\frac{3M}{5}$ (d) $\frac{4M}{6}$
- 155.** If m_A gram of a metal A displaces m_B gram of another metal B from its salt solution and if the equivalent mass are E_A and E_B respectively then equivalent mass of A can be expressed as:
- (a) $E_A = \frac{m_A}{m_B} E_B$ (b) $E_A = \frac{m_A}{E_B} m_B$ (c) $E_A = \frac{m_B}{m_A} E_B$ (d) $E_A = \sqrt{\frac{m_A}{m_B}} E_B$

156. Hydrazine reacts with KIO_3 in presence of HCl as



The equivalent masses of N_2H_4 and KIO_3 respectively are:

- (a) 8 and 53.5 (b) 16 and 53.5 (c) 8 and 35.6 (d) 8 and 87
157. What will be the normality of a solution obtained by mixing 0.45 N and 0.60 N NaOH in the ratio 2 : 1 by volume ?
 (a) 0.4 N (b) 0.5 N (c) 1.05 N (d) 0.15 N
158. A solution containing 2.7×10^{-3} mol of A^{2+} ions required 1.6×10^{-3} mole of MnO_4^- for the oxidation of A^{2+} to AO_3 the medium used is :
 (a) neutral (b) acidic (c) strong basic (d) none of these
159. H_2O_2 is used as bleaching reagent because on dissociation it gives oxygen ($\text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{O} + \frac{1}{2}\text{O}_2$).
 "Chachi 420" used H_2O_2 solution to bleach her hair and she required 2.24 L O_2 gas at 1 atm and 273 K. She has a H_2O_2 solution labelled as '5.6 V' then what volume of such solution must she required to bleach her hair ?
 (a) 200 mL (b) 300 mL (c) 400 mL (d) 500 mL
160. 1.25 g of a solid dibasic acid is completely neutralised by 25 mL of 0.25 molar $\text{Ba}(\text{OH})_2$ solution. Molecular mass of the acid is :
 (a) 100 (b) 150 (c) 120 (d) 200
161. 10 mL of N-HCl , 20 mL of $\text{N}/2 \text{H}_2\text{SO}_4$ and 30 mL of $\text{N}/3 \text{HNO}_3$ are mixed together and volume made to one litre. The normality of H^+ in the resulting solution is :
 (a) $3\text{N}/100$ (b) $\text{N}/10$ (c) $\text{N}/20$ (d) $\text{N}/40$
162. 0.45 g of an acid of mol. mass 90 was neutralised by 20 mL of 0.54 N caustic potash (KOH). The basicity of acid is :
 (a) 1 (b) 2 (c) 3 (d) 4
163. A 3.4 g sample of H_2O_2 solution containing $x\%$ H_2O_2 by mass requires x mL of a KMnO_4 solution for complete oxidation under acidic condition. The molarity of KMnO_4 solution is :
 (a) 1 (b) 0.5 (c) 0.4 (d) 0.2
164. What volume of $\text{O}_2(\text{g})$ measured at 1 atm and 273 K will be formed by action of 100 mL of 0.5 N KMnO_4 on hydrogen peroxide in an acid solution ? The skeleton equation for the reaction is

$$\text{KMnO}_4 + \text{H}_2\text{SO}_4 + \text{H}_2\text{O}_2 \rightarrow \text{K}_2\text{SO}_4 + \text{MnSO}_4 + \text{O}_2 + \text{H}_2\text{O}$$

 (a) 0.12 litre (b) 0.028 litre (c) 0.56 litre (d) 1.12 litre
165. A sample of 1.0 g of solid Fe_2O_3 of 80% purity is dissolved in a moderately concentrated HCl solution which is reduced by zinc dust. The resulting solution required 16.7 mL of a 0.1 M solution of the oxidant. Calculate the mole of electrons taken up by the oxidant.
 (a) 2 (b) 4 (c) 6 (d) 5
166. KMnO_4 reacts with oxalic acid according to the equation

$$2\text{MnO}_4^- + 5\text{C}_2\text{O}_4^{2-} + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 10\text{CO}_2 + 8\text{H}_2\text{O}$$

 Here, 20 mL of 0.1 M KMnO_4 is equivalent to :
 (a) 120 mL of 0.25 M $\text{H}_2\text{C}_2\text{O}_4$ (b) 150 mL of 0.10 M $\text{H}_2\text{C}_2\text{O}_4$
 (c) 25 mL of 0.20 M $\text{H}_2\text{C}_2\text{O}_4$ (d) 50 mL of 0.20 M $\text{H}_2\text{C}_2\text{O}_4$

- 167.** Ratio of moles of Fe (II) oxidised by equal volumes of equimolar KMnO_4 and $\text{K}_2\text{Cr}_2\text{O}_7$ solutions in acidic medium will be:
(a) 5 : 3 (b) 1 : 1 (c) 1 : 2 (d) 5 : 6
- 168.** The mass of a mixture containing HCl and H_2SO_4 is 0.1 g. On treatment with an excess of an AgNO_3 solution, reacted with this acid mixture gives 0.1435 g of AgCl. Mass % of the H_2SO_4 mixture is :
(a) 36.5 (b) 63.5 (c) 50 (d) None of these
- 169.** A solution of $\text{Na}_2\text{S}_2\text{O}_3$ is standardized iodometrically against 0.167 g of KBrO_3 . This process requires 50 mL of the $\text{Na}_2\text{S}_2\text{O}_3$ solution. What is the normality of the $\text{Na}_2\text{S}_2\text{O}_3$?
(a) 0.2 N (b) 0.12 N (c) 0.72 N (d) 0.02 N
- 170.** 0.80 g of impure $(\text{NH}_4)_2\text{SO}_4$ was boiled with 100 mL of a 0.2 N NaOH solution till all the $\text{NH}_3(\text{g})$ evolved. The remaining solution was diluted to 250 mL. 25 mL of this solution was neutralized using 5 mL of a 0.2 N H_2SO_4 solution. The percentage purity of the $(\text{NH}_4)_2\text{SO}_4$ sample is:
(a) 82.5 (b) 72.5 (c) 62.5 (d) 17.5
- 171.** The NH_3 evolved due to complete conversion of N from 1.12 g sample of protein was absorbed in 45 mL of 0.4 N HNO_3 . The excess acid required 20 mL of 0.1 N NaOH. The % N in the sample is:
(a) 8 (b) 16 (c) 20 (d) 25
- 172.** Find out % of oxalate ion in a given sample of an alkali metal oxalate salt, 0.30 g of it is dissolved in 100 mL water and its required 90 mL of centimolar KMnO_4 solution in acidic medium :
(a) 66% (b) 55% (c) 44% (d) 6.6%
- 173.** 320 mg of a sample of magnesium having a coating of its oxide required 20 mL of 0.1 M hydrochloric acid for the complete neutralisation of the latter. The composition of the sample is:
(a) 87.5% Mg and 12.5% MgO (b) 12.5% Mg and 87.5% MgO
(c) 80% Mg and 20% MgO (d) 20% Mg and 80% MgO
- 174.** The concentration of bivalent lead ions in a sample of polluted water that also contains nitrate ions is determined by adding solid sodium sulphate ($M = 142$) to exactly 500 mL water. Calculate the molarity of lead ions if 0.355 g of sodium sulphate was needed for complete precipitation of lead ions as sulphate.
(a) $1.25 \times 10^{-3} \text{ M}$ (b) $2.5 \times 10^{-3} \text{ M}$ (c) $5 \times 10^{-3} \text{ M}$ (d) None of these
- 175.** What volume of HNO_3 (sp. gravity 1.05 g mL⁻¹ containing 12.6 (w/w) of HNO_3) that reduce into NO is required to oxidise iron 1 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ in acid medium is:
(a) 70 mL (b) 0.57 mL (c) 80 mL (d) 0.65 mL
- 176.** The total volume of 0.1 M KMnO_4 solution that are needed to oxidize 100 mg each of ferrous oxalate and ferrous sulphate in a mixture in acidic medium is:
(a) 1.096 mL (b) 1.32 mL (c) 5.48 mL (d) none of these
- 177.** When 2.5 g of a sample of Mohr's salt reacts completely with 50 mL of $\frac{N}{10}$ KMnO_4 solution. The % purity of the sample of Mohr's salt is :
(a) 78.4 (b) 70 (c) 37 (d) 40
- 178.** 4 mole of a mixture of Mohr's salt and $\text{Fe}_2(\text{SO}_4)_3$ requires 500 mL of 1 M $\text{K}_2\text{Cr}_2\text{O}_7$ for complete oxidation in acidic medium. The mole % of the Mohr's salt in the mixture is:
(a) 25 (b) 50 (c) 60 (d) 75

- 179.** The equivalent mass of a metal is twice to that of oxygen. How many times is the equivalent mass of it's oxide than the equivalent mass of the metal ?
 (a) 1.5 (b) 2 (c) 3 (d) 4
- 180.** A metal oxide has the formula X_2O_3 . It can be reduced by hydrogen to give free metal and water. 0.1596 g of metal oxide requires 6 mg of hydrogen for complete reduction. The atomic mass of the metal (in amu) is:
 (a) 15.58 (b) 155.8 (c) 5.58 (d) 55.8
- 181.** Calculate the mass of anhydrous oxalic acid, which can be oxidised to $CO_2(g)$ by 100 mL of an MnO_4 solution, 10 mL of which is capable of oxidising 50 mL of 1N I₂ to I₂.
 (a) 45 g (b) 22.5 g (c) 30 g (d) 12.25 g
- 182.** A mixture of $NaHC_2O_4$ and $KHC_2O_4 \cdot H_2C_2O_4$ required equal volumes of 0.2 N $KMnO_4$ and 0.12 N NaOH separately. What is the molar ratio of $NaHC_2O_4$ and $KHC_2O_4 \cdot H_2C_2O_4$ in the mixture ?
 (a) 6 : 1 (b) 1 : 6 (c) 1 : 3 (d) 3 : 1
- 183.** Stannous sulphate ($SnSO_4$) and potassium permanganate are used as oxidising agents in acidic medium for oxidation of ferrous ammonium sulphate to ferric sulphate. The ratio of number of moles of stannous sulphate required per mole of ferrous ammonium sulphate to the number of moles of $KMnO_4$ required per mole of ferrous ammonium sulphate, is :
 (a) 2.5 (b) 0.2 (c) 0.4 (d) 2.0
- 184.** If a g is the mass of $NaHC_2O_4$ required to neutralize 100 mL of 0.2 M NaOH and b g that required to reduce 100 mL of 0.02 M $KMnO_4$ in acidic medium, then:
 (a) $a = b$ (b) $2a = b$ (c) $a = 2b$ (d) None of these
- 185.** 2 mole, equimolar mixture of $Na_2C_2O_4$ and $H_2C_2O_4$ required V_1 L of 0.1 M $KMnO_4$ in acidic medium for complete oxidation. The same amount of the mixture required V_2 L of 0.2 M NaOH for neutralization. The ratio of V_1 to V_2 is:
 (a) 1 : 2 (b) 2 : 1 (c) 4 : 5 (d) 5 : 4
- 186.** A mixture containing 0.05 mole of $K_2Cr_2O_7$ and 0.02 mole of $KMnO_4$ was treated with excess of KI in acidic medium. The liberated iodine required 1.0 L of $Na_2S_2O_3$ solution for titration. Concentration of $Na_2S_2O_3$ solution was:
 (a) 0.40 mol L⁻¹ (b) 0.20 mol L⁻¹ (c) 0.25 mol L⁻¹ (d) 0.30 mol L⁻¹
- 187.** 25 mL of 2 N HCl, 50 mL of 4 N HNO_3 and x mL of 2 M H_2SO_4 are mixed together and the total volume is made up to 1 L after dilution. 50 mL of this acid mixture completely reacted with 25 mL of a 1 N Na_2CO_3 solution. The value of x is:
 (a) 250 mL (b) 62.5 mL (c) 100 mL (d) None of these
- 188.** In an iodometric estimation, the following reactions occur
 $2Cu^{2+} + 4I^- \rightarrow Cu_2I_2 + I_2$; $I_2 + 2Na_2S_2O_3 \rightarrow 2NaI + Na_2S_4O_6$
 0.12 mole of $CuSO_4$ was added to excess of KI solution and the liberated iodine required 120 mL of hypo. The molarity of hypo solution was:
 (a) 2 (b) 0.20 (c) 0.1 (d) 1.0
- 189.** 1 g mixture of equal number of mole of Li_2CO_3 and other metal carbonate (M_2CO_3) required 21.6 mL of 0.5 N HCl for complete neutralisation reaction. What is the approximate atomic mass of the other metal ?
 (a) 25 (b) 23 (c) 51 (d) 118

- 190.** 32 g of a sample of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ were dissolved in dilute sulphuric acid and water and its volume was made up to 1 litre. 25 mL of this solution required 20 mL of 0.02 M KMnO_4 solution for complete oxidation. Calculate the mass % of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ in the sample.
(a) 34.75 (b) 69.5
(c) 89.5 (d) None of these
- 191.** In the mixture of $(\text{NaHCO}_3 + \text{Na}_2\text{CO}_3)$, volume of HCl required is x mL with phenolphthalein indicator and y mL with methyl orange indicator in the same titration. Hence, volume of HCl for complete reaction of Na_2CO_3 is:
(a) $2x$ (b) y
(c) $x/2$ (d) $(y - x)$
- 192.** 0.1 g of a solution containing Na_2CO_3 and NaHCO_3 requires 10 mL of 0.01 N HCl for neutralization using phenolphthalein as an indicator. Mass % of Na_2CO_3 in solution is:
(a) 25 (b) 32
(c) 50 (d) None of these
- 193.** A mixture of NaOH and Na_2CO_3 required 25 mL of 0.1 M HCl using phenolphthalein as the indicator. However, the same amount of the mixture required 30 mL of 0.1 M HCl when methyl orange was used as the indicator. The molar ratio of NaOH and Na_2CO_3 in the mixture was:
(a) 2 : 1 (b) 1 : 2 (c) 4 : 1 (d) 1 : 4
- 194.** When 100 mL solution of NaOH and Na_2CO_3 was first titrated with $N/10$ HCl in presence of HPh, 17.5 mL were used till end point is obtained. After this end point MeOH was added and 2.5 mL of same HCl were required to attain new end point. The amount of NaOH in mixture is:
(a) 0.06 g per 100 mL (b) 0.06 g per 200 mL
(c) 0.05 g per 100 mL (d) 0.012 g per 200 mL
- 195.** 1 gram of a sample of CaCO_3 was strongly heated and the CO_2 liberated was absorbed in 100 mL of 0.5 M NaOH solution. Assuming 90% purity for the sample, how many mL of 0.5 M HCl would be required to react with the resulting solution to reach the end point in presence of phenolphthalein?
(a) 73 mL (b) 41 mL
(c) 82 mL (d) 100 mL
- 196.** A sample of pure sodium carbonate 0.318 g is dissolved in water and titrated with HCl solution. A volume of 60 mL is required to reach the methyl orange end point. Calculate the molarity of the acid.
(a) 0.1 M (b) 0.2 M
(c) 0.4 M (d) None of these
- 197.** 10 L of hard water required 5.6 g of lime for removing hardness. Hence temporary hardness in ppm of CaCO_3 is:
(a) 1000 (b) 2000
(c) 100 (d) 1
- 198.** 1 L of pond water contains 20 mg of Ca^{2+} and 12 mg of Mg^{2+} ions. What is the volume of a 2 N Na_2CO_3 solution required to soften 5000 L of pond water?
(a) 500 L (b) 50 L (c) 5 L (d) None of these

- 199.** One litre of a sample of hard water contain 4.44 mg CaCl_2 and 1.9 mg of MgCl_2 . What is the total hardness in terms of ppm of CaCO_3 ?
- (a) 2 ppm (b) 3 ppm
(c) 4 ppm (d) 6 ppm
- 200.** If hardness of water sample is 200 ppm, then select the incorrect statement :
- (a) Mass ratio of CaCO_3 to H_2O is $\frac{0.02}{100}$
(b) Mole ratio of CaCO_3 to H_2O is 3.6×10^{-5}
(c) Mass of CaCO_3 present in hard water is 0.2 g/L
(d) 1 miliequivalent of CaCO_3 present in 1 kg of hard water




Level 2

- A mixture of NH_4NO_3 and $(\text{NH}_4)_2\text{HPO}_4$ contain 30.40% mass per cent of nitrogen. What is the mass ratio of the two components in the mixture ?
 (a) 2 : 1 (b) 1 : 2 (c) 3 : 4 (d) 4 : 1
- What volume of 75% alcohol by weight ($d = 0.80 \text{ g/cm}^3$) must be used to prepare 150 cm^3 of 30% alcohol by mass ($d = 0.90 \text{ g/cm}^3$)?
 (a) 67.5 mL (b) 56.25 mL (c) 44.44 mL (d) None of these
- Calculate the number of millilitres of NH_3 (aq) solution ($d = 0.986 \text{ g/mL}$) contain 2.5% by mass NH_3 , which will be required to precipitate iron as $\text{Fe}(\text{OH})_3$ in a 0.8 g sample that contains 50% Fe_2O_3 .
 (a) 0.344 mL (b) 3.44 mL (c) 17.24 mL (d) 10.34 mL
- In the preparation of iron from haematite (Fe_2O_3) by the reduction with carbon

$$\text{Fe}_2\text{O}_3 + \text{C} \rightarrow \text{Fe} + \text{CO}_2$$
 how much 80% pure iron may be produced from 120 kg of 90% pure Fe_2O_3 ?
 (a) 94.5 kg (b) 60.48 kg (c) 116.66 kg (d) 120 kg
- A mineral consists of an equimolar mixture of the carbonates of two bivalent metals. One metal is present to the extent of 12.5% by mass. 2.8 g of the mineral on heating lost 1.32 g of CO_2 . What is the % by mass of the other metal ?
 (a) 87.5 (b) 35.71 (c) 65.11 (d) 23.21
- 6.2 g of a sample containing Na_2CO_3 , NaHCO_3 and non-volatile inert impurity on gentle heating loses 5% of its mass due to reaction $2\text{NaHCO}_3 \rightarrow \text{Na}_2\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2$. Residue is dissolved in water and formed 100 mL solution and its 10 mL portion requires 7.5 mL of 0.2 M aqueous solution of BaCl_2 for complete precipitation of carbonates.
 Determine mass (in gram) of Na_2CO_3 in the original sample.
 (a) 1.59 (b) 1.06 (c) 0.53 (d) None of these
- Nitric acid can be produced from NH_3 in three steps process given below

$$\begin{aligned} \text{(I)} \quad & 4\text{NH}_3(\text{g}) + 5\text{O}_2(\text{g}) \rightarrow 4\text{NO}(\text{g}) + 6\text{H}_2\text{O}(\text{g}) \\ \text{(II)} \quad & 2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{NO}_2(\text{g}) \\ \text{(III)} \quad & 3\text{NO}_2(\text{g}) + \text{H}_2\text{O}(\text{l}) \rightarrow 2\text{HNO}_3(\text{aq}) + \text{NO}(\text{g}) \end{aligned}$$
 percent yield of 1st, 2nd and 3rd steps are respectively 50%, 60% and 80% respectively then what volume of $\text{NH}_3(\text{g})$ at 1 atm and 0°C required to produced 1575 g of HNO_3 .
 (a) 156.25 (b) 350 L (c) 3500 L (d) None of these
- 1 M NaOH solution was slowly added into 1000 mL of 183.75 g impure H_2SO_4 solution and the following plot was obtained. The percentage purity of H_2SO_4 sample and slope of the curve respectively are:



- (a) 75%, $\frac{1}{3}$ (b) 80%, $\frac{1}{2}$ (c) 80%, 1 (d) None of these
9. MnO_2 on ignition converts into Mn_3O_4 . A sample of pyrolusite having 75% MnO_2 , 20% inert impurities and rest water is ignited in air to constant mass. What is the percentage of Mn in the ignited sample ?
 (a) 24.6% (b) 37% (c) 55.24% (d) 74.05%
10. A 1.0 g sample of a pure organic compound containing chlorine is fused with Na_2O_2 to convert chlorine to NaCl . The sample is then dissolved in water, and the chloride precipitated with AgNO_3 , giving 1.96 g of AgCl . If the molecular mass of organic compound is 147, how many chlorine atoms does each molecule contain ?
 (a) 1 (b) 2 (c) 3 (d) 4
11. A 0.60 g sample consisting of only CaC_2O_4 and MgC_2O_4 is heated at 500°C , converting the two salts of CaCO_3 and MgCO_3 . The sample then weighs 0.465 g. If the sample had been heated to 900°C , where the products are CaO and MgO . What would the mixtures of oxides have weighed ?
 (a) 0.12 g (b) 0.21 g (c) 0.252 g (d) 0.3 g
12. A metal M forms the sulphate $M_2(\text{SO}_4)_3$. A 0.596 gram sample of the sulphate reacts with excess BaCl_2 to give 1.220 g BaSO_4 . What is the atomic mass of M ?
 (Atomic mass : $\text{S} = 32$, $\text{Ba} = 137.3$)
 (a) 26.9 (b) 69.7 (c) 55.8 (d) 23
13. Urea (H_2NCONH_2) is manufactured by passing $\text{CO}_2(\text{g})$ through ammonia solution followed by crystallization. CO_2 for the above reaction is prepared by combustion of hydrocarbons. If combustion of 236 kg of a saturated hydrocarbon ($\text{C}_n\text{H}_{2n+2}$) produces as much CO_2 as required for production of 999.6 kg urea then molecular formula of hydrocarbon is:
 (a) $\text{C}_{10}\text{H}_{22}$ (b) $\text{C}_{12}\text{H}_{26}$ (c) $\text{C}_{13}\text{H}_{28}$ (d) C_8H_{18}
14. 11.6 g of an organic compound having formula $\text{C}_n\text{H}_{2n+2}$ is burnt in excess of $\text{O}_2(\text{g})$ initially taken in a 22.41 litre steel vessel. Before reaction the gaseous mixture was at 273 K with pressure reading 2 atm. After complete combustion and loss of considerable amount of heat, the mixture of product and excess of O_2 had a temperature of 546 K and 4.6 atm pressure. The formula of organic compound is:
 (a) C_2H_6 (b) C_3H_8 (c) C_5H_{12} (d) C_4H_{10}

15. $\text{H}_2\text{O}_2 + 2\text{KI} \xrightarrow{40\% \text{ yield}} \text{I}_2 + 2\text{KOH}$
 $\text{H}_2\text{O}_2 + 2\text{KMnO}_4 + 3\text{H}_2\text{SO}_4 \xrightarrow{50\% \text{ yield}} \text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 3\text{O}_2 + 4\text{H}_2\text{O}$
 150 mL of H_2O_2 sample was divided into two parts. First part was treated with KI and formed KOH required 200 mL of $M/2 \text{ H}_2\text{SO}_4$ for neutralisation. Other part was treated with KMnO_4 yielding 6.74 litre of O_2 at 1 atm. and 273 K. Using % yield indicated find volume strength of H_2O_2 sample used.
 (a) 5.04 (b) 10.08 (c) 3.36 (d) 33.6
16. SO_2Cl_2 (sulphuryl chloride) reacts with water to give a mixture of H_2SO_4 and HCl . What volume of 0.2 M Ba(OH)_2 is needed to completely neutralize 25 mL of $0.2 \text{ M SO}_2\text{Cl}_2$ solution:
 (a) 25 mL (b) 50 mL (c) 100 mL (d) 200 mL
17. 5 g sample contain only Na_2CO_3 and Na_2SO_4 . This sample is dissolved and the volume made up to 250 mL. 25 mL of this solution neutralizes 20 mL of $0.1 \text{ M H}_2\text{SO}_4$. Calculate the % of Na_2SO_4 in the sample.
 (a) 42.4 (b) 57.6 (c) 36.2 (d) none of these
18. 20 mL of 0.2 M NaOH(aq) solution is mixed with 35 mL of 0.1 M NaOH(aq) solution and the resultant solution is diluted to 100 mL. 40 mL of this diluted solution reacted with 10% impure sample of oxalic acid ($\text{H}_2\text{C}_2\text{O}_4$). The mass of impure sample is:
 (a) 0.15 gram (b) 0.135 gram (c) 0.59 gram (d) None of these
19. A silver coin weighing 11.34 g was dissolved in nitric acid. When sodium chloride was added to the solution all the silver (present as AgNO_3) was precipitated as silver chloride. The mass of the precipitated silver chloride was 14.35 g. Calculate the percentage of silver in the coin.
 (a) 4.8% (b) 95.2% (c) 90% (d) 80%
20. Two elements X (at. mass 16) and Y (at. mass 14) combine to form compounds A, B and C. The ratio of different masses of Y which combine with a fixed mass of X in A, B and C is 1 : 3 : 5. If 32 parts by mass of X combines with 84 parts by mass of Y in B, then in C, 16 parts by mass of X will combine with :
 (a) 14 parts by mass of Y (b) 42 parts by mass of Y
 (c) 70 parts by mass of Y (d) 84 parts by mass of Y
21. The conversion of oxygen to ozone occurs to the extent of 15% only. The mass of ozone that can be prepared from 67.2 L of oxygen at 1 atm and 273 K will be :
 (a) 14.4 g (b) 96 g (c) 640 g (d) 64 g
22. RH_2 (ion exchange resin) can replace Ca^{2+} ions in hard water as $\text{RH}_2 + \text{Ca}^{2+} \rightarrow \text{RCa} + 2\text{H}^+$. If 1 L of hard water after passing through RH_2 has $\text{pH} = 3$ then hardness in parts per million of Ca^{2+} is:
 (a) 20 (b) 10 (c) 40 (d) 100
23. 100 cm^3 of a solution of an acid (Molar mass = 98) containing 29.4 g of the acid per litre were completely neutralized by 90.0 cm^3 of aq. NaOH containing 20 g of NaOH per 500 cm^3 . The basicity of the acid is:
 (a) 3 (b) 2 (c) 1 (d) data insufficient

24. 20 mL of 0.1 M solution of compound $\text{Na}_2\text{CO}_3 \cdot \text{NaHCO}_3 \cdot 2\text{H}_2\text{O}$ is titrated against 0.05 M HCl. x mL of HCl is used when phenolphthalein is used as an indicator and y mL of HCl is used when methyl orange is the indicator in two separate titrations. Hence $(y - x)$ is:
- (a) 40 mL (b) 80 mL
(c) 120 mL (d) None of these
25. A sample containing HAsO_2 (mol. mass = 108) and weighing 3.78 g is dissolved and diluted to 250 mL in a volumetric flask. A 50 mL sample (aliquot) is withdrawn with a pipet and titrated with 35 mL of 0.05 M solution of I_2 . Calculate the percentage HAsO_2 in the sample:
- (a) 25% (b) 20%
(c) 10% (d) none of these
26. A mixture of FeO and Fe_2O_3 is completely reacted with 100 mL of 0.25 M acidified KMnO_4 solution. The resultant solution was then treated with Zn dust which converted Fe^{3+} of the solution to Fe^{2+} . The Fe^{2+} required 1000 mL of 0.10 M $\text{K}_2\text{Cr}_2\text{O}_7$ solution. Find out the weight % Fe_2O_3 in the mixture.
- (a) 80.85 (b) 19.15
(c) 50 (d) 89.41
27. To a 10 mL, 1 M aqueous solution of Br_2 , excess of NaOH is added so that all Br_2 is disproportionated to Br^- and BrO_3^- . The resulting solution is free from Br_2 , by extraction and excess of OH^- neutralised by acidifying the solution. The resulting solution is sufficient to react with 2 g of impure CaC_2O_4 ($M = 128$ g/mol) sample. The % purity of oxalate sample is:
- (a) 85.3% (b) 12.5%
(c) 90% (d) 64%
28. 0.10 g of a sample containing CuCO_3 and some inert impurity was dissolved in dilute sulphuric acid and volume made up to 50 mL. This solution was added into 50 mL of 0.04 M KI solution where copper precipitates as CuI and I^- is oxidized into I_3 . A 10 mL portion of this solution is taken for analysis, filtered and made up free I_3 and then treated with excess of acidic permanganate solution. Liberated iodine required 20 mL of 2.5 mM sodium thiosulphate solution to reach the end point. Determine mass percentage of CuCO_3 in the original sample.
- (a) 7.41 (b) 74.1
(c) 61.75 (d) None of these
29. 1 mole of equimolar mixture of ferric oxalate and ferrous oxalate requires x mole of KMnO_4 in acidic medium for complete oxidation. x is:
- (a) 0.5 mole (b) 0.9 mole
(c) 1.2 mole (d) 4.5 mole
30. An impure sample of sodium oxalate ($\text{Na}_2\text{C}_2\text{O}_4$) weighing 0.20 g is dissolved in aqueous solution of H_2SO_4 and solution is titrated at 70°C , requiring 45 mL of 0.02 M KMnO_4 solution. The end point is overrun, and back titration is carried out with 10 mL of 0.1 M oxalic acid solution. Find the % purity of $\text{Na}_2\text{C}_2\text{O}_4$ in sample:
- (a) 75 (b) 83.75
(c) 90.25 (d) None of these

31. 0.5 g mixture of $K_2Cr_2O_7$ and $KMnO_4$ was treated with excess of KI in acidic medium. Iodine liberated required 150 cm^3 of 0.10 N solution of thiosulphate solution for titration. Find the percentage of $K_2Cr_2O_7$ in the mixture:
- (a) 14.64 (b) 34.2
(c) 65.69 (d) 50
32. A 150 mL of solution of I_2 is divided into two unequal parts. I part reacts with hypo solution in acidic medium. 15 mL of 0.4 M hypo was consumed. II part was added with 100 mL of 0.3 M NaOH solution. Residual base required 10 mL of 0.3 M H_2SO_4 solution for complete neutralization. What was the initial concentration of I_2 ?
- (a) 0.08 M (b) 0.1 M
(c) 0.2 M (d) None of these
33. A mixture of H_2SO_4 and $H_2C_2O_4$ (oxalic acid) and some inert impurity weighing 3.185 g was dissolved in water and the solution made up to 1 litre. 10 mL of this solution required 3 mL of 0.1 N NaOH for complete neutralization. In another experiment 100 mL of the same solution in hot condition required 4 mL of 0.02 M $KMnO_4$ solution for complete reaction. The mass % of H_2SO_4 in the mixture was:
- (a) 40 (b) 50
(c) 60 (d) 80
34. During developing of an exposed camera film, one step involves in the following reaction :
- HO  OH $2AgBr$ $2OH$  O $2Ag$ $2H_2O$ $2Br$
- (Quinol)
- Which of the following best describes the role of quinol ?
- (a) It acts as an acid (b) It acts as reducing agent
(c) It acts as oxidant (d) It acts as a base
35. The concentration of an oxalic acid solution is ' x ' mol litre⁻¹. 40 mL of this solution reacts with 16 mL of 0.05 M acidified $KMnO_4$. What is the pH of ' x ' M oxalic acid solution ? (Assume that oxalic acid dissociates completely.)
- (a) 1.3 (b) 1.699
(c) 1 (d) 2



Level 3



Passage 1

Oleum is considered as a solution of SO_3 in H_2SO_4 , which is obtained by passing SO_3 in solution of H_2SO_4 . When 100 g sample of oleum is diluted with desired mass of H_2O then the total mass of H_2SO_4 obtained after dilution is known as % labelling in oleum.


For example, a oleum bottle labelled as '109% H_2SO_4 ' means the 109 g total mass of pure H_2SO_4 will be formed when 100 g of oleum is diluted by 9 g of H_2O which combines with all the free SO_3 present in oleum to form H_2SO_4 as $\text{SO}_3 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4$.

- What is the % of free SO_3 in an oleum that is labelled as '104.5% H_2SO_4 ' ?
(a) 10 (b) 20 (c) 40 (d) None of these
- 9.0 g water is added into oleum sample labelled as "112%" H_2SO_4 then the amount of free SO_3 remaining in the solution is : (STP 1 atm and 273 K)
(a) 14.93 L at STP (b) 7.46 L at STP (c) 3.73 L at STP (d) 11.2 L at STP
- If excess water is added into a bottle sample labelled as "112% H_2SO_4 " and is reacted with 5.3 g Na_2CO_3 , then find the volume of CO_2 evolved at 1 atm pressure and 300 K temperature after the completion of the reaction :
(a) 2.46 L (b) 24.6 L (c) 1.23 L (d) 12.3 L
- 1 g of oleum sample is diluted with water. The solution required 54 mL of 0.4 N NaOH for complete neutralization. The % of free SO_3 in the sample is :
(a) 74 (b) 26 (c) 20 (d) None of these



Passage 2

The strength of H_2O_2 is expressed in several ways like molarity, normality, % (w/V), volume strength, etc. The strength of "10 V" means 1 volume of H_2O_2 on decomposition gives 10 volumes of oxygen at 1 atm and 273 K or 1 litre of H_2O_2 gives 10 litre of O_2 at 1 atm and 273 K. The decomposition of H_2O_2 is shown as under :



H_2O_2 can act as oxidising as well as reducing agent. As oxidizing agent H_2O_2 is converted into H_2O and as reducing agent H_2O_2 is converted into O_2 . For both cases its n -factor is 2.

Normality of H_2O_2 solution = 2 Molarity of H_2O_2 solution

- What is the molarity of "11.2 V" H_2O_2 ?
(a) 1 M (b) 2 M (c) 5.6 M (d) 11.2 M
- What is the percentage strength (% w/V) of "11.2 V" H_2O_2 ?
(a) 1.7 (b) 3.4 (c) 34 (d) None of these

3. 20 mL of H_2O_2 solution is reacted with 80 mL of 0.05 M KMnO_4 in acidic medium then what is the volume strength of H_2O_2 ?
 (a) 2.8 (b) 5.6 (c) 11.2 (d) None of these
4. 40 g $\text{Ba}(\text{MnO}_4)_2$ (mol. mass = 375) sample containing some inert impurities in acidic medium completely reacts with 125 mL of "33.6 V" of H_2O_2 . What is the percentage purity of the sample?
 (a) 28.12% (b) 70.31% (c) 85% (d) None of these




Passage 3

A water is said to be a soft water if it produces sufficient foam with the soap and water that does not produce foam with soap is known as hard water. Hardness has been classified into two types (i) Temporary hardness (ii) Permanent hardness.



Temporary hardness is due to presence of calcium and magnesium bicarbonate. It is simply removed by boiling as





Temporary hardness can also be removed by addition of slaked lime, $\text{Ca}(\text{OH})_2$



Permanent hardness is due to presence of sulphates and chlorides of Ca, Mg, etc. It is removed by washing soda as



Permanent hardness also removed by ion exchange resin process as



The degree of hardness of water is measured in terms of ppm of CaCO_3 . 100 ppm means 100 g of CaCO_3 is present in 10^6 g of H_2O . If any other water sample which contain 120 ppm of MgSO_4 , hardness in terms of CaCO_3 is equal to 100 ppm.

1. One litre of a sample of hard water ($d = 1 \text{ g/mL}$) contains 136 mg of CaSO_4 and 190 mg of MgCl_2 . What is the total hardness of water in terms of CaCO_3 ?
 (a) 100 ppm (b) 200 ppm (c) 300 ppm (d) None of these
2. What is the mass of $\text{Ca}(\text{OH})_2$ required for 10 litre of water remove temporary hardness of 100 ppm due to $\text{Ca}(\text{HCO}_3)_2$?
 (a) 1.62 g (b) 0.74 g (c) 7.4 g (d) None of these
3. A 200 g sample of hard water is passed through the column of cation exchange resin, in which H^+ is exchanged by Ca^{2+} . The outlet water of column required 50 mL of 0.1 M NaOH for complete neutralization. What is the hardness of Ca^{2+} ion in ppm?
 (a) 250 ppm (b) 500 ppm (c) 750 ppm (d) 1000 ppm



Passage 4

$$\text{Equivalent mass} = \frac{\text{Molecular mass / Atomic mass}}{n\text{-factor}}$$

n -factor is very important in redox as well as non-redox reactions. With the help of n -factor we can predict the molar ratio of the reactant species taking part in reactions. The reciprocal of n -factor's ratio of the reactants is the molar ratio of the reactants.

In general n -factor of acid/base is number of moles of H^+/OH^- furnished per mole of acid/base. n -factor of a reactant is number of moles of electrons lost or gained per mole of reactant.

Example 1 :

1. In acidic medium : KMnO_4 ($n = 5$) Mn^{2+}
2. In neutral medium : KMnO_4 ($n = 3$) Mn^{2+}
3. In basic medium : KMnO_4 ($n = 1$) Mn^{6-}

Example 2 : FeC_2O_4 Fe^{3+} 2CO_2

Total number of moles of e^- lost by 1 mole of FeC_2O_4

1 1 2 3

1. n -factor of $\text{Ba}(\text{MnO}_4)_2$ in acidic medium is :
(a) 2 (b) 6 (c) 10 (d) None of these
2. Consider the following reaction.




What is the equivalent mass of H_3PO_2 ? (mol. wt. is M)

- (a) M (b) $M/2$ (c) $M/3$ (d) None of these
3. For the reaction, $\text{Fe}_{0.95}\text{O}$ (molar mass = M) Fe_2O_3 , what is the eq. mass of $\text{Fe}_{0.95}\text{O}$?
(a) $\frac{M}{0.85}$ (b) $\frac{M}{0.95}$ (c) $\frac{M}{0.8075}$ (d) None of these
4. In the reaction, $x\text{VO} + y\text{Fe}_2\text{O}_3 \rightarrow \text{FeO} + \text{V}_2\text{O}_5$, what is the value of x and y respectively?
(a) 1, 1 (b) 2, 3 (c) 3, 2 (d) None of these



Passage 5

Consider the following series of reactions :



1. How much Cl_2 is required to prepare 122.5 g of NaClO_4 by above sequential reactions ?
(a) 284 g (b) 213 g (c) 142 g (d) 71 g

2. How many moles of NaCl will be formed by using 1 mole Cl_2 and other reagents in excess ?
 (a) $\frac{1}{12}$ mole (b) 1.67 mole (c) 1.75 mole (d) 0.75 mole
3. How many moles of NaClO_3 obtained after the completion of reaction by taking 1 mole Cl_2 and other reagents in excess ?
 (a) $\frac{1}{3}$ mole (b) Zero (c) $\frac{1}{4}$ mole (d) 1 mole

One or More Answers is/are Correct



1. 1 g atom of nitrogen represents :
 (a) $6.02 \times 10^{23} \text{ N}_2$ molecules (b) 22.4 litre of N_2 at 1 atm and 273 K
 (c) 11.2 litre of N_2 at 1 atm and 273 K (d) 14 g of nitrogen
2. 1 g molecule of V_2O_5 contains :
 (a) 5 mole of oxygen atom (b) 2 mole of V atom
 (c) 1 mole of oxygen atom (d) 2.5 mole of oxygen atom
3. Select the dimensionless quantity(ies) :
 (a) vapour density (b) molality (c) specific gravity (d) mass fraction
4. Which of the following concentration terms is/are affected by a change in temperature?
 (a) Molarity (b) Molality (c) Normality (d) Specific gravity
5. Which of the following statements regarding the compound A_xB_y is/are correct?
 (a) 1 mole of A_xB_y contains 1 mole of A and 1 mole B
 (b) 1 equivalent of A_xB_y contains 1 equivalent of A and 1 equivalent of B
 (c) 1 mole of A_xB_y contains x moles of A and y moles of B
 (d) equivalent mass of A_xB_y = equivalent mass of A + equivalent mass of B
6. 1 mole of Ba(OH)_2 will exactly neutralize :
 (a) 0.5 mole HCl (b) 1 mole of H_2SO_4 (c) 1 mole of H_3PO_3 (d) 2 mole of H_3PO_2
7. The pair of species having different percentage (mass) of carbon is :
 (a) CH_3COOH and $\text{C}_6\text{H}_{12}\text{O}_6$ (b) CH_3COOH and $\text{C}_2\text{H}_5\text{OH}$
 (c) HCOOCH_3 and HCOOH (d) $\text{C}_2\text{H}_5\text{OH}$ and CH_3OCH_3
8. 30 mL of CH_3OH ($d = 0.8 \text{ g/cm}^3$) is mixed with 60 mL of $\text{C}_2\text{H}_5\text{OH}$ ($d = 0.92 \text{ g/cm}^3$) at 25°C to form a solution of density 0.88 g/cm^3 . Select the correct option :
 (a) Molarity and molality of resulting solution are 6.33 and 13.59 respectively
 (b) The mole fraction of solute and molality are 0.385 and 13.59 respectively
 (c) Molarity and % change in volume are 13.59 and zero respectively
 (d) Mole fraction of solvent and molality are 0.615 and 13.59 respectively
9. Which of the following is/are correct for 17 g/L of H_2O_2 solution?
 (a) Volume strengths is 5.6 at 273 K and 1 atm
 (b) Molarity of solution is 0.5 M
 (c) 1 mL of this solution gives 2.8 mL O_2 at 273 K and 2 atm
 (d) The normality of solution is 2 N

10. Solutions containing 23 g HCOOH is/are :
- 46 g of 70% $\frac{w}{V}$ HCOOH ($d_{\text{solution}} = 1.40 \text{ g/mL}$)
 - 50 g of 10 M HCOOH ($d_{\text{solution}} = 1 \text{ g/mL}$)
 - 50 g of 25% $\frac{w}{w}$ HCOOH
 - 46 g of 5 M HCOOH ($d_{\text{solution}} = 1 \text{ g/mL}$)
11. A sample of H_2O_2 solution labelled as "28 volume" has density of 265 g/L. Mark the correct option(s) representing concentration of same solution in other units :
- $M_{\text{H}_2\text{O}_2} = 2.5$
 - $\% \frac{w}{V} = 17$
 - Mole fraction of $\text{H}_2\text{O}_2 = 0.2$
 - $m_{\text{H}_2\text{O}_2} = 13.88$
12. A mixture of 100 mL of CO, CO_2 and O_2 was sparked. When the resulting gaseous mixture was passed through KOH solution, contraction in volume was found to be 80 mL. The composition of initial mixture may be (in the same order) :
- 30 mL, 60 mL, 10 mL
 - 30 mL, 50 mL, 20 mL
 - 50 mL, 30 mL, 20 mL
 - 20 mL, 70 mL, 10 mL
13. If 1 mole of H_3PO_4 reacts with 1 mole of X(OH)_2 as shown below:
- $$\text{H}_3\text{PO}_4 + \text{X(OH)}_2 \rightarrow \text{XHPO}_4 + 2\text{H}_2\text{O} \text{ then :}$$
- the equivalent mass of base is $\frac{\text{mol. mass}}{2}$
 - the eq. mass of H_3PO_4 is $\frac{98}{3}$
 - the resulting solution requires 1 mole NaOH for complete neutralization
 - minimum 1 mole of X(OH)_2 is required for complete neutralization of XHPO_4
14. In acidic medium dichromate ion oxidizes stannous ion as:
- $$x\text{Sn}^{2+} + y\text{Cr}_2\text{O}_7^{2-} + z\text{H}^+ \rightarrow a\text{Sn}^{4+} + b\text{Cr}^{3+} + c\text{H}_2\text{O}$$
- the value of $x : y$ is 1 : 3
 - the value of $x + y + z$ is 18
 - $a : b$ is 3 : 2
 - the value of $z + c$ is 7
15. When an equimolar mixture of Cu_2S and CuS is titrated with $\text{Ba(MnO}_4)_2$ in acidic medium, the final products contain Cu^{2+} , SO_2 and Mn^{2+} . If the mol. mass of Cu_2S , CuS and $\text{Ba(MnO}_4)_2$ are M_1 , M_2 and M_3 respectively then :
- eq. mass of Cu_2S is $\frac{M_1}{8}$
 - eq. mass of CuS is $\frac{M_2}{6}$
 - eq. mass of $\text{Ba(MnO}_4)_2$ is $\frac{M_3}{5}$
 - Cu_2S and CuS both have same equivalents in mixture

16. Which is/are **INCORRECT** statement ?

- (a) Equivalent mass of H_2PO_3 is 40.5.
- (b) Eq. mass of H_2PO_4 may be equal to molar mass or less than molar mass because it depends on the reaction.
- (c) KMnO_4 has maximum eq. mass in acidic medium.
- (d) Oxidation state of H in MgH_2 is greater than in H_2O_2 .

Match the Column



Column-I and **Column-II** contain four entries in each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I

- (A) 0.5 mole of $\text{SO}_2(\text{g})$
- (B) 1 g of $\text{H}_2(\text{g})$
- (C) 0.5 mole of $\text{O}_3(\text{g})$
- (D) 1 g molecule of $\text{O}_2(\text{g})$

Column-II

- (P) Occupy 11.2 L at 1 atm and 273 K
- (Q) Weighs 24 g
- (R) Total no. of atoms $1.5 N_A$
- (S) Weighs 32 g

2.

Column-I

- (A) 44 g CO_2 gas
- (B) 35.2 g of CH_4
- (C) 48 g of O_3 gas
- (D) 44 g of N_2O gas

Column-II

- (P) 1g molecule
- (Q) N_A molecule
- (R) $22 N_A$ electrons
- (S) 49.28 L at 1 atm and 273 K
- (T) N_A atoms of oxygen

3.

Column-I

[Atomic masses (M)]

Isotope-I	Isotope-II	Average
(A) (z 1)	(z 3)	z
(B) (z 1)	(z 3)	(z 2)
(C) z	3z	2z
(D) (z 1)	(z 1)	z

Column-II

(% composition of heavier isotope)

- (P) 25% by moles
- (Q) 50% by moles
- (R) % by mass dependent on z
- (S) 75% by mass

4.

Column-I

- (A) When Bi_2S_3 converted into Bi^5 and S
 (B) When $\text{Al}_2(\text{Cr}_2\text{O}_7)_3$ reduced into Cr^3 in acidic medium
 (C) When FeS_2 converted into Fe_2O_3 and SO_2
 (D) When $\text{Mn}(\text{NO}_3)_2$ converted into MnO_4^{2-} and NO

Column-II (*n*-factor)

- (P) 18
 (Q) 11
 (R) 2
 (S) 10

5.

Column-I

- (A) Eq. mass $\frac{\text{Molecular mass}}{33}$
 (B) Eq. mass $\frac{\text{Molecular mass}}{27}$
 (C) Eq. mass $\frac{\text{Molecular mass}}{28}$
 (D) Eq. mass $\frac{\text{Molecular mass}}{24}$

Column-II

- (P) When CrI_3 oxidises into $\text{Cr}_2\text{O}_7^{2-}$ and IO_4^-
 (Q) When $\text{Fe}(\text{SCN})_2$ oxidises into Fe^3 , SO_4^{2-} , CO_3^{2-} and NO_3^-
 (R) When NH_4SCN oxidizes into SO_4^{2-} , CO_3^{2-} and NO_3^-
 (S) When As_2S_3 oxidises into AsO_3 and SO_4^{2-}

6.

Column-I

- (A) P_2H_4 PH_3 P_4H_2
 (B) I_2 I IO_3
 (C) MnO_4^- Mn^{2+} H_2O Mn_3O_4 H
 (D) H_3PO_2 PH_3 H_3PO_3

Column-II

- (P) E $\frac{3M}{4}$
 (Q) E $\frac{3M}{5}$
 (R) E $\frac{15M}{26}$
 (S) E $\frac{5M}{6}$

7. A sample of raw material contain NaNO_3 . It contains some NaIO_3 also. The NaIO_3 can be used as a source of iodine, produced in the following reactions:



One litre of sample solution containing 396 g of NaIO_3 is treated with stoichiometric quantity of NaHSO_3 . Now a substantial amount of same solution is added to reaction mixture to bring about the reaction (2).

8.

S


Examine the statements carefully and mark the correct answer according to the instructions given below

- (A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

- For More Material Join: @JEEAdvanced_2024

- 6. STATEMENT-1 :** MnO_2 can act as an oxidizing agent as well as reducing agent.
STATEMENT-2 : Oxidation state of Mn in MnO_2 lies between highest and lowest oxidation state.
- 7. STATEMENT-1 :** Equivalent volume of H_2 is 11.2 L at 1 atm and 273 K.
STATEMENT-2 : $1/2$ mole H_2 has produced when 1 mole of $\text{H}^+(aq)$ accepted 1 mole of e^- .
- 8. STATEMENT-1 :** For the reaction in titration $\text{Na}_2\text{CO}_3 + \text{HCl} \rightarrow \text{NaCl} + \text{NaHCO}_3$, the suitable indicator is phenolphthalein.
STATEMENT-2 : Phenolphthalein provide its colour in acidic medium.
- 9. STATEMENT-1 :** $[\text{Fe}(\text{CN})_6]^{4-} \rightarrow \text{Fe}^{3+} + \text{CO}_2 + \text{NO}_3^-$, the equivalent mass of reactant is 3.74.
STATEMENT-2 : Equivalent mass of reactant $\frac{\text{Mol. mass}}{61}$.
- 10. STATEMENT-1 :** In the balanced redox reaction,

$$x\text{As}_2\text{S}_3 + y\text{NO}_3^- + 4\text{H}_2\text{O} \rightarrow a\text{AsO}_4^{3-} + b\text{NO} + c\text{SO}_4^{2-} + 8\text{H}^+$$
the n -factor of As_2S_3 and NO_3^- is 28 and 3 respectively.
STATEMENT-2 : Molar ratio is reciprocal of n -factor's ratio so $x : y$ is 3 : 28.
- 11. STATEMENT-1 :** In the given reaction, $\text{NaOH} + \text{H}_3\text{PO}_4 \rightarrow \text{NaH}_2\text{PO}_4 + \text{H}_2\text{O}$ equivalent mass of H_3PO_4 is $M/3$
STATEMENT-2 : H_3PO_4 is tribasic acid.
- 12. STATEMENT-1 :** In CrO_5 oxidation number of Cr is +6.
STATEMENT-2 : CrO_5 has butterfly structure in which peroxide bonds are present.





- 13. STATEMENT-1 :** $\text{I}_2 + \text{IO}_3^- + \text{I}^-$, is an example of a disproportionation reaction.
STATEMENT-2 : Oxidation number of I can vary from -1 to $+7$.
- 14. STATEMENT-1 :** Fluorine exists only in -1 oxidation state.
STATEMENT-2 : Fluorine has $2s^2 2p^5$ configuration.
- 15. STATEMENT-1 :** H_2SO_4 can not act as reducing agent.
STATEMENT-2 : Sulphur can not increase its oxidation number beyond $+6$.

Subjective Problems




- What volume of a liquid (in L) will contain 10 mole ? If molar mass of liquid is 280 and its density is 1.4 g/mL.
- 16 g of SO_x gas occupies 5.6 L at 1 atm and 273 K. What will be the value of x ?
- 200 mL of 1M HCl is mixed with 300 mL of 6M HCl and the final solution is diluted to 1000 mL. Calculate molar concentration of $[\text{H}^+]$ ion.
- $\text{N}_2(g)$ reacts with $\text{H}_2(g)$ in either of the following ways depending upon supply of $\text{H}_2(g)$:



If 5 L $\text{N}_2(\text{g})$ and 3 L $\text{H}_2(\text{g})$ are taken initially (at same temperature and pressure), calculate the contraction in volume after the reaction (in L).

5. One commercial system removes SO_2 emission from smoke at 95 °C by the following set of reaction :



How many grams of CaSO_4 may be produced from 3.78 g of SO_2 ?

6. W is the mass of iron (in g) which will be converted into Fe_3O_4 by the action of 18 g of steam on it. What is the value of $W/7$?
- $$3\text{Fe} + 4\text{H}_2\text{O} \rightarrow \text{Fe}_3\text{O}_4 + 4\text{H}_2$$
7. Calculate the total moles of atoms of each element present in 122.5 g of KClO_3 .
8. On dissolving 2.0 g of metal in sulphuric acid, 4.51 g of the metal sulphate was formed. The specific heat of the metal is $0.057 \text{ cal g}^{-1} \text{ }^\circ\text{C}^{-1}$. What is the valency of the metal ?
9. One gram of a metallic chloride was found to contain 0.835 g of chlorine. Its vapour density is 85.5. If its molecular formula is M_xCl_y , then what is value of (x/y) ?
10. 0.7875 g of crystalline barium hydroxide is dissolved in water. For the neutralization of this solution 20 mL of $N/4 \text{ HNO}_3$ is required. How many moles of water of crystallization are present in one mole of this base ? (Given : Atomic mass Ba = 137, O = 16, N = 14, H = 1)
11. 2.0 g of polybasic organic acid (Molecular mass = 600) required 100 mL of a $\frac{M}{6}$ NaOH solution for complete neutralisation. Find the basicity of acid.
12. A mixture contains 1.0 mole each of NaOH, Na_2CO_3 and NaHCO_3 . When half of mixture is titrated with HCl, it required x mole of HCl in presence of phenolphthalein. In another experiment, half of mixture required y mole of same HCl in presence of methyl orange. Find the value of (x/y) .
13. When BrO_3^- ion reacts with Br^- ion in acidic medium, Br_2 is liberated. Calculate the ratio of molecular mass and equivalent mass of KBrO_3 .
14. A volume of 12.5 mL of 0.05 M SeO_2 reacts with 25 mL of 0.1 M CrSO_4 which is oxidised to Cr^{3+} . To what oxidation state was the selenium converted by the reaction ?
15. A 0.276 g impure sample of copper ore is dissolved and Cu^{2+} is titrated with KI solution. I_2 liberated required 40 mL of 0.1 M $\text{Na}_2\text{S}_2\text{O}_3$ solution for titration. What is the % of impurities in the ore ?
16. A sample of 28 mL of $\text{H}_2\text{O}_2(\text{aq})$ solution required 10 mL of 0.1 M $\text{KMnO}_4(\text{aq})$ solution for complete reaction in acidic medium. What is the volume strength of H_2O_2 ?
17. For the redox reaction given below, what is the value of $\frac{x}{z}$?
- $$x\text{NO}_3^- + y\text{As}_2\text{S}_3 + z\text{H}_2\text{O} \rightarrow \dots \text{AsO}_4^{3-} + \dots \text{NO} + \dots \text{SO}_4^{2-} + \dots \text{H}^+$$
18. On heating 0.220 g of a metallic oxide in presence of hydrogen, 0.045 g of water is formed. If the equivalent mass of the metal is E , then what is the value of $E/9$?

19. 10 g mixture of $K_2Cr_2O_7$ and $KMnO_4$ was treated with excess of KI in acidic medium. Iodine liberated required 100 cm^3 of 2.2 N sodium thiosulphate solution for titration. If the mass percent of $KMnO_4$ in the mixture Z , then what is the value of $2Z/5$?
20. In an ore the only oxidisable material is Sn^{2+} . This ore is titrated with a dichromate solution containing $2.5\text{ g } K_2Cr_2O_7$ in 0.50 litre . A 0.40 g of sample of the ore required 10.0 cm^3 of the titrant to reach equivalent point. If the percentage of tin in ore is x , then what is the value of $x/5$? ($K = 39.1$, $Cr = 52$, $Sn = 118.7$)

ANSWERS

Level 1

1. (c)	2. (a)	3. (b)	4. (d)	5. (c)	6. (d)	7. (c)	8. (c)	9. (b)	10. (b)
11. (d)	12. (b)	13. (b)	14. (a)	15. (d)	16. (d)	17. (d)	18. (d)	19. (d)	20. (a)
21. (a)	22. (c)	23. (b)	24. (b)	25. (a)	26. (d)	27. (d)	28. (c)	29. (c)	30. (c)
31. (b)	32. (a)	33. (a)	34. (a)	35. (b)	36. (d)	37. (b)	38. (d)	39. (c)	40. (a)
41. (c)	42. (d)	43. (d)	44. (a)	45. (b)	46. (a)	47. (c)	48. (d)	49. (a)	50. (a)
51. (c)	52. (c)	53. (c)	54. (a)	55. (c)	56. (a)	57. (d)	58. (c)	59. (d)	60. (c)
61. (c)	62. (d)	63. (d)	64. (b)	65. (d)	66. (b)	67. (d)	68. (a)	69. (c)	70. (d)
71. (c)	72. (a)	73. (d)	74. (d)	75. (d)	76. (b)	77. (a)	78. (a)	79. (b)	80. (c)
81. (c)	82. (d)	83. (a)	84. (b)	85. (c)	86. (d)	87. (a)	88. (d)	89. (d)	90. (b)
91. (c)	92. (a)	93. (a)	94. (c)	95. (a)	96. (b)	97. (b)	98. (c)	99. (c)	100. (c)
101. (d)	102. (c)	103. (a)	104. (c)	105. (a)	106. (b)	107. (b)	108. (b)	109. (b)	110. (a)
111. (b)	112. (d)	113. (c)	114. (d)	115. (b)	116. (c)	117. (c)	118. (a)	119. (b)	120. (b)
121. (c)	122. (a)	123. (b)	124. (d)	125. (b)	126. (d)	127. (b)	128. (c)	129. (a)	130. (c)
131. (c)	132. (a)	133. (d)	134. (b)	135. (d)	136. (d)	137. (a)	138. (a)	139. (a)	140. (d)
141. (a)	142. (c)	143. (b)	144. (b)	145. (d)	146. (d)	147. (d)	148. (b)	149. (b)	150. (a)
151. (b)	152. (d)	153. (c)	154. (c)	155. (a)	156. (a)	157. (b)	158. (b)	159. (c)	160. (d)
161. (a)	162. (b)	163. (c)	164. (c)	165. (c)	166. (c)	167. (d)	168. (b)	169. (b)	170. (a)
171. (c)	172. (a)	173. (a)	174. (c)	175. (b)	176. (c)	177. (a)	178. (d)	179. (a)	180. (d)
181. (b)	182. (d)	183. (a)	184. (d)	185. (c)	186. (a)	187. (b)	188. (d)	189. (d)	190. (b)
191. (a)	192. (d)	193. (c)	194. (a)	195. (c)	196. (a)	197. (a)	198. (c)	199. (d)	200. (d)



Level 2

1. (a)	2. (a)	3. (d)	4. (a)	5. (d)	6. (b)	7. (c)	8. (c)	9. (c)	10. (b)
11. (c)	12. (a)	13. (b)	14. (d)	15. (d)	16. (b)	17. (b)	18. (a)	19. (b)	20. (c)
21. (a)	22. (a)	23. (a)	24. (b)	25. (a)	26. (a)	27. (d)	28. (b)	29. (b)	30. (b)
31. (a)	32. (b)	33. (a)	34. (b)	35. (c)					



Level 3

Passage-1	1. (b)	2. (c)	3. (c)	4. (b)
Passage-2	1. (a)	2. (b)	3. (b)	4. (b)
Passage-3	1. (c)	2. (b)	3. (b)	
Passage-4	1. (c)	2. (a)	3. (a)	4. (b)
Passage-5	1. (a)	2. (c)	3. (b)	

One or More Answers is/are correct

1. (c,d) 2. (a,b) 3. (a,c,d) 4. (a,c,d) 5. (b,c,d) 6. (b,c,d) 7. (b,c) 8. (b,d)
 9. (a,b,c) 10. (a,b) 11. (a,c,d) 12. (a,b) 13. (a,c) 14. (b,c,d) 15. (a,b) 16. (a,c,d)

Match the Column

- | | | | |
|---------------|----------|------------|--------------|
| 1. A P, R, S; | B P; | C P, Q, R; | D S |
| 2. A P, Q, R; | B R, S ; | C P, Q; | D P, Q, R, T |
| 3. A P, R; | B Q, R; | C Q, S; | D Q, R |
| 4. A S; | B P; | C Q; | D R |
| 5. A Q; | B P; | C S; | D R |
| 6. A S; | B Q; | C R; | D P |
| 7. A S; | B P; | C Q; | D R |
| 8. A P, R; | B P; | C Q; | D Q, S |

Assertion-Reason Type Questions

1. (A) 2. (B) 3. (C) 4. (A) 5. (D) 6. (A) 7. (A) 8. (C) 9. (D) 10. (A)
 11. (D) 12. (A) 13. (B) 14. (B) 15. (A)

Subjective Problems

1. 2	2. 2	3. 2	4. 6	5. 8	6. 6	7. 5	8. 3	9. 5	10. 8
11. 5	12. 3	13. 5	14. 0	15. 8	16. 1	17. 7	18. 4	19. 6	20. 3

Hints and Solutions

Level 1

1. (c) Number of neutrons $(17 - 8) \times 12 \times 10^{25}$

$$9 \times 2 \times 6 \times 10^{23} \times 100 \times 1800 N_A$$

2. (a) Number of nucleons = mass number.

Molar mass of element

$$3.32 \times 10^{23} \times 6.02 \times 10^{23}$$

$$20 \times \text{mass number.}$$

So, number of nucleons in 2 atoms = 40

3. (b) Moles of PO_4^{3-} $\frac{9.5}{95} \times 0.1 \text{ mol}$

Number of electrons in $0.1 \text{ mol PO}_4^{3-}$

$$50 \times 0.1 \times N_A \times 5 N_A$$

4. (d) 126 amu HNO_3 2 molecules of HNO_3 6 atoms of O

$$\text{Moles of O-atom} = \frac{6}{N_A}$$

5. (c) 96 amu of S^{2-} 3 S^{2-} ions

Charge on 3 S^{2-} ions 3×2 charge of 2 electrons $3 \times 2 \times 1.6 \times 10^{19} \times 9.6 \times 10^{19} \text{ C}$

6. (d) Number of sodium atoms $\frac{46}{23} N_A$ $2 N_A$

Number of calcium atoms = Number of sodium atoms $2 N_A$

$$\text{Mass of Ca atoms} = \frac{2 N_A}{N_A} \times 40 = 80 \text{ g}$$

7. (c) $54 \text{ mL H}_2\text{O}$ $54 \text{ g H}_2\text{O}$

$$\text{Moles of H}_2\text{O} = \frac{54}{18} = 3 \text{ mole}$$

$$\text{Number of neutrons} = 8 \times 3 \times N_A = 24 N_A$$

8. (c) Moles of Mg^{2+} $\frac{48}{24} = 2 \text{ mole}$

$$\text{Number of electrons} = 2 \times 10 \times N_A = 20 N_A$$

9. (b) Moles of D_2O $\frac{5}{20} \times \frac{1}{4}$

$$\text{Number of neutrons} = 10 \times \frac{1}{4} \times N_A = 2.5 N_A$$

10. (b) Mass of one molecule $300 \times 1.66 \times 10^{-24} \text{ g}$
 $4.98 \times 10^{-22} \text{ g}$

11. (d) Moles of $\text{C}_9\text{H}_8\text{O}_4$ $\frac{360 \times 10^{-3}}{180} = 2 \times 10^{-3}$

$$\text{Number of O-atoms} = 2 \times 10^{-3} \times N_A \times 4$$

$$8 \times 6.02 \times 10^{20} \times 4.81 \times 10^{21}$$

12. (b) $M = \frac{wRT}{PV} = \frac{20 \times 0.0821 \times 273}{5.6 \times 1} = 80$

13. (b) 1 mole $(\text{NH}_4)_3\text{PO}_4$

$$12 \text{ mole H-atoms} \quad 4 \text{ mole O-atoms}$$

$$6 \text{ mole H-atoms} \quad 2 \text{ mole O-atoms}$$

14. (a) Moles of O_3 $\frac{3 \times 8.21}{0.0821 \times 300} = 1$

$$\text{Moles of O-atoms} = 3$$

15. (d) Moles of atoms $\frac{3.011 \times 10^{22}}{6.02 \times 10^{23}} = \frac{1}{20}$

$$\text{Atomic mass} = \frac{\text{mass}}{\text{moles}} = \frac{1.15}{1/20} = 23$$

16. (d) Atomic mass of an element

$$x = 6.643 \times 10^{-23} \times N_A \times 40$$

$$\text{number of moles of } x = \frac{20 \times 1000}{40} = 500$$

17. (d) $3.9854 \times 10^{-23} \text{ g of element}$ 1 atom

$$1 \text{ g of element} = \frac{1}{3.9854 \times 10^{-23}}$$

$$2.509 \times 10^{22}$$

18. (d) 4 mole $\text{C}_2\text{H}_4\text{O}_2$ 16 mole H-atoms

$$16 \text{ g hydrogen}$$

19. (d) 12.044×10^{22} molecules of CO_2 0.2 mole

$$\text{CO}_2 \quad 0.4 \text{ mole O-atoms} \quad 0.4 N_A \text{ O-atoms}$$

20. (a) (I) 0.5 mole $\text{O}_3 = 24 \text{ g O}_3$;

(II) 0.5 g atom of oxygen = 8 g

$$\text{(III)} \quad \frac{3.011 \times 10^{23}}{6.022 \times 10^{23}} \times 32 = 16 \text{ g O}_2$$

$$\text{(IV)} \quad \frac{5.6}{22.4} \times 44 \text{ g CO}_2 = 11 \text{ g CO}_2$$

21. (a) Mass of a drop of water 0.0018 g

$$\text{Moles of water} = \frac{0.0018}{18} = 10^{-4} \text{ mole}$$

$$\text{Number of water molecules in one drop} = 10^{-4} N_A$$

$$\text{Number of water molecules in two drops} = 2 \times 10^{-4} \times 6.023 \times 10^{23} = 12.046 \times 10^{19}$$

22. (c) Number of protons in ${}_6\text{C}^{14} = 6$;
Number of neutrons in ${}_6\text{C}^{14} = 8$;

$$\text{As per given new atomic mass of } {}_6\text{C}^{14} = 12 + 4 = 16$$

(As the mass of e^- is negligible as compared to neutron and proton)

$$\% \text{ increase in mass} = \frac{16 - 14}{14} \times 100 = 14.28$$

23. (b) Mass of NaCl 25 $\frac{8.775}{100}$ 2.193 g

$$\text{Number of NaCl units} = \frac{2.193}{58.5} N_A = 2.25 \times 10^{22}$$

24. (b) Moles of $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ $\frac{25.6}{342.3}$ 0.0747

$$\text{Number of H-atoms} = 0.0747 \times 22 \times 6.023 \times 10^{23} = 9.91 \times 10^{23}$$

25. (a) Let there are x -atoms of N

$$\% \text{ of N} = 28.9 \times \frac{x \times 14}{194} \times 100 = x \times 4$$

26. (d) Mass of 1 molecule of water 18 amu
18 1.66×10^{-24} g

$$\text{Volume of 1 molecule} = \frac{\text{mass}}{\text{density}}$$

$$29.88 \times 10^{-24} \times 2.988 \times 10^{23} \text{ mL}$$

27. (d) Volume of gold foil 25 40 0.25 mm^3
 $250 \times 10^3 \text{ cm}^3$

$$\text{Mass of gold foil} = 19.32 \times 250 \times 10^3 \text{ g} = 4.83 \text{ g}$$

$$\text{No. of gold atoms} = \frac{4.83}{197} N_A = 1.47 \times 10^{22}$$

28. (c) Let total mass be

$$\begin{array}{c} 100 \text{ g} \\ \swarrow \quad \searrow \\ \text{N}_2 \quad \text{O}_2 \\ x \quad (100 - x) \end{array}$$

$$M_{\text{Avg.}} = \frac{\text{Total mass}}{\text{Total moles}}$$

$$29 = \frac{100}{\frac{x}{28} + \frac{(100 - x)}{32}} \quad x = 72.41 \%$$

29. (c) $d = \frac{PM}{RT}$

$$1.15 = \frac{\frac{740}{760} M}{0.0821 \times 300}$$

$$M = 29.09. \text{ Let mole fraction of } \text{N}_2 \text{ is } x.$$

$$29.09 = 28x + 32(1 - x);$$

$$\begin{array}{l} x = 0.7275 \\ \text{mass \%} \quad \text{mole \%} \quad \frac{\text{mol. mass of } \text{N}_2}{\text{average mol. mass of air}} \\ = 72.75 \quad \frac{28}{29.09} \quad 70.02 \end{array}$$

30. (c) No. of moles of CO_2 in 100 g mixture

$$\frac{66}{44} = 1.5$$

$$\text{No. of moles of } \text{H}_2 \text{ in 100 g mixture}$$

$$\frac{34}{2} = 17$$

$$M_{\text{average}} = \frac{100}{18.5} = 5.40$$

$$\text{V. D.} = \frac{5.4}{2} = 2.7$$

31. (b) Let 1 mole of mixture has x mole N_2O_4

$$2 \times 27.6 + x(92) = (1 + x)46; x = 0.2$$

32. (a) $PM = dRT$

$$M_{\text{Gas}} = \frac{2 \times 0.0821 \times 600}{2} = 49.26$$

$$\text{At const. } P \text{ and } T \quad d \propto M$$

$$\frac{d_{\text{Gas}}}{d_{\text{Ne}}} = \frac{M_{\text{Gas}}}{M_{\text{Ne}}} = \frac{49.26}{20} = 2.463 \quad 2.5$$

33. (a) Let mole % of ^{26}Mg be x

$$\frac{(21-x)25 + x(26)}{100} = 24.31$$

$$x = 10\%$$

34. (a) Let atomic weight of other isotope is M

$$114.82 = \frac{114.9041 + 95.72M}{100}$$

$$M = 112.94$$

35. (b) Moles of $\text{N}_2 = \frac{3.01}{6.02} = 0.5$

$$\text{Moles of } \text{O}_2 = \frac{32}{32} = 1 \text{ mol}$$

$$\text{Average molar mass} = \frac{5 \times 28 + 32 \times 1}{5 + 1} = \frac{172}{6} = 28.67$$

$$d_{\text{mix}} = \frac{PM_{\text{avg.}}}{RT} = \frac{3 \times 28.67}{\frac{1}{12} \times 860} = 1.2 \text{ g/L}$$

36. (d) Let mole fraction of O_2 is x

$$40 \times 32 + x \times 80 = (1+x) \times 40$$

$$\text{or } x = 5/6$$

$$a:b = x:(1-x) = \frac{5}{6}:\frac{1}{6}$$

When ratio is changed

$$M_{\text{mixture}} = 32 \times \frac{1}{6} + 80 \times \frac{5}{6} = 72$$

41. (c) Weight of calcium = $4 \times \frac{40}{100} = 1.6 \text{ g}$

43. (d) Weight % of $\text{N} = \frac{14}{17} \times 100 = 82.35$

44. (a) Molar mass of $x = 0.444 \times \text{molar mass of } y$

$$\text{Molar mass of } x = \frac{2.96}{12} \times 12$$

$$\text{Molar mass of } y = \frac{2.96}{0.444} \times 12 = 80$$

45. (b) Moles of $\text{Zn} = \frac{9.81}{65} = 0.15$

$$\text{Moles of } \text{Cr} = \frac{1.8 \times 10^{23}}{6.02 \times 10^{23}} = 0.3$$

$$\text{Moles of O} = 0.6$$

$$\text{Zn}_{0.15}\text{Cr}_{0.3}\text{O}_{0.6}$$



46. (a) $n = \frac{\text{Mass}}{\text{Molar mass}}$

$$0.0242 = \frac{1.657}{M} \quad M = 68.47$$

$$1 \times 32 = 68.47$$

$$X = 35.5$$

47. (c) Mass of oxide = Mass of metal

Mass of oxygen

$$2.74 - 1.53 = W_{\text{Oxygen}} = 1.21 \text{ g}$$

$$\text{Moles of V} = \frac{1.53}{51} = 0.03$$

$$\text{Moles of O} = \frac{1.21}{16} = 0.075$$

$$\text{V}_{0.03}\text{O}_{0.075}$$



48. (d)
- | | C | H | N | O |
|-------|---------------------|--------------------|----------------------|---------------------|
| Moles | $\frac{70.6}{12}$ | $\frac{4.2}{1}$ | $\frac{11.8}{14}$ | $\frac{13.4}{16}$ |
| Moles | $\frac{5.88}{0.84}$ | $\frac{4.2}{0.84}$ | $\frac{0.842}{0.84}$ | $\frac{0.84}{0.84}$ |

$$\text{Simple ratio} = 7 : 5 : 1 : 1$$



49. (a) Loss in mass is due to loss of water = 63.1

$$\% \text{ of } \text{H}_2\text{O} = 63 \times \frac{18x}{106 \times 18x} = 100 \times x = 10$$

50. (a) $\text{Sr}(\text{OH})_2 \cdot x\text{H}_2\text{O}$
- | | $\text{Sr}(\text{OH})_2$ | $x\text{H}_2\text{O}$ |
|--------|--------------------------|-----------------------|
| 6.85 g | 3.13 g | 3.72 g |

$$\% \text{ of water} = \frac{3.72}{6.85} \times 100 = \frac{18x}{121 \times 18x} \times 100$$

$$x = 8$$

51. (c) $\% \text{ of O} = \frac{16 \times 27}{(100 + 3 \times 310)} \times 100$
- $$= 41.94\%$$

52. (c) $\text{C}_x\text{H}_y\text{Cl}_z\text{O}_w$

$$\text{Millimoles of C} = \frac{41.21}{44}$$

$$\text{Millimoles of H} = \frac{5.63}{18} = 2$$

29.72 mg of Dieldrin contains $\frac{57.13}{25.31}$ 29.72 mg of AgCl

Millimoles of Cl $\frac{57.13}{25.31} \frac{29.72}{143.5}$

Weight of O 29.72 $\frac{41.21}{44}$ 12 $\frac{5.63}{18}$ 2 1 $\frac{57.13}{25.31} \frac{29.72}{143.5} \frac{35.5}{143.5}$

Weight of O 29.72 [11.239 0.6255 16.59]

Weight of oxygen 1.265 mg

n_C $\frac{41.21}{44}$ $\frac{41.21}{44}$ $\frac{16}{1.265}$ 11.84 12

n_H $\frac{5.63}{9}$ $\frac{5.63}{9}$ $\frac{16}{1.265}$ 7.9 8

n_O $\frac{1.265}{16}$ $\frac{1.265}{16}$ $\frac{16}{1.265}$ 1

n_{Cl} $\frac{57.13}{25.31} \frac{29.72}{143.5}$ $\frac{57.13}{25.31} \frac{29.72}{143.5}$ $\frac{16}{1.265}$ 5.95 6

53. (c) $d \frac{PM}{RT} M \frac{dRT}{P} = \frac{2.28}{1} \frac{0.0821}{1} \frac{300}{1} = 56.15 \text{ g/mol}$

E.F. $\frac{85.7}{12} : \frac{14.3}{1} 7.14 : 14.3 = 1 : 2$; E.F. is CH_2 ; M.F. $(CH_2)_n$

where $n \frac{56.15}{12 + 2} \sim 4$;

M.F. is C_4H_8

54. (a) Moles of C $n_{CO_2} \frac{2.64}{44} 0.06$

mass of C 0.72

Mole of H 2 Moles of H_2O 2 $\frac{1.26}{18} 0.14$ mass of H 0.14

Compound does not contains oxygen.

So EF $C_{0.06} H_{0.14} C_3H_7$

Lowest M.M. 43

55. (c) Formula of sulphate will be $—M SO_4 7H_2O$

% of metal $\frac{M}{M 96 7 18} 100 9.87$

M 24.31

56. (a) C H N
Moles $\frac{9}{12} \frac{1}{1} \frac{3.5}{14} n \frac{108}{54} 2$

$\frac{3}{4} : 1 : \frac{1}{4}$

3 : 4 : 1

- $\text{C}_3\text{H}_4\text{N}$ M.F. $(\text{C}_3\text{H}_4\text{N})_n$
 MF $(\text{C}_3\text{H}_4\text{N})_2$ $\text{C}_6\text{H}_8\text{N}_2$
57. (d)

	I	O
Moles	$\frac{254}{127}$	$\frac{80}{16}$
	2	5









 I_2O_5
58. (c) $A^4 \times B^2 \rightarrow A_2B_4$
 Empirical formula AB_2
59. (d) 100 g compound contain 43.8 g oxygen so 1 g oxygen present with $\frac{100}{43.8}$ g compound
 $\frac{16}{43.8} \times 4$ g or 4 mole oxygen present with $\frac{100}{16} \times 4 \sim 146$ g compound
 mol. mass of compound acid is 146.
60. (c) Mol. mass of $\text{XY}_2 = \frac{5}{0.05} = 100$
 Mol. mass of $\text{X}_2\text{Y}_3 = \frac{85}{3.011 \times 10^{23}} N_A = 170$
 Let molar mass of X and Y are a and b respectively
 $a + 2b = 100$
 $2a + 3b = 170$;
 $a = 40$;
 $b = 30$
61. (c) $\text{CO}_2 = \frac{88}{44} = 2$ mole of CO_2 2 mole of C
 $\text{H}_2\text{O} = \frac{36}{18} = 2$ mole of H_2O 4 mole of H
 Mass of C + Mass of H + Mass of O = 44
 $24 + 4 + x = 44$; $x = 16$
 mole of O = 1 and molecular formula is $\text{C}_2\text{H}_4\text{O}$.
62. (d) M.M. of X_2
 $\frac{\text{WRT}}{\text{PV}} = \frac{40 \times 10^3 \times 0.0821 \times 300}{1 \times 4.92 \times 10^3} = 200$
 Atomic mass of X = 100.

63. (d)

X	Y
$\frac{75.8}{75}$	$\frac{24.2}{16}$
1	$\frac{3}{2}$
2	3

 X_2Y_3
64. (b) $M = \frac{\text{WRT}}{\text{PV}} = \frac{12.4 \times 0.0821 \times 100}{8 \times 0.821} = 124$ P_4
65. (d) % Mn $\frac{55}{55 \times 16x} \times 100 = 64$
 $x = 1.93$
66. (b)

Ti	+	O_2	Ti _{1.44} O
$\frac{1.44}{48}$ mole		$\frac{x}{48(1.44)}$ mole	
$\frac{1.44}{48}$		$\frac{1.44x}{48(1.44)}$	
		$\frac{1}{16}$	
x		1.77 g	
67. (d) 3 molecules of O_2 1 molecules of CS_2
 6 molecules of O_2 2 molecules of CS_2
68. (a) Factor label method.
69. (c) $3\text{Fe}_2\text{O}_3(\text{s}) \rightarrow 2\text{Fe}_3\text{O}_4 + \frac{1}{2}\text{O}_2$
 480 g Fe_2O_3 provide 16 g O_2 . For loss of
 $0.04 \text{ g O}_2 = 0.04 \times \frac{480}{16} = 1.2 \text{ g Fe}_2\text{O}_3$
 % by mass of $\text{SiO}_2 = \frac{0.8}{2.0} \times 100 = 40\%$
70. (d) Mass of S_8 in sample = 160 g;
 Moles of $\text{S}_8 = \frac{160}{32 \times 8} = 0.625$
 Number of moles of O_2 required = 0.625 $\times 8$
 Volume of O_2 required = 22.4 $\times 5$
 Vol. of air required = 22.4 $\times 5 \times \frac{100}{21} = 533.33 \text{ L}$



79. (b) $\text{CO}_2 + 2\text{NaOH} \rightarrow \text{Na}_2\text{CO}_3 + \text{H}_2\text{O}$
 $n_{\text{NaOH}} = 1$; CO_2 present in mixture 0.5 and CO present 0.3 mole
 Moles of CO_2 obtained from CO 0.3, extra moles of NaOH required 0.3 2 0.6 mole
80. (c) Let mass of Ag_2O be x g
 $\text{Ag}_2\text{O} \quad 2\text{Ag} \quad \frac{1}{2}\text{O}_2$
 $232 \text{ g} \quad \frac{1}{2} \quad 32 \quad 16 \text{ g O}_2$
 $x \text{ g} \quad \frac{16}{232} \quad x \quad 0.104 \quad x \quad 1.508 \text{ g}$
 $\% \text{ of Ag}_2\text{O} = \frac{1.508}{1.6} \times 100 = 94.25\%$
81. (c) Number of moles of $\text{Ba}(\text{OH})_2 = \frac{20}{100} \times \frac{342}{171} = 0.4$ mole
 or moles of $\text{OH}^- = 0.4 \times 2 = 0.8$
 Number of moles of $\text{HNO}_3 = 1.2 \times 2 = 2.4$ mole
 Hence, the final solution is acidic due to presence of excess H^+
 $[\text{H}^+] = \frac{(2.4 - 0.8) \times 1000}{1200 + \frac{342}{0.57}} = 0.888 \text{ M}$
82. (d) Total moles of $\text{H}_2\text{SO}_4 = 0.1$ mole
 Total volume $\frac{150}{1.25} + \frac{400}{1.25} = \frac{550}{1.25} = 440$
 $M = \frac{0.1}{440} \times 1000 = \frac{1}{4.4} = 0.227 \text{ M}$
83. (a) $\text{Zn} + 2\text{HCl} \rightarrow \text{ZnCl}_2 + \text{H}_2$;
 moles of H_2 evolved 2
 moles of HCl required 4
 $\frac{V}{36.5} \times \frac{1.2}{4} = \frac{0.365}{4}$; $V = 333.33 \text{ mL}$
84. (b) $\text{C}_2\text{H}_6 + 3.5\text{O}_2 \rightarrow 2\text{CO}_2 + 3\text{H}_2\text{O}$;
 $\text{C}_2\text{H}_4 + 3\text{O}_2 \rightarrow 2\text{CO}_2 + 2\text{H}_2\text{O}$
 Let volume of ethane is x litre, $22.4 - x = 3.5x$ $4 = 3.5x$ $x = 1.14$ litre
 at constant T and P , $V \propto n$;
 Mole fraction of C_2H_6 in mixture $\frac{11.2}{28} = 0.4$
85. (c) $X_{\text{Sn}} = \frac{\frac{12.5}{209} + \frac{119}{207} + \frac{12.5}{119} + \frac{125}{112}}{\frac{12.5}{209} + \frac{119}{207} + \frac{12.5}{119} + \frac{125}{112}} = 0.176$
86. (d) Mass of solution 100 g
 Mass of solute 56 g
 moles of solute $\frac{56}{60} = 0.93$
 Mass of solvent 44 g

$$\text{moles of solvent} \quad \frac{44}{18} \quad 2.44$$

$$X_{\text{Solute}} \quad \frac{0.93}{0.93 + 2.44} \quad \frac{0.93}{3.37} \quad 0.276$$

87. (a) $\frac{\text{mass of Al}_2(\text{SO}_4)_3}{\text{mass of water}} \quad 10^6 \quad 34.2$

1 litre solution contains 1000 g of water

$$\text{In 1 litre solution, mass of Al}_2(\text{SO}_4)_3 \quad \frac{34.2}{10^6} \times 1000 \quad 34.2 \text{ mg}$$

$$\text{molarity of Al}_2(\text{SO}_4)_3 = \frac{34.2 \times 10^{-3}}{342} \text{ M} = 10^{-4} \text{ M}$$



$$10^{-4} \text{ M} \quad 2 \times 10^{-4} \text{ M} \quad 3 \times 10^{-4} \text{ M}$$

$$[\text{SO}_4^{2-}] = 3 \times 10^{-4} \text{ M}$$

88. (d) $M \quad \frac{n_{\text{Solute}}}{\text{Vol of solution (in mL)}} \quad 1000 \quad \dots(1)$

$$m \quad \frac{n_{\text{Solute}}}{\text{Weight of solvent (in g)}} \quad 1000 \quad \dots(2)$$

$$m \quad \frac{n_{\text{Solute}}}{\text{Weight of solution (in g)} - \text{wt of solute (in g)}} \quad 1000$$

$$m \quad \frac{M \quad \text{vol of solution (in mL)} \quad 1000}{1000 \quad (\text{wt of solution (in g)} - \text{wt of solute (in g)})}$$

$$m \quad \frac{M \quad 1000}{\text{Density of solution (g / mL)} \quad 1000 \quad \frac{\text{wt of solute} \quad M_{\text{Solute}} \quad 1000}{M_{\text{Solute}} \quad \text{volume of solution (in mL)}}$$

$$m \quad \frac{M \quad 1000}{1000 \quad M \cdot M_{\text{Solute}}}$$

89. (d) $m \quad \frac{1000 M}{1000 d \quad M \quad M_{\text{Solute}}}$

$$d \quad \frac{1000 \quad 9}{10} \quad 9 \quad 50 \quad \frac{1}{1000} \quad 1.35 \text{ g/mL}$$

90. (b) 1000 mL solution contain 2 mole of ethanol or 1000 mL solution contain 2 mole of ethanol

$$\text{Mass of solvent} \quad 1000 \quad 1.025 \quad 2 \quad 46$$

$$m \quad \frac{2}{1000 \quad 1.025 \quad 2 \quad 46} \quad 1000$$

$$m \quad \frac{2}{933} \quad 1000 \quad 2.143$$

91. (c) Total Cl $0.2 \quad 0.4 \text{ mole (from BaCl}_2) \quad 0.6 \text{ mole}$

$$M_{\text{Cl}} \quad \frac{0.6}{500} \quad 1000 \quad 1.2$$

92. (a) Mass of solution $10 \quad 1.2 \quad 12 \text{ g}$

$$\text{Mass of solute} \quad 12 \quad \frac{37}{100} \quad 4.44 \text{ g}$$

93. (a) The sample of H_2SO_4 is 80% by volume, $W_{\text{H}_2\text{SO}_4} \quad 800 \text{ g in 1000 mL}$

$$\text{Volume of solution} \quad 1000 \text{ mL}$$

$$\text{Mass of solution} = 1000.0 - 1.80 = 1800 \text{ g}$$

$$\text{Mass of water} = \frac{1800}{1000} \times 800 = 1440 \text{ g}$$

$$\text{Molality} = \frac{800}{98} \times \frac{1000}{1000} = 8.16 \text{ mol kg}^{-1}$$

$$94. \text{ (c) Initial molarity } (M_1) = \frac{10}{336} \times \frac{10^3}{500} \times \frac{1000}{0.0595} = 10^3$$

$$\frac{M_1 V_1}{M_2} = \frac{M_2 V_2}{V_2} \Rightarrow \frac{0.0595 \times 10^3 \times 1}{1000} = \frac{5.95 \times 10^8}{M_2}$$

$$95. \text{ (a) Molar mass of } \text{Pb}(\text{NO}_3)_2 = 25 \times 0.15 = 3.75 \text{ m. moles}$$

$$\text{Molar mass of } \text{Al}_2(\text{SO}_4)_3 = \frac{1}{3} \times 3.75 \times M = 20$$

$$M = 0.0625 \times 6.25 \times 10^2 = 3.91 \text{ M}$$

$$96. \text{ (b) m-moles of } \text{HNO}_3 \text{ required} = \frac{250}{1.2} \times \frac{300}{100} = 625$$

$$100 \text{ g solution contains } 63 \text{ g } \text{HNO}_3 \text{ or}$$

$$\frac{100}{1.4} \text{ mL solution contain } 1 \text{ mole } \text{HNO}_3$$

$$\text{molarity of } \text{HNO}_3 \text{ solution}$$

$$\frac{1000}{100} \times \frac{1.40}{14} = 1.4$$

$$14 \times V = 300 \text{ or } V = 21.42 \text{ mL}$$

$$97. \text{ (b) } 50 \text{ mL } \text{BaCl}_2 \text{ (aq) solution contain } 10.4 \text{ g } \text{BaCl}_2$$

$$n_{\text{BaCl}_2} = \frac{10.4}{137 + 71} = 0.05$$

$$\text{moles of } \text{H}_2\text{SO}_4 \text{ in } 100 \text{ mL } \text{H}_2\text{SO}_4 \text{ (aq) solution}$$

$$\frac{9.8}{98} = 0.1$$



$$\text{moles of HCl formed} = 0.1$$

$$[\text{Cl}^-] = \frac{0.1 \times 1000}{50 \times 100} = 0.666 \text{ M}$$

$$98. \text{ (c) No. of moles of HCl are more than NaOH, so final solution becomes acidic.}$$

$$99. \text{ (c) Let volume is } x \text{ mL, } x \times 0.1 \times 2 = 50 \times 0.1 \times 2$$

$$(x - 50) \times 0.05 = 2$$

$$x = 100 \text{ mL}$$

$$100. \text{ (c) } [\text{HCl}] = \frac{1}{5} \times \frac{4V}{V} \times \frac{2}{V} = 1.2 \text{ M}$$

$$101. \text{ (d) } M = \frac{55 \times 0.07 \times 20 + 0.12 \times 25 + 0.15 \times 25}{55 + 20 + 25} = 0.1$$

$$102. \text{ (c) Molarity} = \frac{28}{11.2} \times \frac{2.5}{1000} = 2.5 \text{ M}$$

$$1000 \text{ mL of solution contains } 2.5 \text{ mole } \text{H}_2\text{O}_2$$

$$\text{molality, } m = \frac{2.5}{1250} \times \frac{1000}{85} = 2.15 \text{ m}$$

$$103. \text{ (a) The reaction that takes place is}$$



$$\therefore 143.5 \text{ g of AgCl is produced from } 58.5 \text{ g NaCl}$$

$$14 \text{ g of AgCl will be produced from}$$

$$\frac{58.5}{143.5} \times 14 = 5.70 \text{ g NaCl}$$

$$\text{This is the amount of NaCl in common salt;}$$

$$\% \text{ purity} = \frac{5.70}{6} \times 100 = 95\%$$

$$104. \text{ (c) Let no. of moles of } \text{Al}_2(\text{SO}_4)_3 \text{ in solution is } x$$

$$\frac{x}{2684} \times \frac{1000}{342} = 1$$

$$2\text{Al}_2(\text{SO}_4)_3 \rightarrow 6 \text{ moles } \text{SO}_4^{2-}$$

$$105. \text{ (a) } 0.478 \text{ mL } 0.478 \text{ g of water; } 10^9 \text{ g water contain } 0.10 \text{ g } \text{CHCl}_3$$

$$0.478 \text{ g water contain}$$

$$\frac{0.1}{10^9} \times 0.478 \text{ g } \text{CHCl}_3$$

$$n_{\text{CHCl}_3} = \frac{0.1}{10^9} \times \frac{0.478}{119.5}$$

$$\text{No. of molecules} = \frac{0.1}{10^9} \times \frac{0.478}{119.5} \times N_A$$

$$4 \times 10^{13} \times N_A$$

$$106. \text{ (b) (i) Mass of NaOH} = 50 \times \frac{40}{100} = 20 \text{ g}$$

$$\text{(ii) } \frac{w}{v} \text{ NaOH} = 50 \text{ g NaOH in } 100 \text{ mL}$$

$$\text{So, in } 50 \text{ mL, mass of NaOH will be } 25 \text{ g}$$

$$\text{(iii) } 50 \text{ g of } 15 \text{ M NaOH (d = 1 g/mL)}$$

$$15 \text{ M NaOH} = 15 \text{ moles NaOH in } 1000 \text{ mL solution}$$

$$\text{Mass of solution} = 1000 \times 1 = 1000 \text{ g}$$

$$\text{Mass of NaOH} = 15 \times 40 = 600 \text{ g}$$

$$\text{In } 1000 \text{ g solution mass of NaOH} = 600 \text{ g}$$

$$\text{So, in } 50 \text{ g solution, mass of NaOH}$$

$$\frac{600}{1000} \times 50 = 30 \text{ g}$$

$$\text{Order (iii) > (ii) > (i)}$$

$$107. (b) B \text{ H}_2\text{PtCl}_6 \quad \text{Pt}; \quad \frac{12}{M_B} \quad \frac{410}{195}$$

moles of Pt; Molecular mass of base 58



Apply PoAC for Ag

$$\begin{array}{ccc} 2 & n_{\text{Salt}} & 1 & n_{\text{Ag}} \\ 2 & \frac{1}{216} & A & \frac{0.5934}{108} \end{array} \quad A \quad 148$$

i.e., Molar mass of acid (H_2A) 150,

Let % of C is x

$$\begin{array}{ccccccc} \text{C} & \text{H} & \text{O} & \text{wt. \% of H} & \frac{x}{8} & & \\ \text{Mass } x & \frac{x}{8} & 2x & \text{wt \% of O} & 2 & \% \text{ of} & \\ & & & & & \text{C} & 2x \end{array}$$

$$\begin{array}{ccc} \text{Moles } \frac{x}{12} & \frac{x}{8} & \frac{2x}{16} \\ \frac{x}{3} & \frac{x}{2} & \frac{x}{2} & \text{C}_2\text{H}_3\text{O}_3 & \text{E.F.} \end{array}$$

$$\text{M.F. } (\text{C}_2\text{H}_3\text{O}_3)_n \quad n \quad \frac{\text{M.M.}}{\text{EFM}} \quad \frac{150}{75} \quad 2$$



$$109. (b) \text{ Moles of Ag}_3\text{A} = \frac{\text{moles of Ag}}{3} = \frac{0.607}{M}$$

$$\frac{0.37}{108} \quad \frac{1}{3} \quad M \quad 531$$

mol. mass of H_3A = mol. mass of

$$\begin{array}{ccc} \text{Ag}_3\text{A} & 3 & \text{At. mass of Ag} + 3 \\ \text{H} & 210 & \text{At. mass of} \end{array}$$

110. (a) Equivalents of KOH used by oil

$$[25 \quad 0.40 \quad 8.5 \quad 0.28 \quad 2] \quad 10^3$$



$$\text{Moles of KOH used } 5.24 \quad 10^3$$

Mass of KOH used in milligrams

$$5.24 \quad 10^3 \quad 56 \quad 1000 \quad 223.44$$

$$\text{Saponification number } \frac{223.44}{2}$$

$$146.72$$



$$18 \quad (20 \ 10) \quad x$$

$$x \quad 12 \text{ mL} \quad \text{CO}$$

$$\text{H}_2 \quad 8 \text{ mL}$$

$$V_{\text{CO}} : V_{\text{H}_2}$$

$$12 : 8$$

$$3 : 2$$

112. (d) For 6 mole HCl, 3 mole $\text{H}_2(g)$ is produced

So, For 1 mole HCl, $\frac{1}{2}$ mole H_2 will be



produced

$$\frac{1}{2} \quad 22.4 \text{ L at STP} \quad 11.2 \text{ L H}_2 \text{ gas}$$

113. (c) 100 mL gaseous mixture contain 20 mL C_3H_8

So, volume of CH_4 and CO

$$(100 \ 20) \quad 80 \text{ mL}$$




80 mL (CH_4 and CO) will produce 80 mL CO_2 ; C_3H_8 will produce 3 20

$$60 \text{ mL CO}_2$$

Total CO_2 produced 80 60 140 mL

114. (d) Let the volume of CO, CH_4 and Ne be x, y & z respectively



$$x \quad x/2 \quad x$$



$$y \quad 2y \quad y$$

Remaining volume of O_2 15 $\frac{x}{2}$ 2y 1.5

Volume after reaction :

$$x \quad y \quad 9 \quad \dots(i)$$

$$x \quad y \quad z \quad 40 \quad \dots(ii)$$

by Eq. (i) and (ii)


Volume of CH_4 = 6 mL;

$$\% \text{ of CH}_4 = \frac{6}{40} \quad 100 \quad 15$$

115. (b) Suppose, the volume of propane V L



$$V \quad 5V \quad 3V$$



$$(3 \ V) \quad \frac{13}{2} (3 \ V) \quad 4(3 \ V)$$


$\therefore$ Total volume of CO_2 produced = 10 L;

$$\therefore 3V \ 4(3 \ V) \ 11; \quad V \ 1$$

Volume of butane (3 1) 2 L

Thus, the ratio of volume of butane to propane 2 : 1

116. (c) Hypophosphorous Acid — H_3PO_2




118. (a) S_8 , S_2F_2 , H_2S^2 , H_2SO_4

119. (b) Let oxidation state is x ;

$$(x \ 5 \ 0 \ 1) \ 2 \ 0 \ x \ 1$$

120. (b) The redox reaction is :



121. (c) NH_4NO_3

122. (a) Caro's acid H_2SO_5

= Peroxomono-sulphuric acid

Marshall's acid $\text{H}_2\text{S}_2\text{O}_8$


= Peroxodi-sulphuric acid

The oxidation state in both the acids is +6 each because oxidation state cannot be greater than the number of valence electrons.

123. (b) BaO_2 , KO_2 , O_3 , OF_2

124. (d) KO_3 , Na_2O_2

125. (b) $\text{Ba}(\text{H}_2\text{PO}_2)_2$, Ba^{2+} , H_2PO_2^-



Charge balance $2x + 3(0.96 - x) = 2$, 0

$$x = \frac{11}{12}$$

127. (b) HNO_3 , NO , N_2 , NH_4Cl

128. (c) 1 mole N_2H_4 loses 8 mole e^- ; 1 mole N loses 4 mole of e^-

New oxidation state of N is 2, 4, 2

129. (a) $\text{Cr}_2\text{O}_7^{2-}$, CrO_4^{2-}

130. (c) Mn^{2+} , $(\text{KNO}_3 \ \text{NaOH})$, MnO_4^{2-}

131. (c) Fe^{2+} , MnO_4^- , H^+ , Fe^{3+} , Mn^{2+} , H_2O

132. (a) Both Fe and C are oxidised and MnO_4^- is reduced.

133. (d) H_2O_2 it self be oxidised to O_2 .

134. (b) Metallic iron is oxidised to Fe^{3+} .


137. (a) $2\text{Na}[\text{Ag}(\text{CN})_2] + \text{Zn} \rightarrow$



138. (a) The balanced redox reaction is



139. (a) Balancing the equation, we have



140. (d) CS_2 , 3Cl_2 , CCl_4 , S_2Cl_2

141. (a) Balanced reaction is



Sum of stoichiometric coefficients

$$1 \ 5 \ 1 \ 4 \ 5 \ 16$$

142. (c) If $x = 4$



0.1 mole, 0.3 mole

If $x = 3$



0.1 mole, 0.2 mole

If $x = 2$



0.1 mole, 0.1 mole

So, if $x = 2$, 5.6 g KOH will be required.

143. (b) In acidic medium $\text{MnO}_4^- \rightarrow \text{Mn}^{2+}$

$$n_f = 5, \quad E = \frac{M}{5}$$

144. (b) $n = |1 \ 5 \ 2|$

145. (d) 14 mole HCl loses 6 mole e^- ;

1 mole HCl loses $\frac{6}{14}$ mole e^-

$$\text{eq. mass of HCl} = \frac{M}{\frac{6}{14}}$$

$$\frac{36.5 \times 14}{6} = 85.1$$

146. (d) n -factor $\frac{n_1 \ n_2}{n_1 \ n_2} = \frac{4 \ 2}{4 \ 2} = \frac{4}{3}$

$$\text{Eq. mass} = \frac{\text{mol mass}}{n\text{-factor}} = \frac{M}{4}$$

147. (d) n -factor $|2 \ 2 \ 8 \ 3| = 28$

148. (b) In SCl_2

71 g chlorine combines with 32 g S

So, 35.5 g chlorine will combine with 16 g

So, Eq. mass of S 16 g

As the mass which combines with 35.5 g chlorine is equivalent mass.

149. (b) E_{Element} 4, Formula of chloride $M\text{Cl}_x$


Eq. mass of chloride $E_{\text{Element}} E_{\text{Chlorine}}$

$$\frac{M_{\text{Mass}}}{n_f} E_{\text{Element}} E_{\text{Chlorine}}$$

$$\frac{59.25}{x} \cdot \frac{2}{4} = 35.5$$

$$x = 3$$

150. (a) $\text{K}_2\text{Cr}_2\text{O}_7 + X^{n+} \rightarrow X^{5+}\text{O}_3 + \text{Cr}^{3+}$



151. (b) Normality $\frac{1}{100}$ (Centinormal)

Number of eq. Normality Volume (in L)

$$\text{Number of eq.} \quad \frac{1}{100} \quad \frac{250}{1000} \quad \frac{1}{400}$$


$$\frac{\text{Mass}}{\text{Molar mass}} n_f \quad \frac{1}{400}$$

As R.A. $\text{C}_2\text{O}_4^{2-}$ CO_2

$$\frac{w}{126} \cdot \frac{2}{400} = \frac{1}{400}$$

$$w = 0.1575 \text{ g}$$

152. (d) As R.A.



153. (c) $E_{\text{Chloride}} E_{\text{Element}} E_{\text{Chlorine}}$

$$\frac{E_{\text{Chloride}}}{w} = 35.5$$

Metal is divalent, n_f for chloride 2

$$E \cdot \frac{M}{2} = M \cdot 2(w = 35.5) \quad 2w = 71$$

154. (c) $2\text{BrO}_3 + 12\text{H}^+ + 10\text{Br} \rightarrow 6\text{Br}_2 + 6\text{H}_2\text{O}$
10 mole e^- are required for formation of 6 moles of Br_2

$$n\text{-factor of Br}_2 = \frac{10}{6} = \frac{5}{3}$$

$$\text{eq. wt.} = \frac{\text{mol. wt.}}{n} = \frac{m}{5/3} = \frac{3m}{5}$$

155. (a) Eq. of A Eq. of B

$$\frac{m_A}{E_A} = \frac{m_B}{E_B}$$

156. (a) O.N. of N in N_2H_4 is -2 which changes to 0 in N_2 . Hence, equivalent mass of

$$\text{N}_2\text{H}_4 = \frac{\text{molar mass}}{2 \cdot 2}$$

O.N. of iodine changes from +5 in IO_3^- to 1 in ICl . Hence, equivalent mass of

$$\text{IO}_3 = \frac{\text{molar mass}}{4}$$

$$\text{157. (b) } N_R = \frac{N_1 V_1}{V_1} \cdot \frac{N_2 V_2}{V_2} = \frac{2 \cdot 0.45 \cdot 0.6}{3} = 0.5$$

158. (b) Eq. of A^{2+} Eq. of MnO_4^-

$$2.7 \cdot 10^3 \cdot 3 = 1.6 \cdot 10^3 \cdot n_f$$

$$n_f = 5 \quad \text{Acidic medium.}$$

159. (c) Moles of O_2 gas obtained $\frac{2.24}{22.4} = 0.1$ mole

$$\text{H}_2\text{O}_2 \quad \text{H}_2\text{O} \quad \frac{1}{2} \text{O}_2 \quad \text{Moles of H}_2\text{O}_2 \quad 0.2$$

$$1 \text{ mole} \quad \frac{1}{2} \text{ mole} \quad M \quad V(\text{L}) \quad 0.2$$

$$2 \text{ mole} \quad 1 \text{ mole} \quad \frac{5.6}{11.2} \quad V(\text{L}) \quad 0.2$$

$$2 \text{ mole} \quad 1 \text{ mole} \quad V \quad 0.4 \text{ L} \quad 400 \text{ mL}$$

$$0.2 \text{ mole} \quad 0.1 \text{ mole}$$

160. (d) Eq. of acid Eq. of base

$$1000 \cdot \frac{1.25}{M} = 2 \cdot 25 \cdot 0.25 = 2$$

$$M = 200$$

$$\text{161. (a) } N_{(\text{H}^+)} = \frac{\left| \begin{array}{ccc} 10 & 1 & 20 \\ 1 & 2 & 30 \\ 2 & 3 & 1 \end{array} \right|}{1000} = \frac{3N}{100}$$

162. (b) Eq. of acid Eq. of base

$$\frac{0.45}{90} \cdot n_f = \frac{20 \cdot 0.54}{1000} \quad n_f = 2$$

163. (c) Eq. mass of $\text{H}_2\text{O}_2 = \frac{34}{2} = 17$

$$\text{Eq. of H}_2\text{O}_2 = \frac{3.4}{100} \cdot \frac{x}{17}$$

$$= \text{Eq. of KMnO}_4 \quad x \cdot N \cdot 10^3 = N \cdot 2$$


$$\text{Molarity of KMnO}_4 = \frac{2}{5} = 0.4$$

164. (c) 50 m-eq. $\text{KMnO}_4 = 10$ m-mole KMnO_4

$$\text{or } 25 \text{ m-mole of O}_2 = 25 \cdot \frac{22.4}{10^3}$$

$$0.56 \text{ L}$$

165. (c) Zn will reduce Fe^{3+} to Fe^{2+}



n_f 2 POAC for Fe
 Fe^{2+} O. A. Fe^{3+}

$$2 \frac{0.8}{160} 1 n_{\text{Fe}^{2+}} n_{\text{Fe}^{2+}} 0.01$$

eq. of Fe^{2+} Eq. of O.A.

$$\text{Moles of Fe}^{2+} 1 16.7 0.1 n_f$$

$$0.01 1 \frac{16.7 \cdot 0.1}{1000} n_f$$

$$n_f 6$$

166. (c) Eq. of KMnO_4 20 0.1 5 10

Eq. of oxalic acid 25 0.2 2 10

167. (d) Fe^{2+} MnO_4^- Fe^{3+} Mn^{2+}

$$n_f 1 n_f 5$$

Moles of Fe^{2+} 1 by KMnO_4

moles of MnO_4^- 5 ... (1)



$$n_f 1 n_f 6$$

Moles of Fe^{2+} 1 by $\text{K}_2\text{Cr}_2\text{O}_7$

moles of MnO_4^- 6 ... (2)

(1) (2)

$$\frac{n_1}{n_2} = \frac{5}{6}$$

168. (b) Moles of AgCl moles of AgNO_3

$$= \text{moles of HCl} \frac{0.1435}{143.5} 10^{-3}$$

mass of HCl 0.0365

mass of H_2SO_4 = 0.0635

$$\text{mass \% of H}_2\text{SO}_4 = \frac{0.0635}{0.1} \frac{100}{63.5}$$

169. (b) Eq. mass of KBrO_3 = $\frac{1}{6}$ of its mol. mass

$$\frac{1}{6} 167$$


$$N_{\text{Na}_2\text{S}_2\text{O}_3} \frac{0.167}{167} 6 \frac{1}{0.05} 0.12 \text{ N}$$

170. (a) Total m-eq. of NaOH taken 20

m-eq. of H_2SO_4 = m-eq. of NaOH reacted

$$\frac{5 \cdot 0.2}{25} 250 10$$

m-eq. of NaOH reacted 20 10 10



m-moles of $(\text{NH}_4)_2\text{SO}_4$ reacted = 5

$$\text{mass of } (\text{NH}_4)_2\text{SO}_4 \quad 5 \quad 10^3 \quad 132 \quad 0.66$$

Percentage of $(\text{NH}_4)_2\text{SO}_4$ in sample

$$= \frac{0.66}{0.80} 100 = 82.5$$

171. (c) milli-equivalent of NH_3 reacted with HNO_3

$$45 \cdot 0.4 \cdot 20 \cdot 0.1 = 16$$

$$\frac{W}{17} 1000 16; W_{\text{NH}_3} 0.272 \text{ g;}$$

$$\text{mass of N} 0.272 \frac{14}{17} 0.224$$

$$\% \text{ N in the sample} \frac{0.224}{1.12} 100 20\%$$

$$172. (a) 1000 \frac{W}{24 \cdot 64} 2 \cdot 90 \frac{1}{100} 5;$$

$$W 0.198 \text{ g}$$

% of oxalate ion in a given sample

$$\frac{0.198}{0.3} 100 66$$

173. (a) m-moles of HCl 20 0.1 = 2

m-moles of MgO reacted with HCl $\frac{2}{2}$ 1

mass of MgO present 1 40 mg

$$\% \text{ of MgO} \frac{40}{320} 100 12.5\%$$

174. (c) $\text{Pb}^{2+} + \text{SO}_4^{2-}$ (from Na_2SO_4) PbSO_4

moles of Na_2SO_4 required = moles of Pb^{2+} ions

$$\frac{0.355}{142} 0.0025$$

$$[\text{Pb}^{2+}] \frac{0.0025}{0.50} 5 \cdot 10^{-3} \text{ M}$$

175. (b) Let volume of HNO_3 is V then

$$V 1.05 \frac{12.6}{100} \frac{3}{63} = \frac{1}{278} 1$$

$$V 0.57 \text{ mL}$$

176. (c) m-eq. of KMnO_4 m-eq. of FeC_2O_4

+ m-eq. of FeSO_4

$$(0.1 \cdot 5) V \frac{100}{144} \frac{100}{152}$$

$$3 1$$

$$V 5.48 \text{ mL}$$

177. (a) meq. FeSO_4 $(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$


meq of KMnO_4

$$(n-1)$$

$$\frac{W}{392} \quad 1 \quad 1000 \quad 0.1 \quad 50; \quad W \quad 1.96 \text{ g}$$

Hence, % purity of Mohr's salt


$$\frac{1.96}{2.5} \quad 100 \quad 78.4\%$$



$$\begin{aligned} \text{Equivalents of Fe}^{2+} &= \text{moles of Mohr's salt} \\ &= \text{equivalents of K}_2\text{Cr}_2\text{O}_7 \\ &\quad 500 \quad 10^3 \quad 6 \quad 1 \quad 3.0 \end{aligned}$$

Hence, mole per cent of Mohr's salt

$$\frac{3}{4} \quad 100 \quad 75$$



$$\begin{array}{ccc} \text{Eq. mass of oxide} & E_{\text{Element}} & E_{\text{Oxygen}} \\ & 16 & 8 \quad 24 \end{array}$$

$$\frac{E_{\text{Oxide}}}{E_{\text{Metal}}} \quad \frac{24}{16} \quad \frac{3}{2}$$



i.e., the valency of metal in oxide is 3

$\therefore 6 \text{ mg } 6 \times 10^{-3} \text{ g of H}_2 \text{ reacts with X}_2\text{O}_3$

1 g or 1 equivalent of H₂ reacts with

$$\text{X}_2\text{O}_3 = 0.1596 \text{ g}$$

$$\text{Equivalents of X}_2\text{O}_3 = \frac{0.1596}{6 \times 10^{-3}} = 26.6 \text{ g;}$$

$$\text{Equivalent mass of X}_2\text{O}_3 = 26.6$$


= Equivalent mass of X

+ Equivalent mass of O $E \quad 8$

$$E \quad 8 \quad 26.6; \quad E \quad 18.6$$

$\therefore \text{At. mass} = \text{Eq. mass} \times \text{Valency}$

$$18.6 \quad 3 = 55.8$$




$$N \quad 10 \quad 50 \quad 1 \quad N \quad 5$$

m-eq. of MnO_4^- m-eq. of oxalic acid

$$N_1 V_1 \quad \frac{W}{E} \quad 1000$$

$$5 \quad 100 \quad \frac{W}{(90/2)} \quad 1000 \quad W \quad 22.5 \text{ g}$$



In redox reaction :

when KMnO_4 is used : $0.2 \quad V \quad 2a \quad 4b$.


..(1)

In acid-base neutralization :

when NaOH is used: $0.12 \quad V \quad a \quad 3b \dots(2)$

from (1) and (2) $a \quad 0.06 \quad V$


$$b \quad 0.02 \quad V \text{ or } a:b \quad 3:1$$



Eq. of SnSO_4 eq. of FAS

Moles $2 \quad \text{Moles} \quad 1$

Moles of SnSO_4 required for 1 mole FAS $\frac{1}{2}$



Eq. of KMnO_4 Eq. of FAS

Moles $5 \quad \text{Moles} \quad 1$

Moles of KMnO_4 req. $\frac{1}{5}$

$$\text{Ratio} \quad \frac{1/2}{1/5} \quad \frac{5}{2} \quad 2.5$$



eq. of $\text{NaHC}_2\text{O}_4 = \text{eq. of NaOH}$

$$100 \times 10^{-3} \quad 0.2$$

$$0.02$$

$$a \quad 0.02 \quad 112 = 2.24 \text{ g}$$

eq. mass of NaHC_2O_4 as reductant

$$(\text{C}_2\text{O}_4^{2-} \quad 2\text{CO}_2 + 2e^-) \quad \frac{112}{2} \quad 56$$

eq. of NaHC_2O_4 as reductant = eq. of $\text{KMnO}_4 = 100 \times 10^{-2} \times 0.02 \times 5 = 0.01$

$$b \quad 0.01 \quad 56 = 0.56 \text{ g}$$

$$\text{Hence, } a:b \quad \frac{2.24}{0.56} \quad 4:1 \text{ or } a \quad 4b$$




$$= \text{eq. of KMnO}_4 \quad V \quad 0.1 \quad 5$$

$$2 \times 2 \times 0.5V; \quad V = 8\text{L}$$

eq. of $\text{H}_2\text{C}_2\text{O}_4 = \text{eq. of NaOH}$

$$1 \times 2 \times 0.2 \quad V_2; \quad V_2 \quad 10 \text{ L}$$

$$\text{Hence, } V_1:V_2 \quad 8:10 \quad 4:5$$



$$\text{eq. mass of KMnO}_4 = \frac{\text{molar mass}}{5}$$

eq. of $\text{Na}_2\text{S}_2\text{O}_3 = \text{eq. of I}_2$ liberated

= eq. of $\text{KMnO}_4 + \text{eq. of K}_2\text{Cr}_2\text{O}_7$

$$\text{or } N \quad 1 \quad 0.02 \quad 5 \quad 0.05 \quad 6$$

$$N \quad 0.4 \quad \text{or } M \quad 0.4$$

187. (b) m-eq. of acids in 50 its mL m-eq. of Na_2CO_3

$$25 \quad 2 \quad 50 \quad 4 \quad 2 \quad 2 \quad \times \quad \frac{1000}{50} \quad 25 \quad 1$$

$\times 62.5 \text{ mL}$

188. (d) Moles of pure $\text{CuSO}_4 = 0.12 =$ moles of $\text{Na}_2\text{S}_2\text{O}_3$ reacted with I_2

Moles of $\text{Na}_2\text{S}_2\text{O}_3 = 120 \times 10^{-3} \text{ M}$

$$\frac{0.12}{\text{M}} = 1.0$$

189. (d) Let $x \text{ g}$ of Li_2CO_3 and $(1 - x) \text{ g}$ of M_2CO_3 present in given mixture

2 Total moles of carbonates moles of HCl and

$$2 \quad \frac{x}{74} \quad \frac{1-x}{2M} \quad \frac{x}{60} \quad 21.6 \quad 0.5 \quad 10^{-3}$$

$$\therefore \quad \frac{x}{74} \quad \frac{1-x}{2M} \quad \frac{x}{60} \quad \frac{x}{0.20} \quad \frac{1-x}{118}$$

190. (b) m-eq. of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ in 1 litre

$$20 \quad 0.02 \quad 5 \quad \frac{1000}{25} \quad 80$$

$$\frac{W}{278} \quad 1 \quad 1000 \quad 80 \quad W \quad 22.24$$

mass % of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ in given sample

$$\frac{22.24}{32} \quad 100 \quad 69.5$$

191. (a) Since, phenolphthalein indicates conversion of Na_2CO_3 into NaHCO_3 only hence, $x \text{ mL}$, of HCl will be further required to convert NaHCO_3 to H_2CO_3 . So, total volume of HCl required to convert Na_2CO_3 into H_2CO_3 $x \quad x \quad 2x \text{ mL}$

192. (d) In presence of phenolphthalein ($\text{Na}_2\text{CO}_3 \quad \text{NaHCO}_3$)

50 % No reaction

$$\frac{1}{2} \text{ eq. of } \text{Na}_2\text{CO}_3 \quad \text{eq. of HCl}$$

$$\frac{1}{2} \quad \frac{x}{106} \quad 2 \quad \frac{10 \cdot 0.01}{1000}$$

$$x \quad 0.0106$$

$$\% \text{ of } \text{Na}_2\text{CO}_3 \quad \frac{0.0106}{0.1} \quad 100 \quad 10.6$$

193. (c) Phenolphthalein gives the end point corresponding to the reactions :



$$\text{m-moles of NaOH} + \text{m-moles of } \text{Na}_2\text{CO}_3 \quad \text{m-moles of HCl} \quad 2.5$$

Methyl orange gives the end point corresponding to the reactions :



$$\text{m-moles of NaOH}$$

$$+ \text{m moles of } \text{Na}_2\text{CO}_3 \quad 2$$

$$= \text{m-moles of HCl} \quad 30 \quad 0.1 \quad 3$$

$$\text{m-moles of } \text{Na}_2\text{CO}_3 = 0.5$$

$$\text{m-moles of NaOH} = 2.5 \quad 0.5 \quad 2$$

$$\text{Ratio of m-moles of NaOH and } \text{Na}_2\text{CO}_3$$

$$= \frac{2}{0.5} = 4 : 1$$

194. (a) $\text{NaOH} \quad \text{Na}_2\text{CO}_3$
 $x \text{ g} \quad y \text{ g}$

In presence of HPh

$$\text{Eq. of NaOH} \quad \frac{1}{2} \quad \text{eq. of } \text{Na}_2\text{CO}_3$$

$$\text{Eq. of HCl}$$

$$\frac{x}{40} \quad 1 \quad \frac{1}{2} \quad \frac{y}{106} \quad 2 \quad \frac{17.5}{1000} \quad \frac{1}{10} \quad \dots(1)$$

After this MeOH is added

$$\frac{1}{2} \quad \text{eq. of } \text{Na}_2\text{CO}_3 \quad \text{eq. of HCl}$$

$$\frac{1}{2} \quad \frac{y}{106} \quad 2 \quad 2.5 \quad \frac{1}{10} \quad \frac{1}{1000} \quad \dots(2)$$

Placing the value of Eqn. ... (1)

$$\frac{x}{40} \quad \frac{2.5}{10000} \quad \frac{17.5}{10000}$$

$$\frac{x}{40} \quad 15 \quad 10^4 \quad x \quad 15 \quad 40 \quad 10^4$$

$$x \quad 0.06 \text{ g}$$

195. (c) $\text{CaCO}_3 \quad \text{CaO} \quad \text{CO}_2$

$$\text{Mass of } \text{CaCO}_3 \quad 0.9 \text{ g}$$

$$\text{Moles} \quad 9 \times 10^{-3}$$



$$9 \times 10^{-3} \quad 50 \times 10^{-3} \text{ moles}$$

$$0$$

$$- \quad 32 \times 10^{-3} \quad 9 \times 10^{-3}$$

$$\text{moles} \quad \text{moles}$$



$$\text{Eq. of HCl} \quad \text{Eq. of NaOH} \quad \text{Eq. of } \text{Na}_2\text{CO}_3$$

$$0.5 \quad V \quad 1 \quad 32 \quad 10^3 \quad 1 \quad 9 \quad 10^3 \quad 1$$

$$V \quad 82 \text{ mL}$$



At the equivalent point m-mole of HCl

$$2 \text{ m-mole of Na}_2\text{CO}_3$$

$$60 \quad M \quad 2 \quad \frac{0.318}{106} \quad 1000$$

$$M \quad 0.1$$



$$162 \text{ g} \quad \text{—} \quad 56 \text{ g}$$

$$16.2 \text{ g} \quad \text{—} \quad 5.6 \text{ g}$$

$$162 \text{ g Ca}(\text{HCO}_3)_2 \quad 100 \text{ g CaCO}_3$$

$$16.2 \text{ g Ca}(\text{HCO}_3)_2 \quad 10 \text{ g CaCO}_3$$

$$\text{Hardness} \quad \frac{\text{Mass of CaCO}_3}{\text{Mass of water}} \quad 10^6$$

$$\frac{10}{10^4} \quad 10^6 \quad 1000 \text{ ppm}$$



+ milli-equivalents of Mg^{2+}


= milli-equivalents of Na_2CO_3

$$\frac{20}{20} \quad \frac{12}{12} \quad V \quad 2$$

$$V \quad 1 \text{ mL for 1 L of tap water}$$


For 5000 L of pond water

$$V \quad 5000 \text{ mL or 5 L}$$



Total mass of $\text{CaCO}_3 = 6 \text{ mg}$

$$\text{Hardness in ppm} \quad \frac{6 \quad 10^3}{10^3} \quad 10^6 \quad 6 \text{ ppm}$$



$$10^6 \text{ g of H}_2\text{O} \quad 200 \text{ g of CaCO}_3$$

$$10^3 \text{ g of H}_2\text{O} \quad \frac{200 \quad 10^3}{10^6} \text{ g of CaCO}_3$$

$$\text{Millimoles of CaCO}_3 \quad 2$$

$$\text{Milliequivalents of CaCO}_3 \quad 2 \quad 2 \quad 4$$

Level 2

1. (a) Let wt. of NH_4NO_3 and $(\text{NH}_4)_2\text{HPO}_4$ are x and y gram respectively

$$\frac{x}{80} \quad 2 \quad 14 \quad \frac{y}{132} \quad 2 \quad 14$$

$$x \quad y$$

$$x : y \quad 2 : 1$$

2. (a) Let V mL of alcohol be required

$\therefore$ mass of alcohol is same in both solutions

$$\frac{75}{100} \quad 0.8 \quad V \quad \frac{30}{100} \quad 0.9 \quad 150$$

$$V \quad 67.5 \text{ mL}$$

3. (d) The precipitation reaction is



$$\text{mole of Fe}_2\text{O}_3 \text{ in sample} \quad \frac{0.80 \quad 0.5}{160}$$

$$2.5 \quad 10^3$$

$$\text{mole of Fe}^{3+} \quad 2 \quad 2.5 \quad 10^3 \quad 5 \quad 10^3$$

$$M_{\text{NH}_3} \quad \frac{0.986 \text{ g/mL} \quad 1000 \text{ mL/litre} \quad 0.025}{17}$$

$$M_{\text{NH}_3} \quad 1.45$$

$$3 \text{ moles of Fe}^{3+} = \text{moles of NH}_3$$

$$1.45 \quad V \text{ (inL)}$$

$$V \quad \frac{3 \quad 5 \quad 10^3}{1.45} \quad 10.34 \quad 10^3 \text{ L or } 10.34 \text{ mL}$$

4. (a) Balanced reaction is



Number of moles of Fe_2O_3

$$= \frac{120 \quad 1000}{2 \quad 56 \quad 48} \quad \frac{90}{100}$$

Mass of 80% pure iron produced

$$\frac{120 \quad 1000 \quad 0.9}{2 \quad 56 + 48} \quad \frac{2 \quad 56}{0.8}$$

$$94500 \text{ gram or } 94.5 \text{ kg}$$

5. (d) Let two equimolar carbonates are ACO_3 and BCO_3

$$\text{mass of metal A} = 2.8 \quad 0.125 = 0.35 \text{ g}$$



Total two moles CO_3^{2-} provide 2 moles of CO_2 or 60 g CO_3^{2-} provide 44 g of CO_2

Total mass of CO_3^{2-} required to produce

$$1.32 \text{ g of CO}_2 \quad \frac{60}{44} \quad 1.32 \quad 1.8 \text{ g}$$


$$\text{mass of metal (A + B)} = 2.8 \quad 1.8 = 1 \text{ g}$$

$$\text{mass of B} = 1 \quad 0.35 = 0.65 \text{ g}$$

$$\% \text{ of B} \quad \frac{0.65}{2.80} \quad 100 \quad 23.21$$

6. (b) Loss in mass due to heating $6.2 \quad \frac{5}{100}$

$$= 0.31 \text{ g}$$



$$2 \quad 84 \text{ g} \quad \quad \quad 44 \quad 18 \text{ g loss}$$

$$1 \text{ mole NaHCO}_3 \quad \quad \quad 31 \text{ g loss due to heating}$$

$$0.31 \text{ g loss from } 0.01 \text{ mole of } \text{NaHCO}_3$$

$$\text{Moles of } \text{Na}_2\text{CO}_3 \text{ produced } \frac{0.01}{2} = 0.005$$

$$\text{Total moles of carbonate reacted with } \text{BaCl}_2$$

$$7.5 \times 10^{-2} \times \frac{100}{10} = 0.015$$

$$\text{Moles of carbonates in original sample}$$

$$0.015 - 0.005 = 0.01$$

$$\text{Mass of } \text{Na}_2\text{CO}_3 \text{ in original sample}$$

$$0.01 \times 106 = 1.06 \text{ gram}$$

7. (c) Moles of NO_2 required

$$\frac{1575}{63} \times \frac{3}{2} \times \frac{1}{0.8} = 46.875$$

$$\text{moles of NO required } \frac{46.875}{0.60}$$

$$\text{moles of } \text{NH}_3 \text{ required } \frac{46.875}{0.60} \times \frac{1}{0.50}$$

$$156.25$$

$$\text{Volume of } \text{NH}_3 \text{ at STP required}$$

$$156.25 \times 22.4 = 3500 \text{ L}$$

8. (c) Moles of NaOH consumed when it reacts with $\text{H}^+ = 3 \times 1$

$$\text{moles of } \text{H}_2\text{SO}_4 \text{ present in sample} = 1.5$$

$$\text{mass of } \text{H}_2\text{SO}_4 \text{ in sample} = 1.5 \times 98 = 147 \text{ g}$$

$$\% \text{ purity } \frac{147}{183.75} \times 100 = 80$$

9. (c) Let mass of sample is x g; mass of MnO_2 is $0.75x$ g

$$\text{moles of } \text{MnO}_2 = \frac{0.75x}{87}$$

$$\text{mass of Mn } \frac{0.75x}{87} \times 55 = 0.474x$$



$$\text{mass of } \text{Mn}_3\text{O}_4 \text{ produced}$$

$$\frac{0.75x}{87} \times \frac{1}{3} \times 229 = 0.658x \text{ g}$$

$$\text{Total mass of residue } 0.658x + 0.2x$$

$$(\text{mass of impurities})$$

$$\% \text{ of Mn in residue } \frac{0.474x}{0.858x} \times 100$$

$$55.24$$

10. (b) Moles of AgCl = moles of chloride

$$\frac{1.96}{143.5} = 0.0136$$

$$\text{Moles of organic compound } \frac{1}{147}$$

$$6.8 \times 10^{-3}$$

$$\text{Chlorine atoms in each molecule of organic compound } \frac{0.0136}{6.8 \times 10^{-3}} = 2$$

11. (c) Let x g is mass of CaC_2O_4 and $(0.6 - x)$ g mass of MgC_2O_4



$$\text{mass of } \text{CaCO}_3 \text{ produced } \frac{x}{128} \times 100$$

$$\text{mass of } \text{MgCO}_3 \text{ produced } \frac{(0.6 - x)}{112} \times 84$$

$$\frac{x}{128} \times 100 + \frac{(0.6 - x)}{112} \times 84 = 0.465$$

$$x = 0.48 \text{ g}$$

Due to further heating



$$- \quad \frac{x}{128}$$



$$- \quad \frac{0.6 - x}{112}$$

$$\text{mass of CaO and MgO produced}$$

$$\frac{0.48}{128} \times 56 + \frac{0.12}{112} \times 40 = 0.252 \text{ g}$$

12. (a) Moles of $\text{BaSO}_4 = \frac{1.22}{233.3}$

$$\text{moles of } \text{M}_2(\text{SO}_4)_3 = \frac{1.22}{233.3} \times \frac{1}{3}$$

$$1.743 \times 10^{-3}$$

$$\text{wt. of } \text{M}_2(\text{SO}_4)_3 = 0.596$$


$$1.743 \times 10^{-3} \times (2M + 96 \times 3) = 0.596$$

$$M = 26.9$$

13. (b) $2\text{NH}_3 + \text{CO}_2 \rightarrow \text{NH}_2\text{—CO—NH}_2 + \text{H}_2\text{O}$

$$\text{moles of urea } \frac{999.6 \times 10^3}{60} = 16.66 \times 10^3$$


$$\text{moles of } \text{CO}_2 = 16.66 \times 10^3$$



$$\frac{16.66 \times 10^3}{n} = [12n + (2n - 2)] \times 236 \times 10^3$$

$$n = 12$$

molecular formula of hydrocarbon



Let initial pressure of C_nH_{2n-2} is P then increase in pressure

$$P(2n-1) - 1 = \frac{3n-1}{2}$$

$$\frac{n-1}{2} = P$$


546 K and 4.6 atm 273 K and 2.3 atm;
Increase in pressure 0.3 atm

$$P = \frac{nRT}{V} = \frac{11.6}{M} = \frac{0.0821 \times 273}{22.41}$$

$$\frac{n-1}{2} = \frac{11.6}{(14n-2)} \quad 0.3$$

$$\frac{n-1}{14n-2} = \frac{0.6}{11.6} \quad n = 4$$

Compound is C_4H_{10}



moles of H_2O_2 used in first reaction

$$\frac{0.2}{2} = \frac{1}{0.4} \quad 0.25$$

$$\text{moles of } O_2 \text{ produce} = \frac{6.74}{22.4} = 0.3$$


moles of H_2O_2 used in second reaction

$$\frac{0.3}{3} = \frac{0.2}{0.5}$$

Total mole of H_2O_2 consumed 0.45

$$\text{Molarity of } H_2O_2 = \frac{0.45}{0.15} = 3 M$$

Volume strength 11.2 3 = 33.6 volumes




m-moles of H_2SO_4 produced 5; m-moles of HCl produced 10

m-moles of $Ba(OH)_2$ required

$$5 \text{ (for } H_2SO_4) + \frac{10}{2} \text{ (for } HCl) = 10$$

$$M V = 10 \times 0.2 = V \times 10, V = 50 \text{ mL}$$



m-moles of $Na_2CO_3 =$ m-moles of H_2SO_4

$$20 \quad 0.1 \quad 2$$

m-moles of Na_2CO_3 in 250 mL solution

$$\frac{250}{25} \quad 2 \quad 20$$

$$\text{wt. of } Na_2CO_3 = \frac{20}{25} \times 106 \times 10^{-3} = 2.12 \text{ g}$$

$$\% \text{ of } Na_2CO_3 = \frac{2.12}{5} \times 100 = 42.4$$

$$\% \text{ of } Na_2SO_4 = 100 - 42.4 = 57.6$$

18. (a) M_{NaOH} resultant


$$\frac{20 \times 0.2 \times 35 \times 0.1}{100} = 0.075 M$$

Milli-equivalent of $NaOH =$ milli-equivalent of $H_2C_2O_4$

Let mass of impure sample is x gram

$$40 \times 0.075 = \frac{x \times 0.90}{90} \times 2 \times 1000$$

$$x = 0.15 \text{ gram}$$




POAC on Ag

Mole of Ag in coin = mole of Ag in $AgCl$

$$a = \frac{14.35}{143.5} = 0.1 \text{ mole}$$

$$\text{Mass of } Ag \text{ in coin} = 0.1 \times 108 = 10.8 \text{ g}$$


$$\% \text{ silver in coin} = \frac{10.8}{11.34} \times 100 = 95.2\%$$



16 parts of X combine with 42 parts of Y


For Y 3 parts in compound B , then in compound C 5 parts

42 parts in compound B , then in the compound C $\frac{5}{3} \times 42 = 70$ parts



$$\text{Mole of Ozone formed} = \frac{2}{3} \times \frac{15}{100} \times 3 = 0.3 \text{ mole}$$

$$\text{Mass of Ozone formed} = 0.3 \times 48 \text{ g} = 14.4 \text{ g}$$



$$\text{So } [Ca^{2+}] = \frac{1}{2} \times 10^{-3} M$$

$$\text{Mass of } Ca^{2+} = \frac{10^{-3}}{2} \times 40$$

So, mass of Ca^{2+} ions in 10^6 mL hard water

$$\frac{\frac{1}{2} \times 10^3 \times 40}{10^3} \times 20$$

23. (a) m-eq. of acid = m-eq. of base

$$\frac{N_1 V_1}{n} = \frac{N_2 V_2}{n}$$

$$\frac{29.4}{98} \times 100 = \frac{90}{40} \times \frac{1000}{500}$$

$$n = 3$$

24. (b) When phenolphthalein is used as an indicator : $0.05 \times 20 = 0.1 \times x$; $x = 40$ mL

When methyl orange is used as an indicator:

$$0.05 \times y = 20 \times 0.1 \times 3$$

$$y = 120 \text{ mL}$$

$$y = x = 80 \text{ mL}$$

25. (a) $\text{I}_2 + \text{AsO}_2 + 2\text{H}_2\text{O} \rightarrow \text{HAsO}_2 + 2\text{I}^- + 3\text{H}^+$

$$\text{m-eq. of HAsO}_2 \text{ (in 50 mL)} = \text{m-eq. of I}_2$$

$$= 35 \times 0.05 \times 2 = 3.5$$

$$\text{m-eq. of HAsO}_2 \text{ in 250 mL} = 3.5 \times \frac{250}{50}$$

$$= 17.5$$

mass of HAsO_2 in sample

$$\frac{17.5}{2} \times (108) \times 10^{-3} = 0.945 \text{ g}$$

$$\% \text{ of HAsO}_2 \text{ in the sample} = \frac{0.945}{3.78} \times 100$$

$$= 25\%$$

26. (a) m-eq. of FeO = m-eq. of KMnO_4

$$0.25 \times 5 \times 100 = \frac{0.25 \times 100 \times 5}{1}$$

$$125$$

total m-eq. or m-mole. of Fe^{2+}

$$1000 \times 0.1 \times 6 = 600$$

(from FeO and Fe_2O_3 after reaction with Zn dust)

$$\text{m-mole of Fe}^{2+} \text{ from Fe}_2\text{O}_3 = 600 - 125$$

$$= 475$$

$$\text{m-mole of Fe}_2\text{O}_3 = \frac{475}{2}$$

$$\text{Mass of FeO} = \frac{125 \times (56 \times 16)}{1000} \text{ g} = 9 \text{ g}$$

$$\text{Mass of Fe}_2\text{O}_3 = \frac{475}{2} \times \frac{160}{1000} = 38 \text{ g}$$

$$\% \text{ Fe}_2\text{O}_3 = \frac{38}{38 + 9} \times 100 = 80.85\%$$

27. (d) $3\text{Br}_2 + 6\text{OH}^- \rightarrow 5\text{Br}^- + \text{BrO}_3^- + 3\text{H}_2\text{O}$... (1)



10 m-mole of Br_2 produce $(10/3)$ m-mole BrO_3^- and m-moles of

$$\text{CaC}_2\text{O}_4 \text{ required} = \frac{10}{3} \times 3 = 10$$

$$\text{Mass of CaC}_2\text{O}_4 = 10 \times 10^{-3} \times 128$$

$$\% \text{ purity} = \frac{10 \times 10^{-3} \times 128}{2} \times 100 = 64$$

28. (b) m-moles of KI = 50 $\times$ 0.04 = 2

m-equivalent of $\text{Na}_2\text{S}_2\text{O}_3$ reacted with 10 mL solution = $20 \times 2.5 \times 10^{-3} = 50 \times 10^{-3}$

m-equivalent of I left unreacted in 100 mL solution = $50 \times 10^{-3} - \frac{100}{10} \times 0.5$

m-moles of I which reacts with CuCO_3 = $2 - 0.5 = 1.5$



$$\text{m-moles of Cu}^{2+} = \frac{2}{5} \times 1.5 = 0.6$$


$$\text{weight of CuCO}_3 = 0.6 \times 10^{-3} \times 123.5$$

$$= 0.0741$$

$$\text{weight \% of CuCO}_3 = \frac{0.0741}{0.1} \times 100$$

$$= 74.1$$

29. (b) $\text{Fe}_2(\text{C}_2\text{O}_4)_3 + \text{MnO}_4^- + \text{H}^+ \rightarrow 2\text{Fe}^{3+} + 6\text{CO}_2 + \text{H}_2\text{O}$

$$(n = 6) \quad (n = 5)$$


Total equivalents of $(\text{Fe}_2(\text{C}_2\text{O}_4)_3 + \text{FeC}_2\text{O}_4)$

equivalents of KMnO_4

$$0.5 \times 6 + 0.5 \times 3 = x \times 5$$

$$\text{moles of KMnO}_4 = x \times 0.9$$

30. (b) m-eq. of $\text{Na}_2\text{C}_2\text{O}_4$ = m-eq. of KMnO_4 reacted

total m-eq. of KMnO_4 - excess m-eq. of KMnO_4 reacted with $\text{H}_2\text{C}_2\text{O}_4$

$$45 - 0.02 \times 5 \times 10 \times 0.1 \times 2 = 2.5$$


$$1000 \times \frac{W}{134} \times 2 = 2.5$$

$$W_{\text{Na}_2\text{C}_2\text{O}_4} = 0.1675 \text{ g}$$

% purity of $\text{Na}_2\text{C}_2\text{O}_4$ in sample

$$\frac{0.1675}{0.2} \times 100 = 83.75$$

31. (a) Reactions of $\text{K}_2\text{Cr}_2\text{O}_7$ and KMnO_4 with KI may be given as :



Thus, equivalent mass of $\text{K}_2\text{Cr}_2\text{O}_7$

$$= \frac{294}{6} = 49$$

equivalent mass of $\text{KMnO}_4 = \frac{158}{5} = 31.6$

$\therefore$ m-eq. of $\text{K}_2\text{Cr}_2\text{O}_7$ + m-eq. of KMnO_4
= m-eq. of I_2 = m-eq. of hypo.

Let the mass of $\text{K}_2\text{Cr}_2\text{O}_7 = x$ g;

mass of $\text{KMnO}_4 = (0.5 - x)$ g

$$\frac{x}{49} \times \frac{(0.5 - x)}{31.6} \times 150 \times 10^{-3}$$

or $x = 0.0732$

$$\% \text{ of } \text{K}_2\text{Cr}_2\text{O}_7 = \frac{0.0732}{0.5} \times 100$$

14.64

32. (b) $\text{I}_2 + 2\text{Na}_2\text{S}_2\text{O}_3 \rightarrow 2\text{NaI} + \text{Na}_2\text{S}_4\text{O}_6 \dots$ (i)

m-moles of $\text{Na}_2\text{S}_2\text{O}_3$ consumed

$$15 \times 0.4 = 6 \text{ m-mole}$$

m-moles of I_2 consumed = 3 m-mole



$\dots$ (ii)

m-mol of I_2 reacted with NaOH are

$$\frac{30 \times 2 \times 3}{2} = 12 \text{ m-mole}$$

Total m-mole of I_2 consumed in reaction

(i) and (ii) = 3 + 12 = 15 m-mole

$$\text{Molarity of } \text{I}_2 = \frac{15}{150} = 0.1 \text{ M}$$

33. (a) Let x and y are milli-moles H_2SO_4 and $\text{H}_2\text{C}_2\text{O}_4$ present in the given mixture reacts with base

m-eq. of acids = m-eq. of base;

$$(x + y) \times 2 \times 10^{-3} = 0.1 \times \frac{1000}{10} \text{ or } x + y = 15$$

In second experiment only KMnO_4 reacted with $\text{H}_2\text{C}_2\text{O}_4$ in which Mn^{7+} converted into Mn^{2+} and $\text{C}_2\text{O}_4^{2-}$ converted into CO_2

milli-equivalent of $\text{H}_2\text{C}_2\text{O}_4$

= milli-equivalent of KMnO_4

$$y \times 2 \times 10^{-3} = 0.02 \times 5 \times \frac{1000}{100}$$

$$y = 2$$

$$x = 13, \text{ mass of } \text{H}_2\text{SO}_4 = 13 \times 10^{-3} \times 98$$

$$= 1.274 \text{ g}$$

$$\text{wt. \% of } \text{H}_2\text{SO}_4 \text{ in sample} = \frac{1.274}{3.185} \times 100$$

$$= 40\%$$

35. (c) $\text{KMnO}_4 + \text{C}_2\text{O}_4^{2-} + \text{H}^+ \rightarrow \text{Mn}^{2+} + \text{CO}_2$

$$n_f = 5 \quad n_f = 2$$

$$16 \text{ mL} \quad 40 \text{ mL}$$

$$0.05 \text{ M} \quad X \text{ M}$$

$$\text{m. eq. of } \text{C}_2\text{O}_4^{2-} = \text{m. eq. of } \text{KMnO}_4$$

$$2(40)X = 5 \times 16 \times 0.05$$

$$X = 0.05 \text{ M}$$

$$[\text{H}^+] = 2X = 0.1 \text{ M}$$

$$\text{pH} = 1$$

Level 3

Passage-1

1. (b) $\text{H}_2\text{O} + \text{SO}_3 \rightarrow \text{H}_2\text{SO}_4$;
18 g water combines with 80 g SO_3
4.5 g of H_2O combines with 20 g of SO_3 .
100 g of oleum contains 20 g of SO_3
or 20% free SO_3 .

2. (c) Initial moles of free SO_3 present in oleum

$$\frac{12}{18} \times \frac{2}{3} \text{ moles}$$

moles of water that can combine with SO_3

moles of free SO_3 combined with water

$$\frac{9}{18} \times \frac{1}{2} \text{ mole}$$

$$\text{moles of free } \text{SO}_3 \text{ left} = \frac{2}{3} - \frac{1}{2} = \frac{1}{6} \text{ mole}$$

$$\begin{aligned} \text{volume of free } \text{SO}_3 \text{ at STP} &= \frac{1}{6} \times 22.4 \\ &= 3.73 \text{ L} \end{aligned}$$

3. (c) $\text{Na}_2\text{CO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + \text{H}_2\text{O} + \text{CO}_2$

moles of CO_2 formed = moles of Na_2CO_3

$$\text{reacted} = \frac{5.3}{106} = 0.05$$

volume of CO_2 formed at 1 atm pressure and 300 K 0.05 24.63 1.23 L

4. (b) eq. of H_2SO_4 + eq. of SO_3 eq. of NaOH

$$\frac{x}{98} \cdot 2 + \frac{(1-x)}{80} \cdot 2 = 54 \cdot 0.4 \cdot 10^{-3}$$

$$\% \text{ of free } \text{SO}_3 = \frac{x \cdot 0.74}{1 \cdot 0.74} \cdot 100 = 26\%$$

Passage-2

1. (a) 1 litre of $\text{H}_2\text{O}_2(\text{aq})$ provide 11.2 litre of O_2 at 1 atm and 273 K

$$\text{moles of } \text{O}_2 = \frac{11.2}{22.4} = 0.5$$

$$n_{\text{H}_2\text{O}_2} \text{ required } = 0.5 \cdot 2$$

$$M_{\text{H}_2\text{O}_2} = \frac{n_{\text{H}_2\text{O}_2}}{V_{\text{solution}}} = 1 \text{ M}$$

2. (b) Strength in percentage means how many H_2O_2 present per 100 mL

$$\therefore M = 1 \text{ and mol. mass of } \text{H}_2\text{O}_2 = 34$$

34 g H_2O_2 present per litre of solution or
3.4 g H_2O_2 present per 100 mL of solution.

3. (b) m-eq. of H_2O_2 m-eq. of KMnO_4

$$20 \text{ N } 0.05 \cdot 5 \cdot 80 \text{ N } 1$$

$$N = \frac{\text{volume strength of } \text{H}_2\text{O}_2}{5.6}$$

$$\text{volume strength of } \text{H}_2\text{O}_2 = 5.6$$

4. (b) m-eq. of $\text{Ba}(\text{MnO}_4)_2$ m-eq. of H_2O_2

$$\therefore M = \frac{33.6}{11.2} = 3$$

$$\frac{w}{375} \cdot 10 \cdot 1000 = 3 \cdot 125 \cdot 2; w = 28.125$$

$$\% \text{ purity} = \frac{w}{40} \cdot 100$$

$$\frac{28.125}{40} \cdot 100 = 70.31$$

Passage-3

1. (c) 136 mg of CaSO_4 100 mg of CaCO_3 ;
190 mg of MgCl_2 2 100 mg of CaCO_3
1 litre or 1000 g water contain 300 mg of CaCO_3

10^6 g water contain

$$\frac{300 \cdot 10^3}{1000} \cdot 10^6 = 300 \text{ ppm}$$

2. (b) 10^6 g water contain 100 g CaCO_3

10 L or 10^4 g water contain

$$\frac{100}{10^6} \cdot 10^4 = 1 \text{ g of } \text{CaCO}_3$$

$$100 \text{ g } \text{CaCO}_3 \quad 162 \text{ g of } \text{Ca}(\text{HCO}_3)_2$$

$$\text{or } 1 \text{ g } \text{CaCO}_3 \quad 1.62 \text{ g of } \text{Ca}(\text{HCO}_3)_2$$

or 0.01 mole



moles of $\text{Ca}(\text{OH})_2$ required = 0.01

or mass of $\text{Ca}(\text{OH})_2$ required = 0.74 g

3. (b) m-moles of H present in outlet water

$$= \text{m-moles of OH} = 50 \cdot 0.1 = 5$$

$$\text{m-moles of } \text{Ca}^{2+} \text{ present in hard water} = \frac{5}{2}$$

= 2.5; (1 Ca^{2+} replaced by 2H)

$$\text{no. of mg of } \text{Ca}^{2+} \text{ ion} = 2.5 \cdot 40 = 100$$

200 g sample hard water contain 100 mg of Ca^{2+}

10^6 g sample hard water contain

$$\frac{100}{200} \cdot 10^6 = 500 \text{ ppm}$$

Passage-4

1. (c) n -factor $5 \cdot 2 = 10$

2. (a) H_3PO_2 is a monobasic acid n -factor 1

3. (a) n -factor $3 \cdot \frac{2}{0.95} \cdot 0.95 = 0.85$

$$E = \frac{M}{0.85}$$

4. (b) n -factor of VO 3 ; Fe_2O_3 $1 \cdot 2 = 2$;

x and y are 2 and 3

Passage-5

1. (a) Cl_2 2NaOH NaCl NaClO H_2O

$$4 \text{ mole} \quad \quad \quad 4 \text{ mole}$$

$$3\text{NaClO} \quad 2\text{NaCl} \quad \text{NaClO}_3$$

$$\frac{3}{1} \cdot \frac{4}{3} \quad 4 \text{ mole} \quad \frac{4}{3} \text{ mole}$$

$$4\text{NaClO}_3 \quad 3\text{NaClO}_4 \quad \text{NaCl}$$

$$\frac{4}{3} \text{ mole} \quad 1 \text{ mole}$$

$$\text{Moles of NaClO}_4 \quad \frac{122.5}{122.5} \quad 1 \text{ mole}$$

4 mol Cl_2 284 g Cl_2 will be required.



1 mole 1 mole 1 mole



1 mole $\frac{2}{3}$ 0.67 mole



$\frac{1}{3}$ mole $\frac{1}{4}$ $\frac{1}{3}$ $\frac{1}{12}$ mole 0.083 mole

Total NaCl 1 0.67 0.083 1.75 mole

3. (b) As NaClO_3 is intermediate it will be consumed in the complete reaction.

One or More Answers is/are Correct

1. (c, d)

1g-atom of nitrogen 1 mol N atoms

$\frac{1}{2}$ mol N_2 gas

$\frac{1}{2}$ mol N_2 gas 11.2 L N_2 at STP 14 g N_2

2. (a, b)

1g molecule V_2O_5 1 mole V_2O_5 2 mole V atom 5 mole O atom

5. (b, c, d)



As per equivalent concept

Number of equivalent of A No. of equivalent of B No. of equivalent of A_xB_y .

6. (b, c, d)



Equivalent of Ba(OH)_2 1 2 2

Acids having 2 equivalents will be exactly neutralised by Ba(OH)_2 .

7. (b, c)

Species having same empirical formula (EF) will have same percentage of C.

EF



8. (b, d)

$\therefore$ both are liquid CH_3OH is solute (less amount)

Mass of CH_3OH 30 0.8 24 g,

Mass of $\text{C}_2\text{H}_5\text{OH}$ 60 0.92 55.2 g

Mass of solution 24 55.2 79.2 g

Volume of solution $\frac{79.2}{0.88}$ 90 mL

Molarity $\frac{n_{\text{CH}_3\text{OH}}}{V(\text{L})}$ $\frac{24/32}{90}$ 1000

8.33 mol L^{-1}

Molality $\frac{n_{\text{Solute}}}{w_{\text{Solvent}}(\text{kg})}$ $\frac{24/32}{55.2}$ 1000

13.59

24

Mole fraction of solute $\frac{\frac{24}{32}}{\frac{24}{32} + \frac{55.2}{46}}$ 0.385

Mole fraction of solvent 1 0.385 0.615

9. (a, b, c)

17 g/L H_2O_2 $\frac{17}{34}$ mol L^{-1} H_2O_2

Molarity of H_2O_2 $\frac{1}{2}$

M $\frac{V}{11.2}$ V $\frac{1}{2}$ 11.2 5.6

5.6 volume H_2O_2 means 1 mL will give 5.6 mL at 273 K and 1 atm

$P_1 V_1$ $P_2 V_2$

1 5.6 2 V_2

V_2 2.8 mL

10. (a, b)

(a) 46 g of 70% $\frac{W}{V}$ HCOOH (d_{Solution} 1.4 g/mL)

70% $\frac{W}{V}$ HCOOH 70 g HCOOH in

100 mL solution.

Mass of solution 1.4 100 140 g


So, in 140 g solution, mass of HCOOH 70 g

in 46 g, mass of HCOOH $\frac{70}{140}$ 46 23 g

(b) 10 M HCOOH 10 mole HCOOH in 1000 mL solution mass of solution 1000 g

Mass of HCOOH 10 46 460 g

So in 50 g solution mass of



- (c) $25\% \frac{W}{W} \text{HCOOH}$ 25 g HCOOH in 100 g solution.




So in 50 g solution, mass of HCOOH 12.5 g

$$11. \text{ (a,c,d) } V_{\text{strength}} 56; \\ M \frac{28}{11.2} 2.5$$

1 L contain 2.5 moles of H_2O_2

or 2.5 34 = 85 g H_2O_2

Mass of 1 litre solution 265 g
($\therefore d$ 265 g/L)



12. (a, b)



(A) O_2 is L.R.

So, Volume of CO_2 produced 20 mL

Total volume of CO_2 20 60 80 mL

(B) CO is L.R.

Volume of CO_2 produced 30 mL


Total volume of CO_2 30 50 80 mL

(C) O_2 is L.R.

Volume of CO_2 produced 40 mL

Total volume of CO_2 30 40 70 mL

13. (a, c)



(A) Equivalent weight of base $\frac{\text{Mol. wt}}{2}$


(B) Equivalent weight of H_3PO_4 $\frac{98}{2}$

(C) XHPO_4 NaOH
 $n\text{-factor} = 1 \quad n\text{-factor} = 1$

Moles of NaOH 1 moles of XHPO_4 1

Moles of NaOH 1 1 1

(D)



Moles of X(OH)_2 required $\frac{1}{2}$


14. (b, c, d)



Upon balancing the redox-reaction



15. (a, b)



16. (a, c, d)

Equivalent weight $\frac{\text{Molecular weight}}{n\text{-factor}}$

(A) $n\text{-factor}$ of H_2PO_3 as acid 1
as base 1

In both case, eq. wt. of H_2PO_3 $\frac{81}{1}$ 81

(B) $n\text{-factor}$ of H_2PO_4 as acid 1 or 2
as base 1

(C) In acidic medium

Eq. wt. of KMnO_4 $\frac{\text{Molecular weight}}{5}$

In basic medium

Eq. wt. of KMnO_4 $\frac{\text{Molecular weight}}{1}$

In neutral/slightly basic medium

Eq. wt. of KMnO_4 $\frac{\text{Molecular weight}}{3}$

(D) In MgH_2 , O.S. of H 1

In H_2O_2 , O.S. of H 1

Match the Column

1. (A) 0.5 mole of $\text{SO}_2(\text{g})$ 32 g
weight $1.5 N_A$ atoms
11.2 L at 1 atm and 273 K

(B) 1 g of $\text{H}_2(\text{g})$ 0.5 mole
 H_2 11.2 L at 1 atm and 273 K
 N_A atoms

- (C) 0.5 moles of O_3 $1.5N_A$ atoms 11.2 L at 1 atm and 273 K 24 g weight
 (D) 1 g molecules of O_2 1 mole of O_2 $2N_A$ atom 22.4 L at 1 atm and 273 K 32 g weight

2. (A) Moles of $CO_2(g)$ $\frac{44}{44}$ 1

1 mole $CO_2(g)$ 1 g-molecule $CO_2(g)$

(B) Moles of $CH_4(g)$ $\frac{35.2}{16}$ 2.2 moles

V 2.2 22.4 49.28 L at 1 atm and 273 K

(C) Moles of $O_3(g)$ $\frac{48}{48}$ 1

(D) Moles of $N_2O(g)$ $\frac{44}{44}$ 1

Number of O atoms N_A 1 N_A

3. Use % by moles $\frac{M_{avg}}{M_2} \frac{M_1}{M_1}$ 100

% by mass = % by moles $\frac{M_2}{M_{avg}}$

4. (A) Bi_2S_3 Bi^5 S^0
 n -factor (2 2) (2 3) 10

(B) $Al_2(Cr_2O_7)_3$ Cr^3
 n -factor (3 6) 18

(C) FeS_2 Fe_2O_3 SO_2
 n -factor (1 1) (5 2) 11

(D) $Mn(NO_3)_2$ MnO_4^2 NO^0
 n -factor |(4 1) (3 2)| 2

5. (P) CrI_3 $Cr_2O_7^{2-}$ IO_4^-
 n -factor (3 1) (8 3) 27

(Q) $Fe(SCN)_2$ Fe^3 SO_4^{2-} CO_3^{2-} NO_3^-
 n -factor (1 1) (8 2) (8 2) 33

(R) NH_4SCN SO_4^{2-} CO_3^{2-} NO_3^-
 n -factor (8 2) (8 1) 24

(S) As_2S_3 AsO_3 SO_4^{2-}
 n -factor (2 2) (8 3) 28

7. Moles of $NaIO_3$ $\frac{396}{198}$ 2

(1) IO_3^- $3HSO_3^-$ I^- $3SO_4^{2-}$ $3H^+$
 n -factor $\frac{6}{1}$ $\frac{6}{3}$ $\frac{6}{1}$ $\frac{6}{3}$ $\frac{6}{3}$

(2) $5I^-$ IO_3^- $6H^+$ $3I_2$ $3H_2O$
 n -factor $\frac{5}{5}$ $1, \frac{5}{1}$ $\frac{5}{3}$ $\frac{5}{3}$

(A) n -factor of IO_3^- in reaction (2) 5

(B) Moles of HSO_3^- used in (1) reaction 3
 moles of IO_3^- 3 2 6

(C) Moles of I_2 produced moles of I^- produced
 $\frac{3}{5}$ $2 \frac{3}{5}$ 1.2

(D) Equivalents of IO_3^- used in reaction (2)

Equivalents of I

2 1 2

8. (A) Br_2^0 OH^- BrO_3^- Br^0

Intermolecular, Disproportionation Redox Reaction

(B) $FeCl_2$ $KMnO_4$ HCl $MnCl_2$ $FeCl_3$

Intermolecular Redox Reaction

(C) Ag_2O^0 Ag^0 $\frac{1}{2}O_2^0$

Intermolecular Redox

(D) NH_4^3 NO_3^5 N_2O^1 H_2O

Intermolecular comproportionation Redox Reaction

Assertion-Reason Type Question

5. N M n -factor

and n -factor of H_3PO_3 2 (max).

9. $[Fe(CN)_6]^{4-}$ Fe^{3+} CO_2^4 NO_3^5

n -factor (1 1) (2 6) (8 6) 61

10. $As_2S_3^3$ NO_3^5 AsO_4^{3-} NO^2 SO_4^{2-}

n -factor (2 2) (8 3) n f (3 1)
 28 3

11. $NaOH$ H_3PO_4 NaH_2PO_4 H_2O

n -factor 1 1
 Eq. wt of H_3PO_4 $\frac{M}{1}$

13. I_2^0 IO_3^5 I^1

Subjective Problems

1. Mass of liquid in 10 moles $280 \times 10 = 2800 \text{ g}$

Volume $\frac{2800}{1.4} = 2000 \text{ mL} = 2 \text{ L}$

2. $(M) \frac{16}{5.6} = (22.4)$

$M = 64$

$64 \times 32 = 16x$

$x = 2$

3. $[\text{H}] = \frac{(200 \times 1) + (300 \times 6)}{1000}$

$[\text{H}] = 2$

4. $\text{N}_2(\text{g}) \quad \text{H}_2(\text{g}) \quad \text{N}_2\text{H}_2(\text{l})$
 $5 \text{ L} \quad 3 \text{ L} \quad 2 \text{ L}$

$V = (8) + (2) = 6$

5. $n_{\text{SO}_2} = \frac{3.78}{64} = 0.06$; Let moles of CaSO_4 produced be x
 POAC on S

$\text{SO}_2 \quad \text{CaSO}_4$

$(n_s)_{\text{SO}_2} \quad (n_s)_{\text{CaSO}_4}$

$0.06 \times 1 = x \times 1 \quad x = 0.06$

Mass of $\text{CaSO}_4 = 0.06 \times 136 = 8.16$

6. Moles of $\text{H}_2\text{O} = 1$



Consumed moles of Fe $\frac{3}{4}$

Mass of Fe $\frac{3}{4} \times 56 = 42$

7. $n_{\text{KClO}_3} = \frac{122.5}{122.5} = 1$

Moles of atoms $1 \times 1 + 3 \times 5$

8. Let metal sulfate is $\text{M}_2(\text{SO}_4)_x$
 $x = \text{valency of metal}$

Specific heat $\frac{\text{molar mass}}{6.4}$

Molar mass $\frac{6.4}{0.057} = 112.2 \text{ g/mol}$

POAC on metal

$\frac{2}{112.2} = \frac{4.51}{2(112.2) + (96)(x)} \quad x = 2$

$224.4 + 96x + 506.02$

$x = 3$

9. $\text{M}_x\text{Cl}_y \quad \text{MCl}_y (\because \text{valency of Cl} = 1)$

Molar mass of $\text{M}_x\text{Cl}_y = \frac{85.5 \times 2}{171 \text{ g/mol}}$

$(M)(x) + (35.5)(y) = 171 \quad \dots(1)$

Mass of chlorine $\frac{0.835}{1}$


Mass of metallic chloride $\frac{35.5 y}{M(x) + (35.5)(y)}$

$\frac{35.5 y}{M(x) + (35.5)(y)}$

On solving $y = 4$ and $x = 1$

$x = y = 5$

10. $\text{Ba}(\text{OH})_2 + x\text{H}_2\text{O}(\text{aq}) \rightarrow \text{HNO}_3(\text{aq})$



Eq. of $\text{Ba}(\text{OH})_2 + x\text{H}_2\text{O} = \text{eq. of HNO}_3$

$\frac{0.7875}{(171 + 18x)} = 2 \times \frac{1}{4} \times \frac{20}{1000}$

$x = 8$

11. n -factor of acid $\times \text{Basicity}(n)$

Eq. of acid $\times \text{Eq. of base}$

$\frac{2}{600} \times n = \frac{1}{6} \times \frac{100}{1000} \times 1$

$n = 5$

12. In HPh Eq. of NaOH $\times \text{Eq. of Na}_2\text{CO}_3$

$\times \text{Eq. of HCl}$

$(0.5)(1) + (0.5)(1) + (x)(1)$

$x = 1$

In MeOH Eq. of NaOH $+ \text{Eq. of Na}_2\text{CO}_3$

$\times \text{Eq. of NaHCO}_3 + \text{Eq. of HCl}$

$(0.5)(1) + (0.5)(2) + (0.5)(1) + (y)(1)$

$y = 2$

$x = y = 3$

13. $\text{BrO}_3 \quad \text{Br} \quad \text{H} \quad \text{Br}_2(\text{l})$

n -factor 5

$E_{\text{KBrO}_3} = \frac{M_{\text{KBrO}_3}}{5}$

14. $\text{SeO}_2 \quad \text{CrSO}_4 \quad \text{Se}^x \quad \text{Cr}^3$

Eq. of $\text{SeO}_2 \times \text{Eq. of CrSO}_4$

$(12.5)(0.05)(4 - x) = (25)(0.1)(1)$

$(4 - x) = 4$

$x = 0$

15. $\text{Cu}^{2+} \quad \text{KI} \quad \text{I}_2 \quad \text{Cu}$

n -factor $= 1$

n -factor $= 2$



n -factor $= 2 \quad n$ -factor $= 1$



$$\begin{array}{lll} \text{Eq. of Cu}^2 & \text{Eq. of I}_2 \text{ produced} & \text{Eq. of I}_2 \\ \text{reacted} & \text{Eq. of Na}_2\text{S}_2\text{O}_3 & \\ (x)(1) & \frac{40}{1000} (0.1)(1) & \end{array}$$

$$x \quad 4 \quad 10^3 \text{ mol}$$

$$\text{Mass of pure Cu} \quad 4 \quad 10^3 \quad 63.5 \quad 0.254 \text{ g}$$


$$\% \text{ Purity} \quad \frac{0.254}{0.276} \quad 100 \quad 92\%$$

$$\% \text{ impurity} \quad 8\%$$

$$\begin{array}{ll} \text{Eq. of H}_2\text{O}_2 & \text{Eq. of KMnO}_4 \\ (25 \text{ M } 2) & (10 \text{ 0.1 } 5) \\ M & 0.089 \end{array}$$

$$\text{Volume strength} \quad M \quad 11.2 \quad 1$$

$$\text{Balanced redox reaction}$$



$$\begin{array}{ll} \text{M}_2\text{O}_x \text{ (Metallic oxide)} & \text{H}_2 \\ 0.220\text{g} & \\ & M(s) \quad \text{H}_2\text{O}(l) \\ & \quad \quad 0.045\text{g} \end{array}$$

$$\text{POAC on O}$$

$$\frac{0.220}{2M} \cdot x \quad \frac{0.045}{18} \cdot 1$$

$$3.96x \quad 0.09M \quad 0.72x$$

$$E \quad \frac{M}{x} \quad 36$$

$$\frac{E}{9} \quad 4$$

19.

$$\text{Let mass of K}_2\text{Cr}_2\text{O}_7 \quad a \text{ g}$$

$$\text{Mass of KMnO}_4 \quad (10 - a) \text{ g}$$

$$\text{Eq. of K}_2\text{Cr}_2\text{O}_7 \quad \text{Eq. of KMnO}_4$$

$$\text{Eq. of Na}_2\text{S}_2\text{O}_3$$

$$\frac{9}{294} \cdot 6 \quad \frac{10 - a}{158} \cdot 5 \quad (2.2)(0.1)$$

$$a \quad 8.5 \text{ g}$$

$$\text{Mass \% of KMnO}_4 \quad Z \quad 15$$

$$a \quad 5 \quad 10^4$$

20.

$$\% \text{ of Sn}^{2+} \text{ in sample}$$

$$(x) \quad \frac{5 \cdot 10^4 \cdot 11.8}{0.4} \quad 100$$

$$x \sim 15$$

$$\frac{x}{5} \cdot \frac{15}{5} \quad 3$$


2

ATOMIC STRUCTURE

Atomic Number of an Element

Total number of protons present in the nucleus = Total number of electrons present in the atom

Mass number of an element = Number of protons + Number of neutrons.



e. g., $^{23}_{11}\text{Na}$, $^{35}_{17}\text{Cl}$ and so on.

Terms associated with elements

- ❖ Isotopes : Atoms having same number of protons but different mass number.
- ❖ Isobars : Elements having same mass number.
- ❖ Isotones : Elements having same number of neutrons ($A - Z$).
- ❖ Isoelectronic : Species/elements having same number of electrons.
- ❖ Isosters : Species having same number of atoms and electrons
- ❖ Isodiaphers : Elements having same number of $|N - Z|$ or $|A - 2Z|$
- ❖ Paramagnetic : Species having non-zero unpaired electron.
- ❖ Diamagnetic : Species having zero unpaired electron.

Rutherford's Model

- ❖ Electrons, protons & neutrons are the most important fundamental particles of atoms of all elements (Except hydrogen)
- ❖ ^A_ZX , Mass number (A) Atomic number (Z) number of neutrons (n)
- ❖ $R_N = R_0(A)^{1/3}$, $R_0 = 1.33 \times 10^{-13}\text{cm}$ A mass number, R_N Radius of nucleus


- ❖ $\frac{1}{2} m v^2 = K \frac{q_1 q_2}{r}$; r distance of closest approach, v Velocity of α -particle
- m mass of α -particle
- q_1 charge on α -particle
- q_2 charge on metal foil.

Size of the Nucleus

The volume of the nucleus is very small and is only a minute fraction of the total volume of the atom. Nucleus has a diameter of the order of 10^{-12} to 10^{-13} cm and the atom has a diameter of the order of 10^{-8} cm.

Thus, diameter (size) of the atom is 1,00,000 times the diameter of the nucleus.

Electromagnetic Spectrum



Light

- ❖ Photon is considered as massless bundle of energy.
- ❖ Energy of light $E = mc^2$, where m mass of light particle, c speed of light
- ❖ $E_{\text{Photon}} = h \bar{\nu} = \frac{hc}{\lambda} = \frac{1240 \text{ eV} \cdot \text{nm}}{\lambda (\text{nm})}$
where h Planck constant, λ wavelength of photon, $\bar{\nu}$ wave number.
- ❖ Quantum efficiency or Quantum Yield $= \frac{\text{no. of molecules reacting}}{\text{no. of quanta absorbed}}$

Bohr's Model

- ❖ Electrostatic force $= \frac{K q_1 q_2}{r^2}$ where $K = \frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$

- ❖ Potential energy due to electrostatic force $\frac{Kq_1q_2}{r}$
 q_1 charge of electron, q_2 charge of nucleus
- ❖ Potential due to a charge (Q) particle at a distance (r) $\frac{KQ}{r}$
- ❖ Bohr quantization rule $mvr = n \frac{h}{2\pi} = n\hbar$
- ❖ According to Newtons' second law, in a uniform circular motion resultant of all the forces towards centre must be equal to $\frac{mv^2}{r}$.

$$\frac{Kq_1q_2}{r^2} = \frac{mv^2}{r},$$

where q_1 charge of electron, q_2 charge of nucleus, m mass of electron,
 r radius of Bohr's orbit

- ❖ Total energy of electron in n^{th} Bohr orbit

$$E_n = \frac{E_1}{n^2} Z^2 = \frac{2^2 me^4 K^2}{n^2 h^2} Z^2; \quad E_1 = \frac{2^2 me^4 K^2 Z^2}{h^2}$$

$$E_n = 13.6 \frac{Z^2}{n^2} \text{ eV/atom, where } Z \text{ atomic number of single electron atoms/ion,}$$

n principal quantum number of shell, E_1 total energy of electron in I Bohr orbit.

- ❖ Radius of n^{th} Bohr orbit, $r_n = \frac{h^2}{4\pi^2 e^2 m K} \frac{n^2}{Z} = 0.529 \frac{n^2}{Z} \text{ \AA} = r_1 \frac{n^2}{Z}$,

where r_1 radius of I Bohr orbit.

- ❖ Velocity of electron in n^{th} Bohr orbit, $v_n = \frac{2\pi e^2 K}{h} \frac{Z}{n} = 2.18 \times 10^6 \frac{Z}{n} \text{ m/s} = v_1 \frac{Z}{n} \text{ m/s}$

where v_1 velocity of electron in I Bohr orbit.

- ❖ Revolutions per sec $v/2\pi r = 0.657 \times 10^{16} \frac{Z^2}{n^3}$

- ❖ Time for one revolution $2\pi r/v = 1.52 \times 10^{-16} \frac{n^3}{Z^2}$

- ❖ Energy difference between n_1 and n_2 energy level.

$$E = E_{n_2} - E_{n_1} = 13.6 Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \text{ eV/atom} \quad \text{IE} = \frac{1}{n_1^2} - \frac{1}{n_2^2}$$


where IE = ionization energy of single electron species.

- ❖ Ionization energy $E = E_{\text{G.S.}} - E_{\text{G.S.}}; E_{\text{G.S.}}$ Energy of electron in ground state
- ❖ Total energy of electron in terms of kinetic energy (KE) and potential energy (PE)

$$E_n = \text{KE} + \text{PE} = \text{KE} + \frac{\text{PE}}{2}$$

Spectral Lines

- ❖ Rydberg's Equation $\frac{1}{\lambda} = R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) Z^2$; $R_H = 109700 \text{ cm}^{-1}$ Rydberg constant
- ❖ For First line of a series $n_2 = n_1 + 1$
- ❖ Limiting spectral line (series limit) means $n_2 \rightarrow \infty$
- ❖ H_γ line means $n_2 = n_1 + 1$; also known as line of longest wavelength, least E
- ❖ Similarly H_α line means $n_2 = n_1 + 2$
- ❖ When electrons de-excite from higher energy level (n_2) to ground state in atomic sample, then number of spectral lines observed in the spectrum $\frac{n(n-1)}{2}$
- ❖ When electrons de-excite from higher energy level (n_2) to lower energy level (n_1) in atomic sample, then number of spectral line observed in the spectrum $\frac{(n_2 - n_1)(n_2 - n_1 + 1)}{2}$
- ❖ When electron de-excites from higher energy level (n_2) to lower energy level (n_1) in isolated atom, then number of spectral line observed in the spectrum $(n_2 - n_1)$



Photoelectric Effect

- ❖ When radiation with certain minimum frequency (ν_0 called threshold frequency), strikes the surface of a metal, electrons (called photoelectrons) are ejected from the surface.
- ❖ Kinetic energy of photoelectron $h\nu - w = h\nu - h\nu_0$
where w = work function
 ν_0 = Threshold frequency
- ❖ If $\nu < \nu_0$, then photoelectric effect takes place.
- ❖ Accelerating potential $eV = KE = \frac{1}{2}mv^2$

de-Broglie Hypothesis

- ❖ All material particles possess wave character as well as particle character.
- ❖ $h/mv = h/p$
- ❖ The circumference of the n^{th} orbit is equal to n times of wavelength of moving electron i.e.,
 $2\pi r_n = n\lambda$
 Number of waves = n principal quantum number
- ❖ Wavelength of electron (λ) $= \frac{150}{\sqrt{V(\text{volts})}} \text{ \AA}$
- ❖ Wave nature of electron was confirmed by Davisson and Germer experiment.

Heisenberg Uncertainty Principle

- ❖ According to this principle, "it is impossible to measure simultaneously the position and momentum of a microscopic particle with absolute accuracy".
 If one of them is measured with greater accuracy, the other becomes less accurate.
- ❖ $\Delta x \Delta p = h/4$ or $(\Delta x)(\Delta v) = \frac{h}{4m}$ or $(\Delta x)(\Delta \lambda) = \frac{h}{4m\lambda}$
 where Δx Uncertainty in position, Δp Uncertainty in momentum.
 Δv Uncertainty in velocity, $\Delta \lambda$ Uncertainty in wavelength.
 m mass of microscopic particle, λ Wavelength of microscopic particle
- ❖ Heisenberg replaced the concept of orbit by that of orbital.

Schrodinger Equation

- ❖ Schrodinger equation is central equation of wave mechanics according to following equation.

$$\frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} + \frac{\partial^2 \psi}{\partial z^2} + \frac{8\pi^2m}{h^2}(E - V)\psi = 0$$

Wave function $\psi(x, y, z)$

E Total energy of particle

V Potential energy of particle.

- ❖ A Schrodinger equation leads to infinite solutions.
- ❖ Most of the solution are not realistic (or acceptable). Only few solution can be accepted.
- ❖ Each solution - (x, y, z) correspond to a definite energy state which depends on quantum number n, l & m .

By proper mathematical manipulation the main equation is broken in two parts and solved separately.

(i) Radial part contain only 'r', depends on quantum numbers n & l .

(ii) Angular part contain θ and ϕ , depends on quantum numbers l & m .

Each value of n, l, m contain all the information about that particular quantum state.

- ❖ **Atomic Orbital** : This is a three dimensional space around the nucleus within which the probability of finding the electron is maximum.
- ❖ **Degenerate Orbital** : Orbitals with same value of n and l of same subshell are degenerate orbitals, which have same energy.

For example, $2p_x, 2p_y, 2p_z$ etc.

❖ **Radial Probability Density** $4 r^2 R^2(r)$

It is the probability of finding electron in the region between r and $r + dr$

❖ **Radial Node** : It is zero electron density region. $R^2(r) = 0$ or $R(r) = 0$

❖ **Nodal Point** : It is a point ($r = 0$) where electron density is zero.

❖ **Nodal Planes** : It is plane by which two lobes are separated and electron density is zero here.

Quantum Number

There are four of quantum number which are required in order to explain e^- in atom.

❖ **Principal quantum number (n)** : It determines the size of an orbital. Each value of n represents a shell of orbital. Possible values of $n = 1, 2, 3, 4, \dots$

❖ **Azimuthal quantum number (l)** : It determines shape of an orbital. Each value of l represents a subshell of an orbital. Possible values of $l = 0, 1, 2, \dots (n - 1)$

❖ **Magnetic quantum number (m)** : It decides orientation of orbital in space.

Possible values of $m = -l, -l + 1, \dots, 0, 1, 2, l$


Total $(2l + 1)$ values are possible form for each value of l .

❖ **Spin quantum number (s)** : It is intrinsic property of an electron. The electron has two spin states. Possible values of $s = \frac{1}{2}, -\frac{1}{2}$

Shell, sub-shells and orbitals

Shell (n)	Sub-shells (l)	Orbitals (m)
1	0	0
2	0	0
	1	1, 0
	0	0
3	1	1, 0
	2	2, 1, 0

Various Curves



Important Points on Quantum Number

- ❖ Orbital angular momentum $\frac{h}{2} \sqrt{l(l+1)}$
- ❖ Spin angular momentum $\frac{h}{2} \sqrt{S(S+1)}$
- ❖ Spin Magnetic moment () $\sqrt{n(n+2)}$ B.M. ; n = number of unpaired electron
- ❖ Maximum number of electrons in a shell $2n^2$
- ❖ Maximum number of electrons in a subshell $2(2l+1)$
- ❖ Maximum number of electrons in an orbital 2
- ❖ Total number of orbitals in a subshell $2l+1$
- ❖ Number of subshells in a shell n
- ❖ Number of orbitals in a shell n^2
- ❖ Radial Nodes $(n-l-1)$
- ❖ Angular nodes l
- ❖ Total nodes $(n-1)$
- ❖ Azimuthal quantum number 0 1 2 3 4
- Name of sub-shell s p d f g

Pauli's Exclusion Principle

No two electrons in an atom can have the same set of all the four quantum numbers, *i.e.*, an orbital cannot have more than 2 electrons because three quantum numbers (principal, azimuthal and magnetic) at the most may be same but the fourth will be different, *i.e.*, spins will be in opposite directions.

Aufbau Principle

Electrons are filled in various orbitals in order of their increasing energies. An orbital of lowest energy is filled first. The sequence of orbitals in order of their increasing energy is :

$1s, 2s, 2p, 3s, 3p, 4s, 3d, 4p, 5s, 4d, 5p, 6s, 4f, 5d, 6p, 7s, 5f, 6d, \dots$

The energy of the orbitals is governed by $(n + l)$ rule.

Hund's Rule

No electron pairing takes place in the orbitals in a sub energy shell until each orbital is occupied by an electron with parallel spin. Exactly half filled and fully filled orbitals make the atoms more stable, *i.e.*, $p^3, p^6, d^5, d^{10}, f^7$ and f^{14} configurations are more stable.



Level 1





- Which of the following pair is isodiaphers?
 - ${}^6_{14}\text{C}$ and ${}^{23}_{11}\text{Na}$
 - ${}^{24}_{12}\text{Mg}$ and ${}^{23}_{11}\text{Na}$
 - ${}^4_2\text{He}$ and ${}^{16}_8\text{O}$
 - ${}^{12}_6\text{C}$ and ${}^{15}_7\text{N}$
- Which of the following does not characterise X-rays?
 - The radiation can ionise the gas
 - It causes fluorescence effect on ZnS
 - It is deflected by electric and magnetic fields
 - Its wavelength is shorter than ultraviolet rays
- The ratio of specific charge of a proton and an α -particle is :
 - 2 : 1
 - 1 : 2
 - 1 : 4
 - 1 : 1
- The increasing order for the values of e/m (charge/mass) is :
 - $e, p, n,$
 - $n, p, e,$
 - $n, p, , e$
 - $n, , p, e$
- The mass to charge ratio (m/e) for a cation is $1.5 \times 10^8 \text{ kg/C}$. What is the mass of this cation?
 - $2.4 \times 10^{19} \text{ g}$
 - $2.4 \times 10^{27} \text{ g}$
 - $2.4 \times 10^{24} \text{ g}$
 - None of these
- Rutherford's experiment on scattering of alpha particles showed for the first time that atom has :
 - electrons
 - protons
 - nucleus
 - neutrons
- α -particles are represented by :
 - lithium atoms
 - helium nuclei
 - hydrogen nuclei
 - None of these
- In Bohr's stationary orbits :
 - electrons do not move
 - electrons move emitting radiations
 - energy of the electron remains constant
 - angular momentum of the electron is $h/2$
- On the basis of Bohr's model, the radius of the 3rd orbit is :
 - equal to the radius of first orbit
 - three times the radius of first orbit
 - five times the radius of first orbit
 - nine times the radius of first orbit
- The correct expression derived for the energy of an electron in the n^{th} energy level is for H-atom :
 - $E_n = \frac{2}{n^2} \frac{me^4 K^2}{h^2}$
 - $E_n = \frac{2me^4 K^2}{n^2 h^2}$
 - $E_n = \frac{2}{n^2} \frac{me^2 K^2}{h^2}$
 - $E_n = \frac{2}{n^2} \frac{me^4 K^2}{h^2}$
- Ionization energy for hydrogen atom in ergs, Joules and eV respectively is :
 - $21.8 \times 10^{12}, 218 \times 10^{20}, 13.6$
 - $13.6 \times 218 \times 10^{20}, 21.8 \times 10^{13}$
 - $21.8 \times 10^{20}, 13.6, 21.8 \times 10^{13}$
 - $21.8 \times 10^{13}, 13.6, 21.8 \times 10^{20}$
- For any H like system, the ratio of velocities of electron in I, II & III orbit i.e., $V_1 : V_2 : V_3$ will be:
 - 1 : 2 : 3
 - 1 : 1/2 : 1/3
 - 3 : 2 : 1
 - 1 : 1 : 1

13. The volume of nucleus is about :
 (a) 10^4 times to that of an atom (b) 10^{15} times to that of an atom
 (c) 10^5 times to that of an atom (d) 10^{10} times to that of an atom
14. An electron in an atom jumps in such a way that its kinetic energy changes from x to $\frac{x}{4}$. The change in potential energy will be :
 (a) $\frac{3}{2}x$ (b) $\frac{3}{8}x$ (c) $\frac{3}{4}x$ (d) $\frac{3}{4}x$
15. The potential energy of an electron in the hydrogen atom is 6.8 eV. Indicate in which excited state, the electron is present?
 (a) first (b) second (c) third (d) fourth
16. What is the potential energy of an electron present in N -shell of the Be^3 ion?
 (a) 3.4 eV (b) 6.8 eV (c) 13.6 eV (d) 27.2 eV
17. The kinetic and potential energy (in eV) of electron present in third Bohr's orbit of hydrogen atom are respectively :
 (a) 1.51, 3.02 (b) 1.51, 3.02 (c) 3.02, 1.51 (d) 1.51, 1.51
18. The distance between 4th and 3rd Bohr orbits of He is :
 (a) 2.645×10^{10} m (b) 1.322×10^{10} m (c) 1.851×10^{10} m (d) None of these
19. What atomic number of an element "X" would have to become so that the 4th orbit around X would fit inside the 1 Bohr orbit of H atom ?
 (a) 3 (b) 4 (c) 16 (d) 25
20. The ratio of velocity of the electron in the third and fifth orbit of Li^{2+} would be :
 (a) 3 : 5 (b) 5 : 3 (c) 25 : 9 (d) 9 : 25
21. If radius of second stationary orbit (in Bohr's atom) is R then radius of third orbit will be :
 (a) $R/3$ (b) $9R$ (c) $R/9$ (d) $2.25 R$
22. Which Bohr's orbit of Be^3 has the same orbit radius as that of the ground state of hydrogen atom?
 (a) 3 (b) 2 (c) 4 (d) 5
23. Select the incorrect graph for velocity of e^- in an orbit vs. Z , $\frac{1}{n}$ and n :
- (a)

(b)





(c)

(d)
24. What is the frequency of revolution of electron present in 2nd Bohr's orbit of H-atom?
 (a) $1.016 \times 10^{16} \text{ s}^{-1}$ (b) $4.065 \times 10^{16} \text{ s}^{-1}$ (c) $1.626 \times 10^{15} \text{ s}^{-1}$ (d) $8.2 \times 10^{14} \text{ s}^{-1}$
25. According to Bohr's atomic theory, which of the following relation is correct ?
 (a) Potential energy of electron $\frac{Z^2}{n^2}$
 (b) The product of velocity of electron and principal quantum number (n) Z^2
 (c) Frequency of revolution of electron in an orbit $\frac{Z^2}{n^3}$
 (d) Coulombic force of attraction on the electron $\frac{Z^2}{n^2}$

26. Number of waves produced by an electron in one complete revolution in n^{th} orbit is :
 (a) n (b) n^2 (c) $(n-1)$ (d) $(2n-1)$
27. Which of the following statement does not form part of Bohr's model of the hydrogen atom ?
 (a) Energy of the electrons in the orbit is quantized
 (b) The electron in the orbit which is nearest to the nucleus has the lowest energy
 (c) Electrons revolve in different orbits around the nucleus
 (d) The position and velocity of the electrons in the orbit cannot be determined simultaneously
28. What is the separation energy (in eV) for electron present in the first excited state of Be^3 ?
 (a) 13.6 eV (b) 27.2 eV (c) 40.8 eV (d) 54.5 eV
29. If in Bohr's model, for unielectronic atom, time period of revolution is represented as $T_{n,Z}$ where n represents shell no. and Z represents atomic number then the value of $T_{1,2} : T_{2,1}$ will be :
 (a) 8 : 1 (b) 1 : 8 (c) 1 : 1 (d) 1 : 32
30. Which of the following is discretized in Bohr's theory?
 (a) Potential energy (b) Kinetic energy
 (c) Velocity (d) Angular momentum
31. What is the ratio of time periods (T_1/T_2) in second orbit of hydrogen atom to third orbit of He^+ ion?
 (a) 8/27 (b) 32/27 (c) 27/32 (d) None of these
32. The mass of an electron is m , charge is e and it is accelerated from rest through a potential difference of V volts. The velocity acquired by electron will be :
 (a) $\sqrt{\frac{V}{m}}$ (b) $\sqrt{\frac{eV}{m}}$ (c) $\sqrt{\frac{2eV}{m}}$ (d) zero
33. If the ionization energy of He is 19.6×10^{18} J per atom then the energy of Be^3 ion in the second stationary state is :
 (a) 4.9×10^{18} J (b) 44.1×10^{18} J (c) 11.025×10^{18} J (d) None of these
34. The energy of the second Bohr orbit in the hydrogen atom is 3.41 eV. The energy of the second Bohr orbit of He^+ ion would be :
 (a) 0.85 eV (b) -13.64 eV (c) -1.70 eV (d) -6.82 eV
35. The energy of an electron moving in n^{th} Bohr's orbit of an element is given by $E_n = \frac{-13.6}{n^2} Z^2$ eV/atom (Z = atomic number). The graph of E vs. Z^2 (keeping "n" constant) will be :
 (a)  (b)  (c)  (d) 
36. If ϵ_0 be the permittivity of vacuum and r be the radius of orbit of H-atom in which electron is revolving, then velocity of electron is given by :

$$(a) \propto \frac{e}{\sqrt{4} \, a_0 r m} \quad (b) \propto e \sqrt{4} \, a_0 r m \quad (c) \propto \frac{4 \, a_0 r m}{e} \quad (d) \propto \frac{4 \, a_0 r m}{e^2}$$

37. Which of the following statement(s) is/are consistent with the Bohr theory of the atom (and no others)?
- (1) An electron can remain in a particular orbit as long as it continuously absorbs radiation of a definite frequency.
 - (2) The lowest energy orbits are those closest to the nucleus.
 - (3) All electrons can jump from the *K* shell to the *M* shell by emitting radiation of a definite frequency.
- (a) 1, 2, 3 (b) 2 only (c) 3 only (d) 1, 2
38. The ionization potential for the electron in the ground state of the hydrogen atom is 13.6 eV atom⁻¹. What would be the ionization potential for the electron in the first excited state of Li²⁺?
- (a) 3.4 eV (b) 10.2 eV (c) 30.6 eV (d) 6.8 eV
39. What is the energy content per photon (J) for light of frequency 4.2×10^{14} Hz?
- (a) 2.8×10^{21} (b) 2.5×10^{19} (c) 2.8×10^{19} (d) 2.5×10^{18}
40. Wavelength for high energy EMR transition in H-atom is 91 nm. What energy is needed for this transition?
- (a) 1.36 eV (b) 1240 eV (c) 13 eV (d) 13.6 eV
41. Which graph shows how the energy *E* of a photon of light is related to its wavelengths (λ)?



42. Assume that 2×10^{17} J of light energy is needed by the interior of the human eye to see an object. How many photons of yellow light with 595.2 nm are needed to generate this minimum energy?
- (a) 6 (b) 30 (c) 45 (d) 60
43. Line spectra is characteristic of :
- (a) molecules (b) atoms (c) radicals (d) none of these
44. The spectrum produced from an element is :
- (a) atomic spectrum (b) line spectrum
(c) absorption spectrum (d) any one of the above

45. Electronic transition in He⁺ ion takes from n_2 to n_1 shell such that :



$$2n_2 - 3n_1 = 18$$



$$2n_2 - 3n_1 = 6$$

What will be the total number of photons emitted when electrons transit to n_1 shell ?

- (a) 21 (b) 15 (c) 20 (d) 10
46. Which of the following expressions represents the spectrum of Balmer series (If n is the principal quantum number of higher energy level) in Hydrogen atom ?
- (a) $\bar{\nu} = \frac{R(n-1)(n-1)}{n^2}$ (b) $\bar{\nu} = \frac{R(n-2)(n-2)}{4n^2}$
- (c) $\bar{\nu} = \frac{R(n-2)(n-2)}{n^2}$ (d) $\bar{\nu} = \frac{R(n-1)(n-1)}{4n^2}$
47. Multiple or fine structure of spectral lines is due to :
- (a) presence of main energy levels (b) presence of sub-levels
- (c) presence of electronic configuration (d) is not a characteristics of the atom
48. With increasing principal quantum number, the energy difference between adjacent energy levels in H-atom:
- (a) decreases (b) increases
- (c) remains constant (d) decreases for low value of Z and increases for higher value of Z .
49. Find the value of wave number ($\bar{\nu}$) in terms of Rydberg's constant, when transition of electron takes place between two levels of He⁺ ion whose sum is 4 and difference is 2.
- (a) $\frac{8R}{9}$ (b) $\frac{32R}{9}$ (c) $\frac{3R}{4}$ (d) none of these
50. What is the wavelength in nm of the spectral line associated with a transition from $n = 3$ to $n = 2$ for the Li²⁺ ion?
- (a) 219 (b) 656 (c) 73.0 (d) 486
51. What is the energy (kJ/mol) associated with the de-excitation of an electron from $n = 6$ to $n = 2$ in He⁺ ion?
- (a) 1.36×10^6 (b) 1.36×10^3 (c) 1.16×10^3 (d) 1.78×10^3
52. What is the shortest wavelength line in the Paschen series of Li²⁺ ion?
- (a) $\frac{R}{9}$ (b) $\frac{9}{R}$ (c) $\frac{1}{R}$ (d) $\frac{9R}{4}$
53. What is the maximum wavelength line in the Lyman series of He⁺ ion?
- (a) $3R$ (b) $\frac{1}{3R}$ (c) $\frac{4}{4R}$ (d) None of these
54. Which of the following electron transitions in a hydrogen atom will require the largest amount of energy?
- (a) from $n = 1$ to $n = 2$ (b) from $n = 2$ to $n = 4$
- (c) from $n = 5$ to $n = 1$ (d) from $n = 3$ to $n = 5$
55. Which electronic transition in a hydrogen atom, starting from the orbit $n = 7$, will produce infrared light of wavelength 2170 nm? (Given : $R_H = 1.09677 \times 10^7 \text{ m}^{-1}$)


- (a) $n = 7$ to $n = 6$ (b) $n = 7$ to $n = 5$
 (c) $n = 7$ to $n = 4$ (d) $n = 7$ to $n = 3$
56. Hydrogen atoms in the ground state are excited by monochromatic radiation of wavelength λ . The resulting spectrum consists of maximum 15 different lines. What is the wavelength ? ($R_H = 109737 \text{ cm}^{-1}$)
 (a) $937.3 \text{ \AA}$ (b) $1025 \text{ \AA}$ (c) $1236 \text{ \AA}$ (d) None of these
57. Electromagnetic radiation (photon) with highest wavelength results when an electron in the hydrogen atom falls from $n = 6$ to :
 (a) $n = 1$ (b) $n = 2$ (c) $n = 3$ (d) $n = 5$
58. When an electron jumps from L to K shell -
 (a) energy is absorbed
 (b) energy is released
 (c) energy is neither absorbed nor released
 (d) energy is sometimes absorbed and some times released
59. How do the energy gaps between successive electron energy levels in an atom vary from low to high n values?
 (a) All energy gaps are the same
 (b) The energy gap decreases as n increases
 (c) The energy gap increases as n increases
 (d) The energy gap changes unpredictably as n increases
60. The H-spectrum shows:
 (a) Heisenberg's uncertainty principle (b) diffraction
 (c) polarization (d) presence of quantized energy level
61. Splitting of spectral lines under the influence of magnetic field is called
 (a) Zeeman effect (b) Stark effect
 (c) photoelectric effect (d) None of these
62. The colour of sky is due to
 (a) absorption of light by atmospheric gases (b) transmission of light
 (c) wavelength of scattered light (d) all of the above
63. In photoelectric effect, the kinetic energy of photoelectrons increases linearly with the
 (a) wavelength of incident light (b) frequency of incident light
 (c) velocity of incident light (d) atomic mass of an element
64. Slope of V_0 vs ν curve is (where V_0 = Stopping potential, ν = subjected frequency)
 (a) e (b) $\frac{h}{e}$ (c) (d) h
65. According to Einstein's photoelectric equation, the graph between kinetic energy of photoelectrons ejected and the frequency of the incident radiation is :









66. The photoelectric emission from a surface starts only when the light incident upon the surface has certain minimum :
 (a) intensity (b) wavelength (c) frequency (d) velocity
67. If λ_0 and λ be the threshold wavelength and the wavelength of incident light, the speed of photo-electrons ejected from the metal surface is :
 (a) $\sqrt{\frac{2h}{m}(\frac{1}{\lambda_0} - \frac{1}{\lambda})}$ (b) $\sqrt{\frac{2hc}{m}(\frac{1}{\lambda_0} - \frac{1}{\lambda})}$ (c) $\sqrt{\frac{2hc}{m}(\frac{1}{\lambda} - \frac{1}{\lambda_0})}$ (d) $\sqrt{\frac{2h}{m}(\frac{1}{\lambda} - \frac{1}{\lambda_0})}$
68. A light source of wavelength λ illuminates a metal and ejects photo-electrons with $(K.E.)_{\max} = 1 \text{ eV}$.
 Another light source of wavelength $\frac{\lambda}{3}$, ejects photo-electrons from same metal with $(K.E.)_{\max} = 4 \text{ eV}$.
 Find the value of work function?
 (a) 1 eV (b) 2 eV (c) 0.5 eV (d) None of these
69. Electromagnetic radiation having $310 \text{ \AA}$ is subjected to a metal sheet having work function 12.8 eV. What will be the velocity of photo-electrons having maximum kinetic energy.
 (a) 0, no emission will occur (b) $4.352 \times 10^6 \text{ m/s}$
 (c) $3.09 \times 10^6 \text{ m/s}$ (d) $8.72 \times 10^6 \text{ m/s}$
70. The ratio of slopes of $K_{\max}$ vs. ν and V_0 vs. ν curves in the photoelectric effect gives (ν frequency, $K_{\max}$ maximum kinetic energy, V_0 stopping potential) :
 (a) charge of electron
 (b) Planck's constant
 (c) work function
 (d) the ratio of Planck's constant and electronic charge
71. Radiation corresponding to the transition $n = 4$ to $n = 2$ in hydrogen atoms falls on a certain metal (work function = 2.5 eV). The maximum kinetic energy of the photo-electrons will be :
 (a) 0.55 eV (b) 2.55 eV (c) 4.45 eV (d) None of these
72. Select the incorrect statement.
 (a) K.E. of photo-electron does not depend upon the wavelength of incident radiation
 (b) Photoelectric current depends on intensity of incident radiation and not on frequency
 (c) Stopping potential depends on frequency of radiation and not on intensity
 (d) None of these
73. Which is the de-Broglie equation?
 (a) $h = p\lambda$ (b) $h = p\lambda^{-1}$ (c) $h = p\lambda^2$ (d) $h = p\lambda^{-2}$

74. Which of the following has the largest de Broglie wavelength (all have equal velocity)?
 (a) CO_2 molecule (b) NH_3 molecule (c) Electron (d) Proton
75. The de-Broglie wavelength associated with a particle of mass 10^{-6} kg with a velocity of 10 m s^{-1} is :
 (a) $6.63 \times 10^{-22} \text{ m}$ (b) $6.63 \times 10^{-29} \text{ m}$ (c) $6.63 \times 10^{-31} \text{ m}$ (d) $6.63 \times 10^{-34} \text{ m}$
76. For two particles A and B, curves are plotted $\sqrt{V}$ against de-Broglie wavelengths, where V is the potential on the particles. Which of the following relation is correct about the mass of particles?



- (a) $m_A > m_B$ (b) $m_A < m_B$ (c) $m_A = m_B$ (d) $m_A < m_B$
77. Which of following graphs correctly represents the variation of particle momentum with de-Broglie wavelength ?



78. An excited state of H atom emits a photon of wavelength λ and returns in the ground state. The principal quantum number of excited state is given by :
 (a) $\sqrt{R(\lambda - R)}$ (b) $\sqrt{\frac{R}{(\lambda - R)}}$ (c) $\sqrt{R(\lambda + R)}$ (d) $\sqrt{\frac{R(\lambda + R)}{R}}$
79. A dye absorbs a photon of wavelength λ and re-emits the same energy into two photons of wavelengths λ_1 and λ_2 respectively. The wavelength λ is related with λ_1 and λ_2 as :
 (a) $\frac{1}{\lambda} = \frac{1}{\lambda_1} + \frac{1}{\lambda_2}$ (b) $\frac{1}{\lambda} = \frac{1}{\lambda_1} - \frac{1}{\lambda_2}$ (c) $\frac{1}{\lambda} = \frac{1}{\lambda_1} + \frac{1}{\lambda_2}$ (d) $\frac{1}{\lambda} = \left(\frac{1}{\lambda_1} + \frac{1}{\lambda_2}\right)^2$
80. Be^{3+} and a proton are accelerated by the same potential. Their de-Broglie wavelengths have the ratio (assume mass of proton = mass of neutron) :
 (a) 1 : 2 (b) 1 : 4 (c) 1 : 1 (d) $1 : 3\sqrt{3}$

81. The de-Broglie wavelength of an electron accelerated by an electric potential of V volts is given by :
- (a) $\frac{1.23}{\sqrt{m}}$ (b) $\frac{1.23}{\sqrt{h}} m$ (c) $\frac{1.23}{\sqrt{V}} \text{ nm}$ (d) $\frac{1.23}{V}$
82. An electron travels with a velocity of $x \text{ ms}^{-1}$. For a proton to have the same de-Broglie wavelength, the velocity will be approximately :
- (a) $\frac{1840}{x}$ (b) $\frac{x}{1840}$ (c) $1840 x$ (d) x
83. The momentum (in kg-m/s) of photon having 6 MeV energy is :
- (a) 3.2×10^{-21} (b) 2.0 (c) 1.6×10^{-21} (d) none of these
84. The number of photons of light having wave number ' x ' in 10 J of energy source is :
- (a) $10hc x$ (b) $\frac{hc}{10x}$ (c) $\frac{10}{hc x}$ (d) none of these
85. Which of the following relates to photons both as wave motion and as a stream of particles?
- (a) Interference (b) $E = mc^2$ (c) Diffraction (d) $E = h\nu$
86. If a_0 be the radius of first Bohr's orbit of H-atom, the de-Broglie's wavelength of an electron revolving in the second Bohr's orbit will be :
- (a) $6 a_0$ (b) $4 a_0$ (c) $2 a_0$ (d) None of these
87. Energy required to ionise 2 mole of gaseous He ion present in its ground state is :
- (a) 54.4 eV (b) $108.8 N_A \text{ eV}$ (c) $54.4 N_A \text{ eV}$ (d) 108.8 eV
88. Which of the following is the most correct expression for Heisenberg's uncertainty principle ?
- (a) $x \cdot p \geq \frac{h}{4}$ (b) $x \cdot p \geq \frac{h}{4}$ (c) $x \cdot p \geq \frac{h}{4}$ (d) $x \cdot v \geq \frac{h}{4}$
89. The Heisenberg uncertainty principle can be applied to :
- (a) a cricket ball (b) a football (c) a jet aeroplane (d) an electron
90. The mass of a particle is 10^{-10} g and its radius is $2 \times 10^{-4} \text{ cm}$. If its velocity is 10^6 cm sec^{-1} with 0.0001% uncertainty in measurement, the uncertainty in its position is :
- (a) $5.2 \times 10^{-8} \text{ m}$ (b) $5.2 \times 10^{-7} \text{ m}$ (c) $5.2 \times 10^{-6} \text{ m}$ (d) $5.2 \times 10^{-9} \text{ m}$
91. If an electron is travelling at 200 m/s within 1 m/s uncertainty, what is the theoretical uncertainty in its position in μm (micrometer)?
- (a) 14.5 (b) 29 (c) 58 (d) 114
92. The wave character of moving electron was experimentally verified by :
- (a) de-Broglie (b) A. Einstein (c) Germer (d) Schrodinger
93. "The exact path of electron in 2p-orbital cannot be determined." The above statement is based upon :
- (a) Hund's Rule (b) Bohr's Rule
(c) uncertainty principle (d) aufbau principle
94. Which series of subshells is arranged in the order of increasing energy for multi-electron atoms?
- (a) 6s, 4f, 5d, 6p (b) 4f, 6s, 5d, 6p (c) 5d, 4f, 6s, 6p (d) 4f, 5d, 6s, 6p
95. The correct Schrodinger's wave equation for a electron with total energy E and potential energy V is given by :

- (a) $\frac{1}{x^2} - \frac{1}{y^2} - \frac{1}{z^2} - \frac{8}{mh^2}(E - V) = 0$ (b) $\frac{1}{x^2} - \frac{1}{y^2} - \frac{1}{z^2} - \frac{8m}{h^2}(E - V) = 0$
 (c) $\frac{1}{x^2} - \frac{1}{y^2} - \frac{1}{z^2} - \frac{8}{h^2}m(E - V) = 0$ (d) None of the above

96. Wave mechanical model of the atom depends upon :

- (a) de-Broglie concept of dual nature of electron
 (b) Heisenberg uncertainty principle
 (c) Schrodinger uncertainty principle
 (d) All of these

97. In Schrodinger wave mechanical model, $\psi^2(r, \theta, \phi)$ represents :

- (a) amplitude of electron wave
 (b) probability density of electron
 (c) total probability of finding electron around nucleus
 (d) orbit

98. Radial amplitude of electron wave can be represented by :

- (a) $R(r)$ (b) $R^2(r)$ (c) $4r^2$ (d) $4r^2R^2(r)$

99. Arrange the following orbitals of H-atom in the increasing order of their energy.

$3p_x, 2s, 4d_{xy}, 3s, 4p_z, 3p_y, 4s$

- (a) $2s < 3s < 3p_x < 3p_y < 4s < 4p_z < 4d_{xy}$
 (b) $2s < 3s < 3p_x < 3p_y < 4s < 4p_z < 4d_{xy}$
 (c) $2s < 3s < 3p_x < 3p_y < 4s < 4p_z < 4d_{xy}$
 (d) $2s < 3s < 3p_x < 3p_y < 4s < 4p_z < 4d_{xy}$

100. In a hydrogen atom, which orbital is higher in energy than a 3s-orbital ?

- (a) 2s (b) 3p (c) 3d (d) 4s

101. The radii of maximum probability for 3s, 3p and 3d electrons are in the order :

- (a) $(r_{\max})_{3d} < (r_{\max})_{3p} < (r_{\max})_{3s}$ (b) $(r_{\max})_{3d} < (r_{\max})_{3s} < (r_{\max})_{3p}$
 (c) $(r_{\max})_{3s} < (r_{\max})_{3p} < (r_{\max})_{3d}$ (d) None of these

102. The correct order of closeness of 3s, 3p, 3d orbitals of nucleus is :

- (a) $3d < 3p < 3s$ (b) $3s < 3p < 3d$
 (c) $3s < 3d < 3p$ (d) $3d < 3s < 3p$

103. The correct order of total number of node of atomic orbitals is:

- (a) 4f < 6s < 5d (b) 6s < 5d < 4f
 (c) 4f < 5d < 6s (d) 5d < 4f < 6s

104. If the subsidiary quantum number of a subenergy level is 4, then no. of degenerate orbitals are :

- (a) 3 (b) 5 (c) 9 (d) 11

105. Which two orbitals are located along the axis, and not between the axis?

- (a) d_{xy}, d_{z^2} (b) d_{xy}, p_z (c) d_{yz}, p_x (d) $p_z, d_{x^2 - y^2}$

106. In a set of degenerate orbitals, the electrons distribute themselves to retain similar spins as far as possible. This statement is attributed to :

- (a) Pauli's exclusion principle
 (b) aufbau principles

(c) Hund's Rule

(d) Slater rule

107. Which of the following rules could explain the presence of three unpaired electrons in N-atom?

(a) Hund's rule

(b) aufbau's principles

(c) Heisenberg's uncertainty principle

(d) Pauli's exclusion principle

108. Pauli's exclusion principle states that:

(a) nucleus of an atom contains no negative charge

(b) electrons move in circular orbits around the nucleus

(c) electrons occupy orbitals of lowest energy

(d) All the four quantum numbers of two electrons in an atom cannot be equal.

109. For which of the following sets of quantum numbers, an electron will have the highest energy?

	n	l	m	s		n	l	m	s
(a)	3	2	1	$-1/2$	(b)	4	3	-1	$+1/2$
(c)	4	1	-1	$+1/2$	(d)	5	0	0	$-1/2$

110. Which of the following statements concerning the four quantum numbers is false ?

 (a) n gives idea of the size of an orbital

 (b) l gives the shape of an orbital

 (c) m gives the energy of the electron in the orbital

 (d) s gives the direction of spin of the electron in an orbital

111. Maximum number of electrons in a subshell is given by

(a) $(2l + 1)$	(b) $2(2l + 1)$	(c) $(2l + 1)^2$	(d) $2(2l + 1)^2$
----------------	-----------------	------------------	-------------------

112. In any subshell, the maximum number of electrons having same value of spin quantum number is :

(a) $\sqrt{l(l + 1)}$	(b) $l + 2$	(c) $2l + 1$	(d) $4l + 2$
-----------------------	-------------	--------------	--------------

113. The orbital angular momentum of $3p$ electron is :

(a) $\sqrt{3} h$	(b) $\sqrt{6} h$	(c) zero	(d) $\sqrt{2} \frac{h}{2}$
------------------	------------------	----------	----------------------------

114. The atomic orbitals are progressively filled in order of increasing energy. The principle is called as:

(a) Hund's rule

(b) aufbau principle


(c) exclusion principle

(d) de-Broglie rule

115. The orbital diagram in which both the Pauli's exclusion principle and Hund's rule are violated, is :

(a) $\uparrow\downarrow$ $\downarrow$ $\downarrow$ $\downarrow$	(b) $\uparrow\downarrow$ $\uparrow\downarrow$ $\uparrow$	(c) $\uparrow\downarrow$ $\uparrow\downarrow$ $\uparrow$	(d) $\uparrow\downarrow$ $\uparrow\uparrow$ $\uparrow\downarrow$
---	--	--	--

116. Which of the following elements is represented by the electronic configuration ?



(a) Nitrogen


(b) Fluorine

(c) Oxygen

(d) Neon

117. The ratio of magnetic moments of Fe (III) and Co (II) is :

- (a) $\sqrt{5} : \sqrt{7}$ (b) $\sqrt{35} : \sqrt{15}$ (c) 7 : 3 (d) $\sqrt{24} : \sqrt{15}$

118. If the electronic structure of oxygen atom is written as $1s^2, 2s^2$  it would violate

- (a) Hund's rule (b) Pauli's exclusion principle
(c) both Hund's and Pauli's principles (d) None of these

119. A compound of vanadium has a magnetic moment () of 1.73 BM. If the vanadium ion in the compound is present as V^x , then, the value of x is?

- (a) 1 (b) 2 (c) 3 (d) 4

120. d^6 configuration will result in total spin of:

- (a) $\frac{3}{2}$ (b) $\frac{1}{2}$ (c) 2 (d) 1

121. The probability of finding electrons in d_{xy} orbital is :

- (a) along X- and Y-axis (b) along X- and Z-axis
(c) along Y- and Z-axis (d) at an angle of 45° with X-axis

122. Select correct statement :

- (a) The lower the value of $(n-l)$ for an orbital, the higher is its energy.
(b) If two orbitals have the same value of $(n-l)$, the orbital with higher value of n will have lower energy.
(c) The energy of an electron in a multi-electron atom depends on quantum number n only
(d) The energy of an electron in hydrogen atom depends on quantum number n only

123. Read the following statements and choose the correct option.

(I) If the radius of the first Bohr orbit of hydrogen atom is r , then radius of 2nd orbit of Li^2 would be $4r$

(II) For s -orbital electron, the orbital angular momentum is zero

- (a) only I is correct (b) only II is correct (c) both are correct (d) both are incorrect

124. The quantum numbers of four electrons (e_1 to e_4) are given below :

	n	l	m	s		n	l	m	s
e_1	3	0	0	$+1/2$	e_2	4	0	0	$1/2$
e_3	3	2	2	$-1/2$	e_4	3	1	-1	$1/2$

The correct order of decreasing energy of these electrons is :

- (a) $e_4 \ e_3 \ e_2 \ e_1$ (b) $e_2 \ e_3 \ e_4 \ e_1$ (c) $e_3 \ e_2 \ e_4 \ e_1$ (d) $e_1 \ e_4 \ e_2 \ e_3$

125. The energy of an electron of $2p_y$ orbital is :

- (a) greater than $2p_x$ orbital (b) less than $2p_z$ orbital
(c) equal to $2s$ orbital (d) same as that of $2p_x$ and $2p_z$ orbital

126. The number of unpaired valence electrons in an atom of phosphorus is :

- (a) 0 (b) 2 (c) 3 (d) 4

127. Which quantum number defines the orientation of orbital in the space around the nucleus?

- (a) principal quantum number (n) (b) angular momentum quantum number
(c) magnetic quantum number (m_l) (d) spin quantum number (m_s)

128. What is the maximum number of electrons in a subshell that can have the quantum numbers $n = 3$ and $l = 2$?

- (a) 2 (b) 5 (c) 6 (d) 10

- 129.** Which of the following statements about an electron with $m_l = 2$ is incorrect?
 (a) The electron could be in the third shell (b) The electron is in a non-spherical orbital
 (c) The electron may have $m_s = \frac{1}{2}$ (d) The electron is not in a d -orbital
- 130.** Which of the following set of quantum numbers is impossible for an electron?
 (a) $n = 1, l = 0, m_l = 0, m_s = \frac{1}{2}$ (b) $n = 9, l = 7, m_l = 6, m_s = \frac{1}{2}$
 (c) $n = 2, l = 1, m_l = 0, m_s = \frac{1}{2}$ (d) $n = 3, l = 2, m_l = 3, m_s = \frac{1}{2}$
- 131.** In a $3d$ subshell, all the five orbitals are degenerate. What does it mean?
 (a) All the orbitals have the same orientation.
 (b) All the orbitals have the same shape.
 (c) All the orbitals have the same energy.
 (d) All the orbitals are unoccupied.
- 132.** Which of the following subshell can accommodate as many as 10 electrons?
 (a) $2d$ (b) $3d$ (c) $3d_{xy}$ (d) $3d_{z^2}$
- 133.** Which of the following statements is correct for an electron having azimuthal quantum number $l = 2$?
 (a) The electron may be in the lowest energy shell.
 (b) The electron is in a spherical orbital.
 (c) The electron must have spin $m_s = \frac{1}{2}$
 (d) The electron may have a magnetic quantum number $m_l = 1$
- 134.** Which of the following statements is incorrect?
 (a) The concepts of “penetration” and “shielding” are important in deciding the energetic ordering of orbitals in multi-electron atoms
 (b) A wave-function can have positive and negative values
 (c) “Radial nodes” can appear in radial probability distribution functions
 (d) The shape of an orbital is given by the principal quantum number.
- 135.** For an $4p_y$ orbital, there are nodal planeand azimuthal quantum number l
 (a) 1,0 (b) 0,1 (c) 1,1 (d) 2,1
- 136.** Which of the following statement is correct?
 (a) Number of angular nodes $= n - l - 1$
 (b) Number of radial nodes $= l$
 (c) Total number of nodes $= n - 1$
 (d) All of these
- 137.** Give the correct order of initials **T** (True) or **F** (False) for following statements.
 (I) If electron has zero quantum magnetic number, then it must be present in s -orbital.
 (II) In

↑↓

↑↓		↑↓
----	--	----

 orbital diagram, Pauli's exclusion principle is violated.
 (III) Bohr's model can explain spectrum of the hydrogen atom.

(IV) A d -orbital can accommodate maximum 10 electrons only.

- (a) TTFF (b) FFTF
(c) TFFT (d) FFTT

138. The orbital diagram in which both the Pauli's exclusion principle and Hund's rule are violated is :

- (a) $\uparrow\downarrow \uparrow\uparrow \uparrow \square$ (b) $\uparrow\downarrow \uparrow\downarrow \square \uparrow\downarrow$
(c) $\uparrow\uparrow \uparrow \uparrow \uparrow \uparrow$ (d) $\uparrow\downarrow \uparrow\downarrow \uparrow\downarrow \uparrow$

139. It is not possible to explain the Pauli's exclusion principle with the help of this atom.


- (a) B (b) Be (c) C (d) H

140. The subshell that arises after f subshell is called g subshell.

What is the total number of orbitals in the shell in which the g subshell first occur?

- (a) 9 (b) 16 (c) 25 (d) 36

141. The variation of radial probability density $R^2(r)$ as a function of distance r of the electron from the nucleus for $3p$ -orbital :



142. In iron atom, how many electrons have $n = 3$ and $l = 2$?

- (a) 2 (b) 4 (c) 6 (d) 8

143. For similar orbitals having different values of n :

- (a) the most probable distance increases with increase in n
(b) the most probable distance decreases with increase in n
(c) the most probable distance remains constant with increase in n
(d) none of these

144. If n and l are principal and azimuthal quantum numbers respectively, then the expression for calculating the total number of electrons in any energy level is :

- (a) $\sum_{l=0}^{l=n} 2(2l+1)$ (b) $\sum_{l=1}^{l=n} 2(2l+1)$ (c) $\sum_{l=0}^{l=n} (2l+1)$ (d) $\sum_{l=0}^{l=n-1} 2(2l+1)$

145. Maximum number of nodes are present in :

- (a) $5s$ (b) $5p$
(c) $5d$ (d) All have same number of nodes

146. The possible correct set of quantum numbers for the unpaired electron of Cl atom is :

- (a) $2, 0, 0, \frac{1}{2}$ (b) $2, 1, 1, \frac{1}{2}$ (c) $3, 1, 1, \frac{1}{2}$ (d) $3, 0, \frac{1}{2}$


147. The aufbau principle implies that a new electron will enter an orbital for which :

- (a) n has a lower value (b) l has a lower value
(c) $(n-l)$ value is maximum (d) $(n-l)$ value is minimum

148. The orbital diagram in which aufbau principle is violated, is :

- (a) $\uparrow\downarrow \uparrow\uparrow \uparrow \uparrow$ (b) $\uparrow \uparrow\downarrow \uparrow \uparrow$ (c) $\uparrow\downarrow \uparrow \uparrow \uparrow$ (d) $\uparrow\downarrow \uparrow\downarrow \uparrow\downarrow \uparrow$

149. Consider the following six electronic configurations (remaining inner orbitals are completely filled) and mark the incorrect option.



(a) Stability order : II > I > IV > III

(b) Order of spin multiplicity : IV > III = I > II

(c) V does not violate all the three rules of electronic configuration

(d) If VI represents A then A when kept near a magnet, acts as paramagnetic substance.

150. Which of the following set of quantum numbers shows orbital of highest energy?

(a) $n = 4, l = 0, m = 0, s = \frac{1}{2}$

(b) $n = 2, l = 0, m = 0, s = \frac{1}{2}$

(c) $n = 3, l = 1, m = 1, s = \frac{1}{2}$

(d) $n = 3, l = 2, m = 1, s = \frac{1}{2}$

151. A subshell $n = 5, l = 3$ can accommodate :

(a) 10 electrons

(b) 14 electrons

(c) 18 electrons

(d) None of these

152. In H-atom energy of electron is determined by :

(a) only n

(b) n, l

(c) n, l, m

(d) all the four quantum numbers.

153. How many electron(s) in an atom can have $n = 3, l = 2$?

(a) 1

(b) 2

(c) 5

(d) 10

154. How many electrons in an atom can have $n = 4, l = 2, m = 2$ and $s = \frac{1}{2}$?

(a) 1

(b) 2

(c) 5

(d) 10

155. Not considering the electron spin, the degeneracy of first excited state ($n = 2$) of H-atom is:

(a) 2

(b) 3

(c) 4

(d) 8

156. Not considering the electron spin, the degeneracy of second excited state of H-atom is 9, while the degeneracy of the first excited state of H is:

(a) 1

(b) 2

(c) 3

(d) 4

157. Which orbital has only positive values of wave function at all distances from the nucleus :

(a) 1s

(b) 2s

(c) 2p

(d) 3d

158. Four electrons in an atom have the sets of quantum numbers as given below. Which electron is at the highest energy level?

(a) $n = 4, l = 0, m_l = 0, m_s = 1/2$

(b) $n = 3, l = 0, m_l = 0, m_s = 1/2$

(c) $n = 3, l = 2, m_l = 0, m_s = 1/2$

(d) $n = 4, l = 1, m_l = 1, m_s = 1/2$

159. The set of quantum numbers, $n = 3, l = 2, m_l = 0$

(a) describes an electron in a 2s orbital

(b) is not allowed

(c) describes an electron in a 3p orbital


(d) describes one of the five orbitals of same energy

- 160.** The set of quantum numbers, $n = 2, l = 2, m_l = 0$:
- describes an electron in a $2s$ orbital
 - describes one of the five orbitals of a similar type
 - describes an electron in a $2p$ orbital
 - is not allowed
- 161.** Consider the argon atom. For how many electrons does this atom have $m_l = 1$?
- 1
 - 6
 - 4
 - 2
- 162.** An orbital is occupied by an electron with the quantum numbers $n = 4, l = 1$. How many orbitals of this type are found in a multi-electron atom?
- $4p, 3$
 - $4s, 1$
 - $4d, 5$
 - $4p, 6$
- 163.** Which of the following sets of quantum numbers describes the electron which is removed most easily from a potassium atom in its ground state?
- $n = 3, l = 1, m_l = 1, m_s = \frac{1}{2}$
 - $n = 2, l = 1, m_l = 0, m_s = \frac{1}{2}$
 - $n = 4, l = 0, m_l = 1, m_s = \frac{1}{2}$
 - $n = 4, l = 0, m_l = 0, m_s = \frac{1}{2}$
- 164.** The subshell that arises after f is called the g subshell. How many electrons may occupy the g subshell?
- 9
 - 7
 - 5
 - 18
- 165.** Which of the following electron configurations is correct for iron, (atomic number 26)?
- $[\text{Kr}] 4s^1 3d^6$
 - $[\text{Kr}] 4s^1 3d^7$
 - $[\text{Ar}] 4s^2 3d^6$
 - $[\text{Kr}] 4s^2 3d^6$
- 166.** Which of the following electron configurations is correct for copper, (atomic number 29)?
- $[\text{Ar}] 3d^{10} 4s^1$
 - $[\text{Kr}] 3d^9 4s^1$
 - $[\text{Ar}] 3d^9 4s^2$
 - $[\text{Kr}] 3d^{10} 4s^1$
- 167.** The electronic configurations of ${}_{24}\text{Cr}$ and ${}_{29}\text{Cu}$ are abnormal
- due to extra stability of exactly half filled and exactly fully filled sub shells
 - because they belong to d-block
 - both the above
 - None of the above
- 168.** Which of the following representation of excited states of atoms is impossible?
- $1s^1 2s^1$
 - $[\text{Ne}] 3s^2 3p^3 4s^1$
 - $[\text{Ne}] 3s^2 3p^6 4s^1 3d^6$
 - $1s^2 2s^2 2p^7 3s^2$
- 169.** Among the following representations of excited states of atoms which is impossible?
- $1s^1 2s^1$
 - $[\text{Ne}] 3s^2 3p^3 4s^1$
 - $1s^2 2s^2 2p^4 3s^2$
 - $[\text{Ne}] 3s^2 3p^6 4s^3 3d^2$
- 170.** Among the following series of transition metal ions, the one where all metal ions have same $3d$ electronic configuration is:
- $\text{Ti}^{2+}, \text{V}^{3+}, \text{Cr}^{4+}, \text{Mn}^{5+}$
 - $\text{Ti}^{3+}, \text{V}^{2+}, \text{Cr}^{3+}, \text{Mn}^{4+}$
 - $\text{Ti}^{4+}, \text{V}^{4+}, \text{Cr}^{6+}, \text{Mn}^{7+}$
 - $\text{Ti}^{4+}, \text{V}^{3+}, \text{Cr}^{2+}, \text{Mn}^{3+}$
- 171.** Which of the following has the maximum number of unpaired electrons?
- Mn
 - Ti
 - V
 - Al

172. Which of the following orbitals has two spherical nodes?

- (a) $2s$ (b) $4s$ (c) $3d$ (d) $6f$

173. Wave function of an orbital is plotted against the distance from nucleus. The graphical representation is of:



- (a) $1s$ (b) $2s$ (c) $3s$ (d) $2p$

174. The Schrodinger wave equation for hydrogen atom is

$$2s \quad \frac{1}{4\sqrt{2}} \quad \frac{1}{a_0} \quad 2 \quad \frac{r}{a_0} \quad e^{-r/a_0}$$

where a_0 is Bohr's radius. If the radial node in $2s$ be at r_0 , then r_0 would be equal to :

- (a) $\frac{a_0}{2}$ (b) $2a_0$ (c) $\sqrt{2} a_0$ (d) $\frac{a_0}{\sqrt{2}}$

175. The Schrodinger wave equation for hydrogen atom is

$$(\text{radial}) \quad \frac{1}{16\sqrt{4}} \quad \frac{Z}{a_0} \quad \left[\left(1 - \frac{Zr}{2a_0} + \frac{Z^2 r^2}{8a_0^2} - \frac{Z^3 r^3}{12a_0^3} \right) \right] e^{-Zr/4a_0}$$

where a_0 and Z are the constant in which answer can be expressed and $\frac{2Zr}{a_0}$

minimum and maximum position of radial nodes from nucleus arerespectively.

- (a) $\frac{a_0}{Z}, \frac{3a_0}{Z}$ (b) $\frac{a_0}{2Z}, \frac{a_0}{Z}$ (c) $\frac{a_0}{2Z}, \frac{3a_0}{Z}$ (d) $\frac{a_0}{2Z}, \frac{4a_0}{Z}$



Level 2

- Potential energy of electron present in He is :
 (a) $\frac{e^2}{2 \text{ }_0 r}$ (b) $\frac{3e^2}{4 \text{ }_0 r}$ (c) $\frac{2e^2}{4 \text{ }_0 r}$ (d) $\frac{e^2}{4 \text{ }_0 r^2}$
- A single electron in an ion has ionization energy equal to 217.6 eV. What is the total number of neutrons present in one ion of it?
 (a) 2 (b) 4 (c) 5 (d) 9
- For a hypothetical hydrogen like atom, the potential energy of the system is given by $U(r) = \frac{Ke^2}{r^3}$, where r is the distance between the two particles. If Bohr's model of quantization of angular momentum is applicable then velocity of particle is given by :
 (a) $v = \frac{n^2 h^3}{Ke^2 8 \text{ }^3 m^2}$ (b) $v = \frac{n^3 h^3}{8Ke^2 \text{ }^3 m^2}$ (c) $v = \frac{n^3 h^3}{24Ke^2 \text{ }^3 m^2}$ (d) $v = \frac{n^2 h^3}{24Ke^2 \text{ }^3 m^2}$
- A small particle of mass m moves in such a way that P.E. $= \frac{1}{2} mkr^2$, where k is a constant and r is the distance of the particle from origin. Assuming Bohr's model of quantization of angular momentum and circular orbit, r is directly proportional to :
 (a) n^2 (b) n (c) $\sqrt{n}$ (d) none of these
- A beam of specific kind of particles of velocity 2.1×10^7 m/s is scattered by a gold ($Z = 79$) nuclei. Find out specific charge (charge/mass) of this particle if the distance of closest approach is 2.5×10^{-14} m.
 (a) 4.84×10^7 C/kg (b) 4.84×10^7 C/kg (c) 2.42×10^7 C/kg (d) 3×10^{12} C/kg
- What is the angular velocity () of an electron occupying second orbit of Li^{2+} ion?
 (a) $\frac{8 \text{ }^3 me^4}{h^3} K^2$ (b) $\frac{8 \text{ }^3 me^4}{9h^3} K^2$ (c) $\frac{64}{9} \frac{\text{ }^3 me^4}{h^3} K^2$ (d) $\frac{9 \text{ }^3 me^4}{h^3} K^2$
- The ratio of the radius difference between 4th and 3rd orbit of H-atom and that of Li^{2+} ion is :
 (a) 1 : 1 (b) 3 : 1 (c) 3 : 4 (d) 9 : 1
- The velocity of an electron in excited state of H-atom is 1.093×10^6 m/s. What is the circumference of this orbit?
 (a) 3.32×10^{10} m (b) 6.64×10^{10} m (c) 13.30×10^{10} m (d) 13.28×10^8 m
- The angular momentum of an electron in a Bohr's orbit of He is 3.1652×10^{-34} kg-m²/sec. What is the wave number in terms of Rydberg constant (R) of the spectral line emitted when an electron falls from this level to the first excited state. [Use $h = 6.626 \times 10^{-34}$ Js]
 (a) $3R$ (b) $\frac{5R}{9}$ (c) $\frac{3R}{4}$ (d) $\frac{8R}{9}$
- If radiation corresponding to second line of "Balmer series" of Li^{2+} ion, knocked out electron from first excited state of H-atom, then kinetic energy of ejected electron would be:
 (a) 2.55 eV (b) 4.25 eV (c) 11.25 eV (d) 19.55 eV

11. When an electron makes a transition from $(n-1)$ state to n th state, the frequency of emitted radiations is related to n according to $(n-1)$:

(a) $\nu \propto \frac{2cRZ^2}{n^3}$ (b) $\nu \propto \frac{cRZ^2}{n^4}$
 (c) $\nu \propto \frac{cRZ^2}{n^2}$ (d) $\nu \propto \frac{2cRZ^2}{n^2}$

12. In a collection of H-atoms, all the electrons jump from $n=5$ to ground level finally (directly or indirectly), without emitting any line in Balmer series. The number of possible different radiations is :

(a) 10 (b) 8 (c) 7 (d) 6

13. An electron is allowed to move freely in a closed cubic box of length of side 10 cm. The uncertainty in its velocity will be :

(a) $3.35 \times 10^{-4} \text{ m sec}^{-1}$ (b) $5.8 \times 10^{-4} \text{ m sec}^{-1}$
 (c) $4 \times 10^{-5} \text{ m sec}^{-1}$ (d) $4 \times 10^{-6} \text{ m sec}^{-1}$

14. An element undergoes a reaction as shown :

$X + 2e^- \rightarrow X^{2-}$, energy released = 30.87 eV/atom. If the energy released, is used to dissociate 4 g of H_2 molecules, equally into H and H^* , where H^* is excited state of H atoms where the electron travels in orbit whose circumference equal to four times its de Broglie's wavelength. Determine the least moles of X that would be required :

Given : I.E. of H = 13.6 eV/atom, bond energy of H_2 = 4.526 eV/molecule.

(a) 1 (b) 2 (c) 3 (d) 4

15. If the energy of H-atom in the ground state is E , the velocity of photo-electron emitted when a photon having energy E_p strikes a stationary Li^{2+} ion in ground state, is given by :

(a) $v = \sqrt{\frac{2(E_p - E)}{m}}$ (b) $v = \sqrt{\frac{2(E_p - 9E)}{m}}$
 (c) $v = \sqrt{\frac{2(E_p - 9E)}{m}}$ (d) $v = \sqrt{\frac{2(E_p - 3E)}{m}}$

16. At which temperature will the translational kinetic energy of H-atom equal to that for H-atom of first line Lyman transition? (Given $N_A = 6 \times 10^{23}$)

(a) 780 K (b) $1.32 \times 10^5 \text{ K}$
 (c) $7.84 \times 10^4 \text{ K}$ (d) 1000 K

17. For a 3s-orbital, value of ψ is given by following relation :


$$(3s) \psi = \frac{1}{9\sqrt{3}} \left(\frac{1}{a_0} \right)^{3/2} (6 - 6 \left(\frac{r}{a_0} \right)^2) e^{-r/a_0}; \text{ where } \frac{2rZ}{3a_0}$$

What is the maximum radial distance of node from nucleus ?

(a) $\frac{(3 - \sqrt{3}) a_0}{Z}$ (b) $\frac{a_0}{Z}$
 (c) $\frac{3(3 - \sqrt{3}) a_0}{2Z}$ (d) $\frac{2a_0}{Z}$

18. Monochromatic radiation of specific wavelength is incident on H-atoms in ground state. H-atoms absorb energy and emit subsequently radiations of six different wavelength. Find wavelength of incident radiations:
(a) 9.75 nm (b) 50 nm
(c) 85.8 nm (d) 97.25 nm
19. The energy of a I, II and III energy levels of a certain atom are E , $\frac{4E}{3}$ and $2E$ respectively. A photon of wavelength is emitted during a transition from III to I. What will be the wavelength of emission for transition II to I?
(a) $\frac{\lambda}{2}$ (b) λ (c) 2λ (d) 3λ
20. Calculate the minimum and maximum number of electrons which may have magnetic quantum number, $m = 1$ and spin quantum number, $s = \frac{1}{2}$ in chromium (Cr) :
(a) 0, 1 (b) 1, 2 (c) 4, 6 (d) 2, 3
21. An electron in a hydrogen atom in its ground state absorbs 1.5 times as much energy as the minimum required for it to escape from the atom. What is the velocity of the emitted electron?
(a) 1.54×10^6 m/s (b) 1.54×10^8 m/s
(c) 1.54×10^3 m/s (d) 1.54×10^4 m/s
22. In a measurement of quantum efficiency of photosynthesis in green plants, it was found that 10 quanta of red light of wavelength 6850 Å were needed to release one molecule of O_2 . The average energy storage in this process for 1 mole O_2 evolved is 112 kcal. What is the energy conversion efficiency in this experiment?
Given : 1 cal = 4.18 J; $N_A = 6 \times 10^{23}$; $h = 6.63 \times 10^{-34}$ J.s
(a) 23.5 (b) 26.9
(c) 66.34 (d) 73.1
23. A hydrogen like species (atomic number Z) is present in a higher excited state of quantum number n . This excited atom can make a transition to the first excited state by successive emission of two photons of energies 10.20 eV and 17.0 eV respectively. Alternatively, the atom from the same excited state can make a transition to the second excited state by successive emission of two photons of energy 4.25 eV and 5.95 eV respectively. Determine the value of Z .
(a) 1 (b) 2 (c) 3 (d) 4
24. H-atom is exposed to electromagnetic radiation of 1025.6 Å and excited atom gives out induced radiations. What is the minimum wavelength of these induced radiations?
(a) 102.6 nm (b) 12.09 nm
(c) 121.6 nm (d) 810.8 nm
25. If the lowest energy X-rays have 3.055×10^{-8} m, estimate the minimum difference in energy between two Bohr's orbits such that an electronic transition would correspond to the emission of an X-ray. Assuming that the electrons in other shells exert no influence, at what Z (minimum) would a transition from the second energy level to the first result in the emission of an X-ray?
(a) 1 (b) 2
(c) 3 (d) 4

26. An α -particle having kinetic energy 5 MeV falls on a Cu-foil. The shortest distance from the nucleus of Cu to which α -particle reaches is (Atomic no. of Cu = 29, $K = 9 \times 10^9 \text{ Nm}^2/\text{C}^2$)
 (a) $2.35 \times 10^{-13} \text{ m}$ (b) $1.67 \times 10^{-14} \text{ m}$
 (c) $5.98 \times 10^{-15} \text{ m}$ (d) None of these
27. In the graph between $\sqrt{\nu}$ and Z for the Mosley's equation $\sqrt{\nu} = a(Z - b)$, the intercept OX is -1 on $\sqrt{\nu}$ axis.



- What is the frequency when atomic number (Z) is 51 ?
 (a) 50 s^{-1} (b) 100 s^{-1} (c) 2500 s^{-1} (d) None of these
28. Balmer gave an equation for wavelength of visible region of H-spectrum as $\frac{Kn^2}{n^2 - 4}$.
 Where n = principal quantum number of energy level, K = constant in terms of R (Rydberg constant).
 The value of K in terms of R is:
 (a) R (b) $\frac{R}{2}$ (c) $\frac{4}{R}$ (d) $\frac{5}{R}$
29. The energy of separation of an electron in a hydrogen like atom in excited state is 3.4 eV. The de-Broglie wave length (in $\text{\AA}$) associated with the electron is:
 (Given radius of first orbit of H-atom is $0.53 \text{\AA}$)
 (a) 3.33 (b) 6.66 (c) 13.31 (d) None of these
30. If I excitation energy for the H-like (hypothetical) sample is 24 eV, then binding energy in III excited state is:
 (a) 2 eV (b) 3 eV (c) 4 eV (d) 5 eV



Level 3



Passage 1

Werner Heisenberg considered the limits of how precisely we can measure the properties of an electron or other microscopic particle. He determined that there is a fundamental limit to how closely we can measure both position and momentum. The more accurately we measure the momentum of a particle, the less accurately we can determine its position. The converse is also true. This is summed up in what we now call the Heisenberg uncertainty principle.

The equation is $\Delta x (\Delta mv) \geq \frac{h}{4}$

The uncertainty in the position or in the momentum of a macroscopic object like a baseball is too small to observe. However, the mass of microscopic object such as an electron is small enough for the uncertainty to be relatively large and significant.

- If the uncertainties in position and momentum are equal, the uncertainty in the velocity is :
 (a) $\sqrt{\frac{h}{m}}$ (b) $\sqrt{\frac{h}{2m}}$ (c) $\frac{1}{2m} \sqrt{\frac{h}{m}}$ (d) none of these
- If the uncertainty in velocity and position is same, then the uncertainty in momentum will be :
 (a) $\sqrt{\frac{hm}{4}}$ (b) $m \sqrt{\frac{h}{4m}}$ (c) $\sqrt{\frac{h}{4m}}$ (d) $\frac{1}{m} \sqrt{\frac{h}{4m}}$
- What would be the minimum uncertainty in de-Broglie wavelength of a moving electron accelerated by potential difference of 6 volt and whose uncertainty in position is $\frac{7}{22}$ nm?
 (a) 6.25 Å (b) 6 Å (c) 0.625 Å (d) 0.3125 Å



Passage 2

One of the fundamental laws of physics is that matter is most stable with the lowest possible energy. Thus, the electron in a hydrogen atom usually moves in the $n = 1$ orbit, the orbit in which it has the lowest energy. When the electron is in this lowest energy orbit, the atom is said to be in its ground electronic state. If the atom receives energy from an outside source, it is possible for the electron to move to an orbit with a higher n value, in which case the atom is in an excited state with a higher energy.

The law of conservation of energy says that we cannot create or destroy energy. Thus, if a certain amount of external energy is required to excite an electron from one energy level to another, then that same amount of energy will be liberated when the electron returns to its initial state.

Lyman series is observed when the electron returns to the lowest orbit while Balmer series is formed when the electron returns to second orbit. Similarly, Paschen, Brackett and Pfund series are formed when electrons returns to the third, fourth and fifth orbits from higher energy orbits respectively.

When electrons return from n_2 to n_1 state, the number of lines in the spectrum will equal to

$$\frac{(n_2 - n_1)(n_2 - n_1 - 1)}{2}$$

If the electron comes back from energy level having energy E_2 to energy level having energy E_1 , then the difference may be expressed in terms of energy of photon as :

$$E_2 - E_1 = E = \frac{hc}{\lambda}$$

Since, h and c are constants, E corresponds to definite energy. Thus, each transition from one energy level to another will produce a radiation of definite wavelength. This is actually observed as a line in the spectrum of hydrogen atom.

Wave number of a spectral line is given by the formula

$$\bar{\nu} = R \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

where R is a Rydberg's constant ($R = 1.1 \times 10^7 \text{ m}^{-1}$)

- If the wavelength of series limit of Lyman series for He^+ ion is $x \text{ \AA}$, then what will be the wavelength of series limit of Balmer series for Li^{2+} ion?
 (a) $\frac{9x}{4} \text{ \AA}$ (b) $\frac{16x}{9} \text{ \AA}$ (c) $\frac{5x}{4} \text{ \AA}$ (d) $\frac{4x}{7} \text{ \AA}$
- The emission spectra is observed by the consequence of transition of electrons from higher energy state to ground state of He^+ ion. Six different photons are observed during the emission spectra, then what will be the minimum wavelength during the transition?
 (a) $\frac{4}{27R_H}$ (b) $\frac{4}{15R_H}$ (c) $\frac{15}{16R_H}$ (d) $\frac{16}{15R_H}$
- What transition in the hydrogen spectrum would have the same wavelength as Balmer transition, $n = 4$ to $n = 2$ in the He^+ spectrum?
 (a) $n = 3$ to $n = 1$ (b) $n = 3$ to $n = 2$ (c) $n = 4$ to $n = 1$ (d) $n = 2$ to $n = 1$
- An electron in H-atom in M-shell on de-excitation to ground state gives maximum spectrum lines.
 (a) 10 (b) 6 (c) 2 (d) 1



Passage 3

If hydrogen atoms (in the ground state) are passed through an homogeneous magnetic field, the beam is split into two parts. This interaction with the magnetic field shows that the atoms must have magnetic moment. However, the moment cannot be due to the orbital angular momentum since $l = 0$. Hence one must assume existence of intrinsic angular momentum, which as the experiment shows, has only two permitted orientations.

Spin of the electron produce angular momentum equal to $S = \sqrt{s(s+1)} \frac{h}{2\pi}$ where $S = \frac{1}{2}$.

Total spin of an atom $\frac{n}{2}$ or $\frac{n}{2}$

where n is the number of unpaired electrons.

The substance which contain species with unpaired electrons in their orbitals behave as paramagnetic substances. The paramagnetism is expressed in terms of magnetic moment.

The magnetic moment of an atom

$$\mu_s = \sqrt{s(s+1)} \frac{eh}{2\pi mc} = \sqrt{\frac{n}{2} \left(\frac{n}{2} + 1\right)} \frac{eh}{2\pi mc} = \frac{n}{2} \mu_B$$

$$\mu_s = \sqrt{n(n+2)} \text{ B.M.}$$

n = number of unpaired electrons

1. B.M. (Bohr magneton) $\frac{eh}{4\pi mc}$


If magnetic moment is zero the substances is diamagnetic.

- Which of the following ion has lowest magnetic moment ?
 (a) Fe^{2+} (b) Mn^{2+} (c) Cr^{3+} (d) V^{3+}
- If an ion of ^{25}Mn has a magnetic moment of 3.873 B.M. Then oxidation state of Mn in ion is :
 (a) 2 (b) 3 (c) 4 (d) 5




Passage 4

Ozone in the upper atmosphere absorbs ultraviolet radiation which induces the following chemical reaction



O_2 produced in the above photochemical dissociation undergoes further dissociation into one normal oxygen atom (O) and more energetic oxygen atom O^* .



(O^*) has 1 eV more energy than (O) and normal dissociation energy of O_2 is 480 kJ mol^{-1} .


[1 eV/ Photon = 96 kJ mol^{-1}]

- What is the maximum wavelength effective for the photochemical dissociation of O_2 molecule
 (a) $2440 \text{ \AA}$ (b) $2066.67 \text{ \AA}$ (c) $1000 \text{ \AA}$ (d) $155 \text{ \AA}$
- If dissociation of O_3 into O_2 and O requires 400 kJ mol^{-1} and O_2 produced in this reaction is further dissociated to O and O^* then the total energy required for the dissociation of O_3 into O and O^* is:
 (a) 1168 kJ/mol (b) 976 kJ/mol (c) 880 kJ/mol (d) None of these




Passage 5

The existence of negatively charged particle in an atom was shown by J.J. Thomson as a result of the studies of the passage of electricity through gases at extremely low pressures known as discharge tube experiments. When a high voltage of the order of 10,000 volts or more was impressed across the electrodes, some sort of invisible rays moved from the negative electrode to the positive electrode these rays are called as cathode rays.



Cathode rays experiment



Anode rays experiment

Cathode rays travel in straight path in absence of electrical and magnetic field. Cathode rays consist of material part and charged particles? Cathode rays produce **X-rays** and light is emitted when they strike on ZnS screen. Cathode rays penetrate through thin sheets of aluminium and other metals. They affect the photographic plate and passes heating effect when they strike on metal foil. The ratio of charge to mass i.e. charge/mass is same for all the cathode rays irrespective of the gas used in the tube.

The existence of positively charged particles in an atom was shown by E. Goldstein. He repeated the same discharge tube experiments by using a perforated cathode. It was observed that when a high potential difference was applied between the electrodes, not only cathode rays were produced but also a new type of rays were produced simultaneously from anode moving towards cathode and passed through the holes or canal of the cathode. These termed as canal rays or anode rays.

These rays travel in straight lines and consist of positively charged particles. These rays have kinetic energy and produces heating effect also. The e/m ratio of these rays is smaller than that of electrons. Unlike cathode rays, their e/m value is is dependent upon the nature of the gas taken in the tube. These rays produce flashes of light on ZnS screen and can pass through thin metal foils. They can produce physical and chemical changes and are capable to produce ionisation in gases.

- For cathode rays the value of e/m :
 - is independent of the nature of the cathode and the gas filled in the discharge tube
 - is constant
 - is -1.7588×10^8 coulombs/g
 - all of the above are correct
- Which is not true with respect to cathode rays ?
 - A stream of electrons
 - Charged particles
 - Move with same speed as that of light
 - Can be deflected by the electric field
- Select the incorrect statement.
 - Cathode rays has charge only and no mass
 - Anode rays are deflected by electrical and magnetic field
 - Canal rays is named for beam of positive charged particle
 - Anode rays particles carrying positive charge

One or More Answers is/are Correct







- Select the correct statement(s) :
 - The phenomena of diffraction of light can only be explained by assuming that light behaves as waves
 - de-Broglie postulate the dual character existed with matter
 - In his atomic model Bohr considered electron as a particle
 - Wave nature of electrons was proved when diffraction rings were observed photographically when a stream of protons was passed through a metal foil
- The angular momentum of electron can have the value(s) :
 - $0.5 \frac{h}{2\pi}$
 - $\frac{h}{2\pi}$
 - $\frac{h}{0.5}$
 - $2.5 \frac{h}{2\pi}$
- Select incorrect statement(s) :
 - Only three quantum numbers n , l and m are needed to define an orbital
 - Four quantum numbers are needed for complete description of an electron
 - Two quantum numbers n and l are needed to identify subshell and shape of orbital
 - Splitting of spectrum lines in presence of electric field is known as Zeeman effect
- Select the correct statement(s):
 - An electron near the nucleus is attracted by the nucleus and has a low potential energy
 - According to Bohr's theory, an electron continuously radiate energy if it stays in one orbit
 - Bohr's model could not explain the spectra of multielectron atoms
 - Bohr's model was the first atomic model based on quantisation of energy
- Choose the correct statement(s) :
 - The shape of an atomic orbital depends upon azimuthal quantum number
 - The orientation of an atomic orbital depends upon the magnetic quantum number
 - The energy of an electron in an atomic orbital of multi-electron atom depends upon principal quantum number only
 - The number of degenerate atomic orbitals of one type depends upon the value of azimuthal quantum number


6. For radial probability curves, which of the following is/are correct?
 - (a) The number of maxima in $2s$ orbital are two
 - (b) The number of spherical or radial nodes is equal to $n - l - 1$
 - (c) The number of angular nodes are ' l '
 - (d) $3d_z^2$ has 3 angular nodes
7. Select the correct statement(s):
 - (a) Lower value of quantum number l indicates that there is a higher probability of finding the $3s$ electron close to the nucleus than those of $3p$ and $3d$ electrons
 - (b) Energy of $3s$ orbital is less than for the $3p$ and $3d$ orbitals
 - (c) At the node, the value of the radial function changes from positive to negative
 - (d) The radial function depends upon the quantum numbers n and l
8. Choose the incorrect statement(s):
 - (a) For a particular orbital in hydrogen atom, the wave function may have negative value
 - (b) Radial probability distribution function may have zero value but can never have negative value
 - (c) $3d_{x^2-y^2}$ orbital has two angular nodes and one radial node
 - (d) yz and xz planes are nodal planes for d_{xy} orbital
9. Select the correct statement(s) :
 - (a) Heisenberg's principle is applicable to stationary electron
 - (b) Pauli's exclusion principle is not applicable to photons
 - (c) For an electron the product of velocity and principal quantum number will be independent to principal quantum number
 - (d) Quantum numbers l and m determine the value of angular wave function
10. Choose the correct statements among the following :
 - (a) A node is a point in space where the wave-function has zero amplitude
 - (b) The number of maxima (peaks) in radial probability distribution function is $(n - l)$
 - (c) Radial probability density is $4\pi r^2 R_{n,l}^2(r)$
 - (d) $R_{n,l}^2$ represents probability of finding electron
11. Select the correct statement(s) regarding $3p_y$ orbital :
 - (a) Total number of nodes are 2
 - (b) Number of maxima in the curve $4\pi r^2 R^2$ vs r are two
 - (c) Quantum number n , l and m for an orbital may be 3, 1, -1 respectively
 - (d) The magnetic quantum number may have a positive value
12. Select the correct statement(s) :
 - (a) In wave mechanical model, the energy of electron in the orbital remains constant
 - (b) d_{xy} orbital is lies in yz plane
 - (c) Nodal planes are yz and xy in $d_{x^2-y^2}$ orbital
 - (d) Rest mass of photon is zero and increases with its velocity
13. Hydrogen has :

(a) half filled subshell	(b) half filled shell
(c) one electron in valence shell	(d) half filled orbital

14. Select incorrect statement(s) .
 (a) If the value of $l = 0$, the electron distribution is spherical
 (b) The shape of the orbital is given by magnetic quantum number
 (c) Angular momentum of $1s, 2s, 3s$ orbit electrons are equal
 (d) In an atom, all the electrons travel with the same velocity
15. The radial distribution functions $[P(r)]$ is used to determine the most probable radius, which is used to find the electron in a given orbital. $\frac{dP(r)}{dr}$ for $1s$ -orbital of hydrogen like atom having atomic number Z , is $\frac{dP}{dr} = \frac{4Z^3}{a_0^3} 2r \frac{2Zr^2}{a_0} e^{-2Zr/a_0}$:
 Then which of the following statements is/are correct?
 (a) At the point of maximum value of radial distribution function $\frac{dP(r)}{dr} = 0$; one antinode is present
 (b) Most probable radius of Li^{2+} is $\frac{a_0}{3}$ pm
 (c) Most probable radius of He^+ is $\frac{a_0}{2}$ pm
 (d) Most probable radius of hydrogen atom is a_0 pm
16. Select the correct statement(s) :
 (a) An orbital with $l = 0$ is symmetrical about the nucleus
 (b) An orbital with $l = 1$ is spherically symmetrical about the nucleus
 (c) $3d_{z^2}$ is spherically symmetrical about the z -axis
 (d) All are correct
17. Select the correct statement(s) :
 (a) Radial function $[R(r)]$ is a part of wave function which depends upon quantum number n only
 (b) Angular function depends only on the direction, and is independent to the distance from the nucleus
 (c) $\psi^2(r, \theta, \phi)$ is the probability density of finding the electron at a particular point in space
 (d) Radial distribution function ($4\pi r^2 R^2$) gives the probability of the electron being present at a distance r from the nucleus
18. Which is/are correct graph?



19. Select the correct curve(s) :
 If v = Velocity of electron in Bohr's orbit
 r = Radius of electron in Bohr's orbit
 P.E. = Potential energy of electron in Bohr's orbit
 K.E. = Kinetic energy of electron in Bohr's orbit



20. Select the correct set (s) of quantum numbers
 (a) $n = 3, l = 0, m_l = 1$ (b) $n = 3, l = 3, m_l = 2$
 (c) $n = 3, l = 2, m_l = 2$ (d) $n = 3, l = 1, m_l = 0$
21. Which is/are correct statement?
 (a) Number of subshell present in M-shell 3
 (b) Number of orbitals present in N-shell 16
 (c) Cu ($z = 29$) is paramagnetic
 (d) Zeeman effect explains splitting of spectral lines in magnetic field.
22. In a sample of H-atoms electrons are de-excited from 4th excited state to ground state. Which is/are correct statement?
 (a) No line observed in P-fund series.
 (b) Total ten lines observed in spectrum.
 (c) 4 line in UV-region and 3 line in visible region observed.
 (d) One line observed in Brackett series.

Match the Column

Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.	Column-I	Column-II
	(A) Electron	(P) Negative charge
	(B) Proton	(Q) Positive charge
	(C) Neutron	(R) $1.6 \times 10^{-19} \text{ C}$
	(D) Positron	(S) Chargeless
2.	Column-I	Column-II
	(A) Thomson model of atom	(P) Electrons are present in extra nuclear region
	(B) Rutherford model of atom	(Q) Electron in the atom is described as wave
	(C) Bohr model of atom	(R) Positive charge is accumulated in the nucleus
	(D) Schrodinger model of hydrogen atom	(S) Uniform sphere of positive charge with embedded electrons

3.	Column-I
(A)	Atomic theory of matter
(B)	Quantization of charge
(C)	Quantization of electronic energy level
(D)	Size of nucleus

Column-II
(P) Rutherford scattering experiment
(Q) Muliken's oil drop experiment
(R) Atomic spectra
(S) Law of multiple proportions

4.	Column-I
(A)	$\frac{\text{K.E.}}{\text{P.E.}}$
(B)	$\text{P.E.} - 2 \text{ K.E.}$
(C)	$\frac{\text{P.E.}}{\text{T.E.}}$
(D)	$\frac{\text{K.E.}}{\text{T.E.}}$

Column-II
(P) 2
(Q) $\frac{1}{2}$
(R) -1
(S) 0

5.	Column-I
(A)	Lyman series
(B)	Humphery series
(C)	Paschen series
(D)	Balmer series

Column-II
(P) Visible region
(Q) Ultraviolet region
(R) Infrared region
(S) Far infrared region

6. In case of hydrogen spectrum wave number is given by

$$= R_H \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \quad \text{where } n_1 < n_2$$

Column-I
(A) Lyman series
(B) Balmer series
(C) Pfund series
(D) Brackett series

Column-II
(P) $n_2 = 2$
(Q) $n_2 = 3$
(R) $n_2 = 6$
(S) $n_2 = 5$

7.	Column-I (Shell)
(A)	2nd
(B)	3rd
(C)	4th
(D)	1st

Column-II (Value of l)
(P) 1
(Q) 2
(R) 3
(S) 0

8. In Bohr's model, for a unielectronic atom, following symbols are used

$r_{n,Z}$ Radius of n^{th} orbit with atomic number Z ;

$U_{n,Z}$ Potential energy of electron ; $K_{n,Z}$ Kinetic energy of electron ;

$V_{n,Z}$ Velocity of electron ; $T_{n,Z}$ Time period of revolution

Column-I

(A) $U_{1,2} : K_{1,1}$

(B) $r_{2,1} : r_{1,2}$

(C) $V_{1,3} : V_{3,1}$

(D) $T_{1,2} : T_{2,2}$

Column-II

(P) $1 : 8$

(Q) $-8 : 1$

(R) $9 : 1$

(S) $8 : 1$

9.

Column-I

(A) The radial node of 5s atomic orbital is

(B) The angular node of $3d_{yz}$ atomic orbital is

(C) The sum of angular node and radial node of $4d_{xy}$ atomic orbital

(D) The angular node of 3p atomic orbital is

Column-II

(P) 1

(Q) 4

(R) 2

(S) 3

10.

Column-I

(A) The d-orbital which has two angular nodes

(B) The d-orbital with two nodal surfaces form cones

(C) The orbital without angular node

(D) The orbital which has three angular nodes

Column-II

(P) $3d_{x^2 - y^2}$

(Q) $3d_{z^2}$

(R) $4f$

(S) $3s$

11.

Column-I

(A) Orbital angular momentum of an electron

(B) Angular momentum of an electron in an orbit

(C) Spin angular momentum of an electron

(D) Magnetic moment of atom

Column-II

(P) $\sqrt{s(s-1)} \frac{h}{2}$

(Q) $\sqrt{n(n-2)}$

(R) $\frac{nh}{2}$

(S) $\sqrt{l(l-1)} \frac{h}{2}$

12.

Column-I

- (A) Number of orbitals in the n^{th} shell
 (B) Maximum number of electrons in a subshell
 (C) Number of subshells in n^{th} shell
 (D) Number of orbitals in a subshell

Column-II

- (P) $2(2l - 1)$
 (Q) n
 (R) $2l - 1$
 (S) n^2

13.

Column-I



- (A) $2s$
 (B) $2p_z$
 (C) $4d_{x^2 - y^2}$
 (D) $4d_{z^2}$

Column-II

- (P) $n = 4, l = 2, m = 0$
 (Q) $n = 4, l = 2, m = 2$ or -2
 (R) $n = 2, l = 1, m = 0$
 (S) $n = 2, l = 0, m = 0$

14.

Column-I



- (C) Angular probability depends upon θ and ϕ
 (D) Atleast one angular node is present

Column-II


- (P) $4s$
 (Q) $5p_y$
 (R) $3s$
 (S) $6d_{xy}$

15.

Column-I


- (A) $3s$

Column-II




(B) $3p$ (C) $3d$ (D) $2p$


(Q)



(R)



(S)



Assertion-Reason Type Questions

Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

- (A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** The angular momentum of d -orbitals is $\sqrt{6} \frac{h}{2}$.

STATEMENT-2 : Angular momentum of electron in orbit is $mvr \frac{nh}{2}$.

2. **STATEMENT-1 :** Angular momentum of the electron in the orbit which has four subshell is $\frac{2h}{2}$.

STATEMENT-2 : Angular momentum of electron is quantized.

3. **STATEMENT-1 :** Line emission spectra useful in the study of atomic structure.

STATEMENT-2 : Each element has a unique line emission spectrum.

4. **STATEMENT-1 :** Emitted radiation will fall in visible range when an electron jump from $n = 4$ to $n = 2$ in H-atom.

STATEMENT-2 : Balmer series radiations belong to visible range for hydrogen atom only.

5. **STATEMENT-1 :** Half-filled and fully-filled degenerate orbitals are more stable.
STATEMENT-2 : Extra stability is due to the symmetrical distribution of electrons and exchange energy.
6. **STATEMENT-1 :** The ground state configuration of Cr is $3d^5 4s^1$.
STATEMENT-2 : A set of half-filled orbitals containing one electrons each with their spin parallel provides extra stability.
7. **STATEMENT-1 :** The ground state electronic configuration of nitrogen is
- $\uparrow\downarrow$

$\uparrow\downarrow$

$\uparrow$

$\uparrow$

$\uparrow$
- STATEMENT-2 :** Electrons are filled in orbitals as per aufbau principle, Hund's rule of maximum spin multiplicity and Pauli's principle.
8. **STATEMENT-1 :** An orbital cannot have more than two electrons and they must have opposite spins.
STATEMENT-2 : No two electrons in an atom can have same set of all the four quantum numbers as per Pauli's exclusion principle.
9. **STATEMENT-1 :** Orbital having xz plane as node may be $3d_{xy}$.
STATEMENT-2 : $3d_{xy}$ has zero radial node.
10. **STATEMENT-1 :** The kinetic energy of photo-electrons increases with increase in frequency of incident light where $\nu > \nu_0$.
STATEMENT-2 : Whenever intensity of light is increased the number of photo-electron ejected always increases.
11. **STATEMENT-1 :** Cu^{2+} is a coloured ion.
STATEMENT-2 : Every ion with unpaired electron is coloured.
12. **STATEMENT-1 :** For $n = 3$, l may be 0, 1 and 2 and m may be 0; 0 - 1; 0 - 1 and 2
STATEMENT-2 : For each value of n , there are 0 to $(n - 1)$ possible values of l ; and for each value of l , there are 0 to l values of m .

Subjective Problems



- Given r_{n-1}, r_n, r_{n+1} are Bohr radius for hydrogen atom in $n^{\text{th}}, (n-1)^{\text{th}}$ and $(n+1)^{\text{th}}$ shell respectively. Calculate the value of n .
- The energy of separation of an electron is 30.6 eV moving in an orbit of Li^{2+} . Find out the number of waves made by the electron in one complete revolution in the orbit.
- Calculate the number of waves made by a Bohr electron in one complete revolution in n^{th} orbit of H-atom, if ratio of de-Broglie wavelength associated with electron moving in n^{th} orbit and 2^{nd} orbit is 1.5.
- A certain dye absorbs lights of 400 nm and then fluorescence light of wavelength 500 nm. Assuming that under given condition 40% of the absorbed energy is re-emitted as fluorescence, calculate the ratio of quanta absorbed to number of quanta emitted out.
- A photon of energy 4.5 eV strikes on a metal surface of work function 3.0 eV. If uncertainty in position is $\frac{25}{4} \text{ \AA}$, find the uncertainty in measurement of de-Broglie wavelength (in $\text{\AA}$).

6. Find out the difference in number of angular nodes and number of radial nodes in the orbital to which last electron of chromium present.
7. What is the total number of radial and angular nodes present in 5f orbital ?
8. Infrared lamps are used in restaurants to keep the food warm. The infrared radiation is strongly absorbed by water, raising its temperature and that of the food. If the wavelength of infrared radiation is assumed to be 1500 nm, and the number of quanta of infrared radiation produced per second by an infrared lamp (that consumes energy at the rate of 100 W and is 12% efficient only) is $(x \cdot 10^{19})$, then the value of x is :

(Given : $h = 6.625 \cdot 10^{-34}$ J-s)

9. When an electron makes transition from $(n - 1)$ state to n state the wavelength of emitted radiations is related to $n(n - 1)$ according to n^x .

What is the value of x ?

10. For 3s orbital of hydrogen atom, the normalised wave function is

$$_{3s} \frac{1}{81\sqrt{3}} \frac{1}{a_0} e^{-r/a_0} \left(27 - \frac{18r}{a_0} + \frac{2r^2}{a_0^2} \right) e^{-r/3a_0}$$

If distance between the radial nodes is d , calculate the value of $\frac{d}{1.73 a_0}$.

11. Find the separation between two electrons (in Å) in vacuum, if electrostatic potential energy between these electrons is $7.67 \cdot 10^{-19}$ J.

[Given: $e = 1.6 \cdot 10^{-19}$ C ; $\frac{1}{4\pi\epsilon_0} = 8.85 \cdot 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{m}^{-1}$, $\pi = 3.14$]

12. An α -particle moving with velocity $\frac{1}{30}$ th times of velocity of light. If uncertainty in position is $\frac{3.31}{\text{pm}}$, then minimum uncertainty in kinetic energy is $y \cdot 10^{-16}$ J. Calculate the value of y.

13. In a sample of excited hydrogen atoms electrons make transition from $n = 2$ to $n = 1$. Emitted quanta strike on a metal of work function (ϕ) 4.2 eV.

Calculate the wavelength (in Å) associated with ejected electrons having maximum kinetic energy.

14. For 1s orbital of Hydrogen atom radial wave function is given as:

$$R(r) = \frac{1}{\sqrt{\pi}} \frac{1}{a_0} e^{-r/a_0} \quad (\text{where } a_0 = 0.529 \text{ Å})$$

The ratio of radial probability density of finding electron at $r = a_0$ to the radial probability density of finding electron at the nucleus is given as $(x \cdot e^{-y})$. Calculate the value of $(x - y)$.

15. Calculate the value of A.

$A = \frac{E_{1,2}}{2E_{2,1}}$ where $E_{n,Z}$ = Energy of electron in n^{th} orbit, Z = atomic number of hydrogen like species.

ANSWERS

Level 1

1. (c)	2. (c)	3. (a)	4. (d)	5. (c)	6. (c)	7. (b)	8. (c)	9. (d)	10. (d)
11. (a)	12. (b)	13. (b)	14. (a)	15. (a)	16. (d)	17. (b)	18. (c)	19. (c)	20. (b)
21. (d)	22. (b)	23. (d)	24. (d)	25. (c)	26. (a)	27. (d)	28. (d)	29. (d)	30. (d)
31. (b)	32. (c)	33. (d)	34. (b)	35. (b)	36. (a)	37. (b)	38. (c)	39. (c)	40. (d)
41. (d)	42. (d)	43. (b)	44. (d)	45. (d)	46. (b)	47. (b)	48. (a)	49. (b)	50. (c)
51. (c)	52. (c)	53. (b)	54. (a)	55. (c)	56. (a)	57. (d)	58. (b)	59. (b)	60. (d)
61. (a)	62. (c)	63. (b)	64. (b)	65. (c)	66. (c)	67. (c)	68. (c)	69. (c)	70. (a)
71. (d)	72. (a)	73. (a)	74. (c)	75. (b)	76. (b)	77. (d)	78. (b)	79. (b)	80. (d)
81. (c)	82. (b)	83. (a)	84. (c)	85. (d)	86. (b)	87. (b)	88. (b)	89. (d)	90. (a)
91. (c)	92. (c)	93. (c)	94. (a)	95. (c)	96. (d)	97. (b)	98. (a)	99. (a)	100. (d)
101. (c)	102. (b)	103. (b)	104. (c)	105. (d)	106. (c)	107. (a)	108. (d)	109. (b)	110. (c)
111. (b)	112. (c)	113. (d)	114. (b)	115. (d)	116. (d)	117. (b)	118. (a)	119. (d)	120. (c)
121. (d)	122. (d)	123. (b)	124. (c)	125. (d)	126. (c)	127. (c)	128. (d)	129. (d)	130. (d)
131. (c)	132. (b)	133. (d)	134. (d)	135. (c)	136. (c)	137. (b)	138. (a)	139. (d)	140. (c)
141. (c)	142. (c)	143. (a)	144. (d)	145. (d)	146. (c)	147. (d)	148. (b)	149. (a)	150. (d)
151. (b)	152. (a)	153. (d)	154. (a)	155. (c)	156. (a)	157. (a)	158. (d)	159. (d)	160. (d)
161. (c)	162. (a)	163. (d)	164. (d)	165. (c)	166. (a)	167. (a)	168. (d)	169. (d)	170. (a)
171. (a)	172. (d)	173. (c)	174. (b)	175. (c)					

Level 2

1. (c)	2. (c)	3. (c)	4. (c)	5. (a)	6. (d)	7. (b)	8. (c)	9. (b)	10. (d)
11. (a)	12. (d)	13. (a)	14. (b)	15. (c)	16. (c)	17. (c)	18. (d)	19. (d)	20. (d)
21. (a)	22. (b)	23. (c)	24. (a)	25. (b)	26. (b)	27. (c)	28. (c)	29. (b)	30. (a)



Level 3

Passage-1	1. (c)	2. (a)	3. (c)	
Passage-2	1. (b)	2. (b)	3. (d)	4. (c)
Passage-3	1. (d)	2. (c)		
Passage-4	1. (b)	2. (b)		
Passage-5	1. (d)	2. (c)	3. (a)	

One or More Answers is/are Correct

- | | | | | | |
|---------------|-------------|---------------|---------------|---------------|------------|
| 1. (a,b,c) | 2. (a,b,c) | 3. (d) | 4. (a,c,d) | 5. (a,b,d) | 6. (a,b,c) |
| 7. (a,b,c,d) | 8. (c) | 9. (b,c,d) | 10. (a,b,d) | 11. (a,b,c,d) | 12. (a,d) |
| 13. (a,b,c,d) | 14. (b,c,d) | 15. (a,b,c,d) | 16. (a,c) | 17. (b,c,d) | 18. (b,c) |
| 19. (a,b,c,d) | 20. (c,d) | 21. (a,b,d) | 22. (a,b,c,d) | | |

Match the Column

- | | | | |
|------------------|------------|---------------|-----------|
| 1. A P, R; | B Q, R; | C S; | D Q, R |
| 2. A S; | B P, R; | C P, R; | D P, Q, R |
| 3. A S; | B Q; | C R; | D P |
| 4. A Q; | B S; | C P; | D R |
| 5. A Q; | B S; | C R; | D P |
| 6. A P, Q, R, S; | B Q, R, S; | C R; | D R, S |
| 7. A P, S; | B P, Q, S; | C P, Q, R, S; | D S |
| 8. A Q; | B S; | C R; | D P |
| 9. A Q; | B R; | C S; | D P |
| 10. A P; Q | B Q; | C S; | D R |
| 11. A S; | B R; | C P; | D Q |
| 12. A S; | B P; | C Q; | D R |
| 13. A S; | B R; | C Q; | D P |
| 14. A P; | B P, Q, S; | C Q, S; | D Q, S |
| 15. A Q; | B R; | C S; | D P |

Assertion-Reason Type Questions

1. (B) 2. (B) 3. (A) 4. (A) 5. (A) 6. (A) 7. (A) 8. (A) 9. (B) 10. (C)
11. (C) 12. (A)

Subjective Problems

1.	2	2.	2	3.	3	4.	2	5.	4	6.	2	7.	4	8.	9	9.	3	10.	3
11.	3	12.	5	13.	5	14.	3	15.	8										

Hints and Solutions

Level 1

1. (c) Isodiaphers have same value of $(n - p)$ or $(A - 2Z)$.
2. (c) Travel perpendicular to both electric and magnetic fields.
3. (a) $\frac{(e/m)_p}{(e/m)} = \frac{e/m_p}{2e/4m_p} = \frac{2}{1}$
4. (d) Charge/mass for $n = 0$, for $\frac{2}{4}$,
for $p = \frac{1}{1}$, for $e = \frac{1}{1/1837}$
5. (c) $\frac{m}{e} = 1.5 \times 10^8 \times 10^3 \times \frac{m}{1.6 \times 10^{19}}$
 $m = 2.4 \times 10^{24} \text{ g}$
9. (d) $r = \frac{n^2}{Z}$; $\frac{r_1}{r_2} = \frac{(1)^2}{(3)^2}$
 $r_2 = 9r_1$
12. (b) $v = \frac{Z}{n}$
13. (b) $r_{\text{atom}} = 10^{10} \text{ m}$
 $r_{\text{nucleus}} = 10^5 \text{ m}$
 $\frac{V_{\text{Atom}}}{V_{\text{Nucleus}}} = \frac{4/3 (10^{10})^3}{4/3 (10^{15})^3} = 10^{15}$
14. (a) Change in P.E. $\frac{2x}{4} (2x) = \frac{3}{2} x$
15. (a) $E_n = \frac{1}{2} \text{ P.E.} = \frac{6.8}{2} = 3.4 \text{ eV}$
 $\therefore E_n = \frac{13.6}{n^2} = 3.4$
 $n = 2$ or first excited state
16. (d) Energy of N -shell $\frac{13.6 (4)^2}{(4)^2} = 13.6 \text{ eV}$
P.E. $= 2 E = 2 \times 13.6 = -27.2 \text{ eV}$
17. (b) Total energy of third shell $\frac{13.6}{3^2}$
 $= 1.51 \text{ eV}$
K.E. = Total energy = 1.51 eV
P.E. $= 2 \times \text{T.E.} = 3.02 \text{ eV}$
18. (c) $r = 0.529 \frac{n^2}{Z} \text{ \AA}$;
 $r_4 = r_3 = 0.529 \times \frac{16}{2} \times \frac{9}{2} \text{ \AA} = 1.851 \times 10^{10} \text{ m}$
19. (c) $r_1 = 0.529 \text{ \AA}$; $r_{4(X)} = r_1 \frac{n^2}{Z}$;
 $r_{4(X)} = \frac{0.529 (4)^2}{Z}$; $Z = 16$
20. (b) $\frac{v_1}{v_2} = \frac{n_2}{n_1} \times \frac{5}{3}$
 2
21. (d) $\frac{r_1}{r_2} = \frac{n_1}{n_2} \times \frac{4}{9}$
22. (b) r_1 of H-atom $= 0.529 \text{ \AA}$
 r_n (H like atom) $= \frac{n^2}{Z} r_1$ (H-atom)
 r_n of $\text{Be}^3 = \frac{n^2}{Z} r_1$ (H-atom)
 $= 0.529 \text{ \AA} (Z = 4 \text{ for } \text{Be}^3)$
 $\frac{n^2}{Z} = 0.529 = 0.529 n^2 Z$
 $n^2 = 4 \quad n = 2$
23. (d) $v = 2.188 \times 10^6 Z/n$
24. (d) Frequency of revolution
velocity in second orbit (V_2)
 $\frac{2 r_2}{\frac{1.082 \times 10^6 \text{ ms}^{-1}}{2 (2.12 \times 10^{10}) \text{ m}}} = 8.2 \times 10^{14} \text{ s}^{-1}$
25. (c) $v = \frac{Z}{n}$; $r = \frac{n^2}{Z}$;
frequency of revolution $\frac{v_n}{2 r_n}$;
Coulombic force of attraction $\frac{Ze^2}{(4 \pi \epsilon_0) r^2}$
26. (a) Number of wave in one complete revolution
shell number.

$$28. (d) \text{ For Be}^3 \quad E \quad E_2 \quad 13.6 \frac{Z^2}{n^2}$$

$$13.6 \frac{4^2}{2^2} = 54.4 \text{ eV}$$

$$29. (d) T \quad \frac{n^3}{Z^2}; \quad \frac{T_{1,2}}{T_{2,1}} \quad \frac{1}{4} \quad \frac{1}{8} \quad \frac{1}{32}$$

$$30. (d) mvr \quad \frac{nh}{2}$$

$$31. (b) T \quad \frac{n^3}{Z^2}; \quad \frac{T_1}{T_2} \quad \frac{n_1^3}{Z_1^2} \quad \frac{Z_2^2}{n_2^3}$$

$$\frac{Z^3}{1} \quad \frac{2^2}{3^3} \quad \frac{32}{27}$$

$$32. (c) \text{ When an electron of charge } e \text{ and mass } m \text{ is accelerated with a potential difference of } V \text{ volts K.E. } eV$$

$$\frac{1}{2} m v^2 = eV \text{ or } v^2 = \frac{2 eV}{m}$$

$$v = \sqrt{\frac{2 eV}{m}}$$

$$33. (d) E_H \text{ in first orbit } = \frac{19.6 \times 10^{18}}{4} \text{ J}$$

$$E_{\text{Be}^{3+}} \text{ in second orbit } = \frac{19.6 \times 10^{18}}{4} \times \frac{16}{4}$$

$$19.6 \times 10^{18} \text{ J}$$

$$34. (b) \frac{E_1}{E_2} = \frac{Z_1^2}{Z_2^2} = \frac{1^2}{2^2}$$

$$E_2 = 4E_1 = 13.64 \text{ eV}$$

$$35. (b) \because E_n = \frac{Z^2}{n^2} E_1 \quad E_n = Z^2$$

$$36. (a) \because \frac{mv^2}{r} = \frac{1}{4} \frac{Ze e}{r^2}$$

$$v^2 = \frac{e^2}{4 \times 0.7m} \quad (\text{For hydrogen } Z = 1)$$

$$v^2 = \frac{e^2}{4 \times 0.7m} \quad v = \frac{e}{\sqrt{4 \times 0.7m}}$$

$$38. (c) E = 13.6 Z^2 \frac{1}{n_1^2} - \frac{1}{n_2^2} \text{ eV atom}^{-1}$$

For the ionization of Li^2 ($Z = 3$) from first excited state, $n_1 = 2$ and $n_2 = \infty$.

$$\text{Hence, IP } E = 13.6 \times 3^2 \times \frac{1}{2^2} = \frac{1}{2}$$

$$30.6 \text{ eV}$$

$$39. (c) E = \frac{h\nu}{\lambda} = \frac{6.62 \times 10^{-34} \times 4.2 \times 10^{14}}{2.8 \times 10^{-19} \text{ J}}$$

$$40. (d) E = \frac{1240}{\lambda \text{ nm}} = \frac{1240}{91} = 13.6 \text{ eV}$$

$$41. (d) E = \frac{hc}{\lambda}$$

$$42. (d) E = n \frac{hc}{\lambda}$$

$$\frac{2 \times 10^{17}}{1.6 \times 10^{19}} \text{ eV} \quad n = \frac{1240}{595.2} \frac{\text{eV} \cdot \text{nm}}{\text{nm}}$$

$$n = 60$$

$$45. (d) n_1 = 2$$

$$n_2 = 6$$

$$\text{Total number of photon emitted} = \frac{(6-2)(6-2-1)}{2} = 10$$

$$46. (b) n_1 = 2$$

$$n_2 = n$$

$$\bar{\nu} = R(1)^2 \left[\frac{1}{(2)^2} - \frac{1}{n^2} \right]$$

$$R = \frac{n^2 \cdot 2^2}{4n^2}$$

$$49. (b) n_1 = n_2 = 4; \quad n_2 = n_1 = 2; \quad n_1 = 1, \quad n_2 = 3$$

$$- R(2)^2 \left[\frac{1}{1^2} - \frac{1}{3^2} \right] = \frac{32R}{9}$$

$$50. (c) \frac{91.2}{9 \times \frac{1}{(2)^2} - \frac{1}{(3)^2}} = 91.2 \times \frac{36}{9 \times 5} = 73 \text{ nm}$$

$$51. (c) E = 13.6 (2)^2 \left[\frac{1}{4} - \frac{1}{36} \right] = 96.43$$

$$1.16 \times 10^3 \text{ kJ/mol}$$

$$52. (c) \frac{1}{RZ^2} \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] = R \times 3^2 \left[\frac{1}{3^2} - \frac{1}{2^2} \right]$$

$$R \text{ or } \frac{1}{R}$$

$$53. (b) \frac{1}{RZ^2} \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right] = R \times 2^2 \left[\frac{1}{1^2} - \frac{1}{2^2} \right]$$

$$3R; \quad \frac{1}{3R}$$

54. (a) For the transition $n = 1$ to $n = 2$, the energy change, E is positive, i.e., energy is absorbed. For the transition $n = 5$ to $n = 1$, E is negative, i.e., energy is released.

55. (c) For hydrogen atom

$$\frac{1}{n_1^2} R_H \frac{1}{n_2^2} \frac{1}{n_2^2};$$

$$2170 \text{ nm} \quad 2170 \quad 10^9 \text{ m};$$

$$R_H \quad 1.09677 \quad 10^7 \text{ m}^{-1}$$

$$\frac{10^9}{2170} \quad 1.09677 \quad 10^7 \quad \frac{1}{n_1^2} \quad \frac{1}{7^2}; \quad n_1 = 4$$

So, electronic transition from $n = 7$ to $n = 4$ will produce infrared radiation of wavelength 2170 nm.

56. (a) Total number of spectral lines given by

$$\frac{1}{2} [n - 1] \quad n = 15; \quad n = 6$$

Thus, electron are excited upto 6th energy level from ground state. Therefore,

$$\frac{1}{2} R_H \frac{1}{1^2} \frac{1}{n^2} \quad 109737 \quad \frac{35}{36};$$

$$9.373 \times 10^6 \text{ cm}^{-1} \quad 937.3 \text{ Å}$$

57. (d) $E = \frac{1}{2} m v^2$ for $v_{\max}$, $E_{\min}$

58. (b) $K L M N \dots$ energy increases.

59. (b) $E = 13.6 Z^2 \frac{1}{n^2} \frac{1}{(n - 1)^2}$

63. (b) $h\nu = h\nu_0 + KE$

64. (b) $h\nu = h\nu_0 + eV_0$

$$V_0 = \frac{h\nu}{e} - \frac{h\nu_0}{e}$$

65. (c) $h\nu = h\nu_0 + KE$

67. (c) As per Einstein's equation of photoelectric effect

$$\frac{1}{2} m v^2 = h\nu - h\nu_0 = \frac{hc}{\lambda} - \frac{hc}{\lambda_0}$$

$$v^2 = \frac{2hc}{m} \left(\frac{1}{\lambda} - \frac{1}{\lambda_0} \right);$$

$$v = \frac{2hc}{m} \left(\frac{1}{\lambda} - \frac{1}{\lambda_0} \right)^{1/2}$$

$$\frac{2hc}{m} \left(\frac{1}{\lambda} - \frac{1}{\lambda_0} \right)$$

68. (c) $\frac{hc}{\lambda} = 1 \dots (1)$

$3 \frac{hc}{\lambda} = 4 \dots (2)$

from, eq. (1) and (2) 0.5 eV

69. (c) $\frac{1}{2} m v^2 = \frac{1240 \text{ eV nm}}{31 \text{ nm}} = 40 \text{ eV}$

$$= 27.2 \text{ eV}$$

$$\frac{1}{2} \quad 9.1 \times 10^{-31} \quad v^2 \quad 27.2 \quad 1.6 \times 10^{-19}$$

$v = 3.09 \times 10^6 \text{ m/s}$

70. (a) $h \frac{h}{e} = h_0 \frac{eV_0}{e}; \quad eV_0 = h \frac{h}{e} = h_0 \frac{h}{e}$ or

$$V_0 = \frac{h}{e} \quad \frac{h}{e} = h_0; \quad \text{slope}_1 = \frac{h}{e}$$

Similarly, $h = h_0 \frac{K_{\max}}{K_{\max}}$

or $K_{\max} = h \frac{h}{e};$

slope₂ $h, \frac{\text{slope}_2}{\text{slope}_1} = \frac{h}{h/e} = e$

71. (d) $E_n = \frac{13.6}{n^2} \text{ eV}; \quad E_2 = \frac{13.6}{2^2};$

$$E_4 = \frac{13.6}{4^2} \text{ eV/atom}$$

$$E = E_4 - E_2 = 2.55 \text{ eV}$$

Absorbed energy = work function of metal + K.E.

$$2.55 = 2.5 \text{ K.E.}; \quad \text{K.E.} = 0.05 \text{ eV}$$

74. (c) $\frac{1}{m}$

75. (b) $\frac{h}{mv} = \frac{6.63 \times 10^{-34}}{10^{-6} \times 10} = 6.63 \times 10^{-29} \text{ m}$

76. (b) $\frac{h}{\sqrt{2qVm}}$

77. (d) $\frac{h}{p}$

78. (b) $\therefore \frac{1}{n_1^2} R = \frac{1}{n_2^2} \frac{1}{n_2^2}; \quad n_1 = 1, n_2 = ?;$

$$\frac{1}{n_1^2} R = \frac{1}{1} \frac{1}{n_2^2} \quad n_2^2 = \frac{R}{R - 1}$$

$$n_2 = \sqrt{\frac{R}{R - 1}}$$

79. (b) $E = E_1 - E_2; \quad \frac{hc}{\lambda} = \frac{hc}{\lambda_1} - \frac{hc}{\lambda_2}$

$$\frac{hc}{\lambda} = hc \left(\frac{1}{\lambda_1} - \frac{1}{\lambda_2} \right); \quad \frac{1}{\lambda} = \frac{1}{\lambda_1} - \frac{1}{\lambda_2}$$

80. (d) $p = \frac{h}{\lambda} = \frac{h}{\sqrt{2 eV m_p}}$

$$\text{Be}^3 \quad \frac{h}{\sqrt{2 \times 3 \text{ eV} m_{\text{Be}^{3+}}}} = \frac{h}{\sqrt{2 \times 3 \text{ eV} \times 9 m_p}}$$

Hence, $\frac{\text{Be}^3}{p} = \sqrt{\frac{2 eV m_p}{3 eV \times 9 m_p}} \times \frac{1}{3\sqrt{3}}$

$$81. (c) \frac{h}{\sqrt{2} eV m} \frac{1.23}{\sqrt{V}} \text{ nm}$$

$$82. (b) \frac{h}{m_e x} \frac{h}{m_p V} \frac{h}{1840 m_e V} [m_p \quad 1840 m_e]$$

$$\text{Hence, } V = \frac{x}{1840}$$

$$83. (a) E = (mC) C \text{ or } P = \frac{E}{C}$$

$$\frac{6 \times 10^6}{3 \times 10^8} \frac{1.6 \times 10^{19}}{10^{21}} \text{ kg-m/s}$$

$$84. (c) E = \frac{nhc}{nhc^-} \therefore - \frac{1}{10} \frac{nhc^-}{hc^-} \text{ or } n = \frac{10}{hc^-} \frac{10}{hcx}$$

$$86. (b) \therefore mvr_n = \frac{nh}{2} \text{ and } p = \frac{h}{\lambda}$$

$$\therefore \frac{2r}{\lambda} = n$$

$$\therefore \frac{r}{2} = \frac{4a_0}{2}$$

$$\text{or } \frac{h}{r_2} = \frac{4a_0}{h} \quad r_2 = 4a_0$$

$$87. (b) \text{I.E.} = 13.6 \frac{Z^2}{n^2} \text{ eV} = 13.6 \times 4 = 54.4 \text{ eV}$$

$$\text{for 2 mole } 54.4 \times 2 \times N_A \text{ eV} = 108.8 N_A \text{ eV}$$

$$90. (a) m = 10^{10} \text{ g} = 10^{13} \text{ kg};$$

$$v = \frac{0.0001}{100} \times 10^6 \times 10^2 \times 10^{14} \text{ m sec}^{-1}$$

$$x = p = \frac{h}{4}$$

$$x = \frac{h}{4p} = \frac{h}{4mv}$$

$$x = \frac{6.62 \times 10^{-34}}{4 \times 3.14 \times 10^{13} \times 10^{14}}$$

$$x = \frac{6.62}{12.56} \times \frac{10^{-34}}{10^{27}}; \quad x = 5.2 \times 10^{-8} \text{ m}$$

$$91. (c) x = \frac{h}{4p} = \frac{h}{4mv}$$

$$\frac{6.63 \times 10^{-14}}{4 \times 3.14 \times 9.1 \times 10^{31} \times 1}$$

$$(\therefore v = 1 \text{ m/s})$$

$$x = 58 \text{ m}$$

94. (a) Use $(n - l)$ rule.

101. (c) Radii of maximum probability of electron is smaller for orbital having higher value of 'l' for given same value of n.

103 (b) Total no. of nodes $(n - 1)$

104. (c) $l = 4$;

number of degenerate orbitals $2l + 1 = 9$

109. (b) 4f has the highest energy among 3d, 4f, 4p, 5s orbitals.

110. (c) m orientation of orbital.

112. (c) Maximum number of electrons with same spin is equal to maximum number of orbitals, i.e., $(2l + 1)$.

113. (d) Orbital angular momentum $\sqrt{l(l+1)} \frac{h}{2}$; $l = 1$ for p-orbital.

116. (d) [Ne] $1s^2 2s^2 2p^6$

117. (b) Fe(III)—[Ar] $3d^5$; unpaired electrons = 5;


magnetic moment $\sqrt{5(5+2)} \text{ BM}$

Co(II)—[Ar] $3d^7$; unpaired electrons = 3;

magnetic moment $\sqrt{3(3+2)} \text{ BM}$

119. (d) Given $\sqrt{n(n+2)} = 1.73 \text{ BM}$
(where n is number of unpaired electrons)

$n = 1$; ${}_{23}\text{V } 1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^3$




for $n = 1$ it must release 4 electrons, first two from 4s-orbital and then next two electrons from 3d-orbital. So, $x = 4$.

120. (c)

1	1	1	1	1	1
---	---	---	---	---	---

$|s| = \frac{\text{No. of unpaired } e}{2}$



$$123. (b) I \quad r = 0.529 \frac{n^2}{Z} \text{ \AA}$$

$$II \quad l = \sqrt{l(l+1)} \frac{h}{2}$$

$$l = 0$$

$$124. (c) \quad n \quad l$$

$$e_1 \quad 3$$

$$e_2 \quad 4$$

$$e_3 \quad 5$$

$$e_4 \quad 4$$

$$e_3 \quad e_2 \quad e_4 \quad e_1$$

$$125. (d) E_{2p_x} \quad E_{2p_y} \quad E_{2p_z}$$

$$126. (c) P(33) \quad [Ar] 4s^2 4p^3$$

$$n = 3$$

$$128. (d) 3d\text{-subshell contains maximum 10 electrons.}$$

$$130. (d) |m| = l$$

$$131. (c) \text{ Degenerate orbitals having equal energy.}$$

$$140. (c) \text{ For } f \text{ subshell } l = 3; \quad g \text{ subshell } l = 4$$

$$\text{for principal shell, } l = 4, n = 5$$

$$\text{total no. of orbital in shell } n^2 = 5^2 = 25$$

$$142. (c) Fe(26) [Ar] 3d^6 4s^2$$

$$143. (a) r = n^2$$

$$145. (d) \text{ Total nodes } = n - 1$$

$$146. (c) \text{ Unpaired electron of Cl atom is } 3p^1 \text{ for which } n = 3,$$

$$l = 1, m = 1, 0 \text{ or } +1 \text{ and } s = \frac{1}{2}$$

$$147. (d) Cl(17) 3s^2, 3p^5 \quad \boxed{\uparrow} \quad \boxed{\uparrow} \quad \boxed{\uparrow}$$

$$n = 3, l = 1, m = 1, s = \frac{1}{2}$$

$$149. (a) A : \text{excitation possible only in } d\text{-orbitals}$$

$$B : \text{Spin multiplicity } = 2|S| + 1; |S|$$

$$= \text{total spin}$$

$$C : V \text{ not violated Hund's rule}$$

$$D : A \text{ is paramagnetic due to unpaired } e$$

$$A, B, C \text{ are correct.}$$

$$150. (d) \text{ Orbitals are } 4s, 2s, 3p \text{ and } 3d. \text{ Out of these } 3d \text{ has highest energy.}$$

$$151. (b) 5f \text{ subshell}$$

$$154. (a) \text{ Pauli's exclusion principle}$$

$$155. (c) \text{ For H-atom, energy of } s\text{-orbital} = \text{energy of } 2p\text{-orbital so degeneracy is } 1 + 3 = 4$$

$$156. (a) \text{ For multi electron species (H) first excited state means } e \text{ will transfer from } 1s \text{ to } 2s \text{ subshell \& } 2s \text{ subshell has only one orbital so degeneracy is } 1.$$

$$158. (d) (n - l) \text{ rule}$$

$$160. (d) n - l$$

$$161. (c) 1s^2 2s^2 \quad \boxed{2p^6} \quad 3s^2 \quad \boxed{3p^6}$$

$$\begin{array}{|c|c|c|} \hline \boxed{\uparrow\downarrow} & \boxed{\uparrow} & \boxed{\uparrow} \\ \hline 0 & +1 & -1 \end{array} \quad \begin{array}{|c|c|c|} \hline \boxed{\uparrow\downarrow} & \boxed{\uparrow} & \boxed{\uparrow} \\ \hline 0 & +1 & -1 \end{array}$$

$$163. (d) 4s\text{-orbital}$$

$$164. (d) g\text{-subshell } l = 4, 2(2l + 1) = 18e$$

$$171. (a) Mn(25) \quad [Ar] 3d^5, 4s^2 \quad n = 5$$

$$Ti(22) \quad [Ar] 3d^2, 4s^2 \quad n = 2$$

$$V(23) \quad [Ar] 3d^3, 4s^2 \quad n = 3$$

$$Al(13) \quad [Ne] 3s^2, 3p^1 \quad n = 1$$

$$172. (d) \text{ No. of spherical nodes } = n - l - 1; \text{ for } s, p, d, f \text{ values of } l = 0, 1, 2, 3 \text{ respectively.}$$

$$173. (c) \text{ Radial node } (n - l - 1) = 2$$

$$l = 0, n = 3$$

$$174. (b) \text{ For radial node } 2s \quad 0, 2 \quad \frac{r}{a_0} = 0$$

$$\boxed{r = 2a_0}$$

$$175. (c) \text{ Probability of finding } e \text{ is zero implies that}$$

$$^2_0 \text{ or } ^0_0$$

$$(1) = 0 \quad 1$$

$$\text{or } r_1 \frac{a_0}{2z} \text{ or } (2, 8, 12) = 0$$

$$\text{and } (6)(2) = 0$$

$$6, r \frac{3a_0}{z}$$

$$\text{and } 2, r \frac{a_0}{z}$$

$$r_2 \frac{3a_0}{z}$$

Level 2

1. (c) P.E. $\frac{1}{4} \frac{(Ze)(e)}{r}$
 $\frac{1}{4} \frac{(2e)(e)}{r} = \frac{e^2}{2 \cdot 0 \cdot r}$
2. (c) Ionization energy :
 $217.6 = -13.6 \frac{Z^2}{1^2}; Z = 4$
 So, it is ${}^9_4\text{Be}^3$; no. of neutrons $9 - 4 = 5$
3. (c) $\frac{d[U(r)]}{dr} = \frac{3Ke^2}{r^4}$ Magnitude of the force
 $\frac{3Ke^2}{r^4} = \frac{mv^2}{r}$
 and we know $mvr = \frac{nh}{2}$ or $r = \frac{nh}{2mv}$,
 $3Ke^2 = \frac{8}{n^3h^3} \frac{m^3v^3}{mv^2}$, $v = \frac{n^3h^3}{24Ke^2 \cdot 3m^2}$
4. (c) $\frac{1}{2}$ P.E. = K.E.
 $\frac{1}{2} \frac{1}{2} mkr^2 = \frac{1}{2} mv^2$, $mvr = \frac{nh}{2}$,
 $v^2 = \frac{n^2h^2}{4m^2r^2}$; $r^4 = \frac{n^2h^2}{2m^2k}$
 or $r = \sqrt{\frac{n^2h^2}{2m^2k}}$
5. (a) $\frac{1}{2}mv^2 = \frac{k(q_1)q_2}{r} = \frac{q_2}{m} \frac{r \cdot v^2}{2k \cdot q_1 \cdot Z}$
 $\frac{q_2}{m} = \frac{2.5 \cdot 10^{-14}}{2 \cdot 9 \cdot 10^9} \frac{(2.1 \cdot 10^7)^2}{79 \cdot 1.6 \cdot 10^{-19}}$
 $4.84 \cdot 10^7$ coulomb/kg
6. (d) $v = r \omega$ where $r_n = \frac{n^2h^2}{4me^2ZK}$
 and $v_n = \frac{2Ze^2K}{nh} \frac{nh}{n^2h^2}$;
 $\frac{2Ze^2K}{nh} \frac{nh}{4me^4ZK} = \frac{8me^4Z^2K^2}{n^3h^3}$;
 $\frac{9me^4K^2}{h^3} (\because n=2 \text{ and } Z=3)$
7. (b) $r_n = \frac{v^2}{Z}$; for H, $r_4 = r_3 = 0.529(16-9)$
 $0.529 \cdot 7 \text{ \AA}$
 $r_4 = r_3$ for Li^2 $0.529 \frac{16}{3} - \frac{9}{3}$

$$0.529 \frac{7}{3} \text{ so ratio } \frac{7}{7/3} = 3:1$$

$$8. (c) v_n = 2.186 \cdot 10^6 \frac{Z}{n}$$

$$1.093 \cdot 10^6 = 2.186 \cdot 10^6 \frac{1}{n}; n = 2$$

from Bohr theory we know $2\pi r = n \frac{h}{mv}$
 2π , where $\frac{h}{mv}$

$$\text{or } r = 0.529 \frac{n^2}{Z} = 0.529 \cdot 4 \text{ \AA}$$

Circumference of the orbit

$$2\pi \frac{22}{7} \cdot 0.529 \cdot 4 \cdot 10^{-10}$$

$$13.30 \cdot 10^{-10} \text{ m}$$

$$9. (b) \text{Angular momentum } \frac{nh}{2}$$

$$3.1652 \cdot 10^{-34} = \frac{n \cdot 6.626 \cdot 10^{-34}}{2};$$

$$n = 3$$

$$-RZ^2 \frac{1}{n_1^2} - \frac{1}{n_2^2};$$

$$-RZ^2 \frac{1}{2^2} - \frac{1}{3^2} = \frac{5R}{9}$$

10. (d) Energy of photon corresponding to second line of Balmer series for Li^{2+} ion

$$= (13.6) (3)^2 \frac{1}{2^2} - \frac{1}{4^2}$$

$$= 13.6 \frac{27}{16}$$

Energy needed to eject electron from $n=2$ level in H-atom;

$$= 13.6 \cdot 1^2 - \frac{1}{2^2} = \frac{13.6}{4}$$

K.E. of ejected electron

$$13.6 \frac{9}{16} - \frac{13.6}{4} = 13.6 \frac{27}{16} - \frac{4}{16}$$

$$19.55 \text{ eV}$$

$$11. (a) \frac{1}{n_1^2} - RZ^2 \frac{1}{n_2^2} = \frac{1}{n_1^2} - \frac{1}{n_2^2}, \text{ where } n_1 = n, n_2 = n+1$$


$$\frac{1}{n^2} - RZ^2 \frac{1}{(n+1)^2} = \frac{1}{(n+1)^2}$$

$$\frac{1}{n^2} - \frac{2n-1}{n^2(n+1)^2} = RZ^2$$

Since, $n \gg 1$;

Therefore, $2n-1 \approx 2n$

$$\text{and } \frac{1}{RZ^2} \frac{n^2}{n^2} \frac{2n}{n^2} \\ \frac{2RZ^2}{n^3} \text{ or } \nu = \frac{2cRZ^2}{n^3}$$



Total radiations are = 6

13. (a) If a is side of cube, then $x = a\sqrt{3}$
 $x = 10\sqrt{3} \text{ cm} = 10\sqrt{3} \times 10^{-2} \text{ m}$
 $x = p \frac{h}{4m}; \quad x = m \frac{h}{4m} \quad \nu = \frac{h}{4m} \frac{1}{x}$
 $\nu = \frac{6.63 \times 10^{-34}}{4 \times 3.14 \times 9.1 \times 10^{-31} \times 10 \times \sqrt{3} \times 10^{-2}}$
 $\nu = 3.34 \times 10^{-4} \text{ ms}^{-1}$

14. (b) $2r = 4a_0; n = 4$
 Total energy required = total energy released
 $0 = 2 \times 4.526 N_A - 2 \times 13.6 N_A - 2 \times 13.6 N_A + \frac{1}{16} N_A - 30.87 N_A = 0$
 $x = 2 \text{ moles of } X \text{ required} = 2$

15. (c) Work function for $\text{Li}^{2+} = 9E$.
 $E_p = w + \frac{1}{2}mv^2; \quad E_p = 9E = \frac{1}{2}mv^2$
 $\nu = \sqrt{\frac{2(E_p - 9E)}{m}}$

16. (c) $E = hc\nu = 1.63 \times 10^{-18} \text{ J}$
 where $\nu = R(1)^2 \frac{1}{1^2} - \frac{1}{2^2} = \frac{3}{4}R$
 Translational K.E. of H-atom $= \frac{3}{2} \frac{R}{N_A} T$
 $\frac{3}{2} \times \frac{8.314}{6 \times 10^{23}} T = 1.63 \times 10^{-18}$
 $T = 7.84 \times 10^4 \text{ K}$

17. (c) Radial node occurs where probability of finding e is zero.


$$\frac{2}{6} \times 0 \text{ or } 0; \quad \frac{3}{3} \times \sqrt{3}$$

For max. distance $r = \frac{3}{2} \left(3 - \sqrt{3} \right) a_0$

18. (d) $\frac{n(n-1)}{2} = 6; n = 4$
 $n = 4 \quad E_4 = 0.85 \text{ eV}$
 $n = 1 \quad E_1 = 13.6 \text{ eV}$
 $E = 12.75 \text{ eV}$
 $12.75 \text{ eV} = \frac{1240 \text{ eV} \cdot \text{nm}}{97.25 \text{ nm}}$

19. (d) For II to I transition
 $E = \frac{4E}{3} \quad E = \frac{hc}{\lambda}; \quad \frac{E}{\lambda} = \frac{hc}{\lambda}$

For III to I transition
 $E = 2E \quad E = \frac{hc}{\lambda} \text{ or } E = \frac{hc}{\lambda}$
 $\frac{hc}{3} = \frac{hc}{\lambda} \quad \lambda = 3$



Out of 6 electrons in $2p$ and $3p$ must have one electron with $m = 1$ and $s = \frac{1}{2}$ but in $3d$ -subshell an orbital having $m = 1$ may have spin quantum no. $\frac{1}{2}$ or $-\frac{1}{2}$.

Therefore, minimum and maximum possible values are 2 and 3 respectively.

21. (a) Energy absorbed $13.6 - 1.5 = 20.4 \text{ eV}$ out of this 6.8 eV is converted to K.E.

$$6.8 \text{ eV} = 6.8 \times 1.6 \times 10^{-19} \text{ J};$$

$$6.8 \times 1.6 \times 10^{-19} = \text{K.E.} = \frac{1}{2} mv^2$$

$$\nu = \sqrt{\frac{2KE}{m}} = \sqrt{\frac{2 \times 1.088 \times 10^{-18}}{9.1 \times 10^{-31}}}$$

$$1.54 \times 10^6 \text{ m/s}$$

22. (b) $E = \frac{hc}{\lambda} = 2.9 \times 10^{-19} \text{ J}$

Total energy of 10 quanta

$$10 \times 2.9 \times 10^{19} \times 29 \times 10^{19} \text{ J}$$

Energy stored for process

$$\frac{112 \times 4.18 \times 10^3}{6 \times 10^{23}} \times 7.80 \times 10^{19} \text{ J}$$

$$\% \text{ efficiency} = \frac{7.8 \times 10^{-19}}{29 \times 10^{19}} \times 100 = 26.9\%$$

23. (c) Total energy emitted by photo-electron

$$10.2 + 17 = 27.20 \text{ eV}$$

Since, E_1 Photon of energy emitted through the transition

$$n \rightarrow n-2 \quad \frac{hc}{\lambda} = 27.20 \text{ eV}$$

$$\text{We have } \frac{1}{n} R_H Z^2 \frac{1}{2^2} - \frac{1}{n^2}$$

$$\text{or } \frac{hc}{\lambda} = (hc) R_H Z^2 \left(\frac{1}{2^2} - \frac{1}{n^2} \right)$$

$$27.20 = (hc) R_H Z^2 \left(\frac{1}{4} - \frac{1}{n^2} \right) \quad \dots(1)$$

Similarly, total energy liberated during transition of electron from $n \rightarrow n-3$ is


$$E_2 = \frac{hc}{\lambda} (4.25 + 5.95) = 10.20 \text{ eV}$$

$$10.20 = (hc) R_H Z^2 \left(\frac{1}{9} - \frac{1}{n^2} \right) \quad \dots(2)$$

Dividing Eq. (1) by (2), we get $n = 6$ and putting $n = 6$ in Eq. (1) or (2), we get, $Z = 3$.

24. (a) $E = \frac{hc}{\lambda} = \frac{1240 \text{ eV-nm}}{1025.6 \times 10^{10} \times 10^9}$

$$E = 12.09 \text{ eV}$$



In three different radiations, minimum wavelength for $3 \rightarrow 1$ transition

$$\frac{hc}{\lambda} = E = 12.09 \text{ eV} \quad \lambda = 102.6 \text{ nm}$$

25. (b) $E = \frac{hc}{\lambda} = \frac{(6.63 \times 10^{-34} \text{ J-s})(3.00 \times 10^8 \text{ m/s})}{3.055 \times 10^{-8} \text{ m}}$

$$6.52 \times 10^{-18} \text{ J}$$

$$E_H = \frac{3}{4} (2.176 \times 10^{-18} \text{ J})$$

$$1.63 \times 10^{-18} \text{ J}; \quad E = E_H(Z^2)$$

$$Z^2 = \frac{E}{E_H} = \frac{(6.52 \times 10^{-18})}{(1.63 \times 10^{-18})} = 4;$$

$$Z = 2 \text{ (helium)}$$

26. (b) K.E. $\frac{K \cdot Z e^2}{r}$

$$r = \frac{9 \times 10^9 \times 29 \times 2 \times (1.6 \times 10^{-19})^2}{5 \times 1.6 \times 10^{19} \times 10^6}$$

$$r = \frac{9 \times 10^9 \times 29 \times 2 \times 1.6 \times 10^{-19}}{5 \times 10^6}$$

$$1.67 \times 10^{-14} \text{ m}$$

27. (c) $\sqrt{v} = aZ$ ab

$$\frac{ab}{\sqrt{v}} = \frac{1}{51} \times \frac{a}{1} \times \frac{\tan 45^\circ}{50}$$

$$v = 50^2 \times 2500 \text{ s}^{-1}$$

28. (c) $\frac{1}{R} = \frac{1}{2^2} - \frac{1}{n^2} \Rightarrow R = \frac{n^2}{4n^2}$

$$\frac{4}{R} = \frac{n^2}{4} \quad \dots(1)$$

Given: $k = \frac{n^2}{n^2 - 4} \quad \dots(2)$

Comparing equation (1) and (2) we have

$$K = \frac{4}{R}$$

29. (b) $2 r_n = n \times 2 \times 0.53 \frac{n^2}{z} = n$

$$2 \times 0.53 \frac{n}{z} = n \quad \dots(1)$$

$$E_{\text{sep}} = 3.4 \times 13.6 \frac{z^2}{n^2} = \frac{n}{z} \times 2$$

$$2p \times 0.53 \times 2 \times 6.66 \text{ \AA}$$

30. (a) $\frac{E}{E_{4\text{th}}} = \frac{24}{E_{4\text{th}}} \times \frac{1}{16} = \frac{3/4}{1/16}$

$$E_{4\text{th}} = 2 \text{ eV.}$$

Level 3

Passage-1

1. (c) $x = p \times \frac{h}{4} = p^2 \times \frac{h}{4}$

$$m^2 v^2 = \frac{h}{4} \quad v = \frac{1}{2m} \sqrt{\frac{h}{4}}$$

$$2. (a) \quad x \sqrt{\frac{h}{4m}}; \quad x p \frac{h}{4}$$

$$\sqrt{\frac{h}{4m}} p \frac{h}{4}, \quad p \sqrt{\frac{mh}{4}}$$

$$3. (c) \quad \text{D.B.} \quad \sqrt{\frac{150}{6}} \text{ \AA} = 5 \text{ \AA}$$

$$\text{and} \quad x p \frac{h}{4}; \quad p \frac{h}{4} \text{ or } p \frac{h}{2}$$

$$x \frac{h}{2} \quad \frac{h}{4}$$

$$\frac{1}{2} \frac{10^9}{2} \quad \frac{1}{4} \quad \frac{2.5}{4} 10^{10}$$

$$0.625 \text{ \AA}$$

Passage-2

$$1. (b) \quad \therefore \frac{1}{R_H} Z^2 \left(\frac{1}{n^2} \right) \text{ for Lyman's series}$$

For He ion series limit n ;

$$\frac{1}{R_H} Z^2 \left(\frac{1}{n^2} \right) \quad \frac{1}{R_H} Z^2 \left(\frac{1}{4} \right)$$

Similarly, Balmer series limit for Li^{2+} ion

$$\frac{1}{R_H} Z^2 \left(\frac{1}{n^2} \right) \quad \frac{1}{R_H} Z^2 \left(\frac{1}{4} \right) \quad \frac{4x}{9}$$

$$2. (b) \quad 6 \left(n_2 - n_1 \right) \left(\frac{n_2 - n_1}{2} \right); \quad n_2 = 4, n_1 = 1$$

$$\frac{1}{R_H} Z^2 \left(\frac{1}{n^2} \right) \quad \frac{1}{16}$$

$$R \frac{15}{4}; \quad \frac{4}{15R_H}$$

$$3. (d) \quad \frac{1}{R_H} Z^2 \left(\frac{1}{n_1^2} \right) \quad \frac{1}{n_2^2}$$

$$\frac{1}{R_H} Z^2 \left(\frac{1}{2^2} \right) \quad \frac{1}{4^2}$$

$$R_H \frac{1}{1^2} \quad \frac{1}{2^2} \quad \frac{1}{H}$$

$$\text{for } n_2 = 2 \text{ to } n_1 = 1$$

Passage-3

$$1. (d) \quad \text{Magnetic moment} \quad \sqrt{n(n+2)}$$

$$(A) \quad \text{Fe}^{2+} \quad 3d^6 4s^0 \quad n = 4$$

$$(B) \quad \text{Mn}^{2+} \quad 3d^5 4s^0 \quad n = 5$$

$$(C) \quad \text{Cr}^{3+} \quad 3d^3 4s^0 \quad n = 3$$

$$(D) \quad \text{V}^{3+} \quad 3d^2 4s^0 \quad n = 2$$

Smaller n , smaller

$$2. (c) \quad \sqrt{n(n+2)} \quad 3.873$$

$$n^3$$

$$\text{Mn} \quad 3d^5 \quad 4s^2$$

$$\text{Mn}^{4+} \quad 3d^3 \quad 4s^0$$

Passage-4

1. (b) Total energy required for dissociation of O_2 molecule and then assigning (O^*) 1eV more energy than (O) (480+96) 576 kJ/mol

Maximum wavelength $E = \frac{hc}{\lambda}$;

$$\text{maximum} \quad \frac{hc}{E_{\min}}; \quad \text{max} (\text{\AA}) \quad \frac{12400}{E_{\min}(\text{eV})}$$

$$\frac{12400}{6} \quad 2066.67 \text{ \AA}$$

$$2. (b) \quad \text{O}_3(\text{g}) \quad \text{O}_2(\text{g}) \quad \text{O} \quad \dots (i)$$

$$\text{O}_2(\text{g}) \quad \text{O} \quad \text{O}^* \quad \dots (ii)$$

Energy required for (i) reaction is 400 kJ/mol

Normal dissociation of O_2 required 480 kJ/mol (given)

each (O^*) has 1 eV more energy than (O) and given 1 eV/photon 96 kJ/mol

So total energy required for the dissociation of O_3 into O and O^* is 400 + 480 + 96 = 976 kJ/mol

One or More Answers is/are Correct

2. (a, b, c)

$$mvr = n \frac{h}{2}$$

n integer value 1,2,3,4, ...

15. (a,b,c,d) At the point of maximum value of RDF

$$\frac{dP}{dr} = 0$$

$$2r \frac{2Zr^2}{a_0} = 0; \quad r = \frac{a_0}{Z}$$

where $Z = 3$ for Li^{2+} and $Z = 2$ for the He ;

$Z = 1$ for hydrogen.

18. (b, c)

$$h = W = KE$$

Kinetic energy of electrons does not depend on intensity of light $\frac{h}{\sqrt{2KE m}}$

19. (a, b, c, d)

(a) $v \propto \frac{Z}{n}$ (b) $r \propto \frac{n^2}{Z}$

(c) P.E. $\propto \frac{Z^2}{n^2}$ (d) K.E. $\propto \frac{Z^2}{n^2}$

20. (c, d)

Given $n, l = 0$ to $n-1$ $m_l = -l, \dots, 0, \dots, l$

21. (a, b, d)

(a) No. of subshell $n, n = 3$ (b) No. of orbital $2n^2, n = 4$ (c) Cu $3d^{10}4s^0$ diamagnetic

22. (a, b, c, d)

5 1	5 2	5 3	5 4
			Brackett

4 1	4 2	4 3
		Paschen

3 1	3 2
2 1	Balmer
Lyman	

Match the Column

13. (a) s-orbital $\therefore r = 0, \dots, 0$ and 3 radial nodes(b) 3 radial nodes (s, p, d) $4s, 5p_x, 6d_{xy}$ (c) Angular probability is dependent of θ and ϕ for $5p_y, 6d_{xy}$ (d) At least one angular node $5p_x(1); 6d_{xy}(2)$

Subjective Problems

1. $r_{n-1} = r_{n-1} = 2r_n$
 $0.529 \frac{(n-1)^2}{Z} = 0.529 \frac{(n-1)^2}{Z} = 2 \cdot 0.529 \frac{n^2}{Z}$
 $n^2 = 2n-1 \Rightarrow [n^2 - 2n + 1] = 2n^2 - 4n + 2n^2$
 $n = 2$

2. $E = 30.6 - 13.6 \frac{(3)^2}{(n)^2}$
 $n = 2$ no. of wave in one complete evolution in the orbit.

3. $n = 2$
 2nd orbit $1.5 \frac{n}{2}$

n = 3 no. of wave in one complete revolution in the orbit.

4. $E_{\text{abs}} = \frac{40}{100} E_{\text{Emitted}}$
 $n_{\text{ab}} = \frac{hc}{400} = \frac{40}{100} n_{\text{em}} = \frac{hc}{500}$
 $\frac{n_{\text{ab}}}{n_{\text{em}}} = \frac{400}{500} = \frac{100}{40} = 2$

5. $E_{\text{inc}} = KE$
 $4.5 = 3.0 + KE$
 $KE = 1.5 \text{ eV}$
 $\sqrt{\frac{150}{1.5}} = 10 \text{ \AA}$

$P \propto \frac{1}{4}$
 $\frac{h}{2} \propto \frac{h}{4}$
 $\frac{1}{(10 \cdot 10^{10})^2} = \frac{25}{4} \cdot 10^{10} \cdot \frac{1}{4}$
 $4 \text{ \AA}$

6. Cr $3d^5 4s^1$
 No. of angular nodes = 2
 No. of radial nodes = 3 + 2 + 1 = 0
 Angular node = radial node = 2

7. Total node = Radial node + angular node
 $n = 1 + 1 + 1 = n = 1$

5f $5 = 1 + 4$

8. $P = \frac{hc}{t}$
 $\frac{100}{100} = \frac{12}{100} (x \cdot 10^{19}) = \frac{6.63 \cdot 10^{34} \cdot 3 \cdot 10^8}{1500 \cdot 10^9 \cdot 1}$

9. $\frac{1}{n^2} = RZ^2 \frac{1}{n^2} - \frac{1}{(n-1)^2}$
 $\frac{1}{n^2} = RZ^2 \frac{(n-1)^2 - n^2}{n^2(n-1)^2}$
 $\frac{1}{n^2} = RZ^2 \frac{(n-1-n)(n-1+n)}{n^2(n-1)^2}$
 $\frac{1}{n^2} = RZ^2 \frac{(2n-1)}{n^2(n-1)^2}$
 n^3
 $x = 3$

10. For radial node

$$3s \quad 0 \quad \frac{1}{81\sqrt{3}} \quad \frac{1}{a_0} \quad 27 \quad \frac{18r}{a_0} \quad \frac{2r^2}{a_0^2} \quad e^{-r/3a_0}$$

$$27 \quad \frac{18r}{a_0} \quad \frac{2r^2}{a_0^2} \quad 0$$

$$r \quad \frac{(18a_0)^2}{2} \quad \sqrt{\frac{(18a_0)^2}{2} - 4 \cdot 27a_0^2}$$

$$\frac{18a_0}{4} \quad 10.39a_0$$

distance between to radial node (d)

$$\frac{d}{1.73a_0} \quad \frac{2}{4} \quad \frac{10.39a_0}{1.73a_0}$$

$$d \quad 3$$

$$11. \quad 7.67 \times 10^{19} \quad \frac{(1.6 \times 10^{19})^2}{4 \cdot 3.14 \cdot 8.85 \times 10^{-12} \cdot r}$$

$$r \quad 3.00 \times 10^{10} \text{ m} \quad 3\text{\AA}$$

$$12. \quad d(KE) \quad mv \quad dv \quad mv \quad \frac{h}{4 \cdot m \cdot x}$$

$$\frac{3}{3} \quad \frac{10}{4} \quad \frac{6.62 \times 10^{34}}{3.31 \times 10^{12}}$$

$$5 \quad 10^{16} \text{ J}$$

$$E_{\text{in}} \quad 10.2 \text{ eV}$$

$$4.2 \text{ eV}$$

$$KE_{\text{max}} \quad 10.2 \quad 4.2 \quad 6 \text{ eV}$$

$$e \quad \sqrt{\frac{150}{6}} \text{ \AA} \quad 5\text{\AA}$$

$$14. \quad \frac{\text{Radial probability density at } r}{\text{Radial probability density at } r=0} = \frac{R^2(a_0)}{R^2(0)}$$

$$\text{For 1s orbital: } R(r) = \frac{1}{\sqrt{a_0}} e^{-r/a_0}$$

$$\frac{R^2(a_0)}{R^2(0)} = \frac{(1/a_0^3)e^{-2r/a_0}}{(1/a_0^3)e^0} = e^{-2}$$

$$15. \quad A \quad \frac{E_{1,2}}{2E_{21}} \quad \frac{13.6}{2} \quad \frac{2^2}{1^2} - \frac{2^2}{(13.6)^2} \quad \frac{16}{2} \quad 8$$

GASEOUS STATE

Gaseous State

This state is characterized by sensitivity of volume change with change of pressure and temperature. It is due to large distance between molecules as compared to their own dimensions. There exists weak vander Waal's forces. (but no force in ideal state).

Ideal Gas: A gas with no intermolecular attractions & having very negligible volume occupied by molecules when compared with volume of gas is termed as ideal gas. An ideal gas is a hypothetical gas and none of the gases known is ideal. These gases are called real gases and they may behave like an ideal gas under specific conditions.

Parameter associated with the ideal gas : P, V, T, n
where

P represents pressure exerted by the gas molecules on the walls of the container assuming negligible intermolecular attractions,

V represents free volume available for motion (equal to the volume of the container),

T represents absolute temperature, n represents no. of moles.

❖ Conversion factors

Pressure : atm = 1.013 $\times 10^5$ Pa 760 mm of Hg 760 torr 1.013 bar

Volume : 1 L 1 dm³ 10⁻³ m³ 1000 mL 1000 cm³

Temperature : T_K T_C° 273 $\frac{5}{9} T_F$ 255.22

Instruments for Pressure Measurement

❖ **Barometer:** The instrument used for the measurement of atmospheric pressure is called a barometer.

$$P = \frac{\rho \cdot h \cdot g}{A} \cdot A$$

where ρ density of fluid

h vertical height

g acceleration due to gravity

❖ **Manometer:** The instrument used for the measurement of the pressure of a gas is called a manometer. manometer are of two types

- (i) Open end (ii) Closed end

Experimental Gas Laws

I.	Boyle's law	V	$\frac{1}{P}$	(T & n constant)	$P_1 V_1$	$P_2 V_2$
II.	Charle's law	V	T	(P & n constant)	$\frac{V_1}{T_1}$	$\frac{V_2}{T_2}$
III.	Gay Lussac's law	P	T	(V & n constant)	$\frac{P_1}{T_1}$	$\frac{P_2}{T_2}$
IV.	Avogadro's law	V	n	(T & P constant)	$\frac{V_1}{n_1}$	$\frac{V_2}{n_2}$

Combined Gas law : $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$

Equation of State: $PV = nRT$

$PV = \frac{w}{M} RT$ R is Universal Gas constant,

R 0.0821 atm. L/K. mol 8.314 joule/K. mol 2cal/ K.mol

density of gas (d) $\frac{PM}{RT}$

❖ **Dalton's law of partial pressure:**

$P_{\text{total}}, P_A, P_B, \dots$

P_A, P_B are partial pressures; $P_A = \frac{\text{mole fraction}_A}{\text{Partial pressure}} \times \text{Total pressure}$
and % of gas in mixture $\frac{\text{Partial pressure}}{\text{Total pressure}} \times 100$

$P_{\text{wet gas}} = P_{\text{dry gas}} + P_{\text{H}_2\text{O vapour}}$ (aqueous tension)

❖ **Amagat's law:** At constant temperature and pressure, the total volume of a mixture of gases is equal to the sum of the partial volumes of the constituent gases.

❖ **Relative humidity:**

% Relative humidity $\frac{\text{Partial pressure of H}_2\text{O}(g)}{\text{Vapour pressure of H}_2\text{O}(l)} \times 100$

❖ **Graham's law of Diffusion or Effusion:**

$r \propto \frac{1}{\sqrt{d}}$ or $r \propto \frac{1}{\sqrt{M}}$ $r \propto \frac{P}{\sqrt{M}}$ [For gases effusing at different pressures]

r is rate of diffusion of any gas, d is density at same temperature.

$\frac{r_1}{r_2} = \frac{\sqrt{d_2}}{\sqrt{d_1}} = \frac{\sqrt{M_2}}{\sqrt{M_1}}; \frac{V_1 / t_1}{V_2 / t_2} = \frac{\sqrt{M_2}}{\sqrt{M_1}} \frac{\text{Pressure drop I}}{\text{Pressure drop II}}$

$r = \frac{\text{moles diffused}}{\text{time taken}} = \frac{\text{distance travelled in a narrow tube}}{\text{time taken}}$

❖ **Isotopic Separation Factor:**

After x steps of separation, isotopic separation factor (f) $\frac{(n_1 / n_2)_{\text{final}}}{(n_1 / n_2)_{\text{initial}}} \sqrt{\frac{M_2}{M_1}}^x$

or
$$x = \frac{2 \log f}{\log \frac{M_2}{M_1}}$$

Kinetic Theory of Gases

$$PV = \frac{1}{3} m N u^2 = \frac{1}{3} M u^2 \quad (\text{For 1 mole})$$

Types of speeds:

❖ Mean square speed, $u^2 = \frac{u_1^2 + u_2^2 + \dots + u_N^2}{N}$

❖ Root mean square speed, $u = \sqrt{\frac{3RT}{M}} = \sqrt{\frac{3PV}{nM}} = \sqrt{\frac{3P}{d}}$

❖ Average speed $\frac{u_1 + u_2 + \dots + u_N}{N} = \sqrt{\frac{8RT}{M}} = \bar{u}$

❖ Most probable speed $\sqrt{\frac{2RT}{M}} = u_{\text{mps}}$

Relationship between three types of speeds

❖ Most probable: average : r.m.s. $\sqrt{2} : \sqrt{\frac{8}{\pi}} : \sqrt{3} \quad 1:1.13:1.22$

❖ $u_{\text{rms}} : u_{\text{av}} : u_{\text{mp}}$

❖ Average kinetic energy of a single molecule $\frac{3}{2} \frac{R}{N} T = \frac{3}{2} kT$

where, k Boltzman constant $1.3806 \times 10^{-16} \text{ erg deg}^{-1}$


❖ Total kinetic energy for **one mole** of a gas $\frac{3}{2} RT$.

❖ Kinetic energy of **n moles** of a gas $n \frac{3}{2} RT$

Maxwells speed distribution Law:

$$dN_u = 4 N \frac{M}{2 \pi RT}^{3/2} \exp(-Mu^2 / 2RT) u^2 du$$

$$= 4 N \frac{m}{2 \pi kT}^{3/2} \exp(-mu^2 / 2kT) u^2 du$$



Real Gases

Real gases: Gases which do not obey perfect gas laws over a wide range of pressure and temperature are called real or non-ideal gases. These deviations are due to the interactions of molecules with each other and volume occupied by gas molecules. The repulsive forces between molecules assist expansion and attractive forces assist compression.

❖ Deviation from ideal behaviour

The curve for the real gas has a tendency to coincide with that of an ideal gas at low pressures when the volume is large. At higher pressures, however deviations are observed.

❖ van der Waals equation of state

$$P \frac{an^2}{V^2} (V - nb) = nRT$$

a , b are van der Waals constants; different for each gas

unit of a atm L² mol⁻²; S.I. unit Pa m⁶ mol⁻²



unit of b L mol⁻¹; S.I. unit m³ mol⁻¹

greater the value of ' a ' more easily the gas is liquefiable;

greater the value of ' b ' greater the molecular size,

❖ Compressibility factor

$$Z = \frac{PV}{nRT} = \frac{\text{observed molar volume}}{\text{ideal molar volume}}$$



❖ Interpretation of deviation using van der Waals equation

(i) At low pressure $Z = \frac{PV}{RT} = 1 + \frac{a}{VRT}$

(ii) At high pressure $Z = \frac{PV}{RT} = 1 + \frac{Pb}{RT}$

(iii) At extremely low pressure $Z = \frac{PV}{RT} = 1; PV = nRT$

❖ Virial equation of state for 1 mole of gas

$$Z = \frac{PV}{RT} = 1 + B \frac{1}{V} + C \frac{1}{V^2} + D \frac{1}{V^3} + \dots$$

B second virial coefficient $b = \frac{a}{RT}$ gas dependent

C third virial coefficient b^2

- ❖ **Boyle temperature (T_B):** The temperature at which a real gas obeys Boyle's law (*i.e.*, behaves as an ideal gas) in low pressure range.

$$T_B = \frac{a}{bR}$$

- ❖ **Inversion temperature:** The temperature at which a real gas show no Joule-Thomson effect is called its inversion temperature (T_i)

$$T_i = 2T_B = \frac{2a}{bR}$$

In Joule-Thomson's effect, all gases except H_2 and He show cooling effect while H_2 and He show heating effect.

❖ **Critical constants**

- (i) **Critical temperature (T_c) :** T_c is the maximum temperature at which a gas can be liquefied, *i.e.*, the temperature above which a liquid can not exist.

$$T_c = \frac{8a}{27Rb}$$

- (ii) **Critical pressure (P_c) :** P_c is the minimum pressure required to cause liquefaction at the temperature T_c .

$$P_c = \frac{a}{27b^2}$$

- (iii) **Critical volume (V_c) :** V_c is the volume occupied by one mole of gas at critical temperature and critical pressure.

$$V_c = \frac{3b}{8}$$

- (iv) Compressibility factor under critical state of a gas

$$Z_c = \frac{P_c V_c}{RT_c}$$

Now substituting the values of P_c , V_c and T_c we get


$$Z_c = \frac{3}{8}$$

❖ **The law of corresponding states**

$$p_r = \frac{p}{p_c}, \quad T_r = \frac{T}{T_c} \quad \text{and} \quad V_r = \frac{V}{V_c}$$

$$(p_r - \frac{3}{V_r^2})(3V_r - 1) = \frac{8T_r}{T_c}$$





$$Z = \frac{pV_m}{RT} = \frac{(p_r p_c)(V_r V_c)}{R(T_r T_c)} = \frac{p_c V_c}{T_c} \cdot \frac{p_r V_r}{T_r} = \frac{3P_r V_r}{8T_r}$$









Level 1





- Which one of the following statements is not correct about the three states of matter *i.e.*, solid, liquid and gaseous ?
 - Molecules of a solid possess least energy whereas those of a gas possess highest energy
 - The density of solids is highest whereas that of gases is lowest
 - Gases and liquids possess definite volumes
 - Molecules of a solids possess vibratory motion
- Which of the following curve does not represent Boyle's law?







- A certain sample of gas has a volume of 0.2 litre measured at 1 atm pressure and 0°C. At the same pressure but at 273°C, its volume will be:
 - 0.4 litre
 - 0.8 litre
 - 27.8 litres
 - 55.6 litres
- Among the following curves, which is not according to Charle's law ?





- Initial temperature of an ideal gas is 75°C. At what temperature, the sample of neon gas would be heated to double its pressure, if the initial volume of gas is reduced by 15%?
 - 319°C
 - 592°C
 - 128°C
 - 60°C
- Which is correct curve for Charle's law, when the curve is plotted at 0.821 atm pressure for 10 mole ideal gas?





7. At constant volume, for a fixed number of moles of a gas, the pressure of the gas increases with increase in temperature due to:
- increase in the average molecular speed
 - decrease in rate of collision amongst molecules
 - increase in molecular attraction
 - decrease in mean free path
8. Which is not correct curve for Gay-lussac's law ?







9. Three flasks of equal volumes contain CH_4 , CO_2 and Cl_2 gases respectively. They will contain equal number of molecules if :
- the mass of all the gases is same
 - the mass of all the gas is same but temperature is different
 - temperature and pressure of all the flasks are same
 - temperature, pressure and masses same in the flasks
10. Which is **incorrect** curve for Boyle's law ?










11. "Equal volumes of all gases at the same temperature and pressure contain equal number of particles." This statement is a direct consequence of :
- (a) Avogadro's law (b) Charle's law
(c) ideal gas equation (d) law of partial pressure
12. A 2.24L cylinder of oxygen at 1 atm and 273 K is found to develop a leakage. When the leakage was plugged the pressure dropped to 570 mm of Hg. The number of moles of gas that escaped will be :
- (a) 0.025 (b) 0.050 (c) 0.075 (d) 0.09
13. Which of the following curve is correct for an ideal gas ?







14. In the equation of state of an ideal gas $PV = nRT$, the value of the universal gas constant is not correct :
- (a) $8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ (b) $0.0821 \text{ atm L mol}^{-1} \text{ K}^{-1}$
(c) $0.8314 \text{ bar L mol}^{-1} \text{ K}^{-1}$ (d) $2 \text{ cal mol}^{-1} \text{ K}^{-1}$
15. At 0°C and one atm pressure, a gas occupies 100 cc. If the pressure is increased to one and a half-time and temperature is increased by one-third of absolute temperature, then final volume of the gas will be:
- (a) 80 cc (b) 88.9 cc (c) 66.7 cc (d) 100 cc
16. 10 g of a gas at 1 atm and 273 K occupies 5 litres. The temperature at which the volume becomes double for the same mass of gas at the same pressure is:
- (a) 273 K (b) -273°C (c) 273°C (d) 546°C
17. Which of the following curve does not represent Gay-lusacc's law?



18. If densities of two gases are in the ratio 1 : 2 and their temperatures are in the ratio 2 : 1, then the ratio of their respective molar mass at certain pressure is:
 (a) 1 : 1 (b) 1 : 2 (c) 2 : 1 (d) 4 : 1
19. Two separate bulbs contain ideal gases A and B. The density of gas A is twice that of gas B. The molecular mass of A is half that of gas B. The two gases are at the same temperature. The ratio of the pressure of A to that of gas B is :
 (a) 2 (b) 1/2 (c) 4 (d) 1/4
20. Volume of the air that will be expelled from a vessel of 300 cm^3 when it is heated from 27°C to 37°C at the constant pressure will be:
 (a) 310 cm^3 (b) 290 cm^3 (c) 10 cm^3 (d) 37 cm^3
21. V versus T curves at constant pressure P_1 and P_2 for an ideal gas are shown in fig. Which is correct?
 (a) $P_1 > P_2$ (b) $P_1 < P_2$
 (c) $P_1 = P_2$ (d) All of these
- 
22. Two flasks A and B of 500 mL each are respectively filled with O_2 and SO_2 at 300 K and 1 atm. pressure. The flasks will contain:
 (a) the same number of atoms
 (b) the same number of molecules
 (c) more number of moles of molecules in flask A as compared to flask B
 (d) the same amount of gases
23. 2.8 g of a gas at 1 atm and 273 K occupies a volume of 2.24 litres. The gas can not be:
 (a) O_2 (b) CO (c) N_2 (d) C_2H_4
24. Five grams each of the following gases at 87°C and 750 mm pressure are taken. Which of them will have the least volume ?
 (a) HF (b) HCl (c) HBr (d) HI
25. At what pressure a quantity of gas will occupy a volume of 60 mL, if it occupies a volume of 100 mL at a pressure of 720 mm (while temperature is constant) :
 (a) 700 mm (b) 800 mm (c) 100 mm (d) 1200 mm
26. At 1 atm and 273 K the density of gas, whose molecular weight is 45, is:
 (a) 44.8 g/L (b) 11.4 g/L (c) 2 g/L (d) 3 g/L
27. A small bubble rises from the bottom of a lake, where the temperature and pressure are 8 °C and 6.0 atm, to the water's surface, where the temperature is 25°C and pressure is 1.0 atm. Calculate the final volume of the bubble if its initial volume was 2 mL.
 (a) 14 mL (b) 12.72 mL (c) 11.31 mL (d) 15 mL
28. Argon is an inert gas used in light bulbs to retard the vaporization of the filament. A certain light-bulb containing argon at 1.25 atm and 18°C is heated to 85°C at constant volume. Calculate its final pressure.
 (a) 1.53 atm (b) 1.25 atm (c) 1.35 atm (d) 2 atm
29. Calculate the volume of O_2 at 1 atm and 273 K required for the complete combustion of 2.64 L of acetylene (C_2H_2) at 1 atm and 273 K. $2\text{C}_2\text{H}_2(\text{g}) + 5\text{O}_2(\text{g}) \rightarrow 4\text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$
 (a) 3.6 L (b) 1.056 L (c) 6.6 L (d) 10 L

30. The density of O_2 (g) is maximum at :
 (a) STP (b) 273 K and 2 atm
 (c) 546 K and 1 atm (d) 546 K and 2 atm
31. At 27°C a sample of ammonia gas exerts a pressure of 5.3 atm. What is the pressure when the volume of the gas is reduced to one-tenth of the original value at the same temperature?
 (a) 0.53 atm (b) 5.3 atm (c) 53 atm (d) None of these
32. A certain amount of gas at 25°C and at a pressure of 0.80 atm is kept in a glass vessel. Suppose that the vessel can withstand a pressure of 2.0 atm. How high can you raise the temperature of the gas without bursting the vessel?
 (a) 745°C (b) 472°C (c) 500°C (d) None of these
33. Which one of these graphs for an ideal gas, the arrow indication is incorrectly marked ?



34. The pressure of sodium vapour in a 1.0 L container is 10 torr at 1000°C . How many atoms are in the container?
 (a) 9.7×10^{17} (b) 7.6×10^{19}
 (c) 4.2×10^{17} (d) 9.7×10^{19}
35. An ideal gaseous mixture of ethane (C_2H_6) and ethene (C_2H_4) occupies 28 litre at 1 atm and 273 K. The mixture reacts completely with 128 gm O_2 to produce CO_2 and H_2O . Mole fraction at C_2H_4 in the mixture is :
 (a) 0.6 (b) 0.4 (c) 0.5 (d) 0.8
36. A certain hydrate has the formula $MgSO_4 \cdot xH_2O$. A quantity of 54.2 g of the compound is heated in an oven to drive off the water. If the steam generated exerts a pressure of 24.8 atm in a 2.0 L container at 120°C , calculate x.
 (a) 2 (b) 5 (c) 6 (d) 7
37. Air entering the lungs ends up in tiny sacs called alveoli. From the alveoli, the oxygen diffuses into the blood. The average radius of the alveoli is 0.0050 cm and the air inside contains 14 per cent oxygen. Assuming that the pressure in the alveoli is 1.0 atm and the temperature is 37°C , calculate the number of oxygen molecules in one of the alveoli.
 (a) 6×10^{13} (b) 10^{24} (c) 1.7×10^{22} (d) 1.7×10^{12}

38. Nitrogen forms several gaseous oxides. One of them has a density of 1.33 g/L measured at 764 mmHg and 150°C. Write the formula of the compound.
(a) NO (b) N₂O (c) NO₂ (d) N₂O₅
39. Starting out on a trip into the mountains, you inflate the tires on your automobile to a recommended pressure of 3.21×10^5 Pa on a day when the temperature is 5.0 °C. You drive to the beach, where the temperature is 28.0°C. Assume that the volume of the tire has increased by 3%. What is the final pressure in the tyres?
(a) 350 Pa (b) 3500 Pa (c) 3.5×10^5 Pa (d) None of these
40. A compressed cylinder of gas contains 1.50×10^3 g of N₂ gas at a pressure of 2.0×10^7 Pa and a temperature of 17.1°C. What volume of gas has been released into the atmosphere if the final pressure in the cylinder is 1.80×10^5 Pa? Assume ideal behaviour and that the gas temperature is unchanged.
(a) 1264 L (b) 126 L (c) 12600 L (d) 45 L
41. A high-altitude balloon is filled with 1.41×10^4 L of hydrogen at a temperature of 21°C and a pressure of 745 torr. What is the volume of the balloon at a height of 20 km, where the temperature is -48°C and the pressure is 63.1 torr?
(a) 1.274×10^5 L (b) 1.66×10^5 L (c) 1.66×10^4 L (d) None of these
42. The atmospheric pressure on Mars is 0.61 kPa. What is the pressure in mm Hg?
(a) 0.63 (b) 4.6 (c) 6.3 (d) 3.2
43. The density of liquid gallium at 30°C is 6.095 g/mL. Because of its wide liquid range (30 to 2400°C), gallium is used as a barometer fluid at high temperature. What height (in cm) of gallium will be observed on a day when the mercury barometer reads 740 torr? (The density of mercury is 13.6 g/mL.).
(a) 322 (b) 285 (c) 165 (d) 210
44. A weather balloon is inflated with helium. The balloon has a volume of 100 m³ and it must be inflated to a pressure of 0.10 atm. If 50 L gas cylinders of helium at a pressure of 100 atm are used, how many cylinders are needed? Assume that the temperature is constant.
(a) 2 (b) 3 (c) 4 (d) 1
45. A balloon contains 14.0 L of air at 760 torr. What will be the volume of the balloon when it is taken to a depth of 10 ft. in a swimming pool? Assume that the temperature of the air and water are equal. (density : Hg = 13.6 g/mL.)
(a) 11.0 (b) 11.3 (c) 10 (d) 10.8
46. A 0.50 L container is occupied by nitrogen at a pressure of 800 torr and a temperature of 0°C. The container can only withstand a pressure of 3.0 atm. What is the highest temperature (°C) to which the container may be heated?
(a) 505 (b) 450 (c) 625 (d) 560
47. Equal volumes of oxygen gas and a second gas weigh 1.00 and 2.375 grams respectively under the same experimental conditions. Which of the following is the unknown gas?
(a) NO (b) SO₂ (c) CS₂ (d) CO
48. A high altitude balloon contains 6.81 g of helium in 1.16×10^4 L at -23°C. Assuming ideal gas behaviour, how many grams of helium would have to be added to increase the pressure to 4.0×10^3 atm?
(a) 1.27 g (b) 1.58 g (c) 2.68 g (d) 2.23 g

49. A 4.40 g piece of solid CO_2 (dry ice) is allowed to sublime in a balloon. The final volume of the balloon is 1.00 L at 300 K. What is the pressure (atm) of the gas?
 (a) 0.122 (b) 2.46 (c) 122 (d) 24.6
50. For a closed (not rigid) container containing $n = 10$ moles of an ideal gas fitted with movable, frictionless, weightless piston operating such that pressure of gas remains constant at 0.821 atm. Which graph represents correct variation of $\log V$ vs $\log T$ where V is in litre and T in kelvin.
- (a)

(b)
- (c)


(d)
51. The intercept on y-axis and slope of curve plotted between P/T vs. T (For an ideal gas having 10 moles in a closed rigid container of volume 8.21 L. (P = Pressure in atm and T = Temp. in K, $\log_{10} 2 = 0.30$)) are respectively :
 (a) 0.01, 0 (b) 0.1, 0 (c) 0.1, 1 (d) 10, 1
52. A He atom at 300 K is released from the surface of the earth to travel upwards. Assuming that constant temperature and $g = 9.81 \text{ ms}^{-2}$, how high will it be before coming to the rest?
 (a) 9.53 m (b) 95.3 m (c) 953 m (d) $9.53 \times 10^4 \text{ m}$
53. The density of gas A is twice that to B at the same temperature. The molecular weight of gas B is twice that of A. The ratio of pressure of gas A and B will be :
 (a) 1 : 6 (b) 1 : 1 (c) 4 : 1 (d) 1 : 4
54. Two inflated balloons I and II (thin skin) having volume 600 mL and 1500 mL at 300 K are taken as shown in diagram. If maximum volume of inner and outer balloons are 800 mL and 1800 mL respectively then find the balloon which will burst first on gradual heating.
- (a) inner balloon

(b) outer balloon


(c) both simultaneously

(d) unpredictable


55. An open flask containing air is heated from 300 K to 500 K. What percentage of air will be escaped to the atmosphere, if pressure is keeping constant?
 (a) 80 (b) 40 (c) 60 (d) 20
56. The value of universal gas constant R depends on :
 (a) temperature of gas (b) volume of gas
 (c) number of moles of gas (d) units of volume and pressure
57. A manometer attached to a flask contains with ammonia gas have no difference in mercury level initially as shown in diagram. After sparking into the flask, ammonia is partially dissociated as $2\text{NH}_3(\text{g}) \rightarrow \text{N}_2(\text{g}) + 3\text{H}_2(\text{g})$ now it have difference of 6 cm in mercury level in two columns, what is partial pressure of $\text{H}_2(\text{g})$ at equilibrium?



- (a) 9 cm Hg (b) 18 cm Hg
 (c) 27 cm Hg (d) None of these
58. An ideal gas is collected by downward displacement of water. Select the correct expression for P_{gas} according to the diagram [$d_{\text{Hg}} = 13.6 \text{ g/cm}^3$]:




- (a) $P_{\text{gas}} = P_{\text{atm}} + \frac{h}{13.6}$ (b) $P_{\text{gas}} = P_{\text{atm}} + h d_{\text{g}}$
 (c) $P_{\text{gas}} = P_{\text{atm}} + \frac{h}{13.6}$ (d) none of these
59. A bubble of gas released at the bottom of a lake increases to four times its original volume when it reaches the surface. Assuming that atmospheric pressure is equivalent to the pressure exerted by a column of water 10 m high, what is the depth of the lake?
 (a) 20 m (b) 10 m (c) 30 m (d) 40 m
60. Calculate the number of moles of gas present in the container of volume 10 L at 300 K, if the manometer containing glycerine shows 5 m difference in level as shown diagram.
 (Given : $d_{\text{glycerine}} = 2.72 \text{ g/mL}$; $d_{\text{mercury}} = 13.6 \text{ g/mL}$)



- (a) 0.94 mole (b) 0.49 mole (c) 0.64 mole (d) none of these
61. A rigid vessel of volume 0.50 m^3 containing H_2 at 20.5°C and a pressure of $611 \times 10^3 \text{ Pa}$ is connected to a second rigid vessel of volume 0.75 m^3 containing Ar at 31.2°C at a pressure of $433 \times 10^3 \text{ Pa}$. A valve separating the two vessels is opened and both are cooled to a temperature of 14.5°C . What is the final pressure in the vessels?
- (a) 2×10^5 (b) $3.22 \times 10^5 \text{ Pa}$
 (c) 4840 Pa (d) $4.84 \times 10^5 \text{ Pa}$
62. Two glass bulbs A and B at same temperature are connected by a very small tube having a stop-cock. Bulb A has a volume of 100 cm^3 and contained the gas while bulb B is empty. On opening the stop-cock, the pressure fell down to 20%. The volume of the bulb B is :
- (a) 100 cm^3 (b) 200 cm^3
 (c) 250 cm^3 (d) 400 cm^3
63. A mixture of C_2H_2 and C_3H_8 occupied a certain volume at 80 mm Hg. The mixture was completely burnt to CO_2 and $\text{H}_2\text{O}(l)$. When the pressure of CO_2 was found to be 230 mm Hg at the same temperature and volume, the mole fraction of C_3H_8 in the mixture is :
- (a) 0.125 (b) 0.875
 (c) 0.6 (d) 0.8
64. The total pressure of a mixture of oxygen and hydrogen is 1.0 atm. The mixture is ignited and the water is removed. The remaining gas is pure hydrogen and exerts a pressure of 0.40 atm when measured at the same values of T and V as the original mixture. What was the composition of the original mixture in mole fraction ?
- (a) $x_{\text{O}_2} = 0.2; x_{\text{H}_2} = 0.8$ (b) $x_{\text{O}_2} = 0.4; x_{\text{H}_2} = 0.6$
 (c) $x_{\text{O}_2} = 0.6; x_{\text{H}_2} = 0.4$ (d) $x_{\text{O}_2} = 0.8; x_{\text{H}_2} = 0.2$
65. Two closed vessel A and B of equal volume of 8.21 L are connected by a narrow tube of negligible volume with open valve. The left hand side container is found to contain 3 mole CO_2 and 2 mole of He at 400 K. What is the partial pressure of He in vessel B at 400 K?
- (a) 2.4 atm (b) 8 atm (c) 12 atm (d) None of these
66. At STP, a container has 1 mole of He, 2 mole Ne, 3 mole O_2 and 4 mole N_2 . Without changing total pressure if 2 mole of O_2 is removed. The partial pressure of O_2 will be decreased by :
- (a) 26% (b) 40%
 (c) 58.33% (d) 66.66%

67. 821 mL N_2 (g) was collected over liquid water at 300 K and 1 atm. If vapour pressure of H_2O is 30 torr then moles of N_2 (g) in moist gas mixture is :
 (a) 0.39 (b) 0.032
 (c) 0.96 (d) 0.0013
68. Let p and p_s be the partial pressure of H_2O (g) and vapour pressure of H_2O (l) respectively. Then the % relative humidity is given by:
 (a) $\frac{p_s}{p} \times 100$ (b) $\frac{p}{p_s} \times 100$ (c) $\frac{p_s}{p} \times 100$ (d) $(p - p_s) \times 100$
69. The vapour pressure of water at $80^\circ C$ is 355 mm of Hg. 1 L vessel contains O_2 at $80^\circ C$, which is saturated with water and the total pressure being 760 mm of Hg. The contents of the vessel were pumped into 0.3 L vessel at the same temperature. What is the partial pressure of O_2 ?
 (a) 1350 Hg (b) 2178.3 Hg (c) 121.5 Hg (d) 355 Hg
70. Which of the following gaseous mixture does not follow Dalton's law of partial pressure?
 (a) SO_2 and Cl_2 (b) CO_2 and N_2 (c) CO and CO_2 (d) CO and N_2
71. Equal masses of methane and oxygen are mixed in an empty container at $25^\circ C$. The fraction of the total pressure exerted by oxygen is:
 (a) $\frac{2}{3}$ (b) $\frac{1}{3} \times \frac{273}{298}$ (c) $\frac{1}{3}$ (d) $\frac{1}{2}$
72. A box of 1 L capacity is divided into two equal compartments by a thin partition which are filled with 2g H_2 and 16 g CH_4 respectively. The pressure in each compartment is recorded as P atm. The total pressure when partition is removed will be:
 (a) P (b) $2P$ (c) $P/2$ (d) $P/4$
73. If 10^{-4} dm^3 of water is introduced into a 1.0 dm^3 flask at 300 K, how many moles of water are in the vapour phase when equilibrium is established?
 (Given : Vapour pressure of H_2O at 300 K is 3170 Pa; $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$)
 (a) 1.27×10^{-3} mole (b) 5.56×10^{-3} mole
 (c) 1.53×10^{-2} mole (d) 4.46×10^{-2} mole
74. At room temperature Dalton's law of partial pressure is not applicable to :
 (a) H_2 and N_2 mixture (b) H_2 and Cl_2 mixture
 (c) H_2 and CO_2 mixture (d) none of these
75. 56 g of nitrogen and 96 g of oxygen are mixed isothermally and at a total pressure of 10 atm. The partial pressures of oxygen and nitrogen (in atm) are respectively :
 (a) 4, 6 (b) 5, 5 (c) 2, 8 (d) 6, 4
76. The closed containers of the same capacity and at the same temperature are filled with 44 g of H_2 in one and 44 g of CO_2 in the other. If the pressure of carbon dioxide in the second container is 1 atm, then pressure of hydrogen in the first container would be :
 (a) 1 atm (b) 10 atm (c) 22 atm (d) 44 atm
77. A jar contains a gas and a few drops of water. The pressure in the jar is 830 mm of Hg. The temperature of the jar is reduced by 1%. The vapour pressure of water at two temperatures are 30 and 25 mm of Hg. Calculate the new pressure in jar.
 (a) 792 mm of Hg (b) 817 mm of Hg
 (c) 800 mm of Hg (d) 840 mm of Hg

78. O_2 and SO_2 gases are filled in ratio of 1 : 3 by mass in a closed container of 3 L at temperature of $27^\circ C$. The partial pressure of O_2 is 0.60 atm, the concentration of SO_2 would be
(a) 0.36 mol/L (b) 0.036 mol/L (c) 3.6 mol/L (d) 36 mol/L
79. A gaseous mixture contains three gases A, B and C with a total number of moles of 10 and total pressure of 10 atm. The partial pressure of A and B are 3 atm and 1 atm respectively. If C has molecular weight of 2 g/mol. then, the weight of C present in the mixture will be :
(a) 8 g (b) 12 g (c) 3 g (d) 6 g
80. A rigid container contains 5 mole H_2 gas at some pressure and temperature. The gas has been allowed to escape by simple process from the container due to which pressure of the gas becomes half of its initial pressure and temperature become $(2/3)^{rd}$ of its initial. The mass of gas remaining is :
(a) 7.5 g (b) 1.5 g (c) 2.5 g (d) 3.5 g
81. Pressure of 1 g ideal gas X at 300 K is 2 atm. When 2 g of another gas Y is introduced in the same vessel at same temperature, the pressure become 3 atm. The correct relationship between molar mass of X and Y is :
(a) $M_Y = 2 M_X$ (b) $M_Y = 4 M_X$
(c) $M_X = 4 M_Y$ (d) None of these
82. Dry ice is solid carbon dioxide. A 0.050 g sample of dry ice is placed in an evacuated 4.6 L vessel at $30^\circ C$. Calculate the pressure inside the vessel after all the dry ice has been converted to CO_2 gas.
(a) 6.14 atm (b) 0.614 atm (c) 0.0614 atm (d) 6.14×10^{-3} atm
83. A mixture of helium and neon gases is collected over water at $28.0^\circ C$ and 745 mmHg. If the partial pressure of helium is 368 mmHg, what is the partial pressure of neon?
(Vapour pressure of water at $28^\circ C = 28.3$ mmHg)
(a) 348.7 mmHg (b) 377 mmHg (c) 384.7 mmHg (d) none of these
84. Consider the following apparatus. Calculate the partial pressure of helium after the opening valve. The temperature remains constant at $16^\circ C$.



- (a) 0.164 atm (b) 1.64 atm (c) 0.328 atm (d) 1 atm
85. Oxygen gas generated by the decomposition of potassium chlorate is collected over water. The volume of oxygen collected at $24^\circ C$ and atmospheric pressure of 760 mmHg is 128 mL. Calculate the mass of oxygen gas obtained. The vapour pressure of the water vapour at $24^\circ C$ is 22.4 mmHg.
(a) 1.36 g (b) 1.52 g (c) 0.163 g (d) 1.63 g

86. The quantity $\frac{PV}{k_B T}$ represents the (k_B : Boltzmann constant)
- (a) number of particles of the gas (b) mass of the gas
(c) number of moles of the gas (d) translation energy of the gas
87. Which of the following statements about kinetic energy (K.E.) is true?
- (a) All objects moving with the same velocity have the same K.E.
(b) The K.E. of a body will quadruple if its velocity doubles
(c) As the velocity of a body increases, its K.E. decreases
(d) The K.E. of a body is independent of its mass
88. The Ne atom has 10 times the mass of H_2 molecule. Which of the following statements is true?
- I. At $25^\circ C$ both of them have the same kinetic energy.
II. Ten moles of H_2 would have the same volume as 1 mole of Ne at same temp. and pressure.
III. One mole of Ne exerts the same pressure as one mole of H_2 at STP
IV. A H_2 molecule travels 10 times faster than Ne atom at same temperature.
V. At STP, one litre of Ne has 10 times the density of 1 litre of H_2 .
- (a) II, IV, V (b) I, III, V (c) I, II, III (d) I, II
89. Which of the following is NOT a postulate of the kinetic molecular theory of gases?
- (a) The gas molecules possess a volume that is negligibly small compared to the container
(b) The pressure and volume of a gas are inversely related
(c) Gases consist of discrete particles that are in constant chaotic motion
(d) The average kinetic energy of the molecules is directly proportional to the absolute temperature
90. Which one of the following relationships when graphed does not give a straight line for helium gas?
- I. K.E. and T at constant pressure and volume
II. P v/s V at constant temperature for a constant mass
III. V v/s $1/T$ at constant pressure for a constant mass
- (a) II (b) II and III (c) III (d) I
91. Consider three one-litre flasks labeled A, B and C filled with the gases NO , NO_2 , and N_2O , respectively, each at 1 atm and 273 K. In which flask do the molecules have the highest average kinetic energy?
- (a) Flask C (b) All are the same (c) Flask A (d) None
92. Which of the following statements is false?
- (a) The product of pressure and volume of fixed amount of a gas is independent of temperature.
(b) Molecules of different gases have the same kinetic energy at a given temperature.
(c) The gas equation is not valid at high pressure and low temperature.
(d) The gas constant per molecule is known as Boltzmann's constant.
93. Which is not correct in terms of kinetic theory of gases?
- (a) Gaseous particles are considered as point mass.
(b) The gaseous molecules are in random motion.
(c) When gaseous molecules collide, they lose energy.
(d) When the gas is heated, the molecules move faster.

- 94.** Two flasks *A* and *B* have equal volumes. *A* is maintained at 300 K and *B* at 600 K. *A* contains H_2 gas, *B* has an equal mass of CO_2 gas. Find the ratio of total K.E. of gases in flask *A* to that of *B*.
 (a) 1 : 2 (b) 11 : 1
 (c) 33 : 2 (d) 55 : 7
- 95.** Kinetic energy and pressure of a gas of unit volume are related as :
 (a) $P = \frac{2}{3}E$ (b) $P = \frac{3}{2}E$ (c) $P = \frac{E}{2}$ (d) $P = 2E$
- 96.** Two flask *A* and *B* of equal volumes maintained at temperature 300 K and 700 K contain equal mass of $He(g)$ and $N_2(g)$ respectively. What is the ratio of translational kinetic energy of gas in flask *A* to that of flask *B*?
 (a) 1 : 3 (b) 3 : 1 (c) 3 : 49 (d) None of these
- 97.** Which of the following change is observed occurs when a substance *X* is converted from liquid to vapour phase at the standard boiling point?
 I. Potential energy of the system decreases
 II. The distance between molecules increases
 III. The average kinetic energy of the molecules in both phases are equal
 (a) I only (b) II only
 (c) III only (d) II and III only
- 98.** A mixture of Ne and Ar kept in a closed vessel at 250 K has a total K.E. = 3 kJ. The total mass of Ne and Ar is 30 g. Find mass % of Ne in gaseous mixture at 250 K.
 (a) 61.63 (b) 38.37 (c) 50% (d) 28.3
- 99.** In two vessels of 1 litre each at the same temperature 1g of H_2 and 1g of CH_4 are taken. For these gases:
 (a) V_{rms} values will be same (b) Kinetic energy per mol will be same
 (c) Total kinetic energy will same (d) Pressure will be same
- 100.** Four particles have speed 2,3,4 and 5 cm/s respectively. Their rms speed is:
 (a) 3.5 cm/s (b) $(27/2)$ cm/s (c) $\sqrt{54}$ cm/s (d) $(\sqrt{54}/2)$ cm/s
- 101.** A gaseous mixture contains 4 molecules with a velocity of 6 cm sec⁻¹, 5 molecules with a velocity of 2 cm sec⁻¹ and 10 molecules with a velocity of 3 cm sec⁻¹. What is the rms speed of the gas:
 (a) 2.5 cm sec⁻¹ (b) 1.9 cm sec⁻¹ (c) 3.6 cm sec⁻¹ (d) 4.6 cm sec⁻¹
- 102.** The ratio between the root mean square velocity of H_2 at 50 K and that of O_2 at 800 K is:
 (a) 4 (b) 2 (c) 1 (d) 1/4
- 103.** If C_1, C_2, C_3 represent the speeds of n_1, n_2, n_3 molecules respectively, then the root mean square speed will be:
 (a) $\sqrt{\frac{n_1 C_1^2}{n_1} + \frac{n_2 C_2^2}{n_2} + \frac{n_3 C_3^2}{n_3}}$ (b) $\sqrt{\frac{(n_1 + n_2 + n_3)^2}{n_1 C_1^2 + n_2 C_2^2 + n_3 C_3^2}}$
 (c) $\sqrt{\frac{(n_1 C_1)}{n_1} + \frac{(n_2 C_2)}{n_2} + \frac{(n_3 C_3)}{n_3}}$ (d) $\sqrt{\frac{(n_1 C_1 + n_2 C_2 + n_3 C_3)^2}{n_1 + n_2 + n_3}}$
- 104.** The root mean square speed of hydrogen is $\sqrt{5}$ times than that of nitrogen. If *T* is the temperature of the gas, then :


(a) $T_{\text{H}_2} \quad T_{\text{N}_2}$

(c) $T_{\text{H}_2} \quad T_{\text{N}_2}$

(b) $T_{\text{H}_2} \quad T_{\text{N}_2}$

(d) $T_{\text{H}_2} \quad \sqrt{7} T_{\text{N}_2}$

- 105.** At a definite temperature (T), the distribution of speeds is given by the curve. In the curve points A, B and C indicates the speeds corresponding to :



- (a) most probable, average and root mean square speeds
 (b) average, root mean square and most probable speeds
 (c) root mean square, average and most probable speeds
 (d) most probable, root mean square and average speeds
- 106.** The most probable speed of 8 g of H_2 is 200 ms^{-1} . Average kinetic energy (neglect rotational and vibrational energy) of H_2 gas is :
 (a) 480 J (b) 240 J (c) 120 J (d) none of these
- 107.** At what temperature will average speed of the molecules of the second member of the series C_nH_{2n} be the same to that of Cl_2 at 627°C ?
 (a) 259.4 K (b) 400 K
 (c) 532.4 K (d) None of these
- 108.** If u_{rms} of a gas is $30 R^{1/2} \text{ ms}^{-1}$ at 27°C then the molar mass of gas is :
 (a) 0.02 kg/mol (b) 0.001 kg/mol
 (c) 0.003 kg/mol (d) 1 kg/mol
- 109.** 6×10^{22} gas molecules each of mass 10^{-24} kg are taken in a vessel of 10 litre. What is the pressure exerted by gas molecules? The root mean square speed of gas molecules is 100 m/s .
 (a) 20 Pa (b) $2 \times 10^4 \text{ Pa}$ (c) $2 \times 10^5 \text{ Pa}$ (d) $2 \times 10^7 \text{ Pa}$
- 110.** At what temperature will most probable speed of the molecules of the second member of homologous series $\text{C}_n\text{H}_{2n-2}$ be the same as that of SO_2 at 527°C .
 (a) 500°C (b) 727°C (c) 227°C (d) None of these
- 111.** The root mean square speed of 8 g of He is 300 ms^{-1} . Total kinetic energy of He gas is :
 (a) 120 J (b) 240 J (c) 360 J (d) None of these
- 112.** If T_1 , T_2 and T_3 are the temperatures at which the u_{rms} , u_{average} , u_{mp} of oxygen gas are all equal to 1500 m/s then the correct statement is :
 (a) $T_1 \quad T_2 \quad T_3$ (b) $T_1 \quad T_2 \quad T_3$
 (c) $T_1 \quad T_2 \quad T_3$ (d) None of these
- 113.** The density of a gas filled in electric lamp is 0.75 kg/m^3 . After the lamp has been switched on, the pressure in it increases from $4 \times 10^4 \text{ Pa}$ to $9 \times 10^4 \text{ Pa}$. What is increases in u_{rms} ?
 (a) 100 (b) 200 (c) 300 (d) None of these

- 114.** The root mean square speed of an ideal gas at constant pressure varies with density (d) as :
 (a) d^2 (b) d (c) $\sqrt{d}$ (d) $1/\sqrt{d}$
- 115.** The ratio among most probable speed, mean speed and root mean square speed is given by
 (a) $1 : 2 : 3$ (b) $1 : \sqrt{2} : \sqrt{3}$
 (c) $\sqrt{2} : \sqrt{3} : \sqrt{8}$ (d) $\sqrt{2} : \sqrt{8} : \sqrt{3}$
- 116.** The average speed at temperature $T^\circ\text{C}$ of $\text{CH}_4(g)$ is $\sqrt{\frac{28}{88}} \cdot 10^3 \text{ ms}^{-1}$. What is the value of T ?
 (a) 240.55°C (b) -32.45°C
 (c) 3000°C (d) -24.055°C
- 117.** At what temperature most probable speed of SO_2 molecules have the same value as root mean square speed of O_2 molecules at 300 K ?
 (a) 150 K (b) 600 K
 (c) 750 K (d) 900 K
- 118.** The rms speed of N_2 gas molecules is u . If the temperature is doubled and the nitrogen molecules dissociate into nitrogen atoms, the rms speed becomes:
 (a) $u/2$ (b) $2u$ (c) $4u$ (d) $14u$
- 119.** The rate of diffusion of a gas is proportional to:
 (a) $\frac{P}{\sqrt{M}}$ (b) $\frac{P}{M}$ (c) $\sqrt{\frac{P}{M}}$ (d) $\frac{\sqrt{P}}{M}$
- 120.** At constant volume and temperature conditions, the rates of diffusion r_A and r_B of gases A and B having densities ρ_A and ρ_B are related by the expression :
 (a) $r_A = r_B \cdot (\rho_A / \rho_B)^2$ (b) $r_A = r_B (\rho_B / \rho_A)^{1/2}$
 (c) $r_A = r_B \cdot (\rho_B / \rho_A)^{1/2}$ (d) $r_A = r_B (\rho_A / \rho_B)^{1/2}$
- 121.** What is the ratio of diffusion rate of oxygen to hydrogen?
 (a) $1 : 4$ (b) $4 : 1$ (c) $1 : 8$ (d) $8 : 1$
- 122.** The molecular weight of a gas which diffuses through a porous plug at $1/6^{\text{th}}$ of the speed of hydrogen under identical conditions is:
 (a) 27 (b) 72
 (c) 36 (d) 48
- 123.** $X \text{ mL}$ of H_2 gas effuses through a hole in a container in 5 seconds. The time taken for the effusion of the same volume of the gas specified below under identical condition is:
 (a) $10 \text{ sec} : \text{He}$ (b) $20 \text{ sec} : \text{O}_2$
 (c) $25 \text{ sec} : \text{CO}$ (d) $55 \text{ sec} : \text{CO}_2$
- 124.** At identical temperature and pressure, the rate of diffusion of hydrogen gas is $3\sqrt{3}$ times that of a hydrocarbon having molecular formula $\text{C}_n\text{H}_{2n-2}$. What is the value of n ?
 (a) 1 (b) 4 (c) 3 (d) 8
- 125.** Calculate relative rate of effusion of O_2 to CH_4 from a container containing O_2 and CH_4 in $3 : 2$ mass ratio.
 (a) $\frac{3\sqrt{2}}{4}$ (b) $\frac{3}{4\sqrt{2}}$ (c) $\frac{3}{\sqrt{2}}$ (d) none of these

- 126.** Calculate relative rates of effusion of SO_2 to CH_4 , in the mixture obtained by effusing out a mixture with initial molar ratio $\frac{n_{\text{SO}_2}}{n_{\text{CH}_4}} = \frac{8}{1}$ for three effusing steps.
- (a) 2 : 1 (b) 1 : 4
(c) 1 : 2 (d) none of these
- 127.** A gaseous mixture containing He, CH_4 and SO_2 was allowed to effuse through a fine hole. If mixture contain He, CH_4 and SO_2 in 1 : 2 : 3 mole ratio, calculate the molar ratio of gases effusing out initially.
- (a) 2 : 2 : 3 (b) 6 : 6 : 1 (c) $\sqrt{2} : \sqrt{2} : 3$ (d) 4 : 4 : 3
- 128.** 80 mL of O_2 takes 2 minute to pass through the hole. What volume of SO_2 will pass through the hole in 3 minute?
- (a) $\frac{120}{\sqrt{2}}$ mL (b) $120 \sqrt{2}$ mL (c) $\frac{12}{\sqrt{2}}$ mL (d) None of these
- 129.** When two cotton plugs, one moistened with ammonia and the other with hydrochloric acid, are simultaneously inserted into opposite ends of a glass tube 87.0 cm long, a white ring of NH_4Cl forms where gaseous NH_3 and gaseous HCl first come into contact.
- $\text{NH}_3(\text{g}) + \text{HCl}(\text{g}) \quad \quad \quad \text{NH}_4\text{Cl}(\text{s})$
- At what distance from the ammonia-moistened plug does this occur?
- (a) 51.7 cm from NH_3 end (b) 51.7 cm from HCl end
(c) 43.5 at mid point (d) None of these
- 130.** Dimethyl ether decomposes as
- $\text{CH}_3\text{OCH}_3(\text{g}) \quad \quad \quad \text{CH}_4(\text{g}) + \text{CO}(\text{g}) + \text{H}_2(\text{g})$
- When CH_3OCH_3 decomposes to 20% extent under certain conditions, what is the ratio of diffusion of pure CH_3OCH_3 with methane?
- (a) 0.59 : 1 (b) 1.18 : 1 (c) 2.36 : 1 (d) 1.77 : 1
- 131.** 4 gm of sulphur dioxide gas diffuses from a container in 8 min. Mass of helium gas diffusing from the same container over the same time interval is :
- (a) 0.5 g (b) 1 g (c) 2 g (d) None of these
- 132.** Under identical conditions of pressure and temperature, 4 L of gaseous mixture (H_2 and CH_4) effuses through a hole in 5 min whereas 4 L of a gas X of molecular mass 36 takes to 10 min to effuse through the same hole. The mole ratio of H_2 : CH_4 in the mixture is :
- (a) 1 : 2 (b) 2 : 1 (c) 2 : 3 (d) 1 : 1
- 133.** A balloon weighing 50 kg is filled with 685 kg of helium at 1 atm pressure and 25°C . What will be its pay load if it displaced 5108 kg of air?
- (a) 4373 kg (b) 4423 kg
(c) 5793 kg (d) none of these
- 134.** According to the law of equipartition of energy, the energy associated with each degree of freedom is :
- (a) $\frac{1}{3} K_B T$ (b) $\frac{1}{2} K_B T$
(c) $K_B T$ (d) $\frac{3}{2} K_B T$

- 135.** Calculate (ratio of C_p and C_v) for triatomic linear gas at high temperature. Assume that the contribution of vibrational degree of freedom is 75% :
- (a) 1.222 (b) 1.121
(c) 1.18 (d) 1.33
- 136.** If one mole each of a monoatomic and diatomic gases are mixed at low temperature then C_p/C_v ratio for the mixture is :
- (a) 1.40 (b) 1.428
(c) 1.5 (d) 1.33
- 137.** If one mole of a mono-atomic gas ($\gamma = 5/3$) is mixed with one mole of a diatomic gas ($\gamma = 7/5$), the value of γ for the mixture is :
- (a) 1.4 (b) 1.5
(c) 1.53 (d) 3.07
- 138.** Select the correct statement :
- (a) Internal energy of a real gas at a given temperature increases as the volume increases
(b) Internal energy of an ideal gas at given temperature increase as the volume increases
(c) Internal energy of an ideal gas molecules is not a function of temperature
(d) The internal energy of a real gas at a constant temperature is independent of change in volume
- 139.** Which gas shows real behaviour?
- (a) 16 g O_2 at 1 atm & 273 K occupies 11.2 L
(b) 1 g H_2 in 0.5 L flask exerts pressure of 24.63 atm at 300 K
(c) 1 mole NH_3 at 300 K and 1 atm occupies volume 22.4 L
(d) 5.6 L of CO_2 at 1 atm & 273 K is equal to 11 g
- 140.** Consider a real gas placed in a container. If the intermolecular attractions are supposed to disappear suddenly which of the following would happen?
- (a) Pressure decreases (b) Pressure increases
(c) Pressure remains unchanged (d) Gas collapses
- 141.** The pressure of a real gas is less than the pressure of an ideal gas because of :
- (a) increase in number of collisions (b) finite size of molecule
(c) increase in KE of molecules (d) intermolecular forces of attraction
- 142.** A gas behaves like an ideal gas at :
- (a) high pressure and low temperature (b) low pressure and high temperature
(c) high pressure and high temperature (d) low pressure and low temperature
- 143.** If temperature and volume are same, the pressure of a gas obeying van der Waal's equation is :
- (a) less than that of an ideal gas (b) more than that of an ideal gas
(c) same as that of an ideal gas (d) none of these
- 144.** 1 mole of each of X_1, X_2, X_3 with van der Waal's constants a (in $\text{atm L}^3 \text{mol}^{-2}$) 1.0, 3.8, 2.1 respectively is kept separately in three different vessels of equal volume at identical temperature. Their pressures are observed to P_1, P_2 , and P_3 respectively. On the basis of this data alone, select the correct option (neglect the effect of 'b'):
- (a) $P_1 > P_2 > P_3$ (b) $P_2 > P_1 > P_3$
(c) $P_2 > P_3 > P_1$ (d) $P_1 > P_2 > P_3$

- 145.** A gas obeys the equation of state $P(V - b) = RT$ (The parameter b is a constant). The slope for an isochore will be :
 (a) negative (b) zero
 (c) $R/(V - b)$ (d) R/P
- 146.** van der Waals constant b of helium is 24 mL mol^{-1} . Find molecular diameter of helium.
 (a) $1.335 \times 10^{-10} \text{ cm}$ (b) $1.335 \times 10^{-8} \text{ cm}$
 (c) $2.67 \times 10^{-8} \text{ cm}$ (d) $4.34 \times 10^{-8} \text{ cm}$
- 147.** Which of the following can be most readily liquefied?
 (Given: value of ' a ' for NH_3 $4.17 \text{ atm} \cdot \text{L}^3 \text{mol}^{-2}$, CO_2 $3.59 \text{ atm} \cdot \text{L}^3 \text{mol}^{-2}$, SO_2 $6.71 \text{ atm} \cdot \text{L}^3 \text{mol}^{-2}$, Cl_2 $6.49 \text{ atm} \cdot \text{L}^3 \text{mol}^{-2}$)
 (a) NH_3 (b) Cl_2
 (c) SO_2 (d) CO_2
- 148.** For which of the following gases should the correction for the molecular volume be largest : CO , CO_2 , NH_3 or SF_6 ?
 (a) CO (b) CO_2
 (c) NH_3 (d) SF_6
- 149.** Under which of the following sets of conditions is a real gas expected to deviate from ideal behaviour?
 (I) High pressure, small volume
 (II) High temperature, low pressure
 (III) Low temperature, high pressure
 (a) only I (b) only II (c) only III (d) I and III both
- 150.** For a certain gas which deviates a little from ideal behaviour, a plot between P/V vs P was found to be non-linear. The intercept on y-axis will be :
 (a) $\frac{RT}{M}$ (b) $\frac{M}{RT}$ (c) $\frac{MZ}{RT}$ (d) $\frac{R}{TM}$
- 151.** At low pressure, the van der Waals' equation become :
 (a) $PV_m = RT$ (b) $P(V_m - b) = RT$
 (c) $P = \frac{a}{V_m^2} + \frac{RT}{V_m}$ (d) $P = \frac{RT}{V_m} - \frac{a}{V_m^2}$
- 152.** At low pressure, if $RT = 2\sqrt{aP}$, then the volume occupied by a real gas is :
 (a) $\frac{2RT}{P}$ (b) $\frac{2P}{RT}$ (c) $\frac{RT}{2P}$ (d) $\frac{RT}{P}$
- 153.** For a gas deviation from ideal behaviour is maximum at :
 (a) 0°C and 1.0 atm (b) 100°C and 2.0 atm
 (c) -13°C and 1.0 atm (d) -13°C and 2.0 atm
- 154.** At low pressures, van der Waals' equation is written as $P = \frac{a}{V^2} + \frac{RT}{V}$. The compressibility factor is then equal to:
 (a) $1 - \frac{a}{RTV}$ (b) $1 + \frac{RTV}{a}$
 (c) $1 + \frac{a}{RTV}$ (d) $1 - \frac{RTV}{a}$

155. The compressibility factor for a real gas at high pressure is :

- (a) 1 (b) $1 \frac{Pb}{RT}$
 (c) $1 \frac{Pb}{RT}$ (d) $1 \frac{RT}{Pb}$

156. The compressibility of a gas is greater than unity at 1 atm and 273 K. Therefore :

- (a) V_m 22.4 L (b) V_m 22.4 L
 (c) V_m 22.4 L (d) V_m 44.8 L

157. At 273 K temperature and 9 atm pressure, the compressibility for a gas is 0.9. The volume of 1 milli-mole of gas at this temperature and pressure is :

- (a) 2.24 litre (b) 0.020 mL
 (c) 2.24 mL (d) 22.4 mL

158. The compressibility factor for nitrogen at 330 K and 800 atm is 1.90 and at 570 K and 200 atm is 1.10. A certain mass of N_2 occupies a volume of 1 dm^3 at 330 K and 800 atm. Calculate volume occupied by same quantity of N_2 gas at 570 K and 200 atm :

- (a) 1 L (b) 2 L
 (c) 3 L (d) 4 L

159. Consider the equation $Z = \frac{PV}{RT}$. Which of the following statements is correct?

- (a) When $Z > 1$, real gases are easier to compress than the ideal gas
 (b) When $Z < 1$, real gases get compressed easily
 (c) When $Z < 1$, real gases are difficult to compress
 (d) When $Z > 1$, real gases are difficult to compress

160. What is the compressibility factor (Z) for 0.02 mole of a van der Waals' gas at pressure of 0.1 atm. Assume the size of gas molecules is negligible.

Given : $RT = 20 \text{ L atm mol}^{-1}$ and $a = 1000 \text{ atm L}^2 \text{ mol}^{-2}$

- (a) 2 (b) 1
 (c) 0.02 (d) 0.5

161. Consider the following statements. If the van der Waal's parameters of two gases are given as

	$a/\text{dm}^6 \text{ bar mol}^{-2}$	$b/\text{dm}^3 \text{ mol}^{-1}$
Gas A	6.5	0.055
Gas B	2	0.01

then :

1. Critical volume of A > Critical volume of B
2. Critical pressure A > Critical pressure of B
3. Critical temperature of A > Critical temperature of B

Which of the above statements is/are incorrect?

- (a) 1 alone (b) 1 and 2
 (c) 1, 2 and 3 (d) 2 and 3

162. The van der Waals parameters for gases W, X, Y and Z are

Gas	a (atm L ² mol ⁻²)	b (L mol ⁻¹)
W	4.0	0.027
X	8.0	0.030
Y	6.0	0.032
Z	12.0	0.027

Which one of these gases has the highest critical temperature?

- (a) W (b) X (c) Y (d) Z

163. Pressure remaining the constant, the volume of a given mass of an ideal gas increases for every degree centigrade rise in temperature by definite fraction of its volume at:

- (a) 0°C (b) its critical temperature
(c) absolute zero (d) its Boyle temperature

164. The critical temperature of a substance is:

- (a) the temperature above which the substance undergoes decomposition
(b) the temperature above which a substance can exist only in gaseous state
(c) boiling point of the substance
(d) all are wrong

165. The excluded volume of a gas will be larger, if $\frac{T_C}{P_C}$ is :

- (a) small (b) large
(c) equal to 1 (d) less than unity

166. Select incorrect statement :

- (a) we can condense vapour simply by applying pressure
(b) to liquefy a gas one must lower the temperature below T_C and also apply pressure
(c) at T_C , there is no distinction between liquid and vapour state hence density of the liquid is nearly equal to density of the vapour
(d) However great the pressure applied, a gas cannot be liquified below its critical temp.

167. The correct order of temperature of a real gas is :

- (I) Boyle's temperature
(II) Critical temperature
(III) Inversion temperature
(a) III > I > II (b) I > II > III
(c) II > I > III (d) I > III > II

168. The temperature at which the second virial coefficient of real gas is zero is called :

- (a) Critical temperature (b) Triple point
(c) Boiling point (d) Boyle's temperature

169. The van der Waals' equation for one mole may be expressed as

$$V_m^3 - b \frac{RT}{P} V_m^2 - \frac{aV_m}{P} - \frac{ab}{P} = 0$$

where V_m is the molar volume of the gas. Which of the following is incorrect?

- (a) For a temperature less than T_C , V has three real roots
- (b) For a temperature less than T_C , V has three imaginary roots
- (c) For a temperature equal to T_C all three roots of V are real and identical
- (d) On increasing the temp. ($T > T_C$), the three roots become closer to one another

170. For a gas obeying the van der Waals' equation, at the critical point

- (a) both $\frac{P}{V_T}$ and $\frac{2P}{V^2_T}$ are zero
- (b) only $\frac{2P}{V^2_T}$ is not zero
- (c) $\frac{P}{V_T}$ is zero but $\frac{2P}{V^2_T}$ is non-zero
- (d) $\frac{P}{V_T}$ is non-zero but $\frac{2P}{V^2_T}$ is zero

171. However great the pressure, a gas cannot be liquified above its :

- (a) Boyle temperature
- (b) Inversion temperature
- (c) Critical temperature
- (d) Room temperature

172. The temperature at which real gases obey the ideal gas laws over a wide range of low pressure is called:

- (a) Critical temperature
- (b) Inversion temperature
- (c) Boyle temperature
- (d) Reduced temperature

173. Inversion temperature T_i $\frac{2a}{Rb}$ is defined as the temperature above which if a gas is expanded adiabatically it gets warmed up but if temperature of gas is lower than T_i then it will cool down. What will happen to a gas if it is adiabatically expanded at 600 K if its Boyle's temperature is 290 K?

- (a) Heating
- (b) Cooling
- (c) Constant
- (d) None

174. The van der Waals' equation of law of corresponding states for 1 mole of gas is :

- (a) $P_r \frac{3}{V_r^2} (3V_r - 1) - 8 T_r$
- (b) $P_r \frac{3}{V_r^2} (3V_r - 1) + 8 T_r$
- (c) $P_r \frac{3}{V_r^2} (3V_r - 1) + 8 T_r$
- (d) $P_r \frac{3}{V_r^2} (3V_r - 1) - 8$

175. Calculate the volume occupied by 16 gram O_2 at 300 K and 8.31 MPa if


$$\frac{P_c V_c}{RT_c} = \frac{3}{8} \text{ and } \frac{P_r V_r}{T_r} = 2.21 \text{ (Given : } R = 8.314 \text{ J/K-mol)}$$

- (a) 125.31 mL
- (b) 124.31 mL
- (c) 248.62 mL
- (d) none of these




Level 2

1. Consider the composite system, which is held at 300 K, shown in the following figure. Assuming ideal gas behavior, calculate the total pressure if the barriers separating the compartments are removed. Assume that the volume of the barriers is negligible. (Given : $R = 0.082 \text{ atm L/mol.K}$)



- (a) 1 atm (b) 2 atm (c) 2.3 atm (d) 3.2 atm
2. 11 moles of N_2 and 12 moles of H_2 mixture reacted in 20 litre vessel at 800 K. After equilibrium was reached, 6 mole of H_2 was present. 3.58 litre of liquid water is injected in equilibrium mixture and resultant gaseous mixture suddenly cooled to 300K. What is the final pressure of gaseous mixture? Neglect vapour pressure of liquid solution. Assume (i) all NH_3 dissolved in water (ii) no change in volume of liquid (iii) At 300 K no reaction takes place between N_2 and H_2




- (a) 18.47 atm (b) 60 atm (c) 22.5 atm (d) 45 atm
3. Two vessels connected by a valve of negligible volume. One container (I) has 2.8 g of N_2 at temperature T_1 (K). The other container (II) is completely evacuated. The container (I) is heated to T_2 (K) while container (II) is maintained at $T_2/3$ (K). volume of vessel (I) is half that of vessel (II). If the valve is opened then what is the mass ratio of N_2 in both vessel (W_I/W_{II})?
- (a) 1 : 2 (b) 1 : 3 (c) 1 : 6 (d) 3 : 1
4. A mixture of $\text{NH}_3(\text{g})$ and $\text{N}_2\text{H}_4(\text{g})$ is placed in a sealed container at 300 K. The total pressure is 0.5 atm. The container is heated to 1200 K at which time both substances decompose completely according to the equations



After decomposition is complete, the total pressure at 1200 K is found to be 4.5 atm. Find the mole% of N_2H_4 in the original mixture :

- (a) 20% (b) 25% (c) 50% (d) 75%

5. Correct expression for density of an ideal gas mixture of two gases 1 and 2, where m_1 and m_2 are masses and n_1 and n_2 are moles and M_1 and M_2 are molar masses.
- (a) $d = \frac{(m_1 + m_2)}{(M_1 + M_2)}$ (b) $d = \frac{(m_1 + m_2) P}{(n_1 + n_2) RT}$
- (c) $d = \frac{(n_1 + n_2) P}{(m_1 + m_2) RT}$ (d) None of these
6. Two closed vessel A and B of equal volume containing air at pressure P_1 and temperature T_1 are connected to each other through a narrow open tube. If the temperature of one is now maintained at T_1 and other at T_2 (where $T_1 > T_2$) then that what will be the final pressure?
- (a) $\frac{T_1}{2P_1T_2}$ (b) $\frac{2P_1T_2}{T_1 + T_2}$ (c) $\frac{2P_1T_1}{T_1 + T_2}$ (d) $\frac{2P_1}{T_1 + T_2}$
7. A balloon containing 1 mole air at 1 atm initially is filled further with air till pressure increases to 4 atm. The initial diameter of the balloon is 1 m and the pressure at each stage is proportional to diameter of the balloon. How many moles of air added to change the pressure from 1 atm to 4 atm.
- (a) 80 (b) 257 (c) 255 (d) 256
8. If Pd vs. P (where P denotes pressure in atm and d denotes density in gm/L) is plotted for He gas (assume ideal) at a particular temperature. If $\frac{d}{dP}(Pd)$ at $P = 8.21$ atm is 5, then the temperature will be
- (a) 160 K (b) 320 K (c) 80 K (d) none of these
9. What is the density of wet air with 75% relative humidity at 1 atm and 300 K? Given : vapour pressure of H_2O is 30 torr and average molar mass of air is 29 g mol^{-1} .
- (a) 1.614 g/L (b) 0.96 g/L (c) 1.06 g/L (d) 1.164 g/L
10. 7 moles of a tetra-atomic non-linear gas 'A' at 10 atm and T K are mixed with 6 moles of another gas B at $\frac{T}{3}$ K and 5 atm in a closed, rigid vessel without energy transfer with surroundings. If final temperature of mixture was $\frac{5T}{6}$ K, then gas B is? (Assuming all modes of energy are active)
- (a) monoatomic (b) diatomic (c) triatomic (d) tetra atomic
11. Three closed rigid vessels, A, B and C, which initially contain three different gases at different temperatures are connected by tube of negligible volume, without any energy transfer with surroundings. The vessel A contain 2 mole Ne gas, at 300 K, vessel 'B' contain 2 mole SO_2 gas at 400 K and vessel 'C' contain 3 mole CO_2 gas at temperature 500 K. What is the final pressure (in atm) attained by gases when all valves of connecting three vessels are opened and additional 15.6 kcal heat supplied to vessels through valve. The volume of A, B and C vessel is 2, 2 and 3 litre respectively



Given : $R = 2$ calorie/mol-K; C_v (Ne) $= \frac{3}{2} R$, C_v (CO) $= \frac{5}{2} R$ and C_v (SO₂) $= 3 R$

(a) 73.89 atm (b) 67.31 atm (c) 80 atm (d) none of these

12. Gas molecules each of mass 10^{-26} kg are taken in a container of volume 1 dm^3 . The root mean square speed of gas molecules is 1 km sec^{-1} . What is the temperature of gas molecules.

(Given : $N_A = 6 \times 10^{23}$; $R = 8 \text{ J/mol.K}$)

(a) 298 K (b) 25 K (c) 250 K (d) 2500 K

13. A balloon of diameter 21 meter weighs 100 kg. Calculate its pay-load, if it is filled with He at 1.0 atm and 27°C. Density of air is 1.2 kg m^{-3} . (Given : $R = 0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}$)

(a) 4952.42 kg (b) 4932.42 kg (c) 493.242 kg (d) none of these

14. A given volume of ozonised oxygen (containing 60% oxygen by volume) required 220 sec to effuse while an equal volume of oxygen took 200 sec only under identical conditions. If density of O₂ is 1.6 g/L then find density of O₃.


(a) 1.936 g/L (b) 2.16 g/L (c) 3.28 g/L (d) 2.24 g/L

15. If 250 mL of N₂ over water at 30°C and a total pressure of 740 torr is mixed with 300 mL of Ne over water at 25°C and a total pressure of 780 torr, what will be the total pressure if the mixture is in a 500 mL vessel over water at 35°C.

(Given : Vapour pressure (Aqueous tension) of H₂O at 25°C, 30°C and 35°C are 23.8, 31.8 and 42.2 torr respectively. Assume volume of H₂O(l) is negligible in final vessel)

(a) 760 torr (b) 828.4 torr (c) 807.6 torr (d) 870.6 torr

16. A bulb of constant volume is attached to a manometer tube open at other end as shown in figure. The manometer is filled with a liquid of density $(1/3)^{\text{rd}}$ that of mercury. Initially h was 228 cm.



Through a small hole in the bulb gas leaked assuming pressure decreases as $\frac{dP}{dt} = kP$.

If value of h is 114 cm after 14 minutes, what is the value of k (in hour^{-1})?

[Use : $\ln(4/3) = 0.28$ and density of Hg = 13.6 g/mL .]


(a) 0.6 (b) 1.2 (c) 2.4 (d) none of these

17. A mixture of nitrogen and water vapours is admitted to a flask at 760 torr which contains a sufficient solid drying agent. After long time the pressure attained a steady value of 722 torr.


If the experiment is done at 27°C and drying agent increases in mass by 0.9 gm, what is the volume of the flask? Neglect any possible vapour pressure of drying agent and volume occupied by drying agent.

(a) 443.34 L (b) 246.3 L (c) 12.315 L (d) 24.63 L

18. At room temperature following reaction goes to completion




AB_2 is solid with negligible vapour pressure below 0°C . At 300 K, the AB in the smaller flask exerts a pressure of 3 atm and in the larger flask B_2 exerts a pressure of 1 atm at 400 K when they are separated out by a close valve. The gases are mixed by opening the stop cock and after the end of the reaction the flask are cooled to 250 K



The final pressure is :

- (a) 0.156 atm (b) 0.3125 atm
(c) 0.625 atm (d) 3.2 atm
19. A vessel of uniform cross-section of length 500 cm as shown in figure is divided in two parts by a weightless and frictionless piston one part contains 5 moles of $\text{He}(g)$ and other part 2 moles of $\text{H}_2(g)$ and 4 moles of $\text{O}_2(g)$ added at the same temperature and pressure in which reaction takes place finally vessel cooled to 300 K and 1 atm. What is the length of He compartment? (Assume volume of piston and vol. of $\text{H}_2\text{O}(l)$ formed are negligible)



- (a) 187.5 cm (b) 300 cm (c) 312.5 cm (d) none of these
20. For a real gas (mol. mass = 60) if density at critical point is 0.80 g/cm^3 and its $T_c = \frac{4 \times 10^5}{821} \text{ K}$, then van der Waals' constant a (in $\text{atm L}^2 \text{ mol}^{-2}$) is
(a) 0.3375 (b) 3.375 (c) 1.68 (d) 0.025
21. The van der Waals' constant ' b ' of a gas is $4 \times 10^{-4} \text{ L/mol}$. How near can the centres of the two molecules approach each other? [Use : $N_A = 6 \times 10^{23}$]
(a) 10^{-7} m (b) 10^{-10} m (c) $5 \times 10^{-11} \text{ m}$ (d) $5 \times 10^{-9} \text{ m}$
22. The density of vapour of a substance (X) at 1 atm pressure and 500 K is 0.8 kg/m^3 . The vapour effuses through a small hole at a rate of $4/5$ times slower than oxygen under the same condition. What is the compressibility factor (Z) of the vapour?
(a) 0.974 (b) 1.35 (c) 1.52 (d) 1.22

23. van der Waal's gas equation can be reduced to virial equation and virial equation (in terms of volume) is $Z = A + \frac{B}{V_m} + \frac{C}{V_m^2} + \dots$

where A = first virial coefficient, B = second virial coefficient, C = third virial coefficient.

The third virial coefficient of Hg(g) is $625 \text{ (cm}^3/\text{mol)}^2$. What volume is available for movement of 10 moles He(g) atoms present in 50 L vessel?


- (a) 49.75 L (b) 49.25 L (c) 25 L (d) 50 L
24. If the slope of 'Z' (compressibility factor) vs. 'P' curve is constant slope $\frac{1}{492.6} \text{ atm}^{-1}$ at a particular temperature (300 K) and very high pressure, then calculate diameter of the molecules.

(Given : $N_A = 6.0 \times 10^{23}$, $R = 0.0821 \text{ atm.L mol}^{-1} \text{ K}^{-1}$)

- (a) $7.5 \text{ \AA}$ (b) $5 \text{ \AA}$ (c) $2.5 \text{ \AA}$ (d) $1.25 \text{ \AA}$
25. A graph is plotted between P (atm) vs t ($^\circ\text{C}$) for 10 mol of an ideal gas as follows:


Then slope of curve and volume of container (L) respectively, are:

- (a) 0.1, 8.21 (b) 8.21, 0.1
(c) 27.3, 8.21 (d) 8.21, 27.3



26. For two samples of ideal gases A and B curves are plotted n vs V (volume of container) at 16.42 atm pressure. Then temperature of A and B respectively are:

- (a) $\frac{200}{\sqrt{3}} \text{ K}$, $200\sqrt{3} \text{ K}$ (b) $\frac{200}{\sqrt{3}} \text{ } ^\circ\text{C}$, $(200\sqrt{3}) \text{ } ^\circ\text{C}$
(c) $200\sqrt{3} \text{ K}$, $\frac{200}{\sqrt{3}} \text{ K}$ (d) 200 K , $\frac{\sqrt{3}}{200} \text{ K}$



27. At a constant pressure, what should be the percentage increase in the temperature in kelvin for a 10% increase in volume:

- (a) 10% (b) 20% (c) 5% (d) 50%

28. 6 litre H_2O is placed in a closed room of volume 827 L at the temperature of 300 K. If vapour pressure of liquid water is 22.8 mm of Hg at 300 K and its density is 1 g/cm^3 :

[Given : $R = 0.0821 \text{ atm.L mol}^{-1} \text{ K}^{-1}$, Assuming volume of liquid water to be constant]

Column-I		Column-II	
(P)	Mass of H_2O in gaseous form (in g)	(1)	6
(Q)	Moles of H_2O in gaseous state	(2)	18
(R)	Approximate mass of water left in liquid state (in kg)	(3)	3
(S)	Total number of moles of all atoms in vapour form	(4)	1

Codes :

	P	Q	R	S
(a)	1	2	4	3
(b)	4	3	2	1
(c)	2	4	1	3
(d)	1	2	3	4

29. Match the items of columns I and II.

Column-I		Column-II	
(P)	Z for ideal gas behaviour	(1)	$3/8$
(Q)	Z for real gas at low pressure	(2)	$1 - \frac{Pb}{RT}$
(R)	Z for real gas at high pressure	(3)	1
(S)	Z for critical state	(4)	$1 - \frac{a}{RTV}$

Codes :

	P	Q	R	S
(a)	1	2	4	3
(b)	3	4	2	1
(c)	2	1	4	3
(d)	1	2	3	4

30.

List-I (Conditions for real gas)		List-II	
(P)	If force of attraction among gas particles are negligible	(1)	$PV_m - RT$
(Q)	At 1 atm and 273 K	(2)	$PV_m - RT - \frac{a}{V_m}$
(R)	If the volume of gas particles is negligible	(3)	$P - \frac{a}{V_m^2} (V_m - b) - RT$
(S)	At low pressure and high temperature	(4)	$PV_m - RT - Pb$

Codes :

	P	Q	R	S
(a)	4	1	3	2
(b)	4	3	2	1
(c)	2	1	4	3
(d)	1	2	3	4



Level 3



Passage 1

van der Waals' equation for calculating the pressure of a non ideal gas is

$$P = \frac{an^2}{V^2} (V - nb) - nRT$$

van der Waals' suggested that the pressure exerted by an ideal gas, P_{ideal} , is related to the experimentally measured pressure, P_{real} by the equation

$$P_{\text{ideal}} = P_{\text{real}} + \frac{an^2}{V^2}$$

observed correction
pressure term

Constant 'a' is measure of intermolecular interaction between gaseous molecules that gives rise to nonideal behavior. It depends upon how frequently any two molecules approach each other closely. Another correction concerns the volume occupied by the gas molecules. In the ideal gas equation, V represents the volume of the container. However, each molecule does occupy a finite, although small, intrinsic volume, so the effective volume of the gas becomes $(V - nb)$, where n is the number of moles of the gas and b is a constant. The term nb represents the volume occupied by gas particles present in n moles of the gas.

Having taken into account the corrections for pressure and volume, we can rewrite the ideal gas equation as follows :

$$P = \frac{an^2}{V^2} (V - nb) - nRT$$

corrected corrected
pressure volume

- At relatively high pressures, the van der Waals' equation of state reduces to
 (a) $PV = RT + a/V_m$ (b) $PV = aRT/V_m^2$ (c) $P = RT + a/V_m^2$ (d) $PV_m = RT + Pb$
- For non-zero value of force of attraction between gas molecules at large volume, gas equation will be :
 (a) $PV = nRT + \frac{n^2a}{V}$ (b) $PV = nRT - nbP$ (c) $P = \frac{nRT}{V - b}$ (d) $PV = nRT$
- The van der Waals' constant 'a' for CO_2 gas is greater than that of H_2 gas. Its mean that the
 (a) strength of van der Waals' force of CO_2 gas is less than that of H_2 gas
 (b) strength of van der Waals' force of CO_2 gas is equal to that of H_2 gas
 (c) CO_2 gas can be more easily liquified
 (d) H_2 gas can be more easily liquified


4. Using van der Waals' equation, find the constant ' a ' (in $\text{atm L}^2 \text{mol}^{-2}$) when two moles of a gas confined in 4 L flask exerts a pressure of 11.0 atmospheres at a temperature of 300 K. The value of b is 0.05 L mol^{-1} . ($R = 0.082 \text{ atm.L/K mol}$)
- (a) 2.62 (b) 2.64 (c) 6.24 (d) 6.46



Passage 2

Collision cross-section is an area of an imaginary sphere of radius r around the molecule within which the centre of another molecule cannot penetrate.

The volume swept by a single molecule in unit time is



$$V = (\pi r^2 \bar{u}) \quad \text{where } \bar{u} \text{ is the average speed}$$


If N^* is the number of molecules per unit volume, then the number of molecules within the volume V is

$$N = V N^* = (\pi r^2 \bar{u}) N^*$$

Hence, the number of collision made by a single molecule in unit time will be

$$Z = N = (\pi r^2 \bar{u}) N^*$$

In order to account for the movements of all molecules, we must consider the average velocity along the line of centres of two colliding molecules instead of the average velocity of a single molecule. If it is assumed that, on an average, molecules collide while approaching each other perpendicularly, then the average velocity along their centres is $\sqrt{2} \bar{u}$ as shown below.



Number of collision made by a single molecule with other molecules per unit time is given by

$$Z_1 = \pi r^2 (\bar{u}_{\text{rel}}) N^* = \sqrt{2} \pi r^2 \bar{u} N^*$$

The total number of bimolecular collisions Z_{11} per unit volume per unit time is given by

$$Z_{11} = \frac{1}{2} (Z_1 N^*) \text{ or } Z_{11} = \frac{1}{2} (\sqrt{2} \pi r^2 \bar{u} N^*) N^* = \frac{1}{\sqrt{2}} \pi r^2 \bar{u} N^{*2}$$

If the collision involve two unlike molecules then the number of collisions Z_{12} per unit volume per unit time is given as

$$Z_{12} = \frac{1}{2} \sqrt{\frac{8kT}{\pi}} N_1 N_2$$

where N_1 and N_2 are the number of molecules per unit volume of the two types of molecules, d_{12} is the average diameter of the two molecules and μ is the reduced mass. The mean free path is the average distance travelled by a molecule between two successive collisions. We can express it as follows :

$$\frac{\text{Average distance travelled per unit time}}{\text{No. of collisions made by a single molecule per unit time}} = \frac{\bar{u}}{Z_1}$$

or


$$\frac{\bar{u}}{\sqrt{2} \cdot d_{12}^2 N^*} = \frac{1}{\sqrt{2} \cdot d_{12}^2 N^*}$$

1. For a given gas the mean free path at a particular pressure is :
 - (a) independent to temperature
 - (b) decreases with rise in temperature
 - (c) increases with rise in temperature
 - (d) directly proportional to T^2
 - 2-3 Three ideal gas samples in separate equal volume containers are taken and following data is given :
- | | Pressure | Temperature | Mean free paths | Mol. wt. |
|-------|----------|-------------|-----------------|----------|
| Gas A | 1 atm | 1600 K | 0.16 nm | 20 |
| Gas B | 2 atm | 200 K | 0.16 nm | 40 |
| Gas C | 4 atm | 400 K | 0.04 nm | 80 |
2. Calculate ratio of collision frequencies (Z_{11}) (A : B : C) of following for the three gases.
 - (a) 1 : 2 : 4
 - (b) 4 : 2 : 1
 - (c) 1 : 4 : 16
 - (d) 16 : 4 : 1
 3. Calculate number of collision by one molecule per sec (Z_1).
 - (a) 4 : 1 : 4
 - (b) 1 : 4 : 4
 - (c) 4 : 3 : 2
 - (d) 1 : 2 : 4



Passage 3

A manometer contains a liquid of density 5.44 g/cm^3 is attached to a flask containing gas 'A' as follows




- If the same liquid is used in barometer to measure the atmospheric pressure, then what will be the length of the liquid column, which exerts pressure equal to 1 atm ? (density of Hg = 13.6 g/cm^3)
 (a) 190 cm (b) 76 cm (c) 30.4 cm (d) 266 cm
- The initial pressure of gas A in the flask is:
 (a) 1.5 atm (b) 1 atm (c) 1.3 atm (d) 1.2 atm
- If gas A undergoes 30% trimerisation $[3A(g) \rightleftharpoons A_3(g)]$ then the difference in height of the liquid level in two columns is:
 (a) 38 cm (b) 7.6 cm (c) 3.04 cm (d) 15.1 cm


One or More Answers is/are Correct



- Which of the following curves represent(s) Boyle's law?




- If a gas expands at constant temperature:
 (a) the pressure decreases
 (b) the kinetic energy of the molecules remains the same
 (c) the K.E. of the molecules decrease
 (d) the number of molecules of the gas increase
- Which of the following statements are correct?
 (a) It is not possible to compress a gas at a temperature below T_C
 (b) At a temperature below T_C , the molecules are close enough for the attractive forces to act, and condensation occurs
 (c) No condensation takes place above T_C
 (d) Due to higher kinetic energy of the gas molecules above T_C , it is considered as super critical fluid
- What conclusion would you draw from the following graphs for an ideal gas?




- As the temperature is reduced, the volume as well as the pressure increase
- As the temperature is reduced, the volume becomes zero and the pressure reaches infinity

- (c) As the temperature is reduced, the pressure decrease
(d) A point is reached where, theoretically, the volume become zero
5. At Boyle temperature:
- (a) the effects of the repulsive and attractive intermolecular forces just offset each other
(b) the repulsive intermolecular forces are stronger than the attractive intermolecular forces
(c) the repulsive intermolecular forces are weaker than the attractive intermolecular forces
(d) $b \frac{a}{RT} = 0$
6. Indicate the correct statement for equal volumes of $N_2(g)$ and $CO_2(g)$ at $25^\circ C$ and 1 atm.
- (a) The average translational K.E. per molecule is the same for N_2 and CO_2
(b) The rms speed is same for both N_2 and CO_2
(c) The density of N_2 is less than that of CO_2
(d) The total translational K.E. of both N_2 and CO_2 is the same
7. Which of the following is correct for critical temperature?
- (a) It is the highest temperature at which liquid and vapour can coexist
(b) Beyond this temperature, there is no distinction between the two phases and a gas cannot be liquefied by compression.
(c) At this temperature, the gas and the liquid phases have different critical densities
(d) All are correct
8. Consider the following statement regarding Maxwell's distribution of speeds. The correct statement(s) is/are :
- (a) as temperature increases, the peak (maxima) of a curve is shifted towards right side
(b) as temperature increases, the most probable speed of molecules increases but fraction of molecules of maximum speed decreases
(c) the area under the curve at all the temperatures is the same because it represents the number of gaseous molecules
(d) the fractions of molecules having different speeds are different at a given temperature
9. If a gas expands at a constant pressure by providing heat :
- (a) the temperature increases
(b) the kinetic energy of the gaseous molecules remains same
(c) the kinetic energy of gaseous molecules decreases
(d) the number of molecules of the gas decreases
10. Select the incorrect statement(s) :
- (a) The product of pressure and volume of fixed amount of a gas is independent of temperature
(b) The value of universal gas constant depends upon temperature, volume and number of gaseous molecules
(c) The gas constant also known as Boltzmann's constant
(d) The average kinetic energy of molecules depends only on temperature
11. Following represents the Maxwell distribution curve for an ideal gas at two temperature T_1 and T_2 . Which of the following option(s) is/are true?



- (a) Total area under the two curves is independent to moles of gas
 (b) u_{mp} decreases as temperature decreases
 (c) $T_1 > T_2$ and hence higher the temperature, sharper the curve
 (d) The fraction of molecules having speed u_{mp} decreases as temperature increases
- 12.** Two container each containing liquid water are connected as shown in diagram.




Given that vapour pressure of $H_2O(l)$ at 300 K and 350 K are 22 torr and 40 torr, select correct statement(s) :

- (a) The final pressure in each container if valve is opened while keeping the containers at the given temperature is 22 torr
 (b) The final pressure in each container if valve is opened while keeping the containers at the given temperature is 40 torr
 (c) Mass of $H_2O(l)$ is decreased in container X
 (d) Mass of $H_2O(l)$ is decreased in container Y
- 13.** Select the correct statement(s) :
- (a) At Boyle's temperature a real gas behaves like an ideal gas at low pressure
 (b) Above critical conditions, a real gas is called super critical fluid
 (c) For hydrogen gas 'b' dominates over 'a' at all temperatures
 (d) At high pressure van der Waals' constant 'b' dominates over 'a'
- 14.** Select the correct statement :
- (a) The value of compressibility factor 'Z' for H_2 gas is greater than one at room temperature and pressure
 (b) The real gas behaves as an ideal gas at Boyle's temperature.

(c) For a real gas following van der Waals' equation of state, the expression of critical temperature is $\frac{8a}{27R.b}$

(d) At low pressure, the compressibility factor ' Z ' is $1 - \frac{P.b}{RT}$ for a van der Waals' gas.

15. A open ended mercury manometer is used to measure the pressure exerted by a trapped gas as shown in the figure. Initially manometer shows no difference in mercury level in both columns as shown in diagram.



After sparking 'A' dissociates according to following reaction



If pressure of Gas "A" decreases to 0.9 atm, then (Assume temperature to be constant and is 300 K)

- total pressure increases to 1.3 atm
 - total pressure increases by 0.3 atm
 - total pressure increases by 22.3 cm of Hg
 - difference in mercury level is 228 mm.
16. Select incorrect statements for real gases:
- In low pressure region repulsive forces dominates
 - Volume of gas particles is not negligible in low pressure region
 - Gases behaves as an ideal gas at low pressure and low temperature
 - In high pressure region attractive forces dominates
17. Select correct statements:
- A real gas can be liquified at critical temperature
 - Critical pressure is the maximum pressure at which a substance is present in its liquid state at T_C
 - Ideal gas can be liquified below T_C
 - Critical volume is the molar volume of substance in gaseous state at T_C and P_C
18. Which is/are correct for real gases?
- $\lim_{P \rightarrow 0} (PV_m)$ constant at constant high temperature
 - $\lim_{V_m \rightarrow 0} (PV_m)$ constant at constant low temperature

(c) $\lim_{P \rightarrow 0} \frac{PV_m}{RT} = 1$ at high temperature

(d) $\lim_{V \rightarrow 0} \frac{PV_m}{RT} = R$

19. Select incorrect statement(s)

- (a) At very low pressure real gases show minimum deviation from ideal behaviour.
- (b) The compressibility factor for an ideal gas is zero.
- (c) At Boyle temperature real gas behave as ideal gas in high pressure region.
- (d) Real gas show maximum deviation at high pressure and low temperature.

20. If an ideal gas is heated at constant pressure:

- (a) the volume increases
- (b) the mass of gas remains same
- (c) the kinetic energy of the molecules increases
- (d) attraction forces between gas particles increases

Match the Column



Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column I

- (A) Boyle's temperature
- (B) Inversion temperature
- (C) Critical temperature
- (D) Critical pressure

Column II

- (P) a/Rb
- (Q) $8a/27Rb$
- (R) $2a/Rb$
- (S) $a/27b^2$

2.

Column I

- (A) Root mean square speed
- (B) Most probable speed
- (C) Average speed
- (D) K.E. of gas mol⁻¹

Column II

- (P) $\frac{3}{2}RT$
- (Q) $\sqrt{\frac{8P}{d}}$
- (R) $\sqrt{\frac{3P}{d}}$
- (S) $\sqrt{\frac{2RT}{M}}$

3.

Column I

- (A) If force of attraction among the gas molecules be negligible
 (B) If the volume of the gas molecules be negligible
 (C) At STP
 (D) At low pressure and at high temperature

Column II

- (P) $P \propto \frac{a}{V^2} (V - b) RT$
 (Q) $PV \propto RT \frac{a}{V}$
 (R) $PV \propto RT - Pb$
 (S) $PV \propto RT$

4. Match gases under specified conditions listed in Column I with their properties/laws in Column II.

Column I

- (A) Hydrogen gas ($P = 200 \text{ atm}, T = 273 \text{ K}$)
 (B) Hydrogen gas ($P \sim 0, T = 273 \text{ K}$)
 (C) CO_2 ($P = 1 \text{ atm}, T = 273 \text{ K}$)
 (D) Real gas with very large molar volume

Column II

- (P) Compressibility factor < 1
 (Q) Attractive forces are dominant
 (R) $PV \propto nRT$
 (S) $P(V - nb) \propto nRT$


5.

Column I


- (A) $\frac{1}{V^2}$ vs. P for ideal gas at constant T and n .
 (B) V vs. $\frac{1}{T}$ for ideal gas at constant P and n
 (C) PT vs. T^2 for ideal gas at constant V and n .
 (D) V vs. $\frac{1}{P^2}$ for ideal gas at constant T and n .

Column II


(P)




(Q)




(R)



(S)



6. Match the correct column from list-1 to list-2 on the basis of following Andrew's isotherm of real gas.


**Column-I**

- (A) Substance exist in both liquid and gas state
 (B) Only liquid state exist
 (C) Substance exist in gas state only
 (D) Real gas is called super critical fluid

Column-II

- (P) At AB part
 (Q) At BD part
 (R) At DE part
 (S) At point C
 (T) At GF curve

7.

**Column-I**

- (A) Speed at point 'A' is
 (B) Speed at point 'B' is
 (C) Speed at point 'C' is
 (D) The speed possessed by maximum fraction of the gas particles

Column-II

- (P) $\sqrt{\frac{8RT}{M}}$
 (Q) $\sqrt{\frac{3RT}{M}}$
 (R) U_{rms}
 (S) $\sqrt{\frac{2RT}{M}}$

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below

- (A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE


1. **STATEMENT-1 :** The heat absorbed during the isothermal expansion of an ideal gas against vacuum is zero.
STATEMENT-2 : The volume occupied by the molecules of an ideal gas is zero.
2. **STATEMENT-1 :** A lighter gas diffuses more rapidly than a heavier gas.
STATEMENT-2 : At a given temperature, the rate of diffusion of a gas is inversely proportional to the density.
3. **STATEMENT-1 :** The value of the van der Waals' constant ' a ' is larger for ammonia than for nitrogen.
STATEMENT-2 : Hydrogen bonding is present in ammonia.
4. **STATEMENT-1 :** Helium shows only positive deviation from ideal behaviour at room temperature.
STATEMENT-2 : Helium is an inert gas.
5. **STATEMENT-1 :** CH_4 , CO_2 has value of Z (compressibility factor) less than one at 0°C .
STATEMENT-2 : $Z < 1$ is due to the attractive forces dominate among the molecules.
6. **STATEMENT-1 :** The Joule-Thomson coefficient for an ideal gas is zero.
STATEMENT-2 : There are no intermolecular attractive forces in an ideal gas.
7. **STATEMENT-1 :** The average translational kinetic energy per molecule of the gas per degree of freedom is $\frac{1}{2} kT$.
STATEMENT-2 : For every molecule there are three rotational degree of freedom.
8. **STATEMENT-1 :** On increasing the temperature, the height of the peak of the Maxwell distribution curve decreases.
STATEMENT-2 : The fraction of molecules is very less at the higher speeds.
9. **STATEMENT-1 :** The gases He and H_2 are very different in their behaviour at any temperature and pressure but their compressibility factors are nearly the same at the critical point.
STATEMENT-2 : They have nearly the same critical constant.

- 10. STATEMENT-1 :** Most probable speed is the speed possessed by maximum fraction of molecules at the same temperature.
- STATEMENT-2 :** On collision, more and more molecules acquire higher speed at the same temperature.
- 11. STATEMENT-1 :** Plot of P vs $1/V$ (volume) is a straight line for an ideal gas.
- STATEMENT-2 :** Pressure is directly proportional to volume for an ideal gas.
- 12. STATEMENT-1 :** 1 mol of H_2 and O_2 each occupy 22.7 L of volume at $0^\circ C$ and 1 bar pressure, considering ideal behaviour.
- STATEMENT-2 :** Molar volume for all ideal gases at the same temperature and pressure are equal.
- 13. STATEMENT-1 :** Reacting gases react to form a new gas having pressure equal to the sum of their partial pressure.
- STATEMENT-2 :** Pressure exerted by a mixture of non-reacting gases present in a container is equal to the sum of their partial pressures.
- 14. STATEMENT-1 :** $1/4^{th}$ of the initial mole of the air is expelled, if air present in an open vessel is heated from $27^\circ C$ to $127^\circ C$.
- STATEMENT-2 :** Rate of diffusion of a gas is inversely proportional to the square root of its molecular mass.
- 15. STATEMENT-1 :** Compressibility factor for hydrogen varies with pressure with positive slope at all pressures at $0^\circ C$.
- STATEMENT-2 :** Even at low pressures, repulsive forces dominate in hydrogen gas at $0^\circ C$.
- 16. STATEMENT-1 :** Wet air is heavier than dry air.
- STATEMENT-2 :** The density of dry air is less than the density of water at 1 atm and 273K.

Subjective Problems




- Collapsible balloon is inflated to a volume of 10L at a pressure of 1 atm. When the balloon is immersed to the bottom of a lake, its volume reduce to 1.25L. Assuming atmospheric pressure to be equivalent to 10m column of water and no change in temperature, if the depth of the lake is $x \cdot 10^3$ cm, what is the value of x ?
- A gaseous mixture containing equal moles of H_2 , O_2 and He is subjected to series of effusion steps. The composition (by moles) of effused mixture after 4 effusion steps is $x:1:y$ respectively. Then find the value of $\frac{x}{y}$.
- One mole of a gas changed from its initial state (15L, 2atm) to final state (4L, 10atm) reversibly. If this change can be represented by a straight line in $P-V$ curve maximum temperature (approximate), the gas attained is $x \cdot 10^2 K$. Then find the value of x .
- Two moles of an ideal gas undergoes the following process. Given that $\frac{P}{T}$ is $x \cdot 10^{-y}$, then calculate the value of $(x - y)$




5. 1 mole of a diatomic gas present in 10 L vessel at certain temperature exert a pressure of 0.96 atm. Under similar conditions an ideal gas exerted 1.0 atm pressure. If volume of gas molecule is negligible, then find the value of van der Waals' constant " a " (in $\text{atm L}^2/\text{mol}^2$).
6. The graph of compressibility factor (Z) vs. P for one mole of a real gas is shown in following diagram. The graph is plotted at constant temperature 273 K. If the slope of graph at very high pressure $\frac{dZ}{dP}$ is $\frac{1}{2.8} \text{ atm}^{-1}$, then calculate volume of one mole of real gas molecules (in L/mol)

Given : $N_A = 6 \times 10^{23}$ and $R = \frac{22.4}{273} \text{ L atm K}^{-1} \text{ mol}^{-1}$




7. Under the identical conditions of temperature, the density of a gas X is two times to that of gas Y while molecular mass of gas Y is three times that of X . Calculate the ratio of pressure of X and Y .
8. The time taken for a certain volume of a certain gas to diffuse through a small hole was 2 min. Under similar conditions an equal volume of oxygen took 5.65 min to pass. What is the molecular mass of gas (in amu) ?
9. Excess $\text{F}_2(\text{g})$ reacts at 150°C and 1.0 atm pressure with $\text{Br}_2(\text{g})$ to give a compound BrF_n . If 423 mL of $\text{Br}_2(\text{g})$ at the same temperature and pressure produced 4.2 g of BrF_n , what is n ? [Atomic mass Br = 80, F = 19]
10. Initially bulb "A" contained oxygen gas at 27°C and 950 mm of Hg and bulb "B" contained neon gas at 27°C and 900 mm of Hg. These bulbs are connected by a narrow tube of negligible volume equipped with a stopcock and gases were allowed to mix-up freely. The pressure obtained in the combined system was found to be 910 mm of Hg.



If volume of bulb B was measured to be 10 L, then find the volume of oxygen gas present initially in bulb "A".

11. Air is trapped in a horizontal glass tube by 36 cm mercury column as shown below :



If the tube is held vertical keeping the open end up, length of air column shrink to 19 cm. What is the length (in cm) by which the mercury column shifts down?

12. A flask containing air at 107°C and 722 mm of Hg is cooled to 100 K and 760 mm of Hg. If density in the initial condition 1 g/cm^3 , then what is the final density (g/cm^3)?
13. If an ideal gas at 100 K is heated to 109 K in a rigid container, the pressure increases by $X\%$. What is the value of X ?
14. The van der Waal's constants for a gas are $a = 3.6 \text{ atm L}^2 \text{ mol}^{-2}$, $b = 0.6 \text{ L mol}^{-1}$. If $R = 0.08 \text{ L atm K}^{-1} \text{ mol}^{-1}$ and the Boyle's temperature (K) is T_B of this gas, then what is the value of $\frac{T_B}{15}$?
15. A flask has 10 gas particles, out of which four particles are moving at 7 ms^{-1} and the remaining are moving at the same speed of ' X ' ms^{-1} . If the r.m.s. of the gas is 5 ms^{-1} , what is the value of X ?

ANSWERS

Level 1

1. (c)	2. (c)	3. (a)	4. (d)	5. (a)	6. (b)	7. (a)	8. (d)	9. (c)	10. (c)
11. (a)	12. (a)	13. (c)	14. (c)	15. (b)	16. (c)	17. (a)	18. (a)	19. (c)	20. (c)
21. (b)	22. (b)	23. (a)	24. (d)	25. (d)	26. (c)	27. (b)	28. (a)	29. (c)	30. (b)
31. (c)	32. (b)	33. (b)	34. (b)	35. (a)	36. (d)	37. (d)	38. (c)	39. (c)	40. (a)
41. (a)	42. (b)	43. (c)	44. (a)	45. (d)	46. (a)	47. (c)	48. (d)	49. (b)	50. (a)
51. (b)	52. (d)	53. (c)	54. (b)	55. (b)	56. (d)	57. (a)	58. (a)	59. (c)	60. (a)
61. (d)	62. (d)	63. (b)	64. (a)	65. (b)	66. (c)	67. (b)	68. (b)	69. (a)	70. (a)
71. (c)	72. (a)	73. (a)	74. (b)	75. (d)	76. (c)	77. (b)	78. (b)	79. (b)	80. (a)
81. (b)	82. (d)	83. (a)	84. (a)	85. (c)	86. (a)	87. (b)	88. (b)	89. (b)	90. (b)
91. (b)	92. (a)	93. (c)	94. (b)	95. (a)	96. (b)	97. (d)	98. (d)	99. (b)	100. (d)
101. (c)	102. (c)	103. (a)	104. (c)	105. (a)	106. (b)	107. (c)	108. (d)	109. (b)	110. (c)
111. (c)	112. (b)	113. (b)	114. (d)	115. (d)	116. (b)	117. (d)	118. (b)	119. (a)	120. (b)
121. (a)	122. (b)	123. (b)	124. (b)	125. (b)	126. (c)	127. (d)	128. (a)	129. (a)	130. (c)
131. (b)	132. (d)	133. (a)	134. (b)	135. (c)	136. (c)	137. (b)	138. (a)	139. (c)	140. (b)
141. (d)	142. (b)	143. (a)	144. (c)	145. (c)	146. (c)	147. (c)	148. (d)	149. (d)	150. (a)
151. (c)	152. (c)	153. (d)	154. (a)	155. (b)	156. (a)	157. (c)	158. (d)	159. (c)	160. (d)
161. (c)	162. (d)	163. (a)	164. (b)	165. (b)	166. (d)	167. (a)	168. (d)	169. (b)	170. (a)
171. (c)	172. (c)	173. (a)	174. (a)	175. (b)					

Level 2

1. (b)	2. (c)	3. (c)	4. (b)	5. (b)	6. (b)	7. (c)	8. (a)	9. (d)	10. (b)
11. (a)	12. (c)	13. (b)	14. (d)	15. (d)	16. (b)	17. (d)	18. (c)	19. (c)	20. (b)
21. (b)	22. (c)	23. (a)	24. (b)	25. (a)	26. (a)	27. (a)	28. (c)	29. (b)	30. (b)



Level 3

Passage-1 1. (d) 2. (a) 3. (c) 4. (d)

Passage-2 1. (c) 2. (c) 3. (a)

Passage-3 1. (a) 2. (d) 3. (b)

One or More Answers is/are correct

1. (a,b,c,d) 2. (a,b) 3. (b,c,d) 4. (c,d) 5. (a) 6. (a,c,d) 7. (a,b) 8. (a,b,c,d)
 9. (a) 10. (a,b,c) 11. (a,b,d) 12. (a,d) 13. (a,b,d) 14. (a,b,c) 15. (a,b,d) 16. (a,b,c,d)
 17. (a,d) 18. (a,c) 19. (b,c) 20. (a,b,c)

Match the Column

1. A P; B R; C Q; D S
 2. A R; B S; C Q; D P
 3. A R; B Q; C P; D S
 4. A P, S; B R; C P, Q; D R
 5. A R; B S; C P; D Q
 6. A Q, S; B R; C P, T; D T
 7. A S; B P; C Q, R; D S

Assertion-Reason Type Questions


1. (B) 2. (C) 3. (A) 4. (B) 5. (A) 6. (A) 7. (C) 8. (B) 9. (C) 10. (C)
 11. (C) 12. (A) 13. (D) 14. (B) 15. (A) 16. (D)

Subjective Problems

1. 7	2. 4	3. 7	4. 3	5. 4	6. 2	7. 6	8. 4	9. 5	10. 4
11. 9	12. 4	13. 9	14. 5	15. 3					

Hints and Solutions

Level 1

2. (c) Boyle's Law
Temperature and mass constant
 $PV = K$
 $\log P + \log V = \log K$
3. (a) $\frac{V_1}{T_1} = \frac{V_2}{T_2}$; pressure and mass constant
 $\frac{0.2}{273} = \frac{V_2}{546}$
 $V_2 = 0.4 \text{ L}$
4. (d) Charles's law
Pressure and mass constant
 $\frac{V}{T} = \text{constant}$
 $V = KT$
 $\log V = \log K + \log T$
5. (a) $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$
 $\frac{P}{348} = \frac{(2P)}{T_2} \cdot \frac{(0.85 V)}{592 \text{ K}}$
 $T_2 = 319 \text{ K}$
6. (b) $V = \frac{KT}{P}$
(i) $\frac{V}{T} = \frac{K}{P}$ (ii) $VT = KT^2$
(iii) $\log V = \log K - \log P$
8. (d) $V = \text{constant}$, mass constant
 $\frac{P}{T} = \text{constant}$
 $P = KT$
 $\log P = \log K + \log T$
10. (c) Boyle's law $PV = K$
 $\frac{P}{V} = \frac{K}{V^2}$... (i)
 $PV^2 = KV$... (ii)
 $\log P + 2 \log V = \log K + \log V$... (iii)
12. (a) $P = n, \text{ initial mole } n_1 = \frac{PV}{RT} = 0.1 \text{ mole}$
 $\frac{P_1}{P_2} = \frac{n_1}{n_2}$
- $\frac{1}{0.75} = \frac{0.1}{n_2}$ $n_2 = 0.075$
Escape mole = 0.025
13. (c) For ideal gas, $PV = nRT$
15. (b) $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$
 $\frac{P}{T} = \frac{0.1}{T} = \frac{1.5 P}{\frac{4T}{3}}$
 $V_2 = 88.9 \text{ mL}$
16. (c) $\frac{V_1}{T_1} = \frac{V_2}{T_2}$
17. (a) $P = \frac{KT}{V}$
 $P = K \left(\frac{1}{V} - \frac{1}{V_0} \right)$
18. (a) $PM = dRT$
 $\frac{M_1}{M_2} = \frac{d_1 T_1}{d_2 T_2} = \frac{1}{2} \cdot \frac{2}{1} = \frac{1}{1}$
19. (c) $PM = dRT$
 $\frac{P_A}{P_B} = \frac{d_A}{d_B} = \frac{M_B}{M_A}$
 $\frac{4}{1}$
20. (c) $V = T$
Because temperature by 10 K
expelled air 10 cm^3
21. (b) 
 $PV = nRT$
 $n, V = \text{constant}$
 $P = \frac{nRT}{V}$
 $T_2 = T_1$
 $P_2 = P_1$
22. (b) $PV = \frac{W}{M} RT$

24. (d) $PV = \frac{W}{M} RT$
 $V = \frac{1}{M}$

25. (d) $P_1 V_1 = P_2 V_2$

26. (c) $d = \frac{PM}{RT}$

27. (b) The moles of the gas in the bubble remains constant, so that $n_1 = n_2$. To calculate the final volume, V_2 ,

$$V_2 = V_1 \frac{P_1}{P_2} \frac{T_2}{T_1}$$

$$2.0 \text{ mL} = \frac{6.0 \text{ atm}}{1.0 \text{ atm}} \frac{298 \text{ K}}{281 \text{ K}}$$

29. (c) $n_{\text{C}_2\text{H}_2} = \frac{PV}{RT} = \frac{1 \times 2.64}{0.0821 \times 273} = 0.118 \text{ mole}$

$$n_{\text{O}_2} = \frac{5}{2} n_{\text{C}_2\text{H}_2} = \frac{5}{2} \times 0.118 = 0.294$$


$$V = \frac{nRT}{P} = 6.6 \text{ L}$$

30. (b) $d = \frac{PM}{dRT}$

31. (c) $P_1 V_1 = P_2 V_2$
 $V_1 = V, V_2 = V/10$

32. (b) $\frac{P_1}{T_1} = \frac{P_2}{T_2}$


33. (b)



$$P = \frac{1}{V}$$

$$V_1 = V_2 = V_3 = V_4$$

$$PV = nRT$$



$$P = T$$

$$P = T$$

$$P_1 = P_2 = P_3 = P_4$$

34. (b) $PV = nRT$

$$n = \frac{1 \times 10}{76 \times 0.0821 \times 1273} = 1.26 \times 10^{-3}$$

$$\text{No. of Na atoms} = 1.26 \times 10^{-3} \times 6.02 \times 10^{23} = 7.6 \times 10^{19}$$

35. (a) Moles of C_2H_6 and C_2H_4 are a and b respectively.

$$\frac{a}{b} = \frac{28}{22.4} = 1.25$$



$$\text{Moles of O}_2 = 3.5a + 3b = 4$$

$$\frac{a}{b} = 1.25$$

$$a = 0.5$$

$$b = 0.75$$

36. (d) $n_{\text{H}_2\text{O}} = \frac{PV}{RT} = 1.54$

$$\frac{n_{\text{MgSO}_4}}{n_{\text{H}_2\text{O}}} = \frac{1}{x} = \frac{54.2}{(120 + 18x) \times 1.54}$$

$$x = 7$$

37. (d) Volume of alveoli $\frac{4}{3} R^3$

$$\frac{4}{3} \times 3.14 \times (5 \times 10^{-3})^3 \text{ mL}$$

$$5.23 \times 10^{-7} \text{ mL} = 5.23 \times 10^{-10} \text{ L}$$

$$n_2 = \frac{14}{100} n_{\text{air}} = 0.14 \frac{PV}{RT}$$

$$0.14 \times 1 \times \frac{5.23 \times 10^{-10}}{0.0821 \times 310}$$

$$2.87 \times 10^{12}$$

$$\text{Molecules} = 2.87 \times 6.02 \times 10^{23} = 1.7 \times 10^{12}$$

38. (c) $\frac{PM}{dRT}$
 $M = 46$

39. (c) Because the number of moles is constant.

$$\frac{P_i V_i}{T_i} = \frac{P_f V_f}{T_f}; \quad P_f = \frac{P_i V_i T_f}{V_f T_i}$$

$$P_f = \frac{P_i V_i T_f}{V_f T_i}$$

$$3.21 \times 10^5 \text{ Pa} \times \frac{V_i}{1.03 V_i} = \frac{(273 + 28.0)}{(273 + 5.00)}$$

$$3.50 \times 10^5 \text{ Pa}$$

40. (a) n_{N_2} 53.57 mole

$$\frac{P_1}{P_2} = \frac{n_1}{n_2}$$

$$\frac{P_1}{P_2} = \frac{n_1}{n_2}$$

$$n_2 = \frac{1.8}{200} \times 53.57 = 0.48213$$

$$n_{N_2 \text{ escape}} = 53.087$$

$$\text{volume} = \frac{nRT}{P} = 1264 \text{ L}$$

41. (a) $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$

$$V_2 = \frac{745 \times 1.41 \times 10^4 \times 225}{294 \times 63.1}$$

$$1.274 \times 10^5 \text{ L}$$

42. (b) P in mm of Hg $\frac{0.61 \times 10^3}{1.013 \times 10^5} = 760$

$$4.6 \text{ mm of Hg}$$

43. (c) $h_{\text{Gallium}} = \frac{74 \times 13.6}{6.095} = 165.1$

44. (a) Volume of He at 0.1 atm

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$50 \times 100 \times V_2 = 1$$

$$V_2 = 5000 \text{ L}$$

$$\text{Volume available for balloon}$$

$$5000 - 50 = 4950 \text{ L at 1 atm}$$

$$\text{Volume of balloon} = 100 \text{ m}^3$$

$$100000 \text{ L} = 49500 \text{ L at 0.1 atm}$$

$$\text{So, 2 cylinder required.}$$

45. (d) $P_1 V_1 = P_2 V_2$, $P_2 = \frac{76 \times 30.40 \times 10}{13.6}$

$$V_2 = \frac{76 \times 14}{98.41} = 10.8 \text{ L}$$

46. (a) $\frac{P_1}{T_1} = \frac{P_2}{T_2}$

$$T_2 = 3 \times \frac{76}{80} = 273$$

47. (c) $n_1 = n_2$
 $\frac{1}{32} = \frac{2.375}{M_2}$

$$M_2 = 76$$

48. (d) $P_i = \frac{6.81}{4} = \frac{0.0821 \times 250}{1.1 \times 10^4}$

$$3.012 \times 10^3$$

$$P_f = 4 \times 10^3$$

$$P = 1 \times 10^3$$

$$n_{\text{He-added}} = \frac{10^3 \times 1.16 \times 10^4}{0.0821 \times 250}$$

$$\frac{wt}{n_{\text{He}}} = \frac{4 \times 2.24 \text{ gm.}}{2.46 \text{ atm}}$$

49. (b) $P = \frac{nRT}{V}$

50. (a) $V = \frac{nRT}{P}$

$$\log V = \log T$$

$$\log V = \log T$$

$$\text{slope} = 1$$

$$45$$

51. (b) Intercept on y-axis

$$\log_{10} \frac{nR}{V} = \log_{10} \frac{10 \times 0.0821}{8.21} = 1.0$$

$$\frac{P}{T} \text{ v/s curve } \frac{P}{T} = \frac{nR}{V}$$

$$\text{Intercept} = \frac{nR}{V}$$

$$\frac{10 \times 0.0821}{8.21} = 0.1, \text{ slope} = 0$$

52. (d) $\frac{3}{2} KT = mgh$

$$h = \frac{3}{2} \times \frac{1.38 \times 10^{23} \times 300}{9.81 \times 1.66 \times 10^{-27}}$$

$$9.53 \times 10^4 \text{ m}$$

53. (c) $d_A = 2d_B$; $3M_A = M_B$; $\frac{PM}{P_A} = \frac{PM}{P_B}$ $\frac{dRT}{d_A} = \frac{dRT}{d_B}$

$$\frac{P_A}{P_B} = \frac{M_A}{M_B} = \frac{d_A}{d_B} = \frac{RT}{RT}$$

$$\frac{P_A}{P_B} = \frac{1}{2} = 2$$

$$\frac{P_A}{P_B} = \frac{4}{1}$$

54. (b) Case I—Suppose inner balloon burst first

$$\frac{600}{300} = \frac{800}{T_2} \quad T_2 = 400 \text{ K}$$

Case II—Suppose outer balloon burst first

$$\frac{1500}{300} = \frac{1800}{T_2}; \quad T_2 = 360 \text{ K}$$

55. (b) $V_1 = V, T_1 = 300 \text{ K}, T_2 = 500 \text{ K}, V_2 = ?$

$$\text{At constant pressure } V_1 T_1 = V_2 T_2$$

$$V_2 = \frac{V_1 T_1}{T_2} = \frac{V \times 300}{500} = \frac{3}{5} V$$

$$\text{Volume of air escaped}$$

$$\text{final volume} - \text{initial volume}$$

$$\frac{5V}{3} \quad V \quad \frac{2V}{3}$$

$$\% \text{ of air escaped} = \frac{2V/3}{5V/3} \times 100 = 40\%$$

56. (d) $R \frac{PV}{nT}$

57. (a) $2\text{NH}_3(\text{g}) \quad \text{N}_2(\text{g}) \quad 3\text{H}_2(\text{g})$

Before sparking	76	0	0
After sparking	76	2x	x
at eqm			3x

Increase in pressure $2x$; $x = 3 \text{ cm Hg}$

Partial pressure of $\text{H}_2 = 3 + 3 = 9 \text{ cm Hg}$

58. (a) $P_{\text{Atm}} = \frac{h}{13.6} \quad P_{\text{Gas}} = \text{aq Tension}$

$P_{\text{Gas}} = P_{\text{Atm}} + \text{aq tension} = \frac{h}{13.6}$

$P = \frac{4V_0}{V_0 + h}$

59. (c) $P + \frac{P}{10} h$

$\frac{P_1 V_1}{P} = \frac{P_2 V_2}{P} \quad \frac{h}{10} = V$

$4 = \frac{h}{10}$

$h = 30 \text{ m}$

60. (a) $h_g d_g = h_{\text{Hg}} d_{\text{Hg}}; 5 \times 2.72 = 13.6 h_{\text{Hg}}$

$h_{\text{Hg}} = 1 \text{ m}$

$P_{\text{gas}} = 0.76 \times 1 = 1.76 \text{ m of Hg}$

$= 1760 \text{ mm of Hg}$

$\frac{1760}{760} = 10 = n \times R = 300$


$n = 0.94 \text{ mole}$

61. (d) $n_T = n_1 = n_2$

$$\frac{P}{R} = \frac{1.25}{287.5} = \frac{611 \times 10^3}{293.5 R} = \frac{0.5}{204.5 R} = \frac{0.75}{204.5 R}$$

$P = 4.84 \times 10^5 \text{ Pa}$

62. (d)



$\frac{0.2P}{RT} = \frac{(100 - V)}{RT} = \frac{100P}{RT}$

$0.2(100 - V) = 100$

$V = 400 \text{ cm}^3$

63. (b) Moles of C_2H_2 and C_3H_8 are a and b



$a \quad 2a$



$b \quad 3b$

$2a + 3b = 230$


$a + b = 80$

$b = 70$

$a = 10$

mole fraction $\frac{7}{8} = 0.875$

64. (a) $p = n$



$a \quad b$

$a + 2b = b + b$

$a + b = 1$

$a + 2b = 0.4$

$3b = 0.6$

$b = 0.2$

$a = 0.8$

65. (b) $P_1 = \frac{5}{V} \frac{RT}{V}; \text{ p.pr. of He} = \frac{2}{5} \frac{5RT}{V}$

$$\frac{2 \times 0.0821 \times 400}{8.21} = 8 \text{ atm}$$

66. (c) $P_{\text{O}_2} = \frac{3}{10} P_T;$

After removing 2 mole of O_2 ,

$P_{\text{O}_2} = \frac{1}{8} P_T$

Decrease in partial pressure of O_2

$\frac{3P_T}{10} - \frac{P_T}{8} = \frac{10P_T}{80} = \frac{10 \times 8}{10} = 100$

58.33

67. (b) $P_{\text{N}_2} = \frac{760 \times 30}{730} = \frac{730 \text{ torr.}}{0.821}$

$$n_{\text{N}_2} = \frac{760 \times 0.0821}{760 \times 0.0821 \times 300} = 0.032$$

68. (b) % Relative humidity

$$\frac{\text{Partial pressure of } \text{H}_2\text{O}(\text{g})}{\text{Vapour pressure of } \text{H}_2\text{O}} \times 100$$


at constant temperature

69. (a) Volume of O_2 1 L
and p_{O_2} 760 355 405 mm Hg
Since, the temperature is constant, using Boyle's law we get

$$1 \text{ L } 405 \text{ mm } 0.3 \text{ L } p_{O_2}$$

$$p_{O_2} \quad 1350 \text{ mm Hg}$$

70. (a) Dalton's law is applicable for a non-reacting gaseous mixture.



71. (c) $n_{CH_4} \frac{x}{16} \quad n_{O_2} \frac{x}{32}$

$$\text{mole fraction } O_2 \quad 1/3$$

72. (a) $n_T \quad n_1 \quad n_2 \quad n_1 \quad \frac{P}{R} \quad \frac{1}{T} \quad 1$

$$\frac{P}{RT} \quad 2 \quad P \quad RT$$

$$P \quad RT \quad P$$

73. (a) $n \quad \frac{PV}{RT} \quad \frac{3170}{8.314} \quad \frac{10^3}{300} \quad 1.27 \times 10^3 \text{ mole.}$

74. (b) Dalton's Law is applicable on a non-reacting gas mixture.

75. (d) mole $N_2 \quad \frac{56}{28} \quad 2, \quad \text{mole } O_2 \quad \frac{96}{32} \quad 3$

$$p_{O_2} \quad \frac{2}{5} \quad 10 \quad 4 \text{ atm}$$

$$p_{N_2} \quad \frac{3}{5} \quad 10 \quad 6 \text{ atm}$$

76. (c) $P \quad n$
- $$\frac{P_{H_2}}{P_{CO_2}} \quad \frac{n_{H_2}}{n_{CO_2}} \quad \frac{22}{1}$$

$$P_{H_2} \quad 22 \text{ atm}$$

77. (b) $P_{\text{gas}} \quad P_{\text{dry gas}} \quad P_{\text{moisture}} \text{ at } T \text{ K}$

$$\text{or } P_{\text{dry}} \quad 830 \quad 30 \quad 800$$

$$\text{Now at } T_2 \quad 0.99 T_1;$$

$$\text{at constant volume } \frac{P_1}{T_1} \quad \frac{P_2}{T_2}$$

$$P_{\text{dry}} \quad \frac{800}{T} \quad \frac{0.99 T}{T} \quad 792 \text{ mm}$$

$$P_{\text{gas}} \quad P_{\text{dry}} \quad P_{\text{moisture}}$$

$$792 \quad 25 \quad 817 \text{ mm}$$

78. (b) We have $P_{O_2} \quad X_{O_2} \quad P_{\text{total}}$
- $$0.60 \quad \frac{n_{O_2}}{n_{O_2} + n_{SO_2}} \quad P_{\text{total}}$$

Let no. of moles of O_2 are x ;

$$0.60 \quad \frac{x/32}{x/32 + 3x/64} \quad P_{\text{total}}$$

$$0.60 \quad 0.4 \quad P_{\text{total}}$$

$$P_{\text{total}} \quad \frac{0.60}{0.4} \quad 1.5 \text{ atm}$$

$$\therefore P_{O_2} \quad P_{SO_2} \quad 1.5 \text{ atm}$$

$$P_{SO_2} \quad 1.5 \quad P_{O_2} \quad 1.5 \quad 0.60 \quad 0.9 \text{ atm}$$

$$\text{Concentration of } SO_2 \quad \frac{0.9}{0.0821 \quad 300}$$

$$0.036$$

79. (b) Given $P \quad 10 \text{ atm}$,
total numbers of moles; $n_A \quad n_B \quad n_C \quad 10$

$$P_A \quad 3 \text{ atm}, \quad P_B \quad 1 \text{ atm}$$

$$\therefore P_A \quad x_A \quad P_{\text{(total)}} \quad \frac{n_A}{n_A + n_B + n_C} \quad 10$$

$$\frac{n_A}{10} \quad 10, n_A \quad 3$$

$$\text{Similarly, } P_B \quad x_B \quad P_{\text{(total)}}$$

$$\text{So, } n_B \quad 1$$

$$n_C \quad 10 \quad (n_A \quad n_B) \quad 10 \quad 4 \quad 6$$

$$\text{Mass of C } 6 \quad 2 \quad 12 \text{ g}$$

80. (a) $PV \quad \frac{10}{M} RT \quad \dots(1)$

Let x g of the gas remain in the flask when
final pressure $\frac{P}{2}, \frac{2T}{3}$

$$\frac{P}{2} \quad V \quad \frac{x}{M} \quad R \quad \frac{2T}{3} \quad \dots(2)$$

$$(1)/(2) \quad 2 \quad \frac{10}{x} \quad \frac{3}{2}$$

$$x \quad \frac{3 \cdot 10}{4} \quad 7.5 \text{ g}$$


81. (b) $\frac{P_x}{P_y} \quad \frac{n_x}{n_y}$
- $$\frac{2}{1} \quad \frac{1/m_x}{2/m_y}$$

$$m_y \quad 4m_x$$

82. (d) $P \quad \frac{nRT}{V}$
- $$P \quad \frac{0.05 \quad 0.0821 \quad 303}{44 \quad 4.6} \quad 6.14 \times 10^3 \text{ atm}$$

83. (a) P_T P_{He} P_{Ne} V. PH_2O
 745 368 P_{Ne} 28.3
 P_{Ne} 348.7 mm of Hg

84. (a)



$$n_{He} = \frac{1.2}{0.082} \frac{0.63}{289}, \quad n_{Ne} = \frac{3.4}{0.0821} \frac{2.8}{289}$$

$$P_f = \frac{0.431}{4.6} \frac{0.0821}{289} \quad 2.223$$

$$P_{He} = \frac{0.032}{0.432} \quad 2.223 \quad 0.164 \text{ atm}$$

85. (c) From the total pressure and the vapour pressure of water we can calculate the partial pressure of O_2 .

$$P_{O_2} \quad P_T \quad P_{H_2O}$$

$$760 \quad 22.4$$

$$737.6 \text{ mm Hg}$$

From the ideal gas equation we write

$$m = \frac{PVM}{RT}$$

$$\frac{(0.974 \text{ atm})(0.128 \text{ L})(32.0 \text{ g/mol})}{(0.0821 \text{ L atm/K mol})(273 + 24) \text{ K}}$$

$$0.163 \text{ g}$$

86. (a) $PV = nRT$ R K N_A

$$PV = n N_A \frac{PV}{KT} \text{ no. of particles}$$

87. (b) $KE = \frac{1}{2} Mv^2$

88. (b) K.E is temperature dependent

same moles would have same volume at same P and T

$$\frac{PV}{d} = \frac{nRT}{PM}$$

89. (b) According to Boyle's law at constant temperature and mass

$$\frac{PV}{P} = \frac{K}{V}$$

90. (b) $KE_T = \frac{3}{2} nRT$ at constant P and V

$PV = K$ at constant T and mass

$V = KT$ at constant P and mass

91. (b) $KE_{avg} = \frac{3}{2} KT$

92. (a) $PV = nRT$
 $PV = T$ at given mass.

93. (c) According to kinetic theory of gases collision between molecules are perfectly elastic.

94. (b) $KE_T = \frac{3}{2} nRT = \frac{3}{2} \frac{W}{M} RT$

$$KE = \frac{T}{M}$$

$$\frac{KE_1}{KE_2} = \frac{T_1}{T_2} \frac{M_2}{M_1} = \frac{300}{2} \frac{44}{600} \frac{11}{1}$$

95. (a) $PV = \frac{1}{3} mu^2$
- $$P = \frac{1}{3} \frac{2}{3} \frac{1}{2} mu^2$$
- $$P = \frac{2}{3} E.$$

96. (b) Moles ratio $\frac{n_{He}}{n_{N_2}} = \frac{w/4}{w/28} = \frac{7}{1}$

Ratio of translational K.E. $\frac{n_{He} T_{He}}{n_{N_2} T_{N_2}} = \frac{7}{1} \frac{300}{700} = 3:1$

97. (d) Due to phase transfer distance between particle changes, because temperature remains constant, so KE remains constant.

98. (d) $K = \frac{3}{2} n_T RT$ $n_T = \frac{3000}{3} \frac{2}{250} \frac{2}{8.314}$

$$3 \times 10^3 \times \frac{3}{2} n_T \times 8.314 \times 250$$

$$\frac{x}{20} \times \frac{30}{40} \times \frac{x}{8.314} \times \frac{2}{28.3}$$

99. (b) $KE = \frac{3}{2} RT$

$$U_{rms} = \sqrt{\frac{3RT}{M}}$$

100. (d) $U_{rms} = \sqrt{\frac{U_1^2 + U_2^2 + U_3^2 + U_4^2}{4}} = \sqrt{\frac{54}{4}} = \frac{\sqrt{54}}{2}$

101. (c) $U_{ms} = \sqrt{\frac{4(6)^2 + 5(2)^2 + 10(3)^2}{19}}$

$$3.6 \text{ cm/sec.}$$

$$102. (c) U_{\text{ms}} \sqrt{\frac{3RT}{Mw}}$$

$$\frac{U_{\text{H}_2}}{U_{\text{O}_2}} \sqrt{\frac{50}{2} \cdot \frac{32}{800}} = 1$$

$$104. (c) U_{\text{rms}} \sqrt{\frac{3RT}{M}} \frac{(U_{\text{rms}})_{\text{H}_2}}{(U_{\text{rms}})_{\text{N}_2}} \sqrt{\frac{T_{\text{H}_2}}{M_{\text{H}_2}} \cdot \frac{M_{\text{N}_2}}{T_{\text{N}_2}}};$$

$$\frac{(U_{\text{rms}})_{\text{H}_2}}{(U_{\text{rms}})_{\text{N}_2}} \sqrt{5} \frac{(U_{\text{rms}})_{\text{N}_2}}{(U_{\text{rms}})_{\text{N}_2}}$$

$$\sqrt{5} \sqrt{\frac{T_{\text{H}_2}}{T_{\text{N}_2}} \cdot \frac{28}{2}}$$

$$\frac{\sqrt{5}}{1} \sqrt{\frac{T_{\text{H}_2}}{T_{\text{N}_2}} \cdot 14}$$

$$5 \frac{T_{\text{H}_2}}{T_{\text{N}_2}} = 14$$

$$\frac{T_{\text{N}_2}}{T_{\text{H}_2}} = \frac{5}{14}$$

$$105. (a) U_{\text{rms}} \sqrt{\frac{2RT}{M}} : U_{\text{avg}} \sqrt{\frac{8RT}{M}} : U_{\text{mp}} \sqrt{\frac{2RT}{M}}$$

$$\frac{U_{\text{rms}}}{1.224} : \frac{U_{\text{avg}}}{1.128} : \frac{U_{\text{mp}}}{1}$$

$$\frac{U_{\text{rms}}}{1.224} : \frac{U_{\text{avg}}}{1.128} : \frac{U_{\text{mp}}}{1}$$

$$106. (b) 200 \sqrt{\frac{2RT}{2 \cdot 10^3}} \quad RT = 40$$

$$\text{Average K.E.} = \frac{3}{2} nRT = \frac{3}{2} \cdot \frac{8}{2} = 40 \quad 240 \text{ J}$$

107. (c) Second member of C_nH_{2n} series

$$\text{C}_3\text{H}_6 \quad 42$$

$$\sqrt{\frac{8RT_1}{M_1}} : \sqrt{\frac{8RT_2}{M_2}} = \frac{900}{71} : \frac{T_2}{42}$$

$$T_2 = 532.4 \text{ K}$$

$$108. (d) U_{\text{rms}} \sqrt{\frac{3RT}{M}}$$

$$109. (b) P = \frac{1}{3} \frac{nmc^2}{V}$$

$$\frac{1}{3} \frac{6 \cdot 10^{22} \cdot 10^{-24} \cdot (100)^2}{10 \cdot 10^{-3}}$$

$$2 \cdot 10^4 \text{ Pa}$$

110. (c) Second member is C_3H_4 ;

$$\sqrt{\frac{2RT_1}{M_1}} : \sqrt{\frac{2RT_2}{M_2}}$$

$$T_1 \quad T_2 \quad \frac{M_1}{M_2} \quad 800 \quad \frac{40}{64} \quad \text{K}$$

$$500 \text{ K or } 227^\circ\text{C}$$

$$111. (c) 300 \sqrt{\frac{3RT}{4 \cdot 10^3}}; \quad RT = 120$$

$$\text{Total K.E. of He gas} = \frac{3}{2} nRT$$

$$\frac{3}{2} \cdot \frac{8}{4} = 120 \text{ J}$$

$$360 \text{ J}$$

$$112. (b) U_{\text{mp}} \sqrt{\frac{2RT}{M}}$$

$$U_{\text{avg}} \sqrt{\frac{8RT_2}{M}}$$

$$U_{\text{rms}} \sqrt{\frac{RT_1}{M}}$$

$$T_3 \quad T_2 \quad T_1$$

$$113. (b) U_1 \sqrt{\frac{3P_1}{d_1}}$$

$$U_{\text{rms}} \sqrt{\frac{3}{d}} (\sqrt{P_2} \quad \sqrt{P_1})$$

$$\sqrt{\frac{3}{0.75}} (300 \quad 200)$$

$$\sqrt{4} \quad 100 \quad 200$$

$$114. (d) \text{Rate of diffusion} = \frac{1}{\sqrt{\text{density}}}$$

at constant P and T

$$d \propto \frac{PM}{RT}, \quad d \propto M$$

$$115. (d) : v : u = \sqrt{2} : \sqrt{\frac{8}{3}} : \sqrt{3}$$

$$116. (b) \sqrt{\frac{8RT}{M}} \quad \sqrt{\frac{28}{88}} \quad \sqrt{\frac{7}{22}}$$

$$\frac{8}{16} \cdot \frac{8.314}{10^3} \cdot \frac{T}{22} = 10^6$$

$$T = \frac{1000 \cdot 2}{8.314} = 240.55 \text{ K}$$

$$T \quad ^\circ\text{C} \quad 240.55 \quad 273 \quad 32.45 \quad ^\circ\text{C}$$

$$117. (d) \sqrt{\frac{2RT}{M_{\text{SO}_2}}} : \sqrt{\frac{3RT_1}{M_{\text{O}_2}}} = \frac{2}{64} : \frac{T}{32}$$

$$T = 900 \text{ K}$$

$$118. (b) U_{\text{rms}} \sqrt{\frac{3RT}{M}}$$

$$119. (a) \text{Rate of diffusion} \frac{1}{\sqrt{M}} \quad P$$

$$\text{Rate of diffusion} \frac{P}{\sqrt{M}}$$

120. (b) According to Graham's law

$$r \propto \frac{1}{\sqrt{d}}, \quad \frac{r_1}{r_2} \propto \sqrt{\frac{d_2}{d_1}} \text{ at constant } P \text{ and } T$$

$$121. (a) \frac{r_1}{r_2} \propto \sqrt{\frac{M_2}{M_1}} \text{ at constant } P \text{ and } T$$

$$122. (b) \frac{r_1}{r_{\text{H}_2}} \propto \sqrt{\frac{2}{M_1}} \quad \frac{1}{6}$$

$$M = 72$$

$$123. (b) \frac{r_1}{r_2} \propto \sqrt{\frac{M_2}{M_1}}$$

$$\frac{V_1}{t_1} \propto \frac{t_2}{V_2} \propto \sqrt{\frac{M_2}{M_1}}$$

$$124. (b) \frac{r_{\text{H}_2}}{r_{\text{H.C}}} \propto \sqrt{\frac{M}{2}} \quad 3\sqrt{3}$$

$$\frac{M}{2} = 9 \quad 3$$

$$M = 54$$

$$n = 4$$

$$125. (b) \frac{r_{\text{O}_2}}{r_{\text{CH}_4}} \propto \frac{n_{\text{O}_2}}{n_{\text{CH}_4}} \sqrt{\frac{M_{\text{CH}_4}}{M_{\text{O}_2}}}$$

$$\frac{3}{2} \propto \frac{16}{32} \sqrt{\frac{16}{32}} \quad \frac{3}{4\sqrt{2}}$$

$$126. (c) \frac{n_{\text{CH}_4}}{8/1} \propto \frac{16^{3/2}}{64} \quad \frac{1}{1};$$

$$\frac{r_{\text{SO}_2}}{r_{\text{CH}_4}} \propto \frac{1}{1} \sqrt{\frac{16}{64}} \quad \frac{1}{2}$$

$$127. (d) \frac{n_{\text{He}}}{n_{\text{CH}_4}} \propto \frac{1}{2} \sqrt{\frac{16}{4}} \quad \frac{1}{1}$$


$$\frac{n_{\text{He}}}{n_{\text{SO}_2}} \propto \frac{1}{3} \sqrt{\frac{64}{4}} \quad \frac{4}{3}$$

$$n_{\text{He}} : n_{\text{CH}_4} : n_{\text{SO}_2} = 4 : 4 : 3$$

$$128. (a) \frac{r_1}{r_2} \propto \frac{V_1/t_1}{V_2/t_2} \propto \frac{V_1}{V_2} \cdot \frac{t_2}{t_1} \propto \sqrt{\frac{M_2}{M_1}};$$

$$\frac{80}{V_2} \cdot \frac{3}{2} \propto \sqrt{\frac{64}{32}} \propto \sqrt{2}; \quad V_2 \propto \frac{120}{\sqrt{2}}$$

$$129. (a) \frac{r_{\text{NH}_3}}{r_{\text{HCl}}} \propto \sqrt{\frac{36.5}{17}} \propto \frac{x}{87.5 - x}$$



$$x = 51.7 \text{ cm}$$

$$130. (c) \frac{r_{\text{CH}_3\text{OCH}_3}}{r_{\text{CH}_4}} \propto \sqrt{\frac{M_{\text{CH}_4}}{M_{\text{CH}_3\text{OCH}_3}}} \propto \frac{P_{\text{CH}_3\text{OCH}_3}}{P_{\text{CH}_4}}$$

$$\sqrt{\frac{16}{46}} \propto \frac{0.8}{0.2} \quad 2.36 : 1$$

$$131. (b) \frac{n_1}{n_2} \propto \frac{t_2}{t_1} \propto \sqrt{\frac{M_2}{M_1}}; \quad \frac{w_1}{w_2} \propto \frac{M_2}{M_1} \propto \sqrt{\frac{M_2}{M_1}}$$

$$\frac{w_1}{w_2} \propto \sqrt{\frac{M_1}{M_2}}; \quad \frac{w_1}{4} \propto \sqrt{\frac{4}{64}}$$

$$132. (d) \frac{r_{\text{mix}}}{r_x} \propto \frac{4/5}{4/10} \propto 2 \propto \sqrt{\frac{M_x}{M_{\text{mix}}}}$$

$$M_{\text{mix}} = 9$$

$$M_{\text{mix}} = \frac{M_{\text{H}_2} X_{\text{H}_2} + M_{\text{CH}_4} X_{\text{CH}_4}}{2X_{\text{H}_2} + 16(1 - X_{\text{H}_2})}$$

$$9 = \frac{2 \cdot 2 + 16 \cdot 0.5}{2 \cdot 2 + 16 \cdot 0.5}$$

$$133. (a) \text{Mass of the filled balloon} = 50 + 685 = 735 \text{ kg}$$

$$\text{Pay load} = \text{Mass of displaced air} - \text{Mass of balloon}$$

$$= 5108 - 735 = 4373 \text{ kg}$$

$$134. (b) \text{Energy associated with each degree of freedom is } \frac{1}{2} k_B T$$

$$135. (c) C_v = \frac{3}{2} R \quad R \quad 4R \quad \frac{3}{4} R \quad \frac{11}{2} R;$$

$$C_p = \frac{11}{2} R \quad R \quad \frac{13}{2} R$$

$$\frac{C_p}{C_v} = \frac{13}{11} = 1.18$$

$$136. (c) C_v = \frac{\frac{3}{2}R + \frac{5}{2}R}{2} = 2R$$

$$C_p = \frac{\frac{5}{2}R + \frac{7}{2}R}{2} = 3R$$

$$\frac{C_p}{C_v} = 1.5$$

$$137. (b) C_v = \frac{3}{2}RT \quad \text{for monoatomic gas;}$$

$$C_p = \frac{5}{2}RT$$

$$C_v = \frac{5}{2}RT \quad \text{for diatomic gas}$$

$$C_p = \frac{7}{2}RT$$

Thus, for mixture of 1 mole each,

$$C_v = \frac{\frac{3}{2}RT + \frac{5}{2}RT}{2}$$

$$\text{and } C_p = \frac{\frac{5}{2}RT + \frac{7}{2}RT}{2}$$

$$\text{Thus, } \frac{C_p}{C_v} = \frac{3RT}{2RT} = 1.5$$

138. (a) Internal energy of a substance is a function of temperature and volume (except ideal gas). For ideal gas it is only function of temperature.

139. (c) For ideal gas $PV = nRT$
volume of 1 mole gas at 1 atm pressure and 273 K is 22.4 L and at 300 K and 1 atm pressure volume = 24.63 L.

140. (b) If attraction are disappear than pressure of real gas

$$141. (d) P_{\text{Ideal}} = P_{\text{Real}} + \frac{n^2 a}{V^2}$$

142. (b) Real gas behaves like an ideal gas at low pressure and high temperature at this condition $Z = 1$.

$$143. (a) P_{\text{Ideal}} = P_r + \frac{n^2 a}{V^2}$$

$$144. (c) P_{\text{Real}} = P_i + \frac{n^2 a}{V^2}$$

$$a = P_r$$

$$P_2 = P_3 = P_1$$

$$145. (c) P(V - b) = RT; \quad P = \frac{RT}{(V - b)};$$

$$P = \frac{R}{(V - b)} T$$

146. (c) $b = 4$ volume occupied by molecules of 1 mol of a gas,

$$\text{or } b = 4N_A \frac{4}{3} r^3$$

$$\text{or } r = \frac{3b}{16N_A}^{1/3}$$

$$\frac{3}{16} \frac{24}{6.023 \times 10^{23}}^{1/3} \text{ cm}$$

$$1.355 \times 10^{-8} \text{ cm}$$

Now, molecular diameter, $d = 2r$

$$2.67 \times 10^{-8} \text{ cm}$$

147. (c) Liquifaction I.M.A.F. a

$$148. (d) V_i = V, V_{\text{ex}} = n \cdot b$$

b size

149. (d) A real gas can behave like an ideal gas at low pressure and high temperature.

$$150. (a) P(V - nb) = nRT$$

$$\frac{PV}{nRT} = \frac{Pnb}{nRT} + \frac{d \frac{\text{Mass}}{\text{Volume}}}{n \frac{\text{Mass}}{M}}$$

$$P \frac{\text{Mass}}{d} = nRT + Pnb$$

$$\frac{P}{d} = \frac{n}{\text{Mass}} RT + \frac{Pnb}{\text{Mass}}$$

$$\frac{P}{d} = \frac{RT}{M} + \frac{Pb}{M}$$

$$151. (c) P = \frac{a}{V_m^2} (V_m - b) = RT$$

152. (c) At low pressure,

$$P = \frac{a}{V^2} (V) = RT$$

$$\text{i.e., } PV^2 = RTV + a$$

$$V = \frac{RT}{2P} \frac{\sqrt{R^2 T^2 + 4Pa}}{2P} = \frac{RT}{2P}$$

$$(\because 4aP = R^2 T^2)$$

153. (d) Real gas shows ideal behaviour at low pressure and high temperature.

$$154. (a) \frac{PV}{RT} Z \quad Z - 1 \quad \frac{a}{VRT}$$

155. (b) van der Waals' equation for one mole of a real gas is

$$P \left(\frac{a}{V^2} + (V - b) \right) = RT$$

$$\text{or } PV = RT - Pb + \frac{a}{V}$$

At high pressures, the van der Waals' gas equation reduces to $PV = RT - Pb$

$$\text{or } \frac{PV}{RT} = 1 - \frac{Pb}{RT}$$

$$\text{Compressibility factor } Z = \frac{PV}{RT} = 1 - \frac{Pb}{RT}$$

$$156. (a) \frac{PV}{RT} = 1 \quad RT \text{ constant}$$

$$PV = 1 \quad \text{at 1 atm}$$

$$V_m = 22.4 \text{ L}$$

$$157. (c) Z = \frac{PV}{nRT} = \frac{V}{n} \cdot \frac{0.9}{0.0821} \cdot \frac{273}{9}$$

$$= 2.24 \text{ litre/mol}$$

$$\text{Volume of 1 milli-mole of gas} = 2.24 \text{ mL}$$

$$158. (d) Z = \frac{PV}{nRT}; 1.90 = \frac{1}{n} \cdot \frac{800}{R \cdot 330};$$

$$n = \frac{1}{1.90} \cdot \frac{800}{R \cdot 330}$$

$$Z = 1.10 = \frac{V}{n} \cdot \frac{200}{R \cdot 570};$$

$$1.10 = \frac{V}{800} \cdot \frac{200}{R \cdot 570}$$

$$V = 4 \text{ L}$$

159. (c) $Z > 1$ real gas shows positive deviation from ideal behaviour when repulsive forces dominate.

$$160. (d) 0.1 = \frac{1000}{V^2} \cdot \frac{(0.02)^2}{V} \cdot 20 \cdot 0.02$$

$$0.1 V^2 = 0.4 V \cdot 0.4 \cdot 0$$

$$V^2 = 4V \quad 4 = 0$$

$$V = 2 \text{ L}$$

$$Z = \frac{PV}{nRT} = \frac{0.1}{20} \cdot \frac{2}{0.02} = 0.5$$

$$161. (c) T_C = \frac{8a}{27Rb} \quad P_C = \frac{a}{27b^2} \quad V_C = 3b$$

$$163. (a) V = K(t - C + 273)$$

$$1 \text{ C} \quad V_1 = K(273 - 1) = K \cdot 272 \quad K$$

$$V = K(273) - t + 0 \text{ C}$$

$$2 \text{ C} \quad V_2 = K(273 - 2) = K \cdot 271 \quad 2K$$

164. (b) It is a temperature above which a gas can not be liquified.

$$165. (b) \frac{T_C}{P_C} = \frac{8a}{27Rb} \cdot \frac{27b^2}{a} = \frac{8b}{R}$$

$$V_{\text{Excluded}} = nb$$

$$166. (d) T_{\text{Liquifaction}} = \frac{1}{I.M.A.F} = \frac{1}{a} = \frac{1}{T_C}$$

Critical temperature is a temperature above which a gas can't be liquify.

$$167. (a) T_C = \frac{8a}{27Rb}, \quad T_b = \frac{a}{Rb}, \quad T_i = \frac{2a}{Rb}$$

$$T_i = T_b = T_C$$

$$168. (d) \text{At low pressure } Z = 1 + b \cdot \frac{a}{RT} \cdot \frac{1}{V_m} \text{ at}$$

$$\text{Boyles temperature } b \cdot \frac{a}{RT} = 0$$

169. (b) Below critical temperature a gas can be liquified.

So V_m has three real roots.

170. (a) At critical temperature slope of pressure volume curve is maximum and it is zero.

171. (c) Above critical temperature a gas can't be liquified.

172. (c) At Boyle's temperature and low pressure $Z = 1$.

$$\text{At low pressure } Z = 1 + b \cdot \frac{a}{RT} \cdot \frac{1}{V_m}$$

$$\text{At } T_B \quad a/Rb, \quad Z = 1$$

173. (a) Because $T = T_i$ and if gas expanded adiabatically then T

174. (a) The van der Waals' equation for 1 mole of gas is

$$P \left(\frac{a}{V^2} + (V - b) \right) = RT$$

$$\text{if we put, } P_r = \frac{P}{P_C}, \quad V_r = \frac{V_m}{V_C} \quad \text{and } T_r = \frac{T}{T_C};$$

$$\therefore P_C = \frac{a}{27b^2}, \quad V_C = 3b \quad \text{and } T_C = \frac{8a}{27Rb}$$

$$P_r = \frac{a^2}{27b^2} \cdot \frac{a}{V_r^2(3b)^2} (V_r(3b) - b) = RT_r \cdot \frac{8a}{27Rb}$$

This equation is called van der Waals' equation of law of corresponding states.

175. (b) $\frac{P_r V_r}{T_r} = 2.21$

$$\frac{P}{P_C} \frac{V}{V_C} \frac{T_C}{T} = 2.21$$

$$\frac{PV}{T} \frac{T_C}{P_C V_C} = 2.21$$

$$0.1243 \text{ L} \quad 124.31 \text{ mL}$$

Level 2

1. (b) The number of moles of He, Ne and Xe is given by

$$n_{\text{He}} \frac{PV}{RT} \frac{3}{RT}; \quad n_{\text{Ne}} \frac{PV}{RT} \frac{10}{RT};$$

$$n_{\text{Xe}} \frac{PV}{RT} \frac{1}{RT}$$

$$n \quad n_{\text{He}} \quad n_{\text{Ne}} \quad n_{\text{Xe}} \quad \frac{(3 \quad 10 \quad 1)}{RT}$$

The total pressure is given by

$$P = \frac{(n_{\text{He}} \quad n_{\text{Ne}} \quad n_{\text{Xe}}) RT}{V}$$

$$P_{\text{total}} = \frac{(3 \quad 10 \quad 1)}{RT} \frac{RT}{7} = 2 \text{ atm}$$

2. (c) $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$

Initial moles 11 12 0

at equilibrium 9 6 4

moles of N_2 and H_2 present at equilibrium

15

after addition of water

$\text{NH}_3(\text{g}) \quad \text{H}_2\text{O}(\text{l}) \quad \text{NH}_4\text{OH}(\text{l})$

Volume of vessel available for gaseous


mixture of N_2 and H_2 20 3.58

16.42 L

Pressure exerted by gaseous mixture at

$$300 \text{ K} \quad \frac{15 \quad 0.821 \quad 300}{16.42} \quad 22.5 \text{ atm}$$

3. (c)



Let x mole of N_2 present into vessel II and P is final pressure of N_2

$$P(2V) = xR(T_2/3) \text{ and } P(V) = (0.1 - x)RT_2$$

$$2 \quad \frac{x}{3(0.1 - x)}$$

$$x = 0.6/7 \text{ mole};$$

$$\frac{0.6}{7} \quad 28 \quad 2.4 \text{ g N}_2$$

II has 2.4 g N_2 and I has 0.4 g of N_2 ;

$$\frac{W_I}{W_{II}} = \frac{0.4}{2.4} = 1:6$$

4. (b) Let initial mixture contains n_1 and n_2 moles of NH_3 and N_2H_4 respectively

Total moles of gases originally present

$$n_1 \quad n_2$$

Total moles of gases after decomposition of gases $2n_1 \quad 3n_2$

$$0.5 \quad V \quad (n_1 \quad n_2) R \quad 300$$

$$4.5 \quad V \quad (2n_1 \quad 3n_2) R \quad 1200$$

$$\frac{2n_1 \quad 3n_2}{n_1 \quad n_2} = \frac{9}{4}$$

$$\frac{n_1}{n_2} = \frac{1}{3}$$

$$\frac{n_2}{n_1} = 100 \quad 25\%$$

6. (b) Let $T_1 \quad T_2$; final pressure will be same, Let x mole transfer from A to B vessel.

$$P_A V = (n - x) RT$$

$$\text{and } P_A V = (n - x) RT_2$$

$$x = \frac{n(T_1 - T_2)}{T_1 - T_2};$$

$$\text{finally } P_1 = 2V \quad 2nRT_1; V \quad \frac{nRT_1}{P_1}$$

$$P_A = \frac{nRT_1}{P_1} = n \frac{n(T_1 - T_2)}{(T_1 - T_2)} RT_1$$

$$P_A = \frac{2P_1 T_2}{T_1 - T_2}$$

7. (c) $P = d; \quad P = kd \text{ and } k = \frac{1 \text{ atm}}{1 \text{ metre}}$

$$PV = nRT; \quad kd = \frac{1}{6} d^3 \quad nRT;$$

$$\frac{d_1^4}{d_2^4} = \frac{n_1}{n_2}; \quad \frac{1}{4^4} = n_1/n_2; \quad n_2 = 256$$

$$\text{no. of moles added} = 256 - 1 = 255$$

8. (a) $PM = dRT$

$$Pd = P^2 \frac{M}{RT}$$

$$\frac{d(Pd)}{dP} = \frac{2PM}{RT} = 5$$

$$\frac{0.2}{0.0821} \frac{8.21}{T} \frac{4}{5}$$

$$T = 160 \text{ K}$$

9. (d) % Relative humidity

$$\frac{\text{Partial pressure of H}_2\text{O}}{\text{Vapour pressure of H}_2\text{O}} \times 100$$

$$75 \frac{P_{\text{H}_2\text{O}}}{30} \times 100$$

$$P_{\text{H}_2\text{O}} = 22.5 \text{ torr;}$$

% of H₂O vapour in air

$$\frac{(22.5)}{760} \times 100 = 2.96$$

molar mass of wet air

$$29 \times 97.04 + 2.96 \times 18$$

$$\frac{100}{2814.16 + 53.28} = 28.67$$

$$\text{density of wet air} = \frac{PM}{RT} = \frac{1 \times 28.67}{0.0821 \times 300}$$

$$1.164 \text{ g/L}$$

10. (b) C_v of gas A $\frac{3}{2}R$ $\frac{3}{2}R$ $6R$ $9R$

Let C_v of gas B = X

$$\text{Heat lost by gas A} = 7 \times 9R \times T \times \frac{5T}{6}$$

$$\text{Heat gained by gas B} = 6X \times \frac{5T}{6} \times \frac{T}{3}$$

$$63R \times \frac{T}{6} = 6X \times \frac{T}{2} \Rightarrow X = \frac{7}{2}R;$$

gas is diatomic

11. (a) Total heat received by gas = heat provided

$$2 \times \frac{3}{2}R(T_f - 300) = 2 \times 3R(T_f - 400)$$

$$3 \times \frac{5}{2}R(T_f - 500) = 15600$$

$$T_f = 900 \text{ K}$$

$$P_f = \frac{900 \times 0.0821 \times 7}{7}$$

$$P_f = 73.89 \text{ atm}$$

12. (c) $1000 \sqrt{\frac{3RT}{M}} \times 10^3 = \frac{3 \times 8.314 \times T}{10^{26} \times 6 \times 10^{23}}^{1/2}$

$$T = \frac{10^6 \times 10^{26} \times 6 \times 10^{23}}{3 \times 8} = 250 \text{ K}$$

13. (b) Mass of balloon = 100 kg = $10 \times 10^4 \text{ g}$

$$\text{Volume of balloon} = \frac{4}{3} \pi r^3$$

$$\frac{4}{3} \pi \frac{22}{7} \frac{21^3}{2}$$

$$4851 \text{ m}^3 = 4851 \times 10^3 \text{ L}$$

$$\text{Mass of gas (He) in balloon} = \frac{PVM}{RT}$$

$$\frac{1 \times 4851 \times 10^3 \times 4}{0.082 \times 300}$$

$$78.878 \times 10^4$$

Total mass of gas and balloon

$$78.878 \times 10^4 + 10 \times 10^4$$

$$88.878 \times 10^4 \text{ g or } 888.78 \text{ kg}$$

$$\text{Mass of air displaced} = 1.2 \times 4851$$

$$5821.2 \text{ kg}$$

$$\text{Pay load} = \text{mass of air displaced}$$

$$\text{mass of (balloon + gas)}$$

$$\text{Pay load} = 5821.2 - 888.78 = 4932.42 \text{ kg}$$

14. (d) Let V mL of gas effused

$$\frac{V/220}{V/200} = \sqrt{\frac{d_{\text{O}_2}}{d_{\text{mix}}}} \Rightarrow d_{\text{mix}} = 1.6 \quad (1.1)^2 = 1.936 \text{ g/L}$$

Let density of ozone is d ; In 100 volume ozonised oxygen, 60% O₂ and 40% by volume O₃ is present

$$\text{Mass of mixture} = \text{mass of ozone}$$

$$+ \text{mass of oxygen}$$

$$100 \times 1.936 = 40 \times d + 60 \times 1.6$$

$$\text{density of O}_3 \text{ is } 2.44 \text{ g/L}$$

$$\frac{708.2}{760} = 0.25$$

15. (d) $n_{\text{N}_2} = \frac{760}{0.0821 \times 303} = 9.36 \times 10^3$

$$\frac{756.2}{760} = 0.3$$

$$n_{\text{O}_2} = \frac{0.3}{(0.0821) \times 298}$$

$$0.0122$$

$$n_{\text{total}} \text{ moles} = 0.02156$$

$$\text{pressure in final vessel} = P$$

$$\frac{(n_{\text{total}})RT}{V} = \frac{0.02156 \times 0.0821 \times 308}{0.5}$$

$$P = 1.09 \text{ atm or } 828.4 \text{ torr}$$

$$P_{\text{total}} = P_{(\text{O}_2 + \text{N}_2)} = \text{V.pr. of H}_2\text{O}$$

$$828.4 - 42.2 = 786.2 \text{ torr}$$

16. (b) $P = P_0 e^{kt}$; $P_0 = \frac{228}{3} = 76 \text{ cm Hg}$

$$\text{At } t = 14 \text{ min; } P = \frac{114}{3} \times 76 = 114 \text{ cm Hg}$$

$$152e^{kt} = 114$$

$$k = \frac{1}{t} \ln \frac{152}{114} = \frac{1}{14} \times 0.28$$


$$0.02 \times 60 \text{ hr} = 1.2 \text{ hr}^{-1}$$

$$17. (d) P_{\text{H}_2\text{O}} = 760 - 722 \times \frac{38}{760} \text{ atm; } n_{\text{H}_2\text{O}} = \frac{0.9}{18}$$

$$V = \frac{0.9}{18} \times \frac{0.821}{38} \times 300 = 760; V = 24.63 \text{ L}$$

$$18. (c) n_{AB} = \frac{3 \times 8.21}{0.0821 \times 300} = 1;$$

$$n_{B_2} = \frac{1 \times 8.21}{0.0821 \times 400} = 0.25$$



$$\text{Initial moles} = 1 \quad 0.25$$

$$\text{After reaction} = 0.5 \quad 0$$

$$P_{AB} = (8.21 + 8.21) = 0.5 \times 0.0821 \times 250$$

$$P_{AB} = 0.625 \text{ atm}$$

$$19. (c) \text{Moles of } O_2 \text{ remaining after reaction} = \frac{4}{4} - \frac{1}{1} = 3$$

V is due to same T and P in each compartment after reaction.

Length of He compartment

$$\frac{5}{5} \times 500 = 312.5 \text{ cm}$$

$$20. (b) V_c = \frac{60}{0.80} = 75 \text{ cm}^3 \text{ mol}^{-1};$$

$$b = \frac{V_c}{3} = 25 \text{ cm}^3 \text{ mol}^{-1} = 0.025 \text{ L mol}^{-1}$$

$$T_c = \frac{8a}{27Rb}$$

$$\frac{4 \times 10^5}{821} = \frac{8 \times a}{27 \times 0.0821 \times 0.025}$$

$$a = 3.375$$

$$21. (b) b = 4 \times \frac{4}{3} r^3 N_A;$$

$$4 \times 10^4 = 1000$$

$$4 \times \frac{4}{3} r^3 = 6 \times 10^{23}$$

$$r = 5 \times 10^{-9} \text{ cm}$$

Distance of closest approach = $2r$

$$10^{-8} \text{ cm} \quad \text{or} \quad 10^{-10} \text{ m}$$

$$22. (c) \frac{r_x}{r_{O_2}} = \sqrt{\frac{M_{O_2}}{M_x}} = \frac{4}{5} \Rightarrow \frac{32}{M_x} = \frac{16}{25} \Rightarrow M_x = 50$$

$$d_x = 0.80 \text{ kg/m}^3,$$

$$V_m = \frac{1000}{800} \times 50 = 62.5 \text{ L}$$

$$Z = \frac{PV_m}{RT} = \frac{1 \times 62.5}{0.0821 \times 500} = 1.52$$

$$23. (a) b = 25 \text{ cm}^3/\text{mol}$$

$$b = 0.025 \text{ L/mol}$$

$$V_{\text{Real}} = V_i - nb = 50 - 10 \times 0.025 = 49.75 \text{ L}$$

$$24. (b) Z = 1 - \frac{Pb}{RT} \text{ at high pressure}$$

$$\text{slope} = \frac{b}{RT} = \frac{0.025}{492.6}$$

$$b = \frac{492.6 \times 0.0821}{300} \text{ and}$$

$$b = \frac{4}{3} r^3 \times 4 N_A$$

$$\frac{492.6}{0.0821 \times 300} = 10^3$$

$$r = 2.5 \text{ \AA} \text{ or } d = 5 \text{ \AA}$$

$$26. (a) n = \frac{P}{RT} V$$

$$\text{Slope of line} = \frac{P}{RT} \tan$$

$$T = \frac{P}{R \tan} = \frac{16.42}{0.0821 \tan}$$

$$\text{For B, } T = 200\sqrt{3} \text{ K}$$

$$\text{For A, } T = \frac{200}{\sqrt{3}} \text{ K}$$

Level 3

Passage-1

$$1. (d) p = \frac{a}{V_m^2} (V_m - b) \quad RT$$

$$P = \frac{a}{V_m^2} (V_m - b) \quad P = \frac{a}{V_m^2} p$$

$$P(V_m - b) = RT$$

$$PV_m - RT = Pb$$

$$2. (a) V = \frac{P}{nb}$$

$$V = nb \quad V$$

$$P = \frac{n^2 a}{V^2} \quad V = nRT$$

$$PV = \frac{n^2 a}{V} = nRT$$

$$PV = nRT = \frac{n^2 a}{V}$$

3. (c) van der Waal's constant 'a' I.M.A.F.

So CO_2 gas can be more easily liquified.

4. (d) $P = \frac{n^2 a}{V^2} (V - nb) = nRT$

Passage-2

- 2 & 3. Let A , B and C be collision diameters of A , B and C

$$\frac{kT}{\sqrt{2}} \cdot \frac{1}{2P}, \text{ i.e., } \sqrt{\frac{T}{P}}$$

$$\sqrt{\frac{1600}{0.16}} : \sqrt{\frac{200}{0.16}} : \sqrt{\frac{400}{0.04}}$$

$$4 : 1 : 2$$

2. (c) $Z_{11} = \frac{2P^2}{\sqrt{mT}^{3/2}}$

$$A : B : C = \frac{16}{\sqrt{20}} : \frac{1}{(1600)^{3/2}} :$$

$$\frac{1}{\sqrt{40}} : \frac{4}{(200)^{3/2}} : \frac{4}{\sqrt{80}} : \frac{16}{(400)^{3/2}}$$

$$= 1 : 4 : 16$$

3. (a) $Z_1 = \frac{(P^2)}{\sqrt{MT}}$

$$A : B : C = \frac{0.6}{\sqrt{20}} : \frac{1}{1600} : \frac{1}{\sqrt{40}} : \frac{2}{200}$$

$$: \frac{4}{\sqrt{80}} : \frac{4}{400}$$

$$4 : 1 : 4$$

Passage-3

1. (a) $h_L d_L = \frac{h_{\text{Hg}} d_{\text{Hg}}}{5.44} = 190 \text{ cm}$

2. (d) $P_{\text{Gas}} = P_{\text{Atm}} = P_L = 1 - \frac{38}{190} = 1.2$

3. (b) $3A(g) \rightleftharpoons A_3(g)$
- | | | | |
|----------------------|----------|---------------------|----------|
| $t = 0$ | 1.2 atm | $A_3(g)$ | |
| $t = t_{\text{eq.}}$ | 1.2 0.36 | $\frac{1}{3}(0.36)$ | 0.12 atm |

$$P_T = 1.2 + 0.36 + 0.12 = 0.96 \text{ atm}$$

Pressure difference in column

$$1 - 0.96 = 0.04 \text{ atm}$$

The difference in height of the liquid level in two columns 0.04 190 7.6 cm

One or More Answers is/are Correct

1. (a, b, c, d)

For Boyle's law

T constant

$PV = K$ constant

2. (a, b)

$PV = nRT$

T constant

PV constant
 V P

KE is a function of temperature.

3. (b, c, d)

T_c is a temperature above which a gas can not be liquified at this point, slope of $P-V$ curve is maximum and attraction force is dominating at this point. Both liquid vapour are exist in equilibrium.

4. (c, d)

at constant volume at constant P

$P = T$ $V = T$

$P = KT$ $V = KT$

$P = K(t - C - 273.15)$ $V = K(t - C - 273.15)$

5. (a)

T_B is a temperature at which gas behave like an ideal gas mean attractive and repulsing forces just offset each other.

6. (a, c, d)


$$KE = \frac{3}{2} RT \text{ (1 mole)} U_{\text{rms}} = \sqrt{\frac{3RT}{M_w}}$$

$$d = \frac{PM_w}{RT}$$

7. (a, b)

Refer. Q.3

8. (a, b, c, d)



9. (a) Refer Q.2

10. (a, b, c)

$PV = nRT$

R is constant and independent of P and T

$R = K N_A$

K Boltzmann's constant

11. (a,b,d)

(A) $\therefore$ area under the curve gives fraction of molecules and total area is constant.(B) u_{mp} decreases with decrease in temperature.(C) T_2 is higher temperature

(D) As seen from graph; A, B, D

12. (a, d)

Vapour pressure of liquid is only and only temperature dependent.

13. (a, b, d)

For H_2 gas IMAF 0

$Z = 1$ $Z = 1$ $\frac{Pb}{RT}$

at low pressure $Z = 1$ $Z = 1$ $\frac{a}{RTV_m}$

Real gas behave like an ideal gas at low pressure and high temperature.

14. (a, b, c)

 $T_C = \frac{8a}{27Rb}$ Refer. Q.14

15. (a,b,d)

 P_T (1 3) 1 3 0.1 1.3 atm. P 0.2 atm or 76 0.3 cm of Hg

or 760 0.3 mm of Hg

16. (a, b, c, d)

Refer. Q. No. 13

17. (a, d)

Critical pressure is the minimum pressure at which substance convert into liquid state at critical point.

18. (a,c)

Low P and high temperature gas behaves as an ideal gas. PV constant and $\frac{PV_m}{RT} = 1$.

19. (b, c)

For ideal gas $Z = 1$

at low pressure and Boyle's temperature real gas behave like an ideal gas.

20. (a, b, c)

 $V \propto T$ at constant P $KE = \frac{3}{2} RT$ (1 mole)

Match the Column

5. $y = \frac{1}{V^2}$ or $\sqrt{y} = \frac{1}{V}$

$P \propto x$ and $P \propto \frac{\text{constant}}{V}$

(A) $x = (k)\sqrt{y}$ $y = k^1 x^2$

(B) $V \propto kT$; $y = V$ & $\frac{1}{T} \propto x$ $y = \frac{k}{x}$

(C) $P \propto kT$; $PT \propto kT^2$ or $y \propto kx$

(D) $v \propto \frac{c}{p}$ $y \propto c\sqrt{x}$; $y^2 \propto cx$

Subjective Problems

1. Refer Q.No. 59 Level-1

2. For H_2 and He.

$$f^{4th} = \frac{x/y}{1} \sqrt{\frac{M_{He}}{M_{H_2}}}$$

$$\frac{x}{y} = (\sqrt{2})^4 = 4$$


3. Equation of line

$P = 2 \frac{10}{4} \frac{2}{15} (V - 15)$

$P = 2 \frac{8}{11} (V - 15)$

$P = 2 \frac{8V}{11} - \frac{15 \cdot 8}{11}$

$P = \frac{142}{11} - \frac{8V}{11}$



$f(T) = \frac{1}{nR} \frac{142V}{11} - \frac{8V^2}{11}$

$\frac{d(f(T))}{dV} = \frac{1}{nR} \frac{142}{11} - \frac{8V}{11} = 0$


$V = \frac{142}{11 \cdot 16} = 8.875$

$P = \frac{142}{11} - \frac{8}{11} \cdot \frac{8.875}{16} = \frac{71}{11}$

$$T_{Max} = \frac{PV}{nR} = \frac{71}{11} \cdot \frac{142}{16} \cdot \frac{1}{0.0821} = 700$$

$$7 \cdot 10^2 K$$

$$4. \quad T = 300 \quad \frac{4}{100} \quad \frac{3}{100} (P = 3)$$



$$T = 300 \quad \frac{P}{100} \quad \frac{3}{100}$$

Multiply by 100

$$T = 100 \quad 3 \quad 10^4 \quad P = 3$$

$$P = 3 \quad 3 \quad 10^4 \quad 100T$$

$$\frac{dP}{dT} = 0 \quad 100 \quad 1 \quad 10^2$$

$$x = y = 3$$

$$5. \quad P = \frac{n^2 a}{V^2} \quad P_i$$

$$\frac{n^2 a}{V^2} = 0.04$$

$$a = 4$$

$$6. \quad Z = 1 \quad \frac{Pb}{RT} \quad \text{high pressure}$$

$$\frac{dZ}{dP} = \frac{b}{RT} \quad \frac{1}{2.8}$$

$$b = \frac{RT}{2.8} \quad \frac{22.4}{2.8} \quad 4 \quad N_A \quad \frac{4}{3} \quad R^3$$

$$N_A = \frac{4}{3} \quad R^3 \quad \text{Volume of 1 mole gas}$$

$$\frac{5.6}{2.8} \quad 2$$

$$7. \quad P = \frac{dRT}{M_w}; \quad \frac{P_x}{P_y} = \frac{d_x}{d_y} \quad \frac{M_y}{M_x} \quad 6$$

$$8. \quad \frac{r_x}{r_{O_2}} = \sqrt{\frac{M_{O_2}}{M_x}}$$

$$\frac{V}{2} = \frac{5.65}{V} \quad \sqrt{\frac{32}{M}}$$

$$8 \quad \frac{32}{M}$$

$$M = 4$$

$$9. \quad n_{Br_2} = \frac{1}{0.0821} \quad \frac{0.423}{423} \quad \frac{1}{82.1} \quad \text{mole}$$

POAC on 'Br'

$$\frac{1}{82.1} \quad 2 \quad \frac{4.2}{80} \quad \frac{1}{19n}$$

$$10. \quad \frac{n_T}{910} \quad \frac{n_1}{(10 \quad V)} \quad \frac{n_2}{950 \quad V} \quad \frac{1}{900 \quad 10} \quad \frac{1}{RT} \quad \frac{1}{RT} \quad \frac{1}{RT}$$

$$11. \quad P_f = 1 \quad \frac{36}{76} \quad \frac{112}{76} \quad \text{atm. Final height} \quad 19 \text{ cm}$$

$$P_i = 1 \text{ atm, initial length} \quad h_i \text{ cm}$$

$$\text{Boyle's law} \quad P_i V_i = P_f V_f$$

$$1 \quad h_i A \quad \frac{112}{76} \quad 19A$$

$$h_i = 28 \text{ cm}$$

The length by which the Hg column shifts

$$\text{down} \quad h_i \quad h_f$$

$$28 \quad 19 \quad 9 \text{ cm}$$

$$12. \quad P_i = 722 \text{ mm} \quad P_f = 760 \text{ mm}$$

$$T_i = 107 \quad 273 \quad 380 \text{ K} \quad T_f = 100 \text{ K}$$

$$d_i = 1 \text{ g/cm}^3 \quad d_f = ?$$

$$\frac{d_i T_i R}{P_i} = \frac{d_f T_f R}{P_f}$$

$$d_f = \frac{P_f}{P_i} \quad \frac{T_i}{T_f} \quad d_i \quad \frac{760}{722} \quad \frac{380}{100} \quad 1 \text{ g/cm}^3$$

$$4 \text{ g/cm}^3$$

$$13. \quad V, n \text{ constant.}$$

$$\frac{P_i}{T_i} = \frac{P_f}{T_f} \quad P_f = \frac{T_f}{T_i} P_i \quad \frac{109}{100} P_i$$

$$P_{\text{increases}} = P \quad P_f = P_i \quad \frac{9}{100} P_i$$

% Pressure increases

$$\frac{P_f}{P_i} = 100 \quad \frac{9P_i}{100P_i} = 100\%$$

$$X\% = 9\%$$

$$X = 9$$

$$14. \quad T_B = \frac{a}{Rb} \quad \frac{3.6}{0.08 \quad 0.6} \quad 75 \text{ K}, \quad \frac{75}{15} \quad 5 \text{ K}$$

$$15. \quad 3 \text{ ms}^{-1}$$


$$u_{\text{rms}} = \sqrt{\frac{u_1^2 N_1}{N_1} + \frac{u_2^2 N_2}{N_2}} \quad u_{\text{rms}}^2 = \frac{u_1^2 N_1}{N_1} + \frac{u_2^2 N_2}{N_2}$$

$$25 = \frac{4 \quad 7^2 \quad u_2^2}{10} \quad u_2 = \sqrt{\frac{54}{6}} \quad 3 \text{ ms}^{-1}$$

4

THERMODYNAMICS

- ❖ The subject of **Thermodynamics** deals basically with the interaction of one body with another in terms of quantities of heat & work.
- ❖ A **System** is defined as that macroscopic part of the universe which is at the moment under investigation.
- ❖ **Surroundings** are the rest of the universe outside the system.
- ❖ **Boundary** : The actual or imaginary surface that separates the system from the surroundings.



Thermodynamic Properties

Intensive Properties

do not depend on the amount of matter present in system

Examples : Pressure, Temperature, Density, Melting point, Boiling point etc.

Extensive Properties

depend on the amount of matter present in the system.

Examples : Mass, Volume, Heat Capacity, Enthalpy, Entropy etc.

Thermodynamic Process : The change of a system from one state to another state.

Thermodynamic Processes

Isothermal

temperature remains constant throughout the process
i.e., : $dT=0$

Isobaric

pressure remains constant throughout the process
i.e., : $dP=0$

Isochoric

volume remains constant throughout the process
i.e., : $dV=0$

Adiabatic

system does not exchange heat with surrounding
i.e., : $dq=0$

Polytropic

$$PV^n = \text{constant}$$

❖ **Cyclic Process :** System undergoes a number of different processes and finally returns to its initial state.

❖ **REVERSIBLE PROCESS (QUASI-STATIC) & IRREVERSIBLE PROCESS :**

A process which is carried out so slowly that the system and the surroundings are nearly in equilibrium is known as a **reversible Process**. If this condition does not hold good, the process is said to be, **irreversible**.

In a reversible process the driving force is infinitesimally larger than the opposing force. If the driving force is made infinitely smaller than opposing force, the system can be brought back without producing any permanent change.

❖ **All natural processes are irreversible process.**

Spontaneous Process : Proceeds on its own i.e., without any external help.

❖ **INTERNAL ENERGY (U) :** The sum of all microscopic forms of energy.

$$U = U_{\text{Translational}} + U_{\text{Rotational}} + U_{\text{Vibrational}} + U_{\text{Bonding}} + U_{\text{Electronic}} + \dots$$

❖ U is a state function & an extensive property.

$$U = U_{\text{Final}} - U_{\text{Initial}} ; U = q_v$$

$$❖ dU = \frac{U}{T} dT + \frac{U}{V} dV, dU = nC_{vm}(T_2 - T_1) \text{ at constant volume}$$

Enthalpy

Chemical reactions are generally carried out at constant pressure (atmospheric pressure) so it has been found useful to define a new state function called Enthalpy (H) as :


- ❖ $H = U + PV$; $H = U + (PV)$; at constant pressure $H = U + P \Delta V$; combining with first law. $H = q_p$
- ❖ The difference between H & U becomes significant only when gases are involved (insignificant in solids and liquids)

$$\frac{H}{H} = \frac{U}{U} + \frac{(PV)}{(n_g)RT} = 1 + \frac{nR}{n_g} \frac{T}{T}$$

- ❖ For a given system consider; $H = f(T, P)$; $dH = \left(\frac{\partial H}{\partial T}\right)_P dT + \left(\frac{\partial H}{\partial P}\right)_T dP$
- ❖ For isobaric process : $dP = 0$; $dH = \left(\frac{\partial H}{\partial T}\right)_P dT$; $dH = C_p \cdot dT$; $H = \int C_p \cdot dT$
- ❖ For an ideal gas $H = f(T)$ only
 $\left(\frac{\partial H}{\partial P}\right)_T = 0$ or $dH = C_p \cdot dT$; $H = \int C_p \cdot dT$; for n moles of ideal gas $H = nC_p \cdot dT$

Kirchoff's Equation

- ❖ For chemical reactions $d_r H = \sum_r C_p \cdot dT$
- ❖ ${}_r H_{T_2} - {}_r H_{T_1} = \sum_r C_p (T_2 - T_1)$
- ❖ Similarly ${}_r U_{T_2} - {}_r U_{T_1} = \sum_r C_v (T_2 - T_1)$



❖ ZEROth LAW OF THERMODYNAMICS

Two systems in thermal equilibrium with a third system, are also in thermal equilibrium with each other.


❖ FIRST LAW OF THERMODYNAMICS

It is law of conservation of energy. Mathematically for **closed system** at rest in absence of external fields this law is written as :

$$\Delta U = q + w$$

Where ΔU is change in internal energy of the system, q is the transfer of heat from surroundings to the system and w is the work involved (either done on the system or by the system).

❖ **Sign convention according to IUPAC :**



❖ **HEAT :** The energy that flow into or out of a system because of a difference in temperature between the system and its surrounding.

❖ **TYPES OF WORK :** Two types of work normally come across in chemistry.

1. Mechanical work (e.g., PV work)

2. Electrical work (Non PV work)

❖ *Mechanical work* is involved when a system changes its volume in the presence of an externally applied pressure (i.e., pressure volume work). It is especially important in system containing gases

$$dw = P_{\text{ext}} \cdot dV ; \quad w = \int_{V_1}^{V_2} P_{\text{ext.}} dV$$

❖ **CALCULATION OF WORK DONE IN VARIOUS PROCESS**

❖ **Isochoric process:**

Since $dV = 0$ so $w = 0$
 from 1st law $U = q_v$ $w = q_v$

❖ **Isobaric process:**

$w = P_{\text{ext}}(V_2 - V_1)$
 & $H = q_p$

❖ **Isothermal process:**

Since $dT = 0$ so $dU = 0$ for an ideal gas
 from 1st law $q = w$ 0 or $q = w$

If process is reversible

$$w = nRT \ln \frac{V_2}{V_1} = nRT \ln \frac{P_1}{P_2} \quad \text{or} \quad w = 2.303 nRT \log \frac{V_2}{V_1} = 2.303 nRT \log \frac{P_1}{P_2}$$

If process is irreversible

$$w = P_{\text{ext}} \frac{nRT}{P_2} = \frac{nRT}{P_1}$$

In case of free expansion $P_{\text{ext}} = 0$ so $w = 0$

❖ **Adiabatic process : $q = 0$**

$$U = w = nC_v(T_2 - T_1) = \frac{P_2 V_2 - P_1 V_1}{\gamma - 1}, \text{ where } \gamma = \frac{C_p}{C_v}$$

If process is reversible $T_1 V_1^{\gamma} = T_2 V_2^{\gamma}$


If process is irreversible $\frac{T_2}{1} \frac{T_1}{P_1} = P_{\text{ext}} \cdot \frac{T_2}{P_2} \frac{T_1}{P_1}$

❖ **Cyclic process:**

Work done = Area enclosed in PV -diagram

For clockwise it is +ve

For anti-clockwise it is -ve



❖ Electrical work ($w = V.I.t$) is a type of non PV -work.

Heat Capacity : Between any two temperatures, the quantity of heat required to raise the temperature of the system from the lower to the higher temperature divided by the temperature difference.

❖ $C = dq/dT$.

❖ $q_v = U$

❖ $C_v = \frac{q}{T_2 - T_1} = \frac{U}{T_2 - T_1}$ or $C_v = (U/T)_v$

❖ $C_p = \frac{H}{T}$

❖ $C_{p,m} = C_{v,m} + R$ (for ideal gas)

❖ $C_{p,m}$ and $C_{v,m}$ depends on temperature even for an ideal gas. ($C = a + bT + cT^2$)

❖ $C_{\text{mix}} = \frac{n_1 C_{p1} + n_2 C_{p2} + \dots}{n_1 C_{v1} + n_2 C_{v2} + \dots}$

Reversible Adiabatic process involving ideal gas

❖ $T_1 V_1^{\gamma-1} = T_2 V_2^{\gamma-1}$

❖ $TV^{\gamma-1} = \text{constant}$

❖ $PV^{\gamma} = \text{constant}$

❖ $T P^{1/\gamma} = \text{constant}$

❖ **Molar Specific heat (C_m) of ideal gas in any Reversible Polytropic Process :**

$$C_m = C_{v,m} + \frac{R}{1-x} = \frac{R}{1} + \frac{R}{1-x}$$

Second Law of Thermodynamics

Statements :

- No cyclic engine is possible which take heat from one single source and in a cycle completely convert it into work without producing any change in surroundings.
- In an irreversible process, entropy of universe increases but it remains constant :
in a reversible process

$$S_{\text{sys}} + S_{\text{surr}} = 0 \quad \text{for reversible process}$$

$$S_{\text{sys}} + S_{\text{surr}} > 0 \quad \text{for irreversible process}$$

$$S_{\text{sys}} + S_{\text{surr}} = 0 \quad (\text{In general})$$

(iii) Efficiency of Carnot engine working reversibly is maximum.

Carnot cycle


$$AB \quad \text{Isothermal Reversible Expansion} \quad w_{AB} \quad nRT_2 \ln \frac{V_2}{V_1}$$

$$BC \quad \text{Adiabatic Reversible Expansion} \quad w_{BC} \quad C_V(T_1 - T_2)$$

$$CD \quad \text{Isothermal Reversible Expansion} \quad w_{CD} \quad nRT_1 \ln \frac{V_4}{V_3} \quad T_2 \quad T_1$$

$$DA \quad \text{Adiabatic Reversible Expansion} \quad w_{DA} \quad C_V(T_2 - T_1)$$

$$\text{for Carnot efficiency () calculation} \quad \frac{w_{\text{Total}}}{q_2} \quad \frac{T_2 - T_1}{T_2} \quad \frac{q_2 - q_1}{q_2}$$



Calculation of Entropy change (S or S_{sys})

Ideal gas in any Reversible/Irreversible process from (P_1, V_1, T_1) to (P_2, V_2, T_2)

$$\diamond S = nC_V \ln \frac{T_2}{T_1} + nR \ln \frac{V_2}{V_1} = nC_P \ln \frac{T_2}{T_1} + nR \ln \frac{P_1}{P_2}$$

$\diamond$ Reversible & irreversible isothermal expansion / contraction of an ideal gas

$$S = nR \ln \frac{V_2}{V_1}$$

$$\diamond \text{ Isobaric heating or cooling : } S = nC_P \ln \frac{T_2}{T_1}$$

$$\diamond \text{ Isochoric heating or cooling : } S = nC_V \ln \frac{T_2}{T_1}$$

$\diamond$ Adiabatic process :

$$\diamond S = nC_V \ln \frac{T_2}{T_1} + nR \ln \frac{V_2}{V_1} \text{ for irreversible process}$$

$S = 0$ for reversible adiabatic compression and expansion.

$\diamond$ General reversible heating or cooling of any substance

$$S = \int_{T_1}^{T_2} \frac{dq_{\text{rev}}}{T}$$

$\diamond$ Reversible Phase Transformation

$$(a) S_{\text{Fusion}} = H_{\text{Fusion}} / T_m, \quad T_m \text{ is normal melting point (K)}$$

$$(b) S_{\text{vap}} = H_{\text{vap}} / T_B, \quad T_B \text{ is normal boiling point (K)}$$

$$(c) S_{\text{sub}} = H_{\text{sub}} / T_{\text{sub}}, \quad T_{\text{sub}} \text{ is sublimation temperature (K)}$$

$$(d) S_{\text{Trans}} = H_{\text{Trans}} / T_{\text{Trans}}, \quad T_{\text{Trans}} \text{ is the phase transition temperature (K)}$$

$\diamond$ For chemical reaction

$${}_r S \quad {}_p S \quad {}_f S \text{ (products)} \quad {}_R S \quad {}_f S \text{ (reactants)}$$

Third Law of Thermodynamics

“At absolute zero temperature, the entropy of a perfectly crystalline substance is taken as zero”, which means that at absolute zero every crystalline solid is in a state of perfect order and its entropy should be zero.

By virtue of the third law, the absolute value of entropy (unlike absolute value of enthalpy) for any pure substance can be calculated at room temperature.

$$S_T = S_{0K} + \int_{0K}^T \frac{q_{rev}}{T}, \text{ since } S_{0K} = 0; S_T = \int_{0K}^T \frac{q_{rev}}{T}$$

Absolute entropies of various substances have been tabulated and these value are used to calculate entropy changes for the reactions by the formula;

$$\Delta S = \sum \nu_P S_m^\circ (\text{products}) - \sum \nu_R S_m^\circ (\text{reactants})$$

Variation of S_r with temperature & pressure :
$$dS_r = \frac{C_{p,m} dT}{T}$$

$$S_{T_2} - S_{T_1} = C_{p,m} \ln \frac{T_2}{T_1}$$

Gibbs Free Energy (G) and Spontaneity

- ❖ A new thermodynamic state function G , the Gibbs free energy is defined as :
 $G = H - TS$ or $G = H - T S$ (at constant temperature and pressure)
- ❖ $G < 0$ for spontaneous process
- ❖ $G = 0$ for equilibrium.

Criteria for Spontaneity

For a feasible process $\Delta S_T = \Delta S_{sys} + \Delta S_{surr} > 0$

In absence of non PV work at constant T and P for a feasible process $\Delta G < 0$.

Standard Free Energy Change (ΔG°)

The free energy change for a process at a specified temperature in which the reactants in their standard state are converted to the products in their standard state. It is denoted by ΔG° .

❖ Standard states :

- (a) Standard temperature (any specified temperature)
- (b) For gases/solid/liquid: Standard pressure $P^\circ = 1 \text{ bar}$
- (c) For ion/substance in solution: Standard concentration $(C^\circ) = 1M$

Like the standard enthalpy of formation of an element “the standard free energy of formation of an element in its standard state is zero”. And so;

$$\Delta_r G^\circ = \sum \nu_P \Delta_f G^\circ (\text{products}) - \sum \nu_R \Delta_f G^\circ (\text{reactants})$$

A Thermodynamic Relationship

For any reversible reaction : $\Delta_r G^\circ = -RT \ln Q$.

where Q is reaction quotient.

At equilibrium $\Delta_r G^\circ = 0$, $Q = K$


$$\Delta_r G^\circ = -RT \ln K, \quad \Delta_r G^\circ = -2.303 RT \log K_{eq}.$$

Effect of Temperature : Van't Hoff Equation

$$(a) \frac{d(\ln K)}{dT} = \frac{H}{RT^2} \quad (b) \frac{d(\ln K)}{d\left(\frac{1}{T}\right)} = \frac{H}{R}$$

$$G = H - TS = \ln K = \frac{H}{RT} - \frac{S}{R}$$

Integrated form $\ln \frac{K_2}{K_1} = \frac{H}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$



❖ RELATIONSHIP BETWEEN G & $w_{\text{non-PV}}$

$$dU = dq + dw_{\text{non-PV}}$$

for reversible process at constant T & P

$$dU = pdV + TdS + dw_{\text{non-PV}}$$

$$dH = TdS + dw_{\text{non-PV}}$$

$$(dG_{\text{system}})_{T,P} = dw_{\text{non-PV}}$$

$$(dG_{\text{system}})_{T,P} = (dw_{\text{non-PV}})_{\text{by system}}$$

Means free energy change for a process is equal to the maximum possible work that can be derived from the process i.e.,

- ❖ $G = w_{\text{max}}$ (for a reversible change at constant pressure and temperature)
- ❖ In case of a galvanic cell. $G = -nFE_{\text{cell}}$, where E_{cell} e.m.f. of the cell ; F Faraday constant and n number of electrons being transferred in the chemical process
- ❖ So $G = -nFE_{\text{cell}}^{\circ}$, where E_{cell}° is the standard cell potential.

THERMOCHEMISTRY

❖ Enthalpy of Reaction :

The amount of heat evolved or absorbed when the number of moles of reactants react completely to give the products as given by the balanced chemical equation.

❖ Enthalpy of formation :

The change in heat enthalpy when one mole of a substance is formed directly from its constituent elements at standard states.

(i) For elements, ${}_fH = 0$; e.g., ${}_fH (\text{Zn}, \text{S}) = 0$, ${}_fH (\text{Br}_2, \text{l}) = 0$, ${}_rH (\text{O}_2, \text{g}) = 0$

(ii) If element exists in two or more allotropic forms then for most stable state ${}_fH = 0$
e.g., ${}_fH [\text{C}(\text{s}, \text{graphite})] = 0$, ${}_fH [\text{S}(\text{s}, \text{rhombic})] = 0$

❖ Enthalpy of combustion :

The change in enthalpy when one mole of a substance is completely burnt in oxygen.

❖ Enthalpy of Transition :

The enthalpy change when one mole of one allotropic form changes to another.

For example : $\text{C}(\text{graphite}) \rightarrow \text{C}(\text{diamond})$; $H_{\text{trs}} = 1.90 \text{ kJ mol}^{-1}$.

❖ Enthalpy of fusion :

The change in enthalpy when one mole of a solid is completely converted into liquid state at its melting point.

❖ **Enthalpy of vaporisation :**

The change in enthalpy when one mole of a liquid is completely converted into its vapours at its boiling point.

❖ **Enthalpy of sublimation :**

The change in enthalpy when one mole of a solid is directly converted into vapours at sublimation temperature.

❖ **Enthalpy of solution :**

The change in enthalpy when one mole of a substance is dissolved in excess of water so that further dilution does not involve any heat change.

❖ **Enthalpy of Dilution :**

The change in enthalpy when a solution containing 1 mole of solute is diluted from one concentration to another.

❖ **Enthalpy of Hydration : (For anhydrous salt)**


The enthalpy change when 1 mole anhydrous salt combines with the requisite amount of water to form a new hydrated stable salt. For example, the hydration of anhydrous cupric sulphate is represented by

❖ **Enthalpy of neutralisation :**

Enthalpy of neutralisation is the change in enthalpy when one gram equivalent of an acid is completely neutralised by one gram equivalent of a base in dilute solutions.

❖ **Enthalpy of formation of ions :**

The enthalpy change when one mole of aqueous ions is obtained from element in its standard state as.



By convention, the standard enthalpy of formation of $\text{H}^+(aq)$ is taken to be zero.

❖ **Hess's law of constant heat summation :**

The total heat change in a particular reaction is the same whether the reaction takes place in one step or in a number of steps.



According to Hess's Law $H = H_1 + H_2$.

❖ **Enthalpy of atomisation :**

The change in enthalpy when one mole of substance converts into gaseous atoms.

❖ **Bond enthalpies :**

The average of enthalpies required to dissociate the bond present in different gaseous compounds into free atoms or radicals in the gaseous state.

The bond enthalpy given for any particular pair of atoms is the average value of the dissociation enthalpies of the bond for a number of molecules in which the pair of atoms appears.

❖ **Estimation of Enthalpy of a reaction from bond Enthalpies :**

For the gaseous reaction; $\text{C}_2\text{H}_4(\text{g}) + \text{HCl}(\text{g}) \rightarrow \text{C}_2\text{H}_5\text{Cl}(\text{g})$	
H Energy required to break reactants into gaseous atoms	H Energy released to form products from the gaseous atoms

❖ **Resonance Enthalpy :**



$H_{\text{Resonance}}$	$H_{f, \text{experimental}}$	$H_{f, \text{calculated}}$
	$H_{c, \text{calculated}}$	$H_{c, \text{experimental}}$



Level 1


- Out of molar entropy (I), specific volume (II), heat capacity (III), volume (IV), extensive properties are :
 (a) I, II (b) I, II, IV (c) II, III (d) III, IV
- Out of internal energy (I), boiling point (II), pH (III) and E.M.F. of the cell (IV) intensive properties are :
 (a) I, II (b) II, III, IV (c) I, III, IV (d) All of these
- Thermodynamic equilibrium involves
 (a) chemical equilibrium (b) mechanical equilibrium
 (c) thermal equilibrium (d) all the above simultaneously
- Which has maximum internal energy at 290 K?
 (a) Neon gas (b) Nitrogen gas (c) Ozone gas (d) Equal for all
- A 10 g piece of iron ($C = 0.45 \text{ J/g}^\circ\text{C}$) at 100°C is dropped into 25 g of water ($C = 4.2 \text{ J/g}^\circ\text{C}$) at 27°C . Find temperature of the iron and water system at thermal equilibrium.
 (a) 30°C (b) 33°C (c) 40°C (d) None of these
- When freezing of a liquid takes place in a system it:
 (a) may have $q = 0$ or $q < 0$ depending on the liquid
 (b) is represented by $q = 0$
 (c) is represented by $q < 0$
 (d) has $q = 0$
- Mechanical work is specially important in systems that contain
 (a) gas-liquid (b) liquid-liquid (c) solid-solid (d) amalgam
- Determine which of the following reactions taking place at constant pressure represents system that do work on the surrounding environment
 I. $\text{Ag} (aq) + \text{Cl} (aq) \rightarrow \text{AgCl}(s)$ II. $\text{NH}_4\text{Cl}(s) \rightarrow \text{NH}_3(g) + \text{HCl}(g)$
 III. $2\text{NH}_3(g) + \text{N}_2(g) \rightarrow 3\text{H}_2(g)$
 (a) I (b) III (c) II and III (d) I and II
- Determine which of the following reactions taking place at constant pressure represent surrounding that do work on the system environment
 I. $4\text{NH}_3(g) + 7\text{O}_2(g) \rightarrow 4\text{NO}_2(g) + 6\text{H}_2\text{O}(g)$
 II. $\text{CO}(g) + 2\text{H}_2(g) \rightarrow \text{CH}_3\text{OH}(l)$
 III. $\text{C}(s, \text{graphite}) + \text{H}_2\text{O}(g) \rightarrow \text{CO}(g) + \text{H}_2(g)$
 IV. $\text{H}_2\text{O}(s) \rightarrow \text{H}_2\text{O}(l)$
 (a) III, IV (b) II and III (c) II, IV (d) I and II, IV
- A sample of liquid in a thermally insulated container (a calorimeter) is stirred for 2 hr. by a mechanical linkage to a motor in the surrounding. For this process :
 (a) $w = 0; q = 0; U = 0$ (b) $w = 0; q = 0; U > 0$
 (c) $w = 0; q = 0; U < 0$ (d) $w = 0; q = 0; U = 0$
- A system undergoes a process in which $U = 300 \text{ J}$ while absorbing 400 J of heat energy and undergoing an expansion against 0.5 bar . What is the change in the volume (in L)?
 (a) 4 (b) 5 (c) 2 (d) 3

12. An ideal gas expands against a constant external pressure of 2.0 atmosphere from 20 litre to 40 litre and absorbs 10 kJ of heat from surrounding. What is the change in internal energy of the system? (Given : 1 atm-litre = 101.3 J)
 (a) 4052 J (b) 5948 J (c) 14052 J (d) 9940 J
13. One mole of an ideal gas at 25°C expands in volume from 1.0 L to 4.0 L at constant temperature. What work (in J) is done if the gas expands against vacuum ($P_{\text{external}} = 0$)?
 (a) 4.0×10^2 (b) 3.0×10^2 (c) 1.0×10^2 (d) Zero
14. At 25°C, a 0.01 mole sample of a gas is compressed from 4.0 L to 1.0 L at constant temperature. What is the work done for this process if the external pressure is 4.0 bar?
 (a) 1.6×10^3 J (b) 8.0×10^2 J (c) 4.0×10^2 J (d) 1.2×10^3 J
15. Calculate the work done (in J) when 4.5 g of H_2O_2 reacts against a pressure of 1.0 atm at 25°C
- $$2\text{H}_2\text{O}_2(l) \rightarrow \text{O}_2(g) + 2\text{H}_2\text{O}(l)$$
- (a) 1.63×10^2 (b) 4.5×10^2 (c) 3.2×10^2 (d) 6.1×10^2
16. 2 mole of zinc is dissolved in HCl at 25°C. The work done in open vessel is :
 (a) 2.477 kJ (b) 4.955 kJ (c) 0.0489 kJ (d) None
17. Temperature of 1 mole of a gas is increased by 2°C at constant pressure. Work done is :
 (a) R (b) 2R (c) R/2 (d) 3R
18. A sample of an ideal gas is expanded 1 m^3 to 3 m^3 in a reversible process for which $P = KV^2$, with $K = 6 \text{ bar/m}^6$. Work done by the gas is :
 (a) 5200 kJ (b) 15600 kJ (c) 52 kJ (d) 5267.6 kJ
19. A given mass of gas expands reversibly from the state A to the state B by three paths 1, 2 and 3 as shown in the figure. If w_1 , w_2 and w_3 respectively be the work done by the gas along three paths then:
 (a) $w_1 < w_2 < w_3$
 (b) $w_1 > w_2 > w_3$
 (c) $w_1 < w_2 > w_3$
 (d) $w_2 < w_3 < w_1$
20. Heat absorbed by a system in going through a cyclic process shown in figure is :
 (a) 10^7 J (b) 10^6 J
 (c) 10^2 J (d) 10^4 J




21. In the cyclic process shown in P - V diagram, the magnitude of the work done is :

(a) $\frac{P_2 - P_1}{2} (V_2 - V_1)^2$ (b) $\frac{V_2 - V_1}{2} (P_2 - P_1)^2$
 (c) $\frac{1}{4} (P_2 - P_1) (V_2 - V_1)$ (d) $(V_2 - V_1)^2$



22. An ideal gas is taken around the cycle $ABCA$ as shown in P - V diagram. The net work done during the cycle is equal to :





(a) $12P_1V_1$ (b) $6P_1V_1$
 (c) $5P_1V_1$ (d) P_1V_1











23. An ideal gas is at pressure P and temperature T in a box, which is kept in vacuum with in a large container. The wall of the box is punctured. What happens as the gas occupies entire container?

(a) It's temperature falls (b) Its temperature rises
 (c) Its temperature remains the same (d) Unpredictable



24. In diagrams (1 to 4), variation of volume with changing pressure is shown. A gas is taken along the path $ABCD$. The change in internal energy of the gas will be :



- (a) Positive in all the cases (1) to (4)
 (b) Positive in cases (1), (2), (3) but zero in case (4)
 (c) Negative in cases (1), (2), (3) but zero in case (4)
 (d) Zero in all the cases






25. If the door of a refrigerator is kept open, the room in which the refrigerator is kept
 (a) gets cooled
 (b) gets heated
 (c) neither gets cooled nor gets heated
 (d) gets cooled or heated depending on the initial temperature of the room
26. The temperature of an ideal gas increases in an :
 (a) adiabatic expansion
 (b) isothermal expansion
 (c) adiabatic compression
 (d) isothermal compression
27. For two mole of an ideal gas :
 (a) C_v C_p R (b) C_p C_v $2R$ (c) C_p C_v R (d) C_v C_p $2R$
28. Which of the following expressions is true for an ideal gas ?
 (a) $\frac{V}{T} \Big|_p = 0$ (b) $\frac{P}{T} \Big|_v = 0$ (c) $\frac{U}{V} \Big|_T = 0$ (d) $\frac{U}{T} \Big|_v = 0$
29. Liquefied oxygen at 1 atmosphere is heated from 50 K to 300 K by supplying heat at a constant rate. The graph that correctly shows the relationship between temperature and time is :
 (a)  (b)  (c)  (d) 
30. If w_1, w_2, w_3 and w_4 for an ideal gas are magnitude of work done in isothermal, adiabatic, isobaric and isochoric reversible expansion processes, the correct order will be :
 (a) w_1 w_2 w_3 w_4 (b) w_3 w_2 w_1 w_4
 (c) w_3 w_2 w_4 w_1 (d) w_3 w_1 w_2 w_4
31. For a closed container containing 100 mole of an ideal gas fitted with movable, frictionless, weightless piston operating such that pressure of gas remains constant at 8.21 atm. Which graph represents correct variation of $\log V$ vs. $\log T$ where V is in litre and T in kelvin?
 (a)  (b)  (c)  (d) 
32. A gas expands against a variable pressure given by $P = \frac{20}{V}$ (where P in atm and V in L). During expansion from volume of 1 litre to 10 litre, the gas undergoes a change in internal energy of 400 J. How much heat is absorbed by the gas during expansion?
 (a) 46 J (b) 4660 J (c) 5065.8 J (d) 4260 J
33. 2 mole of an ideal gas at 27°C expands isothermally and reversibly from a volume of 4 litre to 40 litre. The work done (in kJ) by the gas is :
 (a) $w = 28.72$ kJ (b) $w = 11.488$ kJ
 (c) $w = 5.736$ kJ (d) $w = 4.988$ kJ

34. 10 mole of ideal gas expand isothermally and reversibly from a pressure of 10 atm to 1 atm at 300 K. What is the largest mass which can be lifted through a height of 100 meter?
 (a) 31842 kg (b) 58.55 kg (c) 342.58 kg (d) None of these
35. A heat engine carries one mole of an ideal mono-atomic gas around the cycle as shown in the figure. Select the correct option:
 (a) q_{AB} 450 R and q_{CA} 450 R
 (b) q_{AB} 450 R and q_{CA} 225 R
 (c) q_{AB} 450 R and q_{CA} 375 R
 (d) q_{AB} 375 R and q_{CA} 450 R
36. What is the final temperature of 0.10 mole monoatomic ideal gas that performs 75 cal of work adiabatically if the initial temperature is 227°C? (use $R = 2 \text{ cal/K-mol}$)
 (a) 250 K (b) 300 K (c) 350 K (d) 750 K
37. The work done by the gas in reversible adiabatic expansion process is :
 (a) $\frac{P_2 V_2 - P_1 V_1}{1}$ (b) $\frac{nR(T_1 - T_2)}{1}$ (c) $\frac{P_2 V_2 - P_1 V_1}{1}$ (d) None of these
38. During an adiabatic process, the pressure of gas is found to be proportional to the cube of its absolute temperature. The ratio of $(C_{p,m}/C_{v,m})$ for gas is :
 (a) $\frac{3}{2}$ (b) $\frac{5}{3}$ (c) $\frac{7}{2}$ (d) $\frac{4}{3}$
39. A gas expands adiabatically at constant pressure such that $T \propto V^{1/2}$. The value of $(C_{p,m}/C_{v,m})$ of the gas will be :
 (a) 1.30 (b) 1.50 (c) 1.70 (d) 2
40. For a reversible adiabatic ideal gas expansion $\frac{dP}{P}$ is equal to :
 (a) $\frac{dV}{V}$ (b) $\frac{dV}{V}$ (c) $-\frac{1}{\gamma} \frac{dV}{V}$ (d) $\frac{dV}{V}$
41. P-V plot for two gases (assuming ideal) during adiabatic processes are given in the Fig. Plot A and plot B should correspond respectively to:
 (a) He and H_2 (b) H_2 and He
 (c) He and Ne (d) H_2 and Cl_2
42. Calculate the final temperature of a monoatomic ideal gas that is compressed reversibly and adiabatically from 16 L to 2 L at 300 K :
 (a) 600 K (b) 1044.6 K (c) 1200 K (d) 2400 K



43. 5 mole of an ideal gas expand isothermally and irreversibly from a pressure of 10 atm to 1 atm against a constant external pressure of 1 atm. w_{irr} at 300 K is :
 (a) 15.921 kJ (b) 11.224 kJ (c) 110.83 kJ (d) None of these
44. With what minimum pressure (in kPa), a given volume of an ideal gas ($C_{p,m} = 7/2 R$), originally at 400 K and 100 kPa pressure can be compressed irreversibly adiabatically in order to raise its temperature to 600 K :
 (a) 362.5 kPa (b) 275 kPa (c) 437.5 kPa (d) 550 kPa
45. The work done in adiabatic compression of 2 mole of an ideal monoatomic gas against constant external pressure of 2 atm starting from initial pressure of 1 atm and initial temperature of 300 K ($R = 2 \text{ cal/mol-degree}$)
 (a) 360 cal (b) 720 cal (c) 800 cal (d) 1000 cal
46. One mole of an ideal gas $C_{v,m} = \frac{5}{2} R$ at 300 K and 5 atm is expanded adiabatically to a final pressure of 2 atm against a constant pressure of 2 atm. Final temperature of the gas is :
 (a) 270 K (b) 273 K (c) 248.5 K (d) 200 K
47. 10 litre of a non linear polyatomic ideal gas at 127°C and 2 atm pressure is suddenly released to 1 atm pressure and the gas expanded adiabatically against constant external pressure. The final temperature and volume of the gas respectively are.
 (a) $T = 350 \text{ K}; V = 17.5 \text{ L}$ (b) $T = 300 \text{ K}; V = 15 \text{ L}$
 (c) $T = 250 \text{ K}; V = 12.5 \text{ L}$ (d) None of these
48. Calculate average molar heat capacity at constant volume of gaseous mixture contained 2 mole of each of two ideal gases A $C_{v,m} = \frac{3}{2} R$ and B $C_{v,m} = \frac{5}{2} R$:
 (a) R (b) $2R$ (c) $3R$ (d) $8R$
49. 0.5 mole each of two ideal gases A $C_{v,m} = \frac{5}{2} R$ and B ($C_{v,m} = 3R$) are taken in a container and expanded reversibly and adiabatically, during this process temperature of gaseous mixture decreased from 350 K to 250 K. Find H (in cal/mol) for the process :
 (a) 100 R (b) 137.5 R
 (c) 375 R (d) None of these

50. A cyclic process ABCD is shown in P - V diagram for an ideal gas. Which of the following diagram represents the same process?



51. 36 mL of pure water takes 100 sec to evaporate from a vessel and heater connected to an electric source which delivers 806 watt. The $H_{\text{vaporization}}$ of H_2O is :
 (a) 40.3 kJ/mol (b) 43.2 kJ/mol (c) 4.03 kJ/mol (d) None of these
52. For the reaction : $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$:
 (a) $H > U$ (b) $H = U$ (c) $H < U$ (d) None of the above
53. Consider the reaction at 300 K
 $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow 2\text{HCl}(\text{g}); H = 185 \text{ kJ}$
 If 2 mole of H_2 completely react with 2 mole of Cl_2 to form HCl , what is U for this reaction?
 (a) 0 (b) 185 kJ (c) 370 kJ (d) None of these
54. Which of the indicated relationship is correct for the following exothermic reaction carried out at constant pressure?
 $\text{CO}(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow \text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g})$
 (a) $U > H$ (b) $U = H$ (c) $w < 0$ (d) $q < 0$
55. One mole of an ideal gas undergoes a change of state (2.0 atm, 3.0 L) to (2.0 atm, 7.0 L) with a change in internal energy (U) 30 L-atm. The change in enthalpy (H) of the process in L-atm :
 (a) 22 (b) 38 (c) 25 (d) None of these
56. What is the change in internal energy when a gas contracts from 377 mL to 177 mL under a constant pressure of 1520 torr, while at the same time being cooled by removing 124 J heat?
 (a) 40.52 J (b) 83.48 J (c) 248 J (d) None of these
57. For the real gases reaction $2\text{CO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}); H = 560 \text{ kJ}$. In 10 litre rigid vessel at 500 K the initial pressure is 70 bar and after the reaction it becomes 40 bar. The change in internal energy is :
 (a) 557 kJ (b) 530 kJ (c) 563 kJ (d) None of these
58. One mole of a non-ideal gas undergoes a change of state from (1.0 atm, 3.0 L, 200 K) to (4.0 atm, 5.0 L, 250 K) with a change in internal energy (U) 40 L-atm. The change in enthalpy of the process in L-atm :
 (a) 43 (b) 57 (c) 42 (d) None of these
59. Consider the reaction at 300 K
 $\text{C}_6\text{H}_6(\text{l}) + \frac{15}{2}\text{O}_2(\text{g}) \rightarrow 6\text{CO}_2(\text{g}) + 3\text{H}_2\text{O}(\text{l}); H = 3271 \text{ kJ}$
 What is U for the combustion of 1.5 mole of benzene at 27°C ?
 (a) 3267.25 kJ (b) 4900.88 kJ (c) 4906.5 kJ (d) 3274.75 kJ
60. For the reaction; $\text{FeCO}_3(\text{s}) \rightarrow \text{FeO}(\text{s}) + \text{CO}_2(\text{g}); H = 82.8 \text{ kJ}$ at 25°C , what is (E or U) at 25°C ?
 (a) 82.8 kJ (b) 80.32 kJ (c) 2394.77 kJ (d) 85.28 kJ
61. At 5×10^5 bar pressure density of diamond and graphite are 3 g/cc and 2 g/cc respectively, at certain temperature ' T '. Find the value of $U - H$ for the conversion of 1 mole of graphite to 1 mole of diamond at temperature ' T ' :
 (a) 100 kJ/mol (b) 50 kJ/mol (c) 100 kJ/mol (d) None of these
62. Predict which of the following reaction(s) has a positive entropy change?
 I. $\text{Ag}^+(\text{aq}) + \text{Cl}^-(\text{aq}) \rightarrow \text{AgCl}(\text{s})$
 II. $\text{NH}_4\text{Cl}(\text{s}) \rightarrow \text{NH}_3(\text{g}) + \text{HCl}(\text{g})$

- III. $2\text{NH}_3(\text{g}) \quad \text{N}_2(\text{g}) \quad 3\text{H}_2(\text{g})$
 (a) I and II (b) III (c) II and III (d) II
63. Predict which of the following reaction(s) has a negative entropy change?
 I. $\text{CH}_4(\text{g}) \quad 2\text{O}_2(\text{g}) \quad \text{CO}_2(\text{g}) \quad 2\text{H}_2\text{O}(\text{l})$
 II. $\text{NH}_3(\text{g}) \quad \text{HCl}(\text{g}) \quad \text{NH}_4\text{Cl}(\text{s})$
 III. $2\text{KClO}_4(\text{s}) \quad 2\text{KClO}_3(\text{s}) \quad \text{O}_2(\text{g})$
 (a) III (b) II (c) I and II (d) I
64. Which of the following reactions is associated with the most negative change in entropy?
 (a) $2\text{SO}_2(\text{g}) \quad \text{O}_2(\text{g}) \quad 2\text{SO}_3(\text{g})$
 (b) $\text{C}_2\text{H}_4(\text{g}) \quad \text{H}_2(\text{g}) \quad \text{C}_2\text{H}_6(\text{g})$
 (c) $\text{C}(\text{s, graphite}) \quad \text{O}_2(\text{g}) \quad \text{CO}_2(\text{g})$
 (d) $3\text{C}_2\text{H}_2(\text{g}) \quad \text{C}_6\text{H}_6(\text{l})$
65. When two mole of an ideal gas $C_{p,m} = \frac{5}{2}R$ heated from 300 K to 600 K at constant pressure, the change in entropy of gas (ΔS) is :
 (a) $\frac{3}{2}R \ln 2$ (b) $\frac{3}{2}R \ln 2$ (c) $5R \ln 2$ (d) $\frac{5}{2}R \ln 2$
66. Which of the following expression for entropy change of an irreversible process ?
 (a) $dS = \frac{dq}{T}$ (b) $dS = \frac{dq}{T}$ (c) $dS = \frac{dq}{T}$ (d) $dS = \frac{dU}{T}$
67. Which of the following expressions is known as Clausius inequality ?
 (a) $\oint \frac{dq}{T} \leq 0$ (b) $\oint \frac{ds}{T} \leq 0$ (c) $\oint \frac{T}{dq} \leq 0$ (d) $\oint \frac{dq}{T} \leq 0$
68. In problem 65, calculate ΔS_{gas} if process is carried out at constant volume :
 (a) $5R \ln 2$ (b) $\frac{3}{2}R \ln 2$ (c) $3R \ln 2$ (d) $3R \ln 2$
69. If one mole of an ideal gas $C_{p,m} = \frac{5}{2}R$ is expanded isothermally at 300 K until it's volume is tripled, then change in entropy of gas is :
 (a) zero (b) infinity (c) $\frac{5}{2}R \ln 3$ (d) $R \ln 3$
70. In previous problem if expansion is carried out freely ($P_{\text{ext}} = 0$), then ΔS is :
 (a) zero (b) infinity (c) $R \ln 3$ (d) None
71. When one mole of an ideal gas is compressed to half of its initial volume and simultaneously heated to twice its initial temperature, the change in entropy of gas (ΔS) is :
 (a) $C_{p,m} \ln 2$ (b) $C_{v,m} \ln 2$ (c) $R \ln 2$ (d) $(C_{v,m} + R) \ln 2$
72. What is the change in entropy when 2.5 mole of water is heated from 27°C to 87°C? Assume that the heat capacity is constant. ($C_{p,m}(\text{H}_2\text{O}) = 4.2 \text{ J/g}\cdot\text{K}$ and $\ln(1.2) = 0.18$)
 (a) 16.6 J/K (b) 9 J/K (c) 34.02 J/K (d) 1.89 J/K
73. Calculate standard entropy change in the reaction
 $\text{Fe}_2\text{O}_3(\text{s}) \quad 3\text{H}_2(\text{g}) \quad 2\text{Fe}(\text{s}) \quad 3\text{H}_2\text{O}(\text{l})$
 Given : $S_m(\text{Fe}_2\text{O}_3, \text{s}) = 87.4$, $S_m(\text{Fe}, \text{s}) = 27.3$,
 $S_m(\text{H}_2, \text{g}) = 130.7$, $S_m(\text{H}_2\text{O}, \text{l}) = 69.9 \text{ JK}^{-1} \text{ mol}^{-1}$

- (a) $212.5 \text{ JK}^{-1} \text{ mol}^{-1}$ (b) $215.2 \text{ JK}^{-1} \text{ mol}^{-1}$
 (c) $120.9 \text{ JK}^{-1} \text{ mol}^{-1}$ (d) None of these
74. Calculate the entropy change (J/mol K) of the given reaction. The molar entropies [J/K-mol] are given in brackets after each substance.
 $2\text{PbS(s)} [91.2] + 3\text{O}_2(\text{g}) [205.1] \rightarrow 2\text{PbO(s)} [66.5] + 2\text{SO}_2(\text{g}) [248.2]$
 (a) 113.5 (b) 168.3 (c) +72.5 (d) 149.2
75. Given $rS = 266$ and the listed [S_m values] calculate S for $\text{Fe}_3\text{O}_4(\text{s})$:
 $4\text{Fe}_3\text{O}_4(\text{s}) [] + \text{O}_2(\text{g}) [205] \rightarrow 6\text{Fe}_2\text{O}_3(\text{s}) [87]$
 (a) +111.1 (b) +122.4 (c) 145.75 (d) 248.25
76. The entropy change for a phase transformation is :
 (a) $\frac{U}{dT}$ (b) $\frac{T}{H}$ (c) $\frac{H}{T}$ (d) $\frac{H}{T} - \frac{G}{T}$
77. What is the melting point of benzene if $H_{\text{fusion}} = 9.95 \text{ kJ/mol}$ and $S_{\text{fusion}} = 35.7 \text{ J/K-mol}$?
 (a) 278.7°C (b) 278.7 K (c) 300 K (d) 298 K
78. S for freezing of 10 g of $\text{H}_2\text{O(l)}$ (enthalpy of fusion is 80 cal/g) at 0°C and 1 atm is :
 (a) 12.25 J/K (b) 0.244 J/K (c) 2.93 J/K (d) 12.25 J/K
79. Chloroform has $H_{\text{vaporization}} = 29.2 \text{ kJ/mol}$ and boils at 61.2°C . What is the value of $S_{\text{vaporization}}$ for chloroform?
 (a) 87.3 J/mol-K (b) 477.1 J/mol-K (c) 87.3 J/mol-K (d) 477.1 J/mol-K
80. The entropy of vaporization of benzene is $85 \text{ JK}^{-1} \text{ mol}^{-1}$. When 117 g benzene vaporizes at its normal boiling point, the entropy change of surrounding is :
 (a) 85 JK^{-1} (b) $85 \times 1.5 \text{ JK}^{-1}$ (c) $85 \times 1.5 \text{ JK}^{-1}$ (d) None of these
81. Identify the correct statement regarding entropy
 (a) At absolute zero temperature, the entropy of perfectly crystalline substances is ve
 (b) At absolute zero temperature entropy of perfectly crystalline substance is taken to be zero
 (c) At 0°C the entropy of a perfectly crystalline substance is taken to be zero
 (d) At absolute zero temperature, the entropy of all crystalline substances is taken to be zero
82. Calculate S for following process :

$$\begin{array}{cc} X(\text{s}) & X(\text{l}) \\ \text{at } 100 \text{ K} & \text{at } 200 \text{ K} \end{array}$$


 Given : Melting point of $X(\text{s}) = 100 \text{ K}$; $H_{\text{Fusion}} = 20 \text{ kJ/mol}$; $C_{p,m}(X, \text{l}) = 10 \text{ J/mol.K}$
 (a) 26.93 J/K (b) 206.93 J/K (c) 203 J/K (d) 206.93 kJ/K
83. For a perfectly crystalline solid $C_{p,m} = aT^3$, where a is constant. If $C_{p,m}$ is 0.42 J/K mol at 10 K , molar entropy at 20 K is :
 (a) 0.42 J/K mol (b) 0.14 J/K mol (c) 1.12 J/K mol (d) zero
84. Consider the following spontaneous reaction $3X_2(\text{g}) \rightarrow 2X_3(\text{g})$. What are the sign of H , S and G for the reaction?
 (a) +ve, +ve, +ve (b) ve, ve, ve (c) ve, ve, ve (d) ve, ve, ve
85. For the reaction $2\text{H}(\text{g}) \rightarrow \text{H}_2(\text{g})$, the sign of H and S respectively are :
 (a) , (b) +, + (c) , (d) ,

86. Consider the following reaction.



signs of H , S and G for the above reaction will be

- (a) , , (b) , , (c) , , (d) , ,
87. Consider the following reaction at temperature T :



Reaction is supported by :

- (a) entropy (b) enthalpy (c) both (a) & (b) (d) neither
88. For a process to be spontaneous at constant T and P :
- (a) $(\Delta G)_{\text{system}}$ must be negative (b) $(\Delta G)_{\text{system}}$ must be positive
(c) $(\Delta S)_{\text{system}}$ must be positive (d) $(\Delta S)_{\text{system}}$ must be negative
89. For a reaction to occur spontaneously :
- (a) ΔS must be negative (b) $(\Delta H - T \Delta S)$ must be positive
(c) $\Delta H - T \Delta S$ must be negative (d) ΔH must be negative
90. Which of the following conditions regarding a chemical process ensures its spontaneity at all temperature?
- (a) $\Delta H < 0$, $\Delta S < 0$ (b) $\Delta H < 0$, $\Delta S > 0$ (c) $\Delta H > 0$, $\Delta S < 0$ (d) $\Delta H > 0$, $\Delta S > 0$
91. The free energy change $\Delta G < 0$, when
- (a) the system is at equilibrium
(b) catalyst is added
(c) reactants are initially mixed thoroughly
(d) the reactants are completely consumed
92. Which of the following conditions will always lead to a non-spontaneous change?
- (a) ΔH and ΔS both +ve (b) ΔH is -ve and ΔS is +ve
(c) ΔH and ΔS both -ve (d) ΔH is +ve and ΔS is -ve
93. Suppose that a reaction has $\Delta H^\circ = 40$ kJ and $\Delta S^\circ = 50$ J/K. At what temperature range will it change from spontaneous to non-spontaneous?
- (a) 0.8 K to 1 K (b) 799 K to 800 K (c) 800 K to 801 K (d) 799 K to 801 K
94. For isothermal expansion in case of an ideal gas :
- (a) $\Delta G = \Delta S$ (b) $\Delta G = \Delta H$ (c) $\Delta G = T \Delta S$ (d) None of these
95. What is the normal boiling point of mercury?
- Given : $\Delta H_f^\circ(\text{Hg}, l) = 0$; $\Delta S^\circ(\text{Hg}, l) = 77.4$ J/K-mol
- $\Delta H_f^\circ(\text{Hg}, g) = 60.8$ kJ/mol; $\Delta S^\circ(\text{Hg}, g) = 174.4$ J/K-mol
- (a) 624.8 K (b) 626.8 K (c) 636.8 K (d) None of these
96. 18 gm of ice is converted into water at 0°C and 1 atm. The entropies of $\text{H}_2\text{O}(s)$ and $\text{H}_2\text{O}(l)$ are 38.2 and 60 J/mol K respectively. The enthalpy change for this conversion is :
- (a) 5951.4 J/mol (b) 595.14 J/mol (c) 5951.4 J/mol (d) None of these

- 97.** Using the listed [G_f values] calculate G for the reaction :
 $3\text{H}_2\text{S}(\text{g})$ [33.6] $2\text{HNO}_3(\text{l})$ [80.6] $2\text{NO}(\text{g})$ [86.6] $4\text{H}_2\text{O}(\text{l})$ [237.1] $3\text{S}(\text{s})$ [0.0]
 (a) 513.2 (b) 1037.0 (c) +433.4 (d) +225.0
- 98.** From the following H and S values, predict which of reactions I, II, and III would be spontaneous at 25°C.
- | | H (kJ) | S (J/K) |
|------|----------|-----------|
| I. | +10.5 | +30 |
| II. | +1.8 | 113 |
| III. | 126 | +84 |
- (a) III (b) I (c) II and III (d) I and II
- 99.** Calculate H_f for UBr_4 from the G of reaction and the S° values at 298 K.
 $\text{U}(\text{s})$ $2\text{Br}_2(\text{l})$ $\text{UBr}_4(\text{s})$; G 788.6 kJ; S (J/K-mol) 50.3, 152.3, 242.6
 (a) 822.1 kJ/mol (b) 841.2 kJ/mol (c) 775.6 kJ/mol (d) 804.3 kJ/mol
- 100.** The entropies of $\text{H}_2(\text{g})$ and $\text{H}(\text{g})$ are 130.6 and 114.6 J mol⁻¹ K⁻¹ respectively at 298 K. Using the data given below calculate the bond energy of H_2 (in kJ/mol) :
 $\text{H}_2(\text{g})$ $2\text{H}(\text{g})$; G 406.6 kJ
 (a) 377.2 (b) 436.0 (c) 425.5 (d) 430.5
- 101.** Consider the G_f and H_f (kJ/mol) for the following oxides. Which oxide can be most easily decomposed to form the metal and oxygen gas?
 (a) ZnO (G 318.4, H 348.3) (b) Cu_2O (G 146.0, H 168.8)
 (c) HgO (G 58.5, H 90.8) (d) PbO (G 187.9, H 217.3)
- 102.** Which of the following option is correct ?
 (a) $\frac{\ln K_p}{T}$ $\frac{H}{RT^2}$ (b) $\frac{\ln k}{T}$ $\frac{E_a}{RT^2}$
 (c) $\frac{\ln K_p}{T}$ $\frac{U}{RT^2}$ (d) All of these
- 103.** Calculate G (kJ/mol) at 127°C for a reaction with $K_{\text{equilibrium}} = 10^5$:
 (a) 38.294 (b) 16.628 (c) 9.16 (d) None of these
- 104.** When reaction is carried out at standard states then at the equilibrium :
 (a) $H = 0$ (b) $S = 0$
 (c) equilibrium constant $K = 0$ (d) equilibrium constant $K = 1$
- 105.** At 25°C, G for the process $\text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}_2\text{O}(\text{g})$ is 8.6 kJ. The vapour pressure of water at this temperature, is nearly :
 (a) 24 torr (b) 285 torr (c) 32.17 torr (d) 100 torr
- 106.** For the auto-ionization of water at 25°C, $\text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{OH}^-(\text{aq})$ equilibrium constant is 10^{-14} .
 What is G for the process?
 (a) $\sim 8 \times 10^4$ J (b) $\sim 3.5 \times 10^4$ J (c) $\sim 10^4$ J (d) None of these

- 107.** The molar entropies of HI(g) , H(g) and I(g) at 298 K are 206.5, 114.6, and 180.7 $\text{J mol}^{-1} \text{K}^{-1}$ respectively. Using the ΔG° given below, calculate the bond energy of HI.
- $\text{HI(g)} \rightleftharpoons \text{H(g)} + \text{I(g)}; \Delta G^\circ = 271.8 \text{ kJ}$
- (a) 282.4 kJ mol^{-1} (b) 298.3 kJ mol^{-1} (c) 290.1 kJ mol^{-1} (d) 315.4 kJ mol^{-1}
- 108.** Hess's law states that :
- the standard enthalpy of an overall reaction is the sum of the enthalpy changes in individual reactions.
 - enthalpy of formation of a compound is same as the enthalpy of decomposition of the compound into constituent elements, but with opposite sign.
 - at constant temperature the pressure of a gas is inversely proportional to its volume
 - the mass of a gas dissolved per litre of a solvent is proportional to the pressure of the gas in equilibrium with the solution
- 109.** An imaginary reaction $X \rightarrow Y$ takes place in three steps
- $X \rightarrow A, \Delta H = q_1; A \rightarrow B, \Delta H = q_2; B \rightarrow Y, \Delta H = q_3$
- If Hess' law is applicable, then the heat of the reaction ($X \rightarrow Y$) is :
- (a) $q_1 + q_2 + q_3$ (b) $q_2 + q_3 - q_1$ (c) $q_1 + q_2 - q_3$ (d) $q_3 - q_2 - q_1$
- 110.** The enthalpy change for a reaction does not depend upon :
- the physical states of reactants and products
 - use of different reactants for the same product
 - the number of intermediate reaction steps
 - the differences in initial or final temperatures of involved substances
- 111.** The standard enthalpy of formation of gaseous H_2O at 298 K is -241.82 kJ/mol . Calculate ΔH° at 373 K given the following values of the molar heat capacities at constant pressure :
- $\text{H}_2\text{O(g)} = 33.58 \text{ JK}^{-1} \text{mol}^{-1}; \text{H}_2\text{(g)} = 29.84 \text{ JK}^{-1} \text{mol}^{-1}; \text{O}_2\text{(g)} = 29.37 \text{ JK}^{-1} \text{mol}^{-1}$
- Assume that the heat capacities are independent of temperature :
- (a) 242.6 kJ/mol (b) 485.2 kJ/mol (c) 121.3 kJ/mol (d) 286.4 kJ/mol
- 112.** Which of the following value of ΔH_f° represent that the product is least stable ?
- (a) 94.0 kcal mol^{-1} (b) 231.6 kcal mol^{-1} (c) 21.4 kcal mol^{-1} (d) 64.8 kcal mol^{-1}
- 113.** For which of the following substances the enthalpy of formation in the standard state is zero ?
- (a) Sucrose (b) Ethanol (c) Aluminium (d) Calcium chloride
- 114.** Calculate the standard enthalpy of reaction for the following reaction using the listed enthalpies of reaction :
- $3\text{Co(s)} + 2\text{O}_2\text{(g)} \rightarrow \text{Co}_3\text{O}_4\text{(s)}$
- $2\text{Co(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{CoO(s)}; \Delta H_1 = 475.8 \text{ kJ}$
- $6\text{CoO(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{Co}_3\text{O}_4\text{(s)}; \Delta H_2 = 355.0 \text{ kJ}$
- (a) 891.2 kJ (b) 120.8 kJ (c) +891.2 kJ (d) 830.8 kJ
- 115.** From given following equations and ΔH° values, determine the enthalpy of reaction at 298 K for the reaction :
- $\text{C}_2\text{H}_4\text{(g)} + 6\text{F}_2\text{(g)} \rightarrow 2\text{CF}_4\text{(g)} + 4\text{HF(g)}$
- $\text{H}_2\text{(g)} + \text{F}_2\text{(g)} \rightarrow 2\text{HF(g)}; \Delta H_1 = 537 \text{ kJ}$



- (a) 1165 (b) 2486 (c) +1165 (d) +2486



116. Given :




The enthalpy of formation of diamond from graphite is

- (a) 2.0 kJ (b) 1.5 kJ (c) 788 kJ (d) 788 kJ

117. Which of the following equations represents a reaction that provides the enthalpy of formation of CH_3Cl ?



118. Use the given standard enthalpies of formation (in kJ/mol) to determine the enthalpy of reaction of the following reaction :



- (a) 335.8 kJ/mol (b) 873.8 kJ/mol (c) 697.2 kJ/mol (d) 890.4 kJ/mol

119. The standard enthalpy of formation of octane (C_8H_{18}) is 250 kJ/mol. Calculate the enthalpy of combustion of C_8H_{18} . The enthalpy of formation of $\text{CO}_2(\text{g})$ and $\text{H}_2\text{O(l)}$ are 394 kJ/mol and 286 kJ/mol respectively :

- (a) -5200 kJ/mol (b) -5726 kJ/mol (c) -5476 kJ/mol (d) -5310 kJ/mol

120. Determine the enthalpy of formation of $\text{B}_2\text{H}_6(\text{g})$ in kJ/mol of the following reaction :



- (a) 75.6 (b) +75.6 (c) 57.4 (d) 28.4

121. Consider the following reactions :




The heat of formation of CO(g) is :

- (a) $(x - y) \text{ kJ/mol}$ (b) $(x + y) \text{ kJ/mol}$ (c) $(y - x) \text{ kJ/mol}$ (d) None of these

122. If $\Delta_f H^\circ(\text{C}_2\text{H}_4)$ and $\Delta_f H^\circ(\text{C}_2\text{H}_6)$ are x_1 and $x_2 \text{ kcal mol}^{-1}$, then heat of hydrogenation of C_2H_4 is :

- (a) $x_1 - x_2$ (b) $x_1 + x_2$ (c) $x_2 - x_1$ (d) $x_1 - 2x_2$


123. What amount of energy (kJ) is released in the combustion of 5.8 g of $\text{C}_4\text{H}_{10}(\text{g})$?



- (a) 575.6 (b) 287.8 (c) 182 (d) 57.56

- 124.** The enthalpy of the reaction forming PbO according to the following equation is 438 kJ. What heat energy (kJ) is released in formation of 22.3 g PbO(s)?

(Atomic masses : Pb 207, O 16.0)



- (a) 21.9 (b) 28.7 (c) 14.6 (d) 34.2

- 125.** The fat, $\text{C}_{57}\text{H}_{104}\text{O}_6(\text{s})$, is metabolized via the following reaction. Given the enthalpies of formation, calculate the $\Delta_r H$ (kJ) when 1.0 g of this fat reacts.




$\Delta_f H$ ($\text{C}_{57}\text{H}_{104}\text{O}_6, \text{s}$) 70870 kJ/mol; $\Delta_f H$ ($\text{H}_2\text{O}, \text{l}$) 285.8 kJ/mol;

$\Delta_f H$ (CO_2, g) 393.5 kJ/mol


- (a) $\Delta_r H$ 37.98 (b) $\Delta_r H$ 40.4 (c) $\Delta_r H$ 33.4 (d) $\Delta_r H$ 30.2

- 126.** The $\Delta_f H$ of $\text{NH}_3(\text{g})$ is 46 kJ mol⁻¹. The $\Delta_r H$ (in kJ mol⁻¹) of the reaction,



- (a) 46 (b) 46 (c) 92 (d) 92

- 127.** Consider the following reaction:



What is the value of $\Delta_r H$ for second reaction if water vapour instead of liquid water is formed as product?

- (a) 1409 kJ (b) 1629 kJ (c) 1277 kJ (d) None of these


- 128.** The standard enthalpy change for the following reaction is 436.4 kJ :



What is the $\Delta_f H$ of atomic hydrogen (H)?

- (a) 872.8 kJ/mol (b) 218.2 kJ/mol (c) 218.2 kJ/mol (d) 436.9 kJ/mol

- 129.** Determine enthalpy of formation for $\text{H}_2\text{O}_2(\text{l})$, using listed enthalpies of reaction :



- (a) 383 kJ/mol (b) 187 kJ/mol (c) 498 kJ/mol (d) None of these

- 130.** Heat of combustion of ethanol at constant pressure and at temperature T K is found to be q J mol⁻¹. Hence, heat of combustion (in J mol⁻¹) of ethanol at the same temperature at constant volume will be :

- (a) $RT - q$ (b) $(q - RT)$ (c) $q - RT$ (d) $q + RT$

- 131.** Stearic acid [$\text{CH}_3(\text{CH}_2)_{16}\text{CO}_2\text{H}$] is a fatty acid, the part of fat that stores most of the energy. 1.0 g of stearic acid was burned in a bomb calorimeter. The bomb had a heat capacity of 652 J/°C. If the temperature of 500 g water ($c = 4.18 \text{ J/g } ^\circ\text{C}$) increase from 25.0 to 39.3°C, how much heat was released when the stearic acid was burned?

[Given C_p (H_2O) 4.18 J/g°C]

- (a) 39.21 kJ (b) 29.91 kJ (c) 108 kJ (d) 9.32 kJ

- 132.** Gasoline has an enthalpy of combustion 24000 kJ/ gallon. When gasoline burns in an automobile engine, approximately 30% of the energy released is used to produce mechanical work. The remainder is lost as heat transfer to the engine's cooling system. As a start on estimating how much heat transfer is required, calculate what mass of water could be heated from 25°C to 75°C by the combustion of 1.0 gallon of gasoline in an automobile?
(Given : $C(H_2O) = 4.18 \text{ J/g}^\circ\text{C}$)
(a) 34.45 kg (b) 80.383 kg (c) 22 kg (d) 224 kg
- 133.** A 0.05 L sample of 0.2 M aqueous hydrochloric acid is added to 0.05 L of 0.2 M aqueous ammonia in a calorimeter. Heat capacity of entire calorimeter system is 480 J/K. The temperature increase is 1.09 K. Calculate $\Delta_r H$ in kJ/mol for the following reaction :
$$HCl(aq.) + NH_3(aq.) \rightarrow NH_4Cl(aq.)$$

(a) 52.32 (b) 61.1 (c) 55.8 (d) 58.2
- 134.** A coffee cup calorimeter initially contains 125 g of water, at a temperature of 24.2°C. 8 g of ammonium nitrate (NH_4NO_3), also at 24.2°C, is added to the water, and the final temperature is 18.2°C. What is the heat of solution of ammonium nitrate in kJ/mol? The specific heat capacity of the solution is 4.2 J/°C g.
(a) 33.51 kJ/mol (b) 39.5 kJ/mol (c) 32.2 kJ/mol (d) 37.3 kJ/mol
- 135.** Read following statement(s) carefully and select the right option :
(I) The enthalpy of solution of $CaCl_2 \cdot 6H_2O$ in a large volume of water is endothermic to the extent of 3.5 kcal/ mol. If $\Delta_r H = 23.2 \text{ kcal}$ for the reaction,
$$CaCl_2(s) + 6H_2O(l) \rightarrow CaCl_2 \cdot 6H_2O(s)$$

then heat of solution of $CaCl_2$ (anhydrous) in a large volume of water is – 19.7 kcal/mol
(II) For the reaction $2Cl(g) \rightarrow Cl_2(g)$; the sign of $\Delta_r H$ and $\Delta_r S$ are negative.
(a) statement I and II both are wrong (b) both are correct
(c) only I is correct (d) only II is correct
- 136.** If the enthalpy of formation and enthalpy of solution of $HCl(g)$ are 92.3 kJ/mol and 75.14 kJ/mol respectively then find enthalpy of formation of $Cl^- (aq)$:
(a) 17.16 kJ/mol (b) 167.44 kJ/mol (c) 17.16 kJ/mol (d) None of these
- 137.** At 25°C, when 1 mole of $MgSO_4$ was dissolved in water, the heat evolved was found to be 91.2 kJ. One mole of $MgSO_4 \cdot 7H_2O$ on dissolution gives a solution of the same composition accompanied by an absorption of 13.8 kJ. The enthalpy of hydration, i.e., $\Delta_r H_h$ for the reaction
$$MgSO_4(s) + 7H_2O(l) \rightarrow MgSO_4 \cdot 7H_2O(s)$$
 is :
(a) 105 kJ/mol (b) 77.4 kJ/mol (c) 105 kJ/mol (d) None of these
- 138.** The enthalpies of neutralization of a weak base AOH and a strong base BOH by HCl are 12250 cal/mol and 13000 cal/mol respectively. When one mole of HCl is added to a solution containing 1 mole of AOH and 1 mole of BOH, the enthalpy change was 12500 cal/ mol. In what ratio is the acid distribution between AOH and BOH?
(a) 2 : 1 (b) 2 : 3 (c) 1 : 2 (d) None of these
- 139.** The magnitude of enthalpy change for neutralization of the reaction ;
$$MgO(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2O(l)$$
 (Given $\Delta_r H^\circ (aq) : OH^- (aq) = -57 \text{ kJ/mol}$) will be :

- (a) less than 57 kJ mol^{-1} (b) 57 kJ mol^{-1}
 (c) greater than 57 kJ mol^{-1} (d) 57 kJ mol^{-1}

140. Enthalpy of neutralization of HCl by NaOH is 55.84 kJ/mol and by NH_4OH is 51.34 kJ/mol . The enthalpy of ionization of NH_4OH is :

- (a) 107.18 kJ/mol (b) 4.5 kJ/mol (c) 4.5 kJ/mol (d) None of these


141. Which of the following reaction is endothermic ?

- (a) $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$ (b) $\text{Fe} + \text{S} \rightarrow \text{FeS}$
 (c) $\text{NaOH} + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O}$ (d) $\text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$

142. Which of the following is not correct ?

- (a) Dissolution of a salt in excess of water may be endothermic process
 (b) Neutralisation is always exothermic
 (c) The absolute value of enthalpy (H) can be determined by calorimeter
 (d) The heat of reaction at constant volume is denoted by U


143. Substance $\text{A}_2\text{B}(\text{g})$ can undergoes decomposition to form two set of products :



If the molar ratio of $\text{A}_2(\text{g})$ to $\text{A}(\text{g})$ is $5 : 3$ in a set of product gases, then the energy involved in the decomposition of 1 mole of $\text{A}_2\text{B}(\text{g})$ is :

- (a) 48.75 kJ/mol (b) 43.73 kJ/mol (c) 46.25 kJ/mol (d) None of these

144. Boron can undergo the following reactions with the given enthalpy changes :



Assume no other reactions are occurring.

If in a container (operating at constant pressure) which is isolated from the surrounding, mixture of H_2 (gas) and O_2 (gas) are passed over excess of $\text{B}(\text{s})$, then calculate the molar ratio ($\text{O}_2 : \text{H}_2$) so that temperature of the container do not change :

- (a) $15 : 3$ (b) $42 : 1$ (c) $1 : 42$ (d) $1 : 84$

145. The enthalpy change for the following reaction is 368 kJ . Calculate the average O—F bond energy.



146. The enthalpy change for the reaction, $\text{C}_2\text{H}_6(\text{g}) + 2\text{C}(\text{g}) + 6\text{H}(\text{g})$ is $X \text{ kJ}$. The bond energy of C—H bond is :

- (a) $\frac{X}{2}$ (b) $\frac{X}{3}$ (c) $\frac{X}{6}$ (d) data insufficient

- 147.** The table given below lists the bond dissociation energy (E_{diss}) for single covalent bonds formed between C and atoms A, B, D, E.

Bond	E_{diss} (kcal mol ⁻¹)
C — A	240
C — B	382
C — D	276
C — E	486

Which of the atoms has smallest size ?

- (a) D (b) E (c) A (d) D

- 148.** Calculate P—Cl bond enthalpy

Given : $\Delta_f H$ (PCl₃, g) 306 kJ/mol; $H_{\text{atomization}}$ (P, s) 314 kJ/mol;

$\Delta_f H$ (Cl, g) 121 kJ/mol

- (a) 123.66 kJ/mol (b) 371 kJ/mol (c) 19 kJ/mol (d) None of these

- 149.** Calculate the enthalpy for the following reaction using the given bond energies (kJ/mol) :

(C—H 414; H—O 463; H—Cl 431, C—Cl 326; C—O 335)

CH₃ OH(g) HCl(g) CH₃ Cl(g) H₂O(g)

- (a) 23 kJ/mol (b) 42 kJ/mol (c) 59 kJ/mol (d) 511 kJ/mol

- 150.** Based on the values of B.E. given, $\Delta_f H$ of N₂H₄(g) is :

Given : N—N = 159 kJ mol⁻¹; H—H = 436 kJ mol⁻¹

N=N = 941 kJ mol⁻¹; N—H = 398 kJ mol⁻¹

- (a) 711 kJ mol⁻¹ (b) 62 kJ mol⁻¹ (c) 98 kJ mol⁻¹ (d) 711 kJ mol⁻¹



Level 2

- A heating coil is immersed in a 100 g sample of $\text{H}_2\text{O}(l)$ at 1 bar and 100°C in a closed vessel. In this heating process, 60% of the liquid is converted into gaseous form at constant pressure of 1 bar. Densities of liquid and gaseous water under these conditions are 1000 kg/m^3 and 0.60 kg/m^3 respectively. Magnitude of the work done for the process is :
 (a) 4997 J (b) 4970 J (c) 9994 J (d) None of these
- A rigid and insulated tank of 3 m^3 volume is divided into two compartments. One compartment of volume of 2 m^3 contains an ideal gas at 0.8314 MPa and 400 K while the second compartment of volume 1 m^3 contains the same gas at 8.314 MPa and 500 K. If the partition between the two compartments is ruptured, the final temperature of the gas is :
 (a) 420 K (b) 450 K (c) 480 K (d) None of these
- What is the value of change in internal energy at 1 atm in the process?




Given : $C_{v,m}(\text{H}_2\text{O}, l) = 75.0 \text{ JK}^{-1} \text{ mol}^{-1}$; $C_{p,m}(\text{H}_2\text{O}, g) = 33.314 \text{ JK}^{-1} \text{ mol}^{-1}$

H_{vap} at 373 K = 40.7 kJ/mol

- (a) 42.91 kJ/mol (b) 43086 kJ/mol (c) 42.6 kJ/mol (d) 49.6 kJ/mol
- For an ideal gas $\frac{C_{p,m}}{C_{v,m}}$. The molecular mass of the gas is M , its specific heat capacity at constant volume is :
 (a) $\frac{R}{M(1)}$ (b) $\frac{M}{R(1)}$ (c) $\frac{RM}{1}$ (d) $\frac{R}{M(1)}$
- 1 mole of an ideal gas A ($C_{v,m} = 3R$) and 2 mole of an ideal gas B are $C_{v,m} = \frac{3}{2}R$ taken in a container and expanded reversibly and adiabatically from 1 litre to 4 litre starting from initial temperature of 320 K. E or U for the process is :
 (a) 240 R (b) 240 R (c) 480 R (d) 960 R
- Calculate the work done by the system in an irreversible (single step) adiabatic expansion of 2 mole of a polyatomic gas ($\gamma = 4/3$) from 300 K and pressure 10 atm to 1 atm :
 (a) 227 R (b) 205 R (c) 405 R (d) None of these


- One mole of an ideal gas is carried through the reversible cyclic process as shown in figure. The max. temperature attained by the gas during the cycle :

- $\frac{7}{6}R$
- $\frac{12}{49}R$
- $\frac{49}{12}R$
- None of these



8. Two moles of a triatomic linear gas (neglect vibration degree of freedom) are taken through a reversible process ideal starting from A as shown in figure.

The volume ratio $\frac{V_B}{V_A}$ 4. If the temperature at A is 73°C , then:



- (i) Work done by the gas in AB process is :
 (a) 6.16 kJ (b) 308.3 kJ
 (c) 9.97 kJ (d) 0 J
- (ii) Total enthalpy change in both steps is :
 (a) 3000 R (b) 4200 R (c) 2100 R (d) 0
9. A gas $C_{v,m} = \frac{5}{2}R$ behaving ideally was allowed to expand reversibly and adiabatically from 1 litre to 32 litre. Its initial temperature was 327°C . The molar enthalpy change (in J/mol) for the process is :
 (a) 1125 R (b) 675 (c) 1575 R (d) None of these
10. Two mole of an ideal gas is heated at constant pressure of one atmosphere from 27°C to 127°C . If $C_{v,m} = 20 \times 10^2 \text{ J K}^{-1} \text{ mol}^{-1}$, then q and U for the process are respectively :
 (a) 6362.8 J, 4700 J (b) 3037.2 J, 4700 J (c) 7062.8 J, 5400 J (d) 3181.4 J, 2350 J
11. 10 mole of an ideal gas is heated at constant pressure of one atmosphere from 27°C to 127°C . If $C_{v,m} = 21.686 \times 10^3 \text{ J K}^{-1} \text{ mol}^{-1}$, then H for the process is :
 (a) 3000 J (b) 3350 J (c) 3700 J (d) 30350 J
12. For polytropic process $PV^n = \text{constant}$, molar heat capacity (C_m) of an ideal gas is given by :
 (a) $C_{v,m} \frac{R}{(n-1)}$ (b) $C_{v,m} \frac{R}{(1-n)}$ (c) $C_{v,m} R$ (d) $C_{p,m} \frac{R}{(n-1)}$
13. 2 mole of an ideal monoatomic gas undergoes a reversible process for which $PV^2 = C$. The gas is expanded from initial volume of 1 L to final volume of 3 L starting from initial temperature of 300 K. Find H for the process :
 (a) 600 R (b) 1000 R (c) 3000 R (d) None of these
14. Calculate S for 3 mole of a diatomic ideal gas which is heated and compressed from 298 K and 1 bar to 596 K and 4 bar: [Given: $C_{v,m}(\text{gas}) = \frac{5}{2}R$; $\ln(2) = 0.70$; $R = 2 \text{ cal K}^{-1} \text{ mol}^{-1}$]
 (a) 14.7 cal K^{-1} (b) $+14.7 \text{ cal K}^{-1}$ (c) 4.9 cal K^{-1} (d) 6.3 cal K^{-1}
15. One mole of an ideal monoatomic gas at 27°C is subjected to a reversible isentropic compression until final temperature reaches to 327°C . If the initial pressure was 1.0 atm then find the value of $(\ln P_2)$: (Given : $\ln 2 = 0.7$)
 (a) 1.75 atm (b) 0.176 atm (c) 1.0395 atm (d) 2.0 atm
16. Two mole of an ideal gas is expanded irreversibly and isothermally at 37°C until its volume is doubled and 3.41 kJ heat is absorbed from surrounding. S_{total} (system + surrounding) is :
 (a) 0.52 J/K (b) 0.52 J/K (c) 22.52 J/K (d) 0
17. For a perfectly crystalline solid $C_{p,m} = aT^3 + bT$, where a and b are constant. If $C_{p,m}$ is 0.40 J/K mol at 10 K and 0.92 J/K mol at 20 K, then molar entropy at 20 K is :
 (a) 0.92 J/K mol (b) 8.66 J/K mol (c) 0.813 J/K mol (d) None of these

18. Which of the following statement(s) is/are correct?

Statement (i) : The entropy in isolated system with P - V work only, is always maximized at equilibrium.

Statement (ii) : It is possible for the entropy of a closed system to decrease substantially in an irreversible process.

Statement (iii) : Entropy can be created but not destroyed.

Statement (iv) : S_{system} is zero for reversible process in an isolated system.

(a) Statement i, ii, iii

(b) Statement ii, iv

(c) Statement i, ii, iv

(d) All of these

19. Combustion of sucrose is used by aerobic organisms for providing energy for the life sustaining processes. If all the capturing of energy from the reaction is done through electrical process (non P - V work) then calculate maximum available energy which can be captured by combustion of 34.2 gm of sucrose

Given : $H_{\text{combustion}}$ (sucrose) 6000 kJ mol^{-1}

$S_{\text{combustion}}$ 180 J/K-mol and body temperature is 300 K


(a) 600 kJ

(b) 594.6 kJ

(c) 5.4 kJ

(d) 605.4 kJ

20. For the hypothetical reaction



${}_rG$ and ${}_rS$ are 20 kJ/mol and $20 \text{ JK}^{-1} \text{ mol}^{-1}$ respectively at 200 K .

If ${}_rC_p$ is $20 \text{ JK}^{-1} \text{ mol}^{-1}$ then ${}_rH$ at 400 K is :

(a) 20 kJ/mol

(b) 7.98 kJ/mol

(c) 28 kJ/mol

(d) None of these

21. Calculate ${}_fG$ for $(\text{NH}_4\text{Cl}, s)$ at 310 K .

Given : ${}_fH$ $(\text{NH}_4\text{Cl}, s)$ 314.5 kJ/mol ; ${}_rC_p$ 0

$S_{\text{N}_2(g)}$ $192 \text{ JK}^{-1} \text{ mol}^{-1}$; $S_{\text{H}_2(g)}$ $130.5 \text{ JK}^{-1} \text{ mol}^{-1}$;

$S_{\text{Cl}_2(g)}$ $233 \text{ JK}^{-1} \text{ mol}^{-1}$; $S_{\text{NH}_4\text{Cl}(s)}$ $99.5 \text{ JK}^{-1} \text{ mol}^{-1}$

All given data at 300 K .

(a) 198.56 kJ/mol

(b) 426.7 kJ/mol

(c) 202.3 kJ/mol

(d) None of these

22. Using listed informations, calculate ${}_rG$ (in kJ/mol) at 27°C



Given : At 300 K

H_f (kJ/mol) $891, 110.5, 0.0, 393.5$

S (J/K-mol) $102.5, 197.7, 30.0, 213.7$

(a) 214.8

(b) 195.0

(c) 200.3

(d) 256.45

23. Fixed mass of an ideal gas contained in a 24.63 L sealed rigid vessel at 1 atm is heated from 73°C to 27°C . Calculate change in Gibb's energy if entropy of gas is a function of temperature as $S = 2 \times 10^2 T$ (J/K): (Use $1 \text{ atm L} = 0.1 \text{ kJ}$)

(a) 1231.5 J

(b) 1281.5 J


(c) 781.5 J

(d) 0

24. The molar heat capacities at constant pressure (assumed constant with respect to temperature) of A, B and C are in ratio of $3 : 1.5 : 2.0$. The enthalpy change for the exothermic reaction $A + 2B \rightleftharpoons 3C$ at 300 K and 310 K is H_{300} and H_{310} respectively then :

- (a) H_{300} H_{310}
 (b) H_{300} H_{310}
 (c) H_{300} H_{310}
 (d) if $T_2 > T_1$ then $H_{310} > H_{300}$ and if $T_2 < T_1$ then $H_{310} < H_{300}$
- 25.** Determine ΔU at 300 K for the following reaction using the listed enthalpies of reaction :
- | | | | | |
|-----------------------|----------------------------|----------------------------|-------------------------|--------------------------|
| 4CO(g) | $8\text{H}_2\text{(g)}$ | $3\text{CH}_4\text{(g)}$ | $\text{CO}_2\text{(g)}$ | $2\text{H}_2\text{O(l)}$ |
| C (graphite) | $1/2 \text{O}_2\text{(g)}$ | CO(g) | H_1 | 110.5 kJ |
| | CO(g) | $1/2 \text{O}_2\text{(g)}$ | $\text{CO}_2\text{(g)}$ | H_2 282.9 kJ |
| | $\text{H}_2\text{(g)}$ | $1/2 \text{O}_2\text{(g)}$ | $\text{H}_2\text{O(l)}$ | H_3 285.8 kJ |
| C (graphite) | $2\text{H}_2\text{(g)}$ | $\text{CH}_4\text{(g)}$ | H_4 | 74.8 kJ |
- (a) 653.5 kJ (b) 686.2 kJ (c) 747.4 kJ (d) None of these
- 26.** Calculate $\Delta_f H$ (in kJ/mol) for Cr_2O_3 from the $\Delta_f G$ and the S° values provided at 27°C
- | | | | | |
|-----------------------|--|------------------------------------|-------------------------------|---------------|
| 4Cr(s) | $3\text{O}_2\text{(g)}$ | $2\text{Cr}_2\text{O}_3\text{(s)}$ | $\Delta_f G^\circ$ | 2093.4 kJ/mol |
| S° (J/K mol) : | S° (Cr, s) | 24; | S° (O_2 , g) | 205; |
| | S° (Cr_2O_3 , s) | 81 | | |
- (a) 2258.1 kJ/mol (b) 1129.05 kJ/mol (c) 964.35 kJ/mol (d) None of these
- 27.** Calculate the heat produced (in kJ) when 224 gm of CaO is completely converted to CaCO_3 by reaction with CO_2 at 27°C in a container of fixed volume.
- Given : $\Delta_f H^\circ$ (CaCO_3 , s) 1207 kJ/mol; $\Delta_f H^\circ$ (CaO , s) 635 kJ/mol
- $\Delta_f H^\circ$ (CO_2 , g) 394 kJ/mol; [Use $R = 8.3 \text{ JK}^{-1} \text{ mol}^{-1}$]
- (a) 702.04 kJ (b) 721.96 kJ (c) 712 kJ (d) 721 kJ
- 28.** When 1.0 g of oxalic acid ($\text{H}_2\text{C}_2\text{O}_4$) is burned in a bomb calorimeter whose heat capacity is 8.75 kJ/K, the temperature increases by 0.312 K. The enthalpy of combustion of oxalic acid at 27°C is :
- (a) 245.7 kJ/mol (b) 244.452 kJ/mol (c) 241.5 kJ/mol (d) None of these
- 29.** Enthalpy of neutralization of H_3PO_3 acid is 106.68 kJ/mol using NaOH . Enthalpy of neutralization of HCl by NaOH is 55.84 kJ/mol. Calculate $\Delta_{\text{ionization}} H^\circ$ of H_3PO_3 into its ions :
- (a) 50.84 kJ/mol (b) 5 kJ/mol (c) 2.5 kJ/mol (d) None of these
- 30.** The enthalpy of neutralization of a weak monoprotic acid (HA) in 1 M solution with a strong base is 55.95 kJ/mol. If the unionized acid required 1.4 kJ/mol heat for its complete ionization and enthalpy of neutralization of the strong monobasic acid with a strong monoacidic base is 57.3 kJ/mol, what is the % ionization of the weak acid in molar solution ?
- (a) 1% (b) 3.57% (c) 35.7% (d) 10%
- 31.** Determine C—C and C—H bond enthalpy (in kJ/mol)
- Given : $\Delta_f H^\circ$ (C_2H_6 , g) 85 kJ/mol, $\Delta_f H^\circ$ (C_3H_8 , g) 104 kJ/mol
- $\Delta_{\text{sub}} H^\circ$ (C, s) 718 kJ/mol, B.E. (H—H) 436 kJ/mol
- (a) 414, 345 (b) 345, 414 (c) 287, 404.5 (d) None of these
- 32.** Consider the following data : $\Delta_f H^\circ$ (N_2H_4 , l) 50 kJ/mol, $\Delta_f H^\circ$ (NH_3 , g) 46 kJ/mol
- B.E. (N—H) 393 kJ/mol and B.E. (H—H) 436 kJ/mol,
- $\Delta_{\text{vap}} H^\circ$ (N_2H_4 , l) 18 kJ/mol
- The N—N bond energy in N_2H_4 is :
- (a) 226 kJ/mol (b) 154 kJ/mol (c) 190 kJ/mol (d) None of these

33. What is the bond enthalpy of Xe—F bond?



Given : Ionization energy of Xe = 279 kcal/mol

B.E. (F—F) = 38 kcal/mol, Electron affinity of F = 85 kcal/mol


- (a) 24 kcal/mol (b) 34 kcal/mol (c) 8.5 kcal/mol (d) None of these
34. If enthalpy of hydrogenation of $\text{C}_6\text{H}_6(\text{l})$ into $\text{C}_6\text{H}_{12}(\text{l})$ is 205 kJ/mol and resonance energy of

$\text{C}_6\text{H}_6(\text{l})$ is 152 kJ/mol then enthalpy of hydrogenation of  is

Assume H_{vap} of $\text{C}_6\text{H}_6(\text{l})$, $\text{C}_6\text{H}_8(\text{l})$, $\text{C}_6\text{H}_{12}(\text{l})$ all are equal :


- (a) 535.5 kJ/mol (b) 238 kJ/mol (c) -357 kJ/mol (d) 119 kJ/mol
35. The enthalpy of combustion of propane (C_3H_8) gas in terms of given data is :

Bond energy (kJ/mol)



Resonance energy of CO_2 is z kJ/mol and $H_{\text{vaporization}} [\text{H}_2\text{O}(\text{l})]$ is y (kJ/mol)

- (a) $8x_1 - 2x_5 - 5x_2 - 6x_3 - 8x_4 + 4y - 3z$ (b) $6x_1 - x_5 - 5x_2 - 3x_3 - 4x_4 + 4y - 3z$
 (c) $8x_1 - 2x_5 - 5x_2 - 6x_3 - 8x_4 + y - z$ (d) $8x_1 - x_5 - 5x_2 - 6x_3 - 8x_4 + 4y - 3z$
36. The efficiency of an ideal gas with adiabatic exponent ' γ ' for the shown cyclic process would be



- (a) $\frac{(1 - \gamma)(2 \ln 2 - 1)}{1 - (1 - \gamma)2 \ln 2}$ (b) $\frac{(1 - \gamma)(1 - 2 \ln 2)}{(1 - \gamma)2 \ln 2 - 1}$
 (c) $\frac{(2 \ln 2 - 1)(1 - \gamma)}{(1 - \gamma)2 \ln 2 - 1}$ (d) $\frac{(2 \ln 2 - 1)}{(1 - \gamma)}$
37. Initially one mole of ideal gas $C_v = \frac{5}{2}R$ at 1.0 atm and 300 K is put through the following cycle :



Step-I : Heating to twice its initial pressure at constant volume.

Step-II : Adiabatic expansion to its initial temperature.

Step-III : Isothermal compression back to 1.00 atm.

What is the volume at state X ?

- (a) 40.4 L (b) 65.0 L (c) 139 L (d) 4.35 L

38. What is the enthalpy of neutralization of HF against a strong base ?

Given : $\text{H}^+(aq) + \text{OH}^-(aq) \rightarrow \text{H}_2\text{O}(l)$; $\Delta_r H^\circ = -56 \text{ kJ/mol}$

$\Delta_f H^\circ (\text{HF}, aq) = -329 \text{ kJ/mol}$; $\Delta_f H^\circ (\text{H}_2\text{O}, l) = -285 \text{ kJ/mol}$

$\Delta_f H^\circ (\text{F}^-, aq) = -320 \text{ kJ/mol}$

- (a) 17 kJ/mol (b) 38 kJ/mol (c) 47 kJ/mol (d) 43 kJ/mol

39. The molar heat capacities of Iodine vapour and solid are 7.8 and 14 cal/mol respectively. If enthalpy of sublimation of iodine is 6096 cal/mol at 200°C, then what is ΔU (internal energy change) at 250°C in cal/mol

- (a) 5360 (b) 4740 (c) 6406 (d) None of these

40. For the reaction taking place at certain temperature $\text{NH}_2\text{COONH}_4(s) \rightleftharpoons 2\text{NH}_3(g) + \text{CO}_2(g)$, if equilibrium pressure is $3X$ bar then $\Delta_r G^\circ$ would be

- (a) $RT \ln 9 - 3RT \ln X$ (b) $RT \ln 4 - 3RT \ln X$
(c) $3RT \ln X$ (d) None of these



Level 3



Passage 1

The first law of thermodynamics for a closed system is $dU = dq + dw$, where $dw = dw_{pv} + dw_{non-pv}$. The most common type of w_{non-pv} is electrical work. As per IUPAC convention work done on the system is positive.

1. A system generates 50 J of electrical energy, has 150 J of pressure-volume work done on it by the surroundings while releasing 300 J of heat energy. What is the change in the internal energy of the system?
(a) 500 (b) 100 (c) 300 (d) 200
2. A system generates 50 J of electrical energy and delivers 150 J of pressure-volume work against the surroundings while releasing 300 J of heat energy. What is the change in the internal energy of the system?
(a) 100 (b) 400 (c) 300 (d) 500



Passage 2

If the boundary of system moves by an infinitesimal amount, the work involved is given by $dw = P_{ext} dV$

for irreversible process $w = P_{ext} \Delta V$ (where $\Delta V = V_f - V_i$)

for reversible process $P_{ext} = P_{int}$ $dP \sim P_{int}$

so for reversible isothermal process $w = nRT \ln \frac{V_f}{V_i}$

2 mole of an ideal gas undergoes isothermal compression along three different paths :

- (i) reversible compression from $P_i = 2$ bar and $V_i = 8$ L to $P_f = 20$ bar
- (ii) a single stage compression against a constant external pressure of 20 bar, and
- (iii) a two stage compression consisting initially of compression against a constant external pressure of 10 bar until $P_{gas} = P_{ext}$, followed by compression against a constant pressure of 20 bar until $P_{gas} = P_{ext}$.

1. Work done (in bar-L) on the gas in reversible isothermal compression is :
(a) 9.212 (b) 36.848 (c) 18.424 (d) None of these
2. Work done on the gas in single stage compression is :
(a) 36 (b) 72 (c) 144 (d) None of these
3. Total work done on the gas in two stage compression is :
(a) 40 (b) 80 (c) 160 (d) None of these
4. Order of magnitude of work is :
(a) $w_1 < w_2 < w_3$ (b) $w_3 < w_2 < w_1$ (c) $w_2 < w_3 < w_1$ (d) $w_1 < w_2 < w_3$



Passage 3

Standard Gibbs energy of reaction (${}_rG$) at a certain temperature can be computed as ${}_rG = {}_rH - T \cdot {}_rS$ and the change in the value of ${}_rH$ and ${}_rS$ for a reaction with temperature can be computed as follows :

$${}_rH_{T_2} = {}_rH_{T_1} + {}_rC_p (T_2 - T_1)$$

$${}_rS_{T_2} = {}_rS_{T_1} + {}_rC_p \ln \frac{T_2}{T_1}$$

$${}_rG = {}_rH - T \cdot {}_rS$$

and

$$\text{by } {}_rG = RT \ln K_{\text{eq.}}$$

Consider the following reaction : $\text{CO(g)} + 2\text{H}_2\text{(g)} \rightleftharpoons \text{CH}_3\text{OH(g)}$

Given : ${}_fH$ ($\text{CH}_3\text{OH, g}$) 201 kJ/mol; ${}_fH$ (CO, g) 114 kJ/mol

${}_rS$ ($\text{CH}_3\text{OH, g}$) 240 J/K-mol ; ${}_rS$ (H_2, g) 29 JK⁻¹ mol⁻¹

${}_rS$ (CO, g) 198 J/mol-K ; ${}_rC_{p,m}$ (H_2) 28.8 J/mol-K

${}_rC_{p,m}$ (CO) 29.4 J/mol-K; ${}_rC_{p,m}$ (CH_3OH) 44 J/mol-K

and $\ln \frac{320}{300} = 0.06$, all data at 300 K

- ${}_rS$ at 300 K for the reaction is :
(a) 152.6 J/K-mol (b) 181.6 J/K-mol (c) 16 J/K-mol (d) None of these
- ${}_rH$ at 300 K for the reaction is :
(a) 87 kJ/mol (b) 87 kJ/mol (c) 315 kJ/mol (d) 288 kJ/mol
- ${}_rS$ at 320 K is :
(a) 155.18 J/mol-K (b) 150.02 J/mol-K (c) 172 J/mol-K (d) None of these
- ${}_rH$ at 320 K is :
(a) 288.86 kJ/mol (b) 289.1 kJ/mol (c) 87.86 kJ/mol (d) None of these
- ${}_rG$ at 320 K is :
(a) 48295.2 kJ/mol (b) 240.85 kJ/mol (c) 240.85 kJ/mol (d) 81.91 kJ/mol



Passage 4

Enthalpy of neutralization is defined as the enthalpy change when 1 mole of acid/base is completely neutralized by base/acid in dilute solution.

For strong acid and strong base neutralization net chemical change is



${}_rH_{\text{ionization}}$ of aqueous solution of strong acid and strong base is zero.

When a dilute solution of a weak acid or base is neutralized, the enthalpy of neutralization is somewhat less because of the absorption of heat in the ionization of the weak acid or base, for weak acid/base



- If enthalpy of neutralization of CH_3COOH by HCl is -49.86 kJ/mol then enthalpy of ionization of CH_3COOH is :
 (a) 5.98 kJ/mol (b) 5.98 kJ/mol (c) 105.7 kJ/mol (d) None of these
- What is ΔH for complete neutralization of strong diacidic base $\text{A}(\text{OH})_2$ by HNO_3 ?
 (a) 55.84 kJ/mol (b) 111.68 kJ/mol (c) 55.84 kJ/mol (d) None of these
- Under the same conditions how many mL of 0.1 M NaOH and $0.05 \text{ M H}_2\text{A}$ (strong diprotic acid) solution should be mixed for a total volume of 100 mL produce the highest rise in temperature :
 (a) $25 : 75$ (b) $50 : 50$ (c) $75 : 25$ (d) $66.66 : 33.33$



Passage 5

Gibbs Helmholtz equation relates the enthalpy, entropy and free energy change of the process at constant pressure and temperature as

$$\Delta G = \Delta H - T \Delta S \quad (\text{at constant } P, T)$$

In General the magnitude of ΔH does not change much with the change in temperature but the term $T \Delta S$ changes appreciably. Hence in some process spontaneity is very much dependent on temperature and such processes are generally known as entropy driven process.

- For the reaction at 298 K ; $\text{A}_2\text{B}_4 \rightleftharpoons 2\text{AB}_2$
 $\Delta H = -2 \text{ kJ}$ and $\Delta S = 20 \text{ J/K}$ at constant P and T , the reaction will be
 (a) spontaneous and entropy driven (b) spontaneous and enthalpy driven
 (c) non-spontaneous (d) at equilibrium
- When CaCO_3 is heated to a high temperature it decomposes into CaO and CO_2 , however it is quite stable at room temperature. It can be explained by the fact that
 (a) $\Delta_r H$ dominates the term $T \Delta_r S$ at high temperature
 (b) the term $T \Delta_r S$ dominates the $\Delta_r H$ at high temperature
 (c) at high temperature both $\Delta_r S$ and $\Delta_r H$ becomes negative
 (d) thermodynamics can not say anything about spontaneity
- The Dissolution of $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ in a large volume of water is endothermic to the extent of $3.5 \text{ kcal mol}^{-1}$ and ΔH for the reaction is $-23.2 \text{ kcal mol}^{-1}$.




Select the correct statement :

- $\Delta H_{\text{solution}}$ for anhydrous CaCl_2 is -19.7 kcal/mol and the process is enthalpy driven
- $\Delta H_{\text{solution}}$ for anhydrous CaCl_2 is -19.7 kcal/mol and the process is entropy driven
- Dissolution of $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ in water is enthalpy driven process
- The $\Delta_r S$ the reaction $\text{CaCl}_2(s) + 6\text{H}_2\text{O}(l) \rightleftharpoons \text{CaCl}_2 \cdot 6\text{H}_2\text{O}(s)$ is negative

One or More Answers is/are Correct



- Identify the intensive quantities from the following :
 (a) Enthalpy (b) Temperature (c) Pressure (d) Mass
- Identify the extensive quantities from the following :
 (a) Gibb's energy (b) Entropy (c) Refractive index (d) Specific heat
- Identify the state functions from the following :
 (a) Heat (b) Work (c) Enthalpy (d) Enthalpy change
- Which of the following statements is/are correct as per IUPAC sign convention?
 (a) The work done by the system on the surrounding is negative
 (b) The work done by the surrounding on the system is positive
 (c) The heat absorbed by the system from the surrounding is positive
 (d) The heat absorbed by the surrounding from the system is positive
- In an isothermal irreversible expansion of an ideal gas as per IUPAC sign convention :
 (a) $U = 0$ (b) $H = 0$ (c) $w = -nRT \ln \frac{P_1}{P_2}$ (d) $w = q$
- In reversible isothermal expansion of an ideal gas :
 (a) $w = 0$ (b) $U_1 = U_2$ (c) $H_1 = H_2$ (d) $q = nRT \ln \frac{V_2}{V_1}$
- An adiabatic process is that process in which :
 (a) energy is transferred as heat
 (b) no energy is transfer red as heat
 (c) $U = w$
 (d) the temp. of gas increases in a reversible adiabatic compression
- In adiabatic process, the work involved during expansion or compression of an ideal gas is given by :
 (a) $nC_v T$ (b) $\frac{nR}{1} (T_2 - T_1)$
 (c) $nRP_{\text{ext}} \frac{T_2 P_1 - T_1 P_2}{P_1 P_2}$ (d) $2.303 RT \log \frac{V_2}{V_1}$
- One mole of an ideal gas is subjected to a two step reversible process (A—B and B—C). The pressure at A and C is same. Mark the correct statement(s) :
 (a) Work involved in the path AB is zero
 (b) In the path AB work will be done on the gas by the surrounding
 (c) Volume of gas at C = 3 volume of gas at A
 (d) Volume of gas at B is 16.42 litres
- Assume ideal gas behaviour for all the gases considered and neglect vibrational degrees of freedom. Separate equimolar samples of Ne, O₂, CO₂ and SO₂ were subjected to a two process as mentioned. Initially all are at same state of temperature and pressure.



- Step I All undergo reversible adiabatic expansion to attain same final volume, which is double the original volume thereby causing the decreases in their temperature.
- Step II After step I all are given appropriate amount of heat isochorically to restore the original temperature.

Mark the correct option(s) :

- (a) Due to step I only, the decrease in temperature will be maximum for Ne
 (b) During step II, heat given will be minimum for SO_2
 (c) There will be no change in internal energy for any of the gas after both the steps of process are completed
 (d) The P - V graph of O_2 and CO_2 will be same
11. Which of the following is/are correct?
 (a) H U (PV) when P and V both changes
 (b) H U P V when pressure is constant
 (c) H U V P when volume is constant
 (d) H U P V V P when P and V both changes
12. H U for the reaction(s) :
 (a) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$ (b) $\text{Ag}_2\text{O}(\text{s}) \rightarrow 2\text{Ag}(\text{s}) + \frac{1}{2}\text{O}_2(\text{g})$
 (c) $\text{CO}(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g})$ (d) $\text{C}(\text{s}) + \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g})$
13. Which of the following conditions may lead to a non-spontaneous change?
 (a) H and S both + ve (b) H ve; S ve
 (c) H ve; S ve (d) H ve; S ve
14. For a process to be spontaneous :
 (a) $(G_{\text{system}})_{T, p} < 0$ (b) $S_{\text{system}} + S_{\text{surrounding}} < 0$
 (c) $S_{\text{system}} - S_{\text{surrounding}} < 0$ (d) $(G_{\text{system}})_{T, p} > 0$
15. The normal boiling point of a liquid 'X' is 400 K. Which of the following statement is true about the process $X(\text{l}) \rightarrow X(\text{g})$?
 (a) at 400 K and 1 atm pressure $G < 0$ (b) at 400 K and 2 atm pressure $G < 0$
 (c) at 400 K and 0.1 atm pressure $G < 0$ (d) at 410 K and 1 atm pressure $G < 0$
16. When ice melts at 1°C :
 (a) an increase in entropy (b) a decrease in enthalpy
 (c) a decrease in free energy (d) process is spontaneous
17. The value of $H_{\text{transition}}$ of C (graphite) $\rightarrow$ C (diamond) is 1.9 kJ/mol at 25°C . Entropy of graphite is higher than entropy of diamond. This implies that :
 (a) C (diamond) is more thermodynamically stable than C (graphite) at 25°C
 (b) C (graphite) is more thermodynamically stable than C (diamond) at 25°C
 (c) diamond will provide more heat on complete combustion at 25°C
 (d) $G_{\text{transition}}$ of C (diamond) $\rightarrow$ C (graphite) is ve
18. Which of the following statement(s) is/are false?
 (a) All adiabatic processes are isentropic (or isentropic) processes
 (b) When $(G_{\text{system}})_{T, p} < 0$; the reaction must be exothermic

- (c) $dG = VdP - SdT$ is applicable for closed system, both PV and non- PV work
- (d) The heat of vaporisation of water at 100°C is 40.6 kJ/mol . When 9 gm of water vapour condenses to liquid at 100°C and 1 atm , then $S_{\text{system}} = 54.42 \text{ J/K}$
- 19.** Which of the following statement(s) is/are true?
- (a) $U = 0$ for combustion of $\text{C}_2\text{H}_6(\text{g})$ in a sealed rigid adiabatic container
- (b) $\Delta_f H^\circ$ (S , monoclinic) $= 0$
- (c) If dissociation energy of $\text{CH}_4(\text{g})$ is 1656 kJ/mol and $\text{C}_2\text{H}_6(\text{g})$ is 2812 kJ/mol , then value of $\text{C}-\text{C}$ bond energy will be 328 kJ/mol
- (d) If $\Delta_f H^\circ(\text{H}_2\text{O}, \text{g}) = 242 \text{ kJ/mol}$; $\Delta_{\text{vap}} H^\circ(\text{H}_2\text{O}, \text{l}) = 44 \text{ kJ/mol}$ then, $\Delta_f H^\circ(\text{OH}^-, \text{aq.})$ will be -142 kJ/mol
- 20.** From the following data, mark the option(s) where $\Delta_r H^\circ$ is correctly written for the given reaction.
- Given : $\Delta_f H^\circ(\text{OH}^-, \text{aq.}) = -230 \text{ kJ/mol}$; $\Delta_f H^\circ(\text{H}_2\text{O}(\text{l})) = -285.8 \text{ kJ/mol}$
- $\Delta_{\text{solution}} H^\circ$ of $\text{HA}(\text{g}) = 70.7 \text{ kJ/mol}$
- $\Delta_{\text{solution}} H^\circ$ of $\text{BOH}(\text{g}) = 20 \text{ kJ/mol}$
- $\Delta_{\text{ionization}} H^\circ$ of $\text{HA} = 15 \text{ kJ/mol}$ and BOH is a strong base.
- | Reaction | $\Delta_r H^\circ (\text{kJ/mol})$ |
|--|------------------------------------|
| (a) $\text{HA}(\text{aq}) + \text{BOH}(\text{aq}) \rightarrow \text{BA}(\text{aq}) + \text{H}_2\text{O}$ | 42.3 |
| (b) $\text{HA}(\text{g}) + \text{BOH}(\text{g}) \rightarrow \text{BA}(\text{aq}) + \text{H}_2\text{O}$ | 93 |
| (c) $\text{HA}(\text{g}) + \text{H}^+(\text{aq}) \rightarrow \text{A}^-(\text{aq})$ | 55.7 |
| (d) $\text{B}^-(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{BOH}(\text{aq})$ | 20 |
- 21.** Select correct statement(s)
- (a) An adiabatic system can exchange energy with its surroundings.
- (b) A thermodynamic property which is intensive is additive.
- (c) Work done may be zero in a cyclic process.
- (d) For a simple compressible substance, the relation $\oint dq = \oint P \cdot dv = 0$ is true for any cycle involving mechanical work only.
- 22.** For an isolated system, the entropy :
- (a) either increases or remains constant
- (b) either decreases or remains constant
- (c) can never decrease
- (d) can never increase
- 23.** The normal boiling point of a liquid X is 400 K . $\Delta_{\text{vap}} H^\circ$ at normal boiling point is 40 kJ/mol . Select correct statement(s) :
- (a) $S_{\text{Vaporisation}} = 100 \text{ J/mol.K}$ at 400 K and 2 atm
- (b) $S_{\text{Vaporisation}} = 10 \text{ J/mol.K}$ at 400 K and 1 atm
- (c) $G_{\text{Vaporisation}} = 0$ at 410 K and 1 atm
- (d) $U = 43.32 \text{ kJ/mol.K}$ at 400 K and 1 atm
- 24.** Select incorrect statement(s)
- (a) A closed system with all adiabatic boundaries must be an isolated system
- (b) Total heat exchange in a cyclic process may be zero
- (c) Entropy of a closed system is maximum at equilibrium
- (d) Molar Gibb's Energy is an extensive property

25. Select correct statement(s) for the reaction $\text{H}_2\text{O}(g) + \text{CO}(g) \rightleftharpoons \text{H}_2(g) + \text{CO}_2(g)$

Substance	$\text{CO}(g)$	$\text{CO}_2(g)$	$\text{H}_2\text{O}(g)$	$\text{H}_2(g)$
${}_fH_{400}(\text{kcal mol}^{-1})$	25	95	55	0
$S_{400}(\text{cal mol}^{-1} \text{K}^{-1})$	45	50	40	30

(a) Reaction is enthalpy driven

(b) Reaction is entropy driven

(c) Reaction is spontaneous at 400 K

(d) Reaction is non-spontaneous at 400 K

Match the Column

Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I

- (A) Reversible cooling of an ideal gas at constant volume
- (B) Reversible isothermal expansion of an ideal gas
- (C) Adiabatic expansion of non-ideal gas into vacuum
- (D) Reversible melting of sulphur at normal melting point

Column-II

- (P) $w = 0; q < 0; U < 0$
- (Q) $w < 0; q < 0; U < 0$
- (R) $w = 0; q < 0; U < 0$
- (S) $w = 0; q = 0; U = 0$

2.

Column-I

- (A) Adiabatic process
- (B) Isothermal process
- (C) Isoenthalpic process
- (D) Isoentropic process

Column-II

- (P) $q = 0$
- (Q) $H = 0$
- (R) $T = 0$
- (S) $S = 0$

3.

Column-I

- (A) Reversible isothermal expansion of an ideal gas
- (B) Reversible adiabatic compression of an ideal gas
- (C) Irreversible adiabatic expansion of an ideal gas
- (D) Irreversible isothermal compression of an ideal gas

Column-II

- (P) $w = 2.303 nRT \log \frac{V_2}{V_1}$
- (Q) $PV = \text{constant}$
- (R) $w = \frac{nR}{(\gamma - 1)} (T_2 - T_1)$
- (S) $H = 0$

4.

Column-I

- (A) $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$
in a closed system
- (B) $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$
in a closed system
- (C) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$
in a closed system
- (D) $\text{CO}(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightleftharpoons \text{CO}_2(\text{g})$
in an isolated system

Column-II

- (P) $H = U$
- (Q) $H = U = 0$
- (R) $H = U$
- (S) $U = 0$

5.

Column-I (Process)

- (A) Reversible isothermal compression of an ideal gas
- (B) Isothermal free expansion ($P_{\text{ext}} = 0$) of an ideal gas
- (C) Reversible adiabatic expansion of an ideal gas
- (D) Reversible ideal gas expansion

**Column-II
(Entropy Change)**

- (P) $S_{\text{system}} = 0$
- (Q) $S_{\text{system}} > 0$
- (R) $S_{\text{system}} < 0$
- (S) Information insufficient

6.

Column-I (Process)

- (A) Reversible isothermal ideal gas expansion
- (B) Reversible adiabatic ideal gas compression
- (C) Adiabatic free expansion ($P_{\text{ext}} = 0$) of an ideal gas
- (D) Irreversible isothermal ideal gas compression

**Column-II
(Entropy change)**

- (P) $S_{\text{surrounding}} = 0$
- (Q) $S_{\text{surrounding}} > 0$
- (R) $S_{\text{surrounding}} < 0$
- (S) $S_{\text{system}} = 0$

7.

Column-I

- (A) $(\Delta G_{\text{system}})_{T, P} = 0$
- (B) $S_{\text{system}} - S_{\text{surrounding}} > 0$
- (C) $S_{\text{system}} - S_{\text{surrounding}} < 0$
- (D) $(\Delta G_{\text{system}})_{T, P} < 0$

Column-II

- (P) Process is in equilibrium
- (Q) Process is nonspontaneous
- (R) Process is spontaneous
- (S) System is unable to do useful work

8.

Column-I

- (A) $(G_{\text{system}})_{T, P}$
- (B) Work done in reversible isothermal ideal gas expansion
- (C) G for reversible isothermal expansion of an ideal gas
- (D) S_{gas} for isothermal expansion of an ideal gas

Column-II

- (P) $nR \ln \frac{V_2}{V_1}$
- (Q) $nRT \ln \frac{P_2}{P_1}$
- (R) nFE
- (S) $nR \ln \frac{P_1}{P_2}$

9.

Column-I

- (A) Heating of an ideal gas at constant pressure
- (B) Compression of liquid at constant temperature
- (C) Reversible process for an ideal gas at constant temperature
- (D) Adiabatic free expansion of an ideal gas

Column-II

- (P) $H = nC_{p, m} \Delta T$
- (Q) $U = 0$
- (R) $G = V \Delta P$
- (S) $G = nRT \ln \frac{P_2}{P_1}$

10.

Column-I

(Sign of H and S respectively)

- (A) $+$ & $+$
- (B) $+$ & $-$
- (C) $+$ & $+$
- (D) $-$ & $-$

Column-II

(Nature of reaction)

- (P) Spontaneous only at low temperature
- (Q) Spontaneous only at high temperature
- (R) Spontaneous at all temperature
- (S) Non-spontaneous at all temperature

11.

Column-I

- (A) Reversible adiabatic compression
- (B) Reversible vaporisation of liquid
- (C) $2N_2(g) \rightarrow N_2(g)$
- (D) $MgCO_3(s) \rightarrow MgO(s) + CO_2(g)$

Column-II

- (P) $S_{\text{system}} = 0$
- (Q) $S_{\text{system}} = 0$
- (R) $S_{\text{surrounding}} = 0$
- (S) $S_{\text{surrounding}} = 0$

12.

Column-I

- (A) $H(aq)$
 (B) $H(g)$
 (C) $H_2(g)$
 (D) $C(s, \text{diamond})$

Column-II

- (P) ${}_fH \quad 0$
 (Q) ${}_fH \quad 0$
 (R) ${}_fG \quad 0$
 (S) ${}_fS \quad 0$

13.

Column-I

(Partial derivative)

- (A) $\frac{U}{T}_V$
 (B) $\frac{H}{T}_P$
 (C) $\frac{G}{T}_P$
 (D) $\frac{G}{P}_T$

Column-II

(Thermodynamic variable)

- (P) C_P
 (Q) C_V
 (R) S
 (S) V

14.

Column-I

- (A) $C(s, \text{graphite}) \quad O_2(g) \quad CO_2(g)$
 (B) $C(s, \text{graphite}) \quad C(g)$
 (C) $CO(g) \quad \frac{1}{2} O_2(g) \quad CO_2(g)$
 (D) $CH_4(g) \quad C(g) \quad 4H(g)$

Column-II

- (P) $H_{\text{combustion}}$
 (Q) $H_{\text{formation}}$
 (R) $H_{\text{atomization}}$
 (S) $H_{\text{sublimation}}$

15.

Column-I

- (A) $O_2(g)$
 (B) $O_3(g)$
 (C) $Br_2(g)$
 (D) $H_2O(l)$

Column-II

- (P) ${}_fH \quad \text{ve}; \quad {}_fS \quad \text{ve}$
 (Q) ${}_fH \quad \text{ve}; \quad {}_fS \quad \text{ve}$
 (R) ${}_fH \quad \text{ve}; \quad {}_fS \quad \text{ve}$
 (S) ${}_fH \quad 0; \quad {}_fS \quad 0$

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

- (A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** Heat and work are “definite quantities”.
STATEMENT-2 : Heat and work are not properties of a system. Their values depend on the path of the process and vary accordingly.
2. **STATEMENT-1 :** There is no change in internal energy for an ideal gas at constant temperature.
STATEMENT-2 : Internal energy of an ideal gas is a function of temperature only.
3. **STATEMENT-1 :** The heat absorbed during the isothermal expansion of an ideal gas against vacuum is zero.
STATEMENT-2 : The volume occupied by the molecules of an ideal gas is zero.
4. **STATEMENT-1 :** H and U are the same for the reaction, $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$,
STATEMENT-2 : All reactants and products are gases where all gases are ideal.
5. **STATEMENT-1 :** The magnitude of the work involved in an isothermal expansion is greater than that involved in an adiabatic expansion.
STATEMENT-2 : P V curve (P on y -axis and V on x -axis) decrease more rapidly for reversible adiabatic expansion compared to reversible isothermal expansion starting from same initial state.
6. **STATEMENT-1 :** Entropy change in reversible adiabatic expansion of an ideal gas is zero.
STATEMENT-2 : The increase in entropy due to volume increase just compensate the decrease in entropy due to fall in temperature.
7. **STATEMENT-1 :** There cannot be chemical equilibrium in an open system.
STATEMENT-2 : There is no fixed mass in an open system.
8. **STATEMENT-1 :** The standard free energy changes of all spontaneously occurring reactions are negative.
STATEMENT-2 : The standard free energies of the elements in their standard states at 1 bar and 298 K are taken as zero.
9. **STATEMENT-1 :** Enthalpy and entropy of any elementary substance in the standard states are taken as zero.
STATEMENT-2 : At absolute zero, particles of the perfectly crystalline substance become completely motionless.
10. **STATEMENT-1 :** A reaction which is spontaneous and accompanied by decrease of randomness must be exothermic.
STATEMENT-2 : All exothermic reactions are accompanied by decrease of randomness.

- 11. STATEMENT-1 :** Many endothermic reactions that are not spontaneous at room temperatures become spontaneous at high temperature.
STATEMENT-2 : H of the endothermic reaction may increase with increase in temperature.
- 12. STATEMENT-1 :** Decrease of free energy during the process under constant temperature and pressure provides a measure of its spontaneity.
STATEMENT-2 : A spontaneous change must have +ve sign of ΔS_{system} .
- 13. STATEMENT-1 :** $|\Delta_f H|$ of $(\text{H}_2\text{O}, l)$ $>$ $|\Delta_f H|$ of $(\text{H}_2\text{O}, g)$
STATEMENT-2 : $\Delta H_{\text{condensation}}$ is negative.
- 14. STATEMENT-1 :** All combustion reactions are exothermic.
STATEMENT-2 : Enthalpies of products are greater than enthalpies of reactants ($\sum \nu_p \Delta_f H(P) > \sum \nu_R \Delta_f H(R)$)
- 15. STATEMENT-1 :** Enthalpy of neutralization of CH_3COOH by NaOH is less than that of HCl by NaOH .
STATEMENT-2 : Enthalpy of neutralization of CH_3COOH is less because of the absorption of heat in the ionization process.
- 16. STATEMENT-1 :** Internal energy of a real gas may change during expansion at constant temperature.
STATEMENT-2 : Internal energy of a real gas is a function of temperature and volume.
- 17. STATEMENT-1 :** Work is a state function which is expressed in joule.
STATEMENT-2 : Work appears only at the boundary of the system.
- 18. STATEMENT-1 :** The expansion of a gas into an evacuated space takes place non-spontaneously.
STATEMENT-2 : A process in which all steps cannot be retraced by themselves is called a spontaneous process.

Subjective Problems



- A perfect gas undergoes a reversible adiabatic expansion from (300 K, 200 atm) to (90 K, 10 atm). Find the atomicity of gas.
- 5 mole of an ideal gas at temp. T are compressed isothermally from 12 atm. to 24 atm. Calculate the value of $10r$
 Where, $r = \frac{\text{Work done along reversible process}}{\text{Work done along single step irreversible process}}$ (Given : $\ln 2 = 0.7$)
- A diatomic ideal gas is expanded according to PV^3 constant, under very high temperature (Assume vibration mode active). Calculate the molar heat capacity of gas (in cal / molK) in this process.
- A heat engine is operating between 500 K to 300 K and it absorbs 10 kcal of heat from 500 K reservoir reversibly per cycle. Calculate the work done (in kcal) per cycle.
- In a carnot cycle involving ideal non-linear triatomic gas, if during adiabatic expansion volume increases from 2L to 16L and heat absorbed during isothermal expansion is 8 kcal, then calculate magnitude of work done by carnot engine (in kcal).

6. Molar heat capacities at constant pressure for A, B and C are 3, 1.5 and 2 J/K mol. The enthalpy of reaction and entropy of reaction, $A \rightarrow 2B + 3C$ are 20 kJ/mol and 20 J/K mol at 300 K. Calculate ΔG (in kJ/mol) for the reaction,



7. Standard molar enthalpy of combustion of glucose is -2880 kJ. If only 25% of energy is available for muscular work and 1.0 km walk consumes 90 kJ of energy, what maximum distance (in km) a person can walk after eating 90 g of glucose.

8. Given $C_2H_2(g) + H_2(g) \rightarrow C_2H_4(g)$; $\Delta H = 175 \text{ kJ mol}^{-1}$


$$\Delta H_f(C_2H_4, g) = 50 \text{ kJ mol}^{-1}; \Delta H_f(H_2O, l) = 280 \text{ kJ mol}^{-1}; \Delta H_f(CO_2, g) = 390 \text{ kJ mol}^{-1}$$

If ΔH is enthalpy of combustion (in kJ mol⁻¹) of $C_2H_2(g)$, then calculate the value of $\left| \frac{\Delta H}{257} \right|$.

9. The integral enthalpies of solution of anhydrous $CuSO_4(s)$ and hydrated $CuSO_4 \cdot 5H_2O(s)$ are 70 kJ per mol and 10 kJ per mol respectively. Determine the magnitude of enthalpy of hydration of 0.1 mole anhydrous $CuSO_4(s)$ as $CuSO_4(s) + 5H_2O(l) \rightarrow CuSO_4 \cdot 5H_2O(s)$
10. If enthalpy of neutralisation of HCl by NaOH is 57 kJ mol⁻¹ and with NH_4OH is 50 kJ mol⁻¹. Calculate enthalpy of ionisation of $NH_4OH(aq)$.
11. Lattice energy of NaCl(s) is 790 kJ mol⁻¹ and enthalpy of hydration is 785 kJ mol⁻¹. Calculate enthalpy of solution of NaCl(s).
12. x g sample of NH_4NO_3 is decomposed in a Bomb calorimeter. The temperature of calorimeter increase by 4 °C. The heat capacity of the system is 1.25 kJ/°C. Calculate the value of x . Given molar heat of decomposition of NH_4NO_3 at constant volume is 400 kJ mol⁻¹.
13. A heat engine operating between 227°C and 77°C absorbs 10 kcal of heat from the 227°C reservoir reversibly per cycle. Calculate total work done (in kcal) in two cycles.
14. Calculate work done in chemical reaction (in kcal)
 $A(s) + 3B(g) \rightarrow C(l)$ at 227°C at 1 atm in closed vessel.
15. One mole ideal monoatomic gas is heated according to path AB and AC.

If temperature of state B and state C are equal.

Calculate $\frac{q_{AC}}{q_{AB}}$ 10.



ANSWERS

Level 1

1. (d)	2. (b)	3. (d)	4. (c)	5. (a)	6. (c)	7. (a)	8. (c)	9. (d)	10. (d)
11. (c)	12. (b)	13. (d)	14. (d)	15. (a)	16. (b)	17. (b)	18. (a)	19. (b)	20. (c)
21. (c)	22. (c)	23. (c)	24. (d)	25. (b)	26. (c)	27. (b)	28. (c)	29. (c)	30. (d)
31. (a)	32. (c)	33. (b)	34. (b)	35. (c)	36. (a)	37. (a)	38. (a)	39. (b)	40. (b)
41. (b)	42. (c)	43. (b)	44. (b)	45. (b)	46. (c)	47. (a)	48. (b)	49. (c)	50. (c)
51. (a)	52. (b)	53. (d)	54. (b)	55. (b)	56. (b)	57. (b)	58. (b)	59. (b)	60. (b)
61. (a)	62. (c)	63. (c)	64. (d)	65. (c)	66. (a)	67. (a)	68. (c)	69. (d)	70. (c)
71. (d)	72. (c)	73. (b)	74. (b)	75. (c)	76. (c)	77. (b)	78. (d)	79. (a)	80. (b)
81. (b)	82. (b)	83. (c)	84. (d)	85. (d)	86. (b)	87. (b)	88. (a)	89. (b)	90. (b)
91. (a)	92. (d)	93. (d)	94. (c)	95. (b)	96. (a)	97. (a)	98. (a)	99. (a)	100. (b)
101. (c)	102. (d)	103. (a)	104. (d)	105. (a)	106. (a)	107. (b)	108. (a)	109. (b)	110. (c)
111. (a)	112. (d)	113. (c)	114. (a)	115. (b)	116. (a)	117. (c)	118. (b)	119. (c)	120. (c)
121. (c)	122. (c)	123. (b)	124. (a)	125. (a)	126. (c)	127. (c)	128. (b)	129. (b)	130. (a)
131. (a)	132. (b)	133. (a)	134. (a)	135. (b)	136. (b)	137. (a)	138. (a)	139. (c)	140. (b)
141. (a)	142. (c)	143. (b)	144. (d)	145. (a)	146. (d)	147. (b)	148. (a)	149. (a)	150. (b)



Level 2

1. (c)	2. (c)	3. (c)	4. (a)	5. (d)	6. (c)	7. (c)	8. i (c), ii (b)	9. (c)	10. (a)
11. (d)	12. (b)	13. (b)	14. (d)	15. (a)	16. (b)	17. (c)	18. (d)	19. (d)	20. (a)
21. (a)	22. (d)	23. (c)	24. (c)	25. (d)	26. (b)	27. (a)	28. (d)	29. (b)	30. (b)
31. (b)	32. (c)	33. (b)	34. (d)	35. (a)	36. (a)	37. (c)	38. (c)	39. (b)	40. (d)



Level 3

Passage-1	1. (d)	2. (d)				
Passage-2	1. (b)	2. (c)	3. (b)	4. (c)		
Passage-3	1. (c)	2. (a)	3. (d)	4. (c)	5. (d)	
Passage-4	1. (a)	2. (b)	3. (b)			
Passage-5	1. (a)	2. (b)	3. (a)			

One or More Answers is/are Correct

- | | | | | |
|-------------|-------------|---------------|-------------|-------------|
| 1. (b,c) | 2. (a,b) | 3. (c) | 4. (a,b,c) | 5. (a,b,d) |
| 6. (b,c,d) | 7. (b,c,d) | 8. (a,b,c) | 9. (c,d) | 10. (a,c,d) |
| 11. (a,b,c) | 12. (a,c) | 13. (a,c,d) | 14. (b,d) | 15. (a,b,c) |
| 16. (a,c,d) | 17. (b,c,d) | 18. (a,b,c,d) | 19. (a,b,c) | 20. (a,b,c) |
| 21. (a,c,d) | 22. (a,c) | 23. (a,c) | 24. (a,c,d) | 25. (a,c) |

Match the Column

- | | | | | | | | |
|-------|-------|---|-------|---|-------|---|------|
| 1. A | P; | B | S; | C | R; | D | Q |
| 2. A | P; | B | R; | C | Q; | D | S |
| 3. A | P, S; | B | Q,R; | C | R; | D | S |
| 4. A | R; | B | Q; | C | P; | D | S |
| 5. A | Q; | B | P; | C | R; | D | S |
| 6. A | Q; | B | P, S; | C | P; | D | R |
| 7. A | P, S; | B | R; | C | Q,S; | D | Q,S |
| 8. A | R; | B | Q; | C | Q; | D | P, S |
| 9. A | P; | B | R; | C | Q, S; | D | Q,S |
| 10. A | P; | B | R; | C | Q; | D | S |

11. A S; B P, R; C Q; D P, R
 12. A P, R; B Q; C P, R; D Q, S
 13. A Q; B P; C R; D S
 14. A P, Q; B Q, R, S; C P; D R
 15. A S; B R; C P; D Q

Assertion-Reason Type Questions

1. (D) 2. (A) 3. (B) 4. (B) 5. (A) 6. (A) 7. (A) 8. (B) 9. (D) 10. (C)
 11. (B) 12. (C) 13. (A) 14. (C) 15. (A) 16. (A) 17. (D) 18. (D)

Subjective Problems

1.	1	2.	7	3.	6	4.	4	5.	4	6.	7	7.	4	8.	5	9.	8	10.	7
11.	5	12.	1	13.	6	14.	3	15.	8										

Hints and Solutions

Level 1

1. (d) Heat capacity and volume are extensive properties.
2. (b) Boiling point, pH and emf are intensive property.
3. (d) If system is subjected to chemical equilibrium, mechanical equilibrium and thermal equilibrium then system is consider in thermodynamic equilibrium.
4. (c) Ozone have maximum internal energy

$$E = \frac{3RT}{2} + \frac{3RT}{2} = (3N - 6)RT$$
 For non-linear molecule

$$E = \frac{3RT}{2} + \frac{2RT}{2} = (3N - 5)RT$$
 For linear molecule.
5. (a) Heat lost by iron + Heat gained by water = 0
 $10 \times 0.45 (T - 373) + 25 \times 4.2 (T - 300) = 0$
 $T = 303 \text{ K or } 30^\circ\text{C}$
6. (c) In freezing heat released so $q > 0$
7. (a) $w = -P \Delta V$
8. (c) $W = n_g RT$ ve
 II. $n_g = 2$, III. $n_g = 2$
9. (d) $w = P_{\text{ext}} \Delta V = n_g RT$ and n_g is ve for I and III
 also V ve for IV
10. (d) $q = 0$, $U = q + w$
 $w = 0$ so $U = 0$
11. (c) $U = q + w$
 $300 = 400 + 0.5 \times V \times 100$
 $V = 2 \text{ L}$
12. (b) $U = q + w$
 $10 = 1000 + 2 \times (20) \times 101.3$
 5948 J
13. (d) $w = -P \Delta V = 0$
14. (d) $w = -P \Delta V = -4 \times 10^5 \times (3 - 1) \times 10^{-3} = -0.8 \times 10^3 \text{ J}$
15. (a) $w = -n_g RT \ln \frac{V_2}{V_1} = -8.314 \times 298 \times \ln \frac{4.5}{2}$

$$w = 163.9 \text{ J}$$

$$16. (b) w = -P \Delta V = -8.314 \times 298 \times \ln \frac{1000}{2} = 4.955 \text{ kJ}$$

$$17. (b) w = nR(T_2 - T_1), w = 1R(2) = 2R$$

$$18. (a) dw = P \cdot dV$$

$$w_{\text{rev}} = \int P \cdot dV$$

$$= \int \frac{nRT}{V} \cdot dV = nRT \ln \frac{V_2}{V_1}$$

$$w = 5200 \text{ kJ}$$

so, work done by the gas is 5200 kJ.

$$19. (b) w = \text{Area of } P-V \text{ curve}$$

$$20. (c) \text{ For cyclic process : } dU = 0$$

$$w = -q$$

Net work done by system = Area of the circle

21. (c) Process direction in $V-P$ diagram is clockwise so process direction in $P-V$ diagram is anti-clockwise.

$$\text{Net work done by system} = \text{Area} = \frac{(P_2 - P_1)(V_2 - V_1)}{2}$$

22. (c) Work done in the cyclic process
 Area bounded (ABCA) = $5P_1V_1$

$$23. (c) w = -P \Delta V = 0$$

$$U = q + w = 0$$

Temperature remains constant.


24. (d) Zero


Internal energy is state function.

$$27. (b) C_p - C_v = nR = 2R$$

28. (c) No intramolecular force act between gaseous molecule.

29. (c)





31. (a) $V = \frac{nRT}{P} = \frac{100 \times 0.0821 \times T}{8.21}$

$\log V = \log T$

32. (c) $dw = P \cdot dV$

$w = 20 \frac{dV}{V} = 20 \ln \frac{V_2}{V_1}$

$w = 46.06 \text{ L-atm} = 4665.8 \text{ J}$

$U = q = w = 400 \text{ J} = 4665.8 \text{ J}$

$q = 5065.8 \text{ J}$

33. (b) $w = nRT \ln \frac{V_2}{V_1}$

$w = 2 \times 8.314 \times 10^3 \times 300 \ln \frac{40}{4}$

$w = 11.488 \text{ J}$

34. (b) $w = nRT \ln \frac{P_1}{P_2}$

$10 \times 8.314 \times 300 \ln \frac{10}{1}$

57441.42 J

$w = mgh$

$m = 9.81 \times 100 = 57441.42$

$m = 58.55 \text{ kg}$

35. (c) At constant volume $U = q_v = nC_{v,m} T$
at constant pressure $H = q_p = nC_{p,m} T$

36. (a) $U = w = 75 \times 0.1 \times \frac{3}{2} \times 2 [T_2 - 500]$

$T_2 = 250 \text{ K}$

37. (a) $U = w = \frac{P_2 V_1 - P_1 V_1}{1}$

38. (a) $T^{-1} \cdot P = \text{constant}$ or $P = T^{-1}$

$\therefore \frac{P}{T^3} = \text{constant}$
 $\frac{1}{1} = 3; \frac{3}{2}$

39. (b) $T = \frac{1}{\sqrt{V}}; TV^{1/2} = \text{constant}$

For adiabatic process,
 $TV^{\frac{1}{2}} = \text{constant}$

$1 \frac{1}{2}, \frac{3}{2}$

40. (b) $PV = \text{constant}, P = \frac{1}{V} \frac{dV}{dP} = 0$

41. (b) $PV = \text{constant}$ for isothermal process
 $PV = \text{constant}$ for adiabatic process so more is the value of γ , more is decrease in pressure.

42. (c) $T \cdot V^{\gamma-1} = \text{constant}$

$\frac{T_2}{T_1} = \frac{V_1}{V_2}^{\gamma-1}$

$T_2 = 300 \times \left(\frac{16}{2}\right)^{\frac{5}{3}-1}$

1200 K

43. (b) $dw = P_{\text{ext}} dV$

$w_{\text{irr}} = P_{\text{ext}} \frac{nRT}{P_2} = \frac{nRT}{P_1}$

$w_{\text{irr}} = 1 \times (5 \times 8.314 \times 300) = 1 \times \frac{1}{10}$

$w_{\text{irr}} = 11.224 \text{ kJ}$

44. (b) For an adiabatic irreversible compression,
 $U = w$

$nC_{v,m} (T_2 - T_1) = P_{\text{ext}} (V_2 - V_1)$
here $P_{\text{ext}} = P_2$

$nC_{v,m} (T_2 - T_1) = P_2 \frac{nRT_2}{P_2} - \frac{nRT_1}{P_1}$

$\frac{5}{2} R (T_2 - T_1) = P_2 R \frac{T_2}{P_2} - \frac{T_1}{P_1}$

$P_2 = 275 \text{ kPa}$

45. (b) $w = U = n \int_{T_1}^{T_2} C_{v,m} dT$

$w = 2 \times \frac{R}{1} (T_2 - T_1)$

$\frac{T_2 - T_1}{1} = P_{\text{ext}} \frac{T_2}{P_2} - \frac{T_1}{P_1}$

$\frac{T_2 - 300}{\frac{5}{3}} = 2 \times \frac{T_2}{2} - \frac{300}{1}$

$1.5 T_2 - 450 = T_2 - 600 \quad T_2 = 420 \text{ K}$

$$w = 2 \times \frac{2}{5} \times \frac{1}{3} (420 - 300) = 720$$

46. (c) $\therefore q = 0$

$$nC_{v,m} (T_2 - T_1) = P_{\text{ext}} \frac{nRT_2}{P_2} - \frac{nRT_1}{P_1}$$

$$C_{v,m} [T_2 - T_1] = P_{\text{ext}} R \frac{T_1}{P_1} - \frac{T_2}{P_2}$$

$$\frac{5}{2} R [T_2 - 300] = 2 R \left[\frac{300}{5} - \frac{T_2}{2} \right]$$

$$T_2 = 248.5 \text{ K}$$

47. (a) $U = w$

$$n \frac{R}{1} (T_2 - T_1) = P_{\text{ext}} \frac{nRT_2}{P_2} - \frac{nRT_1}{P_1}$$

$$3(T_2 - T_1) = 1 \left[\frac{T_2}{1} - \frac{T_1}{2} \right]$$

$$3T_2 - 3T_1 = T_2 - \frac{T_1}{2}$$

$$4T_2 = \frac{7T_1}{2}$$

$$T_2 = \frac{7T_1}{8} = \frac{7 \times 400}{8} = 350 \text{ K}$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow \frac{2 \times 10}{400} = \frac{1 \times V_2}{350} \Rightarrow V_2 = 17.5 \text{ L}$$

48. (b) Average $C_{v,m}$

$$\frac{n_1 C_{v,m_1} + n_2 C_{v,m_2}}{n_1 + n_2}$$

$$= \frac{2 \times \frac{3}{2} R + 2 \times \frac{5}{2} R}{2 + 2} = 2R$$

49. (c) $H = (n_1 C_{p,m_1} + n_2 C_{p,m_2}) T$

$$0.5 \times \frac{7}{2} R + 0.5 \times 4R = (100)$$

$$375 R$$

50. (c) AB : Isobaric expansion
 BC : Isothermal expansion
 CD : Isochoric
 DA : Isothermal compression

51. (a) 1 watt = 1 J/sec
 Total heat supplied for 36 mL H_2O

$$\frac{806 \times 100}{36} = 2238.9 \text{ J}$$

$$\frac{80600}{36} = 2238.9 \text{ J}$$

$$H_{\text{vap}} = \frac{80600}{36} = 2238.9 \text{ J}$$

$$40300 \text{ J or } 40.3 \text{ kJ/mol}$$

52. (b) $H = U + n_g RT$, $n_g = 1$, $H = U$

53. (d) $n_g = 0$, $H = U$
 For 2 mole $U = 370 \text{ kJ}$

54. (b) $H = U + n_g RT$, $n_g = 2$, $H = U$
 $w = n_g RT$, $w = 0$, q ve

55. (b) At constant pressure, $H = U + P \Delta V$
 $30 \times 2 \times 4 = 240 \text{ J}$
 38 atm-L

56. (b) $H = U + P \Delta V$
 $124 = U + \frac{1520}{760} (177 - 377)$
 $10^3 \times 101.3$
 $U = 83.48 \text{ J}$

57. (b) At constant volume $H = U + V \Delta P$
 $560 = U + 10 \times (30 - 0.1)$
 $U = 530 \text{ kJ}$

58. (b) When both P and V are changing
 $H = U + (PV)$
 $U = (P_2 V_2 - P_1 V_1)$
 $H = 40 \times (20 - 3) = 720 \text{ J}$
 57 L-atm

59. (b) For combustion of 1 mole of benzene

$$n_g = 1.5$$

$$H = U + n_g RT$$

$$3271 = U + \frac{1.5 \times 8.314 \times 300}{1000}$$

$$U = 3267.25 \text{ kJ}$$

For 1.5 mole of combustion of benzene
 $U = 3267.25 \times 1.5 = 4900.88 \text{ kJ}$

60. (b) $H = E + n_g RT$
 $82.8 = U + \frac{1 \times 8.314 \times 298}{1000}$
 $U = 80.32$

61. (a) C (graphite) $\rightarrow$ C (diamond)

$$H = U + P \Delta V$$

$$V_m (\text{diamond}) = \frac{12}{3} \text{ mL}$$

$$V_m (\text{graphite}) = \frac{12}{2} \text{ mL}$$

$$H = U + (500 \times 10^3 - 10^5 \text{ N/m}^2) \left(\frac{12}{3} - \frac{12}{2} \right) \times 10^{-6}$$

$$= -100 \text{ kJ/mol}$$

$U = H + 100 \text{ kJ/mol}$

62. (c) n_g is +ve
63. (b) n_g ve, S ve
64. (d) n_g ve, S ve
 (a) n_g 1 (b) n_g 1
 (c) n_g 0 (d) n_g 3
65. (c) $S = nC_{p,m} \ln \frac{T_2}{T_1}$
 $2 \frac{5}{2} R \ln \frac{600}{300}$
 $5 R \ln 2$
68. (c) $S_{\text{gas}} = nC_{v,m} \ln \frac{T_2}{T_1}$
 $2 \frac{5}{2} 1 R \ln 2$
 $3 R \ln 2$
69. (d) $S = nR \ln \frac{V_2}{V_1} = 1R \ln \frac{3V}{V} = R \ln 3$
70. (c) S will be same because entropy is a state function.
71. (d) $S = nC_{v,m} \ln \frac{T_2}{T_1} = nR \ln \frac{V_2}{V_1}$
 $C_{v,m} \ln 2 = R \ln \frac{1}{2}$
 $S = (C_{v,m} + R) \ln 2$
72. (c) $S = nC_{p,m} \ln \frac{T_2}{T_1}$
 $2.5 \cdot 18 \cdot 4.2 \ln \frac{360}{300}$
 34.02 J/K
73. (b) ${}_rS = \sum \nu_p S (\text{Products}) - \sum \nu_R S (\text{Reactants})$
 $(2 \cdot 27.3 + 3 \cdot 69.9) - (87.4 + 3 \cdot 130.7)$
 $215.2 \text{ JK}^{-1} \text{ mol}^{-1}$
74. (b) ${}_rS = (2 \cdot 248.2) - (2 \cdot 66.5)$
 $(2 \cdot 91.2) - 3(205.1)$
 168.3 J
75. (c) $266 \cdot 6 \cdot 87 \cdot 205 \cdot 4x$
 $x = 145.75$
76. (c) $S = \frac{H}{T}$
77. (b) at M.P. $S_{\text{fusion}} = \frac{H_{\text{fusion}}}{T}$
 $T = \frac{H}{S} = \frac{9.95 \cdot 1000}{35.7} = 278.7 \text{ K}$
78. (d) $S_{\text{freezing}} = -\frac{H_{\text{fusion}}}{T}$
- $S = \frac{80 \cdot 10}{273} = 4.18$
 12.25 J/K
79. (a) $S = \frac{H}{T_{\text{B.P.}}}$
80. (b) $n_{\text{C}_6\text{H}_6} = \frac{117}{78} = 1.5$
 $S_{\text{system}} = 1.5 \cdot 85 \text{ J/K}$
 $S_{\text{surrounding}} = 1.5 \cdot 85 \text{ J/K}$
81. (b) According to 3rd law of thermodynamics at absolute zero temperature entropy of perfectly crystalline substance is taken to be zero.
 $X(s) \xrightarrow{S} X(l)$
82. (b) $S_1 \nearrow$
 $X(l) \xleftarrow{S_2}$
 $S = \frac{20 \cdot 1000}{100} = 1 \cdot \frac{200}{100} C_{p,m} \frac{dT}{T}$
 $S = 200 \cdot 10 \ln 2 = 200 \cdot 10 \cdot 0.693,$
 $S = 206.93 \text{ JK}^{-1}$
83. (c) $0.42 = a(10)^3$ $a = 0.42 \cdot 10^{-3}$
 $S_m = \int_0^{20} \frac{20 C_{p,m}}{T} dT$
 $\int_0^{20} aT^2 dT = \frac{a}{3} [20^3 - 0]$
 1.12 J/K-mol
84. (d) For spontaneous reaction G ve
 S ve (n_g ve)
 H ve
 $G = H - TS$
85. (d) H ve ($H-H$ bond formed)
 S ve (n_g 1)
86. (b) H ve (Combustion reaction)
 S ve (n_g ve)
 $G = H - TS$ ve
87. (b) For ${}_rG$ ve, ${}_rH$ should be ve and S should be +ve
88. (a) For spontaneous reaction
 $(G_{\text{system}})_{T,P}$ ve
89. (b) $(G)_{\text{system}} = H - TS$
 and $H - TS = 0$
90. (b) $G = H - TS$, H ve, S ve
91. (a) At equilibrium $G = 0$
92. (d) $G = H - TS$ ve
 for spontaneous change $H = 0$, $S = 0$

93. (d) G_{799} 40 1000 799 (50),
 G_{799} 50 J spontaneous
 G_{801} 40 1000 801 (50)
 G_{801} 50 J non-spontaneous
94. (c) For ideal gas isothermal expansion $H = 0$
 $G = T \Delta S$
 where $\Delta S = nR \ln \frac{V_2}{V_1}$
95. (b) $\text{Hg}(l) \rightleftharpoons \text{Hg}(g)$,
 rS 174.4 77.4 97 J/K-mol
 $\therefore G = H - TS = 0$
 $T = \frac{H}{S} = \frac{60.8}{0.0001} = 626.8 \text{ K}$
96. (a) At equilibrium $H = T \Delta S$
 $\text{H}_2\text{O}(s) \rightleftharpoons \text{H}_2\text{O}(l)$
 $S_{\text{H}_2\text{O}(l)} - S_{\text{H}_2\text{O}(s)} = 21.8 \text{ J/mol-K}$
 $H = 273 (21.8) = 5951.4 \text{ J/mol}$
97. (a) $G = 2 \times 86.6 + 4 (237.1) - 3 \times 0 - 3 (33.6) - 2 (80.6) = 513.2$
98. (a) $G = H - TS$
 G_{I} 10.5 1000 298 30 1560
 G_{II} 1.8 1000 298 113 35474
 G_{IV} 126 1000 298 84 ve
 III is spontaneous
99. (a) $G = H - TS$
 $S = 242.6 + 50.3 (2 \times 152.3)$, $S = 112.3$
 $G = 788.6 + 1000 \times 112.3$
 $H = 788.6 + 1000 \times 298 = 112.3$
 $H = 822.1 \text{ kJ mol}^{-1}$
100. (b) $G = H - TS$
 $406.6 + 1000 \times H = 298 (2 \times 114.6 + 130.6)$
 $H = 406.6 + 1000 \times 298 (98.6)$
 $H = 436 \text{ kJ mol}^{-1}$
101. (c) $G = RT \ln K$; K stability of reactant
102. (d) $-\frac{1}{T} \ln K_p = \frac{H}{RT^2}$
 $-\frac{1}{T} \ln K = \frac{E_a}{RT^2}$
103. (a) $G = RT \ln K = 8.314 \times 10^3 \times 400 \ln 10^5$
 $G = 38.294 \text{ kJ}$
104. (d) At standard state on equilibrium $G = 0$
 $G = 0 = RT \ln K = K = 1$
105. (a) At equilibrium $G = RT \ln K_{\text{eq}}$
 where $K_{\text{eq}} = (P_{\text{H}_2\text{O}}/P)$ and $P = 1 \text{ bar}$
106. (a) $G = RT \ln K_w$
 $8.314 \times 298 \times 2.303 (14) \sim 80000 \text{ J/mol}$
107. (b) $S = 206.5 + 114.6 \times 180.7 + 88.8$
 $G = H - TS$
 $H = 271.8 + 298 \times 88.8 \times 10^{-3}$
 $H = 298.3$
108. (a) According to Hess's law the standard enthalpy of overall reaction is sum of the enthalpy change in individual reaction.
109. (b) $X + A \rightleftharpoons q_1$
 $q_2 + A \rightleftharpoons B$
 $B + Y \rightleftharpoons q_3$
 $q_2 + X + Y \rightleftharpoons q_1 + q_3$, $H = q_2 + q_1 + q_3$
110. (c) The change in enthalpy does not depend upon number of intermediate steps.
111. (a) $rC_p = 10.945$ for
 $\text{H}_2(g) + \frac{1}{2} \text{O}_2(g) \rightleftharpoons \text{H}_2\text{O}(g)$
 According to Kirchhoff's law
 $rH_{T_2} = rH_{T_1} + rC_p [T_2 - T_1]$
 $rH_{T_2} = (241.82) + [10.94 \text{ J} (373 - 298)/100]$
 $rH_{T_2} = 242.6 \text{ kJ/mol}$
112. (d) fH stability
113. (c) $(fH)_{\text{Al(s)}} = 0$
114. (a) $H = \frac{3}{2} H_1 - \frac{H_2}{2}$
 $713.7 - 177.5 = 891.2 \text{ kJ}$
115. (b) $H = 2 H_1 - 2 H_2 + H_3$
116. (a) $\text{C}(\text{graphite}) \rightleftharpoons \text{C}(\text{diamond})$
 $H = 395 - 393 = 2$
117. (c) $\text{C}(s) + \frac{3}{2} \text{H}_2(g) + \frac{1}{2} \text{Cl}_2(g) \rightleftharpoons \text{CH}_3\text{Cl}(g)$
118. (b) $rH = (fH)_P - (fH)_R$
 $rH = 113 + 3 (269) + 46.2 - 873.8 \text{ kJ}$
119. (c) $\text{C}_8\text{H}_{18}(g) + \frac{25}{2} \text{O}_2(g) \rightleftharpoons 8 \text{CO}_2(g) + 9 \text{H}_2\text{O}(l)$

$$_rH \quad 8 \quad (394) \quad 9 \quad (286) \quad (250) \\ - 5476 \text{ kJ/mol}$$

$$120.(c) \quad 1941 \quad 1273 \quad 3(241.8) \quad ({}_fH)B_2H_6 \\ ({}_fH)B_2H_6 \quad 1273 \quad 725.4 \quad 1941 \quad 57.4$$

$$121.(c) \quad C(s) \quad O_2(g) \quad CO_2(g); \\ {}_rH_1 \quad x \text{ kJ/mol} \quad \dots(1)$$

$$CO(g) \quad 1/2 O_2(g) \quad CO_2(g) \\ {}_rH_2 \quad y \text{ kJ/mol} \quad \dots(2)$$

Equation (1) (2)

$$C(s) \quad \frac{1}{2} O_2(g) \quad CO(g) \\ {}_fH \quad (y - x) \text{ kJ/mol}$$

$$122.(c) \quad C_2H_4(g) \quad H_2(g) \quad C_2H_6(g)$$

$${}_rH \quad x_2 \quad x_1 \\ 123.(b) \quad Q \quad \frac{5.8}{58} \quad \frac{5756}{2} \quad 287.8 \text{ kJ}$$

$$124.(a) \quad Q \quad \frac{1}{2} \quad 438 \quad \frac{22.3}{223} \quad 21.9$$

$$125.(a) \quad {}_rH \quad 57 \quad (393.5) \quad 52 \quad (285.8) \\ (70870)$$

$$\frac{33578.9 \text{ kJ/mol}}{884} \quad 37.98 \text{ kJ/mol}$$

$$126.(c) \quad {}_rH \quad 46 \quad 2 \quad 92$$

$$127.(c) \quad 2CH_3OH \quad 3O_2(g) \quad 4H_2O(l) \quad 2CO_2(g) \\ 1453$$

$$\frac{4 \quad 44 \quad 4H_2O(l) \quad 4H_2O(g)}{2CH_3OH \quad 3O_2(g) \quad 4H_2O(g) \quad 2CO_2(g)} \\ 1277$$

$$128.(b) \quad ({}_fH) \quad \frac{H}{2} \quad 1277 \quad 218.2 \text{ kJ/mol}$$

$$129.(b) \quad \text{For } H_2(g) \quad O_2(g) \quad H_2O_2(l) \\ {}_fH \quad (H_2O_2, l) \quad {}_rH_3 \quad \frac{{}_rH_2}{2} \quad \frac{{}_rH_1}{2}$$

$$130.(a) \quad C_2H_5OH(l) \quad 3O_2(g) \quad 2CO_2(g) \quad 3H_2O(l) \\ n_g \quad 2 \quad 3 \quad 1 \\ H \quad U \quad n_g RT \\ U \quad H \quad RT \quad RT \quad q$$

$$131.(a) \quad q_{\text{reaction}} \quad q_{\text{bomb}} \quad q_{\text{water}} \\ q_{\text{reaction}} \quad (C(\text{bomb}) \quad (m_{\text{water}} \quad c)) \quad T \\ (652 \quad 500 \quad 4.18) \quad 14.3 \\ 39210 \text{ J} \quad \text{or} \quad 39.21 \text{ kJ}$$

$$132.(b) \quad q \quad m \quad c \quad T, \quad m \quad q/(c \quad T) \\ (24 \quad 10^6 \quad 0.7)/(4.18 \quad 50) \\ 80383 \text{ g} \quad \text{or} \quad 80.383 \text{ kg}$$

$$133.(a) \quad m \text{ mole of acid} \quad 0.05 \quad 0.2 \quad 0.01 \\ {}_rH \quad \frac{480 \quad 1.09}{0.01 \quad 1000} \quad 52.32 \text{ kJ/mol}$$

$$134.(a) \quad H_{\text{solution}} \quad \frac{(125 \quad 8) \quad 4.2 \quad 6}{8} \quad 80 \\ 33516 \text{ J/mol} \quad \text{or} \quad 33.51 \text{ kJ/mol}$$

$$135.(b) \quad (I) \quad CaCl_2(s) \quad aq \quad CaCl_2(aq) \quad q \\ CaCl_2(aq) \quad 6H_2O(l) \quad CaCl_2 \quad 6H_2O(s) \\ 23.2 \\ CaCl_2(s) \quad 6H_2O \quad CaCl_2 \quad 6H_2O(s) \quad q \\ 23.2$$

$$3.5 \quad q \quad 23.2$$

$$q \quad 19.7 \\ (II) \quad 2Cl(g) \quad Cl_2(g)$$

$$H \quad \text{ve,} \quad S \quad \text{ve.}$$

$$136.(b) \quad HCl(g) \quad aq \quad H(aq.) \quad Cl(aq.); \\ {}_rH \quad 75.14 \\ 75.14 \quad {}_fH(H, aq) \quad {}_fH(Cl, g) \\ {}_fH(HCl, g) \\ \therefore {}_fH(H, aq) \quad 0 \\ {}_fH(Cl, aq) \quad 75.14 \quad 92.3 \\ 167.44 \text{ kJ/mol}$$

$$137.(a) \quad \text{Given that} \\ MgSO_4(s) \quad nH_2O \quad MgSO_4(nH_2O); \\ {}_rH_1 \quad 91.2 \text{ kJ/mol} \quad \dots(i) \\ MgSO_4 \quad 7H_2O(s) \quad (n-7)H_2O \\ MgSO_4(nH_2O) \\ {}_rH_2 \quad 13.8 \text{ kJ/mol} \quad \dots(ii)$$

Equation (i) (ii)

$$\text{or } H_{\text{hyd}} \quad {}_rH_1 \quad {}_rH_2 \\ 91.2 \text{ kJ/mol} \quad 13.8 \text{ kJ/mol} \\ 105 \text{ kJ/mol}$$

$$138.(a) \quad 12250x \quad 13000(1-x) \quad 12500 \\ 750x \quad 500 \quad x \quad 2/3 \quad \text{and} \quad y \quad 1/3 \\ \text{So,} \quad \frac{x}{y} \quad \frac{2}{1}$$

$$139.(c) \quad |\text{Enthalpy of change}| \quad 57 \\ 2H \quad \text{participate in reaction.}$$

$$140.(b) \quad H_{\text{neut.}} \\ H_{\text{ionization}} \quad {}_rH(H \quad OH \quad H_2O)$$

$$51.34 \times 55.84$$

So, $H_{\text{ionization}}$ or $\times 4.5 \text{ kJ/mol}$



142. (c) By calorimetry we can determine change in enthalpy.

143. (b) ${}_fH \frac{5}{8} \times 40 \frac{3}{8} \times 50 = 43.75 \text{ kJ/mol}$

144. (d) No. of moles of O_2 required to supplied 30 kJ heat to second reaction


$$\frac{30}{1260} \times \frac{3}{2} \times \frac{1}{28}$$

$$\text{So } n_{\text{O}_2} : n_{\text{H}_2} = \frac{1}{28} : 3 \text{ or } 1 : 84$$

145. (a) $(\text{O} - \text{F}) \frac{368}{2} \times 184$

146. (d) C—C bond enthalpy is not given.

147. (b) B.E. size



$$306 \text{ (314 - 3 - 121) [B.E. (P - Cl) 3];}$$

$$\text{B.E. (P—Cl)} = 123.66 \text{ kJ/mol}$$



$${}_fH \text{ (B.E.)}_R \text{ (B.E.)}_p$$

$$3(\text{C} - \text{H}) + (\text{C} - \text{O}) + (\text{O} - \text{H}) + (\text{H} - \text{Cl})$$

$$= 3(\text{C} - \text{H}) + (\text{C} - \text{Cl}) + 2(\text{O} - \text{H})$$

$$335 + 463 + 431 + 326 - 2 \times 463 - 23 \text{ kJ}$$



$${}_fH (\text{N}_2\text{H}_4, \text{g})$$

$$(941 + 2 \times 436) - (159 + 4 \times 398)$$

$$= 1813 - 1751 = 62 \text{ kJ mol}^{-1}$$

Level 2

1. (c) $w = P_{\text{ext}} (V_f - V_i)$

$$10^5 \times \frac{60 \times 10^{-3}}{0.60} - \frac{40 \times 10^{-3}}{1000} - \frac{100 \times 10^{-3}}{1000}$$

$$10^5 (100 \times 10^{-3} - 0.04 \times 10^{-3} - 0.1 \times 10^{-3})$$

$$= |w| = 9994 \text{ J}$$

2. (c) Mole of the gas in the first compartment

$$n_1 = \frac{P_1 V_1}{RT_1} = \frac{0.8314 \times 10^6 \times 2}{8.314 \times 400} = 500$$

Similarly, $n_2 = 2000$

The tank is rigid and insulated hence $w = 0$ and $q = 0$ therefore $U = 0$


Let T_f and P_f denote the final temperature and pressure respectively

$$U = n_1 C_{V,m} [T_f - T_1] + n_2 C_{V,m} [T_f - T_2]$$

$$0$$

$$500 (T_f - 400) + 2000 (T_f - 500) = 0$$

$$T_f = 480 \text{ K}$$



$$\downarrow U_2$$



$$C_{V,m} (\text{H}_2\text{O}, \text{g}) = 33.314 + 8.314 = 25 \text{ J/K mol}$$

$$U_2 = H_2 = n_g RT = 37.6$$

$$U_{\text{total}} = U_1 + U_2 + U_3$$

$$= C_{V,m}(\text{l}) T + V_{\text{vap.}} C_{V,m}(\text{g}) T$$

$$= \frac{75}{1000} \times 50 + 37.6 \times \frac{25}{1000} = 42.6 \text{ kJ/mol}$$

4. (a) $\therefore \frac{C_{p,m}}{C_{V,m}}$ and $C_{p,m} = C_{V,m} + R$

$$C_{V,m} = \frac{R}{1}$$

$$C_{V,m} = \frac{C_V}{n} \text{ and } C_V = m c_V$$

$$\frac{R}{1} = \frac{m c_V}{m} = M$$

$$c_V = \frac{R}{(1) M}$$

5. (d) Av. $C_{V,m} = \frac{n_1 C_{V,m_1} + n_2 C_{V,m_2}}{n_1 + n_2} = 2R$

for adiabatic process $dU = dW$

$$\frac{dT}{T} = \frac{R}{C_{V,m}} \frac{dV}{V}$$

$$n_1 C_{V,m_1} \frac{dT}{T} + n_2 C_{V,m_2} \frac{dT}{T} = (n_1 R + n_2 R) \frac{dV}{V}$$

$$\ln \frac{T_2}{T_1} = \frac{1}{2} \ln \frac{V_2}{V_1}$$

$$T_2 = 320 \times \frac{1}{4}^{1/2}$$

$$= 160 \text{ K}$$

$$U = (n_1 C_{V,m_1} + n_2 C_{V,m_2}) T = 960 R$$

$$6. (c) 2 C_{V,m}(T_2 - T_1) = 1 \frac{2RT_2}{1} - \frac{2RT_1}{10}$$

$$\frac{4}{3} \frac{C_{p,m}}{C_{V,m}}$$

$$\text{and } C_{p,m} - C_{V,m} = R$$

$$3R(T_2 - 300) = RT_2 \frac{C_{V,m}}{R} = 3R \frac{300}{10}$$

$$T_2 = 232.5 \text{ K}$$

$$w = U = 2 \cdot 3R(300 - 232.5)$$

$$w = 405 \text{ R}$$

7. (c) Max. temp. attained by gas in between B to C

According to equation of straight line

$$\frac{P}{1} = \frac{4}{4} \frac{V}{2} + \frac{1}{2}$$

$$P = 4 - 3V + 3$$

$$P = 7 - 3V$$

For 1 mole gas

$$\frac{RT}{V} = 7 - 3V; RT = 7V - 3V^2 \quad \dots(1)$$

$$R \frac{dT}{dV} = 7 - 6V = 0$$

$$V = \frac{7}{6} \text{ put in Eq. (1)}$$

$$RT = 7 - 3 \cdot \frac{7}{6} = \frac{7}{6}$$

$$T = \frac{49}{12R}$$

$$8. (i)(c) w = P \Delta V = nR \Delta T = 2 \cdot 8.314 \cdot 600$$

$$9.97 \text{ kJ}$$

$$(ii)(b) H_{\text{total}} = H_{AB} + H_{BC} = nC_{p,m} \Delta T = 0$$

$$2 \cdot \frac{7}{2} R (800 - 200) = 4200 R$$

$$9. (c) \frac{T_2}{T_1} = \frac{V_1}{V_2}$$

$$T_2 = T_1 \frac{1}{32} = 600 \cdot \frac{1}{32} = 18.75 \text{ K}$$

$$H_m = \frac{7}{2} R (150 - 600) = 1575 R$$

$$10. (a) w = nR \Delta T = 2 \cdot 8.314 \cdot 100$$

$$1662.8 \text{ J}$$

$$U = n C_{v,m} dT$$

$$2 \cdot (20 - 10^2 T) dT$$

$$2 \cdot 20 \cdot (T_2 - T_1) = 2 \cdot 10^2 \cdot \frac{(T_2^2 - T_1^2)}{2}$$

$$4700 \text{ J}$$

$$4700 \text{ J} - q = 1662.8 \text{ J}$$

$$q = 6362.8 \text{ J}$$

$$11. (d) C_{p,m} = (21.686 - 8.314) \cdot 10^{-3} \text{ J K}^{-1} \text{ mol}^{-1}$$

$$H = nC_{p,m} \Delta T$$

$$10 \cdot (30 - 10^3 T) dT$$

$$H = 10 \cdot 30 \cdot (T_2 - T_1) = \frac{10^3}{2} [T_2^2 - T_1^2] = 10$$

$$30350 \text{ J}$$

12. (b)

$$nC_{v,m} \frac{dU}{dT} = nC_{v,m} \frac{dq}{dT} = \frac{P}{n} \frac{dV}{dT}$$

$$C_m = C_{v,m} \frac{P}{n} \frac{dV}{dT}$$

...(1)

$$PV^n = K \text{ and } PV = nRT$$

$$KV^{1-n} = nRT$$

$$K(1-n)V^{-n} dV = nRdT$$

$$\frac{dV}{dT} = \frac{nR}{K(1-n)V^{-n}}$$

...(2)

from Eqs. (1) and (2)

$$C_m = C_{v,m} \frac{R}{(1-n)}$$

$$13. (b) P_2 V_2^2 = P_1 V_1^2$$

$$\frac{P_2}{P_1} = \left(\frac{V_1}{V_2} \right)^2$$

$$\text{Now, } \frac{T_2}{T_1} = \frac{P_2 V_2}{P_1 V_1} = \frac{V_1}{V_2} \cdot \frac{1}{3}$$

$$T_2 = \frac{300}{3} = 100 \text{ K}$$

$$H = nC_{p,m} \Delta T$$

$$2 \cdot \frac{5}{2} R (200 \text{ K}) = 1000 R$$

$$14. (d) S = nC_{p,m} \ln \frac{T_2}{T_1} = nR \ln \frac{P_1}{P_2}$$

$$3 \cdot \frac{7}{2} R \ln \frac{596}{298} = 3R \ln \frac{1}{4}$$

$$6.3 \text{ cal K}^{-1}$$

15. (a) For isentropic process $S_{\text{system}} = 0$

$$nC_{p,m} \ln \frac{T_2}{T_1} = nR \ln \frac{P_1}{P_2} = 0$$

$$\ln(P_2) = \frac{5}{2} \ln \frac{600}{300}$$

$$1.75 \text{ atm}$$

$$16. (b) S_{\text{system}} = nR \ln \frac{V_2}{V_1}$$

$$2 \cdot R \ln 2$$

$$S_{\text{surrounding}} = \frac{11.52 \text{ J/K} + \frac{3.41 \times 1000}{310}}{11 \text{ J/K}}$$

$$S_{\text{total}} = 11.52 + 11 + 0.52 \text{ J/K}$$

17. (c) 0.40 aT_1^3 bT_1
 0.40 a (1000) b 10
 0.4 1000a 10b ... (1)

0.92 aT_2^3 bT_2
 0.92 a 8000 20b ... (2)

from Eqs. (1) and (2)

$$a \cdot 2 \cdot 10^5, \quad b \cdot 0.038$$

$$S_m = \frac{aT^3 + bT}{T} dT$$

$$\frac{a[T_2^3 - T_1^3]}{3} + b[T_2 - T_1]$$

$$0.813 \text{ J/K-mol}$$

19. (d) No. of moles of sucrose $\frac{34.2}{342} \cdot 0.1$

(G)_{T,P} useful work done by the system

$$G_H T S$$

$$(6000 \quad 0.1) \frac{180 \cdot 0.1 \cdot 300}{1000}$$

$$605.4 \text{ kJ}$$

20. (a) G_{200} H_{200} T S_{200}
 H_{200} 20 4 16 kJ/mol

$$H_{T_2} \quad H_{T_1} \quad C_p [T_2 - T_1]$$

$$H_{400} \quad H_{200} \quad \frac{20 \cdot 200}{1000} \text{ kJ/mol}$$

$$16 \cdot 4 \cdot 20 \text{ kJ/mol}$$

21. (a) fS (NH₄Cl, s) at 300 K

$$S_{\text{NH}_4\text{Cl}(s)} = \frac{1}{2} S_{\text{N}_2} + 2 S_{\text{H}_2} + \frac{1}{2} S_{\text{Cl}_2}$$

$$374 \text{ JK}^{-1} \text{ mol}^{-1}$$

$$\therefore \quad rC_p = 0$$

$$fS_{310} \quad fS_{300}$$

$$374 \text{ JK}^{-1} \text{ mol}^{-1}$$

$$fH_{310} \quad fH_{300} \quad 314.5$$

$$fG_{310} \quad fH \quad 310 \quad S$$

$$314.5 \quad \frac{310 (374)}{1000}$$

$$198.56 \text{ kJ/mol}$$

22. (d) $rH \quad 3 \quad fH \text{ (CO}_2\text{, s)} \quad 4 \quad fH \text{ (CO}_2\text{, g)}$
 $fH \text{ (Co}_3\text{O}_4\text{, s)} \quad 4 \quad fH \text{ (CO, g)}$
 $rH \quad 241 \text{ kJ/mol}$
 $rS \quad 3 \quad 30 \quad 4 \quad 213.7 \quad 102.5$
 $4 \quad 197.7$

$$51.5 \text{ J/K-mol}$$

$$rG \quad rH \quad T \quad rS$$

$$241 \quad \frac{300}{1000} \quad 51.5$$

$$256.45 \text{ kJ/mol}$$

23. (c) At constant volume, $\frac{P_1}{T_1} = \frac{P_2}{T_2}$

$$P_2 = 1 \cdot \frac{300}{200} = \frac{3}{2}$$

$$\text{and } V_1 = 24.63 \text{ L}$$

for single phase

$$\therefore \quad dG = V dp = S dT$$

$$G_V P (2 \cdot 10^2 T) dT$$

$$1231.5 \quad 200 \quad \frac{10^2 \cdot 50,000}{2}$$

$$781.5 \text{ J}$$

24. (c) $\therefore rC_p = 0, \quad H_{300} \quad H_{310}$

25. (d) $H \quad 3 \quad H_1 \quad H_2 \quad 2 \quad H_3 \quad 3 \quad H_4$
 747.4 kJ

$$H \quad U \quad n_g RT; \text{ where } n_g = 8$$

$$747.4 \quad U \quad \frac{8 \cdot 8.314 \cdot 300}{1000}$$

$$U \quad 727.44 \text{ kJ}$$

26. (b) $rG \quad rH \quad T \quad rS$
 $rS \quad 2 \quad 81 \quad 4 \quad 24 \quad 3 \quad 205 \text{ J/mol}$

$$rH \quad 2258.1 \text{ kJ/mol}$$

$$rH \quad 2 \quad fH \text{ (Cr}_2\text{O}_3\text{, s)}$$

$$fH \text{ (Cr}_2\text{O}_3\text{, s)} \quad \frac{2258.1}{2}$$

$$1129.05 \text{ kJ/mol}$$

27. (a) CaO(s) CO₂(g) CaCO₃(s)

$$fH \quad H_f \text{ (CaCO}_3\text{)} \quad H_f \text{ (CaO)}$$

$$H_f \text{ (CO}_2\text{)}$$

$$1207 \quad (635) \quad (394)$$

$$178 \text{ kJ/mol}$$


$$U \quad H \quad n_g RT$$

$$U \quad 178 \quad \frac{(1) \cdot 8.3 \cdot 300}{1000}$$

$$175.51 \text{ kJ}$$

$$n_{\text{CaO}} = \frac{224}{56} = 4$$

$$q_v = n_r U = 4 \times (175.51) = 702.04 \text{ kJ}$$



$$n_g = \frac{3}{2}$$


$$U_c = - \frac{0.312 \times 8.75}{1} = 90$$

$$245.7 \text{ kJ/mol}$$

$$H = U + n_g RT$$

$$245.7 + \frac{3}{2} \times \frac{8.314 \times 300}{1000}$$

$$241.95 \text{ kJ/mol}$$



$$\begin{array}{ccc} 2\text{H} & 2\text{OH} & 2\text{H}_2\text{O} \\ rH & 55.84 & 2 \times 111.68 \\ 106.68 & \text{ion} H & 55.84 \times 2 \\ \text{ion} H & 5 \text{ kJ/mol} & \end{array}$$



$$\begin{array}{ccc} H_{\text{neutralization}} & H_{\text{ionization}} & rH \\ & (\text{H}^+ \text{OH}^- + \text{H}_2\text{O}) & \end{array}$$

$$55.95 \text{ kJ/mol} \quad H_{\text{ionization}} = 57.3$$

$$H_{\text{ionization}} \text{ for } 1 \text{ M HA} = 1.35 \text{ kJ/mol}$$

% heat utilized by 1 M acid for ionization

$$\frac{1.35}{1.4} \times 100 = 96.43\%$$

so, acid is 100 - 96.43 = 3.57% ionized



$$\begin{array}{ccc} 2 & 3 & \text{sub} \\ rH & & H(\text{C}, s) \\ 3 & \text{B.E.} & (\text{H}-\text{H}) \end{array}$$

$$\text{B.E.}(\text{C}-\text{C})$$

$$+ 6 \text{ B.E.}(\text{C}-\text{H})$$

$$85 \times 2 + 718 \times 3 = 436 \times (x + 6y)$$


$$x + 6y = 2829 \quad \dots(1)$$

Similarly, for $\text{C}_3\text{H}_8(g)$

$$2x + 8y = 4002 \quad \dots(2)$$

from Eqs. (1) and (2)

$$x = 345 \text{ kJ/mol}; y = 414 \text{ kJ/mol}$$




Let B.E. of N-N is x

$$46 \times \frac{x}{2} + \frac{3}{2} \times 436 = 3 \times 393 + x \times 958$$



$$rH = 50 \text{ kJ/mol}$$



$$\text{B.E.}(\text{N}-\text{N})$$

$$2 \text{ B.E.}(\text{H}-\text{H})$$

$$50 + (18 \times 4 + 393 \times y) = (958 \times 2 + 436)$$



$$50 + (1590 + y) = (1830)$$

$$\text{B.E.}(\text{N}-\text{N}) \text{ or } y = 190 \text{ kJ/mol}$$



$$292 \text{ [} 4x - 279 \text{] [} 38 - 85 \text{]}$$

$$x = 34 \text{ kcal/mol}$$



Enthalpy of hydrogenation of

$$\frac{357}{3} = 119 \text{ kJ/mol}$$



$$8 \text{ B.E.}(\text{C}-\text{H})$$

$$cH = 2 \text{ B.E.}(\text{C}-\text{C})$$


$$5 \text{ B.E.}(\text{O}-\text{O})$$

$$6 \text{ B.E.}(\text{C}-\text{O})$$

$$8 \text{ B.E.}(\text{O}-\text{H})$$

$$3 |\text{R.E.}| \text{ of } \text{CO}_2$$

$$4 \text{ vap } H(\text{H}_2\text{O})$$



AB process

$$0 \quad q_1 \quad w_1$$

$$q_1 \quad nR(2T_0) \ln 2$$

BC process $U_2 \quad q_2 \quad w_2$

$$\frac{nR}{(1)} (T_0 \quad 2T_0) \quad q_2 \quad \frac{nRT_0}{V_0} \quad (V_0 \quad 2V_0)$$

CA process


$$U_3 \quad q_3 \quad w_3$$

$$\frac{nR}{(1)} (2T_0 \quad T_0) \quad q_3 \quad 0$$

$$\text{Efficiency} = \frac{\text{Total work done}}{\text{Total heat absorbed}} = \frac{w_1 \quad w_2}{q_1 \quad q_3}$$

$$= \frac{(2RT_0 \ln 2) \quad (nRT_0)}{(2RT_0 \ln 2) \quad \frac{nRT_0}{1}}$$

37. (c) $V_1 \quad 24.63 \text{ L} \quad T_1 \quad 300 \text{ K}; T_2 \quad 600 \text{ K}$



For path (II)

$$T_1 V_1^{-1} \quad T_2 V_2^{-1} = \text{constant}$$

$$\frac{T_1}{T_2} = \frac{V_2}{V_1}$$

$$\frac{600}{300} = \frac{V_2}{V_1} \quad \frac{7}{5} = 1$$

$$(2)^{5/2} = \frac{V_2}{V_1}$$

$$V_2 = 4\sqrt{2} \quad 24.63 \quad 139.3 \text{ L} \quad 139 \text{ L}$$

38. (c)

$$\begin{array}{llll} {}_rH & {}_rH & {}_rH & \\ 56 & 285 & {}_rH & \end{array}$$

$$\begin{array}{llll} & (\text{H}_2\text{O}, l) & (\text{OH}, aq) & 229 \\ & & {}_rH & (\text{OH}, aq); \\ & & {}_rH & (\text{OH}, aq) \end{array} \quad 229$$

$$\begin{array}{llll} \text{HF}(aq) & \text{OH}(aq) & \text{F}(aq) & \text{H}_2\text{O}(l) \\ .H_{\text{neut}} & (320 \quad 285) & (329 \quad 229) & \\ & 47 \text{ kJ/mol} & & \end{array}$$

39. (b) $\text{I}_{2(s)} \quad \text{I}_{2(g)}$

$$\begin{array}{llll} H_2 & H_1 & C_p(T_2 \quad T_1) & \\ C_p & C_p(\text{I}_{2,g}) & C_p(\text{I}_{2,s}) & 6.2 \text{ cal/mole} \end{array}$$

$$H_2 \quad 6096 \quad 6.2(50) \quad 5786 \text{ cal/mole}$$

$$H \quad U \quad 1 \quad 2 \quad 523$$

$$5786 \quad U \quad 1 \quad 2 \quad 523$$

$$U \quad 5786 \quad 1046 \quad 4740 \text{ cal}$$

40. (d) $G \quad RT \ln K_p; K_p \quad (2x)^2 X \quad 4X^3$

$$G \quad RT \ln(4X^3)$$

$$G \quad RT \ln 4 \quad 3RT \ln X$$

Level 3

Passage-1

1. (d) $U \quad q \quad w$

$$U \quad 300 \quad (50 \quad 150) \quad 200$$

2. (d) $U \quad 300 \quad (50 \quad 150) \quad 500$

Passage-2

1. (b) $PV \quad nRT \quad 2 \quad 8 \quad 2 \quad 0.080 \quad T$

$$T \quad 100 \text{ K}$$

$$w_{\text{rev}} = 2.303 \quad n \quad R \quad T \log \frac{P_1}{P_2}$$

$$= 2.303 \quad 2 \quad 0.08 \quad 100 \quad \log \frac{1}{10}$$

$$= 36.848 \text{ bar-L}$$

2. (c) $w_{\text{irr}} \quad P_{\text{ext}} (V_2 \quad V_1)$

$$= 20 \quad \frac{nRT}{P_2} \quad \frac{nRT}{P_1}$$

$$= 144 \text{ bar-L}$$

3. (b) $w_{\text{irr (total)}} \quad w_1 \quad w_2$

$$= 10 \quad \frac{nRT}{10} \quad \frac{nRT}{2}$$

$$= 20 \quad \frac{nRT}{20} \quad \frac{nRT}{10}$$

$$= 5 \quad nRT \quad 80 \text{ bar-L}$$

4. (c) In isothermal compression is

$$w_{\text{One-stage}} \quad w_{\text{Two-stage}} \quad w_{\text{Reversible}}$$

Passage-3

1. (c) ${}_rS \quad 240 \quad 198 \quad 29 \quad 2 \quad 16$

2. (a) ${}_rH \quad 201 \quad 114 \quad 87$

$$320$$

3. (d) ${}_rS \quad 16 \quad {}_rC_p \quad \frac{dT}{T}$

$$300$$

$${}_rS \quad 16 \quad (44 \quad 29.4 \quad 2 \quad 28.8) \ln \frac{320}{300}$$

$${}_rS \quad 16 \quad (43) \ln \frac{320}{300} \quad 18.78$$

4. (c) ${}_rH \quad 87 \quad {}_rC_p \quad dT$

$$300$$

$$87 \text{ C } 4.31 \text{ 70 } 10^3$$

$$87.86 \text{ kJ/mol}$$

$$5. (d) \quad {}_rS \quad S_{\text{CH}_3\text{OH}} \quad S_{\text{CO}} \quad 2 S_{\text{H}_2} \quad - \text{J/K-mol}$$

$${}_rH \quad {}_fH \text{ (CH}_3\text{OH)} \quad {}_fH \text{ (CO)} \\ 2 \quad {}_fH \text{ (H}_2\text{)}$$

$$87 \text{ kJ/mol}$$

$${}_rS_{320} \quad {}_rS_{300} \quad {}_rC_p [T_2 \quad T_1]$$

$$\text{where } {}_rC_p \quad 44 \quad 29.4 \quad 2 \quad 28.8$$

$$43 \text{ J/K-mol}$$

$${}_rS_{320} \quad 16 \quad (43) \ln \frac{320}{320}$$

$$18.58$$

$${}_rH_{320} \quad {}_rH_{300} \quad {}_rC_p [T_2 \quad T_1]$$

$$87 \quad (43) \frac{20}{1000}$$

$$87.86 \text{ kJ/mol}$$

$${}_rG_{320} \quad {}_rH_{320} \quad T \quad {}_rS_{320} \\ 87.86 \quad \frac{320 \quad (18.58)}{1000}$$

$$81.91 \text{ kJ/mol}$$

Passage-4

$$1. (a) \quad 49.86 \quad (\quad {}_{\text{Ion}}H)_{\text{CH}_3\text{COOH}} \quad 55.84$$

$$(\quad {}_{\text{Ion}}H)_{\text{CH}_3\text{COOH}} \quad 5.98 \text{ kJ mol}^{-1}$$

$$2. (b) \quad {}_rH \quad 2 \quad (\quad 55.84) \quad 111.68 \text{ kJ}$$

$$3. (b) \quad 2\text{NaOH} \quad \text{H}_2\text{A} \quad \text{Na}_2\text{A} \quad 2\text{H}_2\text{O}$$

$$\frac{0.1 V_1}{1000} \quad \frac{0.05 V_2}{1000}$$

$$V_1 \quad V_2 \quad 100$$

$$2 \quad \frac{0.05 V_2}{1000} \quad \frac{0.1 V_1}{1000}$$

$$V_1 \quad V_2$$

$$V_1 \quad V_2 \quad 50 \text{ mL}$$

...(1)

Passage-5



$$1. (a) \quad (\quad G)_{PT} \quad 2000 \quad (20 \quad 298)$$

$$3960 \text{ J/mol}$$

$$2. (b) \quad \text{CaCO}_3 \quad \text{CaO} \quad \text{CO}_2 \quad H \quad \text{ve}$$

Reaction becomes spontaneous at high temperature because $T \Delta S$ dominates over H_{rxn} .

$$3. (a)$$



$$x \quad 3.5$$

$$x \quad 3.5 \quad 23.2$$

$$x \quad 19.7$$

$$H \quad x \quad 19.7$$

enthalpy driven

One or More Answers is/are Correct

$$1. (b, c)$$

Intensive properties : Temperature, pressure.

$$2. (a, b)$$

Extensive properties : Gibbs energy, Entropy.

$$3. (c) \text{ State function : Enthalpy}$$

$$4. (a, b, c)$$

Work done by system is negative.

Work done by surrounding is positive.

Heat absorbed by system is positive.

Heat absorbed by surrounding is negative.

$$5. (a, b, d)$$

$$U \quad n \quad C_{vm} dT \quad H \quad C_{pm} dT$$

$$U \quad q \quad w$$

$$6. (b, c, d) \quad U \quad 0 \quad U_2 \quad U_1 \quad 0$$

$$\text{Similarly, } H \quad 0$$

$$q \quad w \quad nRT \ln \frac{V_2}{V_1}$$

$$7. (b, c, d)$$

No heat transfer

$$U \quad q \quad w \quad 0 \quad w \quad w$$

$$U \quad w$$

$$T_2 \quad T_1 \quad \frac{w}{nC_{vm}}$$

$$w \quad \text{ve}$$

$$T_2 \quad T_1$$

$$8. (a, b, c)$$

$$U \quad q \quad w \quad 0 \quad w \quad w$$

$$w \quad n \quad C_{vm} dT$$

$$w \quad \frac{nR}{1} (T_2 \quad T_1)$$

$$w \quad nR P_{\text{ext}} \frac{T_2}{P_2} \frac{T_1}{P_1}$$

9. (c,d)


$$\text{At } V_A \quad \frac{1}{R} \quad \frac{100}{1} \quad 100 R$$

$$V_B \quad \frac{1}{R} \quad \frac{600}{3} \quad \mathbf{200 R}$$

$\therefore V_B > V_A$ so expansion of gas takes place
 $V_B = 200 \times 0.0821 = 16.42 \text{ L}$

10. (a,c,d)

First step is adiabatic ($q = 0$) so $U_1 = w_1$
 Second step is isochoric ($w = 0$)



So, $U_2 = q_2$

$\therefore$ initial and final temp. are same

$$U_{\text{total}} = U_1 = U_2 = 0$$

or $w_1 = q_2 = 0$

Max. work done by the gas, SO_2 is (area) under the curve

so, SO_2 absorbed

$\therefore$ $\text{SO}_2 > \text{CO}_2 > \text{O}_2 > \text{Ne}$
 so, max. decrease in temp. of Ne due to step 1.

11. (a,b,c)

$H = U + PV$ is correct relation.

12. (a, c)

$$H = U + n_g RT$$

(a) $n_g = 2$ $U = H$

(b) $n_g = \frac{1}{2}$ $U = H$

(c) $n_g = \frac{1}{2}$ $U = H$

(d) $n_g = 0$ $U = H$

13. (a, c, d)

$$G = H - TS$$

(a) H ve, S ve at low temperature
 G ve

(b) H ve, S ve; G ve
 at any temperature.

(c) H ve, S ve, G ve at any temperature

(d) H ve, S ve, G ve at high temperature


14. (b, d)

For spontaneous process

$$S_{\text{System}} + S_{\text{Surrounding}} = 0$$

$$(\Delta G_{\text{System}})_{T,P} = 0$$

15. (a, b, c)



$$x(l) \rightleftharpoons x(g) \quad T_b^0 = 400 \text{ K}$$

(a) At 400 K and 1 atm pressure

$$G = 0$$

(b) at 2 atm and 400 K, G ve

(c) at 400 K and 0.1 atm pressure, G ve

(d) at 410 K and 1 atm pressure, G ve

16. (a, c, d)

Entropy intermolecular force

$$H = 0$$

Spontaneous process $G = 0$

17. (b, c, d)

$$C(\text{graphite}) \rightleftharpoons C(\text{diamond})$$

$$G = H - TS$$

$$1.9 - 298(\text{ve}) = \text{ve}$$

Graphite is more stable than diamond thermodynamically

$$rH = (cH)_G - (cH)_D, (cH)_G < (cH)_D$$

$$1.9 - x < y - 1.9 - x$$

Diamond provides more heat on complete combustion in comparison of graphite.

18. (a, b, c, d)

(a) Reversible adiabatic process is iso-entropic process

(b) When $(\Delta G_{\text{System}})_{T,P} = 0$ then process may be endothermic / exothermic



(c) $dG = VdP - SdT$, only PV work consider

$$(d) S = \frac{9}{18} \frac{40.6 \times 1000}{373} = 54.42$$





19. (a, b, c)

$$(a) E = q = W = 0 = 0 = 0$$

$$(b) (\Delta F_H)(S, \text{rhombic}) = 0$$



20. (a, b, c)



21. (a, c, d)

Work is a form of energy exchange between system and surrounding in adiabatic process.

Intensive property is not additive.

In cyclic process work may be zero.

For cyclic process


$$dU = 0 \quad dq = dw \quad dq = PdV = 0$$

22. (a, c,)

Entropy of isolated system increases or remains constant but it can never decreases.

23. (a, c)

$$S_{\text{vap}} = \frac{H}{\text{B.P.}} = \frac{40}{400} = \frac{1000}{100}$$



24. (a, c, d)

(a) In isolated system no change possible

(b) $U \quad q \quad w \quad 0 \quad q \quad w$
w may be zero so q may be zero.

(c) Entropy of a closed system is not maximum at equilibrium.

(d) Intensive properties.

25. (a, c)

H	ve	Enthalpy driven
S	ve	Entropy driven
G	ve	Spontaneous process
G	ve	Non spontaneous process

Subjective Problems

1. For reversible adiabatic process


$$\begin{array}{ccc} P_1^{\gamma} & T_1 & P_2^{\gamma} & T_2 \\ & \frac{1}{T_1} & & \frac{1}{T_2} \\ \frac{T_1}{T_2} & \frac{P_2}{P_1} & ; \frac{300}{90} & \frac{10}{200} ; \\ \frac{10}{3} & \frac{1}{20} & ; \log \frac{10}{3} & \frac{1}{x} \log \frac{1}{20} \\ \frac{1}{1.301} & & 0.4 & \\ \frac{1}{0.6} & & 1.66 & \end{array}$$

Atomicity 1

$$\begin{array}{ccc} 2. & r = \frac{nRT \ln \frac{12}{24}}{P_{\text{ext}} nRT} = \frac{1}{P_2} \frac{1}{P_1} \\ & r = \frac{0.7}{24} \frac{1}{\frac{1}{24} \frac{1}{12}} = \frac{0.7}{24(1)} \\ 3. & C_m = \frac{R}{C_{VM}} = \frac{R}{1-x} = \frac{R}{1-3} \\ & C_m = \frac{3R}{2} = \frac{2R}{2} = R = \frac{R}{2} = 3R = 6 \\ 4. & \frac{w}{10} = 1 \frac{300}{500} = w = 4 \text{ kcal} \\ 5. & \frac{|w|}{8} \end{array}$$

...(i)

- $T_2 = 2 \times 10^1$ $T_1 = 16 \times 10^1$ $\frac{T_1}{T_2} = \frac{1}{2}$
 $\left| \frac{w}{8} \right| = 1 \times \frac{1}{2} \times \frac{1}{2}$ $|w| = \frac{1}{2} \times 8 \times 4$
6. $\frac{1}{2} A$ B $\frac{3}{2} C$
 G H T S
 $(10 \ 1000 \ 300 \ 10) \times 10^3$
 $G = 7 \text{ kJ/mol}$
7. Energy available $\frac{25}{100}$ 2880 $\frac{1}{2}$ 360
 Minimum distance $\frac{1}{90}$ 360 4 km
8. $C_2H_2(g)$ $H_2(g)$ $C_2H_4(g)$
 175 $(fH)_{C_2H_2}$ 50
 $(fH)_{C_2H_2}$ 225
 C_2H_2 $\frac{5}{2} O_2$ $2CO_2$ H_2O
 H 2 (390) 280 225 1285
 $\left| \frac{H}{257} \right|$ $\left| \frac{1285}{257} \right|$ 5
9. $5H_2O$ $CuSO_4(s)$ aq $CuSO_4(aq)$ 70
 $CuSO_4(aq)$ $CuSO_4$ $5H_2O(s)$ aq 10
 $CuSO_4(s)$ $5H_2O$ $CuSO_4$ $5H_2O$ 80
 $|H| = 80$
 For 0.1 mol : 80 0.1 mol : 80 0.1 8 kJ
10. 50 $(f_{ion}H)_{NH_4OH}$ 57
 $(f_{ion}H)_{NH_4OH}$ 7 kJ



- $sol.H$ 790 785 5
12. Q_v 1.25 4 5
 Q_v 5 $\frac{x}{80}$ 400
 $x = 1$
13. $\frac{T_2}{T_1}$ $\left| \frac{w_{Total}}{q_2} \right|$ $\frac{500}{500}$ $\frac{350}{500}$ $\left| \frac{w_{Total}}{10} \right|$
 Work done in one cycle 3
 Work done in two cycles 3 2 6
14. W $n_g RT$ $\frac{(3) \times 2 \times 500}{1000}$ 3
15. Process AC polytropic process ($P \propto V^K$)
 Molar Heat capacity c_m c_v $R/2$ $2R$
 Process AB Isobaric
 c_m c_p $5R/2$
 T_C $n C_m dT$
 $\frac{q_{AC}}{q_{AB}}$ $\frac{T_A}{T_B}$ $\frac{2R}{5R}$ 0.8 :
 $n C_{p,m} \cdot dT$ $\frac{2}{5} R$
 T_A
 $\frac{q_{AC}}{q_{AB}}$ 10 0.8 10 8

CHEMICAL EQUILIBRIUM

Types of Reactions

Reversible		Irreversible
(i)	A reaction in which not only the reactants react to form products but also the products react to form reactants under the same conditions is called reversible reaction.	A reaction which cannot take place in the reverse direction is called an irreversible reaction
(ii)	A few examples are given below $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$ $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$	A few examples are given below $\text{AgNO}_3(\text{aq}) + \text{NaCl}(\text{aq}) \rightarrow \text{AgCl}(\text{s}) + \text{NaNO}_3(\text{aq})$ $\text{BaCl}_2(\text{aq}) + \text{Na}_2\text{SO}_4(\text{aq}) \rightarrow \text{BaSO}_4(\text{s}) + 2\text{NaCl}(\text{aq})$

Characteristics of Equilibrium State

- ❖ A reaction (or a process) is said to be in equilibrium when the rate of forward reaction (process) becomes equal to the rate of backward reaction (process).
- ❖ At equilibrium, the concentration of each of the reactants and products become constant.
- ❖ Chemical equilibrium is dynamic in nature and not static *i.e.*, even after equilibrium is attained, the forward as well as the backward reaction take place but at equal speeds.
- ❖ A chemical equilibrium can be established only if none of the products is allowed to escape out.
- ❖ Chemical equilibrium can be attained from either direction.
- ❖ When reaction attains equilibrium at certain temperature and pressure, $\Delta_r G = 0$
- ❖ Catalyst increases rate of reaction in forward as well as backward direction upto same extent.

Law of Mass Action

It was put forward by Guldberg and Waage. It states that the rate at which a substance reacts is directly proportional to its active mass raised with suitable powers and hence the rate at which substances react together is directly proportional to the product of their active masses. Active mass means molar concentration.

Law of Chemical Equilibrium

For the reaction $aA(aq) + bB(aq) \rightleftharpoons xX(aq) + yY(aq)$

$$K_c = \frac{[X]^x [Y]^y}{[A]^a [B]^b}$$

Where K_c is called equilibrium constant at constant temperature.

Types of Equilibrium Constants

K_c Equilibrium constant in terms of molar concentration $\frac{[X]^x [Y]^y}{[A]^a [B]^b}$

K_p Equilibrium constant in terms of pressure $\frac{p_X^x \cdot p_Y^y}{p_A^a \cdot p_B^b}$

K_x Equilibrium constant in terms of mole fraction $\frac{(X_x)^x (X_y)^y}{(X_A)^a (X_B)^b}$

Unit of Equilibrium Constant

Unit of K_p (atm) ^{n_g} (where n_g = change in gas mole of reaction)

Unit of K_c M ^{n_g}

Unit of K_x Unitless

Relationship between K_p and K_c : $K_p = K_c (RT)^{n_g}$

where R = universal gas constant, T = temperature

Relationship between K_p and K_x : $K_p = K_x (P)^{n_g}$


where P = equilibrium pressure in container.

Types of Chemical Equilibria

- ❖ **Homogeneous equilibrium.** When in an equilibrium reaction, all the reactants and products are present in the same phase (*i.e.* gaseous or liquid), it is called homogeneous equilibrium.





- ❖ **Heterogeneous equilibrium.** When in an equilibrium reaction, the reactants and the products are present in two or more than two phases, it is called a heterogeneous equilibrium.



Expression of K for Equilibrium Reaction

- ❖ **Homogeneous equilibria:** If reactants and products are present in same phase.





Equilibrium constant expression for them are


$$(i) \quad K_c = \frac{[NH_4(aq)][OH^-(aq)]}{[NH_3(aq)]} \quad ; \quad [] \text{ represents concentration in mol/litre at equilibrium.}$$

$$(ii) \quad K_c = \frac{[O_3]^2}{[O_2]^3} ; K_p = \frac{P_{O_3}^2}{P_{O_2}^3} \quad (iii) \quad K_c = \frac{[CO_2]^3[H_2O]^4}{[C_3H_8][O_2]^5} ; K_p = \frac{P_{CO_2}^3 P_{H_2O}^4}{P_{C_3H_8} P_{O_2}^5}$$

Note: Equilibrium constant for gaseous homogeneous equilibrium can be expressed in two ways K_p and K_c . This means value of equilibrium constant depends upon choice of standard state in which concentration of reactants and products are expressed.

❖ Heterogeneous equilibria

If reactants and products are present in two or more phases, the equilibria describing them is called heterogeneous equilibrium.



Equilibrium expression for them can be written as

$$(i) \quad K_c = [Ag^+(aq)][I^-(aq)] \quad (ii) \quad K_p = \frac{1}{P_{CO_2}} ; K_c = \frac{1}{[CO_2(g)]}$$

$$(iii) \quad K_p = P_{Br_2(g)} ; K_c = [Br_2(g)]$$

Predicting the Extent of a Reaction

High value of equilibrium constant indicates that product(s) concentration is high and its low value indicates that concentration of the product(s) in equilibrium mixture is low.

Large values of K_p or K_c (larger than about 10^3), favour the products strongly. For intermediate values of K (approximately in the range of 10^{-3} to 10^3), the concentrations of reactants and products are comparable. Small values of equilibrium constant (smaller than 10^{-3}), favour the reactants strongly.

Predicting the Direction of the Reaction

For this purpose, we calculate the reaction quotient, Q . The reaction quotient is defined in the same way as the equilibrium constant (with molar concentrations to give Q_c , or with partial pressure to give Q_p) at any stage of reaction. For a general reaction



$$Q_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}$$

Then, if $Q_c < K_c$, the reaction will proceed towards direction of reactants.

If $Q_c = K_c$, the reaction will move towards direction of the products.

If $Q_c > K_c$, the reaction mixture is at equilibrium

Important Point about Equilibrium Constant

- ❖ If the reaction is reversed, the value of equilibrium constant is inversed e.g.,

$\text{CH}_3\text{COOH}(aq) + \text{C}_2\text{H}_5\text{OH}(aq) \rightleftharpoons \text{CH}_3\text{COOC}_2\text{H}_5(aq) + \text{H}_2\text{O}(l)$, Equilibrium constant K
 then for $\text{CH}_3\text{COOC}_2\text{H}_5(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{CH}_3\text{COOH}(aq) + \text{C}_2\text{H}_5\text{OH}(aq)$, Equilibrium constant $1/K$

- ❖ If the reaction is divided by 2, equilibrium constant is the square root of the original e.g.,

for $\text{N}_2(g) + 3\text{H}_2(g) \rightleftharpoons 2\text{NH}_3(g)$, Equilibrium constant K

then for $\frac{1}{2}\text{N}_2(g) + \frac{3}{2}\text{H}_2(g) \rightleftharpoons \text{NH}_3(g)$, Equilibrium constant $\sqrt{K}$

If the reaction is multiplied by 2, equilibrium constant is the square of the original.

- ❖ If the reaction is written in two steps, equilibrium constant is equal to the product of the equilibrium constants of the step reactions e.g., if for

$\text{N}_2(g) + 2\text{O}_2(g) \rightleftharpoons 2\text{NO}_2(g)$, Equilibrium constant K

and for the same reaction taking place in steps i.e.,

$\text{N}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{NO}(g)$, Equilibrium constant K_1

and $\text{NO}(g) + \text{O}_2(g) \rightleftharpoons 2\text{NO}_2(g)$, Equilibrium constant K_2

then $K = K_1 \times K_2$

- ❖ Effect of temperature: According to van't Hoff equation, $\log \frac{K_2}{K_1} = \frac{H}{2.303R} \left(\frac{T_2}{T_1 T_2} - \frac{T_1}{T_1 T_2} \right)$

where K_1 and K_2 are the equilibrium constants at temperatures T_1 and T_2 respectively and H is the molar enthalpy change in the temperature range T_1 to T_2 .

For exothermic reaction, as temperature increases K decreases.

For endothermic reaction, as temperature increases K increases.

Relationship Between Degree of Dissociation (α) and Vapour Density

- ❖ For dissociation reaction: $A_n(g) \rightleftharpoons nA(g)$

$$\text{Degree of dissociation } (\alpha) = \frac{\text{No. of moles dissociated}}{\text{Total no. of moles taken}}$$

$$= \frac{D - d}{d(n-1)} = \frac{M_t - M_0}{M_0(n-1)}$$

where M_t theoretical (calculated) molecular mass
 M_0 observed (experimental) molecular mass
 D theoretical vapour density, d observed vapour density.
 n number of moles of product formed from 1 mole reactant

$$\text{Density of gas mixture} = \frac{PM}{RT}$$

- ❖ For polymerization reaction

$nA(g) \rightleftharpoons A_n(g)$, where $n \geq 2$

$$\alpha = \frac{D - d}{d \left(\frac{1}{n} - 1 \right)} = \frac{M_t - M_0}{M_0 \left(\frac{1}{n} - 1 \right)}$$

Le Chatelier's Principle

"If a system in equilibrium is subjected to a change of concentration, temperature or pressure, the equilibrium shifts in a direction so as to undo the effect of the change imposed."

❖ Effect of change in concentration on equilibrium

As we add or remove reactant (or product) the ratio of new concentrations become 'Q' (reaction quotient)

$Q < K$: equilibrium will shift in forward direction.

$Q > K$: equilibrium will shift in backward direction.

❖ Effect of change in pressure

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, the volume decreases proportionately. The equilibrium will shift in the direction in which there is decrease in number of moles *i.e.*, towards the direction in which there is decrease in volume.

❖ Effect of change in pressure on melting point: There are two types of solids:

(i) Solids whose volume decreases on melting, *e.g.*, ice, diamond, carborundum, magnesium nitride and quartz.

Solid (higher volume) $\rightleftharpoons$ Liquid (lower volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

(ii) Solid whose volume increase on melting, *e.g.*, Fe, Cu, Ag, Au, etc.

Solid (lower volume) $\rightleftharpoons$ Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.

❖ Solubility of substances : When solid substances are dissolved in water, either heat is evolved (exothermic) or heat is absorbed (endothermic).

(i) For endothermic solubility process, solubility increase with increase in temperature.

(ii) For exothermic solubility process, decrease with increase in temperature.

❖ Solubility of gases in liquids : When a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.

❖ Effect of temperature: For endothermic reaction as temperature increases reaction shift in forward direction. For exothermic reaction as temperature increases reaction shift in backward direction.

❖ Addition of inert gas

(i) For reactions in which $n_p = n_r$, there is no effect of adding an inert gas at constant volume or at constant pressure on the equilibrium.

(ii) For reactions in which $n_p \neq n_r$, there is no effect of adding inert gas on the equilibrium at constant volume but at constant pressure, equilibrium shifts towards larger mole side.

Applying Le Chatelier's principle, the favourable conditions for the formation of NH_3 by Haber's process




are (i) High temperature (ii) Low pressure (iii) Removal of N_2 and H_2 (iv) Addition of inert gas at constant pressure.

❖ Physical Equilibrium

Only physical changes are involved.

Example : (i) Solid liquid equilibria



At equilibrium: Net rate of conversion of ice into water = Net rate of conversion of water into ice.

For $\text{H}_2\text{O}(l)$ normal freezing point is 273 K.

(ii) Liquid vapour equilibria :



Equilibrium is characterized by constant value of vapour pressure of $\text{H}_2\text{O}(l)$ at certain temperature.

For $\text{H}_2\text{O}(l)$ normal boiling point is 373 K.







Level 1

- A reversible reaction is one which :
 - proceeds in one direction
 - proceeds in both directions
 - proceeds spontaneously
 - all the statements are wrong
- The equilibrium constant K_c for the reaction

$$\text{P}_4(\text{g}) \rightleftharpoons 2\text{P}_2(\text{g})$$
 is 1.4 at 400°C . Suppose that 3 moles of $\text{P}_4(\text{g})$ and 2 moles of $\text{P}_2(\text{g})$ are mixed in 2 litre container at 400°C . What is the value of reaction quotient (Q_c)?
 - $\frac{3}{2}$
 - $\frac{2}{3}$
 - 1
 - None of these
- In a chemical reaction equilibrium is established when :
 - opposing reaction ceases
 - concentrations of reactants and product are equal
 - velocity of opposing reaction is the same as that of forward reaction
 - reaction ceases to generate heat
- The equilibrium constant for a reaction is K , and the reaction quotient is Q . For a particular reaction mixture, the ratio $\frac{K}{Q}$ is 0.33. This means that :
 - the reaction mixture will equilibrate to form more reactant species
 - the reaction mixture will equilibrate to form more product species
 - the equilibrium ratio of reactant to product concentrations will be 3
 - the equilibrium ratio of reactant to product concentrations will be 0.33
- Consider the reaction $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$ for which $K_c = 278 \text{ M}^{-1}$. 0.001 mole of each of the reagents $\text{SO}_2(\text{g})$, $\text{O}_2(\text{g})$ and $\text{SO}_3(\text{g})$ are mixed in a 1.0 L flask. Determine the reaction quotient of the system and the spontaneous direction of the system :
 - $Q_c = 1000$; the equilibrium shifts to the right
 - $Q_c = 1000$; the equilibrium shifts to the left
 - $Q_c = 0.001$; the equilibrium shifts to the left
 - $Q_c = 0.001$; the equilibrium shifts to the right
- In Q. No. 5, if the mixture of gases was allowed to come to equilibrium. The volume of the reaction vessel was then rapidly increased by a factor of two. As a result of the change the reaction quotient (Q_c) would :
 - increase because of the pressure decrease
 - decrease because of the pressure decrease
 - remain the same because the equilibrium constant is independent of volume
 - increase because the reaction is endothermic
- For the reaction $\text{A}(\text{g}) + 3\text{B}(\text{g}) \rightleftharpoons 2\text{C}(\text{g})$ at 27°C , 2 moles of A, 4 moles of B and 6 moles of C are present in 2 litre vessel. If K_c for the reaction is 1.2, the reaction will proceed in :
 - forward direction
 - backward direction
 - neither direction
 - none of these


8. For a reversible gaseous reaction $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ at equilibrium, if some moles of H_2 are replaced by same number of moles of T_2 (T is tritium, isotope of H and assume isotopes do not have different chemical properties) without affecting other parameters, then :
- the sample of ammonia obtained after sometime will be radioactive.
 - moles of N_2 after the change will be different as compared to moles of N_2 present before the change
 - the value of K_p or K_c will change
 - the average molecular mass of new equilibrium will be same as that of old equilibrium
9. For the synthesis of ammonia by the reaction $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$ in the Haber's process, the attainment of equilibrium is correctly predicted by the curve




10. The figure shows the change in concentration of species A and B as a function of time.

The equilibrium constant K_c for the reaction $\text{A}(\text{g}) \rightleftharpoons 2\text{B}(\text{g})$ is:

- $K_c = 1$
- $K = 1$
- $K = 1$
- data insufficient



11. Attainment of the equilibrium $\text{A}(\text{g}) \rightleftharpoons 2\text{C}(\text{g}) + \text{B}(\text{g})$ gave the following graph. Find the correct option. (% dissociation = fraction dissociated $\times 100$)



- (a) At $t = 5$ sec equilibrium has been reached and $K_c = 40$ (mol/litre)²
 (b) At $t = 5$ sec equilibrium has been reached and % dissociation of A is 20%
 (c) At $t = 5$ sec equilibrium has been reached and % dissociation of A is 30%
 (d) None of these

12. Using molar concentrations, what is the unit of K_c for the reaction?



- (a) M^{-2} (b) M^2 (c) M^{-1} (d) M

13. What is the unit of K_p for the reaction?




- (a) atm (b) atm² (c) atm² (d) atm⁻¹


14. What is the equilibrium expression for the reaction $\text{P}_4(s) + 5\text{O}_2(g) \rightleftharpoons \text{P}_4\text{O}_{10}(s)$

- (a) $K_c [\text{O}_2]^5$ (b) $K_c [\text{P}_4\text{O}_{10}] / 5[\text{P}_4][\text{O}_2]$
 (c) $K_c [\text{P}_4\text{O}_{10}] / [\text{P}_4][\text{O}_2]^5$ (d) $K_c = 1 / [\text{O}_2]^5$

15. At 527°C, the reaction given below has $K_c = 4$



What is the K_p for the reaction?




- (a) $16 (800 R)^2$ (b) $\frac{800R}{4}$ (c) $\frac{1}{4 \cdot 800R}$ (d) None of these

16. The equilibrium constant for the reaction $\text{N}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{NO}(g)$ at temperature (T) is $4 \cdot 10^{-4}$. The value of K_c for the reaction $\text{NO}(g) \rightleftharpoons \frac{1}{2}\text{N}_2(g) + \frac{1}{2}\text{O}_2(g)$ at the same temperature is :


- (a) $4 \cdot 10^{-4}$ (b) 50 (c) $2.5 \cdot 10^2$ (d) 0.02

17. The equilibrium constant K_c for the following reaction at 842°C is $7.90 \cdot 10^{-3}$. What is K_p at same temperature?



- (a) $8.64 \cdot 10^{-5}$ (b) $8.26 \cdot 10^{-4}$ (c) $7.90 \cdot 10^{-2}$ (d) $7.56 \cdot 10^{-2}$

18. The equilibrium constant K_p for the following reaction at 191°C is 1.24. What is K_c ?




- (a) 6.7 (b) 0.61 (c) 8.30 (d) 7.6

19. For the equilibrium $\text{SO}_2\text{Cl}_2(g) \rightleftharpoons \text{SO}_2(g) + \text{Cl}_2(g)$, what is the temperature at which $\frac{K_p (\text{atm})}{K_c (\text{M})} = 3$?

- (a) 0.027 K (b) 0.36 K (c) 36.54 K (d) 273 K

- 20.** For the reversible reaction,

$$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$$
 at 500°C , the value of K_p is 1.44×10^{-5} when partial pressure is measured in atmospheres. The corresponding value of K_c with concentration in mole litre⁻¹, is:
 (a) $1.44 \times 10^{-5}/(0.082 \times 500)^2$ (b) $1.44 \times 10^{-5}/(8.314 \times 773)^2$
 (c) $1.44 \times 10^{-5}/(0.082 \times 773)^2$ (d) $1.44 \times 10^{-5}/(0.082 \times 773)^2$
- 21.** For the reaction $\text{CO}(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons \text{COCl}_2(\text{g})$ the value of $\frac{K_c}{K_p}$ is equal to :
 (a) $\sqrt{RT}$ (b) RT (c) $\frac{1}{RT}$ (d) 1.0
- 22.** The concentration of a pure solid or liquid phase is not included in the expression of equilibrium constant because :
 (a) density of solid and liquid are independent of their quantities.
 (b) solids and liquids react slowly.
 (c) solids and liquids at equilibrium do not interact with gaseous phase.
 (d) the molecules of solids and liquids cannot migrate to the gaseous phase.
- 23.** A catalyst is a substance which :
 (a) increases the equilibrium concentration of the product.
 (b) changes the equilibrium constant of the reaction.
 (c) shortens the time to reach equilibrium.
 (d) supplies energy to the reaction.
- 24.** What will be the effect on the equilibrium constant on increasing temperature, if the reaction neither absorbs heat nor releases heat?
 (a) Equilibrium constant will remain constant.
 (b) Equilibrium constant will decrease.
 (c) Equilibrium constant will increase.
 (d) Can not be predicted.
- 25.** The equilibrium constant for the reaction $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$ is 4×10^{-4} at 200 K. In presence of a catalyst, equilibrium is attained ten times faster. Therefore, the equilibrium constant in presence of the catalyst at 200 K is :
 (a) 40×10^{-4} (b) 4×10^{-4}
 (c) 4×10^{-3} (d) difficult to compute without more data
- 26.** For the reaction $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$, the equilibrium constant changes with :
 (a) total pressure (b) catalyst
 (c) concentration of H_2 and I_2 (d) temperature
- 27.** Consider the reactions
 (i) $2\text{CO}(\text{g}) + 2\text{H}_2\text{O}(\text{g}) \rightleftharpoons 2\text{CO}_2(\text{g}) + 2\text{H}_2(\text{g})$; Eqm. Constant K_1
 (ii) $\text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + 3\text{H}_2(\text{g})$; Eqm. Constant K_2



Which of the following relation is correct?

- (a) $K_3 = \frac{K_1}{K_2}$ (b) $K_3 = \frac{K_1^2}{K_2^2}$ (c) $K_3 = K_1 K_2$ (d) $K_3 = \sqrt{K_1} K_2$


28. For the reaction $2\text{NO}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightleftharpoons \text{N}_2\text{O}_5(\text{g})$, if the equilibrium constant is K_p , then the equilibrium constant for the reaction $2\text{N}_2\text{O}_5(\text{g}) \rightleftharpoons 4\text{NO}_2(\text{g}) + \text{O}_2(\text{g})$ would be :

- (a) K_p^2 (b) $\frac{2}{K_p}$ (c) $\frac{1}{K_p^2}$ (d) $\frac{1}{\sqrt{K_p}}$



29. The equilibrium constant (K_c) for the reaction




is 4×10^{-34} at 25°C . What is the equilibrium constant for the reaction?






- (a) 2×10^{-17} (b) 2.5×10^{33} (c) 5×10^6 (d) None of these
30. At a certain temperature, the following reactions have the equilibrium constants as shown below:



What is the equilibrium constant K_c for the reaction at the same temperature ?



- (a) 2.5×10^{76} (b) 4×10^{23} (c) 4×10^{77} (d) None of these
31. Consider the following gaseous equilibria given below :



The equilibrium constant for the reaction, $2\text{NH}_3 + \frac{5}{2}\text{O}_2 \rightleftharpoons 2\text{NO} + 3\text{H}_2\text{O}$ in terms of K_1 , K_2 and K_3 will be :

- (a) $K_1 K_2 K_3$ (b) $\frac{K_1 K_2}{K_3}$ (c) $\frac{K_1 K_3^2}{K_2}$ (d) $\frac{K_2 K_3^3}{K_1}$

32. In the reaction $\text{X}(\text{g}) + \text{Y}(\text{g}) \rightleftharpoons 2\text{Z}(\text{g})$, 2 mole of X, 1 mole of Y and 1 mole of Z are placed in a 10 litre vessel and allowed to reach equilibrium. If final concentration of Z is 0.2 M , then K_c for the given reaction is :

- (a) 1.60 (b) $\frac{80}{3}$ (c) $\frac{16}{3}$ (d) None of these

33. An equilibrium mixture of the reaction $2\text{H}_2\text{S}(\text{g}) \rightleftharpoons 2\text{H}_2(\text{g}) + \text{S}_2(\text{g})$ had 0.5 mole H_2S , 0.10 mole H_2 and 0.4 mole S_2 in one litre vessel. The value of equilibrium constant (K) in mol litre^{-1} is :
 (a) 0.004 (b) 0.008 (c) 0.016 (d) 0.160
34. Given $[\text{CS}_2] = 0.120 \text{ M}$, $[\text{H}_2] = 0.10 \text{ M}$, $[\text{H}_2\text{S}] = 0.20 \text{ M}$ and $[\text{CH}_4] = 8.40 \times 10^{-5} \text{ M}$ for the following reaction at 900°C , at eq. Calculate the equilibrium constant (K_c).

$$\text{CS}_2(\text{g}) + 4\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_4(\text{g}) + 2\text{H}_2\text{S}(\text{g})$$

 (a) 0.0120 (b) 0.0980 (c) 0.280 (d) 0.120
35. The equilibrium constant for the following reaction is 10.5 at 500 K. A system at equilibrium has $[\text{CO}] = 0.250 \text{ M}$ and $[\text{H}_2] = 0.120 \text{ M}$. What is the $[\text{CH}_3\text{OH}]$?

$$\text{CO}(\text{g}) + 2\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_3\text{OH}(\text{g})$$

 (a) 0.03618 (b) 0.0435 (c) 0.546 (d) 0.0499
36. When sulphur (in the form of S_8) is heated at temperature T , at equilibrium, the pressure of S_8 falls by 30% from 1.0 atm, because $\text{S}_8(\text{g})$ is partially converted into $\text{S}_2(\text{g})$. Find the value of K_p for this reaction.
 (a) 2.96 (b) 6.14 (c) 204.8 (d) None of these
37. 9.2 grams of $\text{N}_2\text{O}_4(\text{g})$ is taken in a closed one litre vessel and heated till the following equilibrium is reached $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$
 At equilibrium, 50% $\text{N}_2\text{O}_4(\text{g})$ is dissociated. What is the equilibrium constant (in mol litre^{-1}) (molecular mass of $\text{N}_2\text{O}_4 = 92$)
 (a) 0.1 (b) 0.4 (c) 0.2 (d) 2
38. Two moles of NH_3 when put into a previously evacuated vessel (one litre), partially dissociated into N_2 and H_2 . If at equilibrium one mole of NH_3 is present, the equilibrium constant is :
 (a) $3/4 \text{ mol}^2\text{litre}^{-2}$ (b) $27/64 \text{ mol}^2\text{litre}^{-2}$ (c) $27/32 \text{ mol}^2\text{litre}^{-2}$ (d) $27/16 \text{ mol}^2\text{litre}^{-2}$
39. In the presence of excess of anhydrous SrCl_2 , the amount of water taken up is governed by $K_p = 10^{12} \text{ atm}^{-4}$ for the following reaction at 273 K

$$\text{SrCl}_2 \cdot 2\text{H}_2\text{O}(\text{s}) \rightleftharpoons \text{SrCl}_2(\text{s}) + 2\text{H}_2\text{O}(\text{g})$$

 What is equilibrium vapour pressure (in torr) of water in a closed vessel that contains $\text{SrCl}_2 \cdot 2\text{H}_2\text{O}(\text{s})$?
 (a) 0.001 torr (b) 10^3 torr (c) 0.76 torr (d) 1.31 torr
40. $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}(\text{s}) \rightleftharpoons \text{CuSO}_4(\text{s}) + 5\text{H}_2\text{O}(\text{g})$; $K_p = 4 \times 10^{-4} \text{ atm}^2$. If the vapour pressure of water is 38 torr then percentage of relative humidity is : (Assume all data at constant temperature)
 (a) 4 (b) 10 (c) 40 (d) None of these
41. $\text{NH}_4\text{HS}(\text{s}) \rightleftharpoons \text{NH}_3(\text{g}) + \text{H}_2\text{S}(\text{g})$
 The equilibrium pressure at 25°C is 0.660 atm. What is K_p for the reaction?
 (a) 0.109 (b) 0.218 (c) 1.89 (d) 2.18

42. For the reaction $2A(g) \rightleftharpoons B(g) + 3C(g)$, at a given temperature, $K_c = 16$. What must be the volume of the flask, if a mixture of 2 mole each of A, B and C exist in equilibrium?
 (a) $\frac{1}{4}$ (b) $\frac{1}{2}$ (c) 1 (d) None of these
43. One mole of pure ethyl alcohol was treated with one mole of pure acetic acid at 25°C . One-third of the acid changes into ester at equilibrium. The equilibrium constant for the reaction will be :
 (a) $\frac{1}{4}$ (b) 2 (c) 3 (d) 4
44. $I_2(aq) + I^-(aq) \rightleftharpoons I_3(aq)$. We started with 1 mole of I_2 and 0.5 mole of I^- in one litre flask. After equilibrium is reached, excess of $AgNO_3$ gave 0.25 mole of yellow precipitate. Equilibrium constant is :
 (a) 1.33 (b) 2.66 (c) 2.0 (d) 3.0
45. At 87°C , the following equilibrium is established.

$$H_2(g) + S(s) \rightleftharpoons H_2S(g); K_c = 0.08$$
 If 0.3 mole hydrogen and 2 mole sulphur are heated to 87°C in a 2 L vessel, what will be the concentration of H_2S at equilibrium?
 (a) 0.011 M (b) 0.022 M (c) 0.044 M (d) 0.08 M
46. In the equilibrium $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$, the partial pressure of SO_2 , O_2 and SO_3 are 0.662, 0.10 and 0.331 atm respectively. What should be the partial pressure of oxygen so that the equilibrium concentrations of SO_2 and SO_3 are equal?
 (a) 0.4 atm (b) 1.0 atm (c) 0.8 atm (d) 0.25 atm
47. When heated, ammonium carbamate decomposes as follows :


$$NH_4COONH_2(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$$
 At a certain temperature, the equilibrium pressure of the system is 0.318 atm. K_p for the reaction is :
 (a) 0.128 (b) 0.426 (c) 4.76×10^{-3} (d) None of these
48. In a system $A(s) \rightleftharpoons 2B(g) + 3C(g)$, if the concentration of C at equilibrium is increased by a factor of 2, it will cause the equilibrium concentration of B to change to :
 (a) two times the original value (b) one half of its original value
 (c) $2\sqrt{2}$ times to the original value (d) $\frac{1}{2\sqrt{2}}$ times the original value
49. For $A + B \rightleftharpoons C + D$, if initially the concentration of A and B are both equal but at equilibrium concentration of D will be twice of that of A then what will be the equilibrium constant of reaction ?
 (a) $\frac{4}{9}$ (b) $\frac{9}{4}$ (c) $\frac{1}{9}$ (d) 4
50. The equilibrium constant K_c for the reaction $SO_2(g) + NO_2(g) \rightleftharpoons SO_3(g) + NO(g)$ is 16. If 1 mole of each of all the four gases is taken in 1 dm^3 vessel, the equilibrium concentration of NO would be :
 (a) 0.4 M (b) 0.6 M (c) 1.4 M (d) 1.6 M

51. On increasing the temperature, the rate of a reaction :
- (a) always increases
 - (b) always decreases
 - (c) first increases and then decreases
 - (d) may increase or decrease depending upon the nature of the reaction
52. A catalyst increases the rate of a reaction by :
- (a) increasing the activation energy of the reaction
 - (b) increasing the value of rate constants (k_f and k_b)
 - (c) increasing the enthalpy change of the reaction
 - (d) decreasing the enthalpy change of the reaction
53. At a certain temperature, only 50% HI is dissociated at equilibrium in the following reaction :



The equilibrium constant for this reaction is :


- (a) 0.25
 - (b) 1.0
 - (c) 3.0
 - (d) 0.5
54. The equilibrium constant K_p for the reaction
- $$\text{H}_2(\text{g}) + \text{CO}_2(\text{g}) \rightleftharpoons \text{H}_2\text{O}(\text{g}) + \text{CO}(\text{g})$$
- is 4.0 at 1660°C. Initially 0.80 mole H_2 and 0.80 mole CO_2 are injected into a 5.0 litre flask. What is the equilibrium concentration of $\text{CO}_2(\text{g})$?
- (a) 0.533 M
 - (b) 0.0534 M
 - (c) 0.535 M
 - (d) None of these
55. At 273 K and 1 atm, 10 litre of N_2O_4 decomposes to NO_2 according to equation



What is degree of dissociation (α) when the original volume is 25% less than that of existing volume?

- (a) 0.25
 - (b) 0.33
 - (c) 0.66
 - (d) 0.5
56. The equilibrium constant for the reaction $\text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}_2(\text{g}) + \text{H}_2(\text{g})$ is 5. How many moles of CO_2 must be added to 1 litre container already containing 3 moles each of CO and H_2O to make 2M equilibrium concentration of CO?
- (a) 15
 - (b) 19
 - (c) 5
 - (d) 20
57. A nitrogen-hydrogen mixture initially in the molar ratio of 1 : 3 reached equilibrium to form ammonia when 25% of the N_2 and H_2 had reacted. If the total pressure of the system was 21 atm, the partial pressure of ammonia at the equilibrium was :
- (a) 4.5 atm
 - (b) 3.0 atm
 - (c) 2.0 atm
 - (d) 1.5 atm
58. Ammonia under a pressure of 15 atm at 27°C is heated to 347°C in a closed vessel in the presence of a catalyst. Under the conditions, NH_3 is partially decomposed according to the equation,
- $$2\text{NH}_3 \rightleftharpoons \text{N}_2 + 3\text{H}_2$$
- the vessel is such that the volume remains effectively constant where as pressure increases to 50 atm. Calculate the percentage of NH_3 actually decomposed :
- (a) 65%
 - (b) 61.3%
 - (c) 62.5%
 - (d) 64%

59. 0.1 mole of $\text{N}_2\text{O}_4(\text{g})$ was sealed in a tube under one atmospheric conditions at 25°C . Calculate the number of moles of $\text{NO}_2(\text{g})$ present, if the equilibrium $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ ($K_p = 0.14$) is reached after some time :
- (a) 1.8×10^{-2} (b) 2.8×10^{-2} (c) 0.034 (d) 2.8×10^{-2}
60. 5 moles of SO_2 and 5 moles of O_2 are allowed to react. At equilibrium, it was found that 60% of SO_2 is used up. If the pressure of the equilibrium mixture is one atmosphere, the partial pressure of O_2 is:
- (a) 0.52 atm (b) 0.21 atm (c) 0.41 atm (d) 0.82 atm
61. $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$
For the reaction initially the mole ratio was 1 : 3 of $\text{N}_2 : \text{H}_2$. At equilibrium 50% of each has reacted. If the equilibrium pressure is P , the partial pressure of NH_3 at equilibrium is :
- (a) $\frac{P}{3}$ (b) $\frac{P}{4}$ (c) $\frac{P}{6}$ (d) $\frac{P}{8}$
62. 2.0 mole of PCl_5 were introduced in a vessel of 5.0 L capacity of a particular temperature. At equilibrium, PCl_5 was found to be 35% dissociated into PCl_3 and Cl_2 . The value of K_c for the reaction
- $$\text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons \text{PCl}_5(\text{g})$$
- (a) 1.89 (b) 0.377 (c) 1.33 (d) 13.3
63. At certain temperature compound $\text{AB}_2(\text{g})$ dissociates according to the reaction
- $$2\text{AB}_2(\text{g}) \rightleftharpoons 2\text{AB}(\text{g}) + \text{B}_2(\text{g})$$
- With degree of dissociation α , which is small compared with unity, the expression of K_p , in terms of α and initial pressure P is :
- (a) $P \frac{\alpha^3}{2}$ (b) $P \frac{\alpha^2}{3}$ (c) $P \frac{\alpha^3}{3}$ (d) $P \frac{\alpha^2}{2}$
64. For the reaction $\text{H}_2(\text{g}) + \text{CO}_2(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g})$, if the initial concentration of $[\text{H}_2] = [\text{CO}_2] = 1\text{M}$ and x moles/litre of hydrogen is consumed at equilibrium, the correct expression of K_p is :
- (a) $\frac{x^2}{(1-x)^2}$ (b) $\frac{(1-x)^2}{(1+x)^2}$ (c) $\frac{x^2}{(2-x)^2}$ (d) $\frac{x^2}{1-x^2}$
65. If D_T and D_O are the theoretical and observed vapour densities at a definite temperature and α be the degree of dissociation of a substance, then, α in the terms of D_O , D_T and n (number of moles of products formed from 1 mole reactant) is calculated by the formula :
- (a) $\frac{D_O - D_T}{(1-n)D_T}$ (b) $\frac{D_T - D_O}{(n-1)D_T}$ (c) $\frac{D_T - D_O}{(n-1)D_O}$ (d) $\frac{D - D_T}{(n-1)D_T}$
66. For the dissociation of PCl_5 into PCl_3 and Cl_2 in gaseous phase reaction, if d is the observed vapour density and D the theoretical vapour density with ' α ' as degree of dissociation, variation of D/d with ' α ' is given by?



67. At 27°C and 1 atm pressure, N_2O_4 is 20% dissociation into NO_2 . What is the density of equilibrium mixture of N_2O_4 and NO_2 at 27°C and 1 atm?

- (a) 3.11 g/litre (b) 2.11 g/litre (c) 4.5 g/litre (d) None of these


68. $COCl_2$ gas dissociates according to the equation, $COCl_2(g) \rightleftharpoons CO(g) + Cl_2(g)$. When heated to 700 K the density of the gas mixture at 1.16 atm and at equilibrium is 1.16 g/litre. The degree of dissociation of $COCl_2$ at 700 K is :

- (a) 0.28 (b) 0.50 (c) 0.72 (d) 0.42

69. The degree of dissociation of I_2 molecule of 1000°C and under atmospheric pressure is 40% by volume. If the dissociation is reduced to 20% at the same temp., total equilibrium pressure on the gas is :

- (a) 1.57 atm (b) 2.57 atm (c) 3.57 atm (d) 4.57 atm

70. Determine the value of equilibrium constant (K_c) for the reaction



if 10 moles of A_2 , 15 moles of B_2 and 5 moles of AB are placed in a 2 litre vessel and allowed to come to equilibrium. The final concentration of AB is 7.5 M :

- (a) 4.5 (b) 1.5 (c) 0.6 (d) None of these

71. At 87°C, the following equilibrium is established



If 0.50 mole of hydrogen and 1.0 mole of sulphur are heated to 87°C in 1.0 L vessel, what will be the partial pressure of H_2S at equilibrium?

- (a) 0.966 atm (b) 1.38 atm (c) 0.0327 atm (d) 1 atm

72. Pure PCl_5 is introduced into an evacuated chamber and comes to equilibrium at 247°C and 2.0 atm. The equilibrium gaseous mixture contains 40% chlorine by volume.

Calculate K_p at 247°C for the reaction



- (a) 0.625 atm (b) 4 atm (c) 1.6 atm (d) None of these

73. For the reaction



calculate K_p at 900 K, where the equilibrium steam-hydrogen mixture was 45% H_2 by volume :

- (a) 1.49 (b) 1.22 (c) 0.67 (d) None of these

74. For the reaction $\text{XCO}_3(s) \rightleftharpoons \text{XO}(s) + \text{CO}_2(g)$, $K_p = 1.642 \text{ atm}$ at 727°C . If 4 moles of $\text{XCO}_3(s)$ was put into a 50 litre container and heated to 727°C .

What mole percent of the XCO_3 remains unreacted at equilibrium?

- (a) 20 (b) 25 (c) 50 (d) None of these

75. $\text{Fe}_2\text{O}_3(s)$ may be converted to Fe by the reaction



for which $K_c = 8$ at temp. 720°C .

What percentage of the H_2 remains unreacted after the reaction has come to equilibrium?

- (a) $\sim 22\%$ (b) $\sim 34\%$ (c) $\sim 66\%$ (d) $\sim 78\%$

76. $\text{AB}_3(g)$ dissociates as $\text{AB}_3(g) \rightleftharpoons \text{AB}_2(g) + \frac{1}{2}\text{B}_2(g)$.

When the initial pressure of AB_3 is 800 torr and the total pressure developed at equilibrium is 900 torr, what fraction of $\text{AB}_3(g)$ is dissociated?

- (a) 10% (b) 20% (c) 25% (d) 30%

77. At 1000 K, a sample of pure NO_2 gas decomposes as :



The equilibrium constant K_p is 156.25 atm. Analysis shows that the partial pressure of O_2 is 0.25 atm at equilibrium. The partial pressure of NO_2 at equilibrium is :


- (a) 0.01 (b) 0.02 (c) 0.04 (d) None of these

78. Pure nitrosyl chloride (NOCl) gas was heated to 240°C in a 1.0 L container. At equilibrium the total pressure was 1.0 atm and the NOCl pressure was 0.64 atm. What would be the value of K_p ?

- (a) 1.02 atm (b) $16.875 \times 10^{-3} \text{ atm}$

- (c) $16 \times 10^{-2} \text{ atm}$ (d) None of these

79. At a certain temperature the equilibrium constant K_c is 0.25 for the reaction



If we take 1 mole of each of the four gases in a 10 litre container, what would be equilibrium concentration of $\text{A}_2(g)$?

- (a) 0.331 M (b) 0.033 M (c) 0.133 M (d) 1.33 M

80. At 200°C PCl_5 dissociates as follows :



It was found that the equilibrium vapours are 62 times as heavy as hydrogen. The % degree of dissociation of PCl_5 at 200°C is nearly :

- (a) 10% (b) 42% (c) 50% (d) 68%

81. For the dissociation reaction $\text{N}_2\text{O}_4(g) \rightleftharpoons 2\text{NO}_2(g)$, the degree of dissociation () in terms of K_p and total equilibrium pressure P is :


(a) $\sqrt{\frac{4P}{K_p}}$

(b) $\sqrt{\frac{K_p}{4P}}$


(c) $\sqrt{\frac{K_p}{4P}}$

- (d) None of these

82. Consider the following equilibrium



Then select the correct graph, which shows the variation in concentrations of NO_2 against concentrations of N_2O_4 :



83. The vapour pressure of mercury is 0.002 mm Hg at 27°C . K_c for the process $\text{Hg}(\text{l}) \rightleftharpoons \text{Hg}(\text{g})$ is:

(a) 0.002 (b) 8.12×10^{-5} (c) 6.48×10^{-5} (d) 1.068×10^{-7}

84. Calculate the equilibrium constant (K_c) for the reaction given below, if at equilibrium mixture contains 5.0 mole of A_2 , 3 mole of B_2 and 2 mole of AB_2 at 8.21 atm and 300 K



(a) 1.333 (b) 2.66 (c) 20 (d) None of these

85. For the reaction (1) and (2)



Given, $K_{p1} : K_{p2} = 9 : 1$

If the degree of dissociation of $\text{A}(\text{g})$ and $\text{X}(\text{g})$ be same then the total pressure at equilibrium (1) and (2) are in the ratio :

(a) 3 : 1 (b) 36 : 1 (c) 1 : 1 (d) 0.5 : 1


86. Given the following reaction at equilibrium, $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$. At constant pressure if some inert gas added in to the system, then:

(a) More $\text{NH}_3(\text{g})$ is produced (b) Less $\text{NH}_3(\text{g})$ is produced
(c) No effect on the equilibrium (d) K_p of the reaction is decreased

87. In which of the following equilibrium, change in volume of the system does not alter the number of moles :

(a) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$ (b) $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$
(c) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$ (d) $\text{SO}_2\text{Cl}_2(\text{g}) \rightleftharpoons \text{SO}_2(\text{g}) + \text{Cl}_2(\text{g})$


88. For the reaction



The number of moles of H_2 at equilibrium will increase if :

(a) volume is increased (b) volume is decreased
(c) argon gas is added at constant volume (d) NH_3 is removed

89. The volume of the reaction vessel containing an equilibrium mixture is increased in the following reaction



When equilibrium is re-established :

- (a) the amount of $\text{Cl}_2(\text{g})$ remains unchanged
 - (b) the amount of $\text{Cl}_2(\text{g})$ increases
 - (c) the amount of $\text{SO}_2\text{Cl}_2(\text{g})$ increases
 - (d) the amount of $\text{SO}_2(\text{g})$ decreases
90. Some inert gas is added at constant volume to the following reaction at equilibrium




Predict the effect of adding the inert gas :

- (a) The equilibrium shifts in the forward direction
 - (b) The equilibrium shifts in the backward direction
 - (c) The equilibrium remains unaffected
 - (d) The value of K_p is increased
- 91 Consider the reaction where $K_p = 0.497$ at 500 K







If the three gases are mixed in a rigid container so that the partial pressure of each gas is initially 1 atm, then which is correct observation ?

- (a) More PCl_5 will be produced
 - (b) More PCl_3 will be produced
 - (c) Equilibrium will be established when 50% reaction is complete
 - (d) None of the above
92. The preparation of $\text{SO}_3(\text{g})$ by reaction $\text{SO}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g}) \rightleftharpoons \text{SO}_3(\text{g})$ is an exothermic reaction. If the preparation follows the following temperature-pressure relationship for its % yield, then for temperatures T_1, T_2 and T_3 . The correct option is :



- (a) $T_3 > T_2 > T_1$
 - (b) $T_1 > T_2 > T_3$
 - (c) $T_1 > T_2 > T_3$
 - (d) Nothing could be predicted about temperature through given information
93. An equilibrium mixture at 700 K of $0.05\text{ M N}_2(\text{g})$, $0.3\text{ M H}_2(\text{g})$ and $0.2\text{ M NH}_3(\text{g})$ is present in a container. Now if this equilibrium is disturbed by adding $\text{N}_2(\text{g})$ so that its concentration becomes 0.15 M just after addition then which of the following graphs represents the above situation more appropriately:



94. In a vessel containing N_2 , H_2 and NH_3 at equilibrium, some helium gas is introduced so that total pressure increase while temperature and volume remain constant. According to Le Chatelier's principle, the dissociation of NH_3 :
- (a) increases (b) decreases
(c) remains unaltered (d) changes unpredictably
95. Le-Chatelier principle is not applicable to :
- (a) $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$ (b) $\text{Fe}(\text{s}) + \text{S}(\text{s}) \rightleftharpoons \text{FeS}(\text{s})$
(c) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$ (d) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$
96. Consider the following reactions. In which case the formation of product is favoured by decrease in pressure?
- (1) $\text{CO}_2(\text{g}) + \text{C}(\text{s}) \rightleftharpoons 2\text{CO}(\text{g}); H = 172.5 \text{ kJ}$
(2) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g}); H = 91.8 \text{ kJ}$
(3) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g}); H = 181 \text{ kJ}$
(4) $2\text{H}_2\text{O}(\text{g}) \rightleftharpoons 2\text{H}_2(\text{g}) + \text{O}_2(\text{g}); H = 484.6 \text{ kJ}$
- (a) 2, 3 (b) 3, 4 (c) 2, 4 (d) 1, 4
97. In which of the following reactions, the formation of product is favoured by decrease in temperature?
- (1) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g}); H = 181$
(2) $2\text{CO}_2(\text{g}) \rightleftharpoons 2\text{CO}(\text{g}) + \text{O}_2(\text{g}); H = 566$
(3) $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g}); H = 9.4$
(4) $\text{H}_2(\text{g}) + \text{F}_2(\text{g}) \rightleftharpoons 2\text{HF}(\text{g}); H = 541$
- (a) 1, 2 (b) 2 only (c) 1, 2, 3 (d) 3, 4

98. For which of the following reactions is product formation favoured by low pressure and high temperature?

- (a) $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g}); \quad H \quad 9.4 \text{ kJ}$
 (b) $\text{CO}_2(\text{g}) + \text{C}(\text{s}) \rightleftharpoons 2\text{CO}(\text{g}); \quad H \quad 172.5 \text{ kJ}$
 (c) $\text{CO}(\text{g}) + 2\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_3\text{OH}(\text{l}); \quad H \quad 21.7 \text{ kJ}$
 (d) $3\text{O}_2(\text{g}) \rightleftharpoons 2\text{O}_3(\text{g}); \quad H \quad 285 \text{ kJ}$

99. For which of the following reaction is product formation favoured by low pressure and low temperature?

- (a) $\text{CO}_2(\text{g}) + \text{C}(\text{s}) \rightleftharpoons 2\text{CO}(\text{g}); \quad H \quad 172.5 \text{ kJ}$
 (b) $\text{CO}(\text{g}) + 2\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_3\text{OH}; \quad H \quad 21.7 \text{ kJ}$
 (c) $2\text{O}_3(\text{g}) \rightleftharpoons 3\text{O}_2(\text{g}); \quad H \quad 285 \text{ kJ}$
 (d) $\text{H}_2(\text{g}) + \text{F}_2(\text{g}) \rightleftharpoons 2\text{HF}(\text{g}); \quad H \quad 541 \text{ kJ}$

100. Consider the following reactions at equilibrium and determine which of the indicated changes will cause the reaction to proceed to the right.

- (1) $\text{CO}(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g})$ (add CH_4)
 (2) $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$ (remove NH_3)
 (3) $\text{H}_2(\text{g}) + \text{F}_2(\text{g}) \rightleftharpoons 2\text{HF}(\text{g})$ (add F_2)
 (4) $\text{BaO}(\text{s}) + \text{SO}_3(\text{g}) \rightleftharpoons \text{BaSO}_4(\text{s})$ (add BaO)
 (a) 2, 3 (b) 1, 4 (c) 2, 4 (d) 2, 3, 4

101. If the pressure in a reaction vessel for the following reaction is increased by decreasing the volume, what will happen to the concentrations of CO and CO_2 ?



- (a) both the $[\text{CO}]$ and $[\text{CO}_2]$ will decrease
 (b) neither the $[\text{CO}]$ nor the $[\text{CO}_2]$ will change
 (c) the $[\text{CO}]$ will decrease and the $[\text{CO}_2]$ will increase
 (d) both the $[\text{CO}]$ and $[\text{CO}_2]$ will increase

102. Consider the following reaction and determine which of the conditions will shift the equilibrium position to the right?




- (a) increasing the temperature (b) increasing the pressure
 (c) adding a catalyst (d) none of the above is correct

103. The conversion of ozone into oxygen is exothermic. Under what conditions is ozone the most stable?




- (a) At low pressure and low temperature (b) At high pressure and high temperature
 (c) At high pressure and low temperature (d) At low pressure and high temperature

104. A system at equilibrium is described by the equation of fixed temperature T .




What effect will be the effect on equilibrium, if total pressure is increased by reducing volume?


- (a) Concentration of $\text{SO}_2\text{Cl}_2(\text{g})$ increases (b) Concentration of $\text{SO}_2(\text{g})$ increases
 (c) Concentration of $\text{Cl}_2(\text{g})$ increases (d) Concentration of all gases increases
- 105.** The reaction $2\text{NO}_2(\text{g}) \rightleftharpoons \text{N}_2\text{O}_4(\text{g})$ is an exothermic equilibrium. This means that :
 (a) equilibration of this gas mixture will be slower at high temperature
 (b) a mole of N_2O_4 will occupy twice the volume of a mole of NO_2 at the same temperature
 (c) the equilibrium will move to the right if an equilibrium mixture is cooled
 (d) the position of equilibrium will move to the left with increasing gas pressure
- 106.** Densities of diamond and graphite are 3.5 and 2.3 g/mL
 $\text{C}(\text{diamond}) \rightleftharpoons \text{C}(\text{graphite}); \quad \Delta_r H \quad 1.9 \text{ kJ/mol}$
 favourable conditions for the formation of graphite are :
 (a) high pressure and low temperature (b) low pressure and high temperature
 (c) high pressure and high temperature (d) low pressure and low temperature
- 107.** For an equilibrium $\text{H}_2\text{O}(\text{s}) \rightleftharpoons \text{H}_2\text{O}(\text{l})$, which of the following statements is true?
 (a) The pressure changes do not affect the equilibrium
 (b) More of ice melts if pressure on the system is increased
 (c) More of liquid freezes if pressure on the system is increased
 (d) The pressure changes may increase or decrease the degree of advancement of the process
- 108.** A pressure cooker reduces cooking time for food because:
 (a) the higher pressure inside the cooker crushes the food material
 (b) cooking involves chemical changes helped by a rise in temperature
 (c) heat is more evenly distributed in the cooking space
 (d) boiling point of water involved in cooking is increased
- 109.** The vapour pressure of a liquid in a closed container depends on :
 (1) temperature of liquid
 (2) quantity of liquid
 (3) surface area of the liquid
 (a) 1 only (b) 2 only (c) 1 and 3 only (d) 1, 2 and 3
- 110.** The pressure on a sample of water at its triple point is reduced while the temperature is held constant. Which phases changes are favoured?
 (I) melting of ice
 (II) sublimation of ice
 (III) vaporization of liquid water
 (a) I only (b) III only (c) II only (d) II and III
- 111.** An exothermic reaction is represented by the graph :



112. An endothermic reaction is represented by the graph :




113. A schematic plot of $\ln K_{eq}$ versus inverse of temperature for a reaction is shown below




the reaction must be :

- (a) Exothermic
 (b) Endothermic
 (c) One with negligible enthalpy change
 (d) Highly spontaneous at ordinary temperature
- 114.** What is the correct relationship between free energy change and equilibrium constant of a reaction?
- (a) $G = RT \ln K$
 (b) $G = -RT \ln K$
 (c) $G = RT \ln K$
 (d) $G = -RT \ln K$
- 115.** For the chemical equilibrium $\text{CaCO}_3(s) \rightleftharpoons \text{CaO}(s) + \text{CO}_2(g)$, $\Delta_r H$ can be determined from which one of the following plots ?



116. K_p has the value of 10^{-6} atm^3 and 10^{-4} atm^3 at 298 K and 323 K respectively for the reaction



$\Delta_r H$ for the reaction is :

- (a) 7.7 kJ/mol
 (b) 147.41 kJ/mol
 (c) 147.41 kJ/mol
 (d) None of these
- 117.** van't Hoff's equation shows the effect of temperature on equilibrium constants K_c and K_p . The K_p varies with temperature according to the relation :

(a) $\log \frac{K_{p2}}{K_{p1}} = \frac{\Delta_r H}{2.303 R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
 (b) $\log \frac{K_{p2}}{K_{p1}} = \frac{\Delta_r H}{2.303 R} \left(\frac{T_2}{T_1 T_2} - \frac{T_1}{T_1 T_2} \right)$

$$(c) \log \frac{K_{p_2}}{K_{p_1}} = \frac{E}{2.303 R} \frac{T_1 - T_2}{T_1 T_2}$$

(d) None of these

118. For a reaction, the value of K_p increases with increase in temperature. The ΔH for the reaction would be :




(a) positive

(b) negative

(c) zero

(d) cannot be predicted

119. The most stable oxides of nitrogen will be :



120. When 1 mole of pure ethyl alcohol ($\text{C}_2\text{H}_5\text{OH}$) is mixed with 1 mole of acetic acid at 25°C , the equilibrium mixture contains $2/3$ mole each of ester and water



The ΔG for the reaction at 298 K is :

(a) 3435 J

(b) 4 J

(c) 3435 J

(d) Zero

121. The value of ΔG for a reaction in aqueous phase having $K_c = 1$, would be :

(a) RT

(b) 1

(c) 0

(d) RT

122. A plot of Gibbs energy of a reaction mixture against the extent of the reaction is :


(a) minimum at equilibrium

(b) zero at equilibrium

(c) maximum at equilibrium

(d) None of these

123. For the reaction at 300 K



What is the value of equilibrium constant?

(a) 0

(b) 1

(c) 10

(d) None of these

124. Solid $\text{Ca}(\text{HCO}_3)_2$ decomposes as



If the total pressure is 0.2 bar at 420 K, what is the standard free energy change for the given reaction ($\Delta_r G^\circ$)?

(a) 840 kJ/mol

(b) 3.86 kJ/mol

(c) 6.98 kJ/mol

(d) 16.083 kJ/mol

125. The standard free energy change of a reaction is $\Delta_r G^\circ = 115 \text{ kJ mol}^{-1}$ at 298 K. Calculate the value of $\log_{10} K_p$ ($R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$)

(a) 20.16

(b) 2.303


(c) 2.016

(d) 13.83




Level 2

1. The following equilibrium constants were determined at 1120 K :




What is the equilibrium constant K_c for the following reaction at 1120 K :



- (a) $3.31 \times 10^{11} \text{ M}^{-1}$ (b) $5.5 \times 10^{10} \text{ M}^{-1}$ (c) $5.51 \times 10^6 \text{ M}^{-1}$ (d) None of these
2. One mole of $\text{N}_2(g)$ is mixed with 2 moles of $\text{H}_2(g)$ in a 4 litre vessel. If 50% of $\text{N}_2(g)$ is converted to $\text{NH}_3(g)$ by the following reaction :




What will be the value of K_c for the following equilibrium?




- (a) 256 (b) 16 (c) $\frac{1}{16}$ (d) None of these

3.



The gas A_2 in the left flask allowed to react with gas B_2 present in right flask as $\text{A}_2(g) + \text{B}_2(g) \rightleftharpoons 2\text{AB}(g); K_c = 4$ at 27°C . What is the concentration of AB when equilibrium is established?

- (a) 1.33 M (b) 2.66 M (c) 0.66 M (d) 0.33 M
4. Assume that the decomposition of HNO_3 can be represented by the following equation



and the reaction approaches equilibrium at 400 K temperature and 30 atm pressure. At equilibrium partial pressure of HNO_3 is 2 atm.

Calculate K_c in $(\text{mol/L})^3$ at 400 K :

(Use : $R = 0.08 \text{ atm-L/mol-K}$)

- (a) 4 (b) 8 (c) 16 (d) 32
5. For the equilibrium




$K_p = 9 \text{ atm}^2$ at 37°C . A 5 litre vessel contains 0.1 mole of $\text{LiCl} \cdot \text{NH}_3$. How many moles of NH_3 should be added to the flask at this temperature to derive the backward reaction for completion?

Use : $R = 0.082 \text{ atm-L/mol K}$

- (a) 0.2 (b) 0.59 (c) 0.69 (d) 0.79

6. Ammonium carbamate dissociates as



In a closed vessel containing ammonium carbamate in equilibrium, ammonia is added such that partial pressure of NH_3 now equals to the original total pressure. Calculate the ratio of partial pressure of CO_2 now to the original partial pressure of CO_2 :

- (a) 4 (b) 9 (c) $\frac{4}{9}$ (d) $\frac{2}{9}$

7. For the reaction $\text{C}_2\text{H}_6(g) \rightleftharpoons \text{C}_2\text{H}_4(g) + \text{H}_2(g)$

K_p is 5×10^{-2} atm. Calculate the mole per cent of $\text{C}_2\text{H}_6(g)$ at equilibrium if pure C_2H_6 at 1 atm is passed over a suitable catalyst at 900 K :

- (a) 20 (b) 33.33 (c) 66.66 (d) None of these

8. $2\text{NOBr}(g) \rightleftharpoons 2\text{NO}(g) + \text{Br}_2(g)$. If nitrosyl bromide (NOBr) is 40% dissociated at certain temp. and a total pressure of 0.30 atm. K_p for the reaction $2\text{NO}(g) + \text{Br}_2(g) \rightleftharpoons 2\text{NOBr}(g)$ is :

- (a) 45 (b) 25 (c) 0.022 (d) 0.25


9. Consider the partial decomposition of A as



At equilibrium 700 mL gaseous mixture contains 100 mL of gas C at 10 atm and 300 K. What is the value of K_p for the reaction?

- (a) $\frac{40}{7}$ (b) $\frac{1}{28}$ (c) $\frac{10}{28}$ (d) $\frac{28}{10}$


10. At a certain temperature and 2 atm pressure equilibrium constant (K_p) is 25 for the reaction



Initially if we take 2 moles of each of the four gases and 2 moles of inert gas, what would be the equilibrium partial pressure of NO_2 ?

- (a) 1.33 atm (b) 0.1665 atm (c) 0.133 atm (d) None of these

11. 0.020 g of selenium vapour at equilibrium occupying a volume of 2.463 mL at 1 atm and 27°C . The selenium is in a state of equilibrium according to reaction



What is the degree of association of selenium?

(At. mass of Se = 79)

- (a) 0.205 (b) 0.315 (c) 0.14 (d) None of these

12. Determine the degree of association (polymerization) for the reaction in aqueous solution, if observed (mean) molar mass of HCHO and $\text{C}_6\text{H}_{12}\text{O}_6$ is 150 :




- (a) 0.50 (b) 0.833 (c) 0.90 (d) 0.96

13. A reaction system in equilibrium according to reaction $2\text{SO}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{SO}_3(g)$ in one litre vessel at a given temperature was found to be 0.12 mole each of SO_2 and SO_3 and 5 mole of O_2 . In another vessel of one litre contains 32 g of SO_2 at the same temperature. What mass of O_2 must be added to this vessel in order that at equilibrium 20% of SO_2 is oxidized to SO_3 ?

- (a) 0.4125 g (b) 11.6 g (c) 1.6 g (d) None of these

14. The equilibrium constant K_p for the following reaction is 4.5



What would be the average molar mass (in g/mol) of an equilibrium mixture of N_2O_4 and NO_2 formed by the dissociation of pure N_2O_4 at a total pressure of 2 atm?

- (a) 69 (b) 57.5 (c) 80.5 (d) 85.5

15. A flask containing 0.5 atm pressure of $\text{A}_2(\text{g})$, some solid AB added into flask which undergoes dissociation according to $2\text{AB}(\text{s}) \rightleftharpoons \text{A}_2(\text{g}) + \text{B}_2(\text{g})$ $K_p = 0.06 \text{ atm}^2$

The total pressure (in atm) at equilibrium is :

- (a) 0.70 (b) 0.6 (c) 0.10 (d) None of these

16. A vessel of 250 litre was filled with 0.01 mole of Sb_2S_3 and 0.01 mole of H_2 to attain the equilibrium at 440°C as




After equilibrium, the H_2S formed was analysed by dissolved it in water and treating with excess of Pb^{2+} to give 1.19 g of PbS as precipitate. What is the value of K_c at 440°C ?

- (a) 1 (b) 2 (c) 4 (d) 8

17. For the reaction $2\text{A}(\text{g}) + \text{B}(\text{g}) \rightleftharpoons \text{C}(\text{g}) + \text{D}(\text{g})$; $K_c = 10^{12}$. If the initial moles of A, B, C and D are 2, 1, 7 and 3 moles respectively in a one litre vessel, what is the equilibrium concentration of A?

- (a) 4×10^{-4} (b) 2×10^{-4} (c) 10^{-4} (d) 8×10^{-4}

18. The equilibrium constant for the following reaction in aqueous solution is 0.90.



How many moles of glycerin should be added per litre of 0.10 M H_3BO_3 so that 80% of the H_3BO_3 is converted to the boric-acid-glycerin complex?

- (a) 4.44 (b) 4.52 (c) 3.6 (d) 0.08

19. Rate of diffusion of ozonized oxygen is $0.4\sqrt{5}$ times that of pure oxygen. What is the per cent degree of association of oxygen assuming pure O_2 in the sample initially?

- (a) 20 (b) 40 (c) 60 (d) None of these

20. One mole of SO_3 was placed in a two litre vessel at a certain temperature. The following equilibrium was established in the vessel



The equilibrium mixture reacts with 0.2 mole KMnO_4 in acidic medium. Hence, K_c is :

- (a) 0.50 (b) 0.25
(c) 0.125 (d) None of these


21. At 800°C , the following equilibrium is established as



The composition of equilibrium may be determined by measuring the rate of effusion of the mixture through a pin hole. It is found that at 800°C and 1 atm mixture effuses 1.6 times as fast as SO_2 effuses under the similar conditions. (At. mass of F = 19). What is the value of K_p (in atm)?

- (a) 0.315 (b) 0.685
(c) 0.46 (d) 1.49


22. The equilibrium constant for the ionization of $\text{RNH}_2(\text{g})$ in water as



is 8×10^{-6} at 25°C . Find the pH of a solution at equilibrium when pressure of $\text{RNH}_2(\text{g})$ is 0.5 bar :

- (a) 12.3 (b) 11.3 (c) 11.45 (d) None

23. Calculate ${}_rG$ for the reaction at 27°C



Given : $P_{\text{H}_2} = 0.5 \text{ bar}$; $[\text{Ag}^+] = 10^{-5} \text{ M}$;



$[\text{H}^+] = 10^{-3} \text{ M}$; ${}_rG [\text{Ag}^+(\text{aq})] = 77.1 \text{ kJ/mol}$

- (a) 154.2 kJ/mol (b) 178.9 kJ/mol
(c) 129.5 kJ/mol (d) None of these

24. When N_2O_5 is heated at certain temperature, it dissociates as $\text{N}_2\text{O}_5(\text{g}) \rightleftharpoons \text{N}_2\text{O}_3(\text{g}) + \text{O}_2(\text{g})$; $K_c = 2.5$. At the same time N_2O_3 also decomposes as : $\text{N}_2\text{O}_3(\text{g}) \rightleftharpoons \text{N}_2\text{O}(\text{g}) + \text{O}_2(\text{g})$. If initially 4.0 moles of N_2O_5 are taken in 1.0 litre flask and allowed to dissociate. Concentration of O_2 at equilibrium is 2.5 M. Equilibrium concentration of N_2O_5 is :

- (a) 1.0 M (b) 1.5 M (c) 2.166 M (d) 1.846 M

25. Two solid compounds X and Y dissociates at a certain temperature as follows



The total pressure of gases over a mixture of X and Y is :

- (a) 4.5 atm (b) 0.45 atm (c) 0.6 atm (d) None of these




Level 3



Passage 1

For a gaseous reaction




equilibrium constants K_c , K_p and K_x are represented by the following relations

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}; \quad K_p = \frac{p_C^c p_D^d}{p_A^a p_B^b} \quad \text{and} \quad K_x = \frac{x_C^c x_D^d}{x_A^a x_B^b}$$

where $[A]$ represents molar concentration of A , p_A represents partial pressure of A and P represents total pressure, x_A represents mole fraction of A

- On the basis of above work-up, select the write option
 (a) $K_p = K_c (RT)^{n_g}$; $K_x = K_p (RT)^{n_g}$ (b) $K_c = K_p (RT)^{n_g}$; $K_x = K_p P^{n_g}$
 (c) $K_c = K_x P^{n_g}$; $K_p = K_x P^{n_g}$ (d) $K_c = K_p (RT)^{n_g}$; $K_x = K_p (RT)^{n_g}$
- For the reaction $\text{SO}_2\text{Cl}_2(g) \rightleftharpoons \text{SO}_2(g) + \text{Cl}_2(g)$, $K_p = K_x$ is obtained at :
 (a) 0.5 atm (b) 0.8 atm (c) 1 atm (d) 2 atm
- For the following equilibrium relation between K_c and K_x (in terms of mole fraction) is



- (a) $K_c = K_x (RT)^{-1}$ (b) $K_c = K_x (RT)$ (c) $K_c = K_x \frac{RT}{P}$ (d) $K_c = K_x \frac{P}{RT}$



Passage 2

Variation of equilibrium constant K with temperature is given by van't Hoff equation

$$\ln K = \frac{{}_rS^\circ}{R} - \frac{{}_rH^\circ}{RT}$$

from this equation, ${}_rH^\circ$ can be evaluated if equilibrium constants K_1 and K_2 at two temperature T_1 and T_2 are known.

$$\log \frac{K_2}{K_1} = \frac{{}_rH^\circ}{2.303 R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

- For an isomerization $X(g) \rightleftharpoons Y(g)$, the temperature dependency of equilibrium constant is given by :

$$\ln K = 2 - \frac{1000}{T}$$

The value of ${}_rS^\circ$ at 300 K is :


- (a) $2R$ (b) $\frac{2}{R}$ (c) $1000 R$ (d) None of these

2. Select the correct statement :

- (a) Value of K_{eq} always increases with increasing temperature
- (b) For exothermic reaction value of K_{eq} increases with decreasing in temperature
- (c) For endothermic reaction value of K_{eq} increases with decreasing in temperature
- (d) For exothermic reaction slope is $(\log K \text{ vs. } 1/T)$ negative

3. Variation of $\log_{10} K$ with $\frac{1}{T}$ is shown by the following graph in which straight line is at 45°

hence H is :



- (a) 4.606 kJ/mol (b) 19.147 kJ/mol (c) 8.314 kJ/mol (d) 10 kJ/mol

4. The equilibrium constant K_p for the following reaction is 1 at 27°C and 4 at 47°C .



For the reaction calculate enthalpy change for the




- (a) -13.31 kcal/mol (b) 13.31 kcal/mol (c) -19.2 kcal/mol (d) 55.63 kcal/mol



Passage 3

N_2O_3 is an unstable oxide of nitrogen and it decomposes into $\text{NO}(g)$ and $\text{NO}_2(g)$ where $\text{NO}_2(g)$ is further dimerise into N_2O_4 as



A flask is initially filled with pure $\text{N}_2\text{O}_3(g)$ having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of $\text{NO}(g)$ was found to be 1.5 bar.

1. The equilibrium partial pressure of $\text{N}_2\text{O}_3(g)$ is:

- (a) 0.5 bar (b) 1.0 bar (c) 1.5 bar (d) 0.1 bar

2. The equilibrium partial pressure of $\text{NO}_2(g)$ is:

- (a) 0.066 bar (b) 0.133 bar (c) 0.423 bar (d) 0.83 bar

3. The value of K_{p_2} is:

- (a) 0.16 bar^{-1} (b) 0.32 bar^{-1} (c) 0.48 bar^{-1} (d) 0.64 bar^{-1}



Passage 4

If a system at equilibrium is subjected to a change of any one of the factors such as concentration, pressure or temperature, the system adjusts itself in such a way so as to minimise the effect of that change.

Effect of change in concentration on equilibrium:

As we add or remove reactant (or product) the ratio of equilibrium concentration become 'Q' (reaction quotient) and depending upon.

$Q < K$: equilibrium will shift in forward direction

$Q > K$: equilibrium will shift in backward direction

Effect of change in pressure :

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, then equilibrium will shift in the direction in which there is decrease in number of moles *i.e.*, towards the direction in which there is decrease in volume.

Effect of change in pressure on melting point : There are two types of solids :

- (a) Solids whose volume decreases on melting, *e.g.*, ice, diamond, carborundum, magnesium nitride and quartz.

Solid (higher volume) $\rightleftharpoons$ Liquid (lower volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

- (b) Solid whose volume increase on melting, *e.g.*, Fe, Cu, Ag, Au, etc.

Solid (lower volume) $\rightleftharpoons$ Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.

- (c) **Solubility of substances :** When solid substance are dissolved in water, either heat is evolved.

For endothermic solubility process solubility increase with increase in temperature.

For exothermic solubility decrease with increase in temperature.

- (d) **Solubility of gases in liquids :** When a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.


Effect of temperature : For endothermic reaction as temperature increases reaction shift in forward direction. For exothermic reaction as temperature increases reaction shift in backward direction.

- A 'X' (g) solute when dissolved in water heat is evolved. Then solubility of 'X' will increase :
 - high temperature, low pressure
 - low temperature, high pressure
 - high temperature, high pressure
 - low temperature, low pressure
- $\text{Fe}(l) \rightleftharpoons \text{Fe}(s)$

Above equilibrium is favoured at :

- high pressure, low temperature
- high pressure, high temperature
- low pressure, high temperature
- low pressure, low temperature

3. For the reaction



If pressure is increased by reducing the volume of the container then :


- (a) total pressure at equilibrium will remain same
- (b) concentration of all the component at equilibrium will change
- (c) concentration of all the component at equilibrium will remain same
- (d) equilibrium will shift in the backward direction

One or More Answers is/are Correct



1. A catalyst :
- (a) increases the average kinetic energy of reacting molecules
 - (b) decreases the activation energy
 - (c) can alter the reaction mechanism
 - (d) can change pre-exponential factor
2. Which of the following is correct about the chemical equilibrium?
- (a) $(\Delta G)_{T,P} = 0$
 - (b) Equilibrium constant is independent of initial concentration of reactants
 - (c) Catalyst has no effect on equilibrium state
 - (d) Reaction stops at equilibrium

3. For the reaction




If ΔG is negligible w.r.t. 1 then degree of dissociation (α) of AB_2 is proportional to :

- (a) $\frac{1}{P}$
- (b) $\frac{1}{V}$
- (c) $\frac{1}{\sqrt{P}}$
- (d) $\sqrt{V}$

4. Consider the reactions given below. In which cases will the reaction proceed toward right by increasing the pressure?

- (a) $4\text{HCl}(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{Cl}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g})$
- (b) $\text{Cl}_2(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons 2\text{HCl}(\text{g}) + \frac{1}{2}\text{O}_2(\text{g})$
- (c) $\text{CO}_2(\text{g}) + 4\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_4(\text{g}) + 2\text{H}_2\text{O}(\text{g})$
- (d) $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$

5. Ammonia is a weak base that reacts with water according to the equation



Select the correct option(s) that can increase the moles of ammonium ion in water:

- (a) Addition of HCl
- (b) Addition of NaOH
- (c) Addition of NH_4Cl
- (d) Addition of H_2O

6. Consider the reaction $2\text{CO}(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{CO}_2(\text{g}) + \text{Heat}$

Under what conditions shift is undeterminable?

- (a) Addition of O_2 and decrease in volume
- (b) Addition of CO and removal of CO_2 at constant volume

- (c) Increase in temperature and decrease in volume
 (d) Addition of CO and increase in temperature at constant volume
7. What will be the effect of addition of catalyst at constant temperature?
 (a) The equilibrium constant will remain constant
 (b) H of the reaction will remain constant
 (c) k_f and k_b will increase upto same extent
 (d) equilibrium composition will change
8. For the reaction $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$, the forward reaction at constant temperature is favoured by:
 (a) introducing an inert gas at constant volume
 (b) introducing chlorine gas at constant volume
 (c) introducing an inert gas at constant pressure
 (d) increasing the volume of the container
9. For the reaction : $\text{Cl}_2(\text{g}) + 3\text{F}_2(\text{g}) \rightleftharpoons 2\text{ClF}_3(\text{g})$; $H = 329 \text{ kJ}$, dissociation of $\text{ClF}_3(\text{g})$ will be favoured by :
 (a) increasing the temperature
 (b) increasing the volume of the container
 (c) adding of F_2 gas
 (d) adding of inert gas at constant pressure
10. Increase in the pressure for the following equilibrium results in the:

$$\text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}_2\text{O}(\text{g})$$
 (a) formation of more $\text{H}_2\text{O}(\text{l})$ (b) formation of more $\text{H}_2\text{O}(\text{g})$
 (c) increase in b.p. of $\text{H}_2\text{O}(\text{l})$ (d) decrease in b.p. of $\text{H}_2\text{O}(\text{l})$
11. Heating a II group metal carbonate leads to decomposition as :

$$\text{BaCO}_3(\text{s}) \rightleftharpoons \text{BaO}(\text{s}) + \text{CO}_2(\text{g})$$
 Equilibrium will shift left
 (a) by addition of $\text{BaO}(\text{s})$ (b) by addition of $\text{CO}_2(\text{g})$
 (c) by decreasing the temperature (d) by decreasing the volume of the vessel
12. $\text{N}_2(\text{g})$ and $\text{H}_2(\text{g})$ are allowed to react in a closed vessel at given temp. and pressure for the formation of $\text{NH}_3(\text{g})$ [$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$ 22.4 kcal]. If $\text{He}(\text{g})$ is added at equilibrium at constant pressure then which is/are correct ?
 (a) Concentration of $\text{N}_2(\text{g})$, $\text{H}_2(\text{g})$ and $\text{NH}_3(\text{g})$ decrease.
 (b) Moles of $\text{NH}_3(\text{g})$ decreases.
 (c) The extent of cooling depends on amount of $\text{He}(\text{g})$ added.
 (d) Concentration of N_2 and H_2 increases and concentration of NH_3 decreases.

Match the Column



Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I

- (A) $\text{CaCO}_3(s) \rightleftharpoons \text{CaO}(s) + \text{CO}_2(g)$
 (B) $\text{CO}(g) + \text{Cl}_2(g) \rightleftharpoons \text{COCl}_2(g)$
 (C) $\text{H}_2(g) + \text{I}_2(g) \rightleftharpoons 2\text{HI}(g)$
 (D) $\text{HCl}(g) \rightleftharpoons \text{H}^+(aq) + \text{Cl}^-(aq)$

Column-II

- (P) K_p , K_c above room temperature
 (Q) K_p , K_c above room temperature
 (R) K_p , K_c above room temperature
 (S) K_p and K_c not defined

2.

Column-I

- (A) $3\text{O}_2(g) \rightleftharpoons 2\text{O}_3(g)$
 (B) $\text{SO}_2(g) + \frac{1}{2}\text{O}_2(g) \rightleftharpoons \text{SO}_3(g)$
 (C) $2\text{HF}(g) \rightleftharpoons \text{H}_2(g) + \text{F}_2(g)$
 (D) $\text{CO}(g) + 3\text{H}_2(g) \rightleftharpoons \text{CH}_4(g) + \text{H}_2\text{O}(g)$

Column-II

- (P) no unit
 (Q) $\text{atm}^{1/2}$
 (R) atm^{-1}
 (S) atm^{-2}

3.

**Column-I
(Reaction)**

- (A) $2X(g) \rightleftharpoons Y(g) + Z(g)$
 (B) $X(g) \rightleftharpoons Y(g) + Z(g)$
 (C) $3X(g) \rightleftharpoons Y(g) + Z(g)$
 (D) $2X(g) \rightleftharpoons Y(g) + 2Z(g)$

**Column-II
(If K_c is negligible w.r.t. 1)**

- (P) $2\sqrt{K_c}$
 (Q) $3\sqrt{K_c}$
 (R) $(2K_c)^{1/3}$
 (S) $\sqrt{K_c}$

4.

Column-I

- (A) $\text{N}_2(g) + 3\text{H}_2(g) \rightleftharpoons 2\text{NH}_3(g)$; H ve
 (B) $\text{N}_2(g) + \text{O}_2(g) \rightleftharpoons 2\text{NO}(g)$; H ve
 (C) $\text{A}(g) + \text{B}(g) \rightleftharpoons 2\text{C}(g) + \text{D}(g)$; H ve
 (D) $\text{PCl}_5(g) \rightleftharpoons \text{PCl}_3(g) + \text{Cl}_2(g)$; H ve

Column-II

- (P) K increases with increase in temperature
 (Q) K decreases with increase in temperature
 (R) Pressure has no effect
 (S) Moles of product increase due to addition of inert gas at constant pressure

5.

Column-I

$$(A) \frac{K_{10} T C}{K_T C} = 2$$

$$(B) \frac{K_{10} T C}{K_T C} = \frac{1}{2}$$



Column-II

(P) Endothermic






(Q) Not affected by pressure

(R) Exothermic

(S) Affected by volume

6.

Column-I



Column-II



(P) Equilibrium shifted in forward direction

(Q) Equilibrium shifted in backward direction


(R) Equilibrium remains unaffected

(S) Theoretically we cannot predict


7.




Column-I
(Reactions)Column-II
(Equilibrium States)

(P)



(Q)





Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

- (A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** The endothermic reactions are favoured at lower temperature and the exothermic reactions are favoured at higher temperature.

STATEMENT-2 : When a system in equilibrium is disturbed by changing the temperature, it will tend to adjust itself so as to overcome the effect of change.

2. **STATEMENT-1 :** The melting point of ice decreases with increase of pressure.

STATEMENT-2 : Ice contracts on melting.

3. **STATEMENT-1 :** The equilibrium of $\text{A}(\text{g}) \rightleftharpoons \text{B}(\text{g}) + \text{C}(\text{g})$ is not affected by changing the volume.

STATEMENT-2 : K_c for the reaction does not depend on volume of the container.

4. **STATEMENT-1 :** For a chemical reaction at initial stage rate of forward reaction (r_f) is greater than rate of reversed reaction (r_b)

STATEMENT-2 : When $r_f = r_b$, chemical reaction is at equilibrium.

5. **STATEMENT-1 :** For the reaction $\text{A}(\text{g}) \rightleftharpoons \text{B}(\text{g}) + \text{C}(\text{g})$, $K_p = 1 \text{ atm}$. If we start with equal moles of all gases at 9 atm of initial pressure, then at equilibrium partial pressure of A increases.


STATEMENT-2 : Reaction quotient $Q_p < K_p$ hence equilibrium shifts in backward direction.

- 6. STATEMENT-1 :** The gas phase reaction $\text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g}) \rightleftharpoons \text{PCl}_5(\text{g})$ shifts to the right on increasing pressure.
- STATEMENT-2 :** When pressure increases, equilibrium shifts towards more number of moles.
- 7. STATEMENT-1 :** For a reaction at equilibrium, the Gibb's free energy of reaction is minimum at constant temp. and pressure.
- STATEMENT-2 :** The Gibb's free energy of both reactants and products increases and become equal at equilibrium.
- 8. STATEMENT-1 :** The physical equilibrium is not static but dynamic in nature.
- STATEMENT-2 :** The physical equilibrium is a state in which two opposing process are proceeding at the same rate.
- 9. STATEMENT-1 :** Equilibrium constant for the reverse reaction is the inverse of the equilibrium constant for the reaction in the forward direction.
- STATEMENT-2 :** Equilibrium constant depends upon the way in which the reaction is written.
- 10. STATEMENT-1 :** If $Q_p < K_p$ reaction moves in direction of reactants.
- STATEMENT-2 :** Reaction quotient is defined in the same way as equilibrium constant at any stage of the reaction.
- 11. STATEMENT-1 :** For the reaction $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$ if the volume of vessel is reduced to half of its original volume, equilibrium concentration of all gases will be doubled.
- STATEMENT-2 :** According to Le-Chatelier's principle, reaction shifts in a direction that tends to minimized the effect of the stress.
- 12. STATEMENT-1 :** The equilibrium constant of the exothermic reaction at high temperature decreases.
- STATEMENT-2 :** Since $\ln \frac{K_2}{K_1} = -\frac{H}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$ and for exothermic reaction, H is ve and thereby; $\frac{K_2}{K_1} < 1$
- 13. STATEMENT-1 :** For the reaction at certain temperature
- $$\text{A}(\text{g}) + \text{B}(\text{g}) \rightleftharpoons \text{C}(\text{g})$$
- there will be no effect by addition of inert gas at constant volume.
- STATEMENT-2 :** Molar concentration of all gases remains constant.
- 14. STATEMENT-1 :** For the physical equilibrium $\text{H}_2\text{O}(\text{s}) \rightleftharpoons \text{H}_2\text{O}(\text{l})$ on increasing temperature and increasing pressure more water will form.
- STATEMENT-2 :** Since forward reaction is endothermic in nature and volume of water is greater than that of the volume of ice.
- 15. STATEMENT-1 :** The catalyst does not alter the equilibrium constant.
- STATEMENT-2 :** Because for the catalysed reaction and uncatalysed reaction ΔH remains same and equilibrium constant depends on ΔH .

Subjective Problems



1. If 50% of CO_2 converts to CO at the following equilibrium :




and the equilibrium pressure is 12 atm. Calculate K_p .

2. Calculate partial pressure of B at equilibrium in the following equilibrium



3. In a gaseous reaction $\text{A} + 2\text{B} \rightleftharpoons 2\text{C} + \text{D}$ the initial concentration of B was 1.5 times that of A . At equilibrium the concentrations of A and D were equal. Calculate the equilibrium constant K_C .

4. For the reaction $\text{A(g)} \rightleftharpoons \text{B(g)}; K_C = 10$




Calculate K_C for the reaction $\text{D(g)} \rightleftharpoons \text{A(g)}$.

5. 5 litre vessel contains 2 moles of each of gases A and B at equilibrium. If 1 mole each of A and B are removed. Calculate K_C for the reaction $\text{A(g)} \rightleftharpoons \text{B(g)}$.

6. Calculate K_p for the reaction $\text{A(g)} \rightleftharpoons \text{B(s)} + 2\text{C(g)}; K_C = 0.2$ at 305 K.

7. A mixture of 3 moles of SO_2 , 4 moles of NO_2 , 1 mole of SO_3 and 4 moles of NO is placed in a 2.0 L vessel.



At equilibrium, the vessel is found to contain 1 mole of SO_2 . Calculate the value of K_C .

8. The density of an equilibrium mixture of N_2O_4 and NO_2 at 1 atm and 373.5 K is 2.0 g/L. Calculate K_C for the reaction $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$


9. If chemical equilibrium is attained at standard states then what is the value of ΔG° ?

10. Calculate the equilibrium concentration ratio of C to A if equimolar ratio of A and B were allowed to come to equilibrium at 300 K.



11. A definite amount of solid NH_4HS is placed in a flask already containing ammonia gas at a certain temperature and 0.1 atm pressure. NH_4HS decomposes to give NH_3 and H_2S and at equilibrium total pressure in flask is 1.1 atm. If the equilibrium constant K_p for the reaction $\text{NH}_4\text{HS(s)} \rightleftharpoons \text{NH}_3(\text{g}) + \text{H}_2\text{S(g)}$ is represented as $z \times 10^{-1}$ then find the value of z .

12. The gaseous reaction : $A(g) + nB(g) \rightleftharpoons mC(g)$ is represented by following curves



What is the value of $n - m$?

ANSWERS

Level 1

1. (b)	2. (b)	3. (c)	4. (a)	5. (b)	6. (a)	7. (a)	8. (a)	9. (a)	10. (a)
11. (b)	12. (b)	13. (b)	14. (d)	15. (c)	16. (b)	17. (d)	18. (d)	19. (c)	20. (d)
21. (b)	22. (a)	23. (c)	24. (a)	25. (b)	26. (d)	27. (d)	28. (c)	29. (d)	30. (c)
31. (d)	32. (c)	33. (c)	34. (c)	35. (a)	36. (a)	37. (c)	38. (d)	39. (c)	40. (c)
41. (a)	42. (b)	43. (a)	44. (a)	45. (a)	46. (a)	47. (c)	48. (d)	49. (d)	50. (d)
51. (a)	52. (b)	53. (a)	54. (b)	55. (b)	56. (b)	57. (b)	58. (b)	59. (c)	60. (c)
61. (a)	62. (d)	63. (a)	64. (a)	65. (c)	66. (a)	67. (a)	68. (c)	69. (d)	70. (a)
71. (a)	72. (c)	73. (a)	74. (d)	75. (b)	76. (c)	77. (b)	78. (b)	79. (c)	80. (d)
81. (b)	82. (b)	83. (d)	84. (b)	85. (b)	86. (b)	87. (a)	88. (a)	89. (b)	90. (c)
91. (a)	92. (b)	93. (a)	94. (c)	95. (b)	96. (d)	97. (d)	98. (b)	99. (c)	100. (a)
101. (d)	102. (d)	103. (b)	104. (d)	105. (c)	106. (d)	107. (b)	108. (d)	109. (a)	110. (d)
111. (c)	112. (b)	113. (a)	114. (b)	115. (a)	116. (c)	117. (b)	118. (a)	119. (a)	120. (c)
121. (c)	122. (a)	123. (b)	124. (d)	125. (a)					

Level 2

1. (a)	2. (c)	3. (c)	4. (d)	5. (d)	6. (c)	7. (c)	8. (a)	9. (c)	10. (c)
11. (b)	12. (d)	13. (b)	14. (b)	15. (a)	16. (a)	17. (a)	18. (b)	19. (c)	20. (c)
21. (d)	22. (b)	23. (c)	24. (d)	25. (b)					



Level 3

Passage-1:	1. (b)	2. (d)	3. (c)		
Passage-2:	1. (a)	2. (b)	3. (b)	4. (a)	
Passage-3:	1. (a)	2. (d)	3. (c)		
Passage-4:	1. (b)	2. (a)	3. (b)		

One or More Answers is/are correct

1. (b,c,d) 2. (a,b,c) 3. (c,d) 4. (a,c) 5. (a,c,d) 6. (c,d) 7. (a,b,c) 8. (c,d)
 9. (a,b,d) 10. (a,c) 11. (b,c,d) 12. (a,b,c)

Match the Column

1. A P; B R; C Q; D S
 2. A R; B Q; C P; D S
 3. A P; B S; C Q; D R
 4. A Q; B P, R; C P, S; D P, S
 5. A P; B R; C S; D Q
 6. A R; B Q; C P; D S
 7. A Q; B S; C P; D S

Assertion-Reason Type Questions

1. (D) 2. (A) 3. (D) 4. (B) 5. (A) 6. (C) 7. (C) 8. (A) 9. (A) 10. (D)
 11. (B) 12. (A) 13. (A) 14. (C) 15. (A)

Subjective Problems

1. 4	2. 2	3. 4	4. 5	5. 1	6. 5	7. 9	8. 2	9. 0	10. 2
11. 3	12. 5								

Hints and Solutions

Level 1

$$2. (b) Q_c = \frac{[P_2(g)]^4}{[P_4(g)]} = \frac{(1)^2}{(3/2)} = \frac{2}{3}$$

$$4. (a) \therefore \frac{K}{Q} = 0.33$$

$$K > Q$$

i.e., reaction moves in backward direction to achieve equilibrium.

5-6. (b,a)

$$Q_c = \frac{[SO_3(g)]^2}{[SO_2(g)]^2 [O_2(g)]^1} = 1000 > K_c$$

i.e., reaction moves backward

also if volume increases pressure decreases

$$Q_c = \frac{\frac{6}{2}^2}{\frac{2}{2} \cdot \frac{4}{2}^3} = \frac{9}{8}$$

$Q_c < K_c$ so reaction will proceed in forward direction.

8. (a) Since D_2 has similar chemical properties as H_2 so upon mixing, NH_2D and NHD_2 are formed.

9. (a) $[N_2]$ and $[H_2]$ start from non-zero value, decrease & then become constant.
 $[NH_3]$ start from zero, increase & then become constant.

$$10. (a) \text{ From graph, } [A]_{eq} = 0.1 M, [B]_{eq} = 0.4 M$$

$$K_c = \frac{[B]^2}{[A]^1} = 1.6$$

$$11. (b) \begin{array}{c} A(g) \rightleftharpoons 2C(g) + B(g) \\ t = 0 \quad 10M \\ t_{eq} \quad (10 - 10) \quad 2(10 -) \quad () \end{array}$$

From graph

$$[A]_{eq} = 10(1 -) = 8$$

$$0.2$$

$$K_c = \frac{[C]^2 [B]^1}{[A]^1}$$

$$4 \text{ (mol/L)}^2$$

$$12. (b) K_c = \frac{[CO]^1 [H_2]^2}{[CH_3OH]^1}$$

$$\text{So units } M^2$$

$$13. (b) K_p = (\text{atm})^{n_g}$$

$$(\text{atm})^2$$

14. (d) Activity of solid component is unit.

$$15. (c) \text{ If, } K_c = \frac{[N_2]^{1/2} [H_2]^{3/2}}{[NH_3]^1} = 4$$

$$\text{then } \frac{[NH_3]^2}{[N_2] [H_2]^3} = (4)^2 = \frac{1}{16}$$

$$\text{also } K_p = K_c (RT)^{n_g}$$

$$16. (b) \text{ For, } N_2 + O_2 \rightleftharpoons 2NO, n_g = 0$$

$$K_c = K_p \cdot K_x = 4 \cdot 10^{-4}$$

$$\text{So for } NO \rightleftharpoons \frac{1}{2} N_2 + \frac{1}{2} O_2$$

$$K_c = (4 \cdot 10^{-4})^{1/2}$$


17-21. (d, d, c, d, b)

$$K_p = K_c (RT)^{n_g}$$


$$24. (a) \ln \frac{K_2}{K_1} = \frac{H}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

25-26. (b,d)

Equilibrium constant is temperature dependant only.




Addition



$$K_3 = \sqrt{K_1} \cdot K_2$$

28-29. (c,d) Similar to Q no. 16

30-31. (c,d) Similar to Q. no. 27



$$t = 0 \quad 0.2M \quad 0.1M \quad 0.1M$$

$$t_{eq} \quad (0.2 - x) \quad (0.1 - x) \quad (0.1 + 2x)$$

$$\therefore [Z] = (0.1 + 2x) = 0.2$$

$$x = 0.05$$

$$\text{Now, } K_c = \frac{[Z]^2}{[X]^1 [Y]^1}$$

$$33. (c) K_c = \frac{[H_2]^2[S_2]}{[H_2S]}$$

$$34. (c) K_c = \frac{[CH_4][H_2S]^2}{[CS_2][H_2]^4}$$

$$35. (a) K_c = \frac{[CH_3OH]}{[CO][H_2]^2}$$

$$36. (a) K_p = \frac{P_{S_2(g)}^4}{P_{S_8(g)}} = \frac{(4 \times 0.3)^4}{0.70} = 2.96$$




$$t = 0 \quad 0.1 M$$

$$t_{eq.} \quad 0.1(1-x) \quad 2(0.1x)$$

$$\therefore$$

$$0.5$$

$$K_c = \frac{[NO_2]^2}{[N_2O_4]}$$



$$t = 0 \quad 2 \text{ moles}$$

$$t_{eq.} \quad 2-2x \quad x \quad 3x$$

$$\text{Given : } (2-2x) = 1$$

$$x = 0.5$$

$$\text{Now } K_c = \frac{[N_2][H_2]^3}{[NH_3]^2}$$

$$39. (c) K_p = \frac{1}{P_{H_2O(g)}^4}$$

$$P_{H_2O(g)} = \frac{1}{K_p^{1/4}}$$

$$(10^{12})^{1/4} = 10^3 \text{ atm}$$

$$K_p = 10^{-3} \times 760 = 0.76 \text{ torr}$$

$$40. (c) P_{H_2O} = 2 \times 10^2 \text{ atm}$$

$$\text{R.H.} = \frac{2 \times 10^2 \times 760}{38} = 100$$

$$40\%$$



$$t = 0 \quad a \text{ moles}$$

$$t_{eq.} \quad (a-x) \quad x \quad x$$

$$\therefore P_{\text{Gas}} = X_{\text{Gas}} P_{\text{Total}}$$

$$P_{NH_3} = \frac{1}{2} \times 0.66 = 0.33 \text{ atm}$$

$$P_{H_2S} = \frac{1}{2} \times 0.66 = 0.33 \text{ atm}$$

$$\text{Now } K_p = (P_{NH_3})^1 (P_{H_2S})^1$$

$$42. (b) K_c = \frac{n_B n_C}{n_A^2} \times \frac{1}{V^2}$$


$$16 \times \frac{2}{2^2} \times \frac{2^3}{V^2}$$

$$V \times \frac{1}{2}$$



$$t = 0 \quad 1 \text{ mole} \quad 1 \text{ mole}$$


$$t_{eq.} \quad (1-x) \quad (1-x)$$



$$\begin{matrix} - & - \\ (x) & (x) \end{matrix}$$

$$x = \frac{1}{3}$$

$$K_c = \frac{[CH_3COOC_2H_5][H_2O]}{[C_2H_5OH][CH_3COOH]}$$



$$t = 0 \quad 1 \text{ mole} \quad 0.5 \text{ mole} \quad -$$

$$t_{eq.} \quad (1-x) \quad (0.5-x) \quad (x)$$

AgNO₃ reacts with I⁻ and forms precipitate.

i.e., AgNO₃ + I⁻ → AgI(s) + NO₃⁻(aq.)

$$(0.5-x) = 0.25$$

$$x = 0.25$$

$$K_c = \frac{[I_3^-]}{[I_2][I^-]}$$




$$8 \times 10^{-2} \times \frac{x}{0.3-x}$$

$$x = 0.022$$

$$[H_2S(g)] = \frac{0.022}{2}$$

$$0.011 M$$




cases



$$P = \frac{0.318}{3} = 0.106$$

$$K_p = 4P^3 = 4.76 \times 10^{-3}$$



and K_c remains constant.



$$t = 0 \quad a \quad a$$

$$t_{eq.} \quad (a-x) \quad (a-x) \quad (x) \quad (x)$$

$$\therefore [D] = 2[A]$$

$$\begin{aligned} & x - 2(a - x) \\ & x - 2a/3 \\ \text{Now } K_c &= \frac{[C][D]}{[A][B]} \end{aligned}$$

50. (d) $Q < K_{eq}$
reaction moves forward
i.e., $\text{SO}_2 + \text{NO}_2 \rightleftharpoons \text{SO}_3 + \text{NO}$

$t = 0$	1	1	1	1
$t_{eq.}$	$(1 - x)$	$(1 - x)$	$(1 + x)$	$(1 + x)$

$$K_c = \frac{[\text{SO}_3][\text{NO}]}{[\text{SO}_2][\text{NO}_2]} = 16$$

$$4 \frac{1 - x}{1 + x}$$

$$x = 0.6 \quad [\text{NO}] = 1 + x = 1.6 M$$

53. (a) $2\text{HI}(g) \rightleftharpoons \text{H}_2(g) + \text{I}_2(g)$

$$t = 0 \quad a M$$

$$t_{eq.} \quad (a - a) \quad \frac{a}{2} \quad \frac{a}{2}$$

Given : 0.5

$$\text{So } K_{eq} = \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2}$$

54. (b) $\text{H}_2(g) + \text{CO}_2(g) \rightleftharpoons \text{H}_2\text{O}(g) + \text{CO}(g)$

$$\text{moles at eqm } 0.8 - x \quad 0.8 - x \quad x \quad x$$

$$\text{conc. at eqm } \frac{0.8 - x}{5} \quad \frac{0.8 - x}{5} \quad \frac{x}{5} \quad \frac{x}{5}$$

$$\therefore n_g = 0$$

$$K_c = \frac{\frac{x^2}{5}}{\frac{(0.8 - x)^2}{5}}$$

$$2 \frac{x}{0.8 - x}$$

$$x = 0.533$$

$$[\text{CO}_2(g)] = \frac{0.8 - 0.533}{5}$$

55. (b) For ideal gas mole % volume %



$$\begin{array}{ccc} \text{Initial moles} & a & 0 \\ \text{at eqm} & a(1 - \alpha) & 2\alpha \end{array}$$

As per given original volume

$$\frac{75}{100} \text{ Volume at eqm}$$

at constant T and P : $V \propto n$

$$a = 0.75 \quad a(1 - \alpha)$$

$$0.33$$

56. (b) $\text{CO} + \text{H}_2\text{O} \rightleftharpoons \text{CO}_2 + \text{H}_2$

$$\begin{array}{cccc} \text{Initial conc.} & 3 & 3 & x & 0 \\ \text{At eqm} & 2 & 2 & (x - 1) & 1 \\ & 5 & \frac{(x - 1)}{4} & 20 & x - 1 \end{array}$$

$$x = 19$$

57. (b) $\text{N}_2(g) + 3\text{H}_2(g) \rightleftharpoons 2\text{NH}_3(g)$

$$\begin{array}{ccc} t = 0 & a \text{ mole} & 3a \\ t_{eq.} & (a - x) & (3a - 3x) \end{array} \quad (2x)$$

$$\begin{array}{ccc} x & a/4 & \\ t_{eq.} & \frac{3a}{4} \text{ mole} & \frac{9a}{4} \text{ mole} \end{array} \quad \frac{a}{2} \text{ mole}$$

$$P_{\text{Gas}} \quad X_{\text{Gas}} \quad P_{\text{Total}}$$

58. (b) $2\text{NH}_3(g) \rightleftharpoons \text{N}_2(g) + 3\text{H}_2(g)$

$$\begin{array}{ccc} t = 0 & a & \\ t_{eq.} & (a - a) & \frac{a}{2} \quad \frac{3a}{2} \end{array}$$

$$\therefore PV = nRT$$

at $t = 0$; $15 \text{ V} = a \cdot R \cdot 300 \quad \dots(1)$

at $t_{eq.}$; $50 \text{ V} = (a - a) \cdot R \cdot 620 \quad \dots(2)$

$$(1) \div (2); \frac{3}{10} = \frac{300}{(1 - \alpha) \cdot 620}$$

59. (c) $\text{N}_2\text{O}_4(g) \rightleftharpoons 2\text{NO}_2(g)$

$$\begin{array}{ccc} t = 0 & 0.1 \text{ mole} & - \\ t_{eq.} & (0.1 - x) & (2x) \end{array}$$

$$K_p = \frac{(P_{\text{NO}_2})^2}{(P_{\text{N}_2\text{O}_4})^1}$$

$$0.14 = \frac{\frac{2x}{0.1 - x}^2}{\frac{0.1 - x}{0.1 - x}^1}$$

60. (c) $2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3$

$$\begin{array}{ccc} t = 0 & 5 \text{ mole} & 5 \text{ mole} \\ t_{eq.} & 5 - 2x & 5 - x \end{array} \quad 2x$$

$$\text{Given : } (2x) = \frac{60}{100} (5)$$

$$x = 1.5$$

$$\text{Now } P_{\text{Gas}} \quad X_{\text{Gas}} \quad P_{\text{Total}}$$


61. (a) Similar to Q. no. 60.

62. (d) $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$

$$\begin{array}{ccc} t = 0 & 0.4 M & (x) \quad (x) \\ t_{eq.} & (0.4 - x) & \end{array}$$

$$x = \frac{35}{100}(0.4)$$

Now $K_c = \frac{[\text{PCl}_3][\text{Cl}_2]}{[\text{PCl}_5]}$



$$t = 0 \quad P$$

$$t_{\text{eq.}} \quad P \quad P \quad P \quad \frac{P}{2}$$

$$K_p = \frac{(P_{\text{AB}})^2 (P_{\text{B}_2})^1}{(P_{\text{AB}_2})^2}$$




$$t = 0 \quad 1M \quad 1M$$

$$t_{\text{eq.}} \quad (1-x) \quad (1-x) \quad (x) \quad (x)$$

$$\therefore n_g = 0$$

$$K_p = K_c = \frac{x^2}{(1-x)^2}$$



$$t = 0 \quad a \text{ mole}$$


$$t_{\text{eq.}} \quad (a-x) \quad na$$

$$M_0 = \frac{a \cdot M_T}{(a-x) + na}$$

$$\frac{M_T - M_0}{(n-1)M_0}$$

$$\therefore D = \frac{M/2}{D_T - D_0}$$

$$\frac{D_T - D_0}{(n-1)D_0}$$



$$\frac{n}{D} = \frac{2}{(2-1)d} = \frac{D}{d} = 1$$

If $\frac{D}{d}$ at x-axis and () at y-axis

then it is a straight line positive slope and negative intercept.




$$\frac{M_T - M_0}{(n-1)M_0}$$

$$0.2 = \frac{92 - M_0}{M_0}$$

$$M_0 = \frac{76.66}{d_{\text{mixture}} \cdot \frac{PM_{\text{mixture}}}{RT}}$$

$$\frac{1}{0.821} = \frac{76.66}{300} \quad 3.11 \text{ g/litre}$$

68. (c) $M_O = \frac{dRT}{P} = \frac{57.47}{99} = 0.72$




$$t = 0 \quad a \text{ mole}$$

$$t_{\text{eq.}} \quad (a-x) \quad (2x)$$

$$K_p = \frac{P_{\text{Gas}} \cdot x_{\text{Gas}} \cdot P_{\text{Total}}^2}{\frac{1}{P_{\text{Total}}}}$$

K_p will be same in both cases.



$$\begin{array}{ccc} \text{Initial conc.} & 10/2 & 15/2 & 5/2 \\ \text{at eqm} & 5-x & 7.5-x & 2.5+2x \\ \therefore & 2.5 & 2x & 7.5 \end{array}$$

$$K_c = \frac{(7.5)^2}{2.5 \cdot 5} = 4.5$$



$$\text{Conc. at eqm} \quad 0.5 \quad x \quad - \quad x$$

$$K_c = \frac{[\text{H}_2\text{S}]}{[\text{H}_2]}$$

$$7 \cdot 10^{-2} = \frac{x}{0.5-x}$$

$$x = 0.0327$$

$$P_{\text{H}_2\text{S}} = \frac{n_{\text{H}_2\text{S}}}{V} \cdot RT$$

$$0.0327 \cdot 0.0821 \cdot 360 = 0.966$$

72. (c) The gaseous mixture contains 40% Cl_2 and 40% PCl_3 , since they are produced in 1 : 1 mole ratio. The PCl_5 % is 20.

For ideal gases mole % = volume %

$$\begin{array}{ccc} P_{\text{Cl}_2} & P_{\text{PCl}_3} & \\ 2 & 0.40 & 0.80 \text{ atm} \\ P_{\text{PCl}_5} & 2 & 0.2 \quad 0.40 \text{ atm} \\ K_p & \frac{P_{\text{PCl}_3} \cdot P_{\text{Cl}_2}}{P_{\text{PCl}_5}} & \\ & \frac{0.80 \cdot 0.80}{0.40} & \\ & = 1.6 \text{ atm} & \end{array}$$

73. (a) For an ideal gas mole % volume %

$$K_p = \frac{P_{\text{H}_2\text{O}}^2}{P_{\text{H}_2}^2} = \frac{(0.55)^2}{(0.45)^2} = 1.49$$

74. (d) Moles of CO_2 present at equilibrium


$$\frac{1.642 - 50}{0.0821 \times 1000} = 1$$

Mole % of XCO_3 decomposed

$$\frac{1}{4} \times 100 = 25\%$$

Hence, 75% remains undecomposed.

75. (b) Let initial moles of $\text{H}_2(\text{g})$ is 1



at eqm — 1 3x — 3x

$$K_c = \frac{\frac{3x}{V}}{\frac{1}{V} \left(\frac{3x}{V}\right)^3}$$

$$8 = \frac{3x}{1 - 3x}$$

$$x = 0.22$$

% of H_2 unreacted

$$\frac{1 - 3 \times 0.22}{1} \times 100 = 34\%$$


76. (c) $\text{AB}_3(\text{g}) \rightleftharpoons \text{AB}_2(\text{g}) + \frac{1}{2}\text{B}_2(\text{g})$

At eqm 800 x x x/2

$$800 - x = x = \frac{x}{2} \Rightarrow 900 - x = 200$$

$$\% \text{ dissociation} = \frac{200}{800} \times 100 = 25\%$$

77. (b) Let P is initial pressure of NO_2



At eqm P 2x 2x x

as per given $x = 0.25$

$$K_p = \frac{(2x)^2 (x)}{(P - 2x)^2}$$

$$156.25 = \frac{(0.5)^2 (0.25)}{P_{\text{NO}_2}^2}$$

$$P_{\text{NO}_2} = 0.02$$

78. (b) $2\text{NOCl}(\text{g}) \rightleftharpoons 2\text{NO}(\text{g}) + \text{Cl}_2(\text{g})$

At eqm P 2x 2x x

$$P - x = 1; P - 2x = 0.64, x = 0.12$$

$$K_p = \frac{(0.24)^2 (0.12)}{(0.64)^2} = 16.875 \times 10^{-3} \text{ atm}$$

$$79. (c) Q_c = \frac{1}{1} \frac{1}{1} = 1$$

$\therefore Q_c < K_c$ so reaction will proceed in backward direction

$$\begin{array}{ccccccc} & \text{A}_2(\text{g}) & \text{B}_2(\text{g}) & \rightleftharpoons & \text{C}_2(\text{g}) & \text{D}_2(\text{g}) \\ \text{Conc eqm} & \frac{1}{10} & \frac{x}{10} & & \frac{1}{10} & \frac{x}{10} \end{array}$$

$$0.25 = \frac{\frac{1}{10} \times \frac{x^2}{10}}{\frac{1}{10} \times \frac{x^2}{10}} \times 0.333$$

$$[\text{A}_2(\text{g})] = \frac{1}{10} \times \frac{x}{10} = \frac{1.333}{10} = 0.133$$

80. (d) Given :

$$M_{\text{Mix}} = 62 M_{\text{H}_2} = 124$$

$$\text{For } \text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$$

$$\frac{(M)_{\text{PCl}_5}}{(2 - 1)(M)_{\text{Mix}}}$$

81. (b) Similar to Q. No. 80

$$82. (b) K_c = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]}$$

If $[\text{NO}_2]$ at y-axis

$[\text{N}_2\text{O}_4]$ at x-axis

$$y = K_c x^2$$

Parabolic curve

$$83. (d) K_c = \frac{[\text{Hg}(\text{g})]; P}{C} = \frac{0.002}{760} \times \frac{1}{0.0821 \times 300}$$

$$= 1.068 \times 10^{-7} \text{ M}$$

$$84. (b) K_c = \frac{n_{\text{AB}_2}^2}{n_{\text{A}_2} n_{\text{B}_2}^2} \times V = \frac{n_{\text{AB}_2}^2}{n_{\text{A}_2} n_{\text{B}_2}^2} \times \frac{n_{\text{total}} RT}{P}$$

$$= \frac{2^2}{5 \times 3^2} \times \frac{10 \times 0.0821 \times 300}{8.21}$$

85. (b) $\text{A}(\text{g}) \rightleftharpoons \text{B}(\text{g}) + \text{C}(\text{g})$

$t = 0$ a mole

$t_{\text{eq.}}$ (a a) (a) (a)

if $P_{\text{Total}} = P_1$

then

$$K_{P_1} = \frac{\frac{1}{1} P_1}{\frac{1}{1} P_1}$$

also

$$X(g) \rightleftharpoons 2Y(g)$$

$t = 0$ b mole

$t_{eq.}$ $(b - b)$ $(2b)$

if $P_{Total} = P_2$

$$K_{P_2} = \frac{\frac{2}{1} P_2}{\frac{1}{1} P_2}$$

Given : $\frac{K_{P_1}}{K_{P_2}} = \frac{1}{9}$

86. (b) Addition of inert gas at constant pressure will increase volume so reaction moves in the direction having more gaseous moles.

87. (a) Volume change has no effect for $n_g = 0$.

88, 89 (a, b)

Volume increase leads the reaction to move towards more gaseous moles.

90. (c) Addition of inert gas at constant volume has no effect.

91. (a) $Q = 1 \text{ atm}$ K_p

92. (b) For exothermic reaction yield on temperature

i.e., $T_1 < T_2 < T_3$

93. (a) $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$

$$(t_{eq})_1 = 0.05M \quad 0.3M \quad 0.2M; K_c = \frac{800}{9}$$

$$(0.05 \quad 0.1)$$

$$(t_{eq})_2 = (0.15 \quad x) \quad (0.3 \quad 3x) \quad (0.2 \quad 2x)$$

$$K_c = \frac{[NH_3]^2}{[N_2][H_2]^3}$$

$$\frac{800}{9} = \frac{(0.2 \quad 2x)^2}{(0.15 \quad x)(0.3 \quad 3x)^3}$$

94. (c) Similar to Q. No. 90.

95. (b) Le-Chatelier's principle is not applicable in solid phase reaction.

96. (d) When pressure decreases Eq. shift toward more gaseous moles.

97. (d) When temperature decreases Eq. shift toward exothermic direction.

98-99 (b,c)

Forward reaction must be exothermic with more gaseous moles at product side.

101. (d) When volume decreases Eq. shift toward less gaseous moles.

103. (b) For O_3 to be stable the reaction shouldn't be favourable in forward direction.

104. (d) Similar to Q. No. 101.

105. (c) Temperature favours exothermic reaction.

106. (d) Graphite has lesser density

more volume

low pressure

also for exothermic low temperature

107. (b) Molar volume of $H_2O(s)$ is more than $H_2O(l)$ so on increasing pressure formation of $H_2O(l)$.

110. (d) Pressure towards more volume.

$$\ln K_p = \frac{H^0}{R} \cdot \frac{1}{T} - \frac{S^0}{R}$$

112. (b) For endothermic; $H > 0$, so slope is negative.

113. (a) For exothermic; $H < 0$, so slope is positive.

115. (a) For dissociation reaction; $H > 0$

$$\ln \frac{K_{P_2}}{K_{P_1}} = \frac{H}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$\ln \frac{10^{-4}}{10^{-6}} = \frac{H}{8.314} \left(\frac{25}{298} - \frac{25}{323} \right)$$

$${}_rH = 147.41 \text{ kJ/mol}$$

117-118 (b, a)

See Q. No. 116

119. (a) The value of K_{eq} is the measure of extent of reaction. For most stable oxide K_{eq} should be minimum.

120 - 125. (c, c, a, b, d, a)

$$G = RT \ln K_{eq}$$

$$G = (H - TS) = RT \ln K_{eq}$$

$$\ln \frac{K_2}{K_1} = \frac{H}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$\ln \frac{2.303}{0.1} = \frac{H}{8.314} \left(\frac{1}{300} - \frac{1}{0.1} \right)$$

$$K = 1$$

124. (d) $P_{total} = 0.2 \text{ bar}$

At equilibrium $P_{CO_2} = P_{H_2O} = 0.1 \text{ bar}$



$$K_p = (0.1)^2 = 0.01$$

$$\ln K_p = \frac{RT}{H}$$

$$8.314 \cdot 420 \ln (0.01)$$


$$16083.6 \text{ J/mol or } 16.083 \text{ kJ/mol}$$

Level 2



$$K_{p2} = (6 \times 10^3)^2 \text{ atm}^{-2}$$


Add (1) and (2)



$$K_p = \frac{10^{14} \times 36 \times 10^6}{36 \times 10^8}$$

 For given reaction $n_g = 1$

$$K_c = K_p(RT)^{n_g} = \frac{36 \times 10^8 \times 0.0821 \times 1120}{3.31 \times 10^{11} M^{-1}}$$




Initial moles	1	2	0
at eqm	1 - x	2 - 3x	2x


 where $x = 0.5$

$$K_c = \frac{(1/4)^2}{\left(\frac{0.5}{4}\right)^2 \left(\frac{0.5}{4}\right)^3} = 256$$

equilibrium constant for the reaction



$$K_c^1 = \frac{1}{\sqrt{K_c}} = \frac{1}{16}$$



Moles at eqm	2 - x	4 - x	2x
--------------	-------	-------	----

$$K_c = \frac{4x^2}{(2-x)(4-x)}$$

$$x = \frac{32}{24} = 1.33 \text{ mole}$$

$$[AB(g)] = \frac{2 \times 1.33}{4} = 0.66 M$$



$$\therefore P_{NO_2} = 4P_{O_2} \text{ and } P_{H_2O} = 2P_{O_2}$$


$$P_{\text{total}} = P_{HNO_3} + 7P_{O_2}$$

$$\frac{30}{P_{O_2}} = \frac{2}{\frac{P_{O_2}}{28}} + \frac{7}{4}$$


$$K_p = \frac{P_{NO_2}^4 \cdot P_{H_2O} \cdot P_{O_2}}{P_{HNO_3}^4}$$

$$K_p = K_c (RT)^{n_g} = K_c (0.08 \times 400)^3$$

$$K_c = \frac{2^{20}}{(32)^3} = 32$$



Therefore,



Initial moles	0.1	a	0
Final moles at eqm	0	(a - 0.2)	0.1


$$K_p = \frac{1}{9} (\text{atm})^2$$

 Let initial moles of NH_3 are a

$$\text{At eqm } K_p = \frac{1}{(P_{NH_3})^2} \text{ or } \frac{1}{9} = \frac{1}{(P_{NH_3})^2}$$

$$P_{NH_3} = 3 \text{ atm}$$

$$\therefore PV = nRT \Rightarrow \frac{3 \times 5}{0.0820 \times 310} = n \Rightarrow n = 0.59 \text{ i.e., } (a - 0.2) = 0.59$$

 initial moles of $NH_3 = 0.79$

 Let partial pressure at equilibrium of CO_2 be P , then $P_{NH_3} = 2P$ and total pressure at equilibrium $= 3P$


$$K_p = (2P)^2 \cdot P = 4P^3 \dots(1)$$

 If NH_3 is added and the pressure of NH_3 after addition at equilibrium is $3P$

$$K_p = 4P^3 = (P_{NH_3})^2 \cdot P_{CO_2}$$

$$P_{CO_2} = \frac{4}{9} P$$

$$\text{ratio of } \frac{P_{CO_2}}{P_{CO_2}} = \frac{4}{9}$$



At eqm	1 - x	x	x
--------	-------	---	---

$$K_p = \frac{P_{C_2H_4} \cdot P_{H_2}}{P_{C_2H_6}} = \frac{x^2}{1-x} = 5 \times 10^{-2}$$

$$x^2 = 0.05 \times \frac{0.05}{\sqrt{(0.05)^2}} = 0$$

$$x = \frac{0.05 \times 0.05}{2 \times 4 \times 0.05} = 0.20 \text{ atm}$$

 Partial pressure of C_2H_6 = mole fraction / total pressure

0.80	mole fraction	1.2
mole % of C_2H_6	0.8	100
	1.2	66.66



Initial pressure	P	0	0
at eqm	$P - 2x$	$2x$	x

where $2x = 0.40 P$


$$x = 0.20 P;$$

$$1.20 P = 0.30$$

$$P = 0.25 \text{ atm}$$


$$K_p = \frac{P_{\text{NO}}^2 P_{\text{Br}_2}}{P_{\text{NOBr}}^2} = \frac{(0.4P)^2 (0.2P)}{(0.6P)^2} = 0.0222$$

K_p for the reaction



$$\text{is } \frac{1}{0.0222} = 45$$


9. (c) $\therefore$ Initially only A is present so at eqm B and C should be present in 2 : 1



At eqm	400 mL	200 mL	100 mL
--------	--------	--------	--------

for ideal gases volume % = mole %

$$K_p = \frac{\left(\frac{200}{700}\right)^2 \left(\frac{100}{700}\right)}{\left(\frac{400}{700}\right)^2} = \frac{10}{28}$$



Initial moles	2	2	2	2
at eqm	$2 - x$	$2 - x$	$2 + x$	$2 + x$

($\therefore Q_p = K_p$)


Total no. of moles of gases at equilibrium

$$K_p = \frac{P_{\text{SO}_3} P_{\text{NO}}}{P_{\text{SO}_2} P_{\text{NO}_2}} = \frac{\left(\frac{2}{10}x\right)^2 P}{\left(\frac{2}{10}\right)^2 P} = 25$$

$$5 \frac{2}{2} x = 1.33$$

$$\text{Partial pressure of NO}_2 = \frac{\frac{2}{10}x}{\frac{0.666}{10}} P_{\text{total}} = 2$$

$$= 0.133 \text{ atm}$$



moles at eqm	$a(1 - \frac{a}{3})$	$\frac{a}{3}$
--------------	----------------------	---------------

$$\frac{\text{Observed no. of moles at equilibrium}}{\text{Initial no. of moles of reactant}} = \frac{M_T}{M_O}$$

$$\frac{a(1 - \frac{a}{3})}{a} = \frac{\frac{M_T}{3}}{M_O} \quad 0.315$$

where molar mass of Se_2

$$(M_T) = 79 \times 2 = 158$$

and molar mass of mixture

$$(M_O) = \frac{wRT}{PV} = \frac{0.02 \times 24.63}{1 \times 2.463 \times 10^{-3}} = 200$$



Conc. at eqm	$C(1 - \frac{C}{6})$	$\frac{C}{6}$
--------------	----------------------	---------------

$$\frac{\text{Observed molar concentration}}{\text{Initial molar concentration}} = \frac{M_T}{M_O}$$

M_T Theoretical molar mass of HCHO


$$\frac{C(1 - \frac{C}{6})}{C} = \frac{\frac{M_T}{6}}{M_O} = \frac{30}{150}$$

$$0.96$$



$$K_c = \frac{(0.12)^2}{(0.12)^2 \times 5} = 0.2$$

Another vessel



moles at eqm	0.5	$2x$	$y - x$	$2x$
--------------	-----	------	---------	------


$$\text{as per given } 2x = \frac{20}{100} \quad 0.5 \quad 0.1$$

$$K_c = \frac{(0.1)^2}{(0.4)^2 (y - 0.05)} = 0.20$$

$$y = 0.3625 \text{ mole}$$

$$\text{mass of O}_2 \text{ added} = 11.6 \text{ g}$$

14. (b) Let initial mole of $\text{N}_2\text{O}_4(g)$ is one



Initial moles	1	0
At equilibrium	1	2

Total no. of moles at equilibrium = 1

$$P_{\text{N}_2\text{O}_4} = \frac{1}{1} P$$

$$P_{\text{NO}_2} = \frac{2}{(1)} P$$

$$\text{Hence, } K_p = \frac{P_{\text{NO}_2}^2}{P_{\text{N}_2\text{O}_4}} = \frac{4^2}{(1-2)^2} P$$

$$4.5 = \frac{4^2}{(1-2)^2} \times 2$$

$$0.6$$


Mole fraction of N_2O_4 :

$$X_{\text{N}_2\text{O}_4} = \frac{1}{1} = 0.25$$

$$X_{\text{NO}_2} = 0.75$$

Average molar mass of mixture

$$0.25 \times 92 + 0.75 \times 46 = 57.5$$




$$0.5 \quad x \quad x$$

$$K_p = \frac{P_{\text{A}_2} P_{\text{B}_2}}{P_{\text{AB}}^2} = \frac{(0.5-x)x}{x^2} = 0.06$$

$$x = 0.1$$

$$P_{\text{total}} = P_{\text{A}_2} + P_{\text{B}_2} = 0.6 + 0.1 = 0.70 \text{ atm}$$



$$0.01 \quad x \quad 0.01 \quad 3x$$

$$2x \quad 3x$$

$$\text{where } 3x = 0.005$$



no. of moles of PbS formed

$$\frac{1.19}{238} = 0.005 \text{ mole}$$

$$\text{At eqm } [\text{H}_2] = \frac{0.005}{250} = \frac{.01}{250} ;$$

$$K_c = \frac{0.005^3}{0.005} = 1$$



$$\begin{array}{cccc} \text{Initial moles} & 1 & 1 & 7 & 3 \\ \text{at eqm} & 2-2x & 1-x & 7+x & 3 \end{array}$$

3 x

Due to very high value of K_c we can assume that reactant is almost converted into products so


$$1-x \approx y; 2-2x \approx 2y \quad x \sim 1$$

$$10^{12} = \frac{8 \times 4}{(2y)^2 y}$$

$$y^3 = 8 \times 10^{-12}$$

equilibrium concentration of

$$\text{A} = 2y = 4 \times 10^{-4}$$



$$\frac{[\text{complex}]}{[\text{H}_3\text{BO}_3]} = \frac{80}{20}$$

$$[\text{glycerine}] = \frac{80}{20 \times 0.9} = 4.44 \text{ M}$$



$$\begin{array}{cccc} \text{At eqm} & 0.1 & x & a & x \\ & & & x & 0.1 \\ & & & & \frac{80}{100} = 0.08; \end{array}$$


$$a = x = 4.44$$

$$a = 4.44 \quad 0.08 \quad 4.52 \text{ M}$$

or Initial moles = 4.52

19. (c) $\frac{r_{\text{mix}}}{r_{\text{O}_2}} = \sqrt{\frac{32}{M_{\text{mix}}}} = 0.4\sqrt{5}$

$$M_{\text{mix}} = 40 \text{ g/mol}$$




$$\begin{array}{ccc} \text{Initial moles} & 1 & 0 \\ \text{at eqm} & 1 & \frac{2}{3} \end{array}$$

$$\frac{32}{40} = \frac{1}{1} = \frac{2}{3}$$

$$0.6$$

$$\% = 60$$



$$\begin{array}{cccc} \text{moles at eqm} & 1 & 2x & 2x & x \end{array}$$

Only SO_2 (O. No. 4) will oxidized

so equivalent of SO_2 = equivalent of KMnO_4

$$2x \times 2 = 0.2 \times 5$$

$$2x = 0.5$$

$$K_c = \frac{0.5^2 \times 0.25}{\frac{2}{2} \times \frac{2}{2}} = 0.125$$



$$\frac{r_{\text{mix}}}{r_{\text{SO}_2}} = \sqrt{\frac{M_{\text{SO}_2}}{M_{\text{mix}}}}$$

$$2.56 = \frac{64}{M_{\text{mix}}}$$

$$M_{\text{mixture}} = 25$$

Let mole fraction of F_2 is x

$$25 = \frac{38 \times x + (1-x) \times 19}{1}$$

$$x = 0.315;$$

$$K_p \frac{P_F^2}{P_{F_2}} \frac{(0.685 P)^2}{(0.315 P)} \sim 1.49 \text{ atm}$$

$$22. (b) 8 \times 10^6 \frac{x^2}{0.5} \times 2 \times 10^3$$

$$\text{pOH } 2.7$$

$$23. (c) \text{ So, } \text{pH } 11.3$$


$$r_G \quad 0 \quad 77.1 \quad 2 \quad 154.2 \text{ kJ/mol}$$

$$Q \frac{[H]^2}{P_{H_2} [Ag]^2}$$



$$\frac{10^6}{0.5 (10^{10})} \quad 2 \times 10^4$$

$$G \quad r_G \quad RT \ln Q$$

$$r_G \quad 154.2 \quad \frac{8.314 \quad 300 \ln (2 \times 10^4)}{1000}$$



25. (b) Let x is partial pressure of A and y is partial pressure of C when both equilibrium simultaneously established in a vessel



$$\frac{K_{p1}}{K_{p2}} \frac{x}{y} \quad x \quad 2y$$

$$K_{p1} \quad x(2x \quad 2y)^2$$

$$x \quad 0.1 \text{ atm};$$

$$y \quad 0.05 \text{ atm}$$

$$\text{Total pressure of gases } P_A \quad P_B \quad P_C$$

$$3(x \quad y)$$

$$= 0.45 \text{ atm.}$$

Level 3


Passage-2 :

1,2,3. (a, b, b)

$$\ln K_p \quad \frac{H}{R} \quad \frac{1}{T} \quad \frac{S}{R}$$

$$4. (a) \ln \frac{K_2}{K_1} \quad \frac{H}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2}$$

Passage-3



$$x \quad y \quad y/2$$

$$P_{NO} \quad x \quad 1.5 \text{ b}$$

$$K_{p1} \frac{P_{NO} \quad P_{NO_2}}{P_{N_2O_3}}$$

$$2.5 \frac{x(x \quad y)}{2 \quad x} \quad \frac{1.5(1.5 \quad y)}{2 \quad 1.5} \quad 4.5 \quad 3y$$


$$y \quad \frac{2}{3} \text{ bar}$$

$$(1) P_{N_2O_3} \quad 2 \quad x \quad 2 \quad 0.5 \quad 0.5 \text{ b}$$

$$(2) P_{NO_2} \quad x \quad y \quad 1.5 \quad \frac{2}{3} \quad \frac{2.5}{3} \text{ b}$$

$$(3) K_{p2} \quad \frac{y}{2(x \quad y)^2} \quad \frac{2/3}{2 \quad \frac{2.5}{3}} \quad 0.48 \text{ bar}^{-1}$$

Passage-4 :



For more solubility forward shift
low temperature (exothermic)
high pressure (less volume)




i.e., $H < 0$ low temperature
less volume high pressure

One or More Answers is/are Correct

6. (c, d) Le Chatelier's principle is not quantitative. If both stress would cause the same direction of shift, the shift is determinable. If the two stresses would cause shifts in opposite directions, no deduction is possible.

Subjective Problems




$$\begin{array}{ccc} t=0 & & P \text{ atm} \\ t_{\text{eq.}} & (P - P/2) & P \end{array}$$

$$P/2 \quad P \quad 12$$

$$P \quad 8 \text{ atm}$$


$$K_p = \frac{(p_{\text{CO}})^1}{(p_{\text{CO}_2})^{1/2}}$$



$$\begin{array}{ccc} t=0 & a & \\ t_{\text{eq.}} & a-p & p \quad 2p \end{array}$$

$$K_p = \frac{(p_B)^1 (p_C)^2}{32 (P)^1 (2P)^2}$$




$$P \quad 2 \text{ atm}$$



$$\begin{array}{ccc} t=0 & 2a & 3a \\ t_{\text{eq.}} & (2a-x) & (3a-2x) \quad (2x) \quad (x) \end{array}$$

$$\text{Given : } (2a-x) \quad x$$

$$K_c = \frac{[C]^2 [D]^2}{[A]^1 [B]^2}$$



Addition

$$\text{D} \rightleftharpoons \text{A}; K_C = \frac{1}{0.01} \cdot \frac{1}{10} \cdot \frac{1}{2}$$



$$(t_{\text{eq.}})_1 \quad 0.4M \quad 0.4M$$

$$K_C = 1$$

 $\therefore K_C$ remains constant

$$\begin{array}{ccc} t=0 & 3 \text{ mole} & 4 & 1 & 4 \\ t_{\text{eq.}} & (3-x) & (4-x) & (1-x) & (4-x) \end{array}$$

$$\text{Given : } (3-x) \quad 1$$

$$K_C = \frac{[\text{SO}_3][\text{NO}]}{[\text{SO}_2][\text{NO}_2]}$$

8. For mixture

$$P.M_{\text{Mix}} = \frac{d.RT \text{ and for } \text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2}{(M)_{\text{N}_2\text{O}_4} (M)_{\text{Mix}}}$$

9. $G = RT \ln K_{eq}$


At standard states all concentration and partial pressure are unity.

$$K_{eq} = 1$$

$$10. G = RT \ln K_{eq} \quad 830 = 2 \cdot 300 \cdot 2303 \log_{10} K_{eq}$$

$$K_{eq} = 4$$


Now



$$\begin{array}{ccc} t=0 & a & a \\ t_{\text{eq.}} & (a-x) & (a-x) \quad (x) \quad (x) \end{array}$$

$$K_C = \frac{[C][D]}{[A][B]}$$

$$x = 2a/3$$



$$\begin{array}{ccc} \text{At eqm.} & - & (0.1 - P) \quad P \\ \text{As per given} & 0.1 & 2P \quad 1.1; \quad P = 0.5 \end{array}$$

$$K_p = \frac{P_{\text{NH}_3} P_{\text{H}_2\text{S}}}{0.6 \cdot 0.5} = \frac{(0.1 - P) P}{3}$$



$$\begin{array}{ccc} t=0 & 8M & 12M \\ t=t_{\text{eq.}} & 8-x & 12-nx \quad mx \end{array}$$

$$\text{Given: } [A]_{\text{eq}} = 6M$$

$$8-x = 6$$

$$x = 2M$$

$$[B]_{\text{eq}} = 8M$$

$$12-nx = 8$$

$$n(2) = 4$$

$$n = 2$$

$$[C]_{\text{eq}} = 6M$$

$$mx = 6$$

$$m(2) = 6$$

$$m = 3$$

$$m : n = 3 : 2 : 5$$

6

IONIC EQUILIBRIUM

Arrhenius Theory

When dissolved in water, the substances which releases

(i) H^+ ions are called acids e.g. HNO_3 , $HClO_4$, CH_3COOH ,

(ii) OH^- ions are called bases, e.g. $NaOH$, KOH , NH_4OH

Bronsted - Lowry Protonic Concept : acid is H^+ donor and base is H^+ acceptor
Conjugate acid and base

To get conjugate acid of a given species add H^+ to it. e.g. conjugate acid of N_2H_4 is $N_2H_5^+$.

To get conjugate base of any species subtract H^+ from it. e.g. Conjugate base of NH_3 is NH_2^- .

HCl is an acid while its conjugate ion Cl^- is a base.

HCO_3^- can donate proton as well as can accept proton hence amphoteric.

Lewis Concept : Acids are substances which accept a lone pair of electrons to form a coordinate bond and bases are the substances which donate a lone pair of electrons to form a coordinate bond.

❖ Lewis-acid is lone pair acceptor ; Lewis-base is lone-pair donor.

Ostwald dilution law: Applicable on weak electrolytes (weak acids/bases)

If α is negligible with respect to 1 ($\alpha < 0.05$), then for mono basic acid and mono acidic base.

$$K_C = C \alpha^2 \quad \text{or} \quad \alpha = \sqrt{\frac{K_C}{C}}$$

So $\alpha = \frac{1}{\sqrt{C}}$ or $\alpha = \frac{1}{\sqrt{V}}$ where, V is the volume of solution


at infinite dilution α reaches its maximum value, unity.

Presence of other solute (Common ion effect)

Degree of dissociation of a weak electrolyte is suppressed by the addition of a substance having an ion common to weak electrolyte is known as common ion effect.

Ionic Product of Water

Pure water is weak electrolyte and dissociated as:



where, K_{eq} is dissociation constant or equilibrium constant of water. Since, dissociation of water is too less and undissociated H_2O can be taken as constant, therefore

$$K_{eq} [\text{H}_2\text{O}] \rightleftharpoons [\text{H}^+][\text{OH}^-]$$

$$K_w = [\text{H}^+][\text{OH}^-]$$

The new constant K_w is known as ionic product of water. The numerical value of K_w increases considerably with temperature. It is 1.0×10^{-14} at 25°C . The variation of ionic product of water with temperature is given by

$$\ln \frac{K_{w2}}{K_{w1}} = \frac{rH}{R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$$

pH CONCEPT

S. Sorenson used a new term pH to express the hydrogen ion concentration.

$$\text{pH} = -\log[\text{H}^+] \text{ or } \text{pH} = -\log[\text{H}_3\text{O}^+] \text{ or } [\text{H}^+] = 10^{-\text{pH}}$$

For simplicity hydrogen ions are simply written as H^+ , with the understanding that H^+ ions in solution are always solvated.

❖ Ionic product (auto dissociation constant) of water

$$K_w = 1 \times 10^{-14} = [\text{H}_3\text{O}^+][\text{OH}^-] \text{ at } 298 \text{ K}$$


❖ K_w increases with increase in T

❖ $\text{pH} + \text{pOH} = \text{p}K_w = 14$ (at 298 K)


❖ For weak acid and its conjugate base

$$\text{or } \frac{K_a K_b}{K_w} = 14 \quad (\text{at } 298 \text{ K})$$

pH calculation of weak acid (HA)



pH calculation of weak base (BOH)



$$[H^+] = \frac{C}{\sqrt{K_a C}}$$

$$pH = \frac{1}{2}(pK_a - \log C)$$

Degree of dissociation

 K_a dissociation constant of weak acid

$$[OH^-] = \frac{C}{\sqrt{K_b C}}$$

$$pOH = \frac{1}{2}(pK_b - \log C)$$

Degree of dissociation

 K_b dissociation constant of weak base**pH Calculation of mixture of two weak acid HA and HB :**
 $[H^+] = \sqrt{K_{a1} C_1 + K_{a2} C_2}$; if C_1 and C_2 of both acids are negligible w.r.t. unity.
 K_{a1} and K_{a2} are dissociation constant of HA and HB respectively. C_1 and C_2 are concentration of HA and HB respectively.**pH calculation of mixture of two weak base AOH and BOH.**
 $[OH^-] = \sqrt{K_{b1} C_1 + K_{b2} C_2}$; if C_1 and C_2 of both bases are negligible w.r.t. unity.
 K_{b1} and K_{b2} are dissociation constant of AOH and BOH respectively. C_1 and C_2 are concentration of AOH and BOH respectively.**Salt Hydrolysis**

- ❖ Hydrolysis is the interaction of cation/anion/both ions of the salt with H_2O
- ❖ Salt of strong acid (SA) and strong base (SB) is not hydrolysed ; solution is neutral, pH = 7 at 25°C.

- ❖ Solution of salt of SA and weak base (WB) is acidic due to hydrolysis of cation

$$h = \text{degree of hydrolysis} = \sqrt{\frac{K_w}{K_b C}} = \sqrt{\frac{K_h}{C}}, \quad K_h \text{ (hydrolysis constant)} = \frac{C h^2}{(1-h)}$$

$$pH = 7 - \frac{1}{2}(pK_b - \log C) \quad \text{At } 25^\circ\text{C}$$

- ❖ Solution of salt of weak acid (WA) and SB is alkaline due to hydrolysis of anion

$$h = \sqrt{\frac{K_w}{K_a C}} = \sqrt{\frac{K_h}{C}}$$

$$pH = 7 + \frac{1}{2}(pK_a - \log C) \quad \text{At } 25^\circ\text{C}$$

- ❖ Solution of salt of WA and WB is neutral ($K_a = K_b$), acidic ($K_a > K_b$) and basic ($K_b > K_a$) due to hydrolysis of both

$$h = \sqrt{\frac{K_w}{K_a K_b}} = \sqrt{K_h} ; \quad pH = 7 + \frac{1}{2}(pK_a - pK_b) \quad \text{At } 25^\circ\text{C}$$

- ❖ For amphoteric salt like HS^- , HCO_3^- , $pH = \frac{pK_1 + pK_2}{2}$ or $[H^+] = \sqrt{K_1 K_2}$

 $(K_1$ first dissociation constant, K_2 second dissociation constant of acid)**Buffer Solutions** are the solutions whose pH does not change significantly on adding a small quantity of strong base or strong acid.

In general buffer solutions are simple (salts of weak acid and weak base), or mixed buffer.

Mixed buffer are of two types :

(A) Acidic buffer solutions : weak acid with its conjugate base :

made by mixing a weak acid with its conjugate base. *e.g.*, CH_3COOH with CH_3COONa , NaHCO_3 and H_2CO_3 , H_3PO_4 and NaH_2PO_4 .

(B) Basic buffer solutions : weak base with its conjugate acid :

$\text{NH}_3(\text{aq})$ with $\text{NH}_4\text{Cl}(\text{aq})$, $\text{RNH}_2(\text{aq.})$ and $\text{RNH}_3\text{Cl}(\text{aq.})$ etc,

If K_a for acid (or K_b for base) is not too high, we may write :

Henderson's Equation

For acidic buffer solution : $\text{pH} = \text{p}K_a + \log \left(\frac{[\text{salt}]}{[\text{acid}]} \right)$

For basic buffer solution : $\text{pOH} = \text{p}K_b + \log \left(\frac{[\text{salt}]}{[\text{base}]} \right)$

❖ For good buffer capacity, [salt] : [acid] ratio should be as close to one as possible. In such a case,

$\text{pH} = \text{p}K_a$. (This also is the case at midpoint of titration)

❖ Buffer capacity
$$\frac{\text{No. of moles of acid (or base) added per litre of buffer solution}}{\text{Change in pH}}$$

Maximum Buffer Capacity

It can be proved that maximum buffer capacity is achieved when the salt and acid or base concentration is equal.

Indicators

Indicator is a substance which indicates the point of equivalence in a titration by undergoing a change in its colour. They are weak acids or weak bases.

Theory of Indicators : The ionized and unionized forms of indicators have different colours. If 90 % or more of a particular form (ionised or unionised) is present, then its colour can be distinctly seen. In general, for an indicator which is weak acid, $\text{HIn} \rightleftharpoons \text{H}^+ + \text{In}^-$, the ratio of ionized to unionized form can be determined from

$$\text{pH} = \text{p}K_a + \log \frac{[\text{In}^-]}{[\text{HIn}]}$$

So, for detectable colour change, $\text{pH} = \text{p}K_a \pm 1$

This roughly gives the range of indicators. Ranges for some popular indicators are

Indicators	pH range	Colour	
		Acid medium	Basic medium
Methyl orange	3.1 - 4.4	red	yellow
Methyl red	4.2 - 6.3	red	yellow
Phenolphthaleine	8.3 - 10	colourless	pink

Equivalence point: The point at which exactly equivalent amounts of titrate and titrant have been mixed.

Acid Base Titration : For choosing a suitable indicator titration curves are of great help. In a titration curve, change in pH is plotted against the volume of alkali to a given acid. Four cases arise.

❖ **Strong acid vs strong base :** The curve is almost vertical over the pH range 3.5-10. This abrupt change corresponds to equivalence point. Any indicator suitable.

- ❖ **Weak acid vs strong base** : Final solution is basic at equivalence point. Vertical region (not so sharp) lies in pH range 6.5-10. So, phenolphthalein is suitable.
- ❖ **Strong acid vs weak base** : Final solution acidic. Vertical point in pH range 3.8-7.2. Methyl red or methyl orange suitable.
- ❖ **Weak acid vs weak base** : No sharp change in pH. No suitable indicator.


Note : At midpoint of titration, $pH = pK_a$, thus by pH measurements, K_a for weak acids (or K_b for weak bases) can be determined.

Polyprotic acids and bases : K_2, K_3 etc. can be safely neglected. If $K_1 \gg K_2 \gg K_3$ and only K_1 plays a significant role in pH calculation.

Solubility (S) and Solubility Product (K_{sp})

This is generally used for sparingly soluble salts,

Let the salt $A_x B_y$ is present in water. Let the solubility of $A_x B_y$, is S



$$K_{sp} = (xS)^x (yS)^y = x^x y^y (S)^{x+y}$$

e.g., For sparingly soluble salts (e.g., $Ag_2C_2O_4$) an equilibrium which exists as



Then

$$K_{sp} = [Ag^+]^2 [C_2O_4^{2-}]$$

Ionic product (Q) : It is defined as product of concentration of all the ions of a sparingly soluble salt at time 't'.

- ❖ **Case-1** : If $Q < K_{sp}$ then solution is unsaturated no precipitation takes place.
- ❖ **Case-2** : If $Q = K_{sp}$ then solution is saturated.
- ❖ **Case-3** : If $Q > K_{sp}$ solution is super saturated, precipitation takes place.

Common ion effect : Suppression of dissociation by adding an ion common with dissociated products. e.g., Solubility of AgCl decreases in presence of NaCl (Cl^- ion common) OR in presence of $AgNO_3$ (Ag^+ ion common)



Solubility in water at eq.:



Solubility in 0.01M NaCl at eq.:





$$K_{sp} = [Ag^+][Cl^-]_{Total}$$

$$K_{sp} = S'(S' + 0.01) \qquad \text{where } S' \ll S$$

Simultaneous solubility : When two sparingly soluble salts are present in water with at least a common ion, then solubility of both salts decreases e.g., AgCl and AgBr, $SrSO_4$ and $BaSO_4$

e.g. Calculate simultaneous solubility of AgCl and AgBr in the solution.

If simultaneous solubility of AgCl and AgBr are x and y respectively.



$$K_{sp1}(AgCl) = [Ag^+]_{Total}[Cl^-]$$

$$K_{sp2}(AgBr) = [Ag^+]_{Total}[Br^-]$$





$$K_{sp1} = (x+y)x$$

$$K_{sp2} = (x+y)y$$

Effect of complex formation and hydrolysis on solubility : Solubility increases due to complex formation and hydrolysis of anion (cation/anion)

Solubility of AgBr in aq. sol. of NH_3 .

Let solubility of AgBr is x



Due to very large value of K_f we can assume $y \approx x$

$$K_{sp} = K_f \frac{x^2}{(z-2x)^2}$$




Level 1


- Morphine ($C_{17}H_{19}NO_3$), which is used medically to relieve pain is a base. What is its conjugate acid?
 (a) $C_{17}H_{18}NO_3^+$ (b) $C_{17}H_{18}NO_3$ (c) $C_{17}H_{20}NO_3$ (d) $C_{17}H_{20}NO_3^+$
- The conjugate base of $H_2PO_4^-$ is :
 (a) H_3PO_4 (b) H_2PO_4 (c) HPO_4^{2-} (d) PO_4^{3-}
- The strongest Bronsted base in the following anion is:
 (a) CN (b) Cl (c) I (d) Br
- Which salt can furnish H in its aqueous solution?
 (a) NaH_2PO_2 (b) Na_2HPO_3 (c) Na_2HPO_4 (d) All of these
- Which is the set of amphiprotic species?
 (a) H_3O^+ , HPO_4^{2-} , HCO_3^- (b) H_2O , HPO_4^{2-} , $H_2PO_2^-$
 (c) $H_2PO_4^-$, $H_2PO_3^-$, H_2O (d) All of these
- The K_a values for HPO_4^{2-} and HSO_3^- are 4.8×10^{-13} and 6.3×10^{-8} respectively. Therefore, it follows the HPO_4^{2-} is a acid than HSO_3^- and PO_4^{3-} is a base than SO_3^{2-} .
 (a) weaker, stronger (b) stronger, weaker (c) weaker, weaker (d) stronger, stronger
- Given the following K_a values, determine which species is the strongest base ?
 HSO_4^- 1.2×10^{-2} , $H_2PO_4^-$ 6.3×10^{-8} , HCO_3^- 4.7×10^{-11}
 (a) CO_3^{2-} (b) H_2SO_4
 (c) SO_4^{2-} (d) HPO_4^{2-}
- Given that K_w for water is $10^{-13} M^2$ at $62^\circ C$, compute the sum of pOH and pH for a neutral aqueous solution at $62^\circ C$:
 (a) 7.0 (b) 13.30
 (c) 14.0 (d) 13.0
- The value of the ion product constant for water, (K_w) at $60^\circ C$ is $9.6 \times 10^{-14} M^2$. What is the $[H_3O^+]$ of a neutral aqueous solution at $60^\circ C$ and an aqueous solution with a pH 7.0 at $60^\circ C$ are respectively?
 (a) 3.1×10^{-8} , acidic (b) 3.1×10^{-7} , neutral
 (c) 3.1×10^{-8} , basic (d) 3.1×10^{-7} , basic
- For pure water :
 (a) pH increases while pOH decreases with rise in temperature
 (b) pH decreases while pOH increases with rise in temperature
 (c) both pH and pOH decreases with rise in temperature
 (d) both pH and pOH increases with rise in temperature
- A beer has a pH of 4.30. What is the $[H_3O^+]$?
 (a) 3.0×10^{-4} (b) 2.0×10^{-4} (c) 2.0×10^{-5} (d) 5.0×10^{-5}

12. The hydrogen ion concentration of the oceans is about $2 \times 10^{-9} M$. What is the pH?
 - (a) 8.85
 - (b) 9.3
 - (c) 7.85
 - (d) 8.7
13. The hydroxide ion concentration of a wine is $8 \times 10^{-11} M$. What is the pH of the wine?
 - (a) 2.10
 - (b) 2.9
 - (c) 3.9
 - (d) 4.9
14. The pH of a solution is 5. To this solution acid was added so that its pH value becomes 2.0. The increase in H^+ concentration is :
 - (a) 100 times
 - (b) 5 times
 - (c) 2.5 times
 - (d) 1000 times
15. A solution has a pH 9. It is 1000 times more basic than the original solution. What was the pH of the original solution?
 - (a) 12
 - (b) 6
 - (c) 9
 - (d) 10
16. Equal volumes of two HCl solutions of pH 3 and pH 5 were mixed. What is the pH of the resulting solution?
 - (a) 3.5
 - (b) 4.0
 - (c) 4.5
 - (d) 3.3
17. pOH of $0.002 M HNO_3$ is :
 - (a) $11 - \log 2$
 - (b) $11 \log 2$
 - (c) $3 - \log 2$
 - (d) None of these
18. Number of equivalents of HCl present in 100 mL of its solution whose pH is 4 :
 - (a) 10^{-4}
 - (b) 10^{-3}
 - (c) 10^{-2}
 - (d) 10^{-5}
19. To a 10 mL of $10^{-3} N H_2SO_4$ solution water has been added to make the total volume of one litre. Its pOH would be :
 - (a) 3
 - (b) 12
 - (c) 9
 - (d) 5
20. The pH of a solution of H_2SO_4 is 1. Assuming complete ionisation, find the molarity of H_2SO_4 solution :
 - (a) 0.1
 - (b) 0.2
 - (c) 0.05
 - (d) 2.0
21. pH of a strong diprotic acid (H_2A) at concentrations :
 - (i) $10^{-4} M$,
 - (ii) $10^{-4} N$
 are respectively :
 - (a) 3.7 and 4.0
 - (b) 4 and 3.7
 - (c) 4 and 4
 - (d) 3.7 and 3.7
22. Calcium hydroxide is a strong base. Compute $[Ca^{2+}]$ and $[OH^-]$ for a solution that is prepared by dissolving 0.60 g of $Ca(OH)_2$ in enough water to make a 1500 mL of solution.
 [Atomic mass : Ca 40, O 16, H 1]
 - (a) 5.4×10^{-3} , 9.1×10^{-13}
 - (b) 5.4×10^{-3} , 1.08×10^{-2}
 - (c) 5.4×10^{-3} , 5.4×10^{-3}
 - (d) 8.1×10^{-3} , 8.1×10^{-3}
23. pH of $10^{-6} M HCl(aq.)$ is :
 - (a) just less than 6
 - (b) exactly equal to 6
 - (c) just greater than 6
 - (d) just less than 7
24. $10^{-5} M NaOH$ solution at $25^\circ C$ is diluted 1000 times. The pH of the resultant solution will :
 - (a) be equal to 8
 - (b) lie between 7 and 8
 - (c) lie between 6 and 7
 - (d) remain unchanged
25. 4.0 g of NaOH and 4.9 g of H_2SO_4 are dissolved in water and volume is made upto 250 mL. The pH of this solution is:
 - (a) 7.0
 - (b) 1.0
 - (c) 2.0
 - (d) 12.0

26. A 25.0 mL sample of 0.10 M HCl is titrated with 0.10 M NaOH. What is the pH of the solution at the points where 24.9 and 25.1 mL of NaOH have been added?
 (a) 3.70, 10.70 (b) 3.30, 10.30 (c) 3.70, 10.30 (d) 3.0, 11.0
27. What is the pH of a solution in which 25.0 mL of 0.1 M NaOH is added to 25 mL of 0.08 M HCl and final solution is diluted to 500 mL?
 (a) 3 (b) 11 (c) 12 (d) 13
28. What is the pH of a solution in which 10.0 mL of 0.010 M $\text{Sr}(\text{OH})_2$ is added to 10.0 mL of 0.010 M HCl?
 (a) 2.30 (b) 1.50 (c) 11.70 (d) 7.00
29. At 90°C, pure water has $[\text{H}^+] = 10^{-6} \text{ M}$. If 100 mL of 0.2 M HCl is added to 200 mL of 0.1 M KOH at 90°C then pH of the resulting solution will be :
 (a) 5 (b) 6 (c) 7 (d) None of these
30. What change will occur for the following reaction if the hypochlorous acid solution is diluted from 0.1 to 0.01 M?



- (a) a decrease in the fraction of acid ionized
 (b) an increase in the fraction of acid ionized
 (c) no change in the fraction of acid ionized
 (d) we can not predict
31. Given K_a values of 5.76×10^{-10} and 4.8×10^{-10} for NH_4^+ and HCN respectively. What is the equilibrium constant for the following reaction?



- (a) 0.83 (b) 1.2 (c) 8.0×10^{-11} (d) 27.6×10^{-10}
32. Which is the strongest acid ($\text{p}K_a$ value is given)?
 (a) HCOOH [3.77] (b) $\text{C}_6\text{H}_5\text{COOH}$ [4.22]
 (c) CH_3COOH [4.7] (d) $\text{CH}_3\text{CH}_2\text{COOH}$ [4.88]
33. Given : Enthalpy of ionization of two acids :
 ΔH° (HCN) 45.2 kJ mol^{-1}
 ΔH° (CH_3COOH) 2.1 kJ mol^{-1}

Which relationship for the two acids is true ?

- (a) $\text{p}K_a(\text{HCN}) < \text{p}K_a(\text{CH}_3\text{COOH})$ (b) $\text{p}K_a(\text{HCN}) > \text{p}K_a(\text{CH}_3\text{COOH})$
 (c) $\text{p}K_a(\text{HCN}) = \text{p}K_a(\text{CH}_3\text{COOH})$ (d) $\text{p}K_a(\text{HCN}) = \frac{45.2}{2.1} \text{p}K_a(\text{CH}_3\text{COOH})$
34. What is the hydronium ion concentration of a 0.25 M HA solution? ($K_a = 4 \times 10^{-8}$)
 (a) 10^{-4} (b) 10^{-5} (c) 10^{-7} (d) 10^{-10}
35. What is the percent dissociation (%) of a 0.01 M HA solution? ($K_a = 10^{-4}$)
 (a) 9.5% (b) 1% (c) 10.5% (d) 17%
36. Given the two concentration of HCN ($K_a = 10^{-9}$) are 0.1 M and 0.001 M respectively. What will be the ratio of degree of dissociation?
 (a) 1 (b) 0.1 (c) 0.003 (d) 0.01

37. A 0.10 M solution of HF is 8.0% dissociated. What is the K_a ?
 (a) 6.4×10^{-4} (b) 8.8×10^{-4} (c) 6.95×10^{-4} (d) 7.6×10^{-4}
38. A weak base MOH of 0.1 N concentration shows a pH value of 9. What is the percentage degree of dissociation of the base?
 (a) 0.01% (b) 0.001% (c) 0.1% (d) 0.02%
39. 0.01 M HA (aq.) is 2% dissociated; $[\text{OH}^-]$ of solution is :
 (a) 2×10^{-4} (b) 10^{-8} (c) 5×10^{-11} (d) 5×10^{-12}
40. If degree of dissociation is 0.01 of decimolar solution of weak acid HA then $\text{p}K_a$ of acid is :
 (a) 2 (b) 3 (c) 5 (d) 7
41. What concentration of HCOO^- is present in a solution of 0.01 M HCOOH ($K_a = 1.8 \times 10^{-4}$) and 0.01 M HCl?
 (a) 1.8×10^{-3} (b) 10^{-2} (c) 1.8×10^{-4} (d) 10^{-4}
42. Chose the correct code

Column-I		Column-II	
(P)	$\text{p}K_b$ of X (K_a of HX 10^{-6})	(1)	6.9
(Q)	pH of 10^{-8} M HCl	(2)	8
(R)	pH of 10^{-2} M acetic acid solution (Take K_a of acetic acid 1.6×10^{-5})	(3)	10.7
(S)	pOH of a solution obtained by mixing equal volumes of solution with pH 3 and 5.	(4)	3.4

Codes :

	P	Q	R	S
(a)	1	2	4	3
(b)	4	3	2	1
(c)	2	1	4	3
(d)	1	2	3	4

43. How much water must be added to 300 mL of 0.2 M solution of CH_3COOH ($K_a = 1.8 \times 10^{-5}$) for the D.O.D. () of the acid to double?
 (a) 600 mL (b) 900 mL (c) 1200 mL (d) 1500 mL
44. What is $[\text{NH}_4^+]$ in a solution that contains 0.02 M NH_3 ($K_b = 1.8 \times 10^{-5}$) and 0.01 M KOH?
 (a) 9×10^{-6} (b) 1.8×10^{-5} (c) 3.6×10^{-5} (d) None of these
45. A hand book states that the solubility of $\text{RNH}_2(\text{g})$ in water at 1 atm and 0°C is 22.41 volumes of $\text{RNH}_2(\text{g})$ per volume of water. ($\text{p}K_b$ of $\text{RNH}_2 = 4$) Find the max. pOH that can be attained by dissolving RNH_2 in water :
 (a) 1 (b) 2 (c) 4 (d) 6

46. The $[H^+]$ of a resulting solution that is 0.01 M acetic acid ($K_a = 1.8 \times 10^{-5}$) and 0.01 M in benzoic acid ($K_a = 6.3 \times 10^{-5}$) :
- (a) 9×10^{-4} (b) 81×10^{-4} (c) 9×10^{-5} (d) 2.8×10^{-3}
47. 6.0 g weak acid HA (mol. mass = 60 g/mol.) is dissolved in water and formed 10 m³ solution. If $K_{a(HA)} = 10^{-9}$, then pOH of solution is : [Given : $\log 4 = 0.6$]
- (a) 6.7 (b) greater than 6.7 and less than 7.0
(c) greater than 7.0 and less than 7.3 (d) greater than 7.3
48. Carbonic acid (H_2CO_3), a diprotic acid has $K_{a_1} = 4.0 \times 10^{-7}$ and $K_{a_2} = 7.0 \times 10^{-11}$. What is the $[HCO_3^-]$ of a 0.025 M solution of carbonic acid?
- (a) 7.8×10^{-3} (b) 6.6×10^{-4} (c) 10^{-10} (d) 1.0×10^{-4}
49. Carbonic acid (H_2CO_3), a diprotic acid has $K_{a_1} = 4.0 \times 10^{-7}$ and $K_{a_2} = 7.0 \times 10^{-11}$. What is the $[CO_3^{2-}]$ of a 0.025 M solution of carbonic acid?
- (a) 5.5×10^{-9} (b) 5.5×10^{-8} (c) 7.0×10^{-9} (d) 7.0×10^{-11}
50. Selenious acid (H_2SeO_3), a diprotic acid has $K_{a_1} = 3.0 \times 10^{-3}$ and $K_{a_2} = 5.0 \times 10^{-8}$. What is the $[OH^-]$ of a 0.30 M solution of selenious acid?
- (a) 2.85×10^{-3} (b) 5.0×10^{-6} (c) 3.5×10^{-12} (d) 3.5×10^{-13}
51. Which of the hydrated species may exist?
I : $H_5O_2^+$ II : H_3O^+ III : H_3O_2 IV : $H_7O_3^+$
- (a) II only (b) I and II (c) I, II and IV (d) I, II, III and IV
52. Consider the following salts. Which one(s) when dissolved in water will produce an acidic solution?
1. NH_4Cl 2. $KHSO_4$ 3. $NaCN$ 4. KNO_3
- (a) 2 and 3 (b) 1 and 2 (c) only 3 (d) 2 and 4
53. Consider the following salts. Which one(s) when dissolved in water will produce a basic solution?
1. $RbClO_4$ 2. $NaNO_2$ 3. NH_4Cl 4. $NaCl$
- (a) 1 and 3 (b) only 2 (c) 1 and 2 (d) 3 and 4
54. At 25°C dissociation constants of acid HA and base BOH in aqueous solution are same. The pH of 0.01 M solution of HA is 5. The pOH of 10^{-4} M solution of BOH at the same temperature is:
- (a) 3.5 (b) 4 (c) 6 (d) None of these
55. Which of the following solutions has the highest pH?
- (a) 0.2 M $HClO_4$ (b) 0.20 M CH_3COOH
(c) 0.020 M HCl (d) 0.2 M $NaCl$
56. pH of solutions of four sodium salts NaW, NaX, NaY and NaZ were found to be 7.0, 9.0, 10.0 and 11.0 respectively. If each solution has concentration 0.1 M, the weakest acid is :
- (a) HW (b) HX (c) HY (d) HZ
57. The pH values 0.1 M solution of $HCOONa$ (I), $HCOOH$ (II), CH_3COONH_4 (III), $NaOH$ (IV), HCl (V), will be in the order :

- (a) $IV > III > I > II > V$ (b) $IV > I > III > II > V$
 (c) $II > III > I > IV > V$ (d) $V > II > III > I > IV$
58. pH of an aqueous NaCl solution at 50°C should be :
 (a) 7 (b) > 7 (c) < 7 (d) 0
59. Upon hydrolysis of sodium carbonate, the reaction takes place between:
 (a) Na and water (b) Na and OH (c) CO_3^{2-} and water (d) CO_3^{2-} and H
60. The solution of blue vitrol in water is acidic because:
 (a) CuSO_4 reacts with water (b) Cu^{2+} reacts with water
 (c) SO_4^{2-} reacts with water (d) CuSO_4 removes OH^- ions from water
61. 1 mL of 0.1 N HCl is added to 999 mL solution of NaCl. The pH of the resulting solution will be :
 (a) 7 (b) 4 (c) 2 (d) 1
62. If a salt of strong acid and weak base hydrolyses appreciably (0.1), which of the following formula is to be used to calculate degree of hydrolysis 'h' ?
 (a) $\sqrt{\frac{K_w}{K_a} a}$ (b) $\sqrt{\frac{K_w}{K_b} a}$ (c) $\sqrt{\frac{K_w}{K_a K_b}}$ (d) None of these
63. The correct formula to calculate the hydroxyl ion concentration of an aqueous solution of NH_4NO_3 is :
 (a) $\sqrt{\frac{C K_w}{K_b}}$ (b) $\sqrt{\frac{K_w K_b}{C}}$ (c) $\sqrt{\frac{C K_w}{K_a}}$ (d) $\sqrt{\frac{K_a K_w}{C}}$
64. $[\text{H}^+] \sqrt{\frac{K_w K_a}{C}}$ is suitable for
 (a) NaCl, NH_4Cl (b) CH_3COONa , NaCN
 (c) CH_3COONa , $(\text{NH}_4)_2\text{SO}_4$ (d) $\text{CH}_3\text{COONH}_4$, $(\text{NH}_4)_2\text{CO}_3$
65. What is the hydrolysis constant of the OCl^- ion? The ionization constant of HOCl is 3.0×10^{-8} .
 (a) 3.33×10^{-8} (b) 3.33×10^{-7} (c) 3.0×10^{-7} (d) 3.33×10^{-6}
66. What is the pH of a 0.10 M $\text{C}_6\text{H}_5\text{O}^-$ solution? The K_a of $\text{C}_6\text{H}_5\text{OH}$ is 1.0×10^{-10} .
 (a) 10.51 (b) 11.04 (c) 11.50 (d) 12
67. Calculate the $[\text{OH}^-]$ in 0.01 M aqueous solution of NaOCN (K_b for OCN 10^{-10}):
 (a) 10^{-6} M (b) 10^{-7} M (c) 10^{-8} M (d) None of these
68. What is the ionization constant of an acid if the hydronium ion concentration of a 0.40 M solution is $1.40 \times 10^{-4} \text{ M}$?
 (a) 1.96×10^{-8} (b) 1.22×10^{-9} (c) 4.90×10^{-8} (d) 1.40×10^{-6}
69. The degree of hydrolysis of 0.1 M RNH_3Cl solution is 1.0%. If the concentration of RNH_3Cl is made 0.4 M, what is the new degree of hydrolysis (in percentage)?
 (a) 0.01 (b) 0.001 (c) 0.2 (d) 0.5
70. % hydrolysis of 0.1M $\text{CH}_3\text{COONH}_4$, when $K_a(\text{CH}_3\text{COOH}) = K_b(\text{NH}_4\text{OH}) = 1.8 \times 10^{-5}$ is:
 (a) 0.55 (b) 7.63 (c) 0.55×10^{-2} (d) 7.63×10^{-3}

71. The enthalpy of neutralisation of four acids HA, HB, HC and HD with NaOH are -13 , -12 , -11 , -10 kcal/mol. Which salt has maximum degree of hydrolysis?
 (a) 1M NaA (b) 1M NaB (c) 1M NaC (d) 1M NaD
72. Calculate $[H^+]$ at equivalent point between titration of 0.1 M, 25 mL of weak acid HA ($K_{a(HA)} = 10^{-5}$) with 0.05 M NaOH solution :
 (a) 3×10^{-9} (b) 1.732×10^{-9} (c) 8 (d) 10
73. When a salt of weak acid and weak base is dissolved in water at 25°C , the pH of the resulting solution will always :
 (a) be 7 (b) be greater than 7
 (c) be less than 7 (d) depend upon K_a and K_b values
74. What will be the pH of an aqueous solution of 1.0 M ammonium formate?
 Given : $pK_a = 3.8$ and $pK_b = 4.8$
 (a) 7.5 (b) 3.4 (c) 6.5 (d) 10.2
75. What will be the pH and % (degree of hydrolysis) respectively for the salt BA of 0.1 M concentration? Given : K_a for HA 10^{-6} and K_b for BOH 10^{-6}
 (a) 5, 1% (b) 7, 10% (c) 9, 0.01% (d) 7, 0.01%
76. The percentage degree of hydrolysis of a salt of weak acid (HA) and weak base (BOH) in its 0.1 M solution is found to be 10%. If the molarity of the solution is 0.05 M, the percentage hydrolysis of the salt should be :
 (a) 5% (b) 10% (c) 20% (d) None of these
77. What is the hydronium ion concentration of a 0.02 M solution of Cu^{2+} solution of copper(II) perchlorate? The acidity constant of the following reaction is 5×10^{-9} .

$$\text{Cu}^{2+}(\text{aq.}) + 2\text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{Cu}(\text{OH})^+(\text{aq.}) + \text{H}_3\text{O}^+(\text{aq.})$$

 (a) 1×10^{-5} (b) 7×10^{-4} (c) 5×10^{-4} (d) 1×10^{-4}
78. What is the acidity constant for the following reaction given that the hydronium ion concentration of a 0.04 M solution of Ni^{2+} solution of nickel(II) perchlorate is 4.5×10^{-6} ?

$$\text{Ni}^{2+}(\text{aq.}) + 2\text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{Ni}(\text{OH})^+(\text{aq.}) + \text{H}_3\text{O}^+(\text{aq.})$$


 (a) 2×10^{-12} (b) 4×10^{-6} (c) 5×10^{-12} (d) 5×10^{-10}
79. Calculate the pH at 25°C of a solution that is 0.10 M in $\text{Fe}(\text{NO}_3)_3$. The acid dissociation constant for the reaction given below is 1.0×10^{-3} .

$$[\text{Fe}(\text{H}_2\text{O})_6]^{3+} + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}_3\text{O}^+(\text{aq.}) + [\text{Fe}(\text{H}_2\text{O})_5(\text{OH})]^{2+}$$


 (a) 2.00 (b) 2.02 (c) 2.30 (d) 2.50
80. Approximate pH of 0.01 M NaHA is calculated by :
 ($K_{a_1} = 10^{-6}$ and $K_{a_2} = 10^{-8}$ are ionization constants of H_2A)
 (a) $\text{pH} = 7 + \frac{pK_{a_1}}{2} + \frac{\log C}{2}$ (b) $\text{pH} = 7 + \frac{pK_{a_1}}{2} - \frac{\log C}{2}$
 (c) $\text{pH} = \frac{pK_{a_1} + pK_{a_2}}{2}$ (d) None of these

- 81.** H_3PO_4 is a weak triprotic acid; approximate pH of 0.1 M $\text{Na}_2\text{HPO}_4(aq.)$ is calculated by :
- (a) $\frac{1}{2} [pK_{a_1} + pK_{a_2}]$ (b) $\frac{1}{2} [pK_{a_2} + pK_{a_3}]$ (c) $\frac{1}{2} [pK_{a_1} + pK_{a_3}]$ (d) $pK_{a_1} + pK_{a_2}$
- 82.** Which of the following is a buffer solution?
- (a) 500 mL of 0.1 N CH_3COOH + 500 mL of 0.1 N NaOH
 (b) 500 mL of 0.1 N CH_3COOH + 500 mL of 0.1 N HCl
 (c) 500 mL of 0.1 N CH_3COOH + 500 mL of 0.2 N NaOH
 (d) 500 mL of 0.2 N CH_3COOH + 500 mL of 0.1 N NaOH
- 83.** If 20 mL of 0.1 M NaOH is added to 30 mL of 0.2 M CH_3COOH ($pK_a = 4.74$), the pH of the resulting solution is :
- (a) 4.44 (b) 9.56 (c) 8.96 (d) 9.26
- 84.** H_2CO_3 - NaHCO_3 found in blood helps in maintaining pH of the blood close to 7.4. An excess of acid entering the blood stream is removed by:
- (a) HCO_3^- (b) H_2CO_3 (c) H^+ ion (d) CO_3^{2-} ion
- 85.** 100 mL of 0.02 M benzoic acid ($pK_a = 4.20$) is titrated using 0.02 M NaOH, pH after 50 mL and 100 mL of NaOH have been added are:
- (a) 3.50, 7 (b) 4.2, 7 (c) 4.2, 8.1 (d) 4.2, 8.25
- 86.** What is the pH of a solution of 0.28 M acid and 0.84 M of its conjugate base if the ionization constant of acid is 4×10^{-4} ?
- (a) 3.88 (b) 3.34 (c) 7 (d) 10.12
- 87.** The toxic compound 2,4-dinitrophenol has $K_a = 10^{-4}$. In an experiment, a buffer solution of 2,4-dinitrophenol was prepared with the pH adjusted to 5. Calculate the ratio of the concentrations of the dissociated ion to the undissociated acid :
- (a) 0.01 (b) 0.1 (c) 10 (d) 100
- 88.** Equilibrium constant for the following reaction is 1×10^9 :
- $$\text{C}_5\text{H}_5\text{N}(aq.) + \text{H}_2\text{O}(l) \rightleftharpoons \text{C}_5\text{H}_5\text{NH}^+(aq.) + \text{OH}^-(aq.)$$
- Determine the moles of pyridinium chloride ($\text{C}_5\text{H}_5\text{N.HCl}$) that should be added to 500 mL solution of 0.4 M pyridine ($\text{C}_5\text{H}_5\text{N}$) to obtain a buffer solution of pH 5 :
- (a) 0.1 mole (b) 0.2 mole (c) 0.3 mole (d) 0.4 mole
- 89.** Which one of the following mixture does not act as a buffer solution?
- (a) Boric acid and borax
 (b) Sodium phosphate & disodium hydrogen phosphate
 (c) Sodium propionate and propionic acid
 (d) Sod. acetate and sodium propionate
- 90.** The acid dissociation constant of uric acid is $K_a = 4.0 \times 10^{-6}$ M. The pH of a sample of urine is 6.0. What is the ratio of concentration of urate ion to uric acid in the urine?
- (a) 2.0 (b) 4.0 (c) 6.0 (d) 0.25
- 91.** CH_3NH_2 (0.12 mole, $pK_b = 3.3$) is added to 0.08 moles of HCl and the solution is diluted to one litre, resulting pH of solution is :
- (a) 10.7 (b) 3.6 (c) 10.4 (d) 11.3

- 92.** An aqueous solution at room temperature contains $0.1\text{ M NH}_4\text{Cl}$ and $0.01\text{ M NH}_4\text{OH}$ ($pK_b = 5$), the pH of the solution is :
 (a) 7.5 (b) 6.8 (c) 6.5 (d) 8.0
- 93.** Which of the following will decrease with dilution at a given temperature?
 (a) pH of 10^{-3} M acetic acid solution
 (b) pH of 10^{-3} M aniline solution
 (c) degree of dissociation of 10^{-3} M acetic acid
 (d) degree of dissociation of 10^{-3} M aniline solution
- 94.** A 1 L solution contains $0.2\text{ M NH}_4\text{OH}$ and $0.2\text{ M NH}_4\text{Cl}$. If 1.0 mL of 0.001 M HCl is added to it, what will be the $[\text{OH}^-]$ of the resulting solution ($K_b = 2 \times 10^{-5}$)
 (a) 2×10^{-5} (b) 5×10^{-10} (c) 2×10^{-3} (d) None of these
- 95.** 0.1 M formic acid solution is titrated against 0.1 M NaOH solution. What would be the difference in pH between $1/5$ and $4/5$ stages of neutralization of acid ?
 (a) $2 \log 3/4$ (b) $2 \log 1/5$ (c) $\log 1/3$ (d) $2 \log 4$
- 96.** In the neutralization process of H_3PO_4 and NaOH , the number of buffers formed will be:
 (a) 3 (b) 1 (c) 2 (d) 4
- 97.** A buffer solution is made up of acetic acid [$pK_a = 5$] having conc. 1.5 M and sodium acetate having conc. 0.15 M . What is the number of OH^- ions present in 1 litre solution?
 (a) $10^{10} N_A$ (b) $10^4 N_A$ (c) $10^3 N_A$ (d) $10^6 N_A$
- 98.** The pH of a solution of $0.10\text{ M CH}_3\text{COOH}$ increases when which of the following substances is added?
 (a) NaHSO_4 (b) HClO_4 (c) KNO_3 (d) K_2CO_3
- 99.** H^+ ion concentration of water does not change by adding :
 (a) CH_3COONa (b) NaNO_3 (c) NaCN (d) Na_2CO_3
- 100.** pK_a of NH_4^+ is 9.26. Hence, effective range for $\text{NH}_4\text{OH} - \text{NH}_4\text{Cl}$ buffer is about pH :
 (a) 8.26 to 10.26 (b) 4.74 to 5.74 (c) 3.74 to 5.74 (d) 8.26 to 9.26
- 101.** 1.0 L solution is prepared by mixing 61 g benzoic acid ($pK_a = 4.2$) with 72 g of sodium benzoate and then 300 mL 1.0 M HBr solution was added. The pH of final solution is:
 (a) 3.6 (b) 3.8 (c) 4.2 (d) 4.8
- 102.** The pH of a solution containing 0.4 M HCO_3^- and 0.2 M CO_3^{2-} is :
 $[K_{a1}(\text{H}_2\text{CO}_3) = 4 \times 10^{-7}; K_{a2}(\text{HCO}_3^-) = 4 \times 10^{-11}]$
 (a) 10.4 (b) 10.1 (c) 6.1 (d) 10.7
- 103.** The pH of the resultant solution of 20 mL of $0.1\text{ M H}_3\text{PO}_4$ and 20 mL of $0.1\text{ M Na}_3\text{PO}_4$ is :
 (a) $pK_{a1} - \log 2$ (b) pK_{a1} (c) pK_{a2} (d) $\frac{pK_{a1} + pK_{a2}}{2}$
- 104.** Which one of the following curves represents the graph of pH during the titration of NaOH and HCl(aq.) ?



105. When 100 mL of 0.1 M NaCN solution is titrated with 0.1 M HCl solution the variation of pH of solution with volume of HCl added will be :



106. The best indicator for the detection of end point in titration of a weak acid and a strong base is :

- (a) Methyl orange (3.1 to 4.4) (b) Methyl red (4.2 to 6.3)
(c) Bromothymol blue (6 to 7.6) (d) Phenolphthalein (8.2 to 10)

107. Select the best indicator from the given table for titration of 20 mL of 0.02 M CH_3COOH with 0.02 M NaOH. Given $\text{p}K_a$ (CH_3COOH) = 4.74





	Indicator	pH range
(I)	Bromothymol blue	6.0–7.6
(II)	Thymolphthalein	9.3–10.5
(III)	Malachite green	11.4–13
(IV)	M-Cresol purple	7.4–9.0

- (a) I (b) II (c) III (d) IV

108. Bromothymol blue is an indicator with a K_a value of 6×10^{-5} . What % of this indicator is in its basic form at a pH of 5 ?

- (a) 40 (b) 85.7 (c) 14.3 (d) 60

- 109.** An acid-base indicator has $K_{\text{HIn}} = 3.0 \times 10^{-5}$. The acid form of the indicator is red and the basic form is blue. The change in $[\text{H}^+]$ required to change the indicator from 75% red to 75% blue is :
 (a) $8 \times 10^{-5} \text{ M}$ (b) $9 \times 10^{-5} \text{ M}$ (c) $1 \times 10^{-5} \text{ M}$ (d) $3 \times 10^{-4} \text{ M}$
- 110.** An acid-base indicator which is a weak acid has a pK_a value 5.45. At what concentration ratio of sodium acetate to acetic acid would the indicator show a colour half-way between those of its acid and conjugate base forms ? [pK_a of acetic acid 4.75]
 (a) 4 : 1 (b) 6 : 1 (c) 5 : 1 (d) 3 : 1
- 111.** A 20.0 mL sample of a 0.20 M solution of the weak diprotic acid H_2A is titrated with 0.250 M NaOH. The concentration of solution at the second equivalent point is :
 (a) 0.10 M NaHA (b) 0.153 M Na_2A (c) 0.10 M Na_2A (d) 0.0769 M Na_2A
- 112.** During the titration of a weak diprotic acid (H_2A) against a strong base (NaOH), the pH of the solution half-way to the first equivalent point and that at the first equivalent point are given respectively by :
 (a) pK_{a_1} and $pK_{a_1} - pK_{a_2}$ (b) $\sqrt{K_{a_1} C}$ and $\frac{pK_{a_1} + pK_{a_2}}{2}$
 (c) pK_{a_1} and $\frac{pK_{a_1} + pK_{a_2}}{2}$ (d) pK_{a_1} and pK_{a_2}
- 113.** In which of the following cases is the solution of AgCl unsaturated?
 (a) $[\text{Ag}^+][\text{Cl}^-] < K_{sp}$ (b) $[\text{Ag}^+][\text{Cl}^-] = K_{sp}$ (c) $[\text{Ag}^+][\text{Cl}^-] > K_{sp}$ (d) $[\text{Ag}^+][\text{Cl}^-] = K_{sp}$
- 114.** When equal volumes of the following solutions are mixed, the precipitation of AgCl ($K_{sp} = 1.8 \times 10^{-10}$) will occur with :
 (a) 10^{-4} M (Ag^+) and 10^{-4} M (Cl^-) (b) 10^{-5} M (Ag^+) and 10^{-5} M (Cl^-)
 (c) 10^{-5} M (Ag^+) and 10^{-6} M (Cl^-) (d) 10^{-10} M (Ag^+) and 10^{-10} M (Cl^-)
- 115.** Choose the correct set of True/False for following statements:
 (i) Silver chloride is more soluble in very concentrated sodium chloride solution than in pure water.
 (ii) The pH of a buffer solution does not change on addition of small amount of an acid or a base.
 (iii) Addition of NH_4Cl does not affect the pH of a solution of NH_4OH
 (iv) Degree of hydrolysis of ammonium acetate does not depend upon the concentration of ammonium acetate solution.
 (v) A mixture of acetic acid and sodium acetate can act as buffer solution.
 (a) TTFTT (b) FTTTF (c) TFTFT (d) FFTTT
- 116.** A 1 litre solution containing NH_4Cl and NH_4OH has hydroxide ion concentration of 10^{-6} mol/litre. Which of the following hydroxides could be precipitated when the solution is added to 1 litre solution of 0.1 M metal ions?
 (I) $\text{Ba}(\text{OH})_2$ ($K_{sp} = 5 \times 10^{-3}$) (II) $\text{Ni}(\text{OH})_2$ ($K_{sp} = 1.6 \times 10^{-16}$)
 (III) $\text{Mn}(\text{OH})_2$ ($K_{sp} = 2 \times 10^{-13}$) (IV) $\text{Fe}(\text{OH})_2$ ($K_{sp} = 8 \times 10^{-16}$)
 (a) I, II, IV (b) IV (c) II and IV (d) II, III, IV

- 117.** 150 mL of 0.0008 M ammonium sulphate is mixed with 50 mL of 0.04 M calcium nitrate. The ionic product of CaSO_4 will be : ($K_{sp} = 2.4 \times 10^{-5}$ for CaSO_4)
- (a) K_{sp} (b) K_{sp} (c) K_{sp} (d) None of these
- 118.** In a saturated solution of AgCl , NaCl is added gradually. The concentration of Ag^+ is plotted against the concentration of Cl^- . The graph appears as :
- (a)  (b)  (c)  (d) 
- 119.** K_{sp} of AgCl is 1×10^{-10} . Its solubility in 0.1 M KNO_3 will be :
- (a) 10^{-5} moles/litre (b) 10^{-5} moles/litre (c) 10^{-5} moles/litre (d) None of these
- 120.** 50 mL of a solution containing 10^{-3} mole of Ag^+ is mixed with 50 mL of a 0.1 M HCl solution. How much Ag^+ remains in solution? (K_{sp} of $\text{AgCl} = 1.0 \times 10^{-10}$)
- (a) 2.5×10^{-9} (b) 2.5×10^{-7} (c) 2.5×10^{-8} (d) 2.5×10^{-10}
- 121.** The solubility of a sparingly soluble salt A_xB_y in water is 'S' moles per litre. The solubility product has the value :
- (a) S^2 (b) $x^y y^x S^{x+y}$ (c) $x^x y^y S^{x+y}$ (d) S^{x+y}
- 122.** What is the molarity of a saturated solution of CaCO_3 ? ($K_{sp} = 2.8 \times 10^{-9}$)
- (a) 2.6×10^{-5} (b) 2.8×10^{-9} (c) 5.2×10^{-5} (d) 5.6×10^{-9}
- 123.** K_{sp} of $\text{Zr}_3(\text{PO}_4)_4$ in terms of solubility (S) is :
- (a) $108 S^7$ (b) $4 S^3$ (c) $6912 S^7$ (d) None of these
- 124.** The solubility of electrolytes MX_1 , MX_2 and MX_3 is 1×10^{-3} moles per litre. Hence their respective solubility products are :
- (a) 10^{-6} , 4×10^{-9} , 27×10^{-12} (b) 10^{-9} , 4×10^{-9} , 32×10^{-12}
- (c) 10^{-9} , 8×10^{-8} , 32×10^{-12} (d) None of these
- 125.** A saturated solution of $\text{Ca}_3(\text{PO}_4)_2$ has $[\text{Ca}^{2+}] = 2 \times 10^{-8} \text{ M}$ and $[\text{PO}_4^{3-}] = 1.6 \times 10^{-5} \text{ M}$. K_{sp} of $\text{Ca}_3(\text{PO}_4)_2$ is :
- (a) 3.2×10^{-13} (b) 3.2×10^{-34} (c) 2.048×10^{-33} (d) None of these
- 126.** Which of the following is most soluble in water?
- (a) $\text{Ba}_3(\text{PO}_4)_2$ ($K_{sp} = 6 \times 10^{-39}$) (b) ZnS ($K_{sp} = 7 \times 10^{-16}$)
- (c) $\text{Fe}(\text{OH})_3$ ($K_{sp} = 6 \times 10^{-38}$) (d) $\text{Ag}_3(\text{PO}_4)$ ($K_{sp} = 1.8 \times 10^{-18}$)
- 127.** Silver ions are added to a solution with $[\text{Br}^-] = [\text{Cl}^-] = [\text{CO}_3^{2-}] = [\text{AsO}_4^{3-}] = 0.1 \text{ M}$. Which compound will precipitate with lowest $[\text{Ag}^+]$?
- (a) AgBr ($K_{sp} = 5 \times 10^{-13}$) (b) AgCl ($K_{sp} = 1.8 \times 10^{-10}$)
- (c) Ag_2CO_3 ($K_{sp} = 8.1 \times 10^{-12}$) (d) Ag_3AsO_4 ($K_{sp} = 1 \times 10^{-22}$)

128. The solubility product of different sparingly soluble salts are given as under :

S. No.	Formula Type	Solubility product
1.	AB	4.0×10^{-20}
2.	A_2B	3.2×10^{-11}
3.	AB_3	2.7×10^{-31}

The correct increasing order of solubility is :

- (a) 1, 3, 2 (b) 2, 1, 3 (c) 1, 2, 3 (d) 3, 1, 2

129. If K_{sp} for $HgSO_4$ is 6.4×10^{-5} , then solubility of this substance in mole per m^3 is :

- (a) 8×10^{-3} (b) 6.4×10^{-5} (c) 8×10^{-6} (d) None of these

130. The solubility of $Ba_3(AsO_4)_2$ (formula mass 690) is 6.9×10^{-2} g/100 mL. What is the K_{sp} ?

- (a) 1.08×10^{-11} (b) 1.08×10^{-13} (c) 1.0×10^{-15} (d) 6.0×10^{-13}

131. The solubility of $AgBrO_3$ (formula mass 236) is 0.0072 g in 1000 mL. What is the K_{sp} ?

- (a) 2.2×10^{-8} (b) 3.0×10^{-10} (c) 3.0×10^{-5} (d) 9.3×10^{-10}

132. The solubility of PbF_2 (formula mass 245) is 0.46 g/L. What is the solubility product?

- (a) 1.1×10^{-10} (b) 2.6×10^{-8} (c) 1.1×10^{-7} (d) 6.8×10^{-9}

133. How many grams of MgC_2O_4 (formula mass 112) will dissolve in 1.5 L of water? ($K_{sp} = 8.1 \times 10^{-5}$)

- (a) 1.0 (b) 1.29 (c) 1.512 (d) 4.65

134. What is the molarity of F^- ions in a saturated solution of BaF_2 ? ($K_{sp} = 1.0 \times 10^{-6}$)

- (a) 1.0×10^{-2} (b) 1.0×10^{-3} (c) 1.26×10^{-2} (d) 6.3×10^{-3}

135. What is the molarity of F^- in a saturated solution of InF_3 ? ($K_{sp} = 7.9 \times 10^{-10}$)

- (a) 2.3×10^{-3} (b) 8.3×10^{-3} (c) 1.0×10^{-3} (d) 7.0×10^{-3}

136. What is the pH of a saturated solution of $Cu(OH)_2$? ($K_{sp} = 2.6 \times 10^{-19}$)

- (a) 6.1 (b) 7.30 (c) 8.42 (d) 7.90

137. The solubility product of $AgCl$ is $10^{-10} M^2$. The minimum volume (in m^3) of water required to dissolve 14.35 mg of $AgCl$ is approximately :

- (a) 0.01 (b) 0.1 (c) 100 (d) 10

138. What is the molar solubility of $Fe(OH)_2$ ($K_{sp} = 8.0 \times 10^{-16}$) at pH 13.0?

- (a) 8.0×10^{-18} (b) 8.0×10^{-15} (c) 8.0×10^{-17} (d) 8.0×10^{-14}

139. What is the minimum pH necessary to cause a precipitate of $Pb(OH)_2$ ($K_{sp} = 1.2 \times 10^{-5}$) to form in a 0.12 M $PbCl_2$ solution?

- (a) 12.4 (b) 10.8 (c) 12.0 (d) 11.1

140. Which of the following would increase the solubility of $Pb(OH)_2$?

- (a) Add hydrochloric acid
 (b) Add a solution of $Pb(NO_3)_2$
 (c) Add a solution of NaOH
 (d) None of the above—the solubility of a compound is constant at constant temperature

- 141.** What is the molar solubility of Ag_2CO_3 ($K_{sp} = 4 \times 10^{-13}$) in $0.1 \text{ M Na}_2\text{CO}_3$ solution?
 (a) 10^{-6} (b) 10^{-7} (c) 2×10^{-6} (d) 2×10^{-7}
- 142.** What is the concentration of Pb^{2+} when PbSO_4 ($K_{sp} = 1.8 \times 10^{-8}$) begins to precipitate from a solution that is 0.0045 M in SO_4^{2-} ?
 (a) $4.0 \times 10^{-8} \text{ M}$ (b) $1.0 \times 10^{-6} \text{ M}$ (c) $2.0 \times 10^{-8} \text{ M}$ (d) $4.0 \times 10^{-6} \text{ M}$
- 143.** What is the concentration of Ba^{2+} when BaF_2 ($K_{sp} = 1.0 \times 10^{-6}$) begins to precipitate from a solution that is 0.30 M F^- ?
 (a) 9.0×10^{-7} (b) 3.3×10^{-5} (c) 1.1×10^{-5} (d) 3.0×10^{-7}
- 144.** Solubility of AgCl in 0.2 M NaCl is x and that in 0.1 M AgNO_3 is y then which of the following is correct?
 (a) $x > y$ (b) $x < y$ (c) $x = y$ (d) We cannot predict
- 145.** What is the molarity of $\text{Fe}(\text{CN})_6^{4-}$ in a saturated solution of $\text{Ag}_4[\text{Fe}(\text{CN})_6]$? ($K_{sp} = 1.6 \times 10^{-41}$)
 (a) 1.6×10^{-8} (b) 5.2×10^{-8} (c) 2.0×10^{-8} (d) 2.3×10^{-9}
- 146.** At 25°C , K_{sp} for PbBr_2 is equal to 8×10^{-5} . If the salt is 80% dissociated, what is the solubility of PbBr_2 in mol/litre?
 (a) $\frac{10^{-4}}{1.6 \times 1.6}^{1/3}$ (b) $\frac{10^{-5}}{1.6 \times 1.6}^{1/3}$ (c) $\frac{10^{-4}}{0.8 \times 0.8}^{1/3}$ (d) $\frac{10^{-5}}{1.6 \times 1.6}^{1/2}$
- 147.** What is the molar solubility of $\text{Mn}(\text{OH})_2$ ($K_{sp} = 4.5 \times 10^{-14}$) in a buffer solution containing equal amounts of NH_4^+ and NH_3 ($K_b = 1.8 \times 10^{-5}$)?
 (a) 3.0×10^{-4} (b) 1.38×10^{-4} (c) 1.38×10^{-3} (d) 7.3×10^{-4}
- 148.** Find moles of NH_4Cl required to prevent $\text{Mg}(\text{OH})_2$ from precipitating in a litre of solution which contains 0.02 mole of NH_3 and 0.001 mole of Mg^{2+} ions.
 Given : $K_b(\text{NH}_3) = 10^{-5}$; $K_{sp}[\text{Mg}(\text{OH})_2] = 10^{-11}$.
 (a) 10^{-4} (b) 2×10^{-3} (c) 0.02 (d) 0.1
- 149.** What mass of AgI will dissolve in 1.0 L of 1.0 M NH_3 ? Neglect change in conc. of NH_3 .
 [Given : $K_{sp}(\text{AgI}) = 1.5 \times 10^{-16}$; $K_f[\text{Ag}(\text{NH}_3)_2] = 1.6 \times 10^7$]; (At. mass $\text{Ag} = 108$; $\text{I} = 127$)
 (a) $4.9 \times 10^{-5} \text{ g}$ (b) 0.0056 g (c) 0.035 g (d) 0.011 g
- 150.** Consider the following statement and select correct option :
 (I) K_{sp} of $\text{Fe}(\text{OH})_3$ in aqueous solution is 3.8×10^{-38} at 298 K . The concentration of Fe^{3+} will increase when $[\text{H}^+]$ ion concentration decreases.
 (II) In a mixture of NH_4Cl and NH_4OH in water, a further amount of NH_4Cl is added. The pH of the mixture will decrease.
 (III) An aqueous solution of each of the following salts (NH_4I , HCOOK) will be basic, acidic respectively.
 (a) only I is correct (b) only II is correct
 (c) only III is correct (d) II and III are correct



Level 2

- Equilibrium constants of T_2O (T or 3H is an isotope of 1H) and H_2O are different at 298 K. Let at 298 K pure T_2O has pT (like pH) is 7.62. The pT of a solution prepared by adding 10 mL of 0.2 M TCl to 15 mL of 0.25 M NaOT is :
 (a) $2 \log 7$ (b) $14 \log 7$ (c) $13.24 \log 7$ (d) $13.24 \log 7$
- Liquid NH_3 dissociates to a slight extent. At a certain temp. its self dissociation constant $K_{SDC}(NH_3) = 10^{-30}$. The number of NH_4^+ ions are present per 100 cm^3 of pure liquid are :
 (a) 10^{15} (b) 6.022×10^8 (c) 6.022×10^7 (d) 6.022×10^6
- To what volume of 10 litre of 0.5 M CH_3COOH ($K_a = 1.8 \times 10^{-5}$) be diluted in order to double the hydroxide ion concentration :
 (a) 20 L (b) 30 L (c) 40 L (d) None of these
- 20 mL of 0.1 M weak acid HA ($K_a = 10^{-5}$) is mixed with solution of 10 mL of 0.3 M HCl and 10 mL of 0.1 M NaOH. Find the value of $[A^-]/([HA] + [A^-])$ in the resulting solution :
 (a) 2×10^{-4} (b) 2×10^{-5} (c) 2×10^{-3} (d) 0.05
- What concentration of FCH_2COOH ($K_a = 2.6 \times 10^{-3}$) is needed so that $[H^+] = 2 \times 10^{-3}$?
 (a) $2 \times 10^{-3} \text{ M}$ (b) $2.6 \times 10^{-3} \text{ M}$ (c) $5.2 \times 10^{-3} \text{ M}$ (d) $3.53 \times 10^{-3} \text{ M}$
- Calculate the ratio of $[HCOO^-]$ and $[F^-]$ in a mixture of 0.2 M $HCOOH$ ($K_a = 2 \times 10^{-4}$) and 0.1 M HF ($K_a = 6.6 \times 10^{-4}$) :
 (a) 1 : 6.6 (b) 1 : 3.3 (c) 2 : 3.3 (d) 3.3 : 2
- If first dissociation of $X(OH)_3$ is 100% where as second dissociation is 50% and third dissociation is negligible then the pH of $4 \times 10^{-3} \text{ M } X(OH)_3$ is :
 (a) 11.78 (b) 10.78 (c) 2.5 (d) 2.22
- H_3A is a weak triprotic acid ($K_{a1} = 10^{-5}$, $K_{a2} = 10^{-9}$, $K_{a3} = 10^{-13}$)
 What is the value of pX of 0.1 M H_3A (aq.) solution? where $pX = -\log X$ and $X = \frac{[A^{3-}]}{[HA^2-]}$
 (a) 7 (b) 8 (c) 9 (d) 10
- Calcium lactate is a salt of weak organic acid and strong base represented as $Ca(LaC)_2$. A saturated solution of $Ca(LaC)_2$ contains 0.6 mole in 2 litre solution. pOH of solution is 5.60. If 90% dissociation of the salt takes place then what is pK_a of lactic acid?
 (a) $2.8 \log(0.54)$ (b) $2.8 \log(0.54)$ (c) $2.8 \log(0.27)$ (d) None of these
- What is the concentration of CH_3COOH (aq.) in a solution prepared by dissolving 0.01 mole of $NH_4^+CH_3COO^-$ in 1 L H_2O ? [$K_a(CH_3COOH) = 1.8 \times 10^{-5}$; $K_b(NH_4OH) = 1.8 \times 10^{-5}$]
 (a) 5.55×10^{-5} (b) 0.10 (c) 6.4×10^{-4} (d) 5.55×10^{-3}
- K_a for the reaction; $Fe^{3+}(\text{aq.}) + H_2O(l) \rightleftharpoons Fe(OH)^{2+}(\text{aq.}) + H_3O^+(\text{aq.})$ is 6.5×10^{-3} . What is the max. pH value which could be used so that at least 80% of the total iron (III) in a dilute solution exists as Fe^{3+} ?
 (a) 2 (b) 2.41 (c) 2.79 (d) 1.59

12. $\text{Fe}(\text{OH})_2$ is diacidic base has $K_{b1} = 10^{-4}$ and $K_{b2} = 2.5 \times 10^{-6}$





What is the concentration of $\text{Fe}(\text{OH})_2$ in 0.1 M $\text{Fe}(\text{NO}_3)_2$ solution?

- (a) 4×10^{-9} (b) 2.5×10^{-6} (c) 10^{-10} (d) 10^{-14}
13. How many gm of solid KOH must be added to 100 mL of a buffer solution to make the pH of solution 6.0, if it is 0.1 M each w.r.t. acid HA and salt K A.
[Given : $\text{p}K_a(\text{HA}) = 5$]
(a) 0.458 (b) 0.327 (c) 5.19 (d) None of these
14. Fixed volume of 0.1 M benzoic acid ($\text{p}K_a = 4.2$) solution is added into 0.2 M sodium benzoate solution and formed a 300 mL, resultant acidic buffer solution. If pH of this buffer solution is 4.5 then find added volume of benzoic acid :
(a) 100 mL (b) 150 mL (c) 200 mL (d) None of these
15. A 1.025 g sample containing a weak acid HX (mol. mass = 82) is dissolved in 60 mL water and titrated with 0.25 M NaOH. When half of the acid was neutralised the pH was found to be 5.0 and at the equivalence point the pH is 9.0. Calculate mass percentage of HX in sample :
(a) 50% (b) 75% (c) 80% (d) None of these
16. Which of the following expression for % dissociation of a monoacidic base (BOH) in aqueous solution at appreciable concentration is not correct?
(a) $100 \sqrt{\frac{K_b}{c}}$ (b) $\frac{1}{1 + 10^{(\text{p}K_b - \text{pOH})}}$
(c) $\frac{K_w [\text{H}^+]}{K_b + K_w}$ (d) $\frac{K_b}{K_b + [\text{OH}^-]}$
17. A solution of weak acid HA was titrated with base NaOH. The equivalent point was reached when 40 mL of 0.1 M NaOH has been added. Now 20 mL of 0.1 M HCl were added to titrated solution, the pH was found to be 5.0. What will be the pH of the solution obtained by mixing 20 mL of 0.2 M NaOH and 20 mL of 0.2 M HA?
(a) 7 (b) 9 (c) 11 (d) None of these
18. A buffer solution 0.04 M in Na_2HPO_4 and 0.02 M in Na_3PO_4 is prepared. The electrolytic oxidation of 1.0 milli-mole of the organic compound RNHOH is carried out in 100 mL of the buffer. The reaction is
$$\text{RNHOH} + \text{H}_2\text{O} \rightarrow \text{RNO}_2 + 4\text{H}^+ + 4\text{e}^-$$

The approximate pH of solution after the oxidation is complete is :
[Given : for H_3PO_4 , $\text{p}K_{a1} = 2.2$; $\text{p}K_{a2} = 7.20$; $\text{p}K_{a3} = 12$]
(a) 6.90 (b) 7.20 (c) 7.5 (d) None of these
19. When a 20 mL of 0.08 M weak base BOH is titrated with 0.08 M HCl, the pH of the solution at the end point is 5. What will be the pOH if 10 mL of 0.04 M NaOH is added to the resulting solution?
[Given : $\log 2 = 0.30$ and $\log 3 = 0.48$]
(a) 5.40 (b) 5.88 (c) 4.92 (d) None of these

20. Calculate approximate pH of the resultant solution formed by titration of 25 mL of 0.04 M Na_2CO_3 with 50 mL of 0.025 M HCl. [Given : $\text{p}K_{a_1}$ 6.4 and $\text{p}K_{a_2}$ 10.3 for H_2CO_3]
- (a) 5.92 (b) 6.88 (c) 6.4 (d) 5.88
21. In the titration of a solution of a weak acid HA and NaOH, the pH is 5.0 after 10 mL of NaOH solution has been added and 5.60 after 20 mL NaOH has been added. What is the value of $\text{p}K_a$ for HA?
- (a) 5.15 (b) 5.3 (c) 5.6 (d) None of these
22. 50 mL of 0.05 M Na_2CO_3 is titrated against 0.1 M HCl. On adding 40 mL of HCl, pH of the solution will be [Given : For H_2CO_3 , $\text{p}K_{a_1}$ 6.35, $\text{p}K_{a_2}$ 10.33; $\log 3$ 0.477, $\log 2$ 0.30]
- (a) 6.35 (b) 6.526 (c) 8.34 (d) 6.173
23. 10 mL of 0.1 M tribasic acid H_3A is titrated with 0.1 M NaOH solution. What is the ratio of $\frac{[\text{H}_3\text{A}]}{[\text{A}^{3-}]}$ at 2nd equivalence point? [Given : K_{a_1} 10^{-3} , K_{a_2} 10^{-8} , K_{a_3} 10^{-12}]
- (a) $\sim 10^{-4}$ (b) $\sim 10^{-4}$ (c) $\sim 10^{-7}$ (d) $\sim 10^{-6}$
24. A_3B_2 is a sparingly soluble salt of molar mass M (g mol^{-1}) and solubility x g litre^{-1} . The ratio of the molar concentration of B^{3-} to the solubility product of the salt is :
- (a) $108 \frac{x^5}{M^5}$ (b) $\frac{1}{108} \frac{M^4}{x^4}$ (c) $\frac{1}{54} \frac{M^4}{x^4}$ (d) None of these
25. A solution is 0.10 M $\text{Ba}(\text{NO}_3)_2$ and 0.10 M $\text{Sr}(\text{NO}_3)_2$. If solid Na_2CrO_4 is added to the solution, what is $[\text{Ba}^{2+}]$ when SrCrO_4 begins to precipitate?
- $[K_{sp}(\text{BaCrO}_4) 1.2 \times 10^{-10}; K_{sp}(\text{SrCrO}_4) 3.5 \times 10^{-5}]$
- (a) 7.4×10^{-7} (b) 2.0×10^{-7} (c) 6.1×10^{-7} (d) 3.4×10^{-7}
26. A solution is 0.01 M KI and 0.1 M KCl. If solid AgNO_3 is added to the solution, what is the $[\text{I}^-]$ when AgCl begins to precipitate?
- $[K_{sp}(\text{AgI}) 1.5 \times 10^{-16}; K_{sp}(\text{AgCl}) 1.8 \times 10^{-10}]$
- (a) 3.5×10^{-7} (b) 6.1×10^{-8} (c) 2.2×10^{-7} (d) 8.3×10^{-8}
27. A solution of 0.1 M in Cl^- and 10^{-4} M CrO_4^{2-} . If solid AgNO_3 is gradually added to this solution, what will be the concentration of Cl^- when Ag_2CrO_4 begins to precipitate?
- $[K_{sp}(\text{AgCl}) 10^{-10} \text{ M}^2; K_{sp}(\text{Ag}_2\text{CrO}_4) 10^{-12} \text{ M}^3]$
- (a) 10^{-6} M (b) 10^{-4} M (c) 10^{-5} M (d) 10^{-9} M
28. If 500 mL of 0.4 M AgNO_3 is mixed with 500 mL of 2 M NH_3 solution then what is the concentration of $\text{Ag}(\text{NH}_3)^+$ in solution?
- Given : $K_{f_1}[\text{Ag}(\text{NH}_3)] 10^3; K_{f_2}[\text{Ag}(\text{NH}_3)_2] 10^4$
- (a) $3.33 \times 10^{-7} \text{ M}$ (b) $3.33 \times 10^{-5} \text{ M}$ (c) $3 \times 10^{-4} \text{ M}$ (d) 10^{-7} M
29. The simultaneous solubility of AgCN ($K_{sp} 2.5 \times 10^{-16}$) and AgCl ($K_{sp} 1.6 \times 10^{-10}$) in 1.0 M $\text{NH}_3(aq.)$ are respectively : [Given : $K_f[\text{Ag}(\text{NH}_3)_2] 10^7$]
- (a) 0.037, 5.78×10^{-8} (b) 5.78×10^{-8} , 0.037
(c) 0.04, 6.25×10^{-8} (d) 1.58×10^{-3} , 1.26×10^{-5}

30. There exist an equilibrium between solid SrSO_4 and Sr^{2+} and SO_4^{2-} ion in aqueous medium. The possible equilibrium states are shown in figure as thick line. Now, if equilibrium is disturbed by addition of (a) $\text{Sr}(\text{NO}_3)_2$ and (b) K_2SO_4 and dotted line represent approach of system towards equilibrium. Match the columns given below :



(I) addition of $\text{Sr}(\text{NO}_3)_2$

(II) addition of K_2SO_4

(a) (I) (iii), (II) (iv)

(b) (I) (iv), (II) (v)

(c) (I) (vi), (II) (v)

(d) (I) (iv), (II) (vi)

31. Solubility of AgCN is maximum in :

(a) acidic buffer solution

(b) basic buffer solution

(c) in pure water

(d) equal in all solution

32. $\text{AgBr}(s) + 2\text{S}_2\text{O}_3^{2-}(aq.) \rightleftharpoons \text{Ag}(\text{S}_2\text{O}_3)_2^{3-}(aq.) + \text{Br}^-(aq.)$

[Using : $K_{sp}(\text{AgBr}) = 5 \times 10^{-13}$, $K_f(\text{Ag}(\text{S}_2\text{O}_3)_2^{3-}) = 5 \times 10^{13}$]

What is the molar solubility of AgBr in $0.1 \text{ M Na}_2\text{S}_2\text{O}_3$?

(a) 0.5 M

(b) 0.45 M

(c) 0.045 M

(d) None of these

33. What is $[\text{Ag}^+]$ in a solution made by dissolving both Ag_2CrO_4 and $\text{Ag}_2\text{C}_2\text{O}_4$ until saturation is reached with respect to both salts?

$[K_{sp}(\text{Ag}_2\text{C}_2\text{O}_4) = 2 \times 10^{-11}, K_{sp}(\text{Ag}_2\text{CrO}_4) = 2 \times 10^{-12}]$

(a) 2.80×10^{-4}

(b) 7.6×10^{-5}

(c) 6.63×10^{-6}

(d) 3.52×10^{-4}

34. What is the minimum pH required to prevent the precipitation of ZnS in a solution that is 0.01 M ZnCl_2 and saturated with $0.10 \text{ M H}_2\text{S}$?

[Given : $K_{sp} = 10^{-21}$, $K_{a1} = K_{a2} = 10^{-20}$]



(a) 0

(b) 1

(c) 2

(d) 4

35. The $\text{Al}(\text{OH})_3$ is involved in the following two equilibria,



Which of the following relationship is correct at which solubility is minimum?

(a) $[\text{OH}^-] = \frac{K_{sp}^{1/3}}{K_c}$

(b) $[\text{OH}^-] = \frac{K_c}{K_{sp}^{1/4}}$

(c) $[\text{OH}^-] = \sqrt{\frac{K_{sp}}{K_c}}$

(d) None of these



Level 3



Passage 1

Solution of a weak acid and its anion (that is, its conjugate base) or of a base and its common cation are buffered. When we add a small amount of acid or base to any one of them, the pH of solution changes very little. pH of buffer solution can be computed as

$$\text{for acidic buffer : } \text{pH} = \text{p}K_a + \log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$$

$$\text{for basic buffer : } \text{pOH} = \text{p}K_b + \log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$$

It is generally accepted that a solution has useful buffer capacity (pH change resisting power) provided that the value of [salt or conjugate base]/[acid] for acidic buffer lies within the range of 1 : 10 to 10 : 1. Buffer capacity is maximum when [conjugate base] [acid]

- One litre of an aqueous solution contains 0.15 mole of CH_3COOH ($\text{p}K_a = 4.8$) and 0.15 mole of CH_3COONa . After the addition of 0.05 mole of solid NaOH to this solution, the pH will be :
(a) 4.5 (b) 4.8 (c) 5.1 (d) 5.4
- Calculate the pH of a solution made by adding 0.01 mole of HCl in 100 mL of a solution which is 0.2 M in NH_3 ($\text{p}K_b = 4.74$) and 0.3 M in NH_4^+ :
(Assuming no change in volume)
(a) 5.34 (b) 8.66 (c) 7.46 (d) None of these
- Useful buffer range of weak acid HA ($K_a = 10^{-5}$) is :
(a) 5 to 7 (b) 4 to 6 (c) 3 to 5 (d) None of these
- Select correct statement :
(a) When we add small amount of NaOH in acidic buffer solution, pOH of solution is increases
(b) When we add small amount of NaOH in basic buffer solution, pH of solution is increases
(c) When we add small amount of water in acidic buffer solution, pH of solution is decreases
(d) When 100 mL of 0.2 M CH_3COOH react with 200 mL of 0.1 M NaOH buffer solution is formed



Passage 2

Hydrolysis is an acid-base reaction of a cation or anion or both ions of a salt with water. Resultant solution of hydrolysis may be acidic, basic or neutral. The anion A^- which is a weaker base than OH^- and which has its conjugate acid HA stronger than water but weaker than H_3O^+ shows the phenomenon of hydrolysis Ex. : CH_3COO^- , CN^- , NO_2^- , etc.

The cation B^+ which is a weaker acid than H_3O^+ and which has its conjugate base BOH stronger than water but weaker than OH^- shows the phenomenon of hydrolysis Ex. : NH_4^+ , $\text{C}_6\text{H}_5\text{NH}^+$, N_2H_5^+ etc.

The hydrolysis constant of anion and cation are given by

$$A^-(aq.) + H_2O(l) \rightleftharpoons HA(aq.) + OH^-(aq.)$$

$$K_h = \frac{K_w}{K_a} = \frac{[HA(aq.)][OH^-(aq.)]}{[A^-(aq.)]}$$

$$B^+(aq.) + H_2O(l) \rightleftharpoons BOH(aq.) + H^+(aq.)$$


$$K_h = \frac{K_w}{K_b} = \frac{[BOH(aq.)][H^+(aq.)]}{[B^+(aq.)]}$$

- Which of the following statement is true?
 - Weaker the acid, greater will be hydrolysis of its anion
 - Weaker the base, greater will be hydrolysis of its cation
 - Both (a) and (b)
 - None of these
- Select the correct statement :
 - KCl undergoes hydrolysis
 - $K_h = K_b(A^-)$ and $K_h = K_a(B^+)$
 - 0.1 M solution of NaCN is acidic
 - resultant solution of equal volume of 0.1 M NH_3 and 0.1 M HCl is basic
- When pure ammonium chloride is dissolved in pure water, the pH of the resulting solution is not 7. This is because :
 - ammonium ions accept protons from water molecules leaving free OH^- ions in solution
 - ammonium ions donate protons to water molecules forming H_3O^+ ions in solution
 - ammonium ions combine with water molecule to give the weak base, ammonium hydroxide
 - chloride ion made the solution acidic
- Calculate percentage degree of hydrolysis in a 0.1 M solution of CH_3COONa .
 $(K_a \text{ of } CH_3COOH = 10^{-5})$
 - 0.1
 - 0.01
 - 10^{-4}
 - None of these



Passage 3

Acid-base indicators are either weak organic acids or weak organic bases. Indicator change colour in dilute solution when the hydronium ion concentration reaches a particular value. For example, phenolphthalein is a colourless substance in any aqueous solution with a pH less than 8.3. In between the pH range 8.3 to 10, transition of colour (colourless to pink) takes place and if pH of solution is greater than 10 solution is dark pink. Considering an acid indicator HIn, the equilibrium involving it and its conjugate base In^- can be represented as :




pH of solution can be computed as :

$$pH = pK_{In} + \log \frac{[In^-]}{[HIn]}$$

In general, transition of colour takes place in between the pH range $pK_{In} \pm 1$.

- An indicator is a weak acid and pH range is 4.0 to 6.0. If indicator is 50% ionized in a given solution then what is the ionization constant of the acid?
 (a) 10^{-4} (b) 10^{-5} (c) 10^{-6} (d) None of these
- Select the correct statement(s) :
 (a) At midway in the transition of an acidic indicator, $\text{pH} = \text{p}K_{\text{In}}$
 (b) Methyl orange (3.1 to 4.4) is a suitable indicator for titration of weak acid and strong base
 (c) Bromothymol blue (6.0 to 7.6) is a good indicator for titration of HCl and NaOH
 (d) Thymol blue (1.2 to 2.8) is a very good indicator for titration of 100 mL of 0.1 M NH_4OH ($\text{p}K_b = 4.74$) and 0.1 M HCl
- Following is the titration curve of two acids HA and HB (5 milli-moles each) titrated against strong base NaOH (0.1 M)



What is $\text{p}K_a$ for HB acid ?


- What is the equilibrium constant for the reaction :

$$\text{HB}(\text{aq.}) + \text{NaA}(\text{aq.}) \rightleftharpoons \text{HA}(\text{aq.}) + \text{NaB}(\text{aq.})?$$
 (a) 10 (b) 0.1 (c) 10^{-7} (d) 10^{-11}
- Calculate the pH at equivalent point when HB is titrated with NaOH.
 (a) 8.75 (b) 8.85 (c) 9.0 (d) None of these
- Which of the following indicator is most suitable for titration of HB with strong base :
 (a) Phenolphthalein (8.3 – 10) (b) Bromothymol blue (6 – 7.6)
 (c) Methyl red (4.2 – 6.3) (d) Malachite green (11.4 – 13)



Passage 4

Consider a saturated solution of silver chloride that is in contact with solid silver chloride. The solubility equilibrium can be represented as



Where K_{sp} is called the solubility product constant or simply the solubility product. In general, the solubility product of a compound is the product of the molar concentrations of the constituent ions, each raised to the power of its stoichiometric coefficient in the equilibrium equation.

For concentrations of ions that do not necessarily correspond to equilibrium conditions we use the reaction quotient (Q) which is called the ion or ionic product (Q), to predict whether a precipitate will form. Note that Q has the same form as K_{sp} .

The possible relationships between Q and K_{sp} are

$Q < K_{sp}$	Unsaturated solution
$Q = K_{sp}$	Saturated solution
$Q > K_{sp}$	Supersaturated solution; precipitate will form

- Will a precipitate form if 50 cm^3 of 0.01 M AgNO_3 and 50 cm^3 of $2 \times 10^{-5} \text{ M NaCl}$ are mixed?
[Given : $K_{sp}(\text{AgCl}) = 10^{-10} \text{ M}^2$]
 (a) Yes
 (b) No
 (c) Ionic product is less than solubility product, hence precipitate will form
 (d) Data insufficient
- Will a precipitate form if 1 volume of 0.1 M Pb^{2+} ion solution is mixed with 3 volume of 0.3 M Cl^- ion solution? [Given : $K_{sp}(\text{PbCl}_2) = 1.7 \times 10^{-5} \text{ M}^3$]
 (a) Yes
 (b) No
 (c) Ionic product is less than solubility product, hence precipitate will form
 (d) Data insufficient
- At 25°C , will a precipitate of $\text{Mg}(\text{OH})_2$ form when a 0.0001 M solution of $\text{Mg}(\text{NO}_3)_2$ is adjusted to a pH of 9.0? At what minimum value of pH will precipitation start?
[Given : $K_{sp}(\text{Mg}(\text{OH})_2) = 10^{-11} \text{ M}^3$]
 (a) No, pH 3.5 (b) No, pH 10.5 (c) No, pH 6.0 (d) Yes, pH 8.5
- Determine the molar solubility of MgF_2 from its solubility product $K_{sp} = 4 \times 10^{-9}$:
 (a) 10^{-3} (b) 6.32×10^{-5} (c) 2×10^{-5} (d) None of these
- The solubility product of ferric hydroxide in aqueous solution is 6×10^{-38} at 298 K . The solubility of Fe^{3+} ion will increase when the :
 (a) pH is increased
 (b) pH is 7.0
 (c) pH is decreased
 (d) saturated solution is exposed to the atmosphere

One or More Answers is/are Correct



- Which is/are wrong statement(s)?
 - Arrhenius acids are also Bronsted acids but all Arrhenius bases are not Bronsted bases
 - All Lewis bases are Bronsted bases
 - All Bronsted acids are Lewis acids
 - Conjugate base of a strong acid is weak
- Which of the following are conjugate acid-base pairs ?

(a) HCO_3^- , CO_3^{2-}	(b) $\text{C}_6\text{H}_5\text{NH}_3^+$, $\text{C}_6\text{H}_5\text{NH}_2$
(c) $\text{H}_2\text{C}_2\text{O}_4$, HC_2O_4^-	(d) OH^- , H^+
- Which are the set of amphoteric species?

(a) H_2O , H_2PO_4^- , HPO_4^{2-}	(b) HPO_4^{2-} , HCO_3^-
(c) H_2PO_2^- , H_2PO_3^- , HC_2O_4^-	(d) HPO_3^{2-} , H_2O , CO_3^{2-}
- Which of the following statements is/are not correct?
 - A substance which can provide OH^- in aqueous medium is a base
 - A substance which can accept a pair of electron is a base
 - A substance which can accept a proton in aqueous medium is a base
 - A substance which can donate a pair of electron is a base
- If degree of ionization (α) of a weak electrolyte AB is very less then α is :
 - directly proportional to the square root of volume of solution
 - inversely proportional to the dilution
 - inversely proportional to the square root of concentration
 - directly proportional to concentration
- Factor influencing the degree of ionization of a weak electrolyte is :

(a) dilution	(b) temperature
(c) presence of other ions	(d) nature of solvent
- Which of the following statement(s) is/are correct about the ionic product of water?
 - K_i (ionization constant of water) $< K_w$ (ionic product of water)
 - $\text{p}K_i = \text{p}K_w$
 - At 25°C , $K_i = 1.8 \times 10^{-14}$
 - Ionic product of water at 10°C is 10^{-14}
- Which among the following statement is/are correct?
 - $\text{pH} = \log_{10} [\text{H}_3\text{O}^+]$ for dilute solution
 - pH of H_2O decreases with increase of temperature
 - pH can not more than 14
 - If a solution is diluted ten times, its pH always increases by 1
- If concentration of two weak acids are different and D.O.D. (α) are very less then their relative strength can be compared by :

(a) $\frac{[\text{H}^+]}{[\text{H}]}$	(b) $\frac{1}{\alpha}$	(c) $\frac{C_1}{C_2}$	(d) $\frac{K_{a1} C_1}{K_{a2} C_2}$
---------------------------------------	------------------------	-----------------------	-------------------------------------

10. If concentration of two weak bases are same and D.O.D. () are very less then their relative strength can be compared by :
- (a) $\frac{[\text{OH}^-]_1}{[\text{OH}^-]_2}$ (b) $\frac{K_{b_1}}{K_{b_2}}$ (c) $\frac{1}{2}$ (d) $\frac{\sqrt{K_{b_1}}}{\sqrt{K_{b_2}}}$
11. Which of the following expressions is/are true?
- (a) $[\text{H}^+] = [\text{OH}^-] = \sqrt{K_w}$ for a neutral solution
 (b) $[\text{OH}^-] = \sqrt{K_w}$ for an acidic solution
 (c) $\text{pH} + \text{pOH} = 14$ at all temperature
 (d) $[\text{OH}^-] = 10^{-7} \text{ M}$ for a neutral solution at 25°C
12. If K_{a_1} , K_{a_2} and K_{a_3} be the first, second and third dissociation constant of H_3PO_4 and $K_{a_1} > K_{a_2} > K_{a_3}$ which is/are correct :
- (a) $[\text{H}^+] \sim \sqrt{K_{a_1} [\text{H}_3\text{PO}_4]}$ (b) $[\text{H}^+] \sim [\text{HPO}_4^{2-}]$
 (c) $K_{a_2} \sim [\text{HPO}_4^{2-}]$ (d) $[\text{HPO}_4^{2-}] = [\text{PO}_4^{3-}]$
13. Which of the following mixtures constitute a buffer?
- (a) $\text{HCOOH} + \text{HCOONa}$ (b) $\text{Na}_2\text{CO}_3 + \text{NaHCO}_3$
 (c) $\text{NaCl} + \text{HCl}$ (d) $\text{NH}_4\text{Cl} + (\text{NH}_4)_2\text{SO}_4$
14. Which of the following mixtures can act as a buffer?
- (a) $\text{NaOH} + \text{HCOONa}$ (1 : 1 molar ratio) (b) $\text{HCOOH} + \text{NaOH}$ (2 : 1 molar ratio)
 (c) $\text{NH}_4\text{Cl} + \text{NaOH}$ (2 : 1 molar ratio) (d) $\text{HCOOH} + \text{NaOH}$ (1 : 1 molar ratio)
15. Which of the following will function as buffer?
- (a) $\text{NaCl} + \text{NaOH}$ (b) Borax + boric acid
 (c) $\text{NaH}_2\text{PO}_4 + \text{Na}_2\text{HPO}_4$ (d) $\text{NH}_4\text{Cl} + \text{NH}_4\text{OH}$
16. Which of the following statements is/are correct?
- (a) The conjugate acid of NH_2^- is NH_3
 (b) Solubility product constant increases with increase in concentration of ions
 (c) On diluting a buffer solution pH change is negligible
 (d) In alkaline buffer solution, if some HCl is added, its $[\text{OH}^-]$ will increase
17. Degree of hydrolysis () for a salt of strong acid and weak base is :
- (a) independent of dilution (b) increases with dilution
 (c) increases with decrease in K_b (d) increases with increase in temperature
18. The compound whose 0.1 M solution is acidic :
- $[\text{p}K_a \text{ of } \text{HCOOH} = 3.75, \text{p}K_b \text{ of } \text{NH}_4\text{OH} = 4.74]$
- (a) Ammonium formate (b) Ammonium sulphate
 (c) Ammonium chloride (d) Sodium formate
19. Formic acid is a weak acid and hydrochloric acid is a strong acid. It shows that the :
- (a) $[\text{OH}^-]$ of 0.01 M HCl (aq.) will be less than that of 0.01 M HCOOH (aq.)
 (b) solution containing 0.1 M NaOH (aq.) and 0.1 M HCOONa (aq.) is a buffer solution
 (c) pH of 10^{-9} M HCl (aq.) will be approximately 7 at 25°C
 (d) pH of a solution formed by mixing equimolar quantities of HCOOH and HCl will be less than that of a similar solution formed from HCOOH and HCOONa

- 20.** If you have a saturated solution of CaF_2 , then :
- (a) $[\text{Ca}^{2+}] = (K_{sp}/4)^{1/3}$ (b) $2 [\text{Ca}^{2+}] = [\text{F}^-]$
 (c) $[\text{Ca}^{2+}] = 2 [\text{F}^-]$ (d) $[\text{Ca}^{2+}] = \sqrt{K_{sp}}$
- 21.** Which is/are correct statement(s) about the solubility of AgCl(s) .?
 Given : $K_{sp}(\text{AgCl}) = 10^{-10}$; $K_f[\text{Ag}(\text{NH}_3)_2] = 10^8$.
- (a) Solubility of AgCl in pure water is 10^{-5} gm/litre
 (b) Solubility of AgCl in 2 M KBr is 10^{-5} mol/litre
 (c) Solubility of AgCl in 2 M AgNO_3 is $5 \times 10^{-11}\text{ M}$
 (d) Solubility of AgCl in 2 M NH_3 is 0.166 M
- 22.** H_2A is a weak diprotic acid. If the pH of 0.1 M H_2A solution is 3 and concentration of A^{2-} is 10^{-12} at 25°C .
 Select correct statement(s)
- (a) $[\text{H}^+]_{\text{total}} = [\text{H}^+]$ from first step of ionization of acid H_2A
 (b) Concentration of OH^- in solution is 10^{-3} M
 (c) The value of K_{a1} is nearly 10^{-5}
 (d) $pK_{a2} - pK_{a1} = 9$
- 23.** Which is/are correct statement(s)?
- (a) $\text{CH}_3\text{COONH}_4$ have greater degree of hydrolysis in 0.2 M solution in comparison of 0.4 M solution.
 (b) Anions which are weaker base than OH^- , do not hydrolyse
 (c) The CH_3COO^- have greater degree of hydrolysis in comparison of HCOO^- when their salt solution have equal conc.
 (d) SO_4^{2-} hydrolyses but HSO_4^- does not undergo hydrolysis
- 24.** 0.01 M NH_4Cl (aq) solution at 25°C has:
- (a) $[\text{Cl}^- (\text{aq})] = 10^{-2}\text{ M}$ (b) $[\text{NH}_4^+ (\text{aq})] = 10^{-2}\text{ M}$
 (c) $\text{pOH} > 7$ (d) $[\text{H}^+] = 10^{-7}\text{ M}$
- 25.** In an acidic indicator HIn has ionization constant is 10^{-8} . The acid form of indicator is yellow and alkaline form is red. Which is correct statement ?
 (Given : $\log 2 = 0.3$, $\log 3 = 0.48$)
- (a) The pH range of indicator is 7 to 9
 (b) Change in pH is 0.96 when 75% yellow colour change to 75% red colour
 (c) This indicator is suitable for the titration of strong acid vs. strong base
 (d) pH of indicator is 8.3 when ratio of acid form to alkaline form is 2.

Match the Column



Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I

- (A) HCl
- (B) NH_3
- (C) H_2O
- (D) CN

Column-II

- (P) Bronsted base
- (Q) Bronsted acid
- (R) Arrhenius acid
- (S) Lewis base in adduct displacement reaction

2.

Column-I

- (A) Conjugate acid-base pair
- (B) Acid-base adduct
- (C) An acid-base reaction
- (D) Proton donation

Column-II

- (P) Bronsted-Lowry concept
- (Q) Lewis concept
- (R) Arrhenius concept
- (S) K_a K_b K_w

3.

Column-I

- (A) $\text{Fe}(\text{NO}_3)_2(aq.)$
- (B) $\text{KClO}_4(aq.)$
- (C) $\text{HCOONa}(aq.)$
- (D) $\text{NH}_4\text{CN}(aq.)$

Column-II

- (P) Only cationic hydrolysis
- (Q) Only anionic hydrolysis
- (R) Both cationic as well as anionic hydrolysis
- (S) No hydrolysis

4.

Column-I

- (A) Salt of weak acid and weak base
- (B) Salt of weak acid and strong base
- (C) Salt of strong acid and strong base
- (D) Salt of strong acid and weak base

Column-II

- (P) $\text{pH} = \frac{1}{2} [\text{p}K_w + \text{p}K_a + \log C]$
- (Q) $\text{pH} = \frac{1}{2} [\text{p}K_w + \text{p}K_a + \text{p}K_b]$
- (R) $\text{pH} = \frac{1}{2} [\text{p}K_w + \text{p}K_b + \log C]$
- (S) $\text{pH} = \frac{1}{2} [\text{p}K_w]$

5.	Column-I	Column-II
	(A) Salt of weak acid and weak base ($pK_a \quad pK_b$)	(P) pH of solution at 25°C less than 7
	(B) Salt of weak acid and strong base	(Q) pH of solution at 25°C greater than 7
	(C) Salt of strong acid and strong base	(R) pH of solution at 25°C equal to 7
	(D) Salt of strong acid and weak base	(S) pH cannot be find until the value of K_a/K_b is given

6.	Column-I	Column-II
	(A) pH of 0.1 M HA ($pK_a \quad 5$) and 0.01 M NaA	(P) 4
	(B) pH of 0.1 M BOH ($pK_b \quad 6$) and 0.1 M BCl	(Q) 7
	(C) pH of 0.1 M salt of weak acid ($pK_a \quad 5$) and weak base ($pK_b \quad 7$)	(R) 6
	(D) pH of 500 litre of 0.02 M HNO_3 and 500 litre 0.01 M $\text{Sr}(\text{OH})_2$	(S) 8

7.	Column-I	Column-II
	(A) CH_3COOH ($pK_a \quad 4.74, 0.1 \text{ M}$) CH_3COONa (0.1 M)	(P) Acidic buffer at its maximum capacity
	(B) CH_3COOH (0.1 M) + HCl (0.1 M)	(Q) Buffer solution
	(C) CH_3COOH ($pK_a \quad 4.74, 0.1 \text{ M}$) + NH_4OH ($pK_b \quad 4.74, 0.1 \text{ M}$)	(R) $\text{pH} < 7$ at 25°C
	(D) CH_3COONa (300 mL of 0.1 M) + HCl (100 mL of 0.1 M)	(S) $\text{pH} \quad 7$ at 25°C

8.	Column-I	Column-II
	(A) Titration of a strong acid with strong base	(P) Methyl orange (3.1 4.4)
	(B) Titration of weak acid with strong base	(Q) Methyl red (4.2 6.3)
	(C) Titration of strong acid with weak base	(R) Phenolphthalein (8.3 10)
	(D) Titration of weak acid with weak base	(S) No general indicator is suitable

9. When we titrate sodium carbonate solution (in beaker) with hydrochloric acid

Column-I	Column-II
(A) At the start of titration	(P) Buffer solution of HCO_3^- and CO_3^{2-}
(B) Before the first equivalent point	(Q) Buffer solution of H_2CO_3 and HCO_3^-
(C) At the first equivalent point	(R) Amphoteric anion, $\text{pH} = \frac{1}{2}(\text{p}K_{a_1} + \text{p}K_{a_2})$
(D) Between the first and second equivalent points	(S) Hydrolysis of CO_3^{2-}

10.

Column-I	Column-II
(A) Mercurous iodide	(P) 108 S^5
(B) Aluminium phosphate	(Q) 4 S^3
(C) Calcium phosphate	(R) S^2
(D) Zirconium phosphate	(S) 6912 S^7

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

- (A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** All strong monoprotic acid with same concentration in dilute solution show same pH.

STATEMENT-2 : Water shows levelling effect.

2. **STATEMENT-1 :** If water is heated to 50°C then pH will increase.

STATEMENT-2 : K_w increases with increase in temperature.

3. **STATEMENT-1 :** Addition of $\text{HCl}(\text{aq.})$ to $\text{HCOOH}(\text{aq.})$ decrease the dissociation of $\text{HCOOH}(\text{aq.})$

STATEMENT-2 : Due to common ion effect of H^+ , dissociation of HCOOH decrease.

4. **STATEMENT-1 :** pH of 10^{-7} M HCl is less than 7 at 25°C .


STATEMENT-2 : At very low concentration of HCl , contribution of H^+ from water is considerable.

- 5. STATEMENT-1 :** The dissociation constants of weak diprotic acid are in the order of $K_{a_1} > K_{a_2}$.
- STATEMENT-2 :** Removal of H⁺ from anion is difficult as compared to neutral molecule.
- 6. STATEMENT-1 :** When 0.1 M weak diprotic acid H₂A dissociates with its dissociation constants $K_{a_1} = 10^{-3}$ and $K_{a_2} = 10^{-8}$, then $[A^{2-}]$ is almost equal to 10^{-3} M.
- STATEMENT-2 :** Since $K_{a_2} \ll K_{a_1}$ for 0.1 M H₂A, so $[A^{2-}]$ is negligible w.r.t. $[HA^-]$.
- 7. STATEMENT-1 :** pH value of acidic buffer solution change, if buffer solution is diluted upto very larger extent.
- STATEMENT-2 :** $[H^+]$ decrease due to change in concentration as well as K_a and decrease in concentration is more as compared to increase in K_a .
- 8. STATEMENT-1 :** In a titration of weak acid with strong base, the pH at the half equivalence point is pK_a .
- STATEMENT-2 :** At half equivalence point, it will form acidic buffer at its maximum capacity where $[acid] = [salt]$.
- 9. STATEMENT-1 :** In the titration of Na₂CO₃ with HCl using methyl orange indicator, the volume of the acid required at the equivalence point is twice that of the acid required using phenolphthalein as indicator.
- STATEMENT-2 :** Two moles of HCl are required for the complete neutralisation of one mole of Na₂CO₃.
- 10. STATEMENT-1 :** In the acid-base titration involving strong base and weak acid, methyl red can be used as an indicator.
- STATEMENT-2 :** Methyl red changes its colour in the pH range 4.2 to 6.3.
- 11. STATEMENT-1 :** Sparingly soluble salts AB and XY₂ with the same solubility product, will have different solubility.
- STATEMENT-2 :** Solubility of sparingly soluble salt depend upon solubility product.
- 12. STATEMENT-1 :** Solubility product of BaF₂ will increase on dilution.
- STATEMENT-2 :** Solubility of BaF₂ will change on changing temperature.
- 13. STATEMENT-1 :** Solubility of sparingly soluble salt decreases due to common ion effect.
- STATEMENT-2 :** Solubility product constant does not depend on common ion effect.
- 14. STATEMENT-1 :** Solubility of AgCl in NH₃(aq.) is greater than in pure water.
- STATEMENT-2 :** When AgCl dissolve in NH₃(aq.), complex ion formation $[Ag(NH_3)_2]^+$ takes place and solubility equilibria of AgCl shifted in forward direction.
- 15. STATEMENT-1 :** Solubility of AgCN in acidic solution is greater than that in pure water.
- STATEMENT-2 :** Solubility equilibria of AgCN in acidic solution is shifted in forward direction due to formation of HCN.

Subjective Problems




- Calculate pOH of 0.1M aq. solution of weak base BOH ($K_b = 10^{-7}$) at 25 °C.
- pH of 0.01M aq. solution of HA is 4. Find the value of pK_a of HA at 25 °C.
- Calculate approximate pH of 10^{-10} M NaOH at 25 °C.
- Calculate pH of a resultant solution of 25 mL of 0.1M HCl, 50 mL of 0.02 M HNO_3 and 25 mL of 0.1M NaOH.
- Calculate pH of a resultant solution of 0.1M HA ($K_a = 10^{-6}$) and 0.45 M HB ($K_a = 2 \times 10^{-6}$) at 25 °C.
- 0.16 g N_2H_4 ($K_b = 4 \times 10^{-6}$) are dissolved in water and the total volume made up to 500 mL. Calculate the percentage of N_2H_4 that has reacted with water in this solution
- Calculate pH of a buffer solution that contains 0.1M NH_4OH ($K_b = 10^{-5}$) and 0.1M NH_4Cl .
- Calculate the ratio of sodium formate and formic acid ($K_a = 2 \times 10^{-4}$) in a buffer solution of pH 4.3.
- What is the pOH of 0.1M KB (salt of weak acid and strong base) at 25 °C ? (Given : pK_b of B = 7)
- A certain weak acid has $K_a = 10^{-5}$. If the equilibrium constant for its reaction with a strong base is represented as 1×10^y then find the value of y.
- If solubility of AgCl in 0.2 M solution of AgNO_3 is represented as $y \times 10^{-10}$ then find the value of y.
(Given : $K_{sp}(\text{AgCl}) = 10^{-10}$)
- When one litre of a saturated solution of PbCl_2 (mol. mass = 278) is evaporated, the residue is found to weight 2.78 g. If K_{sp} of PbCl_2 is represented as $y \times 10^{-6}$ then find the value of y.
- A solution is saturated in SrCO_3 and SrF_2 . The CO_3^{2-} was found to be 10^{-3} mol/L. If the concentration of F^- in solution is represented as $y \times 10^{-2}$ M then what is the value of y ?
[Given : $K_{sp}(\text{SrCO}_3) = 2.5 \times 10^{-10}$; $K_{sp}(\text{SrF}_2) = 10^{-10}$]
- 10 mL of H_2A (weak diprotic acid) solution is titrated against 0.1M NaOH. pH of the solution is plotted against volume of strong base added and following observation is made.



If pH of the solution at first equivalence point is pH_1 and at second equivalence point is pH_2 . Calculate the value of $(\text{pH}_2 - \text{pH}_1)$ at 25°C
Given: For H_2A , $pK_{a1} = 4.6$ and $pK_{a2} = 8$, $\log 25 = 1.4$

15. Amongst the following, the total number of compounds whose aqueous solution turns red litmus paper blue is:



Given:

Acid	Ka_1	Ka_2	Ka_3
H_3PO_4	10^{-3}	10^{-8}	10^{-12}
H_2CO_3	10^{-6}	10^{-11}	—
$\text{H}_2\text{C}_2\text{O}_4$	10^{-2}	10^{-5}	—

ANSWERS

Level 1

1. (d)	2. (c)	3. (a)	4. (c)	5. (c)	6. (a)	7. (a)	8. (d)	9. (d)	10. (c)
11. (d)	12. (d)	13. (c)	14. (d)	15. (b)	16. (d)	17. (a)	18. (d)	19. (c)	20. (c)
21. (a)	22. (b)	23. (a)	24. (b)	25. (a)	26. (c)	27. (b)	28. (c)	29. (b)	30. (b)
31. (b)	32. (a)	33. (b)	34. (a)	35. (a)	36. (b)	37. (c)	38. (a)	39. (c)	40. (c)
41. (c)	42. (c)	43. (b)	44. (c)	45. (b)	46. (a)	47. (c)	48. (d)	49. (d)	50. (d)
51. (d)	52. (b)	53. (b)	54. (c)	55. (d)	56. (d)	57. (b)	58. (c)	59. (c)	60. (b)
61. (b)	62. (d)	63. (b)	64. (b)	65. (b)	66. (c)	67. (a)	68. (c)	69. (d)	70. (a)
71. (d)	72. (b)	73. (d)	74. (c)	75. (b)	76. (b)	77. (a)	78. (d)	79. (b)	80. (c)
81. (b)	82. (d)	83. (a)	84. (a)	85. (c)	86. (a)	87. (c)	88. (b)	89. (d)	90. (b)
91. (c)	92. (d)	93. (b)	94. (a)	95. (d)	96. (a)	97. (a)	98. (d)	99. (b)	100. (a)
101. (a)	102. (b)	103. (c)	104. (b)	105. (c)	106. (d)	107. (d)	108. (b)	109. (a)	110. (c)
111. (d)	112. (c)	113. (a)	114. (a)	115. (a)	116. (c)	117. (a)	118. (c)	119. (a)	120. (a)
121. (c)	122. (c)	123. (c)	124. (a)	125. (c)	126. (d)	127. (a)	128. (a)	129. (d)	130. (b)
131. (d)	132. (b)	133. (c)	134. (c)	135. (d)	136. (d)	137. (a)	138. (d)	139. (c)	140. (a)
141. (a)	142. (d)	143. (c)	144. (c)	145. (d)	146. (a)	147. (b)	148. (b)	149. (d)	150. (b)

Level 2

1. (d)	2. (c)	3. (c)	4. (a)	5. (d)	6. (c)	7. (a)	8. (d)	9. (a)	10. (a)
11. (d)	12. (c)	13. (a)	14. (b)	15. (c)	16. (c)	17. (b)	18. (c)	19. (b)	20. (b)
21. (b)	22. (d)	23. (c)	24. (c)	25. (d)	26. (d)	27. (a)	28. (b)	29. (a)	30. (b)
31. (a)	32. (c)	33. (d)	34. (b)	35. (d)					



Level 3

Passage-1:	1. (c)	2. (b)	3. (b)	4. (b)				
Passage-2:	1. (c)	2. (b)	3. (b)	4. (b)				
Passage-3:	1. (b)	2. (a,c)	3. (c)	4. (a)	5. (b)	6. (a)		
Passage-4:	1. (a)	2. (a)	3. (b)	4. (a)	5. (c)			

One or More Answer is/are Correct

1. (b,c) 2. (a,b,c) 3. (a,b) 4. (b) 5. (a,c) 6. (a,b,c,d) 7. (a,b) 8. (a,b)
 9. (a,c) 10. (a,c,d) 11. (a,b,d) 12. (a,c) 13. (a,b) 14. (b, c) 15. (b, c, d) 16. (a, c)
 17. (b, c, d) 18. (a, b, c) 19. (a, c, d) 20. (a, b) 21. (b,c,d) 22. (a,c) 23. (c,d) 24. (b,c,d)
 25. (a,b,c)

Match the Column

1. A Q, R; B P, S; C P Q, S; D P, S
 2. A P, S; B Q; C P, Q, R; D P, R
 3. A P; B S; C Q; D R
 4. A Q; B P; C S; D R
 5. A R; B Q, S; C R; D P, S
 6. A P; B S; C R; D Q
 7. A P, Q, R; B R; C Q, S; D Q, R
 8. A P, Q, R; B R; C P, Q; D S
 9. A S; B P; C R; D Q
 10. A Q; B R; C P; D S

Assertion-Reason Type Questions

1. (A) 2. (D) 3. (A) 4. (A) 5. (A) 6. (D) 7. (A) 8. (A) 9. (B) 10. (D)
 11. (B) 12. (D) 13. (B) 14. (A) 15. (A)


Subjective Problems

1. 4	2. 6	3. 7	4. 2	5. 3	6. 2	7. 9	8. 4	9. 4	10. 9
11. 5	12. 4	13. 2	14. 4	15. 6					

Hints and Solutions

Level 1

1,2 (d, c)



If, HX acid

then, X^- conjugate base

3. (a) Among HF, HCl, HBr and HI,

HF is weakest acid

So, F⁻ is strongest base.

4. (c) H_3PO_2 and H_3PO_3 are mono and dibasic acids. Hence, NaH_2PO_2 and Na_2HPO_3 are normal salts, while Na_2HPO_4 is acid-salt as one proton is there in the compound.

7. (a) For a conjugate acid-base pair,

$$K_a \cdot K_b = K_w$$

8. (d) pH pOH pK_w

9. (d) In neutral solution

$$[H^+] = \sqrt{K_w}$$

10. (c) As temperature increases, $[H^+]$ and $[OH^-]$ both increase.

11-12-14. (d, d, d)

$$pH = -\log_{10}[H^+]$$

13. (c) $[H^+] = \frac{K_w}{[OH^-]}$ and $pH = -\log_{10}[H^+]$

15. (b) $[H^+] = 10^{-9}$

If it is 1000 times more basic then,

$$[H^+]_{\text{original}} = 10^{-6} M$$

16. (d) When equal volumes are taken, the concentration becomes half.

$$[H^+] = \frac{1}{2} (10^{-3} + 10^{-5})$$

$$= 505 \times 10^{-6}$$

$$pH = 3.3$$

17. (a) $[H^+] = 0.002 M$

$$pH = \log 2 + 3$$

$$= 3 + \log 2$$

$$pOH = 11 - \log 2$$

18. (d) pH 4, $[H^+] = 10^{-4}$ mol/litre

Equivalent of HCl in 100 mL solution

$$\frac{10^{-4} \times 100}{1000} = 10^{-5}$$

$$\begin{aligned} 19. (c) \quad & \frac{N_1 V_1}{10^{-3}} = \frac{N_2 V_2}{10^{-5}} \\ & 10^{-3} \times 10 = N_2 \times 1000 \\ & N_2 = 10^{-5} \\ & pH = 5 \end{aligned}$$

$$\text{and } pOH = 14 - 5 = 9$$

20. (c) pH 1 $[H^+] = 0.1 M$

For $H_2SO_4(aq.)$

$$[H^+] = 2 [Acid]$$

$$\begin{aligned} 21. (a) \quad & H_2A(aq.) \xrightarrow{100\%} 2H^+(aq.) + A^{2-}(aq.) \\ & pH = -\log [H^+] \end{aligned}$$

$$22. (b) [Ca(OH)_2(aq.)] = \frac{0.60}{74} \times \frac{1000}{1500}$$

$$= 5.40 \times 10^{-3} \quad Ca(OH)_2(aq.) \rightleftharpoons Ca^{2+}(aq.) + 2OH^-(aq.)$$

$$[Ca^{2+}] = 5.40 \times 10^{-3}$$

$$[OH^-] = 2 \times 5.40 \times 10^{-3} = 1.08 \times 10^{-2}$$

23-24. (a, b)

At these concentration consider ionization of water also.

$$\begin{aligned} 25. (a) \quad & H_2SO_4 + 2NaOH \rightleftharpoons Na_2SO_4 + 2H_2O \\ & 0.05 \text{ mole} \quad 0.1 \text{ mole} \quad \quad \quad 0.05 \quad 0.1 \end{aligned}$$

Aqueous solution of SASB type salt

i.e., pH 7

26. (c) HCl NaOH NaCl H_2O

$$(i) 2.5 \text{ m mole} \quad 2.49 \text{ mole}$$

$$[H^+] = \frac{0.01}{50} = 2 \times 10^{-4} M$$

$$pH = 4 - \log 2 = 3.7$$

$$(ii) 2.5 \text{ m mole} \quad 2.51 \text{ M mole}$$

$$[OH^-] = \frac{0.01}{50} = 2 \times 10^{-4} M$$


$$pOH = 3.7$$

27. (b) Milli-equivalents of HCl 25 0.08 2

Milli-equivalents of NaOH 2.5


$$\text{remaining } [OH^-] = \frac{2.5 - 2}{500} = 10^{-3}$$

$$[H^+] = 10^{-11} \text{ or } pH = 11$$



29. (b) 20 meq. of HCl reacts completely with 20 meq. of KOH. Hence, the final solution is neutral. Thus, at 90°C , $\text{pH} = 6$.

30. (b) In aqueous reaction, upon dilution, the reaction moves towards more equation moles.



$$K_{eq} = \frac{K_1}{K_2}$$


32. (a) $\therefore$ Strength of acid $\frac{1}{\text{p}K_a}$

Formic acid will be the strongest acid.

33. (b) More H_{ion} weaker is the acid.

34. (a) $\sqrt{\frac{K_a}{C}} = 0.05$

i.e., $[\text{H}^+] = \sqrt{K_a C}$



$$K_a = \frac{Cx^2}{C(1-x)}$$

$$10^{-4} = \frac{(0.01)^2}{1-x}$$

$$\frac{0.095}{9.5} = x$$

36. (b) $\sqrt{\frac{K_a}{C}}$

37. (c) $\therefore 0.05$


$$K_a = \frac{C^2}{1}$$

38. (a) $\text{pH} = 9$; $[\text{OH}^-] = 10^{-5}$; $C = 10^{-5}$
 10^{-4}

$$\% \text{ dissociation} = \frac{10^{-4}}{10^{-5}} \times 100 = 0.01\%$$

39. (c) $[\text{H}^+] = C$ and $[\text{OH}^-] = \frac{K_w}{[\text{H}^+]}$

40. (c) is negligible w.r.t. 1 so $K_a = C^2$
 $0.1 = (0.01)^2 = 10^{-5}$
 $\text{p}K_a = 5$



neglect x w.r.t. 0.01 due to common ion effect

$$K_a = x$$

$$x \sim [\text{HCOO}^-] = 1.8 \times 10^{-4}$$

42. (c) (P) $(\text{p}K_a)_{\text{HX}} = (\text{p}K_b)_X = \text{p}K_w$
 (Q) Very dilute solution

(R) $\text{pH} = \frac{1}{2} \text{p}K_a + \frac{1}{2} \log_{10} C$

(S) $[\text{H}^+]_1 = 10^{-3} M$, $[\text{H}^+]_2 = 10^{-5} M$
 $[\text{H}^+]_{\text{Final}} = \frac{M_1 V_1}{V_1 + V_2} + \frac{M_2 V_2}{V_1 + V_2}$

43. (b) $\therefore$ is negligible w.r.t. 1

$$K_a = \frac{C_1^2}{C_2} = \frac{C_2^2}{2}$$

$$C_2 = C_1 \cdot \frac{1}{2}$$

$$0.2 = \frac{1}{4} \times 0.05$$

$$\frac{C_1 V_1}{300} = \frac{C_2 V_2}{0.2 \times 0.05 + V_2}$$

$$\frac{300}{V_2} = \frac{0.2 \times 0.05}{1200 \text{ mL}}$$

$$\text{Volume of H}_2\text{O added} = \frac{1200 \times 300}{900} = 400 \text{ mL}$$

44. (c) Due to low value of K_b and common ion effect we can neglect x w.r.t. 0.01 when $x = C$

$$K_b = \frac{x \cdot 0.01}{0.02}$$

$$x = \frac{[\text{NH}_4^+]}{1} = \frac{3.6 \times 10^{-5}}{22.41}$$

45. (b) $n_{\text{RNH}_2(\text{g})} = \frac{1}{0.0821} \times \frac{22.41}{273} = 1$

$$[\text{RNH}_2(\text{aq.})] = \frac{n_{\text{RNH}_2}}{\text{Volume of solution}} = 1$$

is negligible w.r.t. 1 so

$$\text{pOH} = \frac{1}{2} [\text{p}K_b - \log C] = 2$$

46. (a) is negligible w.r.t. 1 for both acid so

$$[\text{H}^+] = \sqrt{\frac{K_{a1} C_1}{1} + \frac{K_{a2} C_2}{1}}$$

$$[\text{H}^+] = \sqrt{\frac{1.8 \times 10^{-5} \times 0.01}{1} + \frac{6.3 \times 10^{-5} \times 0.01}{1}}$$

$$[\text{H}^+] = \sqrt{81 \times 10^{-8} + 9 \times 10^{-4}}$$

$$47. (c) [HA] = \frac{60}{10000} \times 10^{-5} M$$

$$= \frac{\sqrt{10^{-9}}}{\sqrt{10^{-5}}} = 0.01 \text{ (negligible w.r.t. 1)}$$

$$[H^+] = \sqrt{10^{-9} \times 10^{-5}} = \sqrt{10^{-14}}$$

$$= 10^{-7} M, \quad \therefore [H^+] = 10^{-6} M$$

So we should consider $[H^+]$ from H_2O
 $[H^+]_{\text{total}} = 2 \times 10^{-7}$ without considering
 common ion effect $pH = 6.7$, so $pOH = 7.3$
 but due to C.I. effect H^+ from H_2O will be
 less

so $pH > 6.7$

pOH of solution is exist between 7.0 and
 7.3.

48. (d) and 49. (d)

$$K_{a1} = \frac{[H^+]_{\text{total}} [HCO_3^-]}{[H_2CO_3]}$$

$\therefore K_{a1}, K_{a2}$ and concentration is
 appreciable so we can neglect w.r.t. 1

$$\frac{[H^+]_{\text{total}}}{[HCO_3^-]} \sim \frac{[HCO_3^-]}{\sqrt{4 \times 10^{-7} \times 0.025}} = 10^{-4}$$

and $[CO_3^{2-}] \sim K_{a2}$

50. (d) $[H^+]_{\text{total}} \sim [H^+]$ from first step

$$K_{a1} = \frac{C}{(1-x)}; \text{ is not negligible w.r.t. 1 so}$$

after solving quadratic equation 0.095

$$[OH^-] = \frac{10^{-14}}{0.095 \times 0.3}$$

$$= 3.5 \times 10^{-13}$$

52. (b) Salt of SAWB type.

53. (b) Salt of WASB type.

$$54. (c) \quad pH = \frac{1}{2} pK_a - \frac{1}{2} \log C$$

$$= 5 - \frac{1}{2} pK_a - \frac{1}{2} \log 2$$

$$pK_a = 8 \quad pK_b = 8;$$

$$pOH = \frac{1}{2} pK_b - \frac{1}{2} \log C = 4 - \frac{1}{2} \log 2 = 6$$

56. (d) All are WASB salts,

$$\text{So, } [OH^-] = \sqrt{\frac{K_w C}{K_a}}$$

For weakest acid, K_a should be minimum.

$[OH^-]$ minimum

$[H^+]$ maximum

pH minimum

57. (b) I WASB salt

II WA

III WAWB salt

IV SB

V SA

58. (c) As temperature pK_w

pH also

$$61. (b) [H^+]_{\text{final}} = \frac{1 \times 0.1}{(1 - 999)} = 10^{-4} M$$

pH = 4

65. (b) Similar to Q. No.7

$$66. (c) (K_b)_{C_6H_5O} = \frac{K_w}{10^{-10}}$$

$$[OH^-] = \sqrt{K_b \cdot c}$$

$$67. (a) K_{a(HOCN)} = \frac{10^{-14}}{10^{-10}} = 10^{-4}$$

$$K_h = K_{b(OCN)} = \frac{10^{-10}}{\sqrt{\frac{K_h}{C}} \sqrt{\frac{10^{-10}}{0.01}}} = 10^{-4}$$

$$\therefore 0.1; K_h \cdot c^2$$

$$[OH^-] = c; [OH^-] = 0.01 \times 10^{-4} = 10^{-6} M$$

68. (c) $[H^+] = C$

$$0.05$$

$$K_a = \frac{C}{1-x}$$

$$69. (d) \frac{1}{2} \sqrt{\frac{C_2}{C_1}} = \frac{0.01}{2} \sqrt{\frac{0.4}{0.1}}$$

$$= 0.005 \text{ or } \% = 0.5$$

$$70. (a) \text{ For WAWB salt, } \sqrt{\frac{K_w}{K_a K_b}} = \frac{1}{1 - C} \sqrt{K_h}$$

$$71. (d) \sqrt{\frac{K_w}{K_a C}}$$

|Enthalpy of neutralisation| $K_a \cdot h$

$$72. (b) [NaA] = \frac{25 \times 0.1}{25 \times 50} = \frac{2.5}{75}$$

$$[H^+] = \sqrt{\frac{K_w K_a}{C}} = 1.732 \times 10^{-9}$$

$$74. (c) pH = \frac{1}{2} (pK_w - pK_a - pK_b) = 6.5$$

$$75. (b) \therefore pH = \frac{1}{2} pK_w - \frac{1}{2} pK_a - \frac{1}{2} pK_b = 7$$

$$\text{and } \sqrt{\frac{K_w}{K_a K_b}} \sqrt{\frac{10^{-14}}{10^{-12}}} = 0.1$$

or % 10

76. (b) does not depend on concentration in this case.

$$77. (a) \sqrt{\frac{K_h}{C}}; [\text{H}_3\text{O}^+] = C$$

$$\sqrt{K_h C} \sqrt{0.02 \times 5 \times 10^{-9}} = 10^{-5}$$

78. (d) $\text{Ni}^{2+}(\text{aq.}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{Ni}(\text{OH})^+(\text{aq.}) + \text{H}^+(\text{aq.})$

$$t = 0 \quad 0.04 \text{ M}$$

$$t_{\text{eq.}} \quad (0.04 - x) \quad \quad \quad (x) \quad \quad \quad (x)$$

$$x = 4.5 \times 10^{-6}$$

$$K_a = \frac{x^2}{(0.04 - x)}$$

79. (b) is not negligible w.r.t. 1 so after solving quadratic equation 0.095

$$[\text{H}^+] = C = 0.0095; \text{pH} = 2.02$$

80. (c) For amphoteric salt,

$$81. (b) \text{pH} = \frac{pK_{a_2} + pK_{a_3}}{2}$$

82. (d) Similar to Q. No. 83

83. (a)

	$\text{CH}_3\text{COOH} + \text{NaOH}$	$\text{CH}_3\text{COONa} + \text{H}_2\text{O}$
initial	6	2
milli-		
moles		
after	4	—
reaction		2

$$\text{pH} = \text{p}K_a + \log \frac{[\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]} = 4.44$$

85. (c) $\text{HX} + \text{NaOH} \rightleftharpoons \text{NaX} + \text{H}_2\text{O}$

(I) 2m mole 1m mole — —

(2 1) — 1

$$\text{pH} = \text{p}K_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$$

	HX	NaOH	NaX	H_2O
(II)	2m mole	2m mole	—	—
	2	2	2	

$$[\text{NaX}] = 0.01 \text{ M}$$

$$\text{pH} = 7 + \frac{1}{2} [pK_a + \log C]$$

86, 87, 90 (a, c, b)

For ABS

$$\text{pH} = \text{p}K_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$$

88. (b) For BBS

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{Salt}]}{[\text{Base}]}$$

91. (c) $\text{CH}_3\text{NH}_2 + \text{HCl} \rightleftharpoons \text{CH}_3\text{NH}_3\text{Cl}$

initial moles 0.12 0.08

after reaction 0.04 — 0.08

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{CH}_3\text{NH}_3^+]}{[\text{CH}_3\text{NH}_2]}$$


$$3.3 = \log \frac{0.08}{0.04} = 3.6$$

$$\text{pH} = 10.4$$

92. (d) $\text{pOH} = \text{p}K_b + \log \frac{[\text{NH}_4^+]}{[\text{NH}_4\text{OH}]}$ 6

$$\text{pH} = 14 - 6 = 8.0$$

94. (a) BBS



Addition of HCl takes equilibrium to right

NH_4OH	H^+	$\text{NH}_4^+ + \text{H}_2\text{O}$
0.2 mole	10^{-6}	0.2
$(0.2 - 10^{-6})$	—	$(0.2 + 10^{-6})$

$$[\text{OH}^-] = K_b \frac{[\text{Base}]}{[\text{Salt}]}$$

95. (d) $\text{HCOOH} + \text{NaOH} \rightleftharpoons \text{HCOONa} + \text{H}_2\text{O}$

(I) a a/5

$$(a - a/5) \quad \quad \quad (a/5)$$

$$\text{pH}_1 = \text{p}K_a + \log \frac{[\text{HCOOH}]}{[\text{HCOONa}]}$$

(II) a 4a/5

(a - 4a/5) — 4a/5

$$\text{pH}_2 = \text{p}K_a + \log \frac{[\text{HCOOH}]}{[\text{HCOONa}]}$$

$$\text{pH} = \text{pH}_2 - \text{pH}_1$$

97. (a) Henderson equation is

$$\text{pH} = \text{p}K_a + \log \frac{[\text{salt}]}{[\text{acid}]} = 4$$

$$\text{pOH} = 10, [\text{OH}^-] = 10^{-10}$$

Number of OH^- ions $10^{-10} N_A$;

where N_A Avogadro's number.

98. (d) pH upon addition of base.

99. (b) SASB salt.

100. (a) $pK_b(\text{NH}_4\text{OH}) = 14 - pK_a(\text{NH}_4^+) = 4.74$

So, buffer range,

$$pK_b \pm 1$$

101 (a) Mol of $\text{C}_6\text{H}_5\text{COOH} = \frac{61}{122} = 0.5$;

Mol of $\text{C}_6\text{H}_5\text{COONa} = \frac{72}{144} = 0.5$

$$\text{pH} = \text{p}K_a + \log \frac{(\text{C}_6\text{H}_5\text{COONa})}{(\text{C}_6\text{H}_5\text{COOH})};$$

$$\text{pH} = 4.2 + \log \frac{0.2}{0.8} = 3.6$$

$$102. (b) \text{pH} = \text{p}K_{a_2} + \log \frac{[\text{CO}_3^{2-}]}{[\text{HCO}_3^-]}$$

$$(11 + \log 4) + \log \frac{0.2}{0.4} = 10.1$$

$$103. (c) \text{H}_3\text{PO}_4 \quad \text{PO}_4^{3-} \quad \text{H}_2\text{PO}_4^- \quad \text{HPO}_4^{2-}$$

$$\text{initial} \quad 20 \quad 0.1 \quad 20 \quad 0.1$$

milli
moles

$$\text{after reaction} \quad - \quad - \quad 2 \quad 2$$

Buffer solution of H_2PO_4^- (acid) and HPO_4^{2-} (conjugate base) is formed;

$$\text{pH} = \text{p}K_{a_2} + \log \frac{[\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]} = \text{p}K_{a_2}$$

106. (d) At eq. point, WASB salt forms means base solution

So, $\text{CTR} = 7$

107. (d) At equivalent point


$$\text{pH} = \frac{1}{2} [\text{p}K_w + \text{p}K_a + \log C]$$

$$\frac{1}{2} [14 + 4.74 + 2] = 8.37$$

$$\text{Where } [\text{Salt}] = \frac{\text{milli-moles of acid}}{\text{Total volume}}$$


$$\frac{20 \times 0.02}{20 + 20} = 0.01$$

For best indicator pH at equivalent point should lie between colour transition range of indicator.



$$K_a = \frac{[\text{H}^+][\text{In}^-]}{[\text{HIn}]}$$

$$6 \times 10^{-5} = \frac{10^{-5}(x)}{(100 - x)}$$



$$\text{Case I : } 3 \times 10^{-5} = \frac{[\text{H}^+]_1 \times 0.25}{0.75}$$

$$\text{Case II : } 3 \times 10^{-5} = \frac{[\text{H}^+]_2 \times 0.75}{0.25}$$

$$[\text{H}^+]_1 = 9 \times 10^{-5} \quad [\text{H}^+]_2 = 10^{-5}$$

Change in $[\text{H}^+] = 8 \times 10^{-5}$

$$110. (c) \text{pH} = \text{p}K_{\text{In}} + \log \frac{[\text{In}^-]}{[\text{HIn}]} = \text{p}K_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$$



For second eq. pt

$$(n)_{\text{NaOH}} = 2(n)_{\text{H}_2\text{A}} \quad [\text{Na}_2\text{A}] = \frac{0.4}{(32 + 250)}$$

$$0.25 \times V = 2(0.2 \times 20)$$

$$V = 32 \text{ mL}$$



$$(I) \quad a \quad a/2$$

$$(a \quad a/2) \quad a/2$$

WA its conjugate ABS

$$\text{pH} = \text{p}K_{a_1} + \log \frac{[\text{NaHA}]}{[\text{H}_2\text{A}]} = \text{p}K_{a_1}$$

$$(II) \quad a \quad a$$

$$- \quad - \quad (a)$$

aq. solution of amphiprotic salt

$$\text{pH} = \frac{\text{p}K_{a_1} + \text{p}K_{a_2}}{2}$$

114. (a) When equal volume mixed concentration become half also, for ppt,

$$[\text{Ag}^+][\text{Cl}^-] = K_{sp}$$

116. (c) When 1 litre each solution are mixed


$$[\text{OH}^-] = 10^{-6} \text{ M (Buffer solution)}$$

$$M^n = 0.05 \text{ M}$$

$$\text{for } Q = [0.5][10^{-6}]^2 = 5 \times 10^{-14}$$

$$Q < K_{sp} \text{ for } \text{Fe}^{2+} \text{ and } \text{Ni}^{2+}$$

So $\text{Fe}(\text{OH})_2$ and $\text{Ni}(\text{OH})_2$ are ppt.



$$0.04 \times 50 = M_2 \times 200 \quad M_2 = 0.01$$

Similarly, for $[\text{SO}_4^{2-}]$

$$0.0008 \times 150 = M_2 \times 200$$

$$M_2 = 0.0006$$

$$\text{ionic product (Q)} = [\text{Ca}^{2+}][\text{SO}_4^{2-}]$$

$$6 \times 10^{-8} < \text{So } Q < K_{sp}$$

118. (c) For AgCl

$$K_{sp} = [\text{Ag}^+][\text{Cl}^-]$$

$$y \quad x$$

xy constant

i.e., Rectangular hyperbolic


119. (a) KNO_3 does not effect the equilibrium.

$$120. (a) [\text{Ag}^+] = \frac{10^{-3}}{50 + 50} = 1000 \times 0.01$$

$$[\text{Cl}^-] = \frac{0.1}{50} - \frac{50}{50} = 0.05$$

$$Q = [\text{Ag}^+][\text{Cl}^-] = 5 \times 10^{-4}$$


$$\therefore Q < K_{sp}$$



$$\begin{array}{l} \text{at given condition} \quad 0.01 \quad 0.05 \\ \text{after precipitation} \quad x \quad \sim 0.04 \end{array}$$

$$K_{sp} = [\text{Ag}^+][\text{Cl}^-] = 10^{-10} \quad x \cdot 0.04$$

$$x = \frac{10^{-10}}{0.04} = 2.5 \times 10^{-9}$$



$$t = 0 \quad a$$


$$t_{eq.} \quad (a-x) \quad (x) \quad (x-y)$$

$$K_{sp} = [A^y]^x [B^x]^y$$

$$(x-y)^x (y-x)^y$$

122. (c) Molarity of saturated solution

Solubility



$$t = 0 \quad a$$

$$t_{eq.} \quad (a-3s) \quad 3s \quad 4s$$

$$K_{sp} = [\text{Zr}^{4+}]^3 [\text{PO}_4^{3-}]^4$$

124. (a) The corresponding values of solubility products of the given electrolytes are :

$$\begin{array}{ccc} \text{MX}_1 & \text{MX}_2 & \text{MX}_3 \\ S^2 \cdot 10^{-6}, & 4S^3 \cdot 4 \cdot 10^{-9}, & 27S^4 \cdot 27 \cdot 10^{-12} \end{array}$$



126. (d) For most soluble salt, solubility should be maximum.

$$\text{127. (a)} \quad (A) [\text{Ag}^+] = \frac{K_{sp}}{[\text{Br}^-]} = 5 \times 10^{-12} M$$

$$(B) [\text{Ag}^+] = \frac{K_{sp}}{[\text{Cl}^-]} = 1.8 \times 10^{-7} M$$


$$(C) [\text{Ag}^+] = \sqrt{\frac{K_{sp}}{[\text{CO}_3^{2-}]}} = 9 \times 10^{-6} M$$

$$(D) [\text{Ag}^+] = \frac{K_{sp}}{[\text{ASO}_4^{3-}]} = 10^{-7} M$$

$$\text{128. (a)} \quad \text{Solubility of } AB = \sqrt{K_{sp}} = 2 \times 10^{-10}$$

$$\text{Solubility of } A_2B = \sqrt[3]{\frac{K_{sp}}{4}} = 2 \times 10^{-4}$$

$$\text{Solubility of } AB_3 = \sqrt[4]{\frac{K_{sp}}{27}} = 10^{-8}$$



$$S = \sqrt{6.4 \times 10^{-5}} = 8 \times 10^{-3} \text{ mol/L}$$


$$8 \text{ mol/m}^3$$

$$\text{130. (b)} \quad S = \frac{6.9 \times 10^{-2} \text{ mol/100 mL}}{690} = 10^{-3} \text{ mol/L}$$

$$\text{For } \text{Ba}_3(\text{AsO}_4)_2, K_{sp} = 108 S^5$$

131-132-133. (d, b, c)


Similar to Q.No. 130



$$S \quad 2S$$


$$K_{sp} = 4S^3; \quad S = \frac{10^{-6} \cdot 1/3}{4} = 0.63 \times 10^{-2}$$

$$\text{Molarity of F} = 2S = 2 \cdot 0.63 \times 10^{-2} = 1.26 \times 10^{-2}$$



$$t_{eq} \quad S \quad 3S$$

$$K_{sp} = (S)^1 (3S)^3$$



$$S \quad 2S$$

$$K_{sp} = 4S^3$$

$$S = 4 \times 10^{-7}$$

$$[\text{OH}^-] = 2S = 8 \times 10^{-7}$$


$$\text{pOH} = 6.1 \quad \text{pH} = 7.90$$



$$10^{-5} = \frac{14.35 \times 10^{-3}}{143.5}$$

$$V (\text{in litre})$$

$$V = 10 \text{ litre or } 0.01 \text{ m}^3$$



$$t_{eq.} \quad S \quad 10^{-1} M$$

$$K_{sp} = [\text{Fe}^{2+}][\text{OH}^-]^2$$


$$8 \times 10^{-16} = (S)^1 (10^{-1})^2$$

139. (c) To precipitate

$$[\text{Pb}^{2+}][\text{OH}^-]^2 = K_{sp}$$

$$[\text{OH}^-]_{\text{Min}} = \sqrt{\frac{K_{sp}}{[\text{Pb}^{2+}]}}$$

141. (a) Let solubility of Ag_2CO_3 in presence of Na_2CO_3 is x ,



$$2x \quad (x - 0.1) \sim 0.1$$

$$K_{sp} = [\text{Ag}^+]^2 [\text{CO}_3^{2-}]$$

$$4 \times 10^{-13} = (2x)^2 \cdot 0.1$$


$$x = 10^{-6}$$

142-143. (d, c)

Similar to Q.No. 139

144. (c) More common ion more in solubility.

$$145. (d) K_{sp} = 256 \times 10^{-5}$$



$$K_{sp} = \frac{[\text{Pb}^{2+}][\text{Br}^-]^2}{(0.8 \times 10^{-5})(1.6 \times 10^{-5})^2}$$

$$S = \frac{10^{-4}}{1.6 \times 1.6}$$

147. (b) For basic buffer solution

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{NH}_4^+]}{[\text{NH}_3]} = \text{p}K_b$$

$$\frac{[\text{OH}^-]}{K_{sp}} = \frac{[\text{Mn}^{2+}][\text{OH}^-]^2}{[\text{Mn}^{2+}]}$$

$$1.38 \times 10^{-4} M$$

$$148. (b) [\text{OH}^-] = \sqrt{\frac{K_{sp}}{[\text{Mg}^{2+}]}} = \sqrt{\frac{10^{-11}}{10^{-3}}} = 10^{-4}$$



$$\text{pOH} = \text{p}K_b + \log \frac{[\text{NH}_4\text{Cl}]}{[\text{NH}_3]}$$

$$4.5 = \log \frac{[\text{NH}_4\text{Cl}]}{[\text{NH}_3]}$$


$$0.10 = \frac{[\text{NH}_4\text{Cl}]}{[\text{NH}_3]}$$

$$\text{moles of NH}_4\text{Cl required} = 0.1 \times 0.02 = 2 \times 10^{-3}$$

149. (d) Due to higher value of K_f , mostly Ag converted into complex.



overall reaction is



$$K_{eq} = \frac{K_{sp}}{K_f} = \frac{x^2}{1^2}$$

$$x = 4.9 \times 10^{-5} \text{ mol/litre}$$

So mass of AgI required is

$$4.9 \times 10^{-5} \times 235 = 0.011 \text{ g}$$

Level 2



$$\text{Initial} \quad 2 \quad 3.75$$

milli-moles

milli-moles of remaining NaOT = 1.75


$$[\text{OT}^-] = \frac{1.75}{25} = 7 \times 10^{-2} \text{ moles}$$

$$\text{pOT} = 2 - \log 7$$

$$\text{pT} + \text{pOT} = 7.62 - 2$$

$$\text{pT} = 15.24 - 2 - \log 7$$

$$13.24 - \log 7$$



$$K_{SIC} = \frac{[\text{NH}_4^+][\text{NH}_2^-]}{[\text{NH}_3]^2} = 10^{-15} M$$

$$\therefore \text{Number of NH}_4^+ \text{ ions in } 1000 \text{ cm}^3 = 10^{-15} \times 6.022 \times 10^{23}$$

$$\text{Number of NH}_4^+ \text{ ions in } 100 \text{ cm}^3 = 10^{-15} \times 6.022 \times 10^{23} \times \frac{100}{1000}$$

$$6.022 \times 10^7$$



$$\text{Now } [\text{H}^+] = \frac{3 \times 10^{-3}}{2} = 1.5 \times 10^{-3}$$

$$(1.5 \times 10^{-3}) = \sqrt{K_a \times C_2}$$


$$C_2 = 0.125$$

milli-moles of acid remain constant


$$C_1 V_1 = C_2 V_2$$

$$10 \times 0.5 = 0.125 \times V_2$$

$$V_2 = 40 \text{ litre}$$



$$[\text{HA}] = \frac{20 \times 0.1}{20 + 20} = 0.05$$



Due to common ion effect neglect x w.r.t. 0.05

$$K_a = \frac{(0.05 - x) \times x}{(0.05 + x)} \sim x$$

$$x = 10^{-5}$$

$$\frac{[\text{A}^-]}{[\text{HA}]} = \frac{x}{0.05} = 2 \times 10^{-4}$$





at eqm. $C(1-x) \quad C \quad C$

$$K_a = \frac{[\text{H}^+][\text{FCH}_2\text{COO}^-]}{[\text{FCH}_2\text{COOH}]}$$

$$2.6 \times 10^{-3} = \frac{(2 \times 10^{-3})^2}{[\text{FCH}_2\text{COOH}]}$$

$$\text{at eqm. } [\text{FCH}_2\text{COOH}] = 1.53 \times 10^{-3}$$

$$\text{Total concentration} = 2 \times 10^{-3} + 1.53 \times 10^{-3} = 3.53 \times 10^{-3}$$



$\therefore$ is very less for both acid so

$$K_a(\text{HCOOH}) = \frac{x(x-y)}{C_1}$$


$$K_a(\text{HF}) = \frac{y(x-y)}{C_2}$$

$$\frac{x}{y} = \frac{K_a(\text{HCOOH}) C_1}{K_a(\text{HF}) C_2}$$


$$\frac{[\text{HCOO}^-]}{[\text{F}^-]} = \frac{x}{y}$$

$$\frac{2 \times 10^{-4}}{6.6 \times 10^{-4}} = \frac{0.2}{0.1} \times \frac{2}{3.3}$$

7. (a) First dissociation



Second dissociation :



Total $[\text{OH}^-] = 4 \times 10^{-3} + 2 \times 10^{-3} = 6 \times 10^{-3}$

$\text{pOH} = 3 - \log 6 \sim 2.22$

$\text{pH} = 11.78$

8. (d) is negligible w.r.t. 1

$[\text{H}^+]$ mainly from first step

$$[\text{H}^+] = \sqrt{K_{a1} C} = \sqrt{10^{-5} \times 0.1} = 10^{-3}$$

$$K_{a3} = \frac{[\text{H}^+][\text{A}^{3-}]}{[\text{HA}^{2-}]} = 10^{-13} = \frac{10^{-3}[\text{A}^{3-}]}{[\text{HA}^{2-}]}$$

$$X = \frac{[\text{A}^{3-}]}{[\text{HA}^{2-}]} = 10^{-10};$$

$$\text{pX} = 10$$




Initial concentration $0.6/2 = 0.3 \text{ M}$

After dissociation $[\text{LaC}^-] = 2 \times 0.3 = 0.6$
 0.54 M


$$\text{pH} = \frac{1}{2} [\text{pK}_w - \text{pK}_a - \log (\text{LaC}^-)]$$

$$14 - 5.6 = \frac{1}{2} [14 - \text{pK}_a - \log (0.54)]$$

$$\text{pK}_a = 2.8 - \log (0.54)$$



at equilibrium $C(1) \rightleftharpoons C(1) + C$



$$K_h = \frac{K_w}{K_a K_b} = \frac{10^{-14}}{(1.8 \times 10^{-5})^2}$$

$$= \frac{10^{-14}}{3.24 \times 10^{-10}} = 3.08 \times 10^{-5}$$

$$= \frac{10^{-14}}{1.8 \times 10^{-5}} = 5.55 \times 10^{-10}$$

$$[\text{CH}_3\text{COOH}] = C$$

$$0.01 \times 0.0055 = 5.55 \times 10^{-5}$$




$$K_a = \frac{[\text{Fe(OH)}^{2+}][\text{H}_3\text{O}^+]}{[\text{Fe}^{3+}]}$$

$$6.5 \times 10^{-3} = \frac{0.20}{0.80} [\text{H}_3\text{O}^+]$$

$$[\text{H}_3\text{O}^+] = 2.6 \times 10^{-2}; \text{pH} = 1.59$$

12. (c) Due to larger difference between K_{h1} and K_{h2} and common ion effect we can assume that y is negligible w.r.t. x



at equilibrium $C \rightleftharpoons x + x + x$



at equilibrium $x \rightleftharpoons y + y + x$

$$K_{h2} = \frac{K_w}{K_{b1}} = \frac{10^{-14}}{10^{-4}} = 10^{-10}$$

$$\therefore \frac{x}{y} = \frac{x}{y} \times \frac{y}{y}$$


$$K_{h2} = \frac{y}{\text{Fe(OH)}_2} = 10^{-10}$$

13. (a) Let x milli-moles of KOH is added

$$6 - 5 = \log \frac{s}{a} \times \frac{x}{x}; \frac{s}{a} \times \frac{x}{x} = 10$$

$$\text{or } \frac{10}{10} \times \frac{x}{x} = 10 \times x = 8.18$$

$$\text{Mass} = 8.18 \times 10^{-3} \times 56 = 0.458 \text{ gm}$$



$$\frac{[\text{C}_6\text{H}_5\text{COO}^-]}{[\text{C}_6\text{H}_5\text{COOH}]} = 2$$

Let volume of acid is $V \text{ mL}$

$$\frac{0.2 (300 - V)}{0.1 V} = 2 \quad V = 150 \text{ mL}$$

15. (c) When half acid is neutralized $\text{pH} = \text{p}K_a$ and at the equivalent point

$$\text{pH} = \frac{1}{2} [\text{p}K_w - \text{p}K_a - \log C]$$

$$9 = \frac{1}{2} [14 - 5 - \log C] \Rightarrow C = 0.1$$

Let V mL of NaOH is used in titration milli eq. of NaOH = 0.25 V milli eq. of salt formed

$$\frac{0.25}{V} \times \frac{V}{60} \times 0.1 = \frac{V}{40} \times 0.1$$

milli equivalent or milli-moles of acid = 10

Mass of acid = $10 \times 10^{-3} \times 82 = 0.82$ gm

Mass % of acid = $\frac{0.82}{1.025} \times 100 = 80\%$

16. (c) $\text{BOH(aq.)} \rightleftharpoons \text{B(aq.)} + \text{OH}^-(\text{aq.})$

At equilibrium : $\frac{[\text{B}][\text{OH}^-]}{[\text{BOH}]}$

$$K_b = \frac{[\text{B}][\text{OH}^-]}{[\text{BOH}]}$$

neglect $\text{w.r.t. } 1$ as concentration is appreciable, $\% = 100 \sqrt{\frac{K_b}{c}}$

Total dissolved base present in solution as BOH and B

$$\text{So } \frac{[\text{B}]}{[\text{B}] + [\text{BOH}]}$$

$$\frac{1}{1 + \frac{[\text{BOH}]}{[\text{B}]}} = \frac{1}{1 + \frac{[\text{OH}^-]}{K_b}}$$

$$\frac{K_b}{K_b + [\text{OH}^-]} = \frac{K_b [\text{H}^+]}{K_b [\text{H}^+] + K_w}$$

also $\text{pOH} = -\log [\text{OH}^-]; [\text{OH}^-] = 10^{-\text{pOH}}$

$$\frac{K_b}{K_b + 10^{-\text{pOH}}} = \frac{1}{1 + 10^{(\text{p}K_b - \text{pOH})}}$$

17. (b) $\text{HA} + \text{NaOH} \rightarrow \text{NaA} + \text{H}_2\text{O}$

milli-moles of salt NaA or A = 40 $\times$ 0.1 = 4

Now

	A	H	HA
Initial milli-moles	4	2	
Final milli-moles	2	—	2

Acidic buffer solution is formed and $[\text{A}^-] = [\text{HA}]$

$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]} = \text{p}K_a = 5$$

Now $\text{HA} + \text{NaOH} \rightarrow \text{NaA} + \text{H}_2\text{O}$


hydrolysis of A will take place

$$[\text{NaA}] = \frac{\text{milli moles of acid}}{\text{Total volume}} = \frac{20}{20} \cdot \frac{0.2}{20} = 0.1$$

$$\text{pH} = \frac{1}{2} (\text{p}K_w - \text{p}K_a - \log C) = \frac{1}{2} [14 - 5 - 1] = 9$$

18. (c) Moles of H^+ produced 4×10^{-3}

$$[\text{H}^+] = \frac{4 \times 10^{-3}}{0.1} = 0.04 \text{ M}$$




Initial milli-moles

0.02 0.04

Final milli-moles

— 0.02 0.02



Initial milli-moles

0.06 0.02 —

Final milli-moles

0.04 — 0.02

Finally buffer solution of H_2PO_4^- (acid) and HPO_4^{2-} (conjugate base) is formed

$$\text{pH} = \text{p}K_{a_2} + \log \frac{[\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]} = 7.2 + \log \frac{0.04}{0.02} = 7.5$$

19. (b)



at equilibrium point $N_1 V_1 = N_2 V_2$; $V_2 = 20 \text{ mL}$

$$[\text{BCl}] = \frac{20}{20} \cdot \frac{0.08}{20} = 0.04$$

$$\text{pH} = \frac{1}{2} [\text{p}K_w - \text{p}K_b - \log (C)]$$

$$\text{p}K_b = 5.4$$

B OH BOH; Basic Buffer is formed

Initial milli-moles

1.6 0.4

Final milli-moles

1.2 — 0.4

$$\text{pOH} = \text{p}K_b + \log \frac{[B]}{[BOH]}$$

$$5.4 + \log \frac{1.2}{0.4} = 5.4 + 0.48$$

$$\text{pOH} = 5.88$$

20. (b)



Initial milli-moles

25 0.04 50 0.025 1

Final milli-moles

— 0.25



Initial milli-moles

1 0.25

Final milli-moles

0.75 — 0.25

$$\text{pH} = \text{p}K_{a_1} + \log \frac{[\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]} = 6.4 + \log \frac{0.75}{0.25} = 6.88$$

21. (b) Let a is initial milli-moles of HA and molarity of NaOH solution is x



In both case solution is acidic, so acidic buffer solution is formed.

$$5 - \text{p}K_a = \log \frac{10x}{a - 10x}$$

$$\begin{aligned}
 5.60 & \quad pK_a \quad \log \frac{20x}{a - 20x} \\
 0.60 & \quad \log \frac{20x}{(a - 20x)} \quad \frac{(a - 10x)}{10x} \\
 4 & \quad \frac{2(a - 10x)}{(a - 20x)} \quad a - 30x \\
 5 & \quad pK_a \quad \log \frac{10x}{20x}
 \end{aligned}$$

or

$$pK_a \quad 5.3$$

22. (d)

	CO_3^{2-}	H	HCO_3^-
Initial milli-moles	50 0.05	40 0.1	—
Final milli-moles	—	1.5	2.5
	HCO_3^-	H	H_2CO_3
Initial milli-moles	2.5	1.5	—
Final milli-moles	1	—	1.5

$$\text{pH} = pK_{a_1} + \log \frac{[\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]} = 6.173$$

23. (c) At second equivalence point the only species present at appreciable concentration is HA^{2-}

So,


$$\text{pH} = \frac{pK_{a_2} + pK_{a_3}}{2} = \frac{8 + 12}{2} = 10$$

$$[\text{H}^+] = 10^{-10}$$

and

$$\frac{[\text{H}_3\text{A}]}{[\text{A}^{3-}]} = \frac{[\text{H}^+]^3}{K_{a_1}K_{a_2}K_{a_3}} = 10^{-7}$$

24. (c) Solubility $S = \frac{x}{M}$



$$\begin{aligned}
 & \quad \quad \quad 3S \quad \quad \quad 2S \\
 & \quad \quad \quad K_{sp} \quad (3S)^3 (2S)^2 = 108S^5 \\
 & \quad \quad \quad \frac{[\text{B}^{3+}]}{K_{sp}} = \frac{2S}{108S^5} = \frac{1}{54} \frac{M^4}{x^4}
 \end{aligned}$$

25. (d)

$$K_{sp}(\text{SrCrO}_4) = [\text{Sr}^{2+}][\text{CrO}_4^{2-}]$$

$$[\text{CrO}_4^{2-}] = \frac{3.5 \times 10^{-5}}{0.1} = 3.5 \times 10^{-4}$$

$$K_{sp}(\text{BaCrO}_4) = [\text{Ba}^{2+}][\text{CrO}_4^{2-}]$$

$$[\text{CrO}_4^{2-}]_{\text{total}} \sim [\text{CrO}_4^{2-}] \text{ from } \text{SrCrO}_4$$

$$[\text{Ba}^{2+}] = \frac{1.2 \times 10^{-10}}{3.5 \times 10^{-4}} = 3.4 \times 10^{-7}$$

27. (a) $[\text{Ag}^+]$ required for precipitation of AgCl

$$[\text{Ag}^+] = \frac{K_{sp}(\text{AgCl})}{[\text{Cl}^-]} = 10^{-9}$$

$[\text{Ag}^+]$ required for precipitation of Ag_2CrO_4


$$\sqrt{\frac{K_{sp}(\text{Ag}_2\text{CrO}_4)}{[\text{CrO}_4^{2-}]}} = \sqrt{\frac{10^{-12}}{10^{-4}}} = 10^{-4}$$

At this point, the concentration of Cl^- ion in the solution can be calculated from $K_{sp}(\text{AgCl})$

$$[\text{Cl}^-] = \frac{K_{sp}(\text{AgCl})}{[\text{Ag}^+]} = \frac{10^{-10}}{10^{-4}} = 10^{-6} \text{ M}$$


28. (b) After mixing $[\text{Ag}^+] = 0.2 \text{ M}$; $[\text{NH}_3] = 1 \text{ M}$

Due to very high value of K_f , Ag^+ mainly converted into complex



Initial concentration 0.2 1

At equilibrium x 0.6 ~ 0.2




At equilibrium 0.2 - y y 0.6 - y

~ 0.2 ~ 0.6


$$\frac{1}{K_{f2}} = \frac{y \cdot 0.6}{0.2} = \frac{0.1}{10^4}; y = [\text{Ag}(\text{NH}_3)^+] = 3.33 \times 10^{-5} \text{ M}$$

29. (a)



$$K_1 = \frac{1.6 \times 10^{-10} \times 10^7}{1.6 \times 10^{-3}}$$

$$\frac{[\text{Ag}(\text{NH}_3)_2^+][\text{Cl}^-]}{[\text{NH}_3]^2}$$



$$K_2 = \frac{2.5 \times 10^{-16} \times 10^7}{2.5 \times 10^{-9}}$$

$$\frac{[\text{Ag}(\text{NH}_3)_2^+][\text{CN}^-]}{[\text{NH}_3]^2}$$




$$\frac{[\text{Cl}^-]}{[\text{CN}^-]} = \frac{1.6 \times 10^{-3}}{2.5 \times 10^{-9}} = 6.4 \times 10^5$$

$$K_1 = \frac{x^2}{(1-2x)^2} = \frac{x}{1-2x} = 0.04$$

$$x = 0.037$$

$$[\text{CN}^-] = \frac{0.037}{6.4 \times 10^5} = 5.78 \times 10^{-8}$$



32. (c)



$$0.1 - 2x \quad x \quad x$$

Value K_f is very higher so we can assume almost Ag^+ converts into complex

33. (d)



$$\frac{K_{sp1}}{K_{sp2}} = \frac{x}{y} \cdot \frac{2}{2} \cdot \frac{10^{-12}}{10^{-11}} = \frac{x}{y} \cdot 0.1$$

$$2 \cdot 10^{-11} (2x - 2y)^2 y$$

$$2 \cdot 10^{-11} \cdot 4.84y^3$$

$$y = 1.6 \cdot 10^{-4};$$

$$x = 0.16 \cdot 10^{-4}$$

$$\text{Total } [\text{Ag}^+] = 2x + 2y$$

$$(2 \cdot 0.16 + 2 \cdot 1.6) \cdot 10^{-4}$$

$$3.52 \cdot 10^{-4}$$

 34. (b) $K_{sp} [\text{Zn}^{2+}][\text{S}^{2-}]$

$$[\text{S}^{2-}] = \frac{10^{-21}}{0.01} = 10^{-19}$$

$$\text{for } K_{a1} K_{a2} \frac{[\text{H}^+]^2 [\text{S}^{2-}]}{[\text{H}_2\text{S}]}$$

$$10^{-20} \frac{[\text{H}^+]^2 \cdot 10^{-19}}{0.1} \quad [\text{H}^+] = 0.1$$

$$\text{or } \text{pH} = 1$$

 35. (d) Dissolved $\text{Al}(\text{OH})_3$ present in solution as $\text{Al}^{3+}(aq.)$ as well as $[\text{Al}(\text{OH})_4]^{-}(aq.)$

$$\text{S} = [\text{Al}^{3+}(aq.)] + [\text{Al}(\text{OH})_4^{-}(aq.)]$$

$$S = \frac{K_{sp}}{[\text{OH}^-]^3} + K_c [\text{OH}^-]$$

for minimum solubility

$$\frac{dS}{d(\text{OH}^-)} = 0$$

$$\text{or } \frac{K_{sp}}{[\text{OH}^-]^4} - K_c = 0$$

$$[\text{OH}^-] = \left(\frac{3K_{sp}}{K_c} \right)^{1/4}$$

Level 3

Passage-1

$$2. (b) [\text{HCl}] = \frac{0.01}{0.1} = 0.1 \text{ M}$$

$$\text{pOH} = 4.74 \quad \log \frac{0.3}{0.2} = \frac{0.1}{0.1} \quad 5.34$$

$$\text{pH} = 14 - 5.34 = 8.66$$

Passage-3

 3. (c) When acid is half neutralized $\text{pH} = \text{p}K_a$

$$\text{p}K_a = 5$$

$$4. (a) K_{eq} = \frac{K_a(\text{HB})}{K_a(\text{HA})} = \frac{10^{-5}}{10^{-6}} = 10$$

 5. (b) When $V_{\text{NaOH}} = 0 \text{ mL}$; $\text{pH} = 3$
 or $[\text{H}^+] = 0.001$

$$K_{a(\text{HB})} = \frac{C}{(1 - \alpha)} = \frac{(0.001)}{1} = 10^{-5}$$

$$0.01$$

$$C = \frac{0.001}{0.01} = 0.1$$

$$C = \frac{n_{\text{HB}}}{V}$$

$$0.1 = \frac{5}{V}; V = 50 \text{ mL}$$

$$[\text{NaB}] = \frac{\text{milli-moles of acid}}{\text{total volume}}$$

$$\frac{5}{50 + 50} = 0.05$$

$$\text{pH} = \frac{1}{2} [\text{p}K_w + \text{p}K_a + \log C]$$

$$\frac{1}{2} [14 - 5 + \log (0.05)]$$

$$8.85$$


 6. (a) For best indicator, $\text{pH} \sim \text{p}K_{\text{In}}$ of an indicator as well as pH transition range of the indicator must coincide with the steep portion of the titration curve.

One or More Answers is/are Correct

21. (b,c,d)




$$(a) S = \sqrt{K_{sp}} = 10^{-5} \text{ mol lit}^{-1}$$

$$(b) s = 10^{-5} \text{ mol lit}^{-1}$$



$$S = 2S$$

$$10^{-10} = (S + 2S)S \quad S = \frac{10^{-10}}{2} = 5 \cdot 10^{-11}$$



Subjective Problems

- $[\text{OH}^-] = \sqrt{K_b \cdot c}$
- $[\text{H}^+] = 10^{-4} \text{ M}$; C
 $\therefore 0.05 = K_a \cdot \frac{C}{1}$
- Consider ionisation of water.
- $\text{H}^+ + \text{OH}^- \rightleftharpoons \text{H}_2\text{O}$
 (2.5 - 1) mole (2.5) m mole
 L.R.
 $[\text{H}^+]_{\text{Final}} = \frac{(3.5 - 2.5)}{25 - 50 - 25} = 10^{-3} \text{ M}$
- $[\text{H}^+]_{\text{Final}} = \sqrt{K_{a1} \cdot c_1 \cdot K_{a2} \cdot c_2}$
- $C = \frac{0.16/32}{500/1000} \text{ M}$
 $\sqrt{K_b \cdot c}$

7. $pOH = pK_b + \log \frac{[NH_4Cl]}{[NH_4OH]}$
8. $pH = pK_a + \log \frac{[Salt]}{[Acid]}$
9. For WASB salt
 $[OH^-] = \sqrt{\frac{K_w \cdot c}{K_a}}$
10. $HX(aq.) \rightleftharpoons H^+(aq.) + X^-(aq.); K_a = 10^{-5}$
 $H^+(aq.) + OH^-(aq.) \rightleftharpoons H_2O(l); K_w = 10^{-14}$
 Addition
 $HX(aq.) + OH^-(aq.) \rightleftharpoons H_2O(l) + X^-(aq.)$
 $K_{eq} = \frac{K_a \cdot K_w}{K_a}$
11. $AgCl(s) \rightleftharpoons Ag^+(aq) + Cl^-(aq)$
 $K_{sp} = [Ag^+][Cl^-]$
12. Upon evaporation of saturated solution, all water will be escaped and only dissolved $PbCl_2$ will be left.
 i.e., Solubility $= \frac{2.78}{278} \times 10^2 \text{ mol/L}$
 for $PbCl_2$ $K_{sp} = 4s^3$
 $y \times 10^{-6} = 4(10^{-2})^3$
 $y = 4$
13. Let's assume simultaneous solubility of $SrCO_3$ as $x \text{ M}$ while SrF_2 as $y \text{ M}$
 $SrCO_3(s) \rightleftharpoons Sr^{2+}(aq) + CO_3^{2-}(aq)$
 At eq. $(x - y) \text{ M} \quad \text{M} \quad ; K_{sp1} = (x - y)x$
 $SrF_2(s) \rightleftharpoons Sr^{2+}(aq) + 2F^-(aq) ; 2.5 \times 10^{-10} = (x - y)x \quad \dots(1)$
 At eq. $(y - x) \text{ M} \quad 2y \text{ M} \quad ; K_{sp2} = (x - y)(2y)^2$
 $10^{-10} = 4y^2(x - y) \quad \dots(2)$
 $2.5 = \frac{x}{4y^2}$
 $10y^2 = x \times 10^{-3}$
 $y^2 = 10^{-4}$
 $y = 10^{-2} \text{ M} \quad [F^-] = 2y = 2 \times 10^{-2} \text{ M}$
14. $H_2CO_3 + NaOH \rightleftharpoons NaHCO_3 + H_2O$
 $pH_1 = \frac{1}{2}(pK_{a1} + pK_{a2}) = \frac{1}{2}(4.6 + 8) = 6.3$
 $\frac{N_1 V_1}{N_1} = \frac{N_2 V_2}{N_2}$
 $\frac{10}{10} = \frac{0.1}{20}$
 $N_1 = 0.2$
 $NaHCO_3 + NaOH \rightleftharpoons Na_2CO_3 + H_2O$
 $c = \frac{2m \text{ mole}}{50 \text{ mL}} = \frac{1}{25} \text{ M}$
 $pH_2 = pH_1 = 10.3 - 6.3 = 4$
15. $NaCN; Na_3PO_4; Na_2CO_3; NaHCO_3; Na_2C_2O_4; Na_2HPO_4$

7

CHEMICAL KINETICS & NUCLEAR CHEMISTRY

Rate of Appearance or Disappearance of a Substance C

(i) Average $\frac{[c]}{t}$

(ii) Instantaneous $\frac{d[c]}{dt}$

❖ Expressions of the rate:

For a general reaction: $aA + bB \rightarrow cC + dD$,

Rate of disappearance of A $\frac{d[A]}{dt}$; Rate of disappearance of B $\frac{d[B]}{dt}$

Rate of appearance of C $\frac{d[C]}{dt}$ & Rate of appearance of D $\frac{d[D]}{dt}$

The positive sign shows that concentrations of C and D increases with time and the negative sign is indicating that concentrations of A and B decrease with time.

Instantaneous rate of reaction : $\frac{1}{a} \frac{d[A]}{dt} = \frac{1}{b} \frac{d[B]}{dt} = \frac{1}{c} \frac{d[C]}{dt} = \frac{1}{d} \frac{d[D]}{dt}$

Average rate of reaction: $\frac{1}{a} \frac{[A]}{t} = \frac{1}{b} \frac{[B]}{t} = \frac{1}{c} \frac{[C]}{t} = \frac{1}{d} \frac{[D]}{t}$

Rate of reaction (ROR) $= \frac{\text{Rate of disappearance of reactant}}{\text{Stoichiometry coefficient of reactant}} = \frac{\text{Rate of appearance of product}}{\text{Stoichiometry coefficient of product}}$

❖ **Units of Rate of Reaction:** $\text{mol L}^{-1} \text{s}^{-1}$ or $\text{mol L}^{-1} \text{min}^{-1}$ (concentration time⁻¹).

Rate Law and Rate Constant

$$\text{Rate} = k[A]^a[B]^b \text{ or } \text{Rate} = k[A]^a[B]^b$$

The constant of proportionality, k is known as the rate constant (specific reaction rate) and may be defined as the rate at unit concentrations of the reactants.

- ❖ k depends on the temperature and is independent of the concentration of the reactants.
- ❖ At a fixed temperature, k is a constant and is a characteristic of the reaction. Larger value of k indicates fast reaction and small k indicates slow reactions.

Molecularity

It is always a whole number (not zero) and never a fraction. Its value does not exceed 3 & it has no meaning for a complex reaction.

Order of Reaction

$$\text{Rate of reaction} = k[A]^m[B]^n$$

Order of reaction $m + n$ & the order w.r.t. A, B are m, n respectively.

- ❖ The order of reaction is obtained from the experimentally determined rate and may be zero, integral or a fraction. In a multi-step complex reaction, the order of the reaction can be determined with the help of slowest step, which is called rate determining step.
- ❖ In elementary reaction stoichiometric coefficient of reactants is equal to order of reaction : $m + n = a + b$.

Order	Differential rate law	Integrated rate law	Straight line plot	Half life	Units of k	Example
0	$d[R]/dt = k$	$kt = [R]_0 - [R]$	$[R]_0$ vs t	$[R]_0 / 2k$	conc $\times$ time ⁻¹ or mole $L^{-1} s^{-1}$	$H_2(g) + Cl_2(g) \xrightarrow{h\nu} 2HCl(g)$ $2HI(g) \xrightarrow[Au \text{ surface}]{} H_2(g) + I_2(g)$
1	$d[R]/dt = k[R]$	$kt = \ln\{[R]_0/[R]\}$	$\ln [R]$ vs t	$(\ln 2)/k$ or $\frac{0.693}{k}$	time ⁻¹ or s^{-1}	Decomposition of H_2O_2 in aqueous solution Radioactive disintegration
2	$d[R]/dt = k[R]^2$	$kt = \frac{1}{[R]} - \frac{1}{[R]_0}$	$\frac{1}{[R]}$ vs t	$\frac{1}{k[R]_0}$	$\text{mol}^{-1} L s^{-1}$	$CH_3COOC_2H_5 + NaOH \rightarrow CH_3COONa + C_2H_5OH$ $C_2H_4 + H_2 \xrightarrow{100^\circ C} C_2H_6$

Important Graphical Representation

where a_0 Initial Concentration of reactant, r rate of reaction

❖ Some useful relationships between times for different fractions of reaction of first order to complete

$$\begin{aligned}
 t_{3/4} \text{ or } t_{75\%} &= 2t_{1/2} \\
 t_{87.5\%} &= 3t_{1/2} \\
 t_{99.9\%} &= 10t_{1/2}
 \end{aligned}$$

❖ Amount of the substance left after n half-lives $\frac{A_0}{2^n}$

 n^{th} Order Reaction


A Product

$$\begin{aligned}
 k &= \frac{1}{n-1} \frac{1}{(a-x)^{n-1}} - \frac{1}{a^{n-1}} \quad [n=1, n \text{ order}] \\
 t_{1/2} &= \frac{1}{k(n-1)} \frac{2^{n-1}-1}{a^{n-1}}
 \end{aligned}$$

❖ **Reversible reaction** $A \xrightleftharpoons[k_b]{k_f} B$


$$(k_f + k_b) \frac{1}{t} \ln \frac{x_{eq}}{x_{eq}-x} \quad ; \quad \text{At equilibrium } k_{eq} = \frac{k_f}{k_b} = \frac{[B]}{[A]} = \frac{x_{eq}}{a-x_{eq}}$$

Parallel/Side Reactions



$$\% \text{ Yield of B} = \frac{k_1}{k_1 + k_2} \times 100 \quad \text{and} \quad \% \text{ Yield of C} = \frac{k_2}{k_1 + k_2} \times 100$$

Series/Consecutive Reactions



$$t_{\max} = \frac{1}{k_1 - k_2} \ln \frac{k_1}{k_2}$$

$$\text{Temperature coefficient ()} = \frac{k_T + 10}{k_T}$$

- ❖ For general chemical reactions with rise in temperature by 10°C , the rate constant is nearly doubled.

Arrhenius Equation

$$k = A e^{-E_a/RT}$$


where k = Rate constant, A = Pre-exponential factor, T = Temperature in kelvin,
 E_a = Activation Energy

- ❖ The factor $e^{-E_a/RT}$ represents fraction of molecules that have kinetic energy greater than E_a

❖ Logarithmic expression

$$\log_{10} \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$


- ❖ Energy profile for exothermic process & endothermic process:



- ❖ Threshold Energy = Activation Energy = Energy possessed by reactant molecules.

❖ **Activation energy diagram for two step endothermic reaction:**

- (A) (i) I step-Slow (ii) II step-Fast
 (B) (i) Rapid equilibrium (ii) Slow



Factors affecting reaction rates : Concentration of reactants and Reaction temperature. Besides these, presence of catalyst and surface area (if a reactant or a catalyst is a solid) exposure to radiation also affect the reaction rates.

Nuclear Chemistry

Nuclear chemistry is the study of the properties and reactions of atomic nuclei. In nuclear reaction only the nuclides (nuclide : the nucleus of a specific isotope) participates and overall (mass + energy) remains conserved.

Radioactivity

The property of a nucleus emitting radiations like α , β and γ is known as radioactivity and the substance possessing the property is called a radioactive substance. The emission of these particles takes place because of the instability of the nucleus. Radioactivity is a property of nucleus.

Properties of α , β -particles and γ -rays.

Properties	Alpha	Beta	Gamma
Nature	Fast moving He nuclei (He^2)	Fast moving electrons	High energy electromagnetic
Representation	${}_2^4\text{He}$ or	${}_1^0e$ or ${}_1^0\beta$	or ${}_0^0\gamma$
Charge	2 unit (+ve)	1 unit (-ve)	No charge
Mass	6.65×10^{-24} g/particle	9.11×10^{-28} g/particle	0
Velocity	1/10 of light (min.)	nearly velocity of light	Same as light waves (max.)
Relative penetrating power	1	100	10000
Relative ionising power	10000	100	1
Effect of electromagnetic field	Deflected toward -ve electrode	Deflected toward +ve electrode	No effect
Kinetic energy	High	Low	-
Effect on ZnS Screen	Maximum effect	Little effect	-


Types of Radioactive Decay

1. **α -decay** causes decrease of atomic number of 2 units and mass number by 4 units, e. g.,




All nucleides with atomic number greater than 83 are beyond the band of stability and are radioactive.

2. **-decay** causes increase of atomic number by 1 unit and no change in mass number, e. g.,



A neutron is converted to proton in this process.




Such emission occurs for the nuclei lying above the stability belt.


3. **-ray** emission (${}_0^0\gamma$) causes no change in atomic number and mass number, since it represents the energy lost, e. g.,



4. **Positron Emission** : Causes decrease the atomic number by 1 unit, e. g.,



A proton is converted to neutron and positron in this process.




Such emission occurs for the nuclei lying below the stability belt.


- ❖ Positron (${}_1^0e$) is a particle having the same mass as an electron, but positively charged.

5. **Electron-capture** :

Causes decrease the atomic number by 1 unit, e. g.,



A proton is converted to neutron in this process.



Electron capture occurs with the nuclei lying below the stability belt, in which an electron from the K-shell is captured by the nucleus.

Group Displacement Law (Soddy-Fajans)


In an α -particle emission, the resulting element has a mass number less by four units and atomic number less by two units and so lies two places to the left in the periodic table.

In a β -particle emission the resulting element has the same mass number but has an atomic number greater by one unit and so lies one place to the right in the periodic table.

Neutron/proton ratio and stability zone

- ❖ For atomic number < 20 , most stable nuclei have $n : p$ ratio nearly $1 : 1$ (except H & Ar)
- ❖ For n/p ratio > 1.5 , nucleus is unstable. Largest stable nucleus is $^{208}_{82}\text{Pb}$ for which n/p ratio is 1.53.
- ❖ For atomic number > 82 , there are no stable nuclei.

Note: Heaviest stable nuclide is $^{208}_{82}\text{Pb}$ $t_{1/2}$ for $^{208}_{82}\text{Pb}$ 1.9 10^{19} yr., which is α -emitter reference. Pierre De Marcillae exp det.... nature, 422, 876-878, 2003



Magic numbers and nuclear stability

Nuclei with 2, 8, 20, 28, 50, 82 or 126 protons or neutrons are unusually stable and have a larger number of stable isotopes than neighboring nuclei in the periodic table. These numbers are called magic numbers. They are supposed to represent completely filled nuclear shells of energy levels.


- ❖ Nuclei with magic number of protons as well as neutrons have notably high stabilities. [e.g., ^4_2He , $^{16}_8\text{O}$, $^{40}_{20}\text{Ca}$ and $^{208}_{82}\text{Pb}$].

Even Odd Theory

Most naturally occurring nuclides have even numbers of neutrons and even numbers of protons. 165 such stable nuclei are known. There exist 55 known nuclei with even number of protons and odd number of neutrons, and 50 known stable nuclei with odd number of protons and even number of neutrons. On the other hand, the number of known stable nuclei having odd numbers of both neutrons and protons is only 5.

Artificial nuclear reactions

The first artificial transmutation was carried out by Rutherford in 1919 who bombarded nitrogen gas with alpha particles and obtained hydrogen and oxygen.



1. Alpha particle induced or (α , n) type reactions :



Since α -particle is used and a neutron is produced, the reaction may be termed as (α , n) reaction. In another α -bombardment nuclear reaction, the isotope ($^{30}_{15}\text{P}$) produced is itself radioactive.



2. Deuteron-induced or (D,) type reaction :



$${}_{7}^{14}\text{N} \quad {}_{1}^{1}\text{H} \quad {}_{8}^{15}\text{O}$$
$${}_{15}^{31}\text{P} \quad {}_0^1n \quad {}_{15}^{32}\text{P}$$

Hence $\frac{dN}{dt} = N$ or $N = N_0 e^t$

$$A \quad A \quad B \quad B \quad C$$

Number of nuclei of B is max. at $t_{\max}$; $t_{\max} = \frac{1}{(\lambda_1 - \lambda_2)} \ln \frac{\lambda_1}{\lambda_2}$

$$\frac{dN_B}{dt} = -k_A N_A - k_B N_B.$$

Secular equilibrium occurs when $\frac{dN_B}{dt} = 0$ or $\frac{N_B}{N_A} = \frac{\lambda_A}{\lambda_B}$

❖ Parallel decay

A	1	B
	2	C

$$\begin{array}{rcc} \% \text{ of } B & \frac{1}{1 \quad 2} & 100 \end{array}$$

$$\begin{array}{rcc} \% \text{ of } C & \frac{2}{1 \quad 2} & 100 \end{array}$$

Applications of Radioactivity :

- ❖ **Age of Minerals and Rocks :** We can determine the age of the rock by measuring the ratio of ${}_{92}^{238}\text{U}$ and ${}_{82}^{206}\text{Pb}$.



At time t w gram y gram

Age of Rock, $t = \frac{1}{\lambda} \ln \frac{w_0}{w}$ where $w_0 = w + x$,

w_0 Initial mass of ${}_{92}^{238}\text{U}$; x mass of ${}_{92}^{238}\text{U}$ disintegrated. $x = \frac{238}{206} y$

- ❖ **Radiocarbon Dating :** ${}^{14}_6\text{C}$ is used to date historical artifacts.

Age of wood artifacts, $t = \frac{1}{\lambda} \ln \frac{A_0}{A}$ (where $t_{1/2}$ of ${}^{14}_6\text{C}$ = 5770 years)


A_0 Activity of fresh wood of plant/tree. ; A Activity of old wood.

- ❖ Uses in Medicines and other Areas

Nuclear Fission

In a nuclear fission reaction, a heavy nucleus splits up into two main fragments of lighter nuclei and several neutrons.

Of the three natural isotopes of uranium (${}_{92}^{238}\text{U}$, ${}_{92}^{235}\text{U}$ and ${}_{92}^{234}\text{U}$) the ${}_{92}^{235}\text{U}$ nucleus undergoes nuclear fission when bombarded with slow neutrons. The ${}_{92}^{236}\text{U}$ formed breaks up in several different ways, for example :



- ❖ The key to the liberation of energy in the nuclear fission reaction is the production of two or more neutrons per reaction initiated by one neutron.
- ❖ Atom bomb is based on nuclear fission.

A nuclear fission chain reaction has three general steps :

1. Initiation : The reaction of a single atom is needed to start the chain reaction. Fission of ${}_{92}^{235}\text{U}$ is initiated by the absorption of a neutron.

2. Propagation : This part of the process repeats itself over and over, with each step yielding more product. The fission of ${}_{92}^{236}\text{U}$ releases neutrons that initiate the fission of other uranium atoms.

3. Termination : Eventually, the chain will end. Termination could occur if the reactant (${}_{92}^{235}\text{U}$) is used up, or if the neutrons that continue the chain escape from the sample without being captured by ${}_{92}^{235}\text{U}$.

Nuclear Fusion

Just as the fission of heavy nuclei is accompanied by mass losses resulting into the liberation of large amounts of energy, the fusion of light nuclei is accompanied by mass losses and the evolution of large quantities of energy. Some such reactions and energy release in each process are shown below,

Fusion reaction				Mass loss	Energy released (kJ mol ⁻¹)
${}^2_1\text{H}$	${}^2_1\text{H}$	${}^4_2\text{He}$		0.026	2.3×10^9
${}^2_1\text{H}$	${}^3_1\text{H}$	${}^4_2\text{He}$	1_0n	0.018	1.79×10^9

Compared with fission reactions, fusion reactions have the advantage that large amounts of highly radioactive nuclides are not obtained as by-products which may pose problem of safe storage.

- ❖ Very high temperatures is required for a fusion reaction.
- ❖ Hydrogen bomb is based on nuclear fusion.
- ❖ **Einstein's mass-energy equation** : Energy changes in nuclear reactions can determined by using Einstein's mass-energy equation $E = mc^2$

Where E is the energy equivalent of mass m and c is the velocity of light.

For a change of 1 amu (atomic mass unit), the corresponding energy change, $E = 931 \times 10^6 \text{ eV} = 931 \text{ MeV}$, i.e., a mass of 1 amu is equivalent to 931 MeV of energy.

Mass Defect

It has invariably been found that the actual mass of an isotope of an element is less than the sum of masses of the protons, neutrons and electrons present in it. This difference is called mass defect. The mass defect is nothing, but the loss of mass during the formation of the nucleus of the isotope.

- ❖ **Mass defect in nuclear reaction** :

m = mass of nuclei of reactants – mass of nuclei of products.

Energy liberated in nuclear reaction : $E = m \cdot c^2$.

- ❖ **Mass defect in an isotope formation** :

Let m_p, m_n and m_e are the respective masses of proton, neutron and electron. Then, the calculated mass of this isotope,

$$M = Zm_p + Zm_e + (A - Z)m_n + Zm_H + (A - Z)m_n$$

($\because m_p + m_e$ = Mass of hydrogen atom m_H)

Let M = Actual atomic mass as determined experimentally.

then, m = Mass defect $M - M$

Binding Energy

Loss of mass during the formation of the nucleus from nucleons is converted into energy. The release of energy imparts stability to the nucleus. The energy released when constituent nucleons combine to form a nucleus, is called binding energy of the nucleus. In other words, energy equal to binding energy will be needed to break up the nucleus into its constituent nucleons. Consequently, the greater the binding energy, the more stable is the nucleus.

Binding Energy and Nuclear Stability

$$\text{B.E. per nucleon} = \frac{\text{B.E.}}{\text{No. of nucleons}}$$

Binding energy per nucleon is a direct indicator of its nuclear stability. Higher the binding energy per nucleon of an isotope, greater is its nuclear stability.



Level 1

- The differential rate law equation for the elementary reaction $A + 2B \xrightarrow{k} 3C$, is :
 - $\frac{d[A]}{dt} = \frac{d[B]}{dt} = \frac{d[C]}{dt} = k[A][B]^2$
 - $\frac{d[A]}{dt} = \frac{1}{2} \frac{d[B]}{dt} = \frac{1}{3} \frac{d[C]}{dt} = k[A]^2[B]$
 - $\frac{d[A]}{dt} = \frac{1}{2} \frac{d[B]}{dt} = \frac{1}{3} \frac{d[C]}{dt} = k[A][B]^2$
 - None of these
- The rate of a reaction is expressed in different ways as follows:





$$\frac{1}{2} \frac{d[C]}{dt} = \frac{1}{3} \frac{d[D]}{dt} = \frac{1}{4} \frac{d[A]}{dt} = \frac{d[B]}{dt}$$
 The reaction is:
 - $4A + B \rightarrow 2C + 3D$
 - $B + 3D \rightarrow 4A + 2C$
 - $A + B \rightarrow C + D$
 - $B + D \rightarrow A + C$
- In the reaction, $A + 2B \rightarrow 6C + 2D$, if the initial rate $\frac{d[A]}{dt}$ at $t = 0$ is $2.6 \times 10^{-2} \text{ M sec}^{-1}$, what will be the value of $\frac{d[B]}{dt}$ at $t = 0$?
 - $8.5 \times 10^{-2} \text{ M sec}^{-1}$
 - $2.5 \times 10^{-2} \text{ M sec}^{-1}$
 - $5.2 \times 10^{-2} \text{ M sec}^{-1}$
 - $7.5 \times 10^{-2} \text{ M sec}^{-1}$
- For the reaction $2A + B \rightarrow 3C$; if $\frac{d[A]}{dt} = k_1[A]^2$; $\frac{d[B]}{dt} = k_2[A]^2$; $\frac{d[C]}{dt} = k_3[A]^2$ the correct relation between k_1 , k_2 and k_3 is :
 - $k_1 = k_2 = k_3$
 - $2k_1 = k_2 = 3k_3$
 - $4k_1 = k_2 = 3k_3$
 - $\frac{k_1}{2} = k_2 = \frac{k_3}{3}$
- The rate constant of n^{th} order has units :
 - $\text{litre}^{1-n} \text{ mol}^{1-n} \text{ sec}^{-1}$
 - $\text{Mol}^{1-n} \text{ litre}^{1-n} \text{ sec}^{-1}$
 - $\text{Mol}^{1-n^2} \text{ litre}^{n^2} \text{ sec}^{-1}$
 - $\text{Mole}^{1-n} \text{ litre}^{n-1} \text{ sec}^{-1}$
- Which of the following statement is incorrect?
 - Unit of rate of disappearance is Ms^{-1}
 - Unit of rate of reaction is Ms^{-1}
 - Unit of rate constant k depends upon order
 - Unit of k for first order reaction is Ms^{-1}
- Which of the following relation is correct for k_f and k_b in an equilibrium process that contains equal moles of reactants and products.
 - $k_f = k_b$
 - $k_f < k_b$
 - $k_f > k_b$
 - we cannot predict
- Listed in the table are forward and reverse rate constants for the reaction

$$2\text{NO}(\text{g}) \rightleftharpoons \text{N}_2(\text{g}) + \text{O}_2(\text{g})$$

Temperature (K)	$k_f (\text{M}^{-1} \text{s}^{-1})$	$k_b (\text{M}^{-1} \text{s}^{-1})$
1400	0.29	1.1×10^6
1500	1.3	1.4×10^5

Select the correct statement :

- (a) Reaction is exothermic and value of equilibrium constant (K_{eq}) at 1400 K is 3.79×10^6
 (b) Reaction is endothermic and value of K_{eq} at 1400 K is 2.63×10^5
 (c) Reaction is exothermic and value of K_{eq} at 1400 K is 2.63×10^5
 (d) Reaction is endothermic and value of K_{eq} at 1500 K is 9.28×10^4
9. The rate constant of a reaction depends on
 (a) temperature
 (b) pressure
 (c) extent of reaction
 (d) initial concentration of the reactant
10. In the following reaction, how is the rate of appearance of the underlined product related to the rate of disappearance of the underlined reactant
- $$\text{BrO}_3(\text{aq}) + 5\text{Br}(\text{aq}) + 6\text{H}(\text{aq}) \rightarrow 3\text{Br}_2(\text{l}) + 3\text{H}_2\text{O}(\text{aq})$$
- (a) $\frac{d[\text{BrO}_3^-]}{dt} = \frac{d[\text{Br}_2]}{dt}$
 (b) $\frac{1}{3} \frac{d[\text{BrO}_3^-]}{dt} = \frac{d[\text{Br}_2]}{dt}$
 (c) $\frac{d[\text{BrO}_3^-]}{dt} = \frac{1}{3} \frac{d[\text{Br}_2]}{dt}$
 (d) None of these
11. Consider a reaction $\text{A}(\text{g}) \xrightarrow{k = 0.1 \text{ M min}^{-1}} 2\text{B}(\text{g})$. If initial concentration of A is 0.5M then select correct graph.



12. Which of the following statements is incorrect?

- (a) A second order reaction must be a bimolecular elementary reaction
- (b) A bimolecular elementary reaction must be a second order reaction
- (c) Zero order reaction must be a complex reaction
- (d) First order reaction may be complex or elementary reaction


13. The molecularity of a complex reaction given below is :




- (a) 1
- (b) 2
- (c) 3
- (d) has no meaning

14. Decomposition of $\text{NH}_4\text{NO}_2(\text{aq})$ into $\text{N}_2(\text{g})$ and $2\text{H}_2\text{O}(\text{l})$ is first order reaction.


Which of the following graph is correct?



15. Decomposition of $\text{HI}(\text{g})$ on Gold surface is zero order reaction. Initially few moles of H_2 are present in container then which of the following graph is correct?



16. Consider the plots for the types of reaction





These plots respectively correspond to the reaction orders :


- (a) 0, 2, 1
- (b) 0, 1, 2
- (c) 1, 1, 2
- (d) 1, 0, 2

17. If decomposition reaction $\text{A}(\text{g}) \rightarrow \text{B}(\text{g})$ follows first order kinetics then the graph of rate of formation (R) of B against time t will be:






18. Consider the plots, given below, for the types of reaction




These plots respectively correspond to the reaction orders :

- (a) 0, 1, 2 (b) 1, 2, 0 (c) 1, 0, 2 (d) None of these
19. For a zero order reaction, the plot of conc. (a) vs time is linear with
- (a) ve slope and zero intercept (b) ve slope and non-zero intercept
(c) +ve slope and zero intercept (d) +ve slope and non-zero intercept

20. What will be the order of reaction for a chemical change having $\log t_{1/2}$ vs $\log a$? (where a = initial concentration of reactant; $t_{1/2}$ half-life)




21. A graph between $\log t_{1/2}$ and $\log a$ (abscissa), a being the initial concentration of A in the reaction For reaction A → Product, the rate law is :




- (a) $\frac{d[A]}{dt} = K$ (b) $\frac{d[A]}{dt} = K[A]$
(c) $\frac{d[A]}{dt} = K[A]^2$ (d) $\frac{d[A]}{dt} = K[A]^3$

22. For the reaction A → B, for which graph between half life ($t_{1/2}$) and initial concentration (a) of the reactant is as given below



Hence graph between $\frac{d[A]}{dt}$ and time will be:



23. For the ideal gaseous reaction, the rate is generally expressed in terms of $\frac{dP}{dt}$ instead of $\frac{dC}{dt}$ or $\frac{dn}{dt}$ (where $C = \frac{n}{V}$ is concentration and n the no. of moles). What is the relation among these three expressions if T and V are constant?

- (a) $\frac{dC}{dt} = \frac{dn}{dt} = \frac{dP}{dt}$ (b) $\frac{dC}{dt} = \frac{1}{V} \frac{dn}{dt} = \frac{1}{RT} \frac{dP}{dt}$
 (c) $RT \frac{dC}{dt} = \frac{dn}{dt} = \frac{dP}{dt}$ (d) None of these


24. $A_2 + B_2 \rightarrow 2AB$; R.O.R. $k[A_2]^a[B_2]^b$

Initial $[A_2]$	Initial $[B_2]$	R.O.R. (r) Ms^{-1}
0.2	0.2	0.04
0.1	0.4	0.04
0.2	0.4	0.08

Order of reaction with respect to A_2 and B_2 are respectively :

- (a) $a = 1, b = 1$ (b) $a = 2, b = 0$ (c) $a = 2, b = 1$ (d) None
25. For a reaction the initial rate is given as: $R_0 = k[A_0]^2[B_0]$. By what factor, the initial rate of reaction will increase if initial concentration of A is taken 1.5 times and of B is tripled?
 (a) 4.5 (b) 2.25 (c) 6.75 (d) None of these
26. For $A_{(s)} + B_{(s)} \rightarrow C_{(s)}$; rate $k[A]^{1/2}[B]^2$, if initial concentration of A and B are increased by factors 4 and 2 respectively, then the initial rate is changed by the factor:
 (a) 4 (b) 6 (c) 8 (d) None of these
27. Reaction $A \rightarrow B$ follows second order kinetics. Doubling the concentration of A will increase the rate of formation of B by a factor of :
 (a) $1/4$ (b) $1/2$ (c) 2 (d) 4

28. The reaction of A_2 and B_2 follows the equation



The following data were observed

$[A_2]_0$	$[B_2]_0$	Initial rate of appearance of $AB(g)$ (in $M s^{-1}$)
0.10	0.10	2.5×10^{-4}
0.20	0.10	5×10^{-4}
0.20	0.20	10×10^{-4}

The value of rate constant for the above reaction is:

- (a) 2.5×10^{-4} (b) 2.5×10^{-2} (c) 1.25×10^{-2} (d) None of these
29. Above given question, the value of rate constant for appearance of $AB(g)$ is :
 (a) 2.5×10^{-4} (b) 2.5×10^{-2} (c) 1.25×10^{-2} (d) None of these
30. The following data pertain to reaction between A and B

S. No	[A] mol L^{-1}	[B] mol L^{-1}	Rate $\text{mol L}^{-1} \text{sec}^{-1}$
I	1×10^{-2}	2×10^{-2}	2×10^{-4}
II	2×10^{-2}	2×10^{-2}	4×10^{-4}
III	2×10^{-2}	4×10^{-2}	8×10^{-4}

Which of the following inference(s) can be drawn from the above data

- (a) Rate constant of the reaction 10^{-4}
 (b) Rate law of the reaction is $k[A][B]$
 (c) Rate of reaction increases four times on doubling the concentration of both the reactant,
 Select the correct answer

codes

- (a) a, b and c (b) a and b (c) b and c (d) c alone

31. The unit of rate constant of elementary reaction depends upon the :

- (a) temperature of the reaction (b) concentration of reactants
 (c) activation energy of the reaction (d) molecularity of the reaction

32. Select the rate law that corresponds to the data shown for the reaction $A + B \rightarrow C$

Exp.	[A]	[B]	Rate
1.	0.012	0.035	0.10
2.	0.024	0.070	0.80
3.	0.024	0.035	0.10
4.	0.012	0.070	0.80

- (a) Rate $k[B]^3$ (b) Rate $k[B]^4$ (c) Rate $k[A][B]^3$ (d) Rate $k[A]^2[B]^2$

33. An elementary reaction between A and B is a second order reaction. Which of the following rate equations must be correct?

- (a) $r = k[A]^2[B]^0$ (b) $r = k[A]^{3/2}[B]^{1/2}$ (c) $r = k[A]^0[B]^2$ (d) $r = k[A][B]$

34. If 'a' is the initial concentration of the reactant, the half-life period of the reaction of n^{th} order is inversely proportional to :
 (a) a^{n-1} (b) a^n (c) a^{1-n} (d) a^{n-1}
35. Which of the following expressions is correct for zero order and first order respectively [where a is initial concentration]?
 (a) $t_{1/2} \propto a; t_{1/2} \propto \frac{1}{a}$ (b) $t_{1/2} \propto a; t_{1/2} \propto a^0$
 (c) $t_{1/2} \propto a^0; t_{1/2} \propto a$ (d) $t_{1/2} \propto a; t_{1/2} \propto \frac{1}{a^2}$
36. The unit of rate constant of zero order and first order chemical reactions are respectively :
 (a) $\text{mol L}^{-1} \text{s}^{-1}, \text{mol L}^{-1} \text{s}^{-1}$ (b) $\text{s}^{-1}, \text{mol L}^{-1} \text{s}^{-1}$
 (c) $\text{mol L}^{-1} \text{s}^{-1}, \text{s}^{-1}$ (d) None of these
37. The units of rate of reaction and rate constant are same for a :
 (a) zero order reaction (b) first order reaction
 (c) second order reaction (d) third order reaction
38. $\text{CH}_3\text{COOC}_2\text{H}_5(\text{aq}) + \text{H}_2\text{O}(\text{l}) \xrightarrow{\text{H}^+(\text{aq})} \text{CH}_3\text{COOH}(\text{aq}) + \text{C}_2\text{H}_5\text{OH}(\text{aq})$. What type of reaction is this?
 (a) Unimolecular elementary (b) Pseudo first order
 (c) Zero order (d) Second order
39. When ethyl acetate was hydrolysed in presence of 0.1 M HCl, the rate constant was found to be $5.4 \times 10^{-5} \text{s}^{-1}$. But in presence of 0.1 M H_2SO_4 the rate constant was found to be $6.25 \times 10^{-5} \text{s}^{-1}$. Thus it may be concluded that :
 (a) H_2SO_4 furnishes more H^+ than HCl
 (b) H_2SO_4 furnishes less H^+ than HCl
 (c) both have the same strength
 (d) will depend on concentration of ethyl acetate
40. For an elementary reaction $2A \rightarrow B + A_2B$ if the volume of vessel is quickly reduced to half of its original volume then rate of reaction will
 (a) remain unchanged (b) increase four times
 (c) increase eight times (d) decrease eight times
41. In the reaction $A + B \rightarrow C$, rate constant is 0.001 Ms^{-1} . If we start with 1 M of A then conc. of A and B after 10 minutes are respectively :
 (a) 0.5 M, 0.5 M (b) 0.6 M, 0.4 M (c) 0.4 M, 0.6 M (d) none of these
42. For a reaction $A \xrightarrow{k_r, 0.6 \text{ M min}^{-1}} 2B$ starting with 1 M of 'A' only, concentration of B (in M) after 100 sec. and 200 sec. is respectively ?
 (a) 2 and 4 (b) 1 and 2 (c) 2 and 3 (d) None of these
43. Half-life ($t_{1/2}$) and completion time (T) of the above reaction ($A \rightarrow 2B$) are :
 (a) 500 min, 750 min (b) 500 sec, 750 sec (c) 50 sec, 100 sec (d) None of these

44. Column I

Column II

P. Zero order reaction

1. $t_{1/2} \propto \frac{1}{[A]_0}$

Q. First order reaction

2. $t_{100\%} \propto [A]_0 / k$

R. Second order reactions

3. Involves at least two reactants

S. Pseudo unimolecular reaction

4. $[A] = [A]_0 e^{-kt}$

Code:

	P	Q	R	S
(a)	2	1	4	2
(b)	2	4	1	3
(c)	2	1	3	4
(d)	3	2	1	4

45. For the zero order reaction $A \rightarrow B + C$; initial concentration of A is 0.1 M. If $[A] = 0.08 \text{ M}$ after 10 minutes, then its half-life and completion time are respectively :

- (a) 10 min; 20 min (b) $2 \times 10^3 \text{ min}$; $4 \times 10^3 \text{ min}$
 (c) 25 min, 50 min (d) 250 min, 500 min

46. For an elementary reaction, $X(g) + Y(g) \rightarrow Z(g)$

the half life period is 10 min. In what period of time would the concentration of X be reduced to 10% of original concentration ?

- (a) 20 Min. (b) 33 Min (c) 15 Min (d) 25 Min

47. In the presence of an acid, the initial concentration of cane sugar was reduced from 0.20 to 0.10 molar in 5 hours and from 0.2 to 0.05 molar in 10 hours. The reaction is of :

- (a) Zero order (b) First order (c) Second order (d) Third order

48. A first order reaction is 75% completed in 100 minutes. How long time will it take for its 87.5% completion?

- (a) 125 min (b) 150 min (c) 175 min (d) 200 min

49. The rate constant for a first order reaction whose half life is 480 sec, is :

- (a) $1.44 \times 10^{-3} \text{ sec}^{-1}$ (b) 1.44 sec^{-1} (c) $0.72 \times 10^{-3} \text{ sec}^{-1}$ (d) $2.88 \times 10^{-3} \text{ sec}^{-1}$

50. Rate constant $k = 2.303 \text{ min}^{-1}$ for a particular reaction. The initial concentration of the reactant is 1 mol/litre then rate of reaction after 1 minute is :

- (a) 2.303 M min^{-1} (b) $0.2303 \text{ M min}^{-1}$ (c) 0.1 M min^{-1} (d) none of these

51. For the reaction $3A(g) \xrightarrow{k} B(g) + C(g)$, k is $10^{-4} \text{ L/mol.min}$.

If $[A] = 0.5 \text{ M}$ then the value of $\frac{d[A]}{dt}$ (in M s^{-1}) is:

- (a) 7.5×10^{-5} (b) 3×10^{-4} (c) 2.5×10^{-5} (d) none of these


52. 99% of a first order reaction was completed in 32 minutes when 99.9% of the reaction will complete :

- (a) 50 min (b) 46 min (c) 48 min (d) 49 min

53. Which of the following represents the expression for $\frac{3}{4}$ th life of first order reaction

- (a) $\frac{k}{2.303} \log 4/3$ (b) $\frac{2.303}{k} \log 3/4$ (c) $\frac{2.303}{k} \log 4$ (d) $\frac{2.303}{k} \log 3$


54. Consider the following first order competing reactions:



If 50% of the reaction of X was completed when 96% of the reaction of Y was completed, the ratio of their rate constants (k_2 / k_1) is:

- (a) 4.06 (b) 0.215 (c) 1.1 (d) 4.65

55. For the reactions



Calculate $\frac{k_I}{K_{II}}$, where k_I and k_{II} are rate constant for the respective reaction.

- (a) 2.303 (b) 1 (c) 0.36 (d) 0.693


56. The decomposition of N_2O_5 in carbon tetrachloride was followed by measuring the volume of O_2 gas evolved: $2N_2O_5(CCl_4) \rightarrow 2N_2O_4(CCl_4) + O_2(g)$. The maximum volume of O_2 gas obtained was 100 cm^3 . In 500 minutes, 90 cm^3 of O_2 were evolved. The first order rate constant (in min^{-1}) for the disappearance of N_2O_5 is:

- (a) $\frac{2.303}{500}$ (b) $\frac{2.303}{500} \log \frac{100}{90}$ (c) $\frac{2.303}{500} \log \frac{90}{100}$ (d) $\frac{100}{10 \cdot 500}$

57. For a first order homogeneous gaseous reaction $A \rightarrow 3B$, if pressure after time t was P_T and after completion of reaction, pressure was P then select correct relation

- (a) $k = \frac{1}{t} \ln \frac{P}{3(P - P_T)}$ (b) $k = \frac{1}{t} \ln \frac{2P}{(P - P_T)}$
 (c) $k = \frac{1}{t} \ln \frac{3P}{2P - P_T}$ (d) $k = \frac{1}{t} \ln \frac{2P}{3(P - P_T)}$


58. The half-life of first order decomposition of NH_4NO_3 is 2.10 hr at 288 K temperature



If 6.2 g of NH_4NO_3 is allowed to decompose, the time required for NH_4NO_3 to decompose 90% is:

- (a) 6.978 hr (b) 0.319 (c) 0.319 hr (d) None of these

59. For a first order homogeneous gaseous reaction, $A \rightleftharpoons 2B + C$ then initial pressure was P_i while total pressure after time ' t ' was P_t . The right expression for the rate constants k in terms of P_i, P_t and t is :
- (a) $k = \frac{2.303}{t} \log \frac{2P_i}{3P_i - P_t}$ (b) $k = \frac{2.303}{t} \log \frac{2P_i}{2P_t - P_i}$
- (c) $k = \frac{2.303}{t} \log \frac{P_i}{P_t}$ (d) none of these
60. The decomposition of azo methane, at certain temperature according to the equation $(CH_3)_2N_2 \rightarrow C_2H_6 + N_2$ is a first order reaction. After 40 minutes from the start, the total pressure developed is found to be 350 mm Hg in place of initial pressure 200 mm Hg of azo methane. The value of rate constant k is :
- (a) $2.88 \times 10^{-4} \text{ sec}^{-1}$ (b) $1.25 \times 10^{-4} \text{ sec}^{-1}$ (c) $5.77 \times 10^{-4} \text{ sec}^{-1}$ (d) None of these
61. The hydrolysis of sucrose was studied with the help of polarimeter and following data were collected
- | | | |
|-----------------------------|------|----------|
| time (min.) | : 0 | 70 |
| observed rotation (degrees) | : 44 | 16.5 -11 |
- The time taken when the reaction mixture will be optically inactive ? (Given: $\ln 2 = 0.7$, $\ln 3 = 1.1$, $\ln 5 = 1.6$)
- (a) 16 min. (b) 69.47 min. (c) 160 min. (d) none of these
62. For a particular reaction with initial conc. of the reactants as a_1 and a_2 , the half-life period are t_1 and t_2 respectively. The order of the reaction (n) is given by :
- (a) $n = 1 - \frac{\log(t_2/t_1)}{\log(a_2/a_1)}$ (b) $n = \frac{\log(t_1/t_2)}{\log(a_2/a_1)}$
- (c) $n = 1 - \frac{\log(t_1/t_2)}{\log(a_2/a_1)}$ (d) none of these
63. The value of $\frac{t_{0.875}}{t_{0.50}}$ for n^{th} order reaction is
- (a) $2^{(2n-2)}$ (b) $2^{(2n-2)-1}$ (c) $\frac{8^{n-1}}{2^{n-1}}$ (d) None of these
64. A B first order reaction A is optical active and B is optically inactive. A series of experiment were conducted on a solution of A
- | | | |
|------------------|----|--------|
| Time | 0 | 60 min |
| optical rotation | 82 | 77 |
- Assume some impurity is present calculate the optical rotation after 5 hours.
(Given $\ln 1.066 = 0.064$, $e^{0.16} = 1.17$)
- (a) 60 (b) 30 (c) 20 (d) 120
65. At 300 K the half-life of a sample of a gaseous compound initially at 1 atm is 100 sec. When the pressure is 0.5 atm the half-life is 50 sec. The order of reaction is :
- (a) 0 (b) 1 (c) 2 (d) 3



concentration of A, is equal to :

- (a) $2 \times 10^3 \text{ M s}^{-1}$ (b) $4 \times 10^3 \text{ M s}^{-1}$ (c) $8 \times 10^3 \text{ M s}^{-1}$ (d) None of these

67. For an endothermic reaction where H represents the enthalpy of reaction in kJ/mol, the minimum value for the energy of activation will be :

- (a) less than H (b) more than H (c) equal to H (d) zero

68. The activation energy of the reaction, $A + B \rightleftharpoons C + D$ is 20 kcal. What would be the activation energy of the following reaction.



- (a) 20 kcal (b) -20 kcal (c) 18 kcal (d) 58 kcal





69. When the activation energies of the forward and backward reactions are equal, then:

- (a) $U = 0$, $S = 0$ (b) $U = 0$; $G = 0$ (c) $S = 0$; $G = 0$ (d) only $U = 0$





70. For an exothermic chemical process occurring in two steps as follows



The progress of reaction can be best described by:




71. Select the correct diagram for an endothermic reaction that proceeds through two steps, with the second step is rate determining :




72. $\frac{k_{35}}{k_{34}} = 1$, this means that
 (a) rate increases with the rise in temperature
 (b) rate decreases with rise in temperature
 (c) rate does not change with rise in temperature
 (d) None of the above
73. The plot of $\ln k$ versus $1/T$ is linear with slope of:
 (a) E_a/R (b) E_a/R (c) $E_a/2.303 R$ (d) $E_a/2.303 R$
74. Rate constant for a chemical reaction taking place at 500K is expressed as $k = A \cdot e^{-1000}$. The activation energy of the reaction is:
 (a) 100 cal/mol (b) 1000 kcal/mol (c) 10^4 kcal/mol (d) 10^6 kcal/mol
75. For a complex reaction $A \xrightarrow{k} \text{products}$
 $E_{a_1} = 180 \text{ kJ/mol}$; $E_{a_2} = 80 \text{ kJ/mol}$; $E_{a_3} = 50 \text{ kJ/mol}$
 Overall rate constant k is related to individual rate constant by the equation $k = \frac{k_1 \cdot k_2}{k_3}^{2/3}$.
 Activation energy (kJ/mol) for the overall reaction is :
 (a) 100 (b) 43.44 (c) 150 (d) 140
76. For reaction $A \xrightarrow{k_1} B$, the rate constant $k_1 = A_1 (e^{-E_{a_1}/RT})$ and for the reaction $X \xrightarrow{k_2} Y$, the rate constant $k_2 = A_2 (e^{-E_{a_2}/RT})$. If $A_1 = 10^9$, $A_2 = 10^{10}$ and $E_{a_1} = 1200 \text{ cal/mol}$ and $E_{a_2} = 1800 \text{ cal/mol}$, then the temperature at which $k_1 = k_2$ is: (Given; $R = 2 \text{ cal/K-mol}$)
 (a) 300K (b) $300 \times 2.303 \text{ K}$ (c) $\frac{300}{2.303} \text{ K}$ (d) None of these
77. The activation energies of the forward and backward reactions in the case of a chemical reaction are 30.5 and 45.4 kJ/mol respectively. The reaction is :
 (a) exothermic (b) endothermic
 (c) neither exothermic nor endothermic (d) independent of temperature
78. A reaction rate constant is given by : $k = 1.2 \times 10^{14} e^{-\frac{25000}{RT}} \text{ sec}^{-1}$. It means :
 (a) $\log k$ versus $\log T$ will give a straight line with a slope as 25000
 (b) $\log k$ versus $\log T$ will give a straight line with a slope as -25000
 (c) $\log k$ versus T will give a straight line with a slope as -25000
 (d) $\log k$ versus $1/T$ will give a straight line
79. The temperature coefficient of a reaction is :
 (a) the rate constant
 (b) the rate constant at a fixed temperature
 (c) the ratio of rate constant at two temperature
 (d) the ratio of rate constant differing by 10°C preferably k_{308}/k_{298}

80. Which graph shows zero activation energy?



81. A homogeneous catalytic reaction takes place through the three alternative plots A, B and C shown in the given figure. Which one of the following indicates the relative ease with which the reaction can take place ?



- (a) A B C (b) C B A (c) A C B (d) A B C

82. A first order reaction is 50% completed in 20 minutes at 27°C and in 5 minutes at 47°C . The energy of activation of the reaction is :

- (a) 43.85 kJ/mol (b) 55.14 kJ/mol (c) 11.97 kJ/mol (d) 6.65 kJ/mol

83. The rate of a reaction gets doubled when temperature increases by 27°C to 37°C . By what factor will it change for the temperature range 17°C to 27°C .

- (a) 1.81 (b) 1.71 (c) 2.1 (d) 2.41

84. Which of the following explains the increase of the reaction rate by catalyst ?

- (a) Catalyst decreases the rate of backward reaction so that the rate of forward reaction increases
- (b) Catalyst provides extra energy to reacting molecules so that they may reduce effective collisions
- (c) Catalyst provides an alternative path of lower activation energy to the reactants
- (d) Catalyst increases the number of collisions between the reacting molecules.

85. Collision theory is satisfactory for:

- (a) First order reactions (b) Zero order reactions
- (c) Bimolecular elementary reactions (d) Any order reactions

86. For the first order reaction $A \rightarrow B + C$, carried out at 27°C . If $3.8 \times 10^{-16}\%$ of the reactant molecules exists in the activated state, the E_a (activation energy) of the reaction is :

- (a) 12 kJ/mol (b) 831.4 kJ/mol (c) 100 kJ/mol (d) 88.57 kJ/mol

87. A catalyst lowers the activation energy for a certain reaction from 83.314 to 75 kJ mol⁻¹ at 500 K. What will be the rate of reaction as compared to uncatalysed reaction? Assume other things being equal.

- (a) Double (b) 28 times (c) 7.38 times (d) 7.38×10^3 times
- 88.** A following mechanism has been proposed for a reaction
 $2A + B + D \rightarrow E$
 $A + B \rightarrow C + D$ (slow)
 $A + C \rightarrow E$ (fast)
 The rate law expression for the reaction by RDS method is:
 (a) $r = k[A]^2[B]$ (b) $r = k[A][B]$ (c) $r = k[A]^2$ (d) $r = k[A][C]$
- 89.** A hypothetical reaction $A_2 + B_2 \rightarrow 2AB$ follows the mechanism as given below
 $A_2 \rightleftharpoons A + A$ (fast)
 $A + B_2 \rightarrow AB + B$ (slow)
 $A + B \rightarrow AB$ (fast)
 The order of the over all reaction by RDS method is
 (a) 2 (b) 1 (c) $\frac{3}{2}$ (d) 0
- 90.** Chemical reaction occurs as a result of collisions between reacting molecules. Therefore, the reaction rate is given by
 (a) total number of collisions occurring in a unit volume per second
 (b) fraction of molecules which possess energy less than the threshold energy
 (c) total number of effective collisions which have enough activation energy
 (d) none of the above
- 91.** Radioactivity is affected by :
 (a) temperature (b) pressure
 (c) electric and magnetic field (d) none of these
- 92.** The radiation from naturally occurring radioactive substance as seen after deflection by a magnetic field in one direction are :
 (a) α -rays (b) β -rays
 (c) both α and β rays (d) either α or β -rays
- 93.** In the radioactive decay
 ${}_Z^AX^A \rightarrow {}_Z^AY^A + {}_Z^AZ^A + {}_Z^AZ^A$ the sequence of the radiation emitted is :
 high energy low energy
 (a) α, β, γ (b) γ, β, α
 (c) β, γ, α (d) γ, α, β
- 94.** A radioactive nuclide emits γ -rays due to the :
 (a) emission of an electron from its orbital
 (b) nuclear energy transition from a higher state to a lower state
 (c) presence of less neutrons than protons
 (d) presence of more neutrons than protons
- 95.** Consider the following decay ${}_Z^AX \rightarrow {}_Z^AY + {}_1^0e$, X is unstable because :
 (a) its nucleus has excess energy (b) $\frac{n}{p}$ ratio is high
 (c) $\frac{n}{p}$ ratio is low (d) none of these

96. Consider the following decay ${}_Z^AX \rightarrow {}_Z^AY + {}_1^0e$ () X is unstable because :
- (a) its nucleus has excess energy (b) $\frac{n}{p}$ ratio is high
(c) $\frac{n}{p}$ ratio is low (d) none of these
97. During β^- -decay :
- (a) $\frac{n}{p}$ ratio decreases (b) $\frac{n}{p}$ ratio increases
(c) $\frac{n}{p}$ ratio remains constant (d) $\frac{n}{p}$ ratio may increase or decrease
98. Which of the following processes causes the emission of X-ray?
- (a) β^- -emission (b) β^+ -emission
(c) (Positron) emission (d) electron capture
99. Which of the following processes result in an increase in the atomic number of a nuclide?
- (a) β^- -emission (b) electron capture (c) β^+ -emission (d) β^- -(Beta)emission
100. is produced when a positron and an electron collide.
- (a) X-ray (b) Neutron (c) γ -radiation (d) Neutrino
101. ${}_{67}^{165}\text{Ho}$ is stable isotope. ${}_{67}^{150}\text{Ho}$ is expected to disintegrate by:
- (a) β^- -emission (b) β^+ -emission (c) positron emission (d) β^- -emission
102. ${}_1^1\text{H}$ is a stable isotope. ${}_1^3\text{H}$ is expected to disintegrated by :
- (a) β^- -emission (b) β^+ -emission (c) positron emission (d) proton emission
103. Emission of β^- -particle is equivalent to :
- (a) increase of one proton only (b) decrease of one neutron only
(c) both (a) and (b) (d) none of these
104. Atoms ${}_7^AX$, ${}_8^BY$ and ${}_9^{17}Z$ are such that ${}_8^BY$ is an isobar of ${}_7^AX$ and atom ${}_9^{17}Z$ is isotone of ${}_8^BY$. Mass no. of X and no. of neutrons in Y are respectively :
- (a) 8, 8 (b) 17, 7 (c) 9, 8 (d) 16, 8
105. ${}_{90}^{234}\text{Th}$ disintegrates to give ${}_{82}^{206}\text{Pb}$ as the final product. Total no. of α and β particles emitted out during this process are :
- (a) 6 (b) 7 (c) 8 (d) 13
106. An isotone of ${}_{32}^{76}\text{Ge}$ is :
- (a) ${}_{32}^{77}\text{Ge}$ (b) ${}_{33}^{77}\text{As}$ (c) ${}_{34}^{77}\text{Se}$ (d) ${}_{36}^{77}\text{Se}$
107. Pair of isobar is :
- (a) ${}_{6}^{13}\text{C}$, ${}_{7}^{13}\text{N}$ (b) ${}_{6}^{13}\text{C}$, ${}_{7}^{14}\text{N}$ (c) ${}_{6}^{14}\text{C}$; ${}_{8}^{15}\text{N}$ (d) none of these
108. Isodiaphers are atoms having:
- (a) n/p same (b) p/n same (c) $(n - p)$ same (d) $(n + p)$ different
109. The 'Group displacement law' was given by:
- (a) Bacqueral (b) Rutherford (c) Madam Curie (d) Soddy and Fajan
110. ${}_3^7\text{Li} + {}_1^1\text{p} \rightarrow X$; Identify X if reaction is $(p, \quad)$ type.
- (a) ${}_4^8\text{Be}$ (b) ${}_2^4\text{He}$ (c) ${}_0^0$ (d) none of these

111. Identify reaction type:



- (a) (d, p) (b) (p, p) (c) (p, d) (d) none of these

112. ${}_{13}^{27}\text{Al} + {}_1^1\text{p} \rightarrow X + {}_0^0$; Identify X if reaction is (p,) type artificial radioactive reaction.

- (a) ${}_{13}^{28}\text{Al}$ (b) ${}_{14}^{27}\text{Si}$ (c) ${}_{14}^{28}\text{Si}$ (d) none of these

113. The number of neutrons accompanying in the formation of ${}_{54}^{139}\text{Xe}$ and ${}_{38}^{94}\text{Sr}$ from the absorption of slow neutron by ${}_{92}^{235}\text{U}$ followed by nuclear fission is :

- (a) 0 (b) 1 (c) 2 (d) 3

114. What will be the product of reaction ${}_{101}^{255}\text{Md}(\alpha, 2n)$?

- (a) ${}_{103}^{256}\text{Lr}$ (b) ${}_{102}^{257}\text{No}$ (c) ${}_{103}^{257}\text{Lr}$ (d) ${}_{82}^{205}\text{Pb}$

115. Complete the following nuclear equation by supplying the symbol for the other product of the fission :



- (a) ${}_{54}^{139}\text{Xe}$ (b) ${}_{54}^{140}\text{Xe}$ (c) ${}_{64}^{104}\text{Gd}$ (d) none of these

116. ${}_{92}^{235}\text{U} + {}_0^1n \rightarrow {}_{56}^{139}\text{Ba} + {}_{36}^{94}\text{Kr} + 3{}_0^1n + 200\text{MeV}$

Total energy released (in MeV) after 5th stage of fission is :

- (a) 48600 (b) 16200 (c) 24200 (d) None of these

117. Proton bombardment on Th^{230} followed by emission of two alpha particles will produce :

- (a) Rn^{232} (b) Ra^{233} (c) Fr^{223} (d) Fr^{222}

118. $X + {}_{82}^{206}\text{Pb} \rightarrow {}_2^4\text{He}$. In this reaction predict the position of group of X :

- (a) II B (b) IV B (c) VI A (d) VI B

119. ${}_{90}\text{Th}$ is a member of III group. After losing α -particle it forms a new element belonging to :

- (a) I group (b) II group (c) III group (d) IV group

120. Alpha decay of ${}_{92}^{238}\text{U}$ forms ${}_{90}^{234}\text{Th}$. What kind of decay from ${}_{90}^{234}\text{Th}$ produces ${}_{89}^{234}\text{Ac}$?

- (a) (b) (c) (positron) (d) β^- -emission

121. ${}_{83}^{214}\text{Bi}$ decays to A by β^- -emission. A then decays to B by beta emission, which further decays to C by another beta emission. Element C decays to D by still another beta emission, and D decays by β^- -emission to form a stable isotope E. What is element E?

- (a) ${}_{81}^{207}\text{Tl}$ (b) ${}_{80}^{206}\text{Hg}$ (c) ${}_{79}^{206}\text{Au}$ (d) ${}_{82}^{206}\text{Pb}$

122. The activity of a radioactive nuclide (X^{100}) is 6.023 curie at a certain time 't'. If its disintegration constant is $3.7 \times 10^4 \text{ s}^{-1}$ the mass of X after t sec is :

- (a) $6.022 \times 10^6 \text{ g}$ (b) 10^{-13} g (c) 10^{-15} g (d) 10^{-17} g


123. Activity of a radioactive substance is A_1 at time t_1 and A_2 at time t_2 ($t_2 > t_1$), then the ratio of $\frac{A_2}{A_1}$ is :

- (a) $e^{-(t_2 - t_1)}$ (b) $e^{(t_1 - t_2)}$ (c) $e^{-(t_1 - t_2)}$ (d) $\frac{t_2}{t_1}$

- 124.** The half-life of $^{14}_6\text{C}$ is 5730 year. What fraction of its original C^{14} would left after 22920 year of storage?
 (a) 0.50 (b) 0.25 (c) 0.125 (d) 0.0625
- 125.** A radioactive sample had an initial activity of 56 dpm. After 69.3 minutes, it was found to have an activity of 28 dpm. Find the number of atoms in a sample having an activity of 100 dpm.
 (a) 693 (b) 100 (c) 1000 (d) 10,000
- 126.** A radioactive sample has initial activity of 28 dpm 30 minutes later its activity 14 dpm. How many atoms of nuclide were present initially?
 (a) 2800 (b) 1212 (c) 528 (d) 2802
- 127.** The value of decay constant of Co^{60} is $2.5 \times 10^{-7} \text{ min}^{-1}$. The activity of 2.0 g of the sample is nearly :
 (a) $5 \times 10^5 \text{ dpm}$ (b) $2.5 \times 10^{10} \text{ dpm}$ (c) $5 \times 10^{15} \text{ dpm}$ (d) 10^{10} dpm
- 128.** Half-life ($t_{1/2}$) for a radioactive decay is 6930 sec. The time required to fall the rate of decay to $\frac{1}{100}$ of its initial value is :
 (a) 69.3 sec (b) 20,000 sec (c) 46060 sec (d) none of these
- 129.** A sample of radioactive substance is found 90% of its initial amount after one day. What % of the original sample can be found after 3 days?
 (a) 81 (b) 72.9 (c) 25 (d) 65.61
- 130.** If time t is required for a radioactive substance to become one third of its initial amount, what fraction would be left after $0.5 t$?
 (a) $\frac{1}{2}$ (b) $\frac{1}{\sqrt{3}}$ (c) $\frac{1}{3}$ (d) $\sqrt{\frac{2}{3}}$
- 131.** The present activity of the hair of Egyption mummy is 1.75 dpm. $t_{1/2}$ of $^{14}_6\text{C}$ is 5770 year and disintegration rate of fresh sample of C^{14} is 14 dpm. Find out age of mummy.
 (a) 23080 year (b) 138480 year (c) 11998.3 year (d) 17310 year
- 132.** The amount of $^{14}_6\text{C}$ isotope in a piece of wood is found to one fourth ($1/4$) of that present in a fresh piece of wood. Calculate the age of the piece of wood ($t_{1/2}$ of $^{14}_6\text{C}$ 5770 years)
 (a) 7999 year (b) 11540 year (c) 16320 year (d) 23080 year
- 133.** A radioactive element undergoing decay is left 20% of its initial weight after certain period of time t . How many such periods should elapse from the start for the 50% of the element to be left over?
 (a) 3 (b) 4 (c) 5 (d) None of these
- 134.** In a sample of wood, the reading of a counter is 32 dpm and in a fresh sample of tree it is 122dpm. Due to error counter gives the reading 2 dpm in absence of ^{14}C . Half life of ^{14}C is 5770 years.
 The approximate age (in years) of wood sample is:
 (a) 7997.2 (b) 57570 (c) 11,540 (d) 15140
- 135.** A 0.50g sample of rock was found to have $2.5 \times 10^{-6} \text{ mol}$ of $^{40}_{19}\text{K}$ ($t_{1/2}$ $1.3 \times 10^9 \text{ yr}$) and $7.5 \times 10^{-6} \text{ mol}$ of $^{40}_{20}\text{Ca}$. How old is the rock ?
 (a) $6.5 \times 10^8 \text{ yr}$ (b) $1.3 \times 10^9 \text{ yr}$ (c) $2.6 \times 10^9 \text{ yr}$ (d) $5.2 \times 10^9 \text{ yr}$

- 136.** Indium-112 is radioactive and has a very short half-life ($t_{1/2}$ 14 min). Its decay constant and average life are respectively :
- (a) 0.0495 min^{-1} , 9.7 min (b) 0.495 min^{-1} , 20.2 min
(c) 9.7 min^{-1} , 20.2 min (d) 0.0495 min^{-1} , 20.2 min
- 137.** The half-life of a radioactive element is 100 minutes. The time interval between the stages to 50% and 87.5% decay will be :
- (a) 100 min (b) 50 min (c) 200 min (d) 25 min
- 138.** The half-life of Tc^{99} is 6.0 hr. The total residual activity in a patient 30 hr after receiving an injection containing Tc^{99} must be more than $0.01 C_i$. What is the maximum activity (in C_i) that the sample injected can have?
- (a) 0.16 (b) 0.32 (c) 0.64 (d) 0.08
- 139.** A pure radio-chemical preparation was observed to disintegrate at the rate of 2140 counts/minutes at 12.35 PM. At 3.55 PM. of the same day, the disintegration rate of the sample was only 535 count/minutes. What is the half-life of the material?
- (a) 50 min (b) 100 min (c) 200 min (d) None of these
- 140.** A certain radioactive isotope ${}^A_Z X$ ($t_{1/2}$ 100 days) decays to ${}^A_Z {}^8 Y$. If 1 mole of ${}^A_Z X$ is kept in sealed container, how much He gas will accumulate at 1 atm and 273 K in 200 days?
- (a) 11.2 litre (b) 33.6 litre (c) 22.4 litre (d) 44.8 litre
- 141.** A radioactive substance decay 25% in 10 minute. If at start there are 4×10^{20} atoms present. After how much time will the number of atoms be reduced to 10^{20} atoms? (given $\ln 3 = 1.098$)
- (a) 10.98 min (b) 21.97 min (c) 48.19 min (d) None of these
- 142.** The time of decay for a nuclear reaction is given by $t = 4 t_{1/2}$. The relation between the mean life (T) and time of decay (t) is given by :
- (a) $2 T \ln 2$ (b) $4 T \ln 2$ (c) $2 T^4 \ln 2$ (d) $\frac{1}{T^2} \ln 2$
- 143.** Two radio isotopes A and B of atomic mass X and Y are mixed in equal amount by mass. After 20 days, their mass ratio is found to be 1 : 4. Isotope A has a half-life of 1 day. The half-life of isotope B is :
- (a) $1.11 \frac{Y}{X} \text{ day}$ (b) $0.11 \frac{X}{Y} \text{ day}$ (c) 0.6237 day (d) 1.10 day
- 144.** Two radioactive nuclides A and B have half-lives 50 min and 10 min respectively. A fresh sample contains the nuclide of B to be eight times that of A. How much time should elapse so that the number of nuclides of A becomes double of B?
- (a) 30 (b) 40 (c) 50 (d) 100
- 145.** A radioactive nuclide is produced at a constant rate of λ per second. It's decay constant is λ . If N_0 be the no. of nuclei at time $t = 0$, then max. no. of nuclei possible are :
- (a) N_0 (b) $\frac{1}{\lambda}$ (c) $N_0 - \frac{1}{\lambda}$ (d) $\frac{1}{\lambda} - N_0$
- 146.** An analysis of the rock shows that the relative number of Sr^{87} and Rb^{87} ($t_{1/2}$ 4.7×10^{10} year) atoms is 0.05. What is the age of the rock? Assume all the Sr^{87} have been formed from Rb^{87} only.
- (a) $7.62 \times 10^9 \text{ year}$ (b) $1.43 \times 10^9 \text{ year}$
(c) $3.28 \times 10^9 \text{ year}$ (d) $4.32 \times 10^8 \text{ year}$

- 147.** There are two radio nuclei A and B . A is a α -emitter and B is β -emitter. Their disintegration constant are in the ratio of $1 : 2$. What should be the number of atoms ratio of A and B at time $t = 0$, so that initially probability of getting of α and β -particles are same.
- (a) $2 : 1$ (b) $4 : 1$ (c) $1 : 2$ (d) $1 : 4$
- 148.** A radioactive substance (parent) decays to its daughter element. The age of radioactive substance (t) is related to the daughter (d)/parent (p) ratio by the equation :
- (a) $t = \frac{1}{\lambda} \ln \left(1 + \frac{p}{d} \right)$ (b) $t = \frac{1}{\lambda} \ln \left(1 + \frac{d}{p} \right)$
- (c) $t = \frac{1}{\lambda} \ln \frac{d}{p}$ (d) $t = \frac{1}{\lambda} \ln \frac{p}{d}$
- 149.** Ac^{227} has a half-life of 22 years. The decays follows two parallel paths



- What are the decay constants (λ) for Th and Fr respectively?
- (a) 0.03087, 0.00063 (b) 0.00063, 0.03087 (c) 0.02, 0.98 (d) None of these
- 150.** $^{218}_{84}\text{Po}$ ($t_{1/2} = 183 \text{ sec}$) decay to $^{214}_{82}\text{Pb}$ ($t_{1/2} = 161 \text{ sec}$) by α -emission, while Pb^{214} is a β -emitter. In an experiment starting with 1 mole of pure Po^{218} , how much time would be required for the number of nuclei of $^{214}_{82}\text{Pb}$ to reach maximum?
- (a) 147.5 (b) 247.5 (c) 182 (d) 304



Level 2


1. The forward rate constant for the elementary reversible gaseous reaction



What is the rate constant for the backward reaction at this temperature if 10^{-4} moles of CH_3 and 10 moles of C_2H_6 are present in a 10 litre vessel at equilibrium.

- (a) $1.57 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$ (b) $1.57 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$
(c) $1.57 \times 10^{11} \text{ L mol}^{-1} \text{ s}^{-1}$ (d) $1.57 \times 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$

2. For a hypothetical reaction,



These reactions are carried simultaneously in a reactor such that temperature is not changing. If rate of disappearance of B is $y \text{ M sec}^{-1}$ then rate of formation (in M sec^{-1}) of Q is:

- (a) $\frac{2}{3}y$ (b) $\frac{3}{2}y$ (c) $\frac{4}{3}y$ (d) $\frac{3}{4}y$

3. The kinetic data for the given reaction $A(g) + 2B(g) \xrightarrow{k} C(g)$ is provided in the following table for three experiments at 300 K

Ex. No.	[A/M]	[B/M]	Initial rate (M sec^{-1})
1.	0.01	0.01	6.930×10^{-6}
2.	0.02	0.01	1.386×10^{-5}
3.	0.02	0.02	1.386×10^{-5}

In another experiment starting with initial concentration of 0.5 and 1 M respectively for A and B at 300 K, find the rate of reaction after 50 minutes from the starting of experiment (in M/sec).

- (a) 6.93×10^{-4} (b) 0.25×10^{-7} (c) 4.33×10^{-5} (d) 3.46×10^{-4}
4. The reaction $A(g) + 2B(g) \rightarrow C(g)$ is an elementary reaction. In an experiment involving this reaction, the initial partial pressures of A and B are $P_A = 0.40 \text{ atm}$ and $P_B = 1.0 \text{ atm}$ respectively. When $P_C = 0.3 \text{ atm}$, the rate of the reaction relative to the initial rate is:
- (a) $\frac{1}{12}$ (b) $\frac{1}{50}$ (c) $\frac{1}{25}$ (d) none of these
5. Which of the following is incorrect statement?
- (a) Stoichiometry of a reaction tells about the order of the elementary reaction
(b) For a zero order reaction, rate and the rate constant are identical.
(c) A zero order reaction is controlled by factors other than concentration of reactants
(d) A zero order reaction is an elementary reaction

6. Two first order reaction have half-lives in the ratio 8 : 1. Calculate the ratio of time intervals $t_1 : t_2$. The time t_1 and t_2 are the time period for $\frac{1}{4}$ th and $\frac{3}{4}$ th completion.


(a) 1 : 0.301 (b) 0.125 : 0.602 (c) 1 : 0.602 (d) none of these

7. Reaction $A + B \rightarrow C + D$ follows rate law, $r = k[A]^{1/2}[B]^{1/2}$. Starting with 1 M of A and B each, what is the time taken for concentration of A become 0.1 M?

[Given $k = 2.303 \times 10^{-2} \text{ sec}^{-1}$].

(a) 10 sec (b) 100 sec (c) 1000 sec (d) 434 sec

8. For the two reactions I: $A \rightarrow B$; II: $C \rightarrow D$ following graph is obtained.



Which of the following is true:

(a) If $[B] = [A]$ then at that time $[D] = 0.75 M$ (b) If $[C] = [A]$ then at that time $[B] = [D]$

(c) $(t_{100\%})_{\text{Reaction I}} = (t_{100\%})_{\text{Reaction II}}$ (d) $[A] = [C]$ at $t = \frac{\sqrt{3}}{2} \text{ min.}$

9. The reaction $A(g) + B(g) \rightarrow 2C(g)$ is a first order reaction with rate constant $2.772 \times 10^{-3} \text{ s}^{-1}$. Starting with 0.1 mole of A in 2 litre vessel, find the concentration of A after 250 sec when the reaction is allowed to take place at constant pressure and at 300 K ?

(a) 0.0125 M (b) 0.025 M (c) 0.05 M (d) none of these

10. For a first order homogeneous gaseous reaction, $A + 2B \rightarrow C$ if the total pressure after time t was P_t and after long time ($t \rightarrow \infty$) was P then k in terms of P_t, P and t is :

(a) $k = \frac{2.303}{t} \log \frac{P}{P - P_t}$

(b) $k = \frac{2.303}{t} \log \frac{2P}{P - P_t}$

(c) $k = \frac{2.303}{t} \log \frac{2P}{3(P - P_t)}$

(d) none of these

11. $A(aq) + B(aq) \rightarrow C(aq)$ is a first order reaction.

Time t
moles of reagent $n_1 \quad n_2$

Reaction progress is measured with the help of titration of reagent 'R'. If all A, B and C react with reagent and have 'n' factors [n factor; eq. mass $\frac{\text{mol. mass}}{n}$] in the ratio of 1 : 2 : 3 with the reagent, the k in terms of t, n_1 and n_2 is :


(a) $k = \frac{1}{t} \ln \frac{n_2}{n_2 - n_1}$

(b) $k = \frac{1}{t} \ln \frac{2n_2}{n_2 - n_1}$


(c) $k \frac{1}{t} \ln \frac{4n_2}{n_2 - n_1}$

(d) $k \frac{1}{t} \ln \frac{4n_2}{5(n_2 - n_1)}$

12. The gaseous decomposition reaction, $A(g) \rightarrow 2B(g) + C(g)$ is observed to first order over the excess of liquid water at 25°C . It is found that after 10 minutes the total pressure of system is 188 torr and after very long time it is 388 torr. The rate constant of the reaction (in hr^{-1}) is : [Given : vapour pressure of H_2O at 25° is 28 torr ($\ln 2 = 0.7, \ln 3 = 1.1, \ln 10 = 2.3$)]
- (a) 0.02 (b) 1.2 (c) 0.2 (d) none of these
13. The reaction, Sucrose $\xrightarrow{H}$ Glucose + Fructose, takes place at certain temperature while the volume of solution is maintained at 1 litre. At time zero the initial rotation of the mixture is 34° . After 30 minutes the total rotation of solution is 19° and after a very long time, the total rotation is -11° . Find the time when solution was optically inactive.
- (a) 135 min (b) 103.7 min (c) 38.7 min (d) 45 min
14. A gaseous compound A reacts by three independent first order processes (as shown in figure) with rate constant 2×10^3 , 3×10^3 and $1.93 \times 10^3 \text{ sec}^{-1}$ for products B, C and D respectively. If initially pure A was taken in a closed container with $P = 8 \text{ atm}$, then the partial pressure of B (in atm) after 100 sec from starting the experiment.




- (a) 0.288 (b) 0.577 (c) 1.154 (d) none of these
15. A compound A dissociates by two parallel first order paths at certain temperature
- | | |
|---|---|
| $A(g) \xrightarrow{k_1(\text{min}^{-1})} 2B(g)$ | $k_1 = 6.93 \times 10^3 \text{ min}^{-1}$ |
| $A(g) \xrightarrow{k_2(\text{min}^{-1})} C(g)$ | $k_2 = 6.93 \times 10^3 \text{ min}^{-1}$ |
- The reaction is started with 1 mole of pure 'A' in 1 litre closed container with initial pressure 2 atm. What is the pressure (in atm) developed in container after 50 minutes from start of experiment?
- (a) 1.25 (b) 0.75 (c) 1.50 (d) 2.50
16. For given hypothetical elementary parallel reaction,



Initially only 2 moles of A are present. The total no. of moles of A, B and C at the end of 75% reaction are:

- (a) 2 (b) 3 (c) 4 (d) 3.5


17. The reaction $\text{cis-X} \xrightleftharpoons[k_b]{k_f} \text{trans-X}$ is first order in both directions. At 25°C, the equilibrium constant is 0.10 and the rate constant $k_f = 3 \times 10^{-4} \text{ s}^{-1}$. In an experiment starting with the pure *cis*-form, how long would it take for half of the equilibrium amount of the *trans*-isomer to be formed?
- (a) 150 sec (b) 200 sec (c) 240 sec (d) 210 sec
18. A hydrogenation reaction is carried out at 500 K. If the same reaction is carried out in the presence of a catalyst at the same rate with same frequency factor, the temperature required is 400 K. What is the activation energy of the reaction, if the catalyst lowers the activation energy barrier by 16 kJ/mol?
- (a) 100 kJ/mol (b) 80 kJ/mol (c) 60 kJ/mol (d) None of these
19. Consider the reaction.



- The rate constant for two parallel reactions were found to be $10^2 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ and $4 \times 10^2 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$. If the corresponding energies of activation of the parallel reaction are 100 and 120 kJ/mol respectively, what is the net energy of activation (E_a) of A?
- (a) 100 kJ/mol (b) 120 kJ/mol (c) 116 kJ/mol (d) 220 kJ/mol
20. A reaction takes place in various steps. The rate constant for first, second, third and fifth steps are k_1, k_2, k_3 and k_5 respectively. The overall rate constant is given by

$$k = \frac{k_2}{k_3} \left(\frac{k_1}{k_5} \right)^{1/2}$$

- If activation energy are 40, 60, 50 and 10 kJ/mol respectively, the overall energy of activation (kJ/mol) is :
- (a) 10 (b) 20 (c) 25 (d) none of these
21. For reaction $A \rightarrow B$, the rate constant $k_1 = A_1 e^{E_{a1}/(RT)}$ and for the reaction $X \rightarrow Y$, the rate constant $k_2 = A_2 e^{E_{a2}/(RT)}$. If $A_1 = 10^8$, $A_2 = 10^{10}$ and $E_{a1} = 600 \text{ cal/mol}$, $E_{a2} = 1800 \text{ cal/mol}$, then the temperature at which $k_1 = k_2$ is (Given : $R = 2 \text{ cal/K-mol}$)
- (a) 1200 K (b) 1200 + 4.606 K (c) $\frac{1200}{4.606} \text{ K}$ (d) $\frac{600}{4.606} \text{ K}$
22. For first order parallel reactions k_1 and k_2 are 4 and 2 min^{-1} respectively at 300 K. If the activation energies for the formation of B and C are respectively 30,000 and 38,314 joule/mol respectively, the temperature at which B and C will be obtained in equimolar ratio is :



(a) 757.48 K

(b) 378.74 K

(c) 600 K

(d) none of these

23. In the series reaction

 $A \xrightarrow{k_1} B \xrightarrow{k_2} C \xrightarrow{k_3} D$, if $k_1 < k_2 < k_3$ then the rate determining step of the reaction is






 (a) A $\rightarrow$ B

 (b) C $\rightarrow$ D

 (c) B $\rightarrow$ C

(d) Any step

24. The following mechanism has been proposed for the exothermic catalyzed complex reaction


 If k_1 is much smaller than k_2 , the most suitable qualitative plot of potential energy (P.E.) versus reaction co-ordinate (R.C.) for the above reaction


$$\begin{array}{c}
 \text{O} \\
 \parallel \\
 \text{CH}_3\text{COH} \quad \text{(I)}
 \end{array}
 \xrightleftharpoons[k_1]{\begin{array}{c} \text{H (fast)} \\ k_1 \end{array}}
 \begin{array}{c}
 \text{O} \quad \text{H} \\
 \parallel \quad | \\
 \text{CH}_3\text{CO} \quad \text{H} \quad \text{(II)}
 \end{array}
 \xrightleftharpoons[\text{C}_2\text{H}_5\text{OH (fast)} k_2]{\begin{array}{c} + \text{C}_2\text{H}_5\text{OH (slow)} k_2 \\ \text{H O C}_2\text{H}_5 \end{array}}
 \begin{array}{c}
 \text{O} \quad \text{H} \\
 \parallel \quad | \\
 \text{CH}_3\text{C} \quad \text{O} \quad \text{H} \\
 | \quad | \\
 \text{H} \quad \text{O} \quad \text{C}_2\text{H}_5 \quad \text{(III)}
 \end{array}
 \xrightleftharpoons[\text{fast } k_3]{\text{fast } k_3}
 \begin{array}{c}
 \text{H} \quad \text{O} \quad \text{H} \\
 | \quad | \quad | \\
 \text{CH}_3\text{C} \quad \text{OH} \\
 | \quad | \\
 \text{O} \quad \text{C}_2\text{H}_5 \quad \text{(IV)}
 \end{array}
 \xrightleftharpoons[\text{H}_2\text{O; slow}]{\begin{array}{c} k_4 \\ \text{H}_2\text{O; slow} \end{array}}
 \begin{array}{c}
 \text{H} \quad \text{O} \quad \text{H} \\
 | \quad | \quad | \\
 \text{CH}_3\text{C} \quad \text{OH} \\
 | \quad | \\
 \text{O} \quad \text{C}_2\text{H}_5 \quad \text{(IV)}
 \end{array}
 \xrightleftharpoons[\text{H}^+; \text{fast}]{\begin{array}{c} k_5 \\ \text{H}^+; \text{fast} \end{array}}
 \begin{array}{c}
 \text{CH}_3 \\
 | \\
 \text{C} \begin{array}{l} \parallel \text{O} \\ \diagdown \text{OC}_2\text{H}_5 \end{array} \quad \text{(VI)}
 \end{array}
 \xrightleftharpoons[\text{H}^+; \text{fast}]{\begin{array}{c} k_5 \\ \text{H}^+; \text{fast} \end{array}}
 \begin{array}{c}
 \text{CH}_3 \\
 | \\
 \text{C} \begin{array}{l} \parallel \text{O} \\ \diagdown \text{OC}_2\text{H}_5 \end{array} \quad \text{(VI)}
 \end{array}
 \xrightleftharpoons[\text{H}_2\text{O; slow}]{\begin{array}{c} k_4 \\ \text{H}_2\text{O; slow} \end{array}}
 \begin{array}{c}
 \text{H} \quad \text{O} \quad \text{H} \\
 | \quad | \quad | \\
 \text{CH}_3\text{C} \quad \text{OH} \\
 | \quad | \\
 \text{O} \quad \text{C}_2\text{H}_5 \quad \text{(IV)}
 \end{array}
 \xrightleftharpoons[\text{H}^+; \text{fast}]{\begin{array}{c} k_5 \\ \text{H}^+; \text{fast} \end{array}}
 \begin{array}{c}
 \text{CH}_3 \\
 | \\
 \text{C} \begin{array}{l} \parallel \text{O} \\ \diagdown \text{OC}_2\text{H}_5 \end{array} \quad \text{(VI)}
 \end{array}$$

Figure 1 consists of four subplots, (a) through (d), each showing a graph of Potential Energy (P.E.) on the vertical axis versus Reaction Coordinate (R.C.) on the horizontal axis. The graphs illustrate the energy profile of a chemical reaction, with peaks labeled I through VI. Subplot (a) is the reference profile. Subplot (b) shows a 10% increase in the heights of peaks I, II, III, IV, and V compared to (a). Subplot (c) shows a 10% decrease in the heights of peaks I, II, III, IV, and V compared to (a). Subplot (d) shows a 20% increase in the heights of peaks I, II, III, IV, and V compared to (a). The minima (I, II, V, VI) are at the same energy levels in all four plots.

- For More Material Join: @JEEAdvanced_2024

28. A radioactive isotope X with half-life of 6.93×10^9 years decay to Y which is stable. A sample of rock from the moon was found to contain both the elements X and Y in the mole ratio 1 : 7. What is the age of the rock?
- (a) 2.079×10^{10} years (b) 1.94×10^{10} years
(c) 1.33×10^9 years (d) 10^{10} years
29. The average (mean) life of a radio nuclide which decays by parallel path is
- $A \xrightarrow{\lambda_1} B; \quad \lambda_1 = 1.8 \times 10^{-2} \text{ sec}^{-1}$
 $2A \xrightarrow{\lambda_2} B; \quad \lambda_2 = 10^{-3} \text{ sec}^{-1}$
- (a) 52.63 sec (b) 500 sec (c) 50 sec (d) none of these
30. The radioactive decay $^{211}_{83}\text{Bi} \rightarrow ^{207}_{81}\text{Tl}$, takes place in 100 L closed vessel at 27°C . Starting with 2 moles of $^{211}_{83}\text{Bi}$ ($t_{1/2} = 130 \text{ sec}$), the pressure development in the vessel after 520 sec will be :
- (a) 1.875 atm (b) 0.2155 atm (c) 0.4618 atm (d) 4.618 atm
31. A fresh radioactive mixture contains short lived species A and B . Both emitting α -particles initially of 8000 α -particles per minute. 20 minutes later, they emits at the rate of 3500 α -particles per minute. If the half-lives of the species A and B are 10 minutes and 500 hours respectively, then the ratio of activities of $A : B$ in the initial mixture was :
- (a) 4 : 6 (b) 6 : 4 (c) 3 : 4 (d) 3 : 1
32. In order to determine the volume of blood in an animal, a 1.0 mL sample of solution of 10^3 dpm of ^3_1H is injected into the animal blood stream. After sufficient time for circulatory equilibrium to be established, 2 mL of blood is found to have activity to 10 dpm. The volume of blood in animal is :
- (a) 199 mL (b) 198 mL (c) 200 mL (d) 20 mL
33. The ratio of activities of two radio nuclides X and Y in a mixture at time $t = 0$ was found to be 4 : 1. After two hours, the ratio of activities become 1 : 1. If the $t_{1/2}$ of radio nuclide X is 20 min then $t_{1/2}$ [in minutes] of radio nuclide Y is :
- (a) 10 (b) 20 (c) 30 (d) 40
34. Find the age of an ancient Egyptian wooden article (in years) from the given information.
- (i) Activity of 1 g of carbon obtained from ancient wooden article = 7 counts/min/g
(ii) Activity of 1 g carbon obtained from fresh wooden sample = 15.4 counts per min/g
(iii) Percentage increase in level of C^{14} due to nuclear explosions in past 100 years is 10%
(iv) $t_{1/2}$ of $^{14}_6\text{C}$ = 5770 years
- (a) 5.770×10^3 (b) 16.87×10^3 (c) 2488 (d) none of these
35. The isotopes ^{238}U and ^{235}U occur in nature in the mass ratio 140 : 1. It is assumed that initially they were found in equal mass. If half life ($t_{1/2}$) of ^{238}U = 4.5×10^9 and $t_{1/2}$ of ^{235}U = 5×10^8 year respectively then the age of earth is ($\log 7 = 0.846$; $\log 2 = 0.3$)
- (a) 4.02×10^9 year (b) 2.01×10^9 year (c) 8.72×10^9 year (d) None of these




Level 3



Passage 1

Two consecutive irreversible first order reactions can be represented by







The rate equation for A is readily integrated to obtain

$$[A]_t = [A]_0 \cdot e^{-k_1 t}; \text{ and } [B]_t = \frac{k_1 [A]_0}{k_2 - k_1} [e^{-k_1 t} - e^{-k_2 t}]$$

1. At what time will B be present in maximum concentration?

- (a) $\frac{k_1}{k_2 - k_1}$ (b) $\frac{1}{k_1 - k_2} \ln \frac{k_1}{k_2}$ (c) $\frac{1}{k_2 - k_1} \ln \frac{k_1}{k_2}$ (d) None of these

2. When $k_1 = 1 \text{ s}^{-1}$ and $k_2 = 500 \text{ s}^{-1}$; select most appropriate graph



3. Select the correct statement for given reaction :

- (a) A decreases linearly
 (b) B rises to a max. and then constant
 (c) B rises to a max. and then falls
 (d) The slowest rate of increases of C occurring where B is max.



Passage 2

Arrhenius studied the effect of temperature on the rate of a reaction and postulated that rate constant varies with temperature exponentially as $k = Ae^{E_a/RT}$. This method is generally used for finding the activation energy of a reaction. Keeping temperature constant, the effect of catalyst on the activation energy has also been studied.

- The pre-exponential factor in the Arrhenius equation of a first order reaction has the unit :
 (a) $\text{mol L}^{-1} \text{s}^{-1}$ (b) $\text{L mol}^{-1} \text{s}^{-1}$ (c) s^{-1} (d) dimensionless
- If x is the fraction of molecules having energy greater than E_a it will be given by :
 (a) $x = \frac{E_a}{RT}$ (b) $\ln x = \frac{E_a}{RT}$ (c) $x = e^{E_a/RT}$ (d) Any of these
- If the rate of reaction doubles for 10°C rise of temperature from 290K to 300K , the activation energy of the reaction will be approximately:
 (a) 40 kcal mol^{-1} (b) 12 kcal mol^{-1} (c) 60 kcal mol^{-1} (d) 70 kcal mol^{-1}




Passage 3

An important parameter of a photochemical reaction is the quantum efficiency or quantum yield (ϕ) which is defined as

$$\frac{\text{moles of the substance reacted}}{\text{moles of photons absorbed}}$$

Absorption of UV radiation decompose acetone according to the reaction



- The quantum yield of a reaction at 330 nm is 0.4 . A sample of acetone absorbs monochromatic radiation at 330 nm at the rate of $7.2 \times 10^3 \text{ Js}^{-1}$ (Given : $N_A = 6 \times 10^{23}$; $h = 6.6 \times 10^{-34}$ in S.I. unit). The rate of formation of $\text{CO}(\text{mol/s})$ is :
 (a) 2×10^8 (b) 8×10^8 (c) 8×10^9 (d) none of these
- If quantum yield is 0.8 then rate of formation of C_2H_6 (mol/s) is :
 (a) 2×10^8 (b) 1.6×10^9 (c) 16×10^9 (d) 8×10^9



Passage 4

Radioactive disintegration is a first order reaction and its rate depends only upon the nature of nucleus and does not depend upon external factors like temperature and pressure. The rate of radioactive disintegration (Activity) is represented as

$$\frac{dN}{dt} = -\lambda N$$

Where λ = decay constant; N = number of nuclei at time t ; N_0 = initial no. of nuclei

The above equation after integration can be represented as

$$\frac{2.303}{t} \log \frac{N_0}{N}$$

- Half-life period of U^{237} is 2.5×10^5 years. In how much time will the amount of U^{237} remaining be only 25% of the original amount?
(a) 2.5×10^5 years (b) 1.25×10^5 years (c) 5×10^5 years (d) 10^6 years
- Calculate the half-life period of a radioactive element which remains only 1/16 of its original amount in 4740 years :
(a) 1185 years (b) 2370 years (c) 52.5 years (d) none of these
- What is the activity in Ci (curie) of 1.0 mole of Plutonium-239? ($t_{1/2} = 24,000$ years)
(a) 1.49 Ci (b) 14.9 Ci (c) 5.513×10^{11} Ci (d) None of these



Passage 5

Size of nucleus was obtained by the equation $r = R_0 A^{1/3}$, where r is the radius of nucleus of mass no. A and R_0 is a constant whose value is equal to 1.5×10^{-15} metre.

(Given : 1 amu = 1.66×10^{-24} g)

- What is the density of a nucleus of mass number A ?
(a) $\frac{4}{3} (1.5 \times 10^{-15})^3 A$ (b) $1.17 \times 10^{17} \text{ kg/cm}^3$ (c) $1.17 \times 10^{17} \text{ kg/m}^3$ (d) None of these
- Nucleus radius of ${}^{12}_6\text{C}$ is 3×10^{-15} metre. What is density ratio of d_C/d_{H_2O} ?
(a) 1.76×10^{17} (b) 1.76×10^{14} (c) 17.6×10^7 (d) 17.6×10^{17}

One or More Answers is/are Correct



- Select the correct statement(s):
(a) Rate constants are never negative
(b) Partial orders are never negative
(c) Molecularity and order of reaction both are equal for elementary reactions
(d) Order of reaction may be change with change in practical conditions (temp. and pressure)
- Select the correct statement(s) :
(a) The rate of reaction decreases with decrease in temperature
(b) The rate of reaction is uniform in zero order reaction
(c) The rate of reaction depends upon the surface area of the solid reactants
(d) Average and instantaneous rate of reaction defined for micro and macro-scopic time interval respectively

3. Select the correct statement(s) :

- (a) The rate law of the elementary reaction; $2A + B \rightarrow C$, must be $r = k[A]^2$
- (b) The rate law for the complex reaction $A + B \rightarrow C$, might not be $r = k[A][B]$
- (c) If the partial orders differ from the stoichiometric coefficients in the balanced reaction, the reaction must be complex
- (d) If the partial orders are equal to corresponding coefficients in the balanced reaction, the reaction must be elementary

4. Select the correct statement(s):

- (a) Every substance that appears in the rate law of reaction must be a reactant or product in that reaction
- (b) If we know the rate law of a reaction; we can deduce its mechanism
- (c) If the reaction has rate $r = k[A][B]^{3/2}$ then reaction may be elementary
- (d) A zero order reaction must be a complex reaction

5. Select the correct statement(s) :

- (a) When T or $E_a \rightarrow 0$ then $k \rightarrow \infty$
- (b) A positive catalyst can change ΔH of the reaction
- (c) A mixture of reactants may be thermodynamically unstable but kinetically stable
- (d) A negative catalyst increases the activation energy of the reaction

6. Consider a reaction $A + B \rightarrow C$, in which both reactants are in the same phase, may be

- (a) unimolecular elementary reaction
- (b) Exothermic
- (c) Heterogeneous
- (d) Photochemical

7. In the following gaseous phase first order reaction



initial pressure was found to be 400 mm of Hg and it changed to 1000 mm of Hg after 20 min. Then


- (a) half life for A is 10 min
- (b) rate constant is 0.0693 min^{-1}
- (c) partial pressure of C at 30 min is 350 mm of Hg
- (d) total pressure after 30 min is 1100 mm of Hg

8. Identify the true statement(s)

- (a) A catalyst is chemically unchanged at the end of a reaction
- (b) A catalyst may appear in the kinetic rate equation of the reaction
- (c) A catalyst will not affect the composition of an equilibrium mixture
- (d) A catalyst cannot cause a non-spontaneous ($\Delta G > 0$) reaction to proceed

9. For the reaction $A \xrightleftharpoons[k_2 \text{ sec}^{-1}]{k_1 \text{ sec}^{-1}} B$ following graph is given,

$k_1 = 4 \times 10^{-2} \text{ sec}^{-1}$. Which is/are correct statement (s) ($\ln 2 = 0.7$, $\ln 8/7 = 0.14$)



- (a) Equilibrium constant is 4.0
 (b) Time taken for the completion of 50% of equilibrium conc. of B is 14 sec.
 (c) Time taken for the completion of 10% of initial conc. of A is 2.8 sec.
 (d) Rate constant of backward reaction is 10^{-2} sec^{-1}
- 10.** Select the correct statement(s) :
- (a) α -particles are simply helium atoms
 (b) γ -rays travel with higher speed as compared to α -particle and have higher ionization power as compared to β -particle
 (c) Loss of α -particles results in the production of isobars
 (d) α -particles are considered as the best bombarding particles
- 11.** Select the correct statement(s) :
- (a) In the reaction ${}_{92}^{235}\text{U} + {}_0^1n \rightarrow {}_{56}^{140}\text{Ba} + 2 {}_0^1n + x$, x is ${}_{36}^{94}\text{Kr}$
 (b) In the reaction ${}_{11}^{23}\text{Na} + z \rightarrow {}_{12}^{23}\text{Mg} + {}_0^1n$, the bombarding particle z is deuteron
 (c) Very large amount of energy is produced during nuclear fission and nuclear fusion
 (d) In a fission reaction, a loss in mass occurs releasing a vast amount of energy
- 12.** Select the correct statement(s):
- (a) SI unit of radioactivity is becquerel (Bq)
 (b) $1 \text{ Ci} = 3.7 \times 10^7 \text{ Bq}$
 (c) ${}^7_3\text{Li} + {}^1_1\text{H} \rightarrow {}^4_2\text{He} + \dots$ is (P , β) type reaction
 (d) The half-life of a particular radioactive isotope is a characteristic constant of that isotope
- 13.** Select the correct statement(s) :
- (a) On bombarding ${}^{14}_7\text{N}$ nuclei with α -particle, the nuclei of the product formed after release of proton would be ${}^{17}_8\text{O}$
 (b) Decay constant does not depend upon temperature
 (c) Nuclide and its decay product after α -emission are called isodiaphers
 (d) Half-life of radium is 1580 years. Its average life will be 1097.22 years
- 14.** In the decay process
- | | | | |
|---|---|---|---|
| A | B | C | D |
|---|---|---|---|
- (a) A and B are isobars
 (b) A and D are isotopes

- (c) B , C and D are isobars
(d) A and C are isotones
- 15.** In electron capture (radioactive process) :
(a) a neutron is formed (b) a proton is consumed
(c) γ -ray emission takes place (d) X-ray emission takes place
- 16.** Select the correct statement (s) for positron emission by unstable nucleus.
(a) X-ray emission takes place (b) A neutron is formed
(c) $\frac{n}{p}$ of daughter nucleus increases (d) A neutron is consumed
- 17.** Select the correct statement (s) :
(a) Mass number remains constant when positron emission takes place
(b) One neutron converts into proton in $(\text{}^0_1e)$ emission process
(c) Activity of a radioactive substance double when temp. increases from 300 K to 310 K
(d) Isodiaphers formed when one alpha particle emitted and isotopes formed when 2 beta particles emitted

Match the Column



Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.	Column-I (A) Unit of k is always equals to (B) Unit of k in zero order (C) Unit of k in first order (D) Unit of k in second order	Column-II (P) 1/time (Q) M/time (R) $\text{Time}^{-1} M^{-1}$ (S) Unit of A (pre-exponential factor)
2.	Column-I (A) Molecularity of a reaction (B) Order of reaction (C) The dissociation of H_2O_2 (aq) is (D) $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \xrightarrow{h\nu} 2\text{HCl}$ is	Column-II (P) 0, 1 Possible (Q) 1, 2 Possible (R) First order reaction (S) Zero order reaction
3.	Column-I (Curve) (A) C vs t (abscissa) for zero order (B) $\log C$ vs t (abscissa) for first order (C) $\frac{dc}{dt}$ vs C for zero order (D) $\ln \frac{dc}{dt}$ vs $\ln C$ for first order	Column-II (Slope) (P) unity (Q) zero (R) k (S) $\frac{k}{2.303}$
4.	Column-I Linear plots (with non zero slope) (A) $\ln \frac{d[A]}{dt}$ vs. $\ln [A]$ (B) $\log_e k$ vs. $\frac{1}{T}$ (C) $\log t_{1/2}$ vs. $\log [A]_0$ (D) $\frac{d[A]}{dt}$ vs. $[A]^2$	Column-II (Order) (P) 2 (Q) $\frac{1}{2}$ (R) 0 (S) 1





$[A]$ = Reactant concentration at time ' t '

k rate constant

T Absolute temperature

5.

Column-I
(First order of reaction A B)

(A) $[B]$ vs t (B) $[A]$ vs t (C) $\frac{1}{[A]}$ vs t (D) $\ln k$ vs $\frac{1}{T}$ **Column-II**

6.

Column-I

(A) Isotones

(B) Isobars

(C) Isotopes

(D) Isodiaphers

Column-II(P) ${}^{234}_{91}\text{Pa}$ and ${}^{234}_{90}\text{Th}$ (Q) ${}^{12}_6\text{C}$ and ${}^{14}_6\text{C}$ (R) ${}^{39}_{19}\text{K}$ and ${}^{19}_9\text{F}$ (S) ${}^{39}_{18}\text{Ar}$ and ${}^{40}_{19}\text{K}$

7.

Column-I(A) α -emission(B) β^- -emission(C) β^+ -emission

(D) (Positron) emission

Column-II

(P) Change in mass no.

(Q) No change in atomic no. and mass no.

(R) Atomic no. decreases

(S) Atomic no. increases

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** Molecularity has no meaning for a complex reaction
STATEMENT-2 : Molecularity defined only for RDS
2. **STATEMENT-1 :** An elementary reaction cannot have fractional order.
STATEMENT-2 : Stoichiometric coefficients in an elementary reaction can be fractional.
3. **STATEMENT-1 :** Concentration of reactant in zero order reaction is constant.
STATEMENT-2 : For zero order reaction $A \xrightarrow{k} B$, successive half life of reaction decreases with the progress of the reaction.
4. **STATEMENT-1 :** Acid catalysed hydrolysis of esters is pseudo first order reaction.
STATEMENT-2 : Water is present in excess in given reaction.
5. **STATEMENT-1 :** The order of reaction can have fractional value.
STATEMENT-2 : For an elementary reaction, the partial orders are determined by the reaction stoichiometry.
6. **STATEMENT-1 :** Catalyst may increase the rate constant to a large extent.
STATEMENT-2 : By using suitable catalyst, we can significantly increase yield.
7. **STATEMENT-1 :** For each 10°C rise of temperature the k is nearly double.
STATEMENT-2 : Energy wise distribution of molecules in a gas sample is an exponential function of temperature so $e^{E_a/RT}$ is doubled.
8. **STATEMENT-1 :** Product is formed only when the required orientation and energy conditions are met.
STATEMENT-2 : All collisions between reactants yield the desired product.
9. **STATEMENT-1 :** The plot of k versus $1/T$ is linear.
STATEMENT-2 : $k = A e^{E_a/(RT)}$
10. **STATEMENT-1 :** For exothermic reaction equilibrium constant decreases with increase in temperature.
STATEMENT-2 : For exothermic reaction rate constant decreases with decrease in temperature.
11. **STATEMENT-1 :** If the activation energy of reaction is zero, temperature will have no effect on the rate constant.
STATEMENT-2 : Lower the activation energy faster is the reaction.
12. **STATEMENT-1 :** Active complex is an intermediate product.
STATEMENT-2 : Active complex is unstable with high vibrational energy.

- 13. STATEMENT-1 :** The pre-exponential factor A has the same units for all reactions.
STATEMENT-2 : $e^{E_a/RT}$ has no unit.
- 14. STATEMENT-1 :** γ -rays have very high penetrating power.
STATEMENT-2 : γ -rays are electromagnetic radiations of high energy.
- 15. STATEMENT-1 :** Nuclide ${}^{30}_{13}\text{Al}$ is less stable than ${}^{40}_{20}\text{Ca}$.
STATEMENT-2 : Nuclide having odd number of protons and neutrons are generally unstable.
- 16. STATEMENT-1 :** Disintegration of ${}^3_1\text{H}$ (tritium) is accompanied by β^- -emission.
STATEMENT-2 : Tritium has high n/p ratio.
- 17. STATEMENT-1 :** The life of radioactive object (organic origin) can found with the help of carbon dating.
STATEMENT-2 : ${}^{14}_6\text{C}$ is a β^- and γ -emitter.
- 18. STATEMENT-1 :** Neutrons are the best bombarding particles.
STATEMENT-2 : Neutrons are neutral particles.
- 19. STATEMENT-1 :** Nucleus does not contain free electrons, yet it emit beta-particles
STATEMENT-2 : At high n/p ratio, one neutron supposed to give 1 proton and 1 e^- (β^-).
- 20. STATEMENT-1 :** Rate of disintegration of thorium increases with the increase in moles of thorium.
STATEMENT-2 : Rate of disintegration does not depend upon temperature, pressure

Subjective Problems



- The rate of decomposition of $\text{NH}_3(\text{g})$ at 10 atm on platinum surface is zero order. What is rate of formation (in $M \text{ min}^{-1}$) of $\text{H}_2(\text{g})$, if rate constant of reaction $2\text{NH}_{3(\text{g})} \rightarrow \text{N}_{2(\text{g})} + 3\text{H}_{2(\text{g})}$ is $2.0 M \text{ min}^{-1}$?
- 5A $\rightarrow$ Product
 In above reaction, half-life period is directly proportional to initial concentration of reactant. The initial rate of reaction is $400 \text{ mol lit}^{-1} \text{ min}^{-1}$.
 Calculate the half-life period (in sec) when initial concentration of reactant is 200 mol lit^{-1} .
- In an elementary reaction $\text{A}(\text{g}) + 2\text{B}(\text{g}) \rightarrow \text{C}(\text{g})$ the initial pressure of A and B are $P_A = 0.40$ atm and $P_B = 0.60$ atm respectively. After time T , if pressure of C is observed 0.1 atm, then find the value of $\frac{r_i \text{ (initial rate of reaction)}}{r_t \text{ (rate of reaction after time } t)}$.
- Carbon monoxide reacts with O_2 to form CO_2 : $2\text{CO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g})$
 Informations about this reaction are given in the table below.

[CO]mol/L	[O ₂]mol/L	Rate of reaction (mol/L.min)
0.02	0.02	4×10^{-5}
0.04	0.02	1.6×10^{-4}
0.02	0.04	8×10^{-5}

What is the value for the rate constant for the reaction in properly related unit ?

5. Half-life for the zero order reaction $A(g) \rightarrow B(g) + C(g)$ and half-life for the first order reaction $X(g) \rightarrow Y(g) + Z(g)$ are equal. If completion time for the zero order reaction is 13.86 min, then calculate the rate constant (in hr^{-1}) for the reaction $X(g) \rightarrow Y(g) + Z(g)$.

6. For any acid catalysed reaction, $A \xrightarrow{H^+} B$ half-life period is independent of concentration of A at given pH. At definite concentration of A half-life time is 10 min at pH 2 and half-life time is 100 min at pH 3. If the rate law expression of reaction is $r = k[A]^x[H^+]^y$ then calculate the value of $(x + y)$.

7. For a reaction, $A \rightleftharpoons B$ equilibrium constant is 1.66 and $k_{\text{forward}} = 0.166 \text{ hr}^{-1}$. Calculate the time (in hours) when concentration of B is 80% of its equilibrium concentration. (Given : $\ln 25 = 3.20$)

8. $A \xrightarrow{k_1} 3B$
 $A \xrightarrow{k_2} 8C$
 $A \xrightarrow{k_3} 3D$; At time $t = 0$, initial mole of A is 1.

Overall half life of the reaction is 15 days. Then calculate the number of mole of C after 45 days if the ratio of $k_1 : k_2 : k_3$ is 4 : 2 : 1

9. Iodine-131 is a radioactive isotope. If 1.0 mg of ^{131}I has an activity of $4.6 \times 10^{12} \text{ Bq}$, what is the half-life of ^{131}I (in days) ?
10. The average life of a radioactive element is 7.2 min. Calculate the time interval (in min.) between the stages of 33.33% and 66.66% decay
11. A, B and C are isodiaphers while C, D and E are isobars. Calculate the difference of protons between A and E $^{206}_{82}\text{A} \quad B \quad C \quad D \quad E$

Given : Isodiaphers and isobars are formed in successive α and β -emission respectively.

12. In the given radioactive disintegration series,



Calculate value of $(n - 2)$.

Where value of n is number of isobars formed in this series, suppose there is successive emission of β -particles.

13. In the given radioactive disintegration series



Calculate difference between number of α and number of β particles emitted in this series.

ANSWERS

Level 1

1. (c)	2. (b)	3. (c)	4. (d)	5. (d)	6. (d)	7. (d)	8. (c)	9. (a)	10. (c)
11. (c)	12. (a)	13. (d)	14. (d)	15. (b)	16. (d)	17. (c)	18. (c)	19. (d)	20. (a)
21. (c)	22. (c)	23. (b)	24. (a)	25. (c)	26. (c)	27. (d)	28. (c)	29. (b)	30. (c)
31. (d)	32. (a)	33. (d)	34. (a)	35. (b)	36. (c)	37. (a)	38. (b)	39. (a)	40. (c)
41. (c)	42. (d)	43. (c)	44. (b)	45. (c)	46. (b)	47. (b)	48. (b)	49. (a)	50. (b)
51. (d)	52. (c)	53. (c)	54. (d)	55. (d)	56. (a)	57. (d)	58. (a)	59. (a)	60. (c)
61. (c)	62. (c)	63. (c)	64. (a)	65. (a)	66. (c)	67. (b)	68. (d)	69. (d)	70. (c)
71. (d)	72. (a)	73. (a)	74. (b)	75. (d)	76. (c)	77. (a)	78. (d)	79. (d)	80. (c)
81. (b)	82. (b)	83. (c)	84. (c)	85. (c)	86. (c)	87. (c)	88. (b)	89. (c)	90. (c)
91. (d)	92. (d)	93. (d)	94. (b)	95. (b)	96. (c)	97. (b)	98. (d)	99. (d)	100. (c)
101. (c)	102. (b)	103. (c)	104. (d)	105. (d)	106. (b)	107. (a)	108. (c)	109. (d)	110. (b)
111. (a)	112. (c)	113. (d)	114. (c)	115. (b)	116. (c)	117. (c)	118. (c)	119. (b)	120. (c)
121. (d)	122. (c)	123. (b)	124. (d)	125. (d)	126. (b)	127. (c)	128. (c)	129. (b)	130. (b)
131. (d)	132. (b)	133. (d)	134. (c)	135. (c)	136. (d)	137. (c)	138. (b)	139. (b)	140. (b)
141. (c)	142. (b)	143. (d)	144. (c)	145. (b)	146. (c)	147. (a)	148. (b)	149. (b)	150. (b)

Level 2

1. (d)	2. (c)	3. (c)	4. (c)	5. (d)	6. (c)	7. (b)	8. (a)	9. (a)	10. (c)
11. (d)	12. (b)	13. (b)	14. (c)	15. (d)	16. (d)	17. (d)	18. (b)	19. (c)	20. (c)
21. (d)	22. (b)	23. (b)	24. (b)	25. (a)	26. (c)	27. (b)	28. (a)	29. (c)	30. (c)
31. (d)	32. (a)	33. (c)	34. (a)	35. (a)					



Level 3

Passage-1 1. (b) 2. (b) 3. (c)

Passage-2 1. (c) 2. (b) 3. (b)

Passage-3 1. (c) 2. (c)

Passage-4 1. (c) 2. (a) 3. (b)

Passage-5 1. (b) 2. (b)

One or More Answers is/are Correct

1. (a,c,d) 2. (a,b,c) 3. (a,b,c) 4. (d) 5. (a,c,d) 6. (b,c,d) 7. (a,b,c,d) 8. (a,b,c,d)
 9. (a,b,c,d) 10. (c) 11. (a,c,d) 12. (a,c,d) 13. (a,b,c) 14. (b,c) 15. (a,b,d) 16. (b,c)
 17. (a,b)

Match the Column

- | | | | |
|---------------|---------------|------------|-----|
| 1. A S; | B Q; | C P; | D R |
| 2. A Q; | B P, Q; | C R; | D S |
| 3. A R; | B S; | C Q; | D P |
| 4. A P, Q, S; | B P, Q, R, S; | C P, Q, R; | D P |
| 5. A P; | B R; | C Q; | D S |
| 6. A S; | B P; | C Q; | D R |
| 7. A P, R; | B S; | C Q; | D R |

Assertion-Reason Type Questions

1. (C) 2. (C) 3. (D) 4. (A) 5. (B) 6. (C) 7. (A) 8. (C) 9. (D) 10. (B)
 11. (B) 12. (D) 13. (D) 14. (A) 15. (B) 16. (A) 17. (C) 18. (A) 19. (A) 20. (B)

Subjective Problems

1. 6	2. 3	3. 3	4. 5	5. 6	6. 2	7. 6	8. 2	9. 8	10. 5
11. 2	12. 7	13. 3							

Hints and Solutions

Level 1

3. (c)

Rate of reaction	Rate of disappearance/appearance		
	Stoichiometric coefficient		
	$\frac{d[A]}{dt}$	$\frac{1}{2} \frac{d[B]}{dt}$	2.6×10^{-2}
	$\frac{1}{2} \frac{d[B]}{dt}$		
5.2×10^{-2}	$\frac{d[B]}{dt}$		

4. (d) Rate of reaction

$$\frac{1}{2} \frac{d[A]}{dt} = \frac{d[B]}{dt} = \frac{1}{3} \frac{d[C]}{dt}$$

or $\frac{k_1[A]^2}{2} = \frac{k_2[A]^2}{1} = \frac{k_3[A]^3}{3}$

$$\frac{k_1}{2} = k_2 = \frac{k_3}{3}$$

5. (d) Rate of reaction $K[\text{conc.}]^n$

$$K = \frac{\text{Rate}}{[\text{conc.}]^n} = \frac{\text{Ms}^{-1}}{\text{M}^n} = \frac{\text{mol L}^{-1} \text{s}^{-1}}{\text{mol}^n \text{L}^n}$$

6. (d) As explain in Q. No. 5

7. (d) $K_{\text{eq}} = \frac{k_f}{k_b} \frac{\text{Conc. of products as per rate law}}{\text{Conc. of reactants as per rate law}}$, stoichiometry of reaction is required.

8. (c) K_{eq} at 1400 K $\frac{0.29}{1.1 \times 10^{-6}} = 2.63 \times 10^5$; K_{eq} at 1500 K 9.28×10^4 ; Value of K_{eq} is reduces with rise in temperature so reaction is exothermic.

9. (a) Arrhenius eqⁿ. Rate constant $K = A e^{-E_a/RT}$

10. (c) As explain in Q. 3

Rate of disappearance	$\frac{d[\text{BrO}_3]}{dt}$
Rate of appearance	$\frac{d[\text{Br}_2]}{dt}$
Rate of reaction	$\frac{d[\text{BrO}_3]}{dt} = \frac{1}{3} \frac{d[\text{Br}_2]}{dt}$

11. (c) $x = kt$

$$t_{\text{completion}} = \frac{a}{k} = \frac{0.5}{0.1} = 5 \text{ min}$$

13. (d) Reaction mechanism required.


14. (d) $(a - x) = A_0 e^{-kt}$ or $A_t = A_0 e^{-kt}$
where A_t (a - x) conc. of reactant remained after time t
 A_0 a Initial concentration of reactant when $t = 0$

17. (c) Rate of reaction decreases with respect to times in 1st order Rx^n .

18. (c) $t_{1/2} = \frac{1}{a^{n-1}}$;

where, a : initial concentration of reactant
 n : order of reaction ($\because n \geq 1$)

19. (d) $a - x = a e^{-kt}$ (a - x)



Slope = -k

where,

a : initial conc. of reactant

x : conc. of reactant convert into product.

20. (a) $t_{1/2} = \frac{a}{2k} \log \frac{a}{a-x}$; $\log(t_{1/2}) = \log k + \log a$

21. (c) $t_{1/2} = \frac{1}{K a}$ for 2nd order ;

where a : initial conc. of reactant

22. (c) $t_{1/2} = \frac{a}{k}$; zero order

23. (b) $C = \frac{n}{V} = \frac{P}{RT}$; $\frac{dC}{dt} = \frac{1}{V} \frac{dn}{dt} = \frac{1}{RT} \frac{dP}{dt}$

24. (a) Rate of reaction $K[A_2]^a[B_2]^b$

0.04	$K(0.2)^a(0.2)^b$	...(i)
0.04	$K(0.1)^a(0.4)^b$	...(ii)
0.08	$K(0.2)^a(0.4)^b$	...(iii)

$$\frac{(i)}{(iii)} = \frac{1}{2} = \frac{1}{2}^b ; b = 1$$

$$\frac{(i)}{(ii)} = 1 = 2^a \cdot \frac{1}{2}^b ; a = 1$$

25. (c) $R_0 \propto K[A_0]^2[B_0]$... (i)
 $R_0 \propto K[1.5A_0]^2[3B_0]$
 $R_0 \propto (1.5)^2 \cdot 3[A_0]^2[B_0]$... (ii)
 From eq. (i) and (ii)
 $R_0 = 6.75 R_0$
26. (c) As explain in Q. 25
27. (d) Rate of reaction $k[A]^2$
 Rate of disappearance of A $-\frac{d[A]}{dt}$
 Rate of appearance of B $\frac{d[B]}{dt}$
 Rate of reaction $-\frac{d[A]}{dt} = \frac{d[B]}{dt} = k[A]^2$
 Rate of appearance of B after double the conc. of A.
 $\frac{d[B]}{dt} = k[2A]^2$
28. (c) Order w.r.t. A = 1; order w.r.t. B = 1
 R.O.R. $\frac{1}{2} \frac{d(AB)}{dt} = k_r[A][B];$
 $\frac{1}{2} (2.5 \cdot 10^{-4}) = k_r (0.1)(0.1)$
 $k_r = 1.25 \cdot 10^{-2}$
29. (b) Rate of appearance of $AB(g)$,
 $\frac{d(AB)}{dt} = k_b[A][B]$
 $k_b = 2.5 \cdot 10^{-2}$
30. (c) As explain in Q. 24
31. (d) For elementary reaction molecularity of reaction = order of reaction.
32. (a) As explain in Q. 24
33. (d) $r = k[A]^a[B]^b$
 Order of reaction $n = a + b = 2$
 In elementary reaction, partial order should be integer.
 So, $a = 1$ and $b = 1$
34. (a) $t_{1/2} = \frac{1}{(n-1)k} \frac{2^{n-1}}{a^{n-1}}$
35. (b) As explain in Q. 18
36. (c) As explain in Q. 5
37. (a) Rate $k[\text{conc.}]^0 = k$;
 for zero order reaction.
38. (b) Conc. of water (solvent(l)) remains constant so rate is independent from conc. of H_2O .
39. (a) $CH_3COOC_2H_5(aq) \xrightarrow{H} H_2O(l)$
 $CH_3COOH(aq) + C_2H_5OH(aq)$
 $r = \frac{(k_{\text{net}})_{H_2SO_4}}{(k_{\text{net}})_{HCl}}$ R.O.R. in presence of
 $H_2SO_4 >$ R.O.R. in presence of HCl
40. (c) $r_1 = k[A]^2[B]; r_2 = k[2A]^2[2B] = 8r_1$
41. (c) $[A]_t = [A] - kt = 1 - 0.001 \cdot 10 = 0.9$
 $[B]_t = 0.4M$
 $[B]_t = 0.001 \cdot 10 = 0.01M$
42. (d) $t_{100\%} = \frac{a}{k} = \frac{1}{0.6} \text{ min} = 100 \text{ sec.}$
 Reaction will be finish in 100 sec., so concentration of B will reach maximum value 2 M after 100 sec.
43. (c) $x = kt$ so $t_{1/2} = \frac{a}{2k}$ and $T = \frac{a}{k}$;
 $t_{1/2} = \frac{1}{2 \cdot 0.001} = 50 \text{ sec}$ and $T = 100 \text{ sec}$
45. (c) $x = 0.1 - 0.08 - 0.02M$;
 $k = \frac{x}{t} = \frac{0.02}{10} = 2 \cdot 10^{-3} M \text{ min}^{-1}$
 $t_{1/2} = \frac{[A]_0}{2K} = \frac{0.1}{2 \cdot 2 \cdot 10^{-3}} = 25 \text{ min}$
 $t_{\text{completion}} = 2 \cdot t_{1/2} = 50 \text{ min.}$
46. (b) Elementary reaction,
 $x(g) + y(g) \rightleftharpoons z(g)$
 Rate $k[x]$, first order
 $k = \frac{1}{t} \ln \frac{[A_0]}{[A_t]}$
 $\frac{\ln 2}{10 \text{ min}} = \frac{1}{t} \ln \frac{100}{10}$
 $\frac{0.3}{10} = \frac{1}{t} \ln 10$
 $t = 33 \text{ min.}$
47. (b) $K_1 = \frac{1}{5} \ln \frac{0.2}{0.1}; K_2 = \frac{1}{10} \ln \frac{0.2}{0.05}; K_1 = K_2$.
48. (b) $K = \frac{1}{100} \ln \frac{100}{25} = \frac{1}{t} \ln \frac{100}{12.5}$
49. (a) $K = \frac{0.693}{t_{1/2}} = \frac{0.693}{480} = 1.44 \cdot 10^{-3} s^{-1}$
50. (b) $k = \frac{2.303}{t} \log \frac{C_{A_0}}{C_A}$;
 $2.303 = 1 \cdot 2.303 \log \frac{C_{A_0}}{C_A}$

$$\frac{C_{A_0}}{C_A} = 10$$

$$C_A = \frac{1}{10} = 0.1$$

$$\text{rate after 1 min } r_1 = k C_A$$

$$2.303 \times 0.1 = 0.2303 \text{ M min}^{-1}$$

51. (d) Reaction is second order; $k = 10^4 \text{ L/mol. min.}$

$$\frac{1}{3} \frac{d[A]}{dt} = k[A]^2$$

$$\frac{d[A]}{dt} = 10^4 \times 3 \times (0.5)^2$$

$$7.5 \times 10^5 \text{ M min}^{-1}$$

$$\frac{7.5 \times 10^5}{60} = 1.25 \times 10^6 \text{ Ms}^{-1}$$

52. (c) $k = \frac{2.303}{t} \log \frac{a}{a-x}$

$$k = \frac{2.303}{32} \log \frac{100}{1} = \frac{2.303}{16}$$

$$t = \frac{2.303}{2.303} \log \frac{100}{0.1} = 48 \text{ min}$$

53. (c) $t_{3/4} = \frac{1}{K} \ln \frac{100}{1/4} = 100$

54. (d) $K_2 = \frac{1}{t} \ln \frac{100}{4} \dots (i)$

$$K_1 = \frac{1}{t} \ln \frac{100}{50} \dots (ii)$$

$$\frac{K_2}{K_1} = \frac{\ln 25}{\ln 2} = 4.65$$

55. (d) $k_I = \frac{0.693}{30}$

$$k_{II} = \frac{a}{2t_{1/2}} = \frac{2}{2 \times t_{1/2}} = \frac{1}{t_{1/2}} = \frac{1}{30}$$

$$\frac{k_I}{k_{II}} = \frac{0.693}{30 \times 1} = 30 \times 0.693$$

56. (a) $2\text{N}_2\text{O}_5(\text{CCl}_4) \rightarrow 2\text{N}_2\text{O}_4(\text{CCl}_4) + \text{O}_2(\text{g})$

$$t = 0 \quad a \text{ mol}$$

$$t = t \quad a - x \quad \frac{x}{2} \text{ mol}$$

$$t \quad 0 \quad \frac{a}{2} \text{ mol}$$

$$a = 100$$

$$x = 90$$

$$K = \frac{1}{t} \ln \frac{a}{a-x} = \frac{2.303}{500}$$

57. (d) $A \rightarrow 3B; P_T = P_0 + 2x$

$$\text{After time } t, P_0 = x + 3x$$

$$x = \frac{P_T - P_0}{2} = k \frac{1}{t} \ln \frac{P_0}{P_0 - x}$$

$$\text{After long time, } 0 = 3P_0$$

$$x = \frac{P_T - P}{2} = k \frac{1}{t} \ln \frac{\frac{P}{3}}{\frac{P}{3} - \frac{P}{6}} = k \frac{1}{t} \ln \frac{P}{\frac{P}{3}}$$

$$P = 3P_0 \quad x = \frac{3P_T - P}{6} = k \frac{1}{t} \ln \frac{P/3}{P/2 - P/6}$$

58. (a) $k = \frac{0.693}{2.1} = 0.33 \text{ hr}^{-1}$

Let t be the time for 90% decomposition,

$$\text{so } a = 100\% \quad x = 90 \text{ or } (a-x) = 10$$

$$k = \frac{2.303}{t} \log \frac{a}{a-x}$$

$$t = \frac{2.303}{0.33} \log \frac{100}{10} ; t = 6.978 \text{ hr}$$

$$n_{\text{NH}_4\text{NO}_3} \text{ taken} = \frac{6.2}{62} = 0.1;$$

as per given no. of moles of N_2O produced

$$0.1 \times \frac{90}{100}$$

Vol. of N_2O produced at STP

$$0.09 \times 22.4 = 2.016 \text{ L}$$

59. (a) $A \rightarrow 2B + C$

$$t = 0 \quad P_i \quad 0 \quad 0$$

$$t = t \quad P_i - y \quad 2y \quad y$$

$$\text{Total pressure } P_t = P_i - y + 2y + y = P_i + 2y$$

$$\frac{P_t - P_i}{2} = y$$

$$K = \frac{2.303}{t} \log \frac{2P_i}{3P_i - P_t}$$

60. (c)

$$k = \frac{2.303}{t} \log \frac{P_i}{P_t} ; (\text{CH}_3)_2\text{N}_2 \rightarrow \text{C}_2\text{H}_6 + \text{N}_2$$

$$t = 0 \quad 200 ; \quad t = 200 \quad x \quad x \quad x$$

$$\text{as per given } 200 \quad x \quad 350 \quad x \quad 150$$

$$k \frac{2.303}{t} \log \frac{200}{200 - 150}$$

$$k = 5.77 \times 10^{-4} \text{ sec}^{-1}$$

$$61. (c) k \frac{1}{t} \ln \frac{r}{r_0} = \frac{r_0}{r_t} \frac{1}{t} \ln \frac{r_0}{r_t} \frac{r}{t}$$

$$k \frac{1}{70} \ln \frac{44}{16.5} = \frac{11}{11}$$

$$k \frac{\ln 2}{70} \text{ when solution is optically inactive}$$

$$r_t = 0$$

$$k \frac{1}{t} \ln \frac{r_0}{r} = \frac{r}{r_0} \frac{1}{t} \ln \frac{r_0}{r}$$

$$\frac{\ln 2}{70} = \frac{1}{t} \ln \frac{44}{11} ;$$

$$\frac{0.7}{70} = \frac{1}{t} \ln 1.6 ; t = 160 \text{ min}$$

$$62. (c) \frac{(t_{1/2})_1}{(t_{1/2})_2} = \frac{a_2}{a_1} \frac{n-1}{n-1} ; n = 1 \frac{\log t_{1/2, 1}}{\log t_{1/2, 2}}$$

$$63. (c) kt = \frac{1}{n-1} \frac{1}{A_t^{n-1}} - \frac{1}{A_0^{n-1}}$$

$$kt_{0.5} = \frac{1}{n-1} \frac{2^{n-1}}{A_0^{n-1}} \quad \dots(1)$$

$$kt_{0.875} = \frac{1}{n-1} \frac{8^{n-1}}{A_0^{n-1}}$$

$$\frac{t_{0.875}}{t_{0.5}} = \frac{8^{n-1}}{2^{n-1}} \frac{1}{1}$$

$$64. (a) kt = \ln \frac{0}{t} ; k = 60 \ln \frac{80}{75}$$

$$k = \frac{1}{60} \ln \frac{80}{75}$$

$$k = 5 \times 10^{-6} \ln \frac{80}{75} ; 300 \frac{1}{60}$$

$$\ln \frac{80}{75} = \ln \frac{80}{r_t} ; \ln \frac{80}{r_t} = 0.32$$

$$r_t = 58$$


optical rotation observed after 5 hrs

$$58 - 2 \times 60$$


$$65. (a) \text{ For zero order } t_{1/2} \propto [\text{conc.}]^0$$

$$66. (c) \frac{1}{2} \frac{dC_B}{dt} = k_1 C_A ; \frac{dC_B}{2t} = 2k_1 C_A$$

$$2 \times 10^{-3} = 2 \times 8 \times 10^{-3} M s^{-1}$$



$$68. (d) E_{a2} = 58$$



70. (c) First step is slow (require more activation energy) second step is fast (less activation energy) and overall reaction exothermic, so product energy level should be less as compared to reactants.

71. (d) Slow step require larger activation energy and product energy level is always more than reactant energy level in endothermic reaction.

$$72. (a) \text{ Rate of Rx}^n = K [\text{Conc.}]^n$$

$$73. (a) K = A e^{-E_a/RT}$$

$$\ln K = \ln A - \frac{E_a}{RT}$$

$$74. (b) K = A e^{-E_a/RT}$$

Given $K = A e^{-1000/RT}$

$\frac{E_a}{RT} = 1000$

$$75. (d) k = \frac{k_1 \cdot k_2}{k_3} \quad ; \quad E = \frac{2}{3} [E_{a1} + E_{a2} - E_{a3}]$$

$\frac{2}{3} [180 + 80 - 50] = 140 \text{ kJ/mol}$

$$76. (c) A_1 e^{-E_{a1}/RT} = A_2 e^{-E_{a2}/RT}$$

$$\frac{A_2}{A_1} = e^{(E_{a2} - E_{a1})/RT}$$

$$10 = \exp \frac{600}{RT} , R = 2 \text{ cal/K-mol}$$

$$\ln 10 = \frac{600}{2T}$$

$$T = \frac{300}{2.303} \text{ K}$$

- $$E_b \quad E_f$$

$$\ln \frac{k_{2(320)}}{k_{1(300)}} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$\ln \frac{K_2}{K_1} = \frac{E_a}{R} \left(\frac{1}{290} - \frac{1}{300} \right)$$

$$\frac{\ln 2}{\ln \frac{K_2}{K_1}} = \frac{10}{\frac{280}{290} - \frac{290}{300}}$$

$$\frac{0.693}{\ln \frac{K_2}{K_1}} = \frac{300}{280}$$

$$\frac{K_2}{K_1} = 1.9$$

- 86.** (c) $e^{\frac{E_a}{RT}}$ $3.8 \frac{10^{16}}{100}$

$$\frac{E_a}{RT} \ln 3.8 \cdot 10^{18}$$

E_a 100 kJ/mol

$\frac{k_2}{k_1}$	exp	$\frac{8.314}{8.314}$	$\frac{1000}{500}$
-------------------	-----	-----------------------	--------------------

exp (2) 7.38

- Rate
- $K[A][B]$

- $$K_2 \frac{[A]^2}{[A_2]} \dots(\text{ii})$$

$$R = K_1 \sqrt{K_2[A_2][B_2]} = K_1 K_2^{1/2}[A_2]^{1/2}[B] K[A_2]^{1/2}[B_2]$$

$$(\because K \leq K_1 K_2^{1/2})$$

- 92.** (d) Due to charge on the and . Rays.

93. (d)
- | | | | |
|--|--|--|--------------------------------------|
| $\begin{matrix} A \\ Z \end{matrix} X$ | $\begin{matrix} A \\ Z \end{matrix} Y$ | $\begin{matrix} 0 \\ 1 \end{matrix}$ | |
| $\begin{matrix} A \\ Z \end{matrix} Y$ | $\begin{matrix} A \\ Z \end{matrix} Z$ | $\begin{matrix} 4 \\ 1 \end{matrix}$ | $\begin{matrix} 4 \\ 2 \end{matrix}$ |
| $\begin{matrix} A \\ Z \end{matrix} Z$ | $\begin{matrix} A \\ Z \end{matrix} Z$ | $\begin{matrix} A \\ Z \end{matrix} Z$ | $\begin{matrix} 0 \\ 0 \end{matrix}$ |
| High energy nucleus | Low energy nucleus | Energy | |

- 95. (b)**
- | | | | |
|---------|---------------------------------|---------------------------------|-----------|
| | $\frac{A}{Z}x$ | $\frac{A}{Z}y$ | 0_1e |
| Neutron | $A \quad Z$ | $A \quad (Z \quad 1)$ | |
| Proton | Z | $Z \quad 1$ | |
| | $\frac{n}{p} \quad \frac{A}{Z}$ | $\frac{A}{Z} \quad \frac{1}{1}$ | |

- 96.** (c) As explain in Q. 95

- 104.** (d) Isotones have same no. of neutron;

17 9 *B* 8

B 16

Isobars have same mass no.

A	B	16
-----	-----	----

no. of neutrons $16 - 8 = 8$

- 105.**(d) ${}_{90}^{234}\text{Th}$ ${}_{82}^{206}\text{Pb}$ x_2^4 2 y_1^0

$$\begin{array}{cccc} 234 & 206 & 4x & 0 \end{array}$$

$x = 7$ -particle

90	82	$2x$	y
----	----	------	-----

γ 6

- 106.(b)** Same number of neutron.

- 107.(a)** Same mass number

- 108.(c)** Same difference between n and p

- 110.**(b) ${}^7_3\text{Li}$ ${}^1_1\text{P}$ ${}^4_2\text{X}$ 4_2 2_2

- 112.(c)** As explain Q. No. 110

- 113.** (d) ${}_{92}\text{U}^{235}$ ${}_0n^1$ ${}_{54}\text{X}^{139}$ ${}_{38}\text{Sr}^{194}$ ${}_0n^1$

- 114.(c)** ${}_{101}^{255}\text{Md}$ $\begin{smallmatrix} 4 \\ 2 \end{smallmatrix}$ 2_0^1n ${}_{103}^{257}\text{Lr}$

- 115.(b)** ${}_{92}^{235}\text{U} + {}_0^1n \rightarrow {}_{38}^{94}\text{Sr} + {}_{54}^{140}\text{Xe} + 2{}_0^1n$


- 116. (c)** $E_{\text{total}} = 200 \frac{(3^5 - 1)}{3 - 1}$

100 242 24200 MeV

- 117.(c)** $^{230}_{90}\text{Th}$ ^1_1P $^{24}_2\text{He}$ $^{223}_{87}\text{Fr}$

- 118.**(c) On emission of one α particle, formed nuclei occupied two position left in periodic table.

120.(c) On emission of positron atomic no. increases mass no. remain constant.



122.(c) $A = N$

$$6.023 \times 3.7 \times 10^{10} = 3.7 \times 10^4 N;$$

$$N = 6.023 \times 10^6 \text{ atoms}$$

$$1 \text{ mole or } 6.023 \times 10^{23} \text{ atoms } 100 \text{ g of } X$$

$$6.023 \times 10^6 \text{ atoms}$$

$$\frac{100}{6.023 \times 10^{23}} \times 6.023 \times 10^6 = 10^{-15} \text{ g}$$

123.(b) $A_1 = A_0 e^{-t_1}$... (i)

$A_2 = A_0 e^{-t_2}$... (ii)

$\frac{A_2}{A_1} = e^{-(t_1 - t_2)}$... (iii)

124.(d) Four half-lives (Total time n half-life so, $n = 4$), hence 0.0625

125.(d) $\frac{0.693}{69.3} \times 10^2 \text{ min}^{-1};$

$$N = \frac{dN}{dt} \times \frac{100}{10^2} = 10,000$$

126.(b) $A_0 = \frac{N_0}{28} \frac{\ln 2}{30} N_0$
 $N_0 = 1212$

129.(b) Equal fraction decay in equal periods of time, fraction of sample remaining after

$$3 \text{ days } (0.9)^3 = 0.729$$

131.(d) $\frac{N_0}{N} = \frac{A_0}{A} = 8;$

$$t = \frac{1}{\lambda} \ln \frac{A_0}{A} = \frac{5770}{0.693} \ln 8$$

$$17310 \text{ year}$$

133.(d) t for 20% left

$$t_1 = \frac{2.303}{\lambda} \log \frac{1}{1 - 0.8}$$

$$\frac{\ln 5}{\lambda}$$

t for 50% left

$$t_2 = \frac{1}{\lambda} \ln 2$$

$$\frac{t_2}{t_1} = \frac{\frac{1}{\lambda} \ln 2}{\frac{\ln 5}{\lambda}} = 0.43$$

134.(c) $t = \frac{t_{0.5}}{0.693} \cdot \ln \frac{r_0}{r} = \frac{5770}{0.693} \ln \frac{120}{30};$
 $t = \frac{5770}{0.693} \times 2 = 11,540 \text{ years}$

135.(c) ${}_{19}^{40}\text{K} \xrightarrow{\beta} {}_{20}^{40}\text{Ca} + {}_1^0e$
 At present 2.5×10^6 7.5×10^6 mole
 moles of ${}_{19}^{40}\text{K}$ consumed 7.5×10^6 mole
 initial mole of ${}_{19}^{40}\text{K}$ was $(7.5 + 2.5) \times 10^6$ mole

$$t = \frac{1}{\lambda} \ln \frac{10 \times 10^6}{2.5 \times 10^6} = \frac{1.3 \times 10^9}{\ln 2} \ln(4)$$

$$1.3 \times 10^9 \times 2 = 2.6 \times 10^9 \text{ yr}$$

136.(d) $t_{\text{avg}} = \frac{1}{\lambda} = \frac{14}{0.693}$

137.(c) $t = |t_1 - 3t_{1/2}| = 200$

138.(b) Total time $nt_{1/2}; n = 5; \frac{\text{Initial activity}}{2^n}$

$$\frac{\text{Initial activity}}{2^5} = 0.32 \text{ } C_i$$

139.(b) $t = \ln \frac{A_0}{A} = \frac{0.693}{t_{1/2}} \times 200 = \ln \frac{2140}{535}$

$$t_{1/2} = 100 \text{ min.}$$

$$\frac{1}{10} \ln \frac{100}{100 - 25}$$

$$t = \frac{1}{\lambda} \ln \frac{N_0}{N}$$

$$t = \frac{10}{\ln \frac{4}{3}} \ln \frac{4 \times 10^{20}}{10^{20}}$$

$$t = 48.19$$

142.(b) $t_{1/2} = \frac{t}{4}; t_{1/2} = T \ln 2$

$$\text{so } \frac{t}{4} = T \ln 2; t = 4T \ln 2$$

143.(d) W_0 initial mass, W mass. after 20 days

$$\frac{2.303}{t} \log \frac{W_0}{W_A};$$

$$_B \frac{2.303}{t} \log \frac{W_0}{W_B}$$

$$_B \frac{2.303}{t} \log \frac{W_B}{W_A}$$

$$(t_{1/2})_B \quad 1.10 \text{ day}$$

$$144. (c) \text{ Given } (n_0)_B = 8 \quad (n_0)_A; (n)_A = 2 \quad (n)_B$$

$$_A \quad _B \frac{2.303}{t} \log \frac{(N_A)_0}{(N_A)} \frac{(N_B)}{(N_B)_0};$$

$$t \frac{\ln \frac{1}{16}}{\frac{0.693}{50} \frac{0.693}{10}}$$

$$= 50 \text{ min}$$

$$145. (b) \text{ as per given } \frac{dN}{dt} \quad N \text{ for max. no. of}$$

nuclei;

$$\frac{dN}{dt} = 0; \quad N, N -$$

$$146. (c) \frac{\text{Atoms of Sr}}{\text{Atoms of Rb}} = 0.05$$

$$\text{or } \frac{\text{Atoms of Sr}}{\text{Atoms of (Sr + Rb)}} = 1.05$$

$$\text{so, } \frac{\text{Initial no. of atoms of Rb}}{\text{Present no. of atoms of Rb}} = 1.05$$

$$t \frac{2.303}{\log (n_0/n)}$$

$$\frac{2.303}{0.693} = 4.7 \quad 10^{10} \log (1.05)$$

$$= 3.28 \quad 10^9 \text{ year}$$

$$147. (a) \quad {}_A N_A \quad {}_B N_B$$

($\therefore$ rate of disintegration are same)

$$\frac{N_A}{N_B} = \frac{B}{A}$$

$$\frac{2}{1}$$

$$148. (b) N = N_0 \cdot e^{-t}; \text{ where } N \text{ Parent remaining}$$

and N_0 Initial parent Parent remaining
(p) + daughter formed (d)

$$p = (p - d) \cdot e^{-t} \text{ or } \ln \frac{(p - d)}{p} = -t$$

$$t \frac{1}{\ln 1} = \frac{d}{p}$$

$$149. (b) \quad 1 \quad 2; \quad \frac{0.693}{22} \text{ and } \frac{1}{2} \frac{2}{98}$$

$$150. (b) \quad {}_{84}^{218}\text{Po} \xrightarrow{1} {}_{82}^{214}\text{Pb} + {}_2^4\text{He} \xrightarrow{2} {}_{83}^{214}\text{Bi} + {}_{-1}^0e$$

Pb^{214} to reach max. no. of nuclei

$$t_{\text{max.}} = \frac{1}{1} \ln \frac{1}{2}$$

$$= 247.5 \text{ sec}$$

$$\text{where } 1 \quad \frac{0.693}{183}; \quad 2 \quad \frac{0.693}{161}$$

Level 2

$$1. (d) \quad K_{\text{eq}} = \frac{k_f}{k_b} \frac{[\text{CH}_3]^2}{[\text{C}_2\text{H}_6]}$$

$$[\text{CH}_3] = \frac{10^{-4}}{10} = 10^{-5} \text{ M}$$

$$\frac{1.57 \cdot 10^{-3}}{k_b} = \frac{(10^{-5})^2}{1}$$

$$k_b = 1.57 \cdot 10^7 \text{ L mol}^{-1} \text{ s}^{-1}$$

$$3. (c) \quad r_1 = k[0.01]^a [0.01]^b = 6.93 \cdot 10^{-6} \dots (i)$$

$$r_2 = k[0.02]^a [0.01]^b = 1.386 \cdot 10^{-5} \dots (ii)$$

$$r_3 = k[0.02]^a [0.02]^b = 1.386 \cdot 10^{-5} \dots (iii)$$

From data $a = 1; b = 0;$

overall order $= 1; k = 6.93 \cdot 10^{-4} \text{ sec}^{-1}$

$$6.93 \cdot 10^{-4} = \frac{1}{50 \cdot 60} \ln \frac{A_0}{A_t};$$

when $[A]_0 = 0.5$

$$A_t = 0.0625$$

$$\text{rate of reaction} = \frac{6.93 \cdot 10^{-4} \cdot 0.0625}{4.33 \cdot 10^{-5} \text{ Ms}^{-1}}$$

$$4. (c) \quad \frac{r_2}{r_1} = \frac{P_{A_2} \cdot P_{B_2}^2}{P_{A_1} \cdot P_{B_1}^2} = \frac{0.1 \cdot (0.4)^2}{0.4 \cdot 1^2}$$

$$\frac{1}{25}$$

$$6. (c) \quad t_1 = \frac{(t_{1/2})_1}{0.693} \ln \frac{1}{(1 - (1/4))}$$

$$t_2 = \frac{(t_{1/2})_2}{0.693} \ln \frac{1}{1 - 3/4}$$

$$\frac{t_1}{t_2} = \frac{8}{1} \frac{\ln (4/3)}{\ln (4)}$$

$$1 : 0.602$$

$$7. (b) \quad \begin{array}{cccc} & A & B & C & D \\ t = 0 & 1 & 1 & & \\ t & 1 & x & 1 & x & x & x \end{array}$$

$$\frac{dx}{dt} = k[A]^{1/2}[B]^{1/2}$$

$$\frac{dx}{dt} = k(1-x)^{1/2}(1-x)^{1/2}$$

or $\frac{dx}{dt} = k(1-x)$

$$t = \frac{1}{k} \ln \frac{1}{1-x};$$

$$t = \frac{2.303}{2.303 \times 10^{-2}} \log \frac{1}{0.1}$$

$$= 100 \text{ sec}$$

8. (a)

(a) True $[B]_t = k_1 t; 0.25 \frac{1}{\sqrt{3}} t; t = 0.25\sqrt{3}$

$$[D]_t = k_{II} t = \sqrt{3} \cdot 0.25 \cdot \sqrt{3} = 0.75 M$$

(b) False if $[C] = [A]$ then at that time $[B] = [D]$

(c) False $t_{100\%} = \frac{a}{k}$ (for zero order)

$$\frac{(t_{100\%})_I}{(t_{100\%})_{II}} = \frac{a_I}{a_{II}} \cdot \frac{k_{II}}{k_I} = \frac{0.5}{1} \cdot \frac{\sqrt{3}}{1/\sqrt{3}} = \frac{3}{2}$$

(d) $[A]_t = [A]_0 - k_1 t$ or $[A]_t = 0.5 - \frac{1}{\sqrt{3}} t$

$$[C]_t = [C]_0 + k_{II} t$$
 or $[C]_t = 1 + \sqrt{3} t$

if $[A] = [C]_t$

$$i.e. 0.5 - \frac{1}{\sqrt{3}} t = 1 + \sqrt{3} t \text{ or } (\sqrt{3} - \frac{1}{\sqrt{3}}) t = 0.5$$

$$t = \frac{\sqrt{3}}{4} \text{ min.}$$

9. (a)

$$t_{1/2} = \frac{0.693}{k}$$

$$= \frac{0.693}{2.772 \times 10^{-3}}$$

$$= 250 \text{ sec}$$

Since volume is changing therefore half-life should be defined on the basis of moles, so moles of A remaining after 250 sec. Also final volume of the container

$$V_f \frac{n_f}{n_i} = V_i \frac{0.05 + 0.15}{0.1} = 2$$

$$= 4 \text{ litre}$$

$$\text{Final concentration of A} = \frac{0.05}{4} = 0.0125 M$$

10. (c)

	A	2B	C
t = 0	P_i	0	0

t	P_i	x	2x	x
t	0		$2P_i$	P_i
P	$3P_i$ or P_i	$\frac{P}{3}$	P_i	$2x$
x	$\frac{P_t - P_i}{2}$			

$$\text{As we know } k = \frac{2.303}{t} \log \frac{P_i}{P_t - x}$$

$$\text{so } k = \frac{2.303}{t} \log \frac{2P}{3(P - P_t)}$$

11. (d) Let n is the moles of reagent 'R' when R is reacted with A at time t = 0

	A	B	C
t = 0	n	0	0
at t	n - x	2x	3x
at t	0	2n	3n
	5n	n ₂	n $\frac{n_2}{5}$
n	4x	n ₁	x $\frac{n_1}{4}$
k	$\frac{2.303}{t} \log \frac{n}{n - x}$		

$$\text{so } k = \frac{1}{t} \ln \frac{4n_2}{5(n_2 - n_1)}$$

12. (b)

	A(g)	2B(g)	C(g)
Let initial pressure	P_0	0	0
After 10 min.	$(P_0 - x)$	2x	x
After long time (t)	0	$2P_0$	P_0
as per given $(P_0 - x) = 2x$	x	vapour	
pressure of H_2O	188		

$$P_0 = 2x \quad 160 \text{ and } 3P_0 = 28 \quad 388$$

$$\text{so, } P_0 = 120 \text{ and } x = 20 \text{ torr}$$

$$k = \frac{1}{t} \ln \frac{P_0}{P_0 - x}$$

$$\frac{1}{10} \ln \frac{120}{100} = \frac{1}{10} (\ln 4 - \ln 3 - \ln 10)$$

$$0.02 \text{ min}^{-1} = 1.2 \text{ hr}^{-1}$$

13. (b)

$$k = \frac{2.303}{t} \log \frac{r}{r_t} \frac{r_0}{r_t}$$

$$\frac{2.303}{30} \log \frac{11}{11} \frac{34}{19}$$

$$= 1.35 \times 10^{-2} \text{ min}^{-1}$$

when solution is optically inactive $r_t = 0$;

$$t \frac{1}{k} \ln \frac{45}{11} = 103.7 \text{ min}$$

14. (c) Overall rate constant

$$k \frac{k_1 k_2}{6.93 \times 10^3} = 100 \text{ sec};$$

$$\text{After half-life, } P_B = \frac{P_C}{k_1} = \frac{P_D}{k_2} = \frac{4 \text{ atm}}{200}$$

$$\frac{P_B}{P_C} = \frac{P_D}{k_1} = \frac{200}{693}$$

$$= 1.154 \text{ atm}$$

15. (d) $A \xrightarrow{k_1} 2B; A \xrightarrow{k_2} C$

$$\frac{d[A]}{dt} = (k_1 + k_2)[A]; \quad \frac{[B]}{[C]} = \frac{2k_1}{k_2} \frac{x}{y}$$

$$\therefore \frac{k_1}{k_2} = \frac{x}{y}$$

$$(k_1 + k_2)t = \ln \frac{A_0}{A_t}$$

$$2 \times 6.93 \times 10^3 = 50 \ln \frac{A_0}{A_t}$$

$$[A]_t = \frac{[A_0]}{2} \quad a_0 \times x \times y = a_0/2 \quad (\because x = y)$$


$x = a_0/4$. Total moles in container

$$a_0 \times x \times y = 2x \times y = a_0 \times x = 1.25 a_0$$

% increase in moles = 25, So final pressure

$$2 \times 1.25 = 2.5 \text{ atm}$$

16. (d) 1 mole of A will form 2 moles of B and C. After completion of reaction when 75% A converted into B and C then total no. of moles = 0.5 + 2 = 1.5 = 3.5 moles



$$\text{Initial } a \quad 0; K_{(aq)} = \frac{k_f}{k_b} = \frac{3 \times 10^{-4}}{0.1} = 3 \times 10^{-3}$$

$$\text{at time } t \quad a - x \quad x$$

$$\text{at eqm} \quad a - x_e \quad x_e$$

$$\text{As we know } (k_f/k_b) = \frac{1}{t} \ln \frac{x_e}{x_e - x}$$

$$\text{given } x = \frac{x_e}{2}$$

$$(k_f/k_b) = \frac{1}{2} \ln 2$$

$$\text{or } (3 \times 10^{-3} = 3 \times 10^{-4}) \frac{0.693}{t}$$

$$t = 210 \text{ sec}$$

18. (b) $k = A e^{-E_a/RT}; A = e^{\frac{E_a}{RT_1}} = A e^{\frac{(E_a - x)}{RT_2}}$

$$\frac{E_a}{500} = \frac{(E_a - 16)}{400} \quad E_a = 80 \text{ kJ/mol}$$

19. (c) For parallel reactions

$$E_a = \frac{k_1 E_{a1}}{k_1 + k_2} = \frac{k_2 E_{a2}}{k_1 + k_2}$$

$$\frac{10^2}{10^2} = \frac{100}{4} = \frac{4}{10^2} = \frac{120}{10^2}$$

$$E_a = 116 \text{ kJ/mol}$$

20. (c) $k = A e^{-E_a/RT}$

Effective overall energy of activation

$$E_a = \frac{E_{a(2)}}{2} = \frac{E_{a(3)}}{2} = \frac{1}{2} E_{a(1)} = \frac{1}{2} E_{a(5)}$$

$$60 = 50 = \frac{1}{2} = 40 = \frac{1}{2} = 10$$

$$25 \text{ kJ/mol}$$

21. (d) $A_1 e^{-E_{a1}/RT} = A_2 e^{-E_{a2}/RT};$

$$\frac{A_2}{A_1} = e^{(E_{a2} - E_{a1})/RT}$$

$$10^2 = \text{Exp} \frac{1200}{RT}; 2 \ln 10 = \frac{1200}{2T}$$

$$T = \frac{600}{4.606} \text{ K}$$

22. (b) $\ln \frac{k_1}{k_2} = \frac{E_1}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right) \dots (i)$

$$\ln \frac{k_2}{k_2} = \frac{E_2}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right) \dots (ii)$$

Solving we get (ii)-(i),

$$\ln \frac{k_2}{k_2} - \ln \frac{k_1}{k_1} = \frac{E_2 - E_1}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$$

(for equimolar formation of B and C, $k_2 = k_1$)

$$\ln \frac{k_1}{k_2} = \frac{8314}{8.314} \left(\frac{T_2 - 300}{300 T_2} \right)$$

$$\ln 2 = (1000) \left(\frac{T_2 - 300}{300 T_2} \right)$$

$$T_2 = 378.74 \text{ K}$$

24. (b) $A \xrightleftharpoons[\text{fast}]{\text{slow}} B$ IAB; So $E_{a(f)}$ is high and $E_{a(b)}$ is low. $k_1 \ll k_2$; So, E_a for this step is very high and for next step is low and overall reaction is exothermic.

25. (a) Activation energy

$$E_I \quad E_{II} \quad E_V \quad E_{VI}$$

$$E_{1(f)} \quad E_{1(b)} \quad E_{2(b)} \quad E_{3(f)} \quad E_{3(b)} \quad E_{4(f)} \quad E_{5(f)} \quad E_{6(b)}$$

$$E_{2(f)} \quad E_{4(b)} \text{ (slow steps so larger)}$$

26. (c) $e^{-\frac{E_a}{RT}} = 100 \cdot 3.8 \cdot 10^{-16}$; $e^{-\frac{E_a}{RT}} = 3.8 \cdot 10^{-18}$
- $$\frac{E_a}{RT} = \ln(3.8 \cdot 10^{-18});$$
- $$\frac{R}{E_a} = 8.314 \text{ and } T = 300$$
- $$E_a = 100 \text{ kJ/mol}$$

27. (b) $t = \ln \frac{A_0}{A}$
- $$\frac{0.693}{30} = 90 \ln \frac{A_0}{A} \quad A_0 = A \cdot 8$$
- $$N_0 = \frac{A_0}{0.693} = \frac{800}{30} = 3.46 \cdot 10^4 \text{ atoms}$$

28. (a) at time $X = \frac{n_0}{x}$; $Y = \frac{n_0}{x}$; $\frac{1}{7}$; $n_0 = \frac{8x}{7}$
- $$\frac{0.693}{6.93 \cdot 10^9} = 10^{10} \text{ years}^{-1}$$
- $$t = \frac{2.303}{\ln 8} \log \frac{w_0}{w_0 - x}$$
- $$\frac{2.303}{10^{10}} \log(8)$$
- $$= 2.079 \cdot 10^{10} \text{ years}$$

29. (c) $\text{net OR } \frac{1}{2} \cdot 1.8 \cdot 10^3 = 2 \cdot 10^3$
- $$\text{Average life } (T_{av}) = \frac{1}{2 \cdot 10^2} = 50 \text{ sec}$$

30. (c) ${}_{83}\text{Bi}^{211} \rightarrow {}_{81}\text{Tl}^{207} + {}_2\text{He}^4$; total time n half-life
- moles of substance left after n halves
- $$\frac{\text{initial moles}}{2^n} = \frac{2}{2^4} = 0.125$$
- moles of He produced
- $$2 \cdot 0.125 = 1.875$$
- Pressure developed due to

$$\text{He} \quad \frac{1.875}{100} \quad \frac{0.0821}{100} \quad \frac{300}{100}$$

$$= 0.4618 \text{ atm}$$

31. (d) Let initial activities of A and B are A_0 and B_0
- ($\therefore$ after 2 half-lives of activity of A will remain $\frac{A_0}{4}$)

$$A_0 = B_0 = 8000 \text{ and also } \frac{A_0}{4} = B_0 = 3500$$

(we can assume that activity of B remains constant due to larger half-life)

$$\text{so } \frac{3A_0}{4} = 4500; A_0 = 6000; B_0 = 2000;$$

$$\frac{A_0}{B_0} = \frac{6000}{2000} = \frac{3}{1}$$

32. (a) No change in activity of sample during establishment of circulatory equilibrium.
- Let volume of blood is V mL, so total vol. $(V + 1)$ mL after injection of sample.
- 2 mL sample has activity of 10 dpm, so $(V + 1)$ mL sample has activity of $\frac{10}{2} (V + 1)$

Since rate is constant so

$$\frac{10}{2} (V + 1) = 1000; V = 199 \text{ mL}$$

33. (c) $\frac{A_{0(X)}}{A_{0(Y)}} = \frac{4}{1}; \frac{A_x}{A_y} = 1$
- $$y = x \cdot \frac{1}{t} \ln \frac{(A_0)_y}{(A_0)_x} = \frac{A_x}{A_y}$$
- $$(y - x)t = \ln \frac{1}{4}; (t_{1/2})_y = 30 \text{ min.}$$

34. (a) Corrected C^{14} content :
- Let initial C^{14} count A_0 ;
- $$A_0 = A_0/10 = 15.4; A_0 = 14$$
- $$t = \frac{1}{\lambda} \log \frac{14}{7} = 5.770 \cdot 10^3 \text{ years}; t = \frac{\ln 2}{\lambda}$$
- $$t_{1/2}; t = 5770 \text{ years}$$

35. (a) For U^{238} ; $t = \ln \frac{n_0}{n_1}$
- For U^{235} ; $t = \ln \frac{n_0}{n_2}$

$$(t_1 - t_2)t = \ln \frac{n_0}{n_1} - \ln \frac{n_0}{n_2};$$

$$\ln 2 \frac{1}{0.5 \times 10^9} \frac{1}{4.5 \times 10^9} t \ln(140)$$

$$t = 4.02 \times 10^9 \text{ year}$$

Level 3

Passage-1

1. (b) For max. con. of B;

$$\frac{d[B]}{dt} = 0; \text{ so } t_{\max} = \frac{1}{k_1 - k_2} \ln \frac{k_1}{k_2}$$

2. (b) $k_1 < k_2$

So rate of appearance of B is much lesser than rate of disappearance of B.

Passage-2

1. (c) $e^{-E_a/RT}$ is fraction of molecules having energy greater than activation energy (E_a), it is unit less. So A has same unit of K.

for 1st order reaction unit of K is s^{-1} .

2. (b) Answer is given in (1)

$$x = e^{-E_a/RT}$$

$$\ln x = \frac{E_a}{RT}$$

3. (b) $\ln \frac{K_2}{K_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$

$$\ln 2 = \frac{E_a}{2 \text{ cal}} \left(\frac{1}{290} - \frac{1}{300} \right)$$

$$E_a = \frac{2.303 \times 0.3 \times 2 \times 290 \times 300}{10} \text{ cal}$$

$$12 \text{ K Cal}$$

Passage-3

1. (c) $E = \frac{nhC}{\lambda}$

$$7.2 \times 10^3 \text{ J/s} = \frac{n \times 6.6 \times 10^{34} \times 3 \times 10^8}{330 \times 10^9}$$

No. of photon

$$\frac{7.2 \times 10^3 \times 330 \times 10^9}{6.6 \times 10^{34} \times 3 \times 10^8} = 120 \times 10^{14}$$

$$\text{Mole of photon} = \frac{120 \times 10^{14}}{6 \times 10^{23}} = 20 \times 10^9$$

Mole of CO formed

$$0.4 \times 20 \times 10^9 = 8 \times 10^9 \text{ mol/s}$$

2. (c) Rate of formation of $C_2H_6 = 0.8 \times 20 \times 10^9$
 $16 \times 10^9 \text{ mol/s}$

Passage-4

1. (c) $\frac{1}{t} \ln \frac{N_0}{N}$

$$\frac{\ln 2}{t_{1/2}} = \frac{1}{t} \ln \frac{N_0}{N}$$

$$\frac{\ln 2}{2.5 \times 10^5} = \frac{1}{t} \ln 4$$

$$t = 2.5 \times 10^5 \times 2 = 5 \times 10^5 \text{ years.}$$

2. (a) $\frac{N}{N_0} = \left(\frac{1}{2} \right)^n$; where n no. of halves;

$$\frac{N}{N_0} = \left(\frac{1}{2} \right)^n$$

$$n = 4$$

$$\text{Total time} = n \times t_{1/2} = 4 \times 1185 \text{ years}$$

3. (b) $\frac{dN}{dt} = \lambda N$

$$\frac{dN}{dt} = \frac{0.693}{24,000} N$$

$$1 \times 6.02 \times 10^{23} \text{ dps or } 14.9 \text{ Ci}$$

Passage-5

1. (b) $r = R_0 A^{1/3}$

Volume of nucleus $\left(\frac{4}{3} \right) \pi r^3$

$$\frac{4}{3} \pi (1.5 \times 10^{-15})^3 A \text{ m}^3$$

$$\text{density } d = \frac{m}{V} = \frac{A \times 1.66 \times 10^{-27} \text{ kg}}{\left(\frac{4}{3} \right) \pi (1.5 \times 10^{-15})^3 A}$$

$$1.17 \times 10^{17} \text{ kg/m}^3$$

2. (b) $d_c = \frac{\text{Mass}}{\text{Volume}} = \frac{12 \times 1.66 \times 10^{-27}}{\left(\frac{4}{3} \right) \pi (3 \times 10^{-15})^3}$

$$1.76 \times 10^{17} \text{ kg/m}^3$$

$$d_{H_2O} = 1000 \text{ kg/m}^3$$

$$\frac{d_c}{d_{H_2O}} = \frac{1.76 \times 10^{17}}{1,000} = 1.76 \times 10^{14}$$

One or More Answers is/are Correct

2. (A) $K = A e^{-E_a/RT}$ as T, K rate of reaction

- (B) Rate of reaction $k[A]^0$ K constant
 (C) As surface area No. of molecules will be more to react per unit time, more surface site available for the reaction.
 (D) Average rate defined for macroscopic time and instantaneous rate defined for microscopic time.
3. For an elementary reaction $aA + bB \rightarrow cC + dD$ rate law is always $r = K[A]^a[B]^b$ but not vice versa.
 For a complex reaction $aA + bB \rightarrow cC + dD$ rate law may or may not be $r = K[A]^a[B]^b$.
4. (A) In a rate law reactant, product or catalyst may be appeared.
 (B) We can not
 (C) In an elementary reaction only positive integer stoichiometric coefficient appears.
 (D) A zero order reaction is always complex
5. (A) $K = A e^{-E_a/RT}$
 As T , K so if T , then K A
 or as E_a , K so if E_a 0, then K A
 (B) Catalyst does not change H of reaction.
 (C) A negative catalyst decrease rate of reaction by increasing activation energy.
6. $A + B \rightarrow C$
 (A) If it is an elementary reaction then it will be bimolecular reaction.
 (B) It may be exothermic eq.
 $NH_3(g) + HCl(g) \rightarrow NH_4Cl(s)$
 (C) It may be heterogeneous
 (D) It may be photochemical reaction
7. $A(g) + 2B(g) \rightarrow C(g)$
 $t = 0$ 400
 $t = 20$ 400 x $2x$ x
 So $400 - x - 2x - x = 1000$
 $x = 300$ mmHg
 $K = \frac{1}{20} \ln \frac{400}{100} = \frac{2 \ln 2}{20} = 0.0693 \text{ min}^{-1}$
 $t_{1/2} = \frac{\ln 2}{K} = \frac{0.693}{0.0693} = 10 \text{ min}$
 $\frac{30}{10} \ln 2 = \ln \frac{400}{400 - x}$
 $8 \frac{400}{400 - x} = 400 - x = 50$
 $x = 350$ mmHg

Partial pressure of C after 30 min 350 mmHg

Total pressure after 30 min

400 $2x$ 400 700 1100 mmHg

9. (a,b,c,d)

$$k = \frac{B_e}{A_e} \frac{1.6}{0.4} = 4; k = 4 \frac{k_1}{k_2} = \frac{4}{k_2} \frac{10^{-2}}{k_2}$$

$$k_2 = 10^{-2}$$


$$t_{0.5x_e} = \frac{1}{k} \ln \frac{x_e}{x_e - x}$$

$$\frac{1}{5 \cdot 10^{-2}} \ln 2 = \frac{70}{2}; t_{0.5x_e} = 14 \text{ min};$$

$$t_{0.1a} = \frac{1}{5 \cdot 10^{-2}} \ln \frac{x_e}{x_e - x} = \frac{1}{5 \cdot 10^{-2}} \ln \frac{1.6}{1.6 - 0.2}$$

$$\frac{100}{5} \ln \frac{16}{14} = 20 \cdot 0.14 = 2.8$$

10. (A) α -particles are just nucleus of helium (${}^4_2\text{He}^{2+}$)



11. (A) ${}^{235}_{92}\text{U} \rightarrow {}^{207}_{82}\text{Pb} + {}^{4}_{2}\text{He} + 6\text{e}^-$
 Atomic no. 92 0 56 (2 0) Z
 $Z = 36$
 Mass no. 235 1 140 (2 1) A
 $A = 236$ 172 94

So X is ${}^{94}_{36}\text{Kr}$

(B) ${}^{23}_{11}\text{Na} \rightarrow {}^{23}_{12}\text{Mg} + {}^0_{-1}e$
 Atomic no. 11 Z 12 0 Z 1
 Mass no. 23 A 23 1 A 1
 $X = {}^1_1\text{H}$ Proton

(C) and (D) both are correct.

12. (A) SI unit of radioactivity is Bq

1 Bq = 1 dps

(B) 1 Ci = 3.7×10^{10} dps or Bq

13. Similar as question 11.

14. ${}^M_Z A \rightarrow {}^M_Z B + {}^M_Z C + {}^M_Z D$

15. (a, b)

${}^1_1\text{P} \rightarrow {}^0_0 n + \text{X-rays}$

17. (a, b)

${}^1_1\text{P} \rightarrow {}^0_0 n + {}^0_{-1}e$ (positron); ${}^1_1\text{P} \rightarrow {}^0_0 n + {}^0_{+1}e$ (Beta)

Activity N , Decay constant does not depend upon temperature.

By the emission of α -particle, isodiaphers are formed.

By the emission of β -particle, isobars are formed.

Match the Column

- (A) K $Ae^{E_a/RT}$, unit of K unit of A
 (B) $K = \frac{\text{Rate}}{(\text{Concentration})^n} = \frac{M/s}{(M)^n} = M^{1-n} s^{-1}$
 $n = 0$ Unit of K $M s^{-1}$
 (C) $n = 1$ Unit of K s^{-1}
 (D) $n = 2$ Unit of K $M^{-1} s^{-1}$
- (A) Molecularity of a reaction never be zero.
 (B) Order of reaction may be 0, 1, 2, 3 fraction
 (C) $H_2O_2 \rightarrow H_2O + \frac{1}{2} O_2$ if its an elementary then it is 1st order reaction.
 (D) It is zero order reaction because reaction rate is depend on due to absorption of photon.
- (A) $C_t = C_0 - Kt$ slope K
 (B) $\log C_t = \log C_0 - \frac{K}{2.303} t$; slope $K/2.303$
 (C) $\frac{dC}{dt} = K[C]$; slope zero
 (D) $\ln \frac{dc}{dt} = \ln K + \ln C$; slope 1
- (A) $\frac{d[A]}{dt}$ vs $[A]$, Let the order of reaction w.r.t. A is x then for non zero slope $[A]^x$, and $x = 0$
 (B) $K = Ae^{E_a/RT}$
 $\ln K = \ln A - \frac{E_a}{RT}$
 It is applicable for all possible order in this slope 0
 (C) $t_{1/2} [A]_0$ it is not possible for 1st order.
 (D) $\frac{d[A]}{dt}$ vs $[A]^2$, $[A]^2$ second order.
- (A) Concentration of product increases w.r.t time.
 (B) Concentration of reactant decreases w.r.t time.
 (D) $K = Ae^{E_a/RT}$
 $\ln K = \ln A - \frac{E_a}{RT}$ slope $-\frac{E_a}{R}$
- (A) Isotone atoms having same number of neutrons (i.e. value of mass number (A) – atomic number (Z)).
 (B) Isobare Atoms has same mass no. and different atomic no.
 (C) Isotope Atoms has same atomic no. and difference mass no.

(D) Isodiaphers Same difference between neutrons and proton.

- (A) ${}_Z^AX$ ${}_Z^{A-4}Y$
 (B) ${}_Z^AX$ ${}_Z^{A-1}Y$, atomic no. increase.
 (C) ${}_Z^AX$ ${}_Z^AY$, no. change in mass no. and atomic no.
 (D) ${}_Z^AX$ ${}_{Z-1}^AY$, Atomic no. decrease.

Subjective Problems

- $2NH_3(g) \rightleftharpoons N_2(g) + 3H_2(g)$
 Rate $K[NH_3]^0$
 Rate 2
 Rate $\frac{1}{3} \frac{[H_2]}{t} = \frac{[H_2]}{t} \times 6$
- $t_{1/2} = \frac{[A]_0}{2K_A} = \frac{[A]_0}{2 \times 5K} = \frac{200 \text{ min.}}{2 \times 5 \times 400}$
 $\frac{60 \text{ sec.}}{20} \times 3$
- | | $A(g)$ | $2B(g)$ | $C(g)$ |
|-------------------|-------------------------------|---------|--------|
| $t = 0$ | 0.4 | 0.6 | |
| $t = t$ | 0.3 | 0.4 | 0.1 |
| r_i | $KP_A^1 P_B^2$ | | |
| r_i | $K \times 0.4 \times (0.6)^2$ | | |
| r_i | $K(0.3)(0.4)^2$ | | |
| $\frac{r_i}{r_i}$ | 3 | | |
- Rate of reaction $K[CO]^x[O_2]^y$
 $4 \times 10^{-5} = K(0.02)^x(0.02)^y \dots(i)$
 $1.6 \times 10^{-4} = K(0.04)^x(0.02)^y \dots(ii)$
 $8 \times 10^{-5} = K(0.02)^x(0.04)^y \dots(iii)$
 $\frac{(i)}{(ii)} = \frac{1}{4} = \frac{1}{2}^x$
 $\frac{(ii)}{(iii)} = \frac{1}{2} = \frac{1}{2}^y$
 $x = 2$
 $\frac{(i)}{(iii)} = \frac{1}{2} = \frac{1}{2}^y$
 $y = 1$
 From eq. (1)
 $4 \times 10^{-5} = K(0.02)^2(0.02)^1$
 $K = \frac{4 \times 10^{-5}}{4 \times 10^{-4} \times 2 \times 10^{-2}} = \frac{10}{2} \times 5$
- For zero order $2(t_{1/2})_0 = t_{\text{Complete}} ;$
 $(t_{1/2})_0 = \frac{1.386}{2} \dots(ii)$

For first order $(t_{1/2})_1 = \frac{0.693}{k_1}$... (ii)

Given $(t_{1/2})_0 = \frac{13.86}{2}$ $(t_{1/2})_1 = \frac{0.693}{k_1}$

$k_1 = 0.1 \text{ min}^{-1} = 0.1 \times 60 \text{ hr}^{-1} = 6 \text{ hr}^{-1}$

6. $r = k[A]^x[H]^y$
Order of reaction w.r.t. A = 1
(x = 1); $r = k[H]^y$ ($\therefore k = k[A]^1$)

$\frac{a/2}{10} = k[10^{-2}]^y$... (i)

$\frac{a/2}{100} = k[10^{-3}]^y$... (ii)


(i)/(ii); $y = 1$;

where a = initial concentration.

7. $(k_f - k_b)t = \ln \frac{x_e}{x_e - x} = \ln \frac{x_e}{x_e - 0.8x_e}$;

$(k_f - k_b)t = \ln \frac{x_e}{0.2x_e} = \ln 5 = 1.6$;

$\frac{k_f}{k_b} = K_c, k_b = 0.1t = 6 \text{ hrs.}$



mole of A remain after 45 days

$\frac{N_0}{2^n} = \frac{N_0}{\frac{T}{2^{t_{1/2}}}} = \frac{1}{2^{45/15}} = \frac{1}{2^3} = \frac{1}{8}$

moles of A convert into product $\frac{7}{8} \text{ mol}$

moles of [C] $\frac{K_2}{K_1 + K_2 + K_3} = \frac{7}{8} \times \frac{8}{1} = 7$

9. $A \xrightarrow{N} 4.6 \times 10^{12} \text{ Bq}$
 $\frac{\ln 2}{t_{1/2}} = \frac{1}{131} \times \frac{10^3}{6} \times 10^{23}$

$t_{1/2} = \frac{0.693 \times 6 \times 10^{20}}{4.6 \times 10^{12} \times 131 \times 3600 \times 24} = 7.98 \text{ day}$

10. $t_{1/2} = 8$
 $\frac{1}{t_{\text{avg}}} = \frac{1}{t} \ln \frac{[A_0]}{[A_t]}$; $\frac{1}{7.2} = \frac{1}{t_1} \ln \frac{100}{100 - \frac{100}{3}}$;
 $\frac{1}{7.2} = \frac{1}{t_2} \ln \frac{100}{100 - \frac{200}{3}}$;

$t_2 - t_1 = 7.2 \ln 3 = 7.2 \ln \frac{3}{2} = 5$

11. $^{206}_{82}A$ $^{202}_{80}B$ $^{298}_{78}C$ $^{198}_{79}D$ $^{198}_{80}E$.

Proton difference in A and E = 2

12. Number of α -particle $\frac{232 - 208}{4} = 6$

$Z_2 - Z_1 = 2x - y$
(where x = number of α and y = number of β)
 $82 - 90 = 12 - y$
 $y = 4$

Series

$^{A}_{Z}P$ $^{A}_{Z-1}Q$ $^{A}_{Z-2}R$ $^{A}_{Z-3}S$ $^{A}_{Z-4}T$

Total number of isobars = 5

$n = 5$, so $n - 2 = 7$

13. Number of α -particle $\frac{235 - 207}{4} = 7$

$Z_2 - Z_1 = 2x - y$
(where x = number of α and y = number of β)
 $82 - 92 = 14 - y$
 $y = 4$

Number of α particle - number of β particle = 3

ELECTROCHEMISTRY

Electrolytic Cell

- ❖ **Electrolysis** : Chemical changes in electrolyte solution by passage of electric current, resulting into deposition of metals or liberation of gases at electrodes is known as electrolysis.
- ❖ **Electrolytic Cell** : This cell converts electrical energy into chemical energy. The entire assembly except that of the external battery is known as the electrolytic cell.

Quantitative Aspect of Electrolysis

Faraday's laws of electrolysis :

- ❖ **First law of electrolysis** : Amount of substance deposited or liberated at an electrode is directly proportional to amount of charge passed (utilized) through the solution.

$$\text{Current efficiency} = \frac{\text{Actual charge utilised in process}}{\text{Charge passed through battery}} \times 100$$

W mass liberated/deposited, Q charge in coulomb, i current, t time
 Z electrochemical equivalent, 1 Faraday = 96500 coulomb

- ❖ **Second law of electrolysis** : When same amount of charge is passed through different electrolytic solutions connected in series then mass of substances deposited or dissolved at anode or cathode are in ratio of their equivalent mass. i.e., $W_1/W_2 = E_1/E_2$


$$\frac{W_1}{W_2} = \frac{E_1}{E_2}$$

Products of Electrolysis

S.No.	Electrolyte	Anode Product	Cathode Product
1.	NaCl (Molten) with Pt electrode	$\text{Cl}_2(g)$	$\text{Na}(l)$
2.	NaCl (aq) with Pt electrode	$\text{Cl}_2(g)$	$\text{H}_2(g)$
3.	$\text{Na}_2\text{SO}_4(aq)$ with Pt electrode	$\text{O}_2(g)$	$\text{H}_2(g)$
4.	$\text{NaNO}_3(aq)$ with Pt electrode	$\text{O}_2(g)$	$\text{H}_2(g)$
5.	$\text{AgNO}_3(aq)$ with Pt electrode	$\text{O}_2(g)$	$\text{Ag}(s)$
6.	$\text{CuSO}_4(aq)$ with inert electrode	$\text{O}_2(g)$	$\text{Cu}(s)$
7.	$\text{CuSO}_4(aq)$ with copper electrode	Cu dissolves	$\text{Cu}(s)$


Galvanic Cell

This cell converts chemical energy into electrical energy.



Galvanic cell is made up of two half cells *i.e.*, anodic and cathodic. Oxidation takes place at anode and reduction at cathode. It is also known as **voltaic cell**. It may be represented as shown in Fig. Zinc rod immersed in ZnSO_4 behaves as anode and copper rod immersed in CuSO_4 behaves as cathode.

❖ **Anode half cell reaction :**



❖ **Cathode half cell reaction :**



❖ **Over all process :**



In galvanic cell like Daniel cell; electrons flow from anode (zinc rod) to the cathode (copper rod) through external circuit; zinc dissolves as Zn^{2+} ; Cu^{2+} ion in the cathode cell picks up two electron and become deposited at cathode.

Salt Bridge

It is U-shaped tube contains saturated solution of inert electrolyte like KCl, KNO_3 , NH_4Cl and NH_4NO_3 etc. in agar-agar gel.

- ❖ Cation and anion of inert electrolyte have same mobility.
- ❖ Ions of inert electrolyte do not mix with electrolytic solution present in the half cell.
- ❖ Ions of inert electrolyte do not participate in electrochemical change.

Function of Salt Bridge


- ❖ It complete the electrical circuit.
- ❖ It maintained the two half cell electrically neutral by the flow of ions.

Representation of a cell (IUPAC conventions) : Let us explain the convention taking the example of Daniel cell.

- (i) Anodic half cell is written on left and cathodic half cell on right hand side.



- (ii) Two half cells are separated by double vertical lines and it indicate salt bridge or any type of porous partition.
- (iii) EMF (electromotive force) may be written on the right hand side of the cell.
- (iv) Single vertical lines indicate the phase separation between electrode and electrolyte solution.



Nernst's Equation

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{2.303RT}{nF} \log Q$$

At 298 K temp. $E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.0591}{n} \log Q$; where Q is reaction quotient.

Cell Thermodynamics

(i) $G = -nFE_{\text{cell}}$

(ii) $G^{\circ} = -nFE_{\text{cell}}^{\circ}$

(iii) Equilibrium constant (K) , $\log K = \frac{n E_{\text{cell}}^{\circ}}{0.0591}$

(iv) Temp. coefficient of cell $\frac{E}{T}_P$


(v) Enthalpy of reaction inside the cell, $H = -nFE_{\text{cell}} - nFT \frac{E}{T}_P$


(vi) Entropy change inside the cell, $S = nF \frac{E}{T}_P$

Concentration Cell

Anode and cathode both are formed by same substance so $E_{\text{cell}} = 0$, such type of cell is known as concentration cell. Concentration gradient may arise either in electrode material or in electrolyte. Thus there are two types of concentration cells.

❖ **Electrode gas conc. cell :**



❖ **Electrolyte concentration cell :**

$$E_{\text{cell}} = \frac{0.0591}{2} \log \frac{C_2}{C_1}$$

Different Type of Electrodes

S.No.	Name of Electrode	Anode	Cathode
1.	Hydrogen electrode	$\text{Pt(s)} \text{H}_2(\text{g}) \text{H}^+(\text{aq})$ $E_{\text{OP}} \quad E_{\text{OP}}^\circ \quad \frac{0.0591}{2} \log \frac{[\text{H}^+]}{P_{\text{H}_2}}$	$\text{H}^+(\text{aq}) \text{H}_2(\text{g}) \text{Pt(s)}$ $E_{\text{RP}} \quad E_{\text{RP}}^\circ \quad \frac{0.0591}{2} \log \frac{P_{\text{H}_2}}{[\text{H}^+]^2}$
2.	Metal-metal ion electrode	$\text{M(s)} \text{M}^{n+}(\text{aq})$ $E_{\text{OP}} \quad E_{\text{OP}}^\circ \quad \frac{0.0591}{n} \log [\text{M}^{n+}]$	$\text{M}^{n+}(\text{aq}) \text{M(s)}$ $E_{\text{RP}} \quad E_{\text{RP}}^\circ \quad \frac{0.0591}{n} \log \frac{1}{[\text{M}^{n+}]}$
3.	Calomel electrode	$\text{Hg(l)}, \text{Hg}_2\text{Cl}_2(\text{s}) \text{Cl}^-(\text{aq})$ $E_{\text{OP}} \quad E_{\text{OP}}^\circ \quad \frac{0.0591}{2} \log \frac{1}{[\text{Cl}^-]^2}$	$\text{Cl}^-(\text{aq}) \text{Hg}_2\text{Cl}_2(\text{s}), \text{Hg(l)}$ $E_{\text{RP}} \quad E_{\text{RP}}^\circ \quad \frac{0.0591}{2} \log [\text{Cl}^-]^2$
4.	Redox electrode	$\text{Pt(s)} \text{Fe}^{2+}(\text{aq}), \text{Fe}^{3+}(\text{aq})$ $E_{\text{OP}} \quad E_{\text{OP}}^\circ \quad \frac{0.0591}{1} \log \frac{[\text{Fe}^{3+}]}{[\text{Fe}^{2+}]}$	$\text{Fe}^{3+}(\text{aq}), \text{Fe}^{2+}(\text{aq}) \text{Pt(s)}$ $E_{\text{RP}} \quad E_{\text{RP}}^\circ \quad \frac{0.0591}{1} \log \frac{[\text{Fe}^{2+}]}{[\text{Fe}^{3+}]}$
5.	Metal insoluble salt anion electrode	$\text{Ag(s)} \text{AgCl(s)} \text{Cl}^-(\text{aq})$ $E_{\text{OP}} \quad E_{\text{OP}}^\circ \quad \frac{0.0591}{1} \log \frac{1}{[\text{Cl}^-]}$	$\text{Cl}^-(\text{aq}) \text{AgCl(s)} \text{Ag(s)}$ $E_{\text{RP}} \quad E_{\text{RP}}^\circ \quad \frac{0.0591}{1} \log [\text{Cl}^-]$

Batteries

Electrochemical cells can be used as **batteries**. Batteries are of two kinds :

- ❖ **Primary battery :** Where the reaction occurs only once and can not be reused once it becomes dead over the course of time. For examples, dry cell (Leclanche cell), Mercury cell.
- ❖ **Secondary battery :** Which can be recharged by passing current through it in the opposite direction so that it can be used again. For example, lead-acid battery, Nickel-cadmium cell.

Fuel Cells

Fuel cells are the galvanic cells that convert the energy of combustion of fuels (e.g., hydrogen, methane, methanol, etc.) directly into electrical energy.


Fuel cells are more efficient thermodynamically, and more of the energy of the reaction can be made available for useful work provided that the supply of reactants is maintained.

Methanol, ethanol, hydrazine, formaldehyde, carbon monoxide can be used as fuels in fuel cells apart from hydrogen.

Corrosion

When the metal exposed to some environment it gets converted to its oxides. The oxidative deterioration of metal is known as corrosion.

Example : Rusting of iron, tarnishing of silver, development of green coating on copper and bronze etc.



Hydrated ferrous oxide ($\text{Fe}_2\text{O}_3 \cdot x\text{H}_2\text{O(s)}$) is known as rust.

❖ Factors which enhance corrosion :

- Presence of impurities in the metal.
- Presence of moisture.
- Presence of electrolyte.

❖ Prevention of corrosion :

- Barrier protection by oil/grease layer, paints or electroplating.
- Sacrificial protection by mixing the metal with more electropositive metal.

Conductance

❖ **Conductance (G):** It is defined as the reciprocal of the electrical resistance i.e., $G = 1/R$. It measures the ease with which the current flows through a conductor.

Unit : Siemens, S or Ω^{-1} .

❖ **Specific resistance or resistivity (ρ) :** The resistivity or specific resistance is defined as the resistance in ohm of a conductor having length equal to 1 cm and area of cross-section equal to 1 cm^2 .

$$R = \frac{\rho l}{A} \text{ or } \rho = R \frac{A}{l} \text{ where } \rho \text{ specific resistance, } l \text{ length of conductor}$$

A area of cross-section of the conductor.

Unit : ohm cm.

❖ **Specific conductance or conductivity (k):** It is defined as the reciprocal of specific resistance.

$$\begin{aligned}
 \text{i.e., } k &= \frac{1}{\rho} \quad R = \frac{l}{A} \quad k = G \frac{l}{A} \\
 k &= \frac{G}{G^*} \quad G^* = \frac{l}{A}, \quad \text{where } G^* \text{ cell constant}
 \end{aligned}$$

When $l = 1 \text{ cm}$ and $A = 1 \text{ cm}^2$; $k = G$

Thus conductivity is the conductance of one centimeter cube of the solution of an electrolyte.

Unit : $\Omega^{-1} \text{cm}^{-1}$, Scm^{-1} .


- ❖ **Molar conductivity (Λ_m):** The conducting power of all the ions produced by dissolving 1 mole of an electrolyte in solution.

Mathematically, $\Lambda_m = \frac{k}{M}$ (S cm² mol⁻¹) where k is conductivity and M is molarity.

Variation of conductivity and molar conductivity with concentration :

For weak and strong electrolytes

- ❖ Conductivity decreases with concentration.
- ❖ Molar conductivity increases with decrease in concentration.



Kohlrausch's Law

According to this law, at infinite dilution, when the dissociation is complete, each ion makes a definite contribution towards molar conductivity of the electrolyte irrespective of the nature of the other ions present.

The molar conductivity of an electrolyte at infinite dilution is the sum of the ionic conductivities of the cations and the anions each multiplied by the number of ions present in one formula unit of the electrolyte

The equivalent conductivity of an electrolyte at infinite dilution is the sum of the equivalent conductivities of the cations and anions.

e.g., $A_x B_y$. $\Lambda_m^0(A_x B_y) = x \Lambda_m^0(A^+) + y \Lambda_m^0(B^-)$
 $\Lambda_{eq}^0(A_x B_y) = \Lambda_{eq}^0(A^+) + \Lambda_{eq}^0(B^-)$

Application

(a) Determination of equivalent/molar conductivities of weak electrolytes at infinite dilution

$$\Lambda_m^0(\text{CH}_3\text{COOH}) = \Lambda_m^0(\text{CH}_3\text{COONa}) - \Lambda_m^0(\text{HCl}) + \Lambda_m^0(\text{NaCl})$$

(b) Determination of degree of dissociation (α) and equilibrium constant (K) of an electrolyte at a given dilution

$$\frac{\text{Molar conductivity at concentration } C}{\text{Molar conductivity at infinite dilution}} = \frac{\Lambda_m}{\Lambda_m^0}$$

$$\text{HA} \rightleftharpoons \text{H}^+ + \text{A}^-$$

Initial conc.	C	0	0
Conc. at equilibrium	$C - C\alpha$	$C\alpha$	$C\alpha$

$$K_a = \frac{C}{1} \cdot \frac{C}{1} = \frac{C^2}{1}$$

(c) Determination of the solubility of a sparingly soluble salt

Since the solution is saturated at infinite dilution 0 and molarity 0 solubility.


$$\text{Solubility (S)} = \frac{k}{1000} \cdot \frac{1000}{M}$$

For sparingly soluble salt $A_x B_y$: Solubility product $K_{sp} = x^x y^y S^{x+y}$



Conductometric Titration

Titration in which end point can be determined by measuring conductivity at each addition of the titre solution (for burette) is called conductometric titration.

Titration of HCl(aq) with NaOH(aq) (Strong acid vs Strong base)



Titration of $\text{CH}_3\text{COOH(aq)}$ with NaOH(aq) (Weak acid vs Strong base)





Level 1

1. A cell reaction would be spontaneous if the cell potential and $\Delta_r G$ are respectively :
(a) positive and negative (b) negative, negative
(c) zero, zero (d) positive, zero
2. Which of the following statement is correct?
(a) Cathode is -ve terminal in both, galvanic and electrolytic cells
(b) Anode is +ve terminal in both, galvanic and electrolytic cells
(c) Cathode and anode are -ve terminal in electrolytic and galvanic cell.
(d) Cathode and anode are +ve terminal in electrolytic and galvanic cell.
3. Electrolytes when dissolved in water dissociate into ions because :
(a) they are unstable.
(b) the water dissolves it.
(c) the force of repulsion increases.
(d) the force of electrostatic attraction is broken down by water.
4. The electric charge required for electrode deposition of one gram-equivalent of a substance is :
(a) one ampere per second (b) 96500 coulombs per second
(c) one ampere for one hour (d) charge on one mole of electrons
5. The amount of an ion liberated on an electrode during electrolysis does not depend upon :
(a) conductance of the solution (b) current strength
(c) time (d) electrochemical equivalent of the element
6. How many electrons are there in one coulomb electricity?
(a) 6.023×10^{23} (b) 1.64×10^{-24} (c) 6.24×10^{18} (d) 6.24×10^{-24}
7. How many coulombs are provided by a current of 0.010 mA in the calculator battery that can operate for 1000 hours?
(a) 1.0 (b) 10 (c) 0.010 (d) 36
8. How many minutes are required to deliver 3.21×10^6 coulombs using a current of 500 A used in the commercial production of chlorine?
(a) 8.3 (b) 5.3×10^4 (c) 6420 (d) 107
9. Passage of a current for 548 seconds through a silver coulometer results in the deposition of 0.746 g of silver. What is the current (in A)?
(a) 1.22 (b) 1.16 (c) 1.07 (d) 1.00
10. Electrolysis can be used to determine atomic masses. A current of 0.550 A deposits 0.55 g of a certain metal in 100 minutes. Calculate the atomic mass of the metal if eq. mass = mol. mass/3
(a) 100 (b) 45.0 (c) 48.25 (d) 144.75
11. Beryllium occurs naturally in the form of beryl. The metal is produced from its ore by electrolysis after the ore has been converted to the oxide and then to the chloride. How many grams of Be(s) is deposited from a BeCl_2 solution by a current of 5.0 A that flows for 1.0 h? (Atomic weight : Be = 9)
(a) 0.840 (b) 1.68 (c) 1.42 (d) 1.08

12. How many minutes will it take to plate out 5.2 g of Cr from a $\text{Cr}_2(\text{SO}_4)_3$ solution using a current of 9.65 A? (Atomic mass : Cr = 52.0)
(a) 200 (b) 50 (c) 100 (d) 103
13. Calculate the current (in mA) required to deposit 0.195 g of platinum metal in 5.0 hours from a solution of $[\text{PtCl}_6]^{2-}$: (Atomic mass : Pt = 195)
(a) 310 (b) 31 (c) 21.44 (d) 5.36
14. How many Faradays are required to reduce 0.25 g of Nb (V) to the metal? (Atomic mass : Nb = 93)
(a) 2.7×10^{-3} (b) 1.3×10^{-2} (c) 2.7×10^{-2} (d) 7.8×10^{-3}
15. One gm metal M^{3+} was discharged by the passage of 1.81×10^{23} electrons. What is the atomic mass of metal?
(a) 33.35 (b) 133.4 (c) 66.7 (d) None of these
16. Total charge required for the oxidation of two moles Mn_3O_4 into MnO_4^{2-} in presence of alkaline medium is :
(a) 5 F (b) 10 F (c) 20 F (d) None of these
17. The electrolytic decomposition of dilute sulphuric acid with platinum electrode, cathodic reaction is :
(a) reduction of H (b) oxidation of SO_4^{2-} (c) reduction SO_3^{2-} (d) oxidation of H_2O
18. Which one of the following metals can not be obtained on electrolysis of aqueous solution of its salts ?
(a) Mg (b) Ag (c) Cu (d) Cr
19. A highly concentrated solution of potassium sulphate in water is electrolysed using inert electrodes. The products at the cathode and anode are respectively.
(a) H_2, O_2 (b) O_2, H_2 (c) O_2, Na (d) $\text{H}_2, \text{S}_2\text{O}_8^{2-}$
20. The passage of current through a solution of certain electrolyte results in the evolution of $\text{H}_2(\text{g})$ at cathode and $\text{Cl}_2(\text{g})$ at anode. The electrolytic solution is :
(a) Water (b) aq. H_2SO_4 (c) aq. NaCl (d) aq. CuCl_2
21. When an aqueous solution of H_2SO_4 is electrolysed, the product at anode is :
(a) H (b) OH (c) SO_4^{2-} (d) O_2
22. An aqueous solution of Na_2SO_4 is electrolysed using Pt electrodes. The products at the cathode and anode are respectively :
(a) H_2, SO_2 (b) O_2, NaOH (c) H_2, O_2 (d) O_2, SO_2
23. The electrolysis of a solution resulted in the formation of $\text{H}_2(\text{g})$ at the cathode and $\text{O}_2(\text{g})$ at the anode. The solution is :
(a) $\text{AgCl}(\text{aq})$ (b) $\text{H}_2\text{SO}_4(\text{aq})$
(c) highly concentrated $\text{NaCl}(\text{aq})$ solution (d) $\text{CuCl}_2(\text{aq})$
24. If Pt is used as cathode in the electrolysis of aqueous NaCl solution, the ion produced at cathode is :
(a) H (b) Na (c) OH (d) Cl
25. A dilute aqueous solution of CuSO_4 is electrolysed using platinum electrodes. The products at the anode and cathode are :
(a) O_2, H_2 (b) H_2, O_2 (c) O_2, Cu (d) $\text{S}_2\text{O}_8^{2-}, \text{H}_2$


26. What products are formed during the electrolysis of concentrated aqueous solution of sodium chloride?
 (I) $\text{Cl}_2(\text{g})$ at anode (II) NaOH as electrolyte (III) $\text{H}_2(\text{g})$ at cathode
 (a) I only (b) I and II only (c) I and III only (d) I, II and III
27. Which of the following aqueous solution produces metal after electrolysis ?
 (a) $\text{K}_2\text{Cr}_2\text{O}_7$ (b) KMnO_4 (c) CH_3COONa (d) CuCl_2
28. How much time is required for complete decomposition of 4 moles of water using 4 ampere?
 (a) $3.86 \times 10^5 \text{ sec}$ (b) $1.93 \times 10^5 \text{ sec}$ (c) 96500 sec (d) 48250 sec
29. An aqueous solution containing 1 M each of Au^{3+} , Cu^{2+} , Ag^+ , Li^+ is being electrolysed by using inert electrodes. The value of standard reduction potentials are :
 $E_{\text{Ag}^+/\text{Ag}} = 0.80 \text{ V}$, $E_{\text{Cu}^{2+}/\text{Cu}} = 0.34 \text{ V}$, and $E_{\text{Au}^{3+}/\text{Au}} = 1.50 \text{ V}$, $E_{\text{Li}^+/\text{Li}} = -3.03 \text{ V}$,
 With increasing voltage, the sequence of deposition of metals on the cathode will be :
 (a) Li, Cu, Ag, Au (b) Cu, Ag, Au (c) Au, Ag, Cu (d) Au, Ag, Cu, Li
30. If 0.50 L of a 0.60 M SnSO_4 solution is electrolysed for a period of 30.0 min using a current of 4.60 A. If inert electrodes are used, what is the final concentration of Sn^{2+} remaining in the solution? [at. mass of Sn = 119]
 (a) 0.342 M (b) 0.544 M (c) 0.389 M (d) 0.514 M
31. A 100.0 mL dilute solution of Ag is electrolysed for 15.0 minutes with a current of 1.25 mA and the silver is removed completely. What was the initial $[\text{Ag}^+]$?
 (a) 2.32×10^{-1} (b) 2.32×10^{-4} (c) 2.32×10^{-3} (d) 1.16×10^{-4}
32. A 250.0 mL sample of a 0.20 M Cr^{3+} is electrolysed with a current of 96.5 A. If the remaining $[\text{Cr}^{3+}]$ is 0.1 M, the duration of process is :
 (a) 25 sec (b) 225 sec (c) 150 sec (d) 75 sec
33. The element indium is to be obtained by electrolysis of a molten halide of the element. Passage of a current of 3.20 A for a period of 40.0 min results in formation of 3.05 g of In. What is the oxidation state of indium in the halide melt? (Atomic mass of In 114.8)
 (a) 3 (b) 2 (c) 5 (d) 1
34. An electrolysis of a oxytungsten complex ion using 1.10 A for 40 min produces 0.838 g of tungsten. What is the charge on tungsten in the material? (Atomic mass of W 184)
 (a) 6 (b) 2 (c) 4 (d) 1
35. In the electrolysis of aqueous NaCl, what volume of $\text{Cl}_2(\text{g})$ is produced in the time that it takes to liberate 5.0 liter of $\text{H}_2(\text{g})$? Assume that both gases are measured at STP
 (a) 5.0 (b) 2.50 (c) 7.50 (d) 10.0
36. How many grams of Cr are deposited in the electrolysis of solution of $\text{Cr}(\text{NO}_3)_3$ in the same time that it takes to deposit 0.54 g of Ag in a silver coulometer arranged in series with the $\text{Cr}(\text{NO}_3)_3$ cell? (Atomic mass : Cr = 52.0; Ag = 108)
 (a) 0.0866 (b) 0.0288 (c) 0.173 (d) 0.220
37. In the electrolysis of a CuSO_4 solution, how many grams of Cu are plated out on the cathode in the time that it takes to liberate 5.6 litre of $\text{O}_2(\text{g})$, measured at 1 atm and 273 K, at the anode?
 (a) 31.75 (b) 14.2 (c) 4.32 (d) None of these
38. Ammonium perchlorate, NH_4ClO_4 , used in the solid fuel in the booster rockets on the space shuttle, is prepared from sodium perchlorate, NaClO_4 , which is produced commercially by the

electrolysis of a hot, stirred solution of sodium chloride. How many faradays are required to produce 1.0 kg of sodium perchlorate?



- (a) 40.3 (b) 18.3 (c) 31.6 (d) 65.3

39. In the commercial preparation of aluminum, aluminum oxide (Al_2O_3) is electrolysed at 1000 °C. How many coulombs of electricity are required to give 54 kg of aluminum? Assume following reaction takes place at cathode :



- (a) 17.3×10^8 (b) 3.21×10^7 (c) 1.82×10^4 (d) 57.6×10^7

40. When molten lithium chloride (LiCl) is electrolysed, lithium metal is formed at the cathode. If current efficiency is 75% then how many grams of lithium are liberated when 1930 C of charge pass through the cell ? (Atomic mass of Li = 7)

- (a) 0.105 (b) 0.120 (c) 0.28 (d) 0.240

41. Sodium metal is produced commercially by the electrolysis of molten sodium chloride and chlorine is produced as a by product. How many litres of chlorine at 1.8 atm and 27 °C will be produced if a current of 1.0×10^3 A is passed through NaCl (l) for 9.65 hr.?

- (a) 2463 (b) 460 (c) 1800 (d) 1231.6


42. $\text{H}_2(\text{g})$ and $\text{O}_2(\text{g})$, can be produced by the electrolysis of water. What total volume (in L) of O_2 and H_2 are produced at 1 atm and 273 K when a current of 30 A is passed through a K_2SO_4 (aq) solution for 193 min.?

- (a) 20.16 (b) 40.32 (c) 60.48 (d) 80.64

43. The cost of 2 Rs/kWh of operating an electric motor for 10 hours takes 10 amp at 110 V is :

- (a) 79200 Rs (b) 22000 Rs (c) 220 Rs (d) 22 Rs

44. A 1 M solution of H_2SO_4 is electrolysed. Select correct statement in respect of products obtained at anode and cathode respectively :



- (a) concentration of H_2SO_4 remain constant; H_2, O_2
 (b) concentration of H_2SO_4 increases; O_2, H_2
 (c) concentration of H_2SO_4 decreases; O_2, H_2
 (d) concentration of H_2SO_4 remains constant; $\text{S}_2\text{O}_8^{2-}, \text{H}_2$

45. Cadmium amalgam is prepared by electrolysis of a solution of CdCl_2 using a mercury cathode. How long should a current of 4 A be passed in order to prepare 10% by mass Cd in Cd—Hg amalgam on cathode of 4.5 g Hg? (atomic mass of Cd = 112)

- (a) 400 sec (b) 215.40 sec (c) 861.6 sec (d) 430.8 sec

46. Use of electrolysis is in :

- (a) electrorefining (b) electroplating
 (c) both (a) and (b) (d) None of these

47. When a solution of AgNO_3 (1 M) is electrolysed using platinum anode and copper cathode, what are the products obtained at two electrodes?



$$E_{\text{Ag}^+|\text{Ag}} \quad 0.8 \text{ volt}$$

- (a) Cu Cu^{2+} at anode; Ag^+ Ag at cathode
 (b) H_2O O_2 at anode; Cu^{2+} Cu at cathode
 (c) H_2O O_2 at anode; Ag Ag at cathode
 (d) NO_3^- NO_2 at anode; Ag Ag at cathode

48. Which of the following statements is correct about Galvanic cell ?

- (a) It converts chemical energy into electrical energy.
 (b) It converts electrical energy into chemical energy.
 (c) It converts metal from its free state to the combined state.
 (d) It converts electrolyte into individual ions.

49. E for $\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-(\text{aq})$ is 1.36 V; E for $\text{Cl}^-(\text{aq}) \rightleftharpoons \frac{1}{2}\text{Cl}_2(\text{g}) + \text{e}^-$ is :

- (a) 1.36 V (b) 1.36 V (c) 0.68 V (d) 0.68 V

50. When two half-cells of electrode potential of E_1 and E_2 are combined to form a half cell of electrode potential E_3 , then

(when n_1, n_2 and n_3 are no. of electrons exchanged in first, second and combined half-cells) :

- (a) $E_3 = E_2 - E_1$ (b) $E_3 = \frac{E_1 n_1 + E_2 n_2}{n_3}$
 (c) $E_3 = \frac{E_1 n_1 - E_2 n_2}{n_3}$ (d) $E_3 = E_1 + E_2$

51. The function of a salt bridge is to :

- (a) maintain electrical neutrality of both half cells
 (b) increase the cell potential at the positive electrode
 (c) decrease the cell potential at the negative electrode
 (d) eliminate the impurities present in the electrolyte

52. Saturated solution of KNO_3 with agar-agar is used to make 'salt bridge' because :

- (a) size of K^+ is greater than that of NO_3^-
 (b) velocity of NO_3^- is greater than that of K^+
 (c) velocities of K^+ and NO_3^- are nearly the same
 (d) both velocity and sizes of K^+ and NO_3^- ions are nearly same


53. A salt bridge may contain :

- (a) a saturated solution of KCl and agar-agar
 (b) a saturated solution of KNO_3 and agar-agar
 (c) a saturated solution of NH_4NO_3 and agar-agar
 (d) all of these

54. The nature of curve of E_{cell} vs. $\log K_c$ is :

- (a) straight line (b) parabola (c) hyperbola (d) elliptical curve

55. Consider the following equations for a cell reaction



then :


(a) $x \propto y, K_1 \propto K_2$

(b) $x \propto 2y, K_1 \propto 2K_2$





(c) $x \propto y, K_1^2 \propto K_2$

(d) $x^2 \propto y, K_1^2 \propto K_2$

56. Which graph correctly correlates E_{cell} as a function of concentrations for the cell




Y-axis : E_{cell} , X-axis : $\log_{10} \frac{[\text{Zn}^{2+}]}{[\text{Ag}^+]^2}$




57. The Nernst equation $E = E^\circ - \frac{RT}{nF} \ln Q$ indicates that the Q will be equal to equilibrium constant K_c when :

(a) $E = E^\circ$ (b) $RT/nF = 1$ (c) $E = \text{zero}$ (d) $E = 1$


58. The cell reaction $2\text{Ag}^+(\text{aq}) + \text{H}_2(\text{g}) \rightleftharpoons 2\text{H}^+(\text{aq}) + 2\text{Ag(s)}$, is best represented by :




59. The cell reaction $\text{Hg}_2\text{Cl}_2(\text{s}) + \text{Cu(s)} \rightleftharpoons \text{Cu}^{2+}(\text{aq}) + 2\text{Cl}^-(\text{aq}) + 2\text{Hg(l)}$, is best represented by :



60. The cell reaction $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14\text{H}^+(\text{aq}) + 6\text{Fe}^{2+}(\text{aq}) \rightleftharpoons 6\text{Fe}^{3+}(\text{aq}) + 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O(l)}$, is best represented by :



61. Select the correct cell reaction of the cell $\text{Ag(s)} | \text{Ag}^+(\text{aq}) || \text{Cu}^{2+}(\text{aq}) | \text{Cu(s)}$:



- (c) $2\text{Ag}(s) + \text{Cu}^{2+}(aq) \rightarrow \text{Cu}(s) + 2\text{Ag}^{+}(aq)$
 (d) $\text{Cu}^{2+}(aq) + 2\text{Ag}(s) \rightarrow 2\text{Ag}^{+}(aq) + \text{Cu}(s)$
- 62.** Select the correct cell reaction of the cell $\text{Pt}(s) | \text{Cl}_2(g) | \text{Cl}^{-}(aq) || \text{Ag}^{+}(aq) | \text{Ag}(s)$:
- (a) $\text{Cl}_2(g) + \text{Ag}(s) \rightarrow \text{Ag}(s) + 2\text{Cl}^{-}(aq)$
 (b) $\text{Cl}_2(g) + \text{Ag}(s) \rightarrow 2\text{Cl}^{-}(aq) + \text{Ag}^{+}(aq)$
 (c) $2\text{Cl}^{-}(aq) + 2\text{Ag}^{+}(aq) \rightarrow 2\text{Ag}(s) + \text{Cl}_2(g)$
 (d) $\text{AgCl}(s) \rightarrow \text{Ag}(s) + \text{Cl}^{-}(aq)$
- 63.** Standard electrode potential of SHE at 298 K is :
 (a) 0.05 V (b) 0.10 V (c) 0.50 V (d) 0.00 V
- 64.** The e.m.f. of the following galvanic cells :
 (a) $\text{Zn} | \text{Zn}^{2+}(1\text{M}) || \text{Cu}^{2+}(1\text{M}) | \text{Cu}$ (b) $\text{Zn} | \text{Zn}^{2+}(0.1\text{M}) || \text{Cu}^{2+}(1\text{M}) | \text{Cu}$
 (c) $\text{Zn} | \text{Zn}^{2+}(1\text{M}) || \text{Cu}^{2+}(0.1\text{M}) | \text{Cu}$ (d) $\text{Zn} | \text{Zn}^{2+}(0.1\text{M}) || \text{Cu}^{2+}(0.1\text{M}) | \text{Cu}$
 are represented by E_1, E_2, E_3 and E_4 respectively. Which of the following statement is true?
 (a) $E_1 > E_2 > E_3 > E_4$ (b) $E_3 > E_2 > E_1 > E_4$
 (c) $E_3 > E_1 > E_4 > E_2$ (d) $E_2 > E_1 > E_4 > E_3$
- 65.** Based on the cell notation for a spontaneous reaction, at the anode :
 $\text{Ag}(s) | \text{AgCl}(s) | \text{Cl}^{-}(aq) || \text{Br}^{-}(aq) | \text{Br}_2(l) | \text{C}(s)$
 (a) AgCl gets reduced (b) Ag gets oxidized
 (c) Br^{-} gets oxidized (d) Br_2 gets reduced
- 66.** Given the listed standard electrode potentials, what is E° for the cell :
 $4\text{BiO}^{+}(aq) + 3\text{N}_2\text{H}_5^{+}(aq) \rightarrow 4\text{Bi}(s) + 3\text{N}_2(g) + 4\text{H}_2\text{O}(l) + 7\text{H}^{+}(aq)$
 $\text{N}_2(g) + 5\text{H}^{+}(aq) + 4e^{-} \rightarrow \text{N}_2\text{H}_5^{+}(aq), E^\circ = 0.23\text{ V}$
 $\text{BiO}^{+}(aq) + 2\text{H}^{+}(aq) + 3e^{-} \rightarrow \text{Bi}(s) + \text{H}_2\text{O}(l), E^\circ = +0.32\text{ V}$
 (a) 0.55 (b) 0.34 (c) 1.88 (d) 0.09
- 67.** What is the standard electrode potential for the reduction of HClO ?
 $\text{HClO}(aq) + \text{H}^{+}(aq) + 2e^{-} \rightarrow \text{Cl}^{-}(aq) + \text{H}_2\text{O}(l)$
 Given : $\text{Cr}^{2+}(aq) \rightarrow \text{Cr}^{3+}(aq) + e^{-}, E^\circ = 0.41\text{ V}$
 $\text{HClO}(aq) + \text{H}^{+}(aq) + 2\text{Cr}^{2+}(aq) \rightarrow 2\text{Cr}^{3+}(aq) + \text{Cl}^{-}(aq) + \text{H}_2\text{O}(l), E^\circ = 1.80$
 (a) 1.39 (b) 1.54 (c) 1.22 (d) 0.90
- 68.** The E° for the following cell is 0.34 V. $\text{In}(s) | \text{In}(\text{OH})_3(aq) || \text{SbO}_2^{-}(aq) | \text{Sb}(s)$.
 Using $E^\circ = 1.0\text{ V}$ for the $\text{In}(\text{OH})_3 | \text{In}$ couple, calculate E° for the $\text{SbO}_2^{-} | \text{Sb}$ half-reaction :
 (a) 1.34 (b) 0.66
 (c) +0.82 (d) 0.66
- 69.** From the following half-cell reactions and their standard potentials, what is the smallest possible standard e.m.f. for spontaneous reactions?
 $\text{PO}_4^{3-}(aq) + 2\text{H}_2\text{O}(l) + 2e^{-} \rightarrow \text{HPO}_3^{2-}(aq) + 3\text{OH}^{-}(aq); E^\circ = 1.05\text{ V}$
 $\text{PbO}_2(s) + \text{H}_2\text{O}(l) + 2e^{-} \rightarrow \text{PbO}(s) + 2\text{OH}^{-}(aq); E^\circ = 0.28\text{ V}$

- $\text{IO}_3^-(\text{aq}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightarrow \text{IO}^-(\text{aq}) + 4\text{OH}^-(\text{aq}); E^\circ = 0.56 \text{ V}$
 (a) +0.00 (b) +0.74 (c) +0.56 (d) +0.28
70. Determine which substance is the best reducing agent in Q. no. 69 :
 (a) HPO_3^{2-} (b) PO_4^{3-} (c) IO^- (d) IO_3^-
71. Which substance is the best oxidizing agent in Q. no. 69?
 (a) IO_3^- (b) IO^- (c) PbO (d) PO_4^{3-}
72. Consider the following half-cell reactions and associated standard half-cell potentials and determine the maximum voltage that can be obtained by combination resulting in spontaneous processes :
- | | |
|---|--|
| $\text{AuBr}_4^-(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s}) + 4\text{Br}^-(\text{aq}); E^\circ = 0.86 \text{ V}$ | |
| $\text{Eu}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Eu}^{2+}(\text{aq}); E^\circ = 0.43 \text{ V}$ | |
| $\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s}); E^\circ = 0.14 \text{ V}$ | |
| $\text{IO}^-(\text{aq}) + \text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{I}^-(\text{aq}) + 2\text{OH}^-(\text{aq}); E^\circ = 0.49 \text{ V}$ | |
- (a) 0.72 (b) 1.54 (c) 1.00 (d) +1.35
73. The position of some metals in the electrochemical series in decreasing electropositive character is $\text{Mg} > \text{Al} > \text{Zn} > \text{Cu} > \text{Ag}$. What will happen if copper spoon is used to stir a solution of aluminium nitrate ?
 (a) The spoon gets coated with aluminium.
 (b) An alloy of aluminium and copper is formed.
 (c) No reaction occurs
 (d) The solution starts turning blue
74. Zn can displace :
 (a) Mg from its aqueous solution (b) Cu from its aqueous solution
 (c) Na from its aqueous solution (d) Al from its aqueous solution
75. Based on the following information arrange four metals A, B, C and D in order of decreasing ability to act as reducing agents :
 (I) Only A, B and C react with 1 M HCl to give $\text{H}_2(\text{g})$
 (II) When C is added to solutions of the other metal ions, metallic B and D are formed
 (III) Metal C does not reduce A^{n+} .
 (a) C A B D (b) C A D B (c) A C D B (d) A C B D
76. When an aqueous solution of CuSO_4 is stirred with a silver spoon then :
 (a) Cu will be formed (b) Ag will be formed
 (c) Cu^{2+} will be deposited (d) None of these
77. Based on the following information arrange four metals, A, B, C and D in order of increasing ability to act as reducing agents :
 (I) Only C reacts with 1 M HCl to give $\text{H}_2(\text{g})$
 (II) When A is added to solution of the other metal ions, metallic D is formed but not B or C
 (a) D A C B (b) A D C B (c) B D A C (d) D A B C
78. In the reaction :
 $4\text{Fe}(\text{s}) + 3\text{O}_2(\text{g}) \rightarrow 4\text{Fe}^{3+}(\text{aq}) + 6\text{O}^{2-}(\text{aq})$


which of the following statement is incorrect ?

- (a) It is a redox reaction (b) Fe is reducing agent
(c) O_2 is an oxidizing agent (d) Fe is reduced to Fe^{3+}

79. Which of the following is displaced by Fe ?

- (a) Ag (b) Zn (c) Na (d) All of these

80. The standard potential at $25^\circ C$ for the following half reactions is given :




When Zinc dust is added to the solution of $MgCl_2$.

- (a) $ZnCl_2$ is formed (b) Mg is precipitated
(c) Zn dissolved in the solution (d) No reaction takes place

81. The element which can displace three other halogens from their compound is :

- (a) F (b) Cl (c) Br (d) I

82. Using the standard half-cell potential listed, calculate the equilibrium constant for the reaction :



- (a) 2.3×10^9 (b) 4.8×10^4 (c) 4.8×10^7 (d) 4.8×10^{11}

83. The E° at $25^\circ C$ for the following reaction is 0.22 V . Calculate the equilibrium constant at $25^\circ C$:



- (a) 2.8×10^7 (b) 5.2×10^8 (c) 5.2×10^6 (d) 5.2×10^3

84. Electrode potential of the half cell $Pt(s) | Hg(l) | Hg_2Cl_2(s) | Cl^-(aq)$ can be increased by :

- (a) Increasing $[Cl^-]$ (b) Decreasing $[Cl^-]$
(c) Increasing $Hg_2Cl_2(s)$ (d) Decreasing $Hg(l)$

85. The equilibrium constant for the following general reaction is 10^{30} . Calculate E° for the cell at 298 K .



- (a) $+0.105 \text{ V}$ (b) $+0.2955 \text{ V}$ (c) 0.0985 V (d) 0.2955 V

86. A solution containing H^+ and D^+ ions is in equilibrium with a mixture of H_2 and D_2 gases at $25^\circ C$. If the partial pressures of both gases are 1.0 atm , find the ratio of $[D^+]/[H^+]$:

(Given : $E^\circ_{D^+/D_2} = 0.003 \text{ V}$)


- (a) 1.23 (b) 1.12 (c) 0.11 (d) 1.0


87. The E° at $25^\circ C$ for the following reaction is 0.55 V . Calculate the ΔG° in kJ/mol :



- (a) 637 (b) 424 (c) 106 (d) 318.5

88. Use the following standard electrode potentials, calculate ΔG° in kJ/mol for the indicated reaction :





- (a) 9.65 (b) 24.3 (c) 48.25 (d) 35.2

89. Consider an electrochemical cell in which the following reaction occurs and predict which changes will decrease the cell voltage :



- (I) decrease the $[\text{Ag}^+]$ (II) increase in $[\text{Fe}^{3+}]$ (III) increase the amount of Ag

- (a) I (b) II and III (c) II (d) I and II

90. Consider the following equation for an electrochemical cell reaction. Which of the following changes in condition will increase the cell voltage?



- (I) addition of concentrated HClO_4 in the cell solution

- (II) increase the pressure of $\text{H}_2(g)$ (III) increase the amount of $\text{Pb}(s)$


- (a) III (b) I and II (c) II and III (d) II

91. The standard electrode potential for the following reaction is +1.33 V. What is the potential at pH 2.0?



- (a) +1.820 V (b) +1.990 V (c) +1.608 V (d) +1.0542 V

92. The standard electrode potential for the following reaction is 0.57 V. What is potential at pH = 12.0?




- (a) 0.17 V (b) 0.21 V (c) 0.39 V (d) +1.95 V

93. $\text{Co}|\text{Co}^{2+}(\text{C}_2)||\text{Co}^{2+}(\text{C}_1)|\text{Co}$; for this cell, G is negative if :

- (a) $\text{C}_2 < \text{C}_1$ (b) $\text{C}_1 > \text{C}_2$ (c) $\text{C}_1 = \text{C}_2$ (d) unpredictable

94. What will be the emf for the given cell?



- (a) $\frac{RT}{F} \ln \frac{P_1}{P_2}$ (b) $\frac{RT}{2F} \ln \frac{P_1}{P_2}$ (c) $\frac{RT}{F} \ln \frac{P_2}{P_1}$ (d) None of these


95. For the electrochemical cell $\text{Pt}(s)|\text{H}_2(g)|\text{H}^+(1M)||\text{Cu}^{2+}(1M)|\text{Cu}(s)$, which one of the following statements is true ?

- (a) H^+ ions are formed at anode and Cu is deposited at cathode.
 (b) H_2 liberated at cathode and Cu is deposited at anode.
 (c) Oxidation occurs at cathode.
 (d) Reduction occurs at anode.

96. In a concentration cell the same reagents are present in both the anode and the cathode compartments, but at different concentrations. Calculate the emf of a cell containing 0.040 M Cr^{3+} in one compartment and 1.0 M Cr^{3+} in the other if Cr electrodes are used in both.

- (a) 0.028 V (b) 0.249 V (c) 0.083 V (d) 0.125 V

97. A 1.0 M solution of Cd^{2+} is added to excess iron and the system is allowed to reach equilibrium. What is the concentration of Cd^{2+} ?



- (a) 0.195 (b) 0.097 (c) 0.053 (d) 0.145
- 98.** The measured voltage for the reaction with the indicated concentrations is 1.50 V. Calculate E .
- $$\text{Cr(s)} + 3\text{Ag}^+(\text{aq}, 0.10\text{ M}) \rightleftharpoons 3\text{Ag(s)} + \text{Cr}^{3+}(\text{aq}, 0.30\text{ M})$$
- (a) 1.35 (b) 1.40 (c) 1.65 (d) 1.55
- 99.** Calculate the standard voltage that can be obtained from an ethane oxygen fuel cell at 25 °C.
- $$\text{C}_2\text{H}_6(\text{g}) + 7/2\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 3\text{H}_2\text{O}(\text{l}); \Delta G^\circ = -1467\text{ kJ/mol}$$
- (a) +0.91 (b) +0.54 (c) +0.72 (d) +1.08
- 100.** $\text{I}_2(\text{s})|\text{I}^-(0.1\text{M})$ half cell is connected to a $\text{H}^+(\text{aq})|\text{H}_2(1\text{ bar})|\text{Pt}$ half cell and e.m.f. is found to be 0.7714 V. If $E_{\text{I}_2/\text{I}^-} = 0.535\text{ V}$, find the pH of $\text{H}^+|\text{H}_2$ half-cell.
- (a) 1 (b) 3 (c) 5 (d) 7
- 101.** What is the SRP for $\text{S}^{2-}|\text{CuS}|\text{Cu}$ half-cell?
- Given: $K_{\text{sp}}(\text{CuS}) = 10^{-35}$ and $E_{\text{Cu}^{2+}/\text{Cu}} = 0.34\text{ V}$
- (a) 1.034 V (b) 1.0 V (c) 0.694 V (d) 0.694 V
- 102.** Given the following standard electrode potentials, the K_{sp} for PbBr_2 is:
- $$\text{PbBr}_2(\text{s}) \rightleftharpoons \text{Pb}^{2+}(\text{aq}) + 2\text{Br}^-(\text{aq}); E^\circ = 0.248\text{ V}$$
- $$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Pb(s)}; E^\circ = 0.126\text{ V}$$
- (a) 7.4×10^{-5} (b) 4.9×10^{-14} (c) 5.2×10^{-6} (d) 2.3×10^{-13}
- 103.** The standard free energy change for the following reaction is -210 kJ/mol. What is the standard cell potential?
- $$2\text{H}_2\text{O}_2(\text{aq}) \rightarrow 2\text{H}_2\text{O(l)} + \text{O}_2(\text{g})$$
- (a) +0.752 (b) +1.09 (c) +0.420 (d) +0.640
- 104.** At equilibrium:
- (a) $E_{\text{cell}}^\circ = 0, \Delta G^\circ = 0$ (b) $E_{\text{cell}} = 0, \Delta G = 0$
- (c) both are correct (d) none is correct
- 105.** The E° at 25 °C for the following reaction at the indicated concentrations is 1.50 V. Calculate the ΔG° in kJ/mol at 25 °C:
- $$\text{Cr(s)} + 3\text{Ag}^+(\text{aq}, 0.1\text{ M}) \rightleftharpoons 3\text{Ag(s)} + \text{Cr}^{3+}(\text{aq}, 0.1\text{ M})$$
- (a) 140.94 (b) 295 (c) 212 (d) 422.83
- 106.** If $E_{\text{Au}^+/ \text{Au}}^\circ$ is 1.69 V and $E_{\text{Au}^{3+}/ \text{Au}}^\circ$ is 1.40 V, then $E_{\text{Au}^+ / \text{Au}^{3+}}^\circ$ will be:
- (a) 0.19 V (b) 2.945 V (c) 1.255 V (d) None of these
- 107.** Consider the following standard electrode potentials and calculate the equilibrium constant at 25 °C for the indicated disproportionation reaction:
- $$3\text{Mn}^{2+}(\text{aq}) \rightleftharpoons \text{Mn(s)} + 2\text{Mn}^{3+}(\text{aq})$$
- $$\text{Mn}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq}); E^\circ = 1.51\text{ V}$$
- $$\text{Mn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mn(s)}; E^\circ = 1.185\text{ V}$$
- (a) 1.2×10^{-43} (b) 2.4×10^{-73} (c) 6.3×10^{-92} (d) 1.5×10^{-62}

- 108.** A galvanic cell is composed of two hydrogen electrodes, one of which is a standard one. In which of the following solutions should the other electrode be immersed to get maximum e.m.f. :
 (a) 0.1 M HCl (b) 0.1 M H₂SO₄ (c) 0.1 M NH₄OH (d) 0.01 M HCOOH
- 109.** Ag|AgCl|Cl⁻(C₂)||Cl⁻(C₁)|AgCl|Ag for this cell G is negative if :
 (a) C₁ < C₂ (b) C₁ > C₂ (c) C₂ < C₁ (d) Both (a) and (c)
- 110.** By how much is the oxidizing power of Cr₂O₇²⁻|Cr³⁺ couple decreased if the H⁺ concentration is decreased from 1 M to 10⁻³ M at 25 °C?
 (a) 0.001 V (b) 0.207 V (c) 0.441 V (d) 0.414 V
- 111.** The temperature coefficient of a cell whose operation is based on the reaction

$$\text{Pb(s)} + \text{HgCl}_2(\text{aq}) \rightleftharpoons \text{PbCl}_2(\text{aq}) + \text{Hg(l)}$$
 is :

$$\frac{dE}{dT}_p = 1.5 \times 10^{-4} \text{ V K}^{-1} \text{ at } 298 \text{ K}$$

 The change in entropy (in J/K mol) during the operation is :
 (a) 8627 (b) 57.9 (c) 28.95 (d) 14.475
- 112.** Thermodynamic efficiency of a cell is given by :
 (a) $\frac{H}{G}$ (b) $\frac{nFE}{G}$ (c) $\frac{nEF}{H}$ (d) nFE°
- 113.** Calculate the value of equilibrium constant (K_f) for the reaction :

$$\text{Zn}^{2+}(\text{aq}) + 4\text{OH}^{-}(\text{aq}) \rightleftharpoons \text{Zn(OH)}_4^{2-}(\text{aq})$$

 Given :

$$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^{-} \rightleftharpoons \text{Zn(s)}; E^\circ = 0.76 \text{ V}$$

$$\text{Zn(OH)}_4^{2-}(\text{aq}) + 2\text{e}^{-} \rightleftharpoons \text{Zn(s)} + 4\text{OH}^{-}(\text{aq}); E^\circ = 1.36 \text{ V}$$


$$2.303 \frac{RT}{F} = 0.06$$

 (a) 10¹⁰ (b) 2 × 10¹⁰ (c) 10²⁰ (d) none of these
- 114.** Which of the following statement is false for fuel cells ?
 (a) They are more efficient (b) They are free from pollution
 (c) They run till reactants are active (d) Fuel burned with O₂
- 115.** When a lead storage battery is charged it acts as :
 (a) a fuel cell (b) an electrolytic cell
 (c) a galvanic cell (d) a concentration cell
- 116.** The metal that forms a self-protecting film of oxide to prevent corrosion is :
 (a) Na (b) Al (c) Cu (d) Au
- 117.** Rusting of iron is catalyzed by which of the following ?
 (a) Fe (b) Zn (c) O₂ (d) H
- 118.** Which of the following is a highly corrosive salt ?
 (a) Hg₂Cl₂ (b) HgCl₂ (c) FeCl₂ (d) PbCl₂
- 119.** The Zn acts as sacrificial or cathodic protection to prevent rusting of iron because :
 (a) E°_{OP} of Zn < E°_{OP} of Fe (b) E°_{OP} of Zn > E°_{OP} of Fe
 (c) E°_{OP} of Zn < E°_{OP} of Fe (d) Zn is cheaper than iron





- 120.** In electrochemical corrosion of metals, the metal undergoing corrosion :
 (a) acts as anode (b) acts as cathode
 (c) undergoes reduction (d) None
- 121.** When an acid cell is charged, then :
 (a) voltage of cell increases (b) resistance of cell increases
 (c) electrolyte of cell dilutes (d) None of these
- 122.** Electrolytic conduction is due to the movement of :
 (a) electrons (b) ions
 (c) atoms (d) electrons as well as ions
- 123.** Molten sodium chloride conducts electricity due to the presence of :
 (a) free electron (b) free ions
 (c) free molecules (d) atoms of sodium and chlorine
- 124.** Pure water does not conduct electricity because it :
 (a) is neutral (b) is readily decomposed
 (c) is almost totally unionized (d) has a low boiling point
- 125.** The relation among conductance (G), specific conductance (κ) and cell constant (l/A) is :
 (a) $G = \frac{l}{A} \kappa$ (b) $G = \frac{A}{l} \kappa$ (c) $G = \frac{l}{A} \kappa$ (d) $G = A l \kappa$
- 126.** If x is specific resistance (in $\Omega^{-1}\text{cm}$) of the electrolyte solution and y is the molarity of the solution, then κ (in $\text{S cm}^2\text{mol}^{-1}$) is given by :
 (a) $\frac{1000x}{y}$ (b) $1000 \frac{y}{x}$ (c) $\frac{1000}{xy}$ (d) $\frac{xy}{1000}$
- 127.** Equivalent conductivity can be expressed in terms of specific conductance (κ) and concentration (N) in gram equivalent dm^{-3} as :
 (a) κN (b) $\frac{1000}{N}$ (c) $\frac{N}{1000}$ (d) $N \cdot 1000$
- 128.** Resistance of a decimolar solution between two electrodes 0.02 meter apart and 0.0004 m^2 in area was found to be 50 ohm. Specific conductance (κ) is :
 (a) 0.1 S m^{-1} (b) 1 S m^{-1} (c) 10 S m^{-1} (d) $4 \cdot 10^{-4} \text{ S m}^{-1}$
- 129.** Resistance of 0.1 M KCl solution in a conductance cell is 300 ohm and conductivity is 0.013 S cm^{-1} . The value of cell constant is :
 (a) 3.9 cm^{-1} (b) 39 m^{-1} (c) 3.9 m^{-1} (d) None of these
- 130.** Dissociation constant of a weak acid (HA) in terms of κ_m and C_m is :
 (a) $K_a = \frac{C_m}{(\kappa_m)^2}$ (b) $K_a = \frac{C_m^2}{\kappa_m(\kappa_m)}$
 (c) $K_a = \frac{C(\kappa_m)^2}{\kappa_m(\kappa_m)}$ (d) None of these
- 131.** When a concentrated solution of an electrolyte is diluted
 (a) its specific conductance increases
 (b) its equivalent conductivity decreases
 (c) its specific conductivity decreases and equivalent conductivity increases
 (d) both specific and equivalent conductivity increase

- 132.** Molar conductivity of a solution of an electrolyte AB_3 is $150 \text{ Scm}^2\text{mol}^{-1}$. If it ionises as $AB_3 \rightarrow A^3 + 3B^-$, its equivalent conductivity will be :
 (a) 150 (in $\text{Scm}^2\text{eq}^{-1}$) (b) 75 (in $\text{Scm}^2\text{eq}^{-1}$)
 (c) 50 (in $\text{Scm}^2\text{eq}^{-1}$) (d) 80 (in $\text{Scm}^2\text{eq}^{-1}$)
- 133.** Equivalent conductivity of $\text{Fe}_2(\text{SO}_4)_3$ is related to molar conductivity by the expression :
 (a) $\lambda_{\text{eq}} = \lambda_m$ (b) $\lambda_{\text{eq}} = \lambda_m/3$ (c) $\lambda_{\text{eq}} = 3 \lambda_m$ (d) $\lambda_{\text{eq}} = \lambda_m/6$
- 134.** The limiting equivalent conductivity of NaCl, KCl and KBr are 126.5, 150.0 and $151.5 \text{ Scm}^2\text{eq}^{-1}$, respectively. The limiting equivalent ionic conductivity for Br^- is $78 \text{ Scm}^2\text{eq}^{-1}$. The limiting equivalent ionic conductivity for Na^+ ions would be :
 (a) 128 (b) 125 (c) 49 (d) 50
- 135.** The specific conductance of a saturated solution of silver bromide is Scm^{-1} . The limiting ionic conductivity of Ag^+ and Br^- ions are x and y , respectively. The solubility of silver bromide in g L^{-1} is : (molar mass of $\text{AgBr} = 188$)
 (a) $\frac{1000}{x+y}$ (b) $\frac{1000}{x-y}$ (c) $\frac{1000}{x+y} \cdot 188$ (d) $\frac{x-y}{188} \cdot 1000$
- 136.** The resistance of 0.1 N solution of formic acid is 200 ohm and cell constant is 2.0 cm^{-1} . The equivalent conductivity (in $\text{Scm}^2\text{eq}^{-1}$) of 0.1 N formic acid is :
 (a) 100 (b) 10 (c) 1 (d) none of these
- 137.** A conductance cell was filled with a 0.02 M KCl solution which has a specific conductance of $2.768 \times 10^{-3} \text{ ohm}^{-1} \text{ cm}^{-1}$. If its resistance is 82.4 ohm at 25°C , the cell constant is :
 (a) 0.2182 cm^{-1} (b) 0.2281 cm^{-1} (c) 0.2821 cm^{-1} (d) 0.2381 cm^{-1}
- 138.** The ionic conductivity of Ba^{2+} and Cl^- at infinite dilution are 127 and $76 \text{ ohm}^{-1} \text{ cm}^2\text{eq}^{-1}$ respectively. The equivalent conductivity of BaCl_2 at infinity dilution (in $\text{ohm}^{-1} \text{ cm}^2\text{eq}^{-1}$) would be :
 (a) 203 (b) 279 (c) 101.5 (d) 139.5
- 139.** Unit of ionic mobility is :
 (a) $\text{m}^2 \text{V}^{-1} \text{s}^{-1}$ (b) $\text{m}^2 \text{V}^{-2} \text{s}^{-1}$ (c) $\text{m}^2 \text{V}^{-1} \text{s}^{-1}$ (d) $\text{m}^2 \text{Vs}^{-1}$
- 140.** λ_{AgCl} can be obtained :
 (a) by extrapolation of the graph λ_{obs} and $\sqrt{C}$ to zero concentration
 (b) by known values of λ_{AgNO_3} , λ_{HCl} and λ_{HNO_3}
 (c) both (a) and (b)
 (d) None of these
- 141.** The conductance of a salt solution (AB) measured by two parallel electrodes of area 100 cm^2 separated by 10 cm was found to be 0.0001 Scm^{-1} . If volume enclosed between two electrode contain 0.1 mole of salt, what is the molar conductivity ($\text{Scm}^2\text{mol}^{-1}$) of salt at same concentration :
 (a) 10 (b) 0.1 (c) 1 (d) none of these





- 142.** The conductivity of a strong electrolyte :
 (a) increases on dilution (b) decreases on dilution
 (c) does not change with dilution (d) depends upon density of electrolytes
- 143.** The increase in equivalent conductivity of a weak electrolyte with dilution is due to :
 (a) increase in degree of dissociation and decrease in ionic mobility
 (b) decrease in degree of dissociation and decrease in ionic mobility
 (c) increase in degree of dissociation and increase in ionic mobility
 (d) decrease in degree of dissociation and increase in ionic mobility
- 144.** Strong electrolytes are those which :
 (a) conduct electricity (b) dissolve readily in water
 (c) dissociate into ions at high dilution (d) completely dissociate into ions
- 145.** The electric conduction of a salt solution in water depends on the :
 (a) size of its molecules (b) shape of its molecules
 (c) size of solvent molecules (d) extent of its ionization
- 146.** A graph was plotted between molar conductivity of various electrolytes (NaCl, HCl and NH_4OH) and $\sqrt{C}$ (in mol L^{-1}). Correct set is :







- (a) I(NaCl), II(HCl), III(NH_4OH)
 (b) I(HCl), II(NaCl), III(NH_4OH)
 (c) I(NH_4OH), II(NaCl), III(HCl)
 (d) I(NH_4OH), II(HCl), III(NaCl)
- 147.** Which of the following is arranged in increasing order of ionic mobility ?
 (a) I Br Cl F (b) F Cl Br I
 (c) F I Cl Br (d) F Cl I Br
- 148.** $\text{HNO}_3(\text{aq})$ is titrated with $\text{NaOH}(\text{aq})$ conductometrically, graphical representation of the titration is :



149. Which of the following plots will be obtained for a conductometric titration of strong acid against a weak base?



150. Conductometric titration curve of an equimolar mixture of HCl and HCN with NaOH(aq) is :





Level 2

1. In the Hall process, aluminium is produced by the electrolysis of molten Al_2O_3 . How many second would it take to produce enough aluminum by the Hall process to make a case of 24 cans of aluminum soft-drink, if each can uses 5.0 g of Al, a current of 9650 amp is employed, and the current efficiency of the cell is 90.0% :

(a) 203.2 (b) 148.14 (c) 333 (d) 6.17

2. 108 g fairly concentrated solution of AgNO_3 is electrolysed by using 0.1 F charge. The mass of resulting solution is :

(a) 94 g (b) 11.6 g (c) 96.4 g (d) None of these

3. The electrolysis of acetate solution produces ethane according to reaction :



The current efficiency of the process is 80%. What volume of gases would be produced at 27 °C and 740 torr, if the current of 0.5 amp is passed through the solution for 96.5 min?

(a) 6.0 L (b) 0.60 L (c) 1.365 L (d) 0.91 L

4. A layer of chromium metal 0.25 mm thick is to be plated on an auto bumper with a total area of 0.32 m^2 from a solution containing CrO_4^{2-} ? What current flow is required for this electroplating if the bumper is to be plated in 60 s ? The density of chromium metal is 7.20 g/cm^3 .

(a) $4.9 \times 10^3 \text{ A}$ (b) $1.78 \times 10^3 \text{ A}$ (c) $5.3 \times 10^4 \text{ A}$ (d) $10.69 \times 10^6 \text{ A}$

5. 100 mL of 0.05 M $\text{CuSO}_4(\text{aq})$ solution was electrolysed using inert electrodes by passing current till the pH of the resulting solution was 2. The solution after electrolysis was neutralized and then treated with excess KI and formed I_2 titrated with 0.04 M $\text{Na}_2\text{S}_2\text{O}_3$. Calculate the required volume (in mL) of $\text{Na}_2\text{S}_2\text{O}_3$:

(a) 112.5 mL (b) 100 mL (c) 125 mL (d) None of these

6. If the equilibrium constant for the reaction $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightleftharpoons \text{H}_2\text{O}(\text{l})$ is 10^{13} at certain temperature then what is the E° for the reaction, $2\text{H}_2\text{O}(\text{l}) + 2e^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$

Given : $\frac{2.303 RT}{F} = 0.066$

(a) 1.230 V (b) 0.858 V (c) 0.80 V (d) 0.8274 V

7. A fuel cell develops an electrical potential from the combustion of butane at 1 bar and 298 K



What is E° of a cell?

(a) 4.74 V (b) 0.547 V (c) 4.37 V (d) 1.09 V


8. The cell $\text{Pt}|\text{H}_2(\text{g}, 0.1 \text{ bar})|\text{H}^+(\text{aq}), \text{pH} = X||\text{Cl}^-(\text{aq}, 1 \text{ M})|\text{Hg}_2\text{Cl}_2|\text{Hg}|\text{Pt}$,

has e.m.f. of 0.5755 V at 25 °C. The SOP of calomel electrode is 0.28V, then pH of solution will be :

(a) 11 (b) 4.5 (c) 5.5 (d) None of these

9. For a cell reaction $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{l})$, $\Delta_r S_{298} = 0.32 \text{ kJ/K}$. What is the value of $\Delta_f H_{298}(\text{H}_2\text{O}, \text{l})$?
- Given : $\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}(\text{l}); E^\circ = 1.23 \text{ V}$
- (a) 285.07 kJ/mol (b) 570.14 kJ/mol
(c) 285.07 kJ/mol (d) None of these
10. What is the potential of an electrode which originally contained 0.1 M NO_3^- and 0.4 M H^+ and which has been treated by 80% of the cadmium necessary to reduce all the NO_3^- to $\text{NO}(\text{g})$ at 1 bar?
- Given : $\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^- \rightarrow \text{NO} + 2\text{H}_2\text{O}; E^\circ = 0.96 \text{ V}; \log 2 = 0.3$
- (a) 0.84 V (b) 1.08 V
(c) 1.23 V (d) 1.36 V
11. The standard reduction potential of normal calomel electrode and reduction potential of saturated calomel electrodes are 0.27 and 0.33 volt respectively. What is the concentration of Cl^- in saturated solution of KCl?
- (a) 0.1 M (b) 0.01 M (c) 0.001 (d) None
12. Determine the potential of the following cell :
- $\text{Pt} | \text{H}_2(\text{g}, 0.1 \text{ bar}) | \text{H}^+(\text{aq}, 10^{-3} \text{ M}) || \text{MnO}_4^-(\text{aq}, 0.1 \text{ M}), \text{Mn}^{2+}(\text{aq}, 0.01 \text{ M}), \text{H}^+(\text{aq}, 0.01 \text{ M}) | \text{Pt}$
- Given : $E^\circ_{\text{MnO}_4^- | \text{Mn}^{2+}} = 1.51 \text{ V}$
- (a) 1.54 V (b) 1.48 V (c) 1.84 V (d) none of these
13. Copper reduces NO_3^- into NO and NO_2 depending upon concentration of HNO_3 in solution. Assuming $[\text{Cu}^{2+}] = 0.1 \text{ M}$, and $P_{\text{NO}} = P_{\text{NO}_2} = 10^{-3} \text{ bar}$, at which concentration of HNO_3 , thermodynamic tendency for reduction of NO_3^- into NO and NO_2 by copper is same?
- [Given : $E^\circ_{\text{Cu}^{2+} | \text{Cu}} = 0.34 \text{ volt}$, $E^\circ_{\text{NO}_3^- | \text{NO}} = 0.96 \text{ volt}$, $E^\circ_{\text{NO}_3^- | \text{NO}_2} = 0.79 \text{ volt}$]
- (a) $10^{1.23} \text{ M}$ (b) $10^{0.56} \text{ M}$ (c) $10^{0.66} \text{ M}$ (d) $10^{0.12} \text{ M}$
14. For the cell, $\text{Pt} | \text{Cl}_2(\text{g}, 0.4 \text{ bar}) | \text{Cl}^-(\text{aq}, 0.1 \text{ M}) || \text{Cl}^-(\text{aq}, 0.01 \text{ M}) | \text{Cl}_2(\text{g}, 0.2 \text{ bar}) | \text{Pt}$
- The measured potential at 298 K is:
- (a) 0.051 V (b) 0.051 V (c) 0.102 V (d) 0.0255 V
15. The chlorate ion can disproportionate in basic solution according to reaction,
- $$2\text{ClO}_3^- \rightleftharpoons \text{ClO}_2^- + \text{ClO}_4^-$$
- What is the equilibrium concentration of perchlorate ions from a solution initially at 0.1 M in chlorate ions at 298 K?
- Given : $E^\circ_{\text{ClO}_4^- | \text{ClO}_3^-} = 0.36 \text{ V}$ and $E^\circ_{\text{ClO}_3^- | \text{ClO}_2^-} = 0.33 \text{ V}$ at 298 K
- (a) 0.019 M (b) 0.024 M (c) 0.1 M (d) 0.19 M

16. A cell diagram shown below contains one litre of buffer solution of $\text{HA}(\text{p}K_a = 4)$ and NaA in both compartments. What is the cell e.m.f.?



- (a) 0.03 V (b) 0.06 V (c) 0.06 V (d) None of these
17. Given the cell : $\text{Cd(s)} | \text{Cd(OH)}_2(\text{s}) | \text{NaOH(aq, 0.01 M)} | \text{H}_2(\text{g, 1 bar}) | \text{Pt(s)}$ with $E_{\text{cell}} = 0.0 \text{ V}$. If $E_{\text{Cd}^{2+}|\text{Cd}} = 0.39 \text{ V}$, then K_{sp} of Cd(OH)_2 is :
 (a) 0.1 (b) 10^{-13} (c) 10^{-15} (d) None of these
18. Calculate the e.m.f. (in V) of the cell :


$$\text{Pt} | \text{H}_2(\text{g}) | \text{BOH(aq)} || \text{HA(aq)} | \text{H}_2(\text{g}) | \text{Pt};$$

$$\begin{array}{ccccccc} & & 0.1 \text{ bar} & 1 \text{ M} & & 0.1 \text{ M} & 1 \text{ bar} \end{array}$$
 Given : $K_a(\text{HA}) = 10^{-7}$, $K_b(\text{BOH}) = 10^{-6}$
 (a) 0.39 V (b) 0.36 V (c) 0.93 V (d) None of these
19. Calculate the potential of a half cell having reaction : $\text{Ag}_2\text{S(s)} + 2\text{e}^- \rightleftharpoons 2\text{Ag(s)} + \text{S}^{2-}(\text{aq})$ in a solution buffered at $\text{pH} = 3$ and which is also saturated with $0.1 \text{ M H}_2\text{S(aq)}$:
 [Given: $K_{\text{sp}}(\text{Ag}_2\text{S}) = 10^{-49}$, $K_{\text{a1}} = 10^{-7}$, $K_{\text{a2}} = 10^{-14}$]
 (a) 1.18 (b) 0.19 (c) 0.19 V (d) none of these
20. The conductivity of 0.1 N NaOH solution is 0.022 S cm^{-1} . When equal volume of 0.1 N HCl solution is added, the conductivity of resultant solution is decreased to 0.0055 S cm^{-1} . The equivalent conductivity in $\text{Scm}^2\text{equivalent}^{-1}$ of NaCl solution is :
 (a) 0.0055 (b) 0.11 (c) 110 (d) None of these
21. In above question after formation of NaCl , further 0.1 N HCl is added, the volume of which is double to that of the first portion added, the conductivity increases to 0.018 S cm^{-1} . The value of $\kappa_{\text{eq}}(\text{HCl})$ is [assume no change in equivalent conductivity of NaCl (aq)]:
 (a) $330 \text{ S cm}^2\text{eq}^{-1}$ (b) $305 \text{ S cm}^2\text{eq}^{-1}$
 (c) $415 \text{ S cm}^2\text{eq}^{-1}$ (d) $360 \text{ S cm}^2\text{eq}^{-1}$
22. Given the following molar conductivities at infinite dilution (Λ_m^∞), HCl , $426 \text{ cm}^2\text{mol}^{-1}$; NaCl , $126 \text{ cm}^2\text{mol}^{-1}$; NaC (sodium crotonate), $83 \text{ cm}^2\text{mol}^{-1}$. What is the dissociation constant of crotonic acid, if the conductivity of a 0.001 M crotonic acid solution is $3.83 \times 10^{-5} \text{ S cm}^{-1}$?
 (a) 10^{-5} (b) 1.11×10^{-5} (c) 1.11×10^{-4} (d) 0.01

23. Equivalent conductivity of BaCl_2 , H_2SO_4 and HCl , are x_1 , x_2 and $x_3 \text{ Scm}^{-1}\text{eq}^{-1}$ at infinite dilution. If conductivity of saturated BaSO_4 solution is $x \text{ Scm}^{-1}$, then K_{sp} of BaSO_4 is :
- (a) $\frac{500x}{(x_1 \quad x_2 \quad 2x_3)}$ (b) $\frac{10^6 x^2}{(x_1 \quad x_2 \quad 2x_3)^3}$
 (c) $\frac{2.5 \quad 10^5 x^2}{(x_1 \quad x_2 \quad x_3)^2}$ (d) $\frac{0.25x^2}{(x_1 \quad x_2 \quad x_3)^2}$
24. The conductivity of $0.001 \text{ M Na}_2\text{SO}_4$ solution is $2.6 \times 10^{-4} \text{ Scm}^{-1}$ and increases to $7.0 \times 10^{-4} \text{ Scm}^{-1}$, when the solution is saturated with CaSO_4 . The molar conductivities of Na^+ and Ca^{2+} are 50 and $120 \text{ Scm}^2\text{mol}^{-1}$, respectively. Neglect conductivity of used water. What is the solubility product of CaSO_4 ?
- (a) 4×10^{-6} (b) 1.57×10^{-3}
 (c) 4×10^{-4} (d) 2.46×10^{-6}
25. The dissociation constant of a weak acid is 1.6×10^{-5} and the molar conductivity at infinite dilution is $380 \times 10^{-4} \text{ S m}^2\text{mol}^{-1}$. If the cell constant is 0.01 m^{-1} then conductance of 0.01 M acid solution is :
- (a) $1.52 \times 10^{-5} \text{ S}$ (b) 1.52 S
 (c) $1.52 \times 10^{-3} \text{ S}$ (d) $1.52 \times 10^{-4} \text{ S}$
26. Three electrolytic cells X, Y, Z containing solution of 1 M each NaCl , AgNO_3 and CuSO_4 respectively are connected in series combination. During electrolysis 21.6 gm of silver deposits at cathode in cell Y . Which is incorrect statement.
- (a) 6.35 gm copper deposits at cathode in cell Z
 (b) 2.24 litre Cl_2 is liberated (at 1 atm and 273 K) at anode in cell X
 (c) 2.24 litre O_2 is liberated (at 1 atm and 273 K) at anode in cell Y
 (d) 2.24 litre H_2 is liberated (at 1 atm and 273 K) at cathode in cell X
27. During electrolysis of $\text{H}_2\text{SO}_4(aq)$ with high charge density, $\text{H}_2\text{S}_2\text{O}_8$ formed as by product. In such electrolysis $22.4 \text{ L H}_2(g)$ and $8.4 \text{ L O}_2(g)$ liberated at 1 atm and 273 K at electrode. The moles of $\text{H}_2\text{S}_2\text{O}_8$ formed is :
- (a) 0.25 (b) 0.50 (c) 0.75 (d) 1.00
28. $\text{Zn}(s) | \text{Zn}(\text{CN})_4^{2-} (0.5 \text{ M}), \text{CN}^- (0.01) || \text{Cu}(\text{NH}_3)_4^{2+} (0.5 \text{ M}), \text{NH}_3 (1 \text{ M}) | \text{Cu}(s)$
 Given: K_f of $\text{Zn}(\text{CN})_4^{2-} \quad 10^{16}$, K_f of $\text{Cu}(\text{NH}_3)_4^{2+} \quad 10^{12}$,
 $E^\circ_{\text{Zn}| \text{Zn}^{2+}} \quad 0.76 \text{ V}$; $E^\circ_{\text{Cu}^{2+}|\text{Cu}} \quad 0.34 \text{ V}$, $\frac{2.303RT}{F} \quad 0.06$
 The emf of above cell is :
- (a) 1.22 V (b) 1.10 V (c) 0.98 V (d) None of these

Level 3

Passage 1



A Galvanic cell consist of three compartment as shown in figure. The first compartment contain $\text{ZnSO}_4(1M)$ and III compartment contain $\text{CuSO}_4(1M)$. The mid compartment contain $\text{NaNO}_3(1M)$. Each. compartment contain 1L solution :

$$E_{\text{Zn}^{2+}/\text{Zn}} = 0.76; E_{\text{Cu}^{2+}/\text{Cu}} = 0.34$$

- The concentration of Zn^{2+} in first compartment after passage of 0.1 F charge will be :
 (a) 1 M (b) 1.05 M (c) 1.025 M (d) 0.5 M
- The concentration of NO_3^- in mid compartment after passage of 0.1 F of charge will be :
 (a) 0.95 M (b) 0.90 M (c) 0.975 M (d) 1.05 M
- The concentration of SO_4^{2-} ion in III compartment after passage of 0.1 F of charge will be :
 (a) 1.05 M (b) 1.025 M (c) 0.95 M (d) 0.975 M

Passage 2

The cell potential (E_{cell}) of a reaction is related as $G = -nF E_{\text{cell}}$, where G represents max. useful electrical work

n = no. of moles of electrons exchanged during the reaction

for reversible cell reaction $dG = \left(\left(\frac{\partial G}{\partial T} \right)_p - nF E_{\text{cell}} \right) dT + \left(\left(\frac{\partial G}{\partial p} \right)_T - nF E_{\text{cell}} \right) dp + \sum \mu_i dn_i$

at constant pressure

$\therefore$ At constant pressure

$$\left(\frac{\partial G}{\partial T} \right)_p = -nF E_{\text{cell}} \quad \dots(1)$$

$$\left(\frac{\partial G}{\partial T} \right)_p = -nF E_{\text{cell}} \quad \dots(2)$$

$\frac{dE_{\text{cell}}}{dT}_p$ is known as temperature coefficient of the e.m.f of the cell

- The temperature coefficient of the e.m.f. of cell, $\frac{dE}{dT}_p$ is given by :
 (a) $\frac{nF}{S}$ (b) $\frac{S}{nF}$ (c) $\frac{S}{nFT}$ (d) nFE
- At 300 K, ΔH for the reaction
 $\text{Zn(s)} + \text{AgCl(s)} \rightarrow \text{ZnCl}_2(\text{aq}) + 2\text{Ag(s)}$ is
 218 kJ/mol while the e.m.f. of the cell was 1.015 V. $\frac{dE}{dT}_p$ of the cell is :
 (a) $4.2 \times 10^{-4} \text{VK}^{-1}$ (b) $3.81 \times 10^{-4} \text{VK}^{-1}$
 (c) 0.11VK^{-1} (d) $7.62 \times 10^{-4} \text{VK}^{-1}$
- Calculate ΔS for the given cell reaction in Q. no. 2 :
 (a) 73.53 J/K mol (b) 83.53 J/K mol (c) 100 J/K mol (d) none of these



Passage 3

Molar conductivity (κ_m) is defined as conducting power of the ions produced by 1 mole of an electrolyte in a solution. $\kappa_m = \frac{\kappa}{C}$ where κ is conductivity (in S cm^{-1}) and C is molar concentration (in mol/cm^3)


The molar conductivity of 0.04 M solution of MgCl_2 is $200 \text{ S cm}^2 \text{ mol}^{-1}$ at 298 K. A cell with electrodes that are 2.0 cm^2 in surface area and 0.50 cm apart is filled with MgCl_2 solution.

- Conductance of MgCl_2 solution is :
 (a) $8 \times 10^{-3} \text{ S}$ (b) 32 S (c) 0.032 S (d) None of these
- How much current will flow when the potential difference between the two electrodes is 5.0 V?
 (a) 156.25 A (b) 0.16 A (c) 160 A (d) None of these



Passage 4

In a hydrogen oxygen fuel cell, electricity is produced. In this process $\text{H}_2(\text{g})$ is oxidised at anode and $\text{O}_2(\text{g})$ reduced at cathode.



4.48 litre H_2 at 1 atm and 273 K oxidised in 9650 sec.

- The current produced is (in amp) :
 (a) 1 A (b) 2 A (c) 4 A (d) 8 A
- The mass of water produced is :
 (a) 7.2 g (b) 3.6 g (c) 1.8 g (d) 0.9 g
- If current produced in fuel cell, is used for the deposition of Cu^{2+} in 1 L, 2 M $\text{CuSO}_4(\text{aq})$ solution for 241.25 sec using Pt electrode, the pH of solution after electrolysis is :
 (a) 1 (b) 2 (c) 3 (d) 4



Passage 5

A saturated solution in $\text{Agx}(K_{sp} = 3 \times 10^{-12})$ and $\text{Agy}(K_{sp} = 10^{-12})$ has conductivity $0.4 \times 10^{-6} \text{ S cm}^{-1}$.

Given : Limiting molar conductivity of Ag^+ $60 \text{ S cm}^2 \text{ mol}^{-1}$

Limiting molar conductivity of x $90 \text{ S cm}^2 \text{ mol}^{-1}$

- The conductivity of y is (in S cm^{-1}) :
 (a) 1.45×10^{-7} (b) 1.45×10^{-5}
 (c) 1.45×10^{-9} (d) None of these
- The limiting molar conductivity of y is (in $\text{S cm}^2 \text{ mol}^{-1}$) :
 (a) 290 (b) 2900 (c) 2.90 (d) None of these

One or More Answers is/are Correct



- If the e.m.f of a galvanic cell is negative, it implies that :
 (a) the cell reaction is spontaneous
 (b) the cell reaction is non-spontaneous
 (c) the cell reaction is exothermic
 (d) the cell is working in reverse direction
- Select correct statement(s) about electrolysis :
 (a) Electric current is used to drive a non-spontaneous reaction
 (b) G is positive for chemical process during electrolysis
 (c) Cations and anions are moved toward the anode and cathode respectively
 (d) Over voltage is generally associated with evolution of O_2 gas
- If the half-cell reaction $A + e^- \rightleftharpoons A^-$ has a large negative reduction potentials, it follows that :
 (a) A is readily reduced (b) A is readily oxidised
 (c) A^- is readily reduced (d) A^- is readily oxidised
- Which of the following statement is correct?
 If $E_{\text{Cu}^{2+}|\text{Cu}} = 0.34 \text{ V}$ and $E_{\text{Sn}^{2+}|\text{Sn}} = 0.136 \text{ V}$, $E_{\text{H}^+|\text{H}_2} = 0.0 \text{ V}$
 (a) Cu^{2+} ions can be reduced by $\text{H}_2(\text{g})$ (b) Cu can be oxidized by H^+
 (c) Sn^{2+} ions can be reduced by H_2 (d) Sn can be oxidized by Cu^{2+}
- The oxidation potential of hydrogen half-cell will be negative if :
 (a) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 1 \text{ M}$ (b) $p(\text{H}_2) = 1 \text{ atm}$ and $[\text{H}^+] = 2 \text{ M}$
 (c) $p(\text{H}_2) = 0.2 \text{ atm}$ and $[\text{H}^+] = 1 \text{ M}$ (d) $p(\text{H}_2) = 0.2 \text{ atm}$ and $[\text{H}^+] = 0.2 \text{ M}$
- Which of the following arrangement will produce oxygen at anode during electrolysis?
 (a) Dilute H_2SO_4 with Pt electrodes
 (b) Fused NaOH with inert electrodes


- (c) Dilute H_2SO_4 with Cu electrodes
 (d) Concentrated aq. NaCl with Pt electrodes
7. When an aqueous concentrated solution of lithium chloride is electrolysed using inert electrodes :
- (a) Cl_2 is liberated at the anode
 (b) Li is deposited at the cathode
 (c) as the current flows, pH of the solution around the cathode remains constant
 (d) as the current flows, pH of the solution around the cathode increases
8. Oxygen and hydrogen gas are produced at the anode and cathode during the electrolysis of fairly concentrated aqueous solution of :
- (a) K_2SO_4 (b) AgNO_3 (c) H_2SO_4 (d) NaOH
9. During the purification of copper by electrolysis :
- (a) the anode used is made of copper ore
 (b) pure copper is deposited on the cathode
 (c) the impurities such as Ag, Au present in solution as ions
 (d) concentration of CuSO_4 solution remains constant during dissolution of Cu
10. When a lead storage battery is discharged :
- (a) SO_2 is evolved
 (b) lead sulphate is produced at both electrodes
 (c) sulphuric acid is consumed
 (d) water is formed
11. Which of the following is characteristic of the cathode in a voltaic cell?
- (a) It may gain weight during reaction
 (b) Electrons flow to it through the external circuit
 (c) It is where oxidation occurs
 (d) It receives electrons from ions in solution
12. In an electrochemical process, a salt bridge is used :
- (a) to maintain electrical neutrality in each solution
 (b) to complete the external circuit so that current can flow for long time
 (c) to mix the solution of anodic and cathodic compartment
 (d) to supply voltage
13. For a reaction in a galvanic cell the value of ΔG at certain temperature is not necessarily equal to :
- (a) nFE (b) $RT \ln K$ (c) $T \Delta S$ (d) zero
14. Standard electrode potential of two half-reactions are given below :
- $$\text{Fe}^{2+} \rightleftharpoons \text{Fe} \quad ; \quad E^\circ = 0.44 \text{ V}$$
- $$\text{Fe}^{3+} \rightleftharpoons \text{Fe}^{2+} \quad ; \quad E^\circ = 0.77 \text{ V}$$

If Fe^{2+} , Fe^{3+} and Fe are kept together :

- (a) the concentration of Fe^{3+} increases (b) the concentration of Fe^{3+} decreases
 (c) the mass of Fe increases (d) the concentration of Fe^{2+} decreases

15. Which of the following statements are correct regarding to galvanic cell?
- A reaction is spontaneous from left to right if $E_{\text{cell}} > 0$
 - A reaction occurs from right to left if $E_{\text{cell}} < 0$
 - If the system is at equilibrium no net reaction occurs
 - E_{cell} is temperature-independent
16. Which of the following are concentration cells?
- $\text{Pt} | \text{H}_2(\text{g}) | \text{HCl} | \text{H}_2(\text{g}) | \text{Pt}$
 $P_1 \qquad P_2$
 - $\text{Cd}, (\text{Hg}) | \text{Cd}^{2+} | (\text{Hg}), \text{Cd}$
 $a_1 \qquad a_2$
 - $\text{Zn}(\text{s}) | \text{Zn}^{2+} || \text{Cu}^{2+} | \text{Cu}$
 $c_1 \qquad c_2$
 - $\text{Ag} | \text{AgCl} | \text{Cl}^-(\text{aq}) || \text{Br}^-(\text{aq}) | \text{AgBr} | \text{Ag}$
 $c_1 \qquad c_2$
17. In electrolyte concentration cell :
- the electrode material and the solution in both half-cells are composed of the same substances
 - only the concentrations of solutions of the same substances is different
 - $E_{\text{cell}} = 0$
 - the Nernst equation reduces to $E_{\text{cell}} = \frac{0.0591}{n} \log Q$ at 25 °C
18. The standard electrode potential of a metal-metal ion ($\text{Ag} | \text{Ag}^+$) and metal-sparingly soluble salt anion ($\text{Ag} | \text{AgCl} | \text{Cl}^-$) are related as :
- $E_{\text{Ag}^+ | \text{Ag}} = E_{\text{Cl}^- | \text{AgCl} | \text{Ag}} - \frac{RT}{F} \ln K_{\text{sp}}$
 - $E_{\text{Cl}^- | \text{AgCl} | \text{Ag}} = E_{\text{Ag}^+ | \text{Ag}} - \frac{RT}{F} \ln K_{\text{sp}}$
 - $E_{\text{Cl}^- | \text{AgCl} | \text{Ag}} = E_{\text{Ag}^+ | \text{Ag}} - \frac{RT}{F} \ln \frac{[\text{Cl}^-]}{K_{\text{sp}}}$
 - $E_{\text{Cl}^- | \text{AgCl} | \text{Ag}} = E_{\text{Ag}^+ | \text{Ag}} - \frac{RT}{F} \ln \frac{K_{\text{sp}}}{[\text{Cl}^-]}$
19. Which of the following units is correctly matched?
- SI units of conductivity is S m^{-1}
 - SI units of molar conductivity is $\text{S cm}^2 \text{mol}^{-1}$
 - SI unit of conductance is S^{-1}
 - All of these
20. Which of the following statements is/are correct?
- The conductance of one cm^3 (or 1 unit³) of a solution is called specific conductance
 - Specific conductance increases while molar conductivity decreases on progressive dilution
 - The limiting equivalent conductivity of weak electrolyte cannot be determined exactly by extrapolation of the plot of κ_{eq} against $\sqrt{c}$
 - The conductance of metals is due to the movement of free electrons
21. Which is/are correct statement ?
- No corrosion takes place in vacuum
 - Corrosion is protected by electroplating
 - During rusting $\text{Fe}_2\text{O}_3 \cdot x\text{H}_2\text{O}$ formed
 - In presence of electrolyte, corrosion takes place with greater rate
22. A dilute solution of KCl was placed between two Pt electrodes 10 cm apart across which a potential difference of 10 volt was applied. Which is/are correct statement (Given : molar conductivity of K⁺ at infinite dilution is $96.5 \text{ S cm}^2 \text{mol}^{-1}$.)

- (a) Ionic mobility of K^+ is $10^{-3} \text{ cm}^2 \text{ sec}^{-1} \text{ volt}^{-1}$.
 (b) The speed of K^+ is $10^{-3} \text{ cm sec}^{-1}$.
 (c) Distance travelled by K^+ in $5 \times 10^3 \text{ sec}$ is 5 cm.
 (d) The potential gradient is 1.0 volt cm^{-1} .



$$pK_b(\text{NH}_4\text{OH}) = 5; pK_a(\text{CH}_3\text{COOH}) = 5; \frac{2.303RT}{F} = 0.06$$

Volume of $0.1M \text{NH}_4\text{OH}$ in anode half cell = 100 mL,
 Volume of $0.1M \text{CH}_3\text{COOH}$ in cathode half cell = 100 mL

Which is/are correct statement ?

- (a) The emf of given cell is 0.48 V.
 (b) The emf of given cell is 0.36 V when 50 mL, $0.1M \text{NaOH}$ added to cathode compartment
 (c) The emf of given cell is 0.36 V when 50 mL $0.1M \text{HCl}$ added to anode compartment
 (d) The emf of given cell is 0.192 V when 100 mL $0.1M \text{NaOH}$ added to anode compartment
24. Given : $\text{Pt}(s) \mid \underset{P_1 \text{ atm}}{\text{Cl}_2(g)} \mid \text{Cl}^-(C_1) \parallel \text{Cl}^-(C_2) \mid \underset{P_2 \text{ atm}}{\text{Cl}_2(g)} \mid \text{Pt}(s)$

Identify in which of following condition working of cell takes place :

- (a) $C_1 = C_2$ and $P_1 = P_2$ (b) $P_2 = P_1$ and $C_1 = C_2$
 (c) $C_1 = C_2$ and $P_1 = P_2$ (d) $P_2 = P_1$ and $C_1 = C_2$
25. 1000 mL $1M \text{CuSO}_4(aq)$ is electrolysed by 9.65 A current for 100 sec using Pt-electrode. Which is/are correct statement ?
- (a) Blue colour intensity decreases during electrolysis.
 (b) Blue colour intensity remains constant if Cu-electrode used.
 (c) pH of solution is 8 after electrolysis.
 (d) 28 mL of CH_4 at 1 atm and 273 K required for its combustion by O_2 , liberated during electrolysis.

Match the Column

Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II

1.

Column-I

- (A) Dilute solution of HCl
 (B) Dilute solution of NaCl
 (C) Concentrated solution of NaCl
 (D) Fairly concentrated solution of AgNO_3

Column-II

- (P) O_2 evolved at anode
 (Q) H_2 evolved at cathode
 (R) Cl_2 evolved at anode
 (S) Ag deposition at cathode

2.	Column-I	Column-II
	(A) If SOP of substance lies between 1.23 to 0.81 V (B) If SOP of substance lies between 0.81 V to -0.40 V (C) If SOP is less than -1.23 V (D) If SOP is greater than 0.40 V	(P) Oxidation of substance is not possible (Q) Oxidation possible only in acidic medium (R) Oxidation possible in any medium (S) Oxidation easily takes place
3.	Column-I	Column-II (SRP)
	(A) $F_2 + 2e \rightleftharpoons 2F^-$ (B) $Cl_2 + 2e \rightleftharpoons 2Cl^-$ (C) $Br_2 + 2e \rightleftharpoons 2Br^-$ (D) $I_2 + 2e \rightleftharpoons 2I^-$	(P) 0.54 (Q) 1.09 (R) 1.36 (S) 2.87
4.	Column-I	Column-II
	(A) $Pt Fe^{3+}, Fe^{2+} $ (B) $Pt H_2 H^+$ (C) $Pt Hg Hg^{2+} $ (D) $Pb PbSO_4 SO_4^{2-} $	(P) Metal-metal ion half-cell (Q) Gas-gas ion half-cell (R) Oxidation-reduction half-cell (S) Metal-sparingly soluble salt half-cell
5.	Column-I (Property)	Column-II (Unit)
	(A) Conductance (B) Conductivity (C) Molar conductivity (D) Resistivity	(P) Sm^{-1} (Q) $S^{-1}m$ (R) Sm^2mol^{-1} (S) S
6.	Column-I (Ion)	Column-II (Molar Conductivity)
	(A) H^+ (B) Na^+ (C) Li (D) Cs	(P) 350 (Q) 50 (R) 39 (S) 77
7.	Column-I	Column-II
	(A) Galvanic cell (B) Electrolytic cell	(P) Used in space craft (Q) No transformation of electrical energy into chemical energy

(C) Dead battery

(D) Fuel cell

(R) Cell reaction is spontaneous

(S) Cell reaction is non-spontaneous

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).
Examine the statements carefully and mark the correct answer according to the instructions given below :

- (A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** E_{cell} is negative for electrolytic cell.
STATEMENT-2 : G is ve for electrolytic cell.
2. **STATEMENT-1 :** When 2 faraday of charge is passed through $0.1 \text{ M H}_2\text{SO}_4(aq)$, 11.2 litre O_2 evolved at STP
STATEMENT-2 : Molecular mass of oxygen is 32.
3. **STATEMENT-1 :** Copper is dissolved at anode and deposited at cathode when Cu electrodes are used and electrolyte is $1 \text{ M CuSO}_4(aq)$ solution.
STATEMENT-2 : SOP of Cu is less than SOP of water and SRP of Cu is greater than SRP of water.
4. **STATEMENT-1 :** 1 coulomb charge deposits 1 g-equivalent of a substance.
STATEMENT-2 : 1 faraday is charge on 1 mole of electrons.
5. **STATEMENT-1 :** If SRP of substance is -0.3 V , its reduction is possible at cathode.
STATEMENT-2 : Reduction potential of water lies between 0 to -0.8274 V at 25°C .
6. **STATEMENT-1 :** If SRP of substance is -0.5 V then reduction of substance is possible only in basic medium.
STATEMENT-2 : SRP of water is -0.8274 V and at reduction potential is zero at $\text{pH} = 7$
7. **STATEMENT-1 :** The voltage of mercury cell remains constant for longer period of time.
STATEMENT-2 : It is because net cell reaction does not involve ions.
8. **STATEMENT-1 :** Lead storage battery is a galvanic cell without salt bridge.
STATEMENT-2 : A secondary cell is rechargeable cell.
9. **STATEMENT-1 :** The SRP of three metallic ions A, B, C are $-0.3, -0.5, 0.8 \text{ volt}$ respectively, so oxidizing power of ions is $C > A > B$.
STATEMENT-2 : Higher the SRP, higher the oxidizing power.
10. **STATEMENT-1 :** If SOP of substance is less than -1.23 V and over voltage $> 0 \text{ V}$, then its oxidation in its aqueous solution is not possible at 298 K .
STATEMENT-2 : Standard reduction potential (SRP) of water is -1.23 V .
11. **STATEMENT-1 :** We cannot add the electrode potential in order to get electrode potential of third electrode if no. of moles of electrons exchanged are not same.

STATEMENT-2 : Electrode potential is an extensive property.

12. STATEMENT-1 : $E_{\text{cell}} = 0$ for a chloride ion concentration cell.

STATEMENT-2 : For this concentration cell where $E_{\text{cell}} = \frac{RT}{nF} \ln \frac{[\text{Cl}^-]_{\text{LHS}}}{[\text{Cl}^-]_{\text{RHS}}}$

13. STATEMENT-1 : If $\frac{dE_{\text{cell}}}{dT} = 0$ for a cell reaction then S is positive.

STATEMENT-2 : $S = nFT \frac{dE}{dT}$

14. STATEMENT-1 : Molar conductivity increases with decrease in concentration for weak electrolytes.

STATEMENT-2 : No. of ions increases and no. of ions per unit volume decreases due to dilution.

15. STATEMENT-1 : Conductivity decreases with the decreases in concentration both the weak and strong electrolytes.

STATEMENT-2 : No. of ions per unit volume linearly decreases in both electrolytes.

Subjective Problems



- How many faradays are required for reduction of 1 mol $\text{C}_6\text{H}_5\text{NO}_2$ into $\text{C}_6\text{H}_5\text{NH}_2$?
- What is the equivalent mass of O_2 in the following reaction, $\text{H}_2\text{O} + \frac{1}{2}\text{O}_2 + 2e^- \rightarrow 2\text{OH}^-$?
- The amount of electricity which releases 2.0 g of gold from a gold salt is same as that which dissolves 0.967 g of copper anode during the electrolysis of copper sulphate solution. What is the oxidation number of gold in the gold ion ? (At. mass of Cu = 63.5; Au = 197)
- When a molten salt was electrolysed for 5 min with 9.65 A current, 0.18 g of the metal was deposited. Calculate the Eq. mass of metal.
- During the electrolysis of a concentrated brine solution, calculate the moles of chlorine gas produced by the passage of 4F electricity.
- Calculate the cell potential (in V) if $G = 96.5 \text{ kJ/mol}$ and $n = 1$.
- If K_c for the reaction $\text{Cu}^{2+}(\text{aq}) + \text{Sn}^{2+}(\text{aq}) \rightleftharpoons \text{Sn}^{4+}(\text{aq}) + \text{Cu}(\text{s})$ at 25 °C is represented as 2.6×10^y then find the value of y .
(Given : $E_{\text{Cu}^{2+}|\text{Cu}} = 0.34 \text{ V}$; $E_{\text{Sn}^{4+}|\text{Sn}^{2+}} = 0.15 \text{ V}$)
- If G for the half cell $\text{MnO}_4^-|\text{MnO}_2$ in an acid solution is xF then find the value of x .
(Given : $E_{\text{MnO}_4^-|\text{Mn}^{2+}} = 1.5 \text{ V}$; $E_{\text{MnO}_2|\text{Mn}^{2+}} = 1.25 \text{ V}$)
- If the equilibrium constant for the reaction $\text{Cd}^{2+}(\text{aq}) + 4\text{NH}_3(\text{aq}) \rightleftharpoons \text{Cd}(\text{NH}_3)_4^{2+}(\text{aq})$ is 10^x then find the value of x .
(Given : $E_{\text{Cd}^{2+}|\text{Cd}} = 0.4 \text{ V}$; $E_{\text{Cd}(\text{NH}_3)_4^{2+}|\text{Cd}} = 0.61 \text{ V}$)

10. At what pH oxidation potential of water is 0.81 V ?
11. The resistance of a conductivity cell containing 0.001 M KCl solution at 298 K is $1500\ \Omega$. What is the cell constant (in mm^{-1}) if the conductivity of 0.001 M KCl solution is $2 \times 10^{-3}\text{ S cm}^{-1}$.
12. Molar conductivities at infinite dilution of KCl, HCl and CH_3COOK are 0.013 , 0.038 and $0.009\text{ S m}^2\text{ mol}^{-1}$ respectively at 291 K . If conductivity of 0.001 M CH_3COOH is $2.72 \times 10^{-3}\text{ S m}^{-1}$ then find % degree of dissociation of CH_3COOH .
13. A solution containing $1\text{ M XSO}_4(\text{aq})$ and $1\text{ M YSO}_4(\text{aq})$ is electrolysed. If conc. of X^{2+} is 10^{-z} M when deposition of Y^{2+} and X^{2+} starts simultaneously, calculate the value of Z .
 Given : $\frac{2.303RT}{F} = 0.06$
 $E^\circ_{\text{X}^{2+}|\text{X}} = 0.12\text{ V}$; $E^\circ_{\text{Y}^{2+}|\text{Y}} = 0.24\text{ V}$
14. Molar conductivity of aqueous solution of HA is $200\text{ S cm}^2\text{ mol}^{-1}$, pH of this solution is 4. Calculate the value of $pK_a(\text{HA})$ at 25°C .
 Given : $\lambda_m(\text{NaA}) = 100\text{ S cm}^2\text{ mol}^{-1}$; $\lambda_m(\text{HCl}) = 425\text{ S cm}^2\text{ mol}^{-1}$;
 $\lambda_m(\text{NaCl}) = 125\text{ S cm}^2\text{ mol}^{-1}$
15. The standard reduction potential of a silver chloride electrode (metal-sparingly soluble salt electrode) is 0.209 V and for silver electrode is 0.80 V . If the moles of AgCl that can dissolve in 10 L of a 0.01 M NaCl solution is represented as 10^{-Z} then find the value of Z .

ANSWERS

Level 1

1. (a)	2. (c)	3. (d)	4. (d)	5. (a)	6. (c)	7. (d)	8. (d)	9. (a)	10. (c)
11. (a)	12. (b)	13. (c)	14. (b)	15. (d)	16. (c)	17. (a)	18. (a)	19. (d)	20. (c)
21. (d)	22. (c)	23. (b)	24. (c)	25. (c)	26. (d)	27. (d)	28. (b)	29. (c)	30. (d)
31. (d)	32. (d)	33. (a)	34. (a)	35. (a)	36. (a)	37. (a)	38. (d)	39. (d)	40. (a)
41. (a)	42. (c)	43. (d)	44. (b)	45. (b)	46. (c)	47. (c)	48. (a)	49. (b)	50. (b)
51. (a)	52. (c)	53. (d)	54. (a)	55. (c)	56. (b)	57. (c)	58. (b)	59. (c)	60. (a)
61. (c)	62. (c)	63. (d)	64. (d)	65. (b)	66. (a)	67. (a)	68. (d)	69. (d)	70. (a)
71. (a)	72. (d)	73. (c)	74. (b)	75. (d)	76. (d)	77. (d)	78. (d)	79. (a)	80. (d)
81. (a)	82. (a)	83. (a)	84. (a)	85. (b)	86. (b)	87. (a)	88. (c)	89. (d)	90. (d)
91. (d)	92. (c)	93. (b)	94. (b)	95. (a)	96. (a)	97. (c)	98. (d)	99. (d)	100. (b)
101. (c)	102. (a)	103. (b)	104. (b)	105. (d)	106. (d)	107. (c)	108. (c)	109. (c)	110. (d)
111. (c)	112. (c)	113. (c)	114. (d)	115. (b)	116. (b)	117. (d)	118. (b)	119. (b)	120. (a)
121. (a)	122. (b)	123. (b)	124. (c)	125. (b)	126. (c)	127. (b)	128. (b)	129. (a)	130. (b)
131. (c)	132. (c)	133. (d)	134. (d)	135. (c)	136. (a)	137. (b)	138. (a)	139. (c)	140. (b)
141. (b)	142. (b)	143. (c)	144. (d)	145. (d)	146. (b)	147. (b)	148. (a)	149. (c)	150. (d)

Level 2

1. (b)	2. (c)	3. (d)	4. (d)	5. (a)	6. (b)	7. (d)	8. (c)	9. (a)	10. (a)
11. (a)	12. (b)	13. (c)	14. (a)	15. (a)	16. (b)	17. (c)	18. (a)	19. (c)	20. (c)
21. (b)	22. (b)	23. (c)	24. (a)	25. (b)	26. (c)	27. (a)	28. (c)		



Level 3

Passage-1: 1. (c) 2. (a) 3. (d)

Passage-2: 1. (b) 2. (b) 3. (a)

Passage-3: 1. (c) 2. (b)

Passage-4: 1. (c) 2. (b) 3. (b)

Passage-5: 1. (a) 2. (a)

One or More Answers is/are correct

1. (b,d) 2. (a,b,d) 3. (d) 4. (a,d) 5. (b,c) 6. (a,b) 7. (a,d) 8. (a, c, d)
 9. (a,b,d) 10. (b,c,d) 11. (a,b) 12. (a) 13. (b,d) 14. (b) 15. (a,b,c) 16. (a,b,d)
 17. (a,b,c,d) 18. (b) 19. (a) 20. (a,c,d) 21. (a,b,c,d) 22. (a,b,c,d) 23. (a,b,c) 24. (a,b)
 25. (a,b,d)

Match the Column

1. A P, Q; B P, Q; C Q, R; D P, S
 2. A Q; B R; C P; D S, R
 3. A S; B R; C Q; D P
 4. A R; B Q; C P; D S
 5. A S; B P; C R; D Q
 6. A P; B Q; C R; D S
 7. A Q, R; B S; C Q; D P, Q, R

Assertion-Reason Type Questions

1. (A) 2. (B) 3. (C) 4. (D) 5. (A) 6. (C) 7. (A) 8. (B) 9. (A) 10. (C)
 11. (C) 12. (B) 13. (C) 14. (A) 15. (C)

Subjective Problems

1. 6	2. 8	3. 3	4. 6	5. 2	6. 1	7. 6	8. 5	9. 7	10. 7
11. 3	12. 8	13. 4	14. 4	15. 7					

Hints and Solutions

Level 1

1. (a) For spontaneous
 $\Delta_r G = -nFE$
 $E > 0$
2. (c) Cathode and anode are () ve terminals in electrolytic and galvanic cell.
3. (d) Electrostatic attraction is broken down by polar solvent (water).
4. (d) $\frac{A}{1 \text{ mole } e^-} = \frac{A}{1 \text{ mole}}$
 Charge of 1 mole electron is required for deposition of 1 equivalent of substance.
5. (a) According to stoichiometry liberated amount of ion depends upon $\therefore w = \frac{EIt}{F}$
 (i) Electrolysis time (ii) Current strength
 (ii) Electrochemical equivalent
6. (c) $1 F = 96500 \text{ C}$,
 $1 \text{ C} = \frac{6.023 \times 10^{23}}{96500} = 6.24 \times 10^{18}$
7. (d) Charge $0.01 \times 10^3 = 1000$ 60 60 36
8. (d) $3.21 \times 10^6 = 500 t$
 $t = \frac{3.21 \times 10^6}{500 \times 60} = 107 \text{ min.}$
9. (a) $W = ZIt$
 $0.746 = \frac{108}{1 \times F} \times 548 \times I$ $I = 1.22$
10. (c) $\frac{0.55}{M} \times 3 = \frac{0.55 \times 100 \times 60}{96500}$
 $M = 48.25 \text{ g/mol}$
11. (a) $w = \frac{9}{2} \times \frac{5}{96500} \times \frac{1 \times 60 \times 60}{195} = 0.840 \text{ g}$
12. (b) $5.2 \times \frac{52}{3} \times \frac{9.65}{96500} \times t \text{ (sec.)}$,
 $t(\text{sec}) = 3000$
 $t(\text{min}) = 50$
13. (c) $\frac{W}{E} = \frac{I \times t}{96500}$; $\frac{0.195}{195} = 4 \times \frac{5 \times 60 \times 60 \times I}{96500}$,
 $I = 21.44$

14. (b) $0.25 \times \frac{93}{5} \times \frac{Q}{F}$
 $Q = 1.3 \times 10^2$
15. (d) When 1 mole of e^- is passed, mass of substance deposited
 $\frac{1 \times 6.023 \times 10^{23}}{1.81 \times 10^{23}} = 3.34 \text{ g eq. wt}$
 atomic mass of metal = $3.34 \times 3 = 10.03 \text{ g}$
16. (c) 1 mole Mn_3O_4 lose $6 \times \frac{8}{3} = 16$ 3 10 mole
 e^- ; so total charge required $2 \times 10 = 20 F$
17. (a) $2\text{H}^+ + 2e^- \rightarrow \text{H}_2$
18. (a) $\therefore E_{\text{Mg}^{2+}|\text{Mg}}^\circ > E_{\text{Cr}^{3+}|\text{Cr}}^\circ > E_{\text{Cu}^{2+}|\text{Cu}}^\circ > E_{\text{Ag}^+|\text{Ag}}^\circ$
19. (d) $\therefore E_{\text{K}^+|\text{K}}^\circ < E_{\text{H}^+|\text{H}_2}^\circ < E_{\text{H}_2\text{O}|\text{O}_2}^\circ < E_{\text{SO}_4^{2-}|\text{S}_2\text{O}_8^{2-}}^\circ$
 (High concentrate solution)
20. (c) $\therefore E_{\text{Na}^+|\text{Na}}^\circ < E_{\text{H}^+|\text{H}_2}^\circ < E_{\text{Cu}^{2+}|\text{Cu}}^\circ < E_{\text{H}^+|\text{H}_2}^\circ$
21. (d) $E_{\text{SO}_4^{2-}|\text{S}_2\text{O}_8^{2-}}^\circ < E_{\text{H}_2\text{O}|\text{O}_2}^\circ$
22. (c) $E_{\text{Na}^+|\text{Na}}^\circ < E_{\text{H}^+|\text{H}_2}^\circ < E_{\text{SO}_4^{2-}|\text{S}_2\text{O}_8^{2-}}^\circ < E_{\text{H}_2\text{O}|\text{O}_2}^\circ$
23. (b) In electrolysis of concentrated Cl^- , $\text{Cl}_2(\text{g})$ is evolved in place of $\text{O}_2(\text{g})$.
25. (c) $E_{\text{Cu}^{2+}|\text{Cu}}^\circ < E_{\text{H}_2\text{O}|\text{H}_2}^\circ < E_{\text{SO}_4^{2-}|\text{S}_2\text{O}_8^{2-}}^\circ < E_{\text{H}_2\text{O}|\text{O}_2}^\circ$
26. (d) Anode: $2\text{Cl}^-(\text{aq}) \rightarrow \text{Cl}_2(\text{g}) + 2e^-$
 Cathode: $2\text{H}_2\text{O} + 2e^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$
 Net reaction
 $2\text{Na}(\text{aq}) + 2\text{Cl}^-(\text{aq}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow \text{Cl}_2(\text{g}) + \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq}) + 2\text{Na}^+(\text{aq})$
27. (d) $\therefore E_{\text{K}^+|\text{K}}^\circ < E_{\text{H}_2\text{O}|\text{H}_2}^\circ < E_{\text{Na}^+|\text{Na}}^\circ < E_{\text{H}_2\text{O}|\text{O}_2}^\circ$;
 $E_{\text{Cu}^{2+}|\text{Cu}}^\circ < E_{\text{H}_2\text{O}|\text{H}_2}^\circ$
28. (b) Let t sec be used; No. of faradays $\frac{4 \times t}{96500}$
 No. of moles of $\text{H}_2\text{O} = 4$, no. of equivalents of $\text{H}_2\text{O} = 4 \times 2$
 $\frac{4 \times t}{96500} = 4 \times 2$
 $t = 1.93 \times 10^5$

29. (c) only Au^{3+} , Ag^+ and Cu^{2+} will deposit at cathode.

Li will not deposit at cathode because SRP of water is 0.8274 V so after Cu^{2+} ; H_2 will evolve at cathode.

30. (d) $\frac{W}{M} = 2 \times \frac{4.6}{96500} \times \frac{30}{60}$

moles of Sn^{2+} reduced = 0.043

Initial moles of Sn^{2+} = 0.5 - 0.6 = 0.30;

remaining moles of Sn^{2+} = 0.30 - 0.043

$$[\text{Sn}^{2+}] = \frac{0.257}{0.5} = 0.514 \text{ M}$$

31. (d) No. of moles of

$$\text{Ag} = \frac{15}{96,500} \times \frac{60}{100} \times \frac{1.25}{10^3} \times \frac{1}{1}$$

$$0.0116 \times 10^3$$

$$[\text{Ag}] = \frac{1.16 \times 10^5}{1000} = 1.16 \times 10^{-4}$$

32. (d) Initial moles of Cr^{3+} = 0.25 - 0.2 = 0.05;
final moles of Cr^{3+} = 0.25 - 0.1 = 0.025
moles of Cr^{3+} reduced,

$$0.05 - 0.025 = 0.025;$$

$$\text{eq. of } \text{Cr}^{3+} \text{ reduced, } 0.025 \times 3 = \frac{t}{96500} \times \frac{96.5}{1000};$$

$$t = 75 \text{ sec}$$

33. (a) $3.05 = \frac{114.8}{n} \times \frac{3.2}{96,500} \times \frac{40}{60}$

34. (a) $\frac{W}{M} = n \times \frac{I}{96500} \times \frac{t}{60} \times \frac{0.838}{184} \times \frac{1.1}{1.1}$

$$n = 6$$

35. (a) No. of equivalent of H_2 = No. of equivalent of Cl_2

$$\frac{5}{22.4} \times \frac{1}{2} = \frac{V_{\text{Cl}_2}}{22.4} \times 2$$

$$V_{\text{Cl}_2(\text{g})} = 5 \text{ L}$$

36. (a) No. of equivalent of Ag = $\frac{0.54}{108} \times 1$ = No. of

$$\text{equivalent of Cr} = \frac{W_{\text{Cr}}}{52} \times 3$$

$$W_{\text{Cr}} = \frac{0.54}{108} \times \frac{52}{3} = 0.0866$$

37. (a) No. of equivalent of Cu = $\frac{5.6}{22.4} \times \frac{1}{2} \times 4$

$$\frac{W_{\text{Cu}}}{63.5} \times 2 = W_{\text{Cu}} \times \frac{31.75}{63.5}$$

38. (d) no. of eq = $\frac{1000}{122.5} \times 8 = 65.3$

39. (d) $54 \times 10^3 \times \frac{27}{3} \times \frac{x}{96,500}$

$$x = 57.9 \times 10^7$$

40. (a) $\frac{W}{7} = \frac{1930}{96500} \times \frac{0.75}{1000}; W = 0.105 \text{ gm}$

41. (a) equivalents of Cl_2 produced = $\frac{1000}{96500} \times \frac{9.65}{3600} \times 3600 = 360$

$$\text{moles of } \text{Cl}_2 = \frac{180}{2463 \text{ L}}$$

$$\text{so } V = \frac{nRT}{P} = \frac{180}{1.8} \times \frac{0.0821}{1} \times 300$$

42. (c) No. of equivalent of H_2 produced = Eq. of O_2

$$\frac{I}{96500} \times \frac{t}{60} = \frac{30}{96500} \times \frac{193}{60}$$

$$= 3.6$$

vol. of O_2 and H_2 produced are

$$3.6 \times 5.6 + 3.6 \times 11.2$$

$$= 60.48 \text{ litre}$$

43. (d) 1 watt = 1 J/sec; 1 kWh = $\frac{V}{1000} \times \frac{I}{1000} \times \frac{t}{1000}$ (in hr)

$$\text{Total kWh} = \frac{10^{-3}}{110} \times \frac{10}{10} \times \frac{10}{11}$$

$$\text{Total cost} = 11 \times 2 \times 22 \text{ Rs.}$$

44. (b) $E_{\text{SO}_4^{2-} | \text{S}_2\text{O}_8^{2-}}^\circ$ $E_{\text{H}_2\text{O} | \text{O}_2}^\circ$

45. (b) 10 g Cd should present with 90 g Hg.
mass of Cd required with 4.5 g of Hg at cathode

$$\frac{10}{90} \times 4.5 = 0.5; (\text{Cd}^{2+} + 2e^- \rightarrow \text{Cd})$$

$$\frac{0.5}{112} \times 2 = \frac{4}{96500} \times \frac{t}{1000}$$

$$t = 215.40 \text{ sec}$$

46. (c) Use of electrolysis is in (i) Electrowinning
(ii) Electroplating

47. (c) $E_{\text{Ag}^+ | \text{Ag}}^\circ$ $E_{\text{H}^+ | \text{H}_2}^\circ$ $E_{\text{Cu}^{2+} / \text{Cu}}^\circ$ $E_{\text{O}_2, \text{H}^+ / \text{H}_2\text{O}}^\circ$

48. (a) Chemical energy into electrical energy.

49. (b) G_1° G_2°
 $2 \times FE_{\text{Cl}^- | \text{Cl}_2}^\circ = 2 \times FE_{\text{Cl}_2 | \text{Cl}}^\circ$
 1.36 V

$$50. (b) \begin{array}{ccc} G_3^\circ & G_1^\circ & G_2^\circ \\ n_3 FE_3^\circ & n_1 FE_1^\circ & n_2 FE_2^\circ \\ E_3^\circ & \frac{n_1 E_1^\circ + n_2 E_2^\circ}{n_3} \end{array}$$

51. (a) Maintain electrical neutrality of both half cells.

52. (c) Velocities of ions are nearly the same.

53. (d) A salt bridge contains strong electrolyte with agar-agar.

$$54. (a) \log K_c = \frac{nFE}{2.303 RT}$$

$$55. (c) \begin{array}{ccc} y & mx & \\ G_1^\circ & 1 & F(x) \quad RT \ln K_1 \\ G_2^\circ & 1 & F(y) \quad RT \ln K_2 \\ G_2^\circ & 2 & G_1^\circ \quad 2RT \ln K_1 \\ K_2 & K_1^2, & x \quad y \end{array}$$

$$56. (b) E = E^\circ - \frac{0.0591}{2} \log \frac{[Zn^{2+}]}{[Ag^+]^2}$$

$$57. (c) \text{When } Q = K_c, E = 0$$

58. (b) $Pt(s) | H_2(g) | H^+(aq) || Ag^+(aq) | Ag(s)$
Anode : $H_2 | H^+$, Cathode : $Ag | Ag^+$

59. (c) $Cu(s) | Cu^{2+}(aq) || Cl^-(aq) | Hg_2Cl_2(s) | Hg(l) | Pt(s)$
Anode : $Cu | Cu^{2+}$
Cathode : $Cl^- | Hg_2Cl_2 | Hg$

60. (a) $Pt(s) | Fe^{2+}(aq), Fe^{3+}(aq) || Cr_2O_7^{2-}(aq), Cr^{3+}(aq) | Pt(s)$
Anode : $Fe^{2+} | Fe^{3+}$
Cathode : $Cr_2O_7^{2-} | Cr^{3+}$

61. (c) Anode : $2Ag(s) \rightleftharpoons 2Ag^+(aq) + 2e^-$
Cathode : $Cu^{2+}(aq) + 2e^- \rightleftharpoons Cu(s)$

Net reaction : $2Ag(s) + Cu^{2+}(aq) \rightleftharpoons 2Ag^+(aq) + Cu(s)$

62. (c) Anode : $Cl_2(g) \rightleftharpoons 2Cl^-(aq) + 2e^-$
Cathode : $2Ag^+(aq) + 2e^- \rightleftharpoons 2Ag(s)$

Net reaction : $Cl_2(g) + 2Ag(s) \rightleftharpoons 2Cl^-(aq) + 2Ag^+(aq)$

63. (d) $E_{H_2|H}^\circ = 0V$

$$64. (d) E = E^\circ - \frac{0.0591}{2} \log \frac{[Zn^{2+}]}{[Cu^{2+}]}$$

As $Q > E$

$Q_1 = 1, Q_2 = 0.1, Q_3 = 10, Q_4 = 1$


E_2, E_1, E_4, E_3

65. (b) Ag becomes oxidized and Br becomes reduced.

$$66. (a) N_2(g) + 5H^+(aq) + 4e^- \rightleftharpoons N_2H_5^+(aq), E = 0.23 V \quad \dots(1)$$

$$BiO^+(aq) + 2H^+(aq) + 3e^- \rightleftharpoons Bi(s) + H_2O(l), E = 0.32 V \quad \dots(2)$$


[Reaction (1)] $\times 3$ [Reaction (2)] $\times 4$, we get




$$G = 3(-4FE) + 4(3FE) = 12FE - 12F(0.23) = 12F(0.32)$$

$$E = 0.23 \quad 0.32 \quad 0.55$$

$$67. (a) Cr^{2+}(aq) + Cr^{3+}(aq) + e^- \rightleftharpoons Cr^{3+}(aq), E = 0.41 V \quad \dots(1)$$



[Reaction (1)] $\times 2$ [Reaction (2)] we get.



$$G = 2FE + 2(-1FE) = 2FE - 2F(1.8) = 2F(0.41)$$

$$E = 1.8 \quad 0.41 \quad 1.39$$

$$68. (d) \begin{array}{ccc} 0.34 & 1 & E_{SbO_2|Sb}^\circ \\ E_{SbO_2|Sb}^\circ & 0.34 & 1 \quad 0.66 \end{array}$$

69. (d) For spontaneous $E > 0$,
minimum possible spontaneous $E = 0.28$

70. (a) For best reducing agent, E_{Red}° is minimum.

71. (a) For best oxidizing agent, E_{Red}° is maximum.

72. (d) Maximum voltage $E = 0.49 \quad 0.86 \quad 1.35$

73. (c) $E_{Al^{3+}|Al}^\circ < E_{Cu^{2+}|Cu}^\circ < E_{Cu|Cu^{2+}}^\circ < E_{Al^{3+}|Al}^\circ$ ve
(Non-spontaneous)

74. (b) $E_{Zn^{2+}|Zn}^\circ < E_{Cu^{2+}|Cu}^\circ$;
 $E_{Cu^{2+}|Cu}^\circ < E_{Zn|Zn^{2+}}^\circ$ ve

75. (d) D does not react with HCl, so not easily oxidized at compared to A, B and C

C reduces B as well as D, but unable to reduce A

so A is oxidized easily as compared to other metals





76. (d) $E_{Cu^{2+}|Cu}^\circ < E_{Ag^+|Ag}^\circ < E_{Cu|Cu^{2+}}^\circ < E_{Ag|Ag^+}^\circ$ ve
(Non-spontaneous)

77. (d) From (I) information, C is strong reducing agent.

From (II) information, A is strong reducing agent than D, B and C reducing agent than A.

78. (d) $4\text{Fe(s)} + 3\text{O}_2\text{(g)} \rightarrow 4\text{Fe}^{3+} + 6\text{O}^{2-}\text{(aq)}$
Fe is reducing agent, O_2 is oxidizing agent.
79. (a) $E^\circ_{\text{Na}|\text{Na}} \quad E^\circ_{\text{Zn}^{2+}|\text{Zn}} \quad E^\circ_{\text{Fe}^{2+}|\text{Fe}} \quad E^\circ_{\text{Ag}|\text{Ag}}$
 $E^\circ_{\text{Fe}|\text{Fe}^{2+}} \quad E^\circ_{\text{Ag}|\text{Ag}} \quad \text{ve} \quad (\text{Spontaneous})$
80. (d) $E^\circ_{\text{Zn}|\text{Zn}^{2+}} \quad E^\circ_{\text{Mg}^{2+}|\text{Mg}} \quad 1.608$
(Non-spontaneous)
81. (a) $E^\circ_{\text{F}_2|\text{F}} \quad E^\circ_{\text{Cl}_2|\text{Cl}} \quad E^\circ_{\text{Br}_2|\text{Br}} \quad E^\circ_{\text{I}_2|\text{I}}$
82. (a) $K = 10^{\frac{2(0.277)}{0.0591}} = 2.3 \times 10^9$
83. (a) $E = \frac{0.0591}{2} \log K_{\text{eq}}; \log K_{\text{eq}} = \frac{2 \times 0.22}{0.0591}$
 $7.44 \text{ or } K_{\text{eq}} \sim 2.8 \times 10^7$
84. (a) $E = E^\circ - \frac{0.0591}{2} \log \frac{1}{[\text{Cl}^-]^2}; \text{ as } [\text{Cl}^-] = 1, E$
85. (b) $K = 10^{\frac{6FE}{2.303RT}}$
 $\log 10^{30} = \frac{6 \times E}{0.0591} \quad E = 0.2955 \text{ V}$
86. (b) $E_{\text{cell}} = \frac{0.0591}{2} \log K_{\text{eq}}$
 $0.003 = \frac{0.0591}{2} \log \frac{[\text{D}^+]^2}{[\text{H}^+]^2}$
 $= \frac{[\text{D}^+]}{[\text{H}^+]} = 1.123$
87. (a) $G = 12 \text{ F} \quad E = 636.9 \text{ kJ} \quad 637 \text{ kJ}$
88. (c) $E_{\text{cell}} \quad E_{\text{RP(RHS)}} \quad E_{\text{RP(LHS)}}$
 $1.61 \quad 1.51 \quad 0.10 \text{ V}$
 $G = nFE = 5 \times 96500 \times 0.10 \text{ J}$
 $G = 48.25 \text{ kJ}$
89. (d) $E = E^\circ - \frac{0.0591}{1} \log \frac{[\text{Fe}^{3+}]}{[\text{Ag}][\text{Fe}^{2+}]}$
as $Q = \frac{[\text{Ag}][\text{Fe}^{3+}]}{[\text{Fe}^{2+}]}, E$
90. (d) $E = E^\circ - \frac{0.0591}{2} \log \frac{[\text{HCl}]^2}{P_{\text{H}_2}}$
as $Q = \frac{(P_{\text{H}_2})^2}{[\text{H}^+]^2}, E$
91. (d) $E_{\text{Cr}_2\text{O}_7^{2-}|\text{Cr}^{3+}} \quad E_{\text{Cr}_2\text{O}_7^{2-}|\text{Cr}^{3+}} \quad \frac{0.0591}{6}$
 $\log \frac{[\text{Cr}^{3+}]^2}{[\text{Cr}_2\text{O}_7^{2-}]} = \frac{1}{[\text{H}^+]^{14}}$

- $E_{\text{Cr}_2\text{O}_7^{2-}|\text{Cr}^{3+}} = 1.33 - \frac{0.0591}{6} \log \frac{1}{(0.01)^{14}}$
 1.0542 V
92. (c) $E = 0.57 - \frac{0.0591}{4} \log \frac{[\text{OH}^-]^6}{[\text{TeO}_3^{2-}]}$
 $0.57 - \frac{0.06}{4} \log (10^{-2})^6 = 0.39$
93. (b) $\text{Co}^{2+}\text{(aq)} \rightleftharpoons \text{Co}^{2+}\text{(aq)}, Q = \frac{C_2}{C_1}$
 $\frac{C_1}{C_2} = 1, G = 0$
94. (b) $\text{H}_2\text{(g)} \rightleftharpoons \text{H}_2\text{(g)}$
 $E = \frac{RT}{F} \ln \frac{P_1}{P_2}$
95. (a) A: $\text{H}_2\text{(g)} \rightleftharpoons 2\text{H}^+\text{(aq)} + 2e^-$
C: $\text{Cu}^{2+}\text{(aq)} + 2e^- \rightleftharpoons \text{Cu(s)}$
96. (a) $E = \frac{0.0591}{3} \log \frac{1}{0.04} = 0.028$
97. (c) $\text{Cd}^{2+}\text{(aq)} + \text{Fe(s)} \rightleftharpoons \text{Cd(s)} + \text{Fe}^{2+}\text{(aq)}$
At eqm. $1 \quad x \quad \quad \quad x$
At equilibrium $E = \frac{0.0591}{2} \log \frac{\text{Fe}^{2+}}{\text{Cd}^{2+}}$
 $0.037 = \frac{0.0591}{2} \log \frac{x}{1-x}$
 $x = [\text{Fe}^{2+}] = 0.947 \text{ M}$
 $[\text{Cd}^{2+}] = 0.053 \text{ M}$
98. (d) $1.50 \quad E = \frac{0.0591}{3} \log \frac{0.3}{(0.1)^3}$
 $E = 1.55$
99. (d) $G = nFE$
 $E = \frac{1467 \times 10^3}{(14 \times 96,500)} = 1.08$
100. (b) The cell reaction is
 $\text{H}_2\text{(g)} + \text{I}_2\text{(s)} \rightleftharpoons 2\text{H}^+\text{(aq)} + 2\text{I}^-\text{(aq)}$
 $0.7714 = 0.535 - \frac{0.0591}{2} \log \frac{[\text{H}^+]^2[\text{I}^-]^2}{P_{\text{H}_2}}$
 $\text{pH} = 3$
101. (c) One design a cell in which the half cell reaction is





from $G_3 \quad G_1 \quad G_2$

$$E_{\text{S}^{2-}|\text{CuS}|\text{Cu}} = E_{\text{Cu}^{2+}|\text{Cu}} - \frac{0.0591}{2} \log K_{sp}$$

$$0.34 + \frac{0.0591}{2} \log (10^{-35})$$

$$0.694 \text{ V}$$



$$G_3 = 2F(0.126) - RT \ln \frac{1}{K_{sp}}$$

$$2F(0.248) - 2F(0.126) - RT \ln \frac{1}{K_{sp}}$$

$$\ln \frac{1}{K_{sp}} = \frac{2F}{RT} [0.248 - 0.126]$$

$$\log \frac{1}{K_{sp}} = \frac{2}{(0.0591)} (0.122) = 4.128$$

$$K_{sp} = 10^{-4.128} \text{ or } 7.4 \times 10^{-5}$$



$$E = 1.09 \text{ V}$$





$$nFE = 2.303 RT \log \frac{[\text{Cr}^{3+}]}{[\text{Ag}^+]^3}$$

$$G = 3 \times 96500 \times 1.50 + 8.314 \times 298$$

$$+ 2.303 \log \frac{(0.1)}{(0.1)^3}$$

$$422838.3 \text{ J or } 422.83 \text{ kJ}$$




$$G_2$$



$$G_3 = G_2 - G_1$$

$$2F \quad E \quad 3F \quad 1.40 \quad 1 \quad 1.69 \quad F; \quad E = 1.255 \text{ V}$$

$$E_{\text{Au}^+|\text{Au}^{3+}} = 1.255 \text{ V}$$



$$E_3 = \frac{E_3}{2} \log K_{\text{eq}}; K_{\text{eq}} \sim 6.3 \times 10^{-92}$$



$$E_{\text{cell}} = \frac{RT}{F} \log \frac{[\text{H}^+]_{\text{RHS}}}{[\text{H}^+]_{\text{LHS}}}$$

for max. emf $[\text{H}^+]_{\text{LHS}}$ should be min. and that is for $\text{NH}_4\text{OH}(aq)$



$$E_{\text{cell}} = \frac{RT}{F} \ln \frac{[\text{Ag}^+]_{\text{RHS}}}{[\text{Ag}^+]_{\text{LHS}}}$$

where $[\text{Ag}^+] = \frac{K_{sp}}{[\text{Cl}^-]}$. For spontaneous

chemical change conc. of Ag^+ of cathodic compartment must be greater than anodic compartment.




$$E_{\text{Cr}_2\text{O}_7^{2-}|\text{Cr}^{3+}} = E_{\text{Cr}_2\text{O}_7^{2-}|\text{Cr}^{3+}}$$


$$\frac{0.0591}{n} \log \frac{[\text{Cr}^{3+}]^2}{[\text{Cr}_2\text{O}_7^{2-}][\text{H}^+]^{14}}$$

$$\frac{0.0591}{6} \log [10^{-3}]^{14} = 0.414 \text{ V}$$

$$\text{Oxidising power reduced} = 0.414 \text{ V}$$



$$28.95 \text{ J/K} \cdot \text{mol}$$



$$0.76 - (1.36) = 0.6$$

$$r_G = RT \ln K_{\text{eq}};$$

$$\text{or } \log K_{\text{eq}} = \frac{nFE}{RT} = \frac{2.303}{20} \times \frac{0.6}{0.06}$$

114. (d) Combustion takes place.

115. (b) $\therefore G = 0$ (Electrolytic cell)

116. (b) When aluminium surfaces are exposed to the atmosphere a thin invisible oxidise skin forms immediately, which protects the metal from further oxidation.

[$\therefore E_{\text{OP}}^{\circ}$ of Al is much longer]

117. (d) Acidic medium.

118. (b) $E_{\text{Hg}^{2+}|\text{Hg}}^{\circ} \quad E_{\text{Hg}_2^{2+}|\text{Hg}}^{\circ} \quad E_{\text{Fe}^{2+}|\text{Fe}}^{\circ} \quad E_{\text{Pb}^{2+}|\text{Pb}}^{\circ}$

119. (b) E_{OP}° of Zn E_{OP}° of Fe

120. (a) Metal acts as anode

121. (a) Voltage of cell increases

122. (b) Movement of ions

123. (b) Free ions

124. (c) Water is weak electrolyte.

125. (b) $G = K \frac{A}{l}$

126. (c) $\frac{1}{M} = \frac{1}{M} \times 1000$
 $\frac{1000}{M} = \frac{1}{M} \times 1000$

127. (b) $\frac{K}{N} = \frac{1000}{N}$

128. (b) $G^* = \frac{l}{A} \times \frac{0.02}{0.0004} = 50 \text{ m}^{-1}$

$$G = \frac{1}{R} \times \frac{1}{50} = 0.02 \text{ S}$$

$$G \cdot G^* = 50 \times 0.02 = 1 \text{ Sm}^{-1}$$

129. (a) Conductance (G) $\frac{1}{R} = \frac{1}{300}$;

$$\therefore G = \frac{l}{A}$$

$$\text{Cell constant } \frac{l}{A} = \frac{0.013}{1} = 300 \quad 3.9 \text{ cm}^{-1}$$

130. (b) $\text{HA(aq)} \rightleftharpoons \text{H}^+(\text{aq}) + \text{A}^-(\text{aq})$
 At eqm. $c(1 - \alpha) \quad c \quad c$

$$K_a = \frac{c \alpha^2}{1 - \alpha}; \text{ where } \frac{m}{m}$$

$$K_a = \frac{c \frac{m}{m}}{1 - \frac{m}{m}} = \frac{c \frac{m}{m}}{m \left(\frac{m}{m} - \frac{m}{m} \right)}$$

131. (c) On dilution number of ions decrease in unit volume hence specific conductance decreases. But separation between ions also increase hence equivalent conductance increases.

132. (c) $\frac{150}{3} = 50 \text{ S cm}^2 \text{ eq}^{-1}$

133. (d) n_{eq} ; when $n_{\text{total cationic/Anionic charge}}$

$$\therefore n = 6; \quad \frac{m}{m} \text{ eq } \frac{m}{m} / 6$$

134. (d) $\frac{m(\text{NaBr})}{m(\text{Na}^+)} = \frac{m(\text{NaCl})}{m(\text{Br}^-)} = \frac{m(\text{KBr})}{m(\text{KCl})}$
 $\frac{126.5}{m(\text{Na}^+)} = \frac{151.5}{m(\text{Br}^-)} = \frac{150}{m(\text{Na}^+)} = 50$

135. (c) $\frac{1000}{M} = \frac{1000}{M} \times \frac{1000}{(x - y)}$

$$\text{Solubility (in gL}^{-1}\text{)} = \frac{1000 \times 188}{x - y}$$

136. (a) $G \cdot G^* = \frac{1}{200} \times 2 = 0.01$;

$$\frac{0.01}{0.1} = \frac{1000}{100}$$

137. (b) $R = \frac{l}{A}$

$$82.4 = \frac{1}{2.768} \times 10^3 \times \frac{l}{A}$$

$$\frac{l}{A} = 0.2281$$

138. (a) Equivalent conductivity of BaCl_2
 $127 \quad 76 \quad 203$

139. (c) Ionic mobility ($\text{m}^2 \text{V}^{-1} \text{s}^{-1}$)

$$\frac{\text{Velocity}}{\text{Voltage / Distance}}$$

140. (b) $\text{AgCl} \quad \text{AgNO}_3 \quad \text{HCl} \quad \text{HNO}_3$

$$\begin{array}{l}
 \text{141. (b) } G^* \frac{l}{a} \frac{10}{100} \quad 0.1; G \quad 0.0001 \text{ S} \\
 V \quad 100 \quad 10 \quad 1000 \text{ cm}^3 \quad 1 \text{ litre} \\
 GG^* \quad 0.1 \quad 0.0001 = 10^{-5} \\
 m \quad \frac{1000}{M} \\
 \frac{(0.1 \quad 0.0001) \quad 1000}{0.1} \quad 0.1 \text{ Scm}^2 \text{mol}^{-1}
 \end{array}$$

142. (b) On dilution, no. of ions decrease in per unit volume.

143. (c) On dilution degree of dissociation of a weak electrolyte increase, hence increase in ionic mobility.

144. (d) Completely dissociate into ions.

145. (d) Electric conduction of a salt solution in water depends on the extent of its ionization.

146. (b) Ionic molar conductivity of H⁺ is very high and NH₄OH is a weak electrolyte.

147. (b) Hydrated size order $\text{F}^- < \text{Cl}^- < \text{Br}^- < \text{I}^-$
 Ionic mobility $\frac{1}{\text{Hydrated size}}$

148. (a) Molar conductivity of H⁺ and OH⁻ are very high as compare to other ions.

Initially conductance of solution sharply decreases due to consumption of free H⁺. After complete neutralization further slightly increases due to presence of OH⁻.

149. (c) Before neutralization conductance of solution sharply decreases due to consumption of free H⁺. After neutralization further conductance almost remains same due to formation of buffer solution.

150. (d) Molar conductivity of H⁺ and OH⁻ are very high as compared to other ions.


Initially conductance of solution sharply decreases due to consumption of free H⁺ then increases due to formation of salt (NaCN) and after complete neutralization further sharply increases due to presence of OH⁻.

Level 2

1. (b) No. of equivalent of aluminum, $\frac{W}{E} = \frac{I}{96500} t$

$$\frac{24}{27} \frac{5}{3} \frac{9650}{96500} \frac{0.9}{t} = \frac{t}{148.14 \text{ sec}}$$

2. (c) At Anode:



Eq. of O₂ evolved Eq. of Ag formed 0.1

Total loss in mass $\frac{0.1 \times 32}{4} = 0.8$ 0.1 108

$0.8 + 10.8 = 11.6$

mass of final solution 108 11.6 96.4 g

3. (d) Equivalents of CO₂ produced $\frac{(I \times t)}{96500}$

$\frac{0.5 \times 0.8 \times 96.5 \times 60}{96500} = 0.024$

moles of CO₂ (n = 1) produced 0.024

moles of C₂H₆ (n = 2) produced

$\frac{0.024}{2} = 0.012$

Total moles of gases produced 0.036

$V_{\text{gases}} = \frac{nRT}{P} = \frac{0.036 \times 0.0821 \times 300}{760}$

0.91 litre

4. (d) Total volume of metal layer

$0.25 \times 0.32 \times 10^{-3}$

$0.08 \times 10^{-3} \text{ m}^3$

Total mass of chromium layer

$0.08 \times 7.20 \times 10^6 \times 10^{-3} = 576 \text{ g}$

$\frac{576}{52} \times 6 = \frac{I \times 60}{96500}$

$I = 10.69 \times 10^4 \text{ A}$

5. (a) Initial m moles of Cu²⁺ = 5;

m -eq. or m -moles of H⁺ produced

$100 \times 10^{-2} = 1$

m -moles of Cu²⁺ converted into

$\text{Cu} \quad \frac{1}{2} = 0.5$

m -moles of Cu²⁺ remaining in solution




$5 - 0.5 = 4.5$



m -moles of Cu²⁺ remaining m -moles of



$4.5 = 0.04 \times V \quad V = 112.5 \text{ mL}$





$${}_rG_3 = G_1 - G_2$$

$$2F E = RT \ln(10^{-26}) - 2F \cdot 0$$

$$E = \frac{0.066}{2} \log(10^{-26}) = 0.858 \text{ V}$$

7. (d) To find n we break the cell reaction into two half cell reactions



$$E = \frac{G}{nF} = \frac{(-2746) - 1000}{26 \cdot 96500} = 1.09 \text{ V}$$

8. (c) For normal calomel electrode $E_{RP} = E_{RP}^0$;

$$E_{\text{cell}} = (0.28 - 0) - \frac{0.0591}{2} \log \frac{[\text{H}^+]^2}{P_{\text{H}_2}}$$

$$0.5755 = 0.28 + 0.0591 \text{ pH} + \frac{0.0591}{2} \log(0.1)$$

$$\text{pH} = 5.5$$

9. (a) We know



$$G_{298} = -nFE = -4 \cdot 96500 \cdot 1.23 = -474.78 \text{ kJ}$$

$${}_rH_{298} = {}_rG_{298} + T \cdot {}_rS_{298}$$

$$= (-474.78) + 298 \cdot (-0.32)$$

$$= -570.14 \text{ kJ/mol}$$

$${}_fH_{298} = -570.14 + 285.07 \text{ kJ/mol}$$

10. (a) After addition of Cd and its oxidation into Cd^{2+} .



$$0.1 - x \quad 0.4 - 4x;$$

where $x = 0.08$

$[\text{NO}_3^-]_{\text{remaining}} = 0.02 \text{ M}; [\text{H}^+]_{\text{remaining}} = 0.08 \text{ M}$

$$E_{\text{NO}_3^-|\text{NO}} = E_{\text{NO}_3^-|\text{NO}}^0 - \frac{0.0591}{3} \log \frac{1}{[\text{NO}_3^-][\text{H}^+]^4}$$

$$0.96 - \frac{0.0591}{3} \log \frac{1}{(0.02)(0.08)^4} = 0.84 \text{ V}$$

11. (a) E for normal calomel electrode E for saturated calomel electrode

$$0.33 = 0.27 - \frac{0.06}{2} \log [\text{Cl}^-]^2; [\text{Cl}^-] = 0.1 \text{ M}$$

12. (b) Anode : $\text{H}_2(\text{g}) \rightarrow 2\text{H}^+(\text{aq}) + 2\text{e}^-$; $E = 0.0 \text{ V}$



$$E_{\text{cell}} = E_{\text{RP(RHS)}} - E_{\text{RP(LHS)}}$$

$$1.51 - \frac{0.06}{5} \log \frac{[\text{Mn}^{2+}]}{[\text{MnO}_4^-][\text{H}^+]^8} = 0 - \frac{0.06}{2} \log \frac{P_{\text{H}_2}}{[\text{H}^+]^2}$$

$$1.51 - \frac{0.06}{5} \log \frac{0.01}{0.1 (10^{-2})^8} = - \frac{0.06}{2} \log \frac{0.1}{(10^{-3})^2}$$

$$= 1.48 \text{ V}$$

13. (c) $3\text{Cu}(\text{s}) + 8\text{H}^+ + 2\text{NO}_3^- \rightarrow 3\text{Cu}^{2+} + 2\text{NO} + 4\text{H}_2\text{O}$



Let concentration of HNO_3 is X so $[\text{H}^+] = X$

and $[\text{NO}_3^-] = X$

or

$$\begin{aligned} E_{\text{NO}_3^-|\text{NO}} &= E_{\text{Cu}^{2+}|\text{Cu}} & E_{\text{NO}_3^-|\text{NO}_2} &= E_{\text{Cu}^{2+}|\text{Cu}} \\ E_{\text{NO}_3^-|\text{NO}} &= E_{\text{NO}_3^-|\text{NO}_2} \\ 0.96 - \frac{0.0591}{3} \log \frac{10^{-3}}{x^5} &= 0.79 - \frac{0.0591}{1} \log \frac{10^{-3}}{x^3} \\ 0.62 &= \frac{0.0591}{6} \log \frac{10^{-9}}{x^{10}} = 0.45 - \frac{0.0591}{2} \log \frac{10^{-9}}{x^6} \\ \log x &= 0.657 \sim 0.66 \text{ or } x = 10^{+0.66} \end{aligned}$$


14. (a) Anode : $2\text{Cl}^-(\text{aq}) \rightarrow \text{Cl}_2(\text{g}) + 2\text{e}^-$



$$E_{\text{cell}} = \frac{0.06}{2} \log \frac{[\text{Cl}^-]_{\text{RHS}}^2 (P_{\text{Cl}_2})_{\text{LHS}}}{[\text{Cl}^-]_{\text{LHS}}^2 (P_{\text{Cl}_2})_{\text{RHS}}} \quad (\because E_{\text{cell}} = 0)$$


$$\frac{0.06}{2} \log \frac{(0.01)^2}{(0.1)^2} = \frac{0.4}{0.2} \quad 0.051 \text{ V}$$

15. (a) $\text{H}_2\text{O}(\text{l}) + \text{ClO}_3^-(\text{aq}) \rightarrow \text{ClO}_4^-(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^-$



$$E_{\text{cell}} = 0.33 - 0.36 = -0.03; \quad E_{\text{cell}} = \frac{RT}{2F} \ln K$$

$$0.03 \quad \frac{0.06}{2} \log K \quad \text{or} \quad K = 0.1$$



$$\frac{x^2}{(0.1 - 2x)^2} = \frac{1}{10}; 3.16x = 0.1 - 2x$$

$$5.16x = 0.1$$

$$x = 0.1/5.16 = 0.0193 \quad 1.9 \times 10^{-2}$$

$$E_{\text{cell}} = \frac{0.06}{1} \log \frac{[\text{H}^+]_{\text{RHS}}}{[\text{H}^+]_{\text{LHS}}}$$

$$E_{\text{cell}} = 0.06 [(\text{pH})_{\text{LHS}} - (\text{pH})_{\text{RHS}}]$$

$$(\text{pH})_{\text{LHS}} - \text{p}K_a = \log \frac{[A^-]}{[HA]}$$

$$4 - \log \frac{1}{0.1} = 5; (\text{pH})_{\text{RHS}} = 4$$

$$E_{\text{cell}} = 0.06 (5 - 4) = 0.06 \text{ V}$$

$$E_{\text{cell}} = E_{\text{H}^+|\text{H}_2} - E_{\text{OH}^-|\text{Cd}(\text{OH})_2|\text{Cd}}$$

$$E_{\text{H}^+|\text{H}_2} - E_{\text{Cd}^{2+}|\text{Cd}}$$

$$E_{\text{cell}} = E - \frac{0.06}{2} \log \frac{[\text{Cd}^{2+}]}{[\text{H}^+]^2}$$

$$E_{\text{cell}} = 0,$$

$$E_{\text{cell}} = 0.03 \log \frac{[\text{Cd}^{2+}][\text{OH}^-]^2}{K_w^2}$$

$$\log \frac{K_{sp}}{K_w^2} = \frac{0.39}{0.03} = 13$$

$$K_{sp} = 10^{13} (10^{-14})^2 = 10^{-15}$$

$$E_{\text{cell}} = \frac{0.06}{2} \log \frac{[\text{H}^+]_{\text{RHS}}^2 [P_{\text{H}_2}]_{\text{LHS}}}{[\text{H}^+]_{\text{LHS}}^2 [P_{\text{H}_2}]_{\text{RHS}}}; \quad \therefore \quad 1$$

$$[\text{OH}^-] = \sqrt{K_b C}; [\text{OH}^-] = 10^{-3}$$

$$[\text{H}^+]_{\text{LHS}} = 10^{-11}$$

$$[\text{H}^+]_{\text{RHS}} = \sqrt{K_a C} = 10^{-4}$$

$$E_{\text{cell}} = \frac{0.06}{2} \log \frac{(10^{-4})^2 \cdot 0.1}{(10^{-11})^2} = 1 - 0.39 \text{ V}$$

16. (b)

or


17. (c)

or

 $\therefore$

18. (a)

or



$$K_{a_1} \cdot K_{a_2} = \frac{[\text{H}^+]^2 [\text{S}^{2-}]}{[\text{H}_2\text{S}]}$$

$$\frac{10^{-21} \cdot 0.1}{[\text{S}^{2-}]} = \frac{(10^{-3})^2 [\text{S}^{2-}]}{10^{-16}}$$

$$E_{\text{S}^{2-}|\text{Ag}_2\text{S}|\text{Ag}} = E_{\text{Ag}^+|\text{Ag}} - \frac{0.06}{2} \log \frac{[\text{S}^{2-}]}{K_{sp}}$$

$$= 0.80 - \frac{0.06}{2} \log \frac{10^{-16}}{10^{-49}}$$

$$= 0.80 - \frac{0.06}{2} \cdot 33 = 0.19 \text{ V}$$

20. (c) Normality of resultant solution

$$\frac{0.1 \text{ V}}{V} = \frac{0.05 \text{ N}}{V}$$

$$\text{eq} = \frac{1000}{1000} \cdot \frac{0.0055}{0.05} = 110 \text{ S cm}^2 \text{eq}^{-1}$$

21. (b) Resultant conc. of NaCl $\frac{0.1 \text{ V}}{4 V} = 0.025 \text{ N}$

and HCl $\frac{0.1 \text{ V}}{4 V} = 0.05 \text{ N}$

$$\text{total solution} = \frac{N(\text{NaCl}) + N(\text{HCl})}{1000}$$

$$= \frac{0.025 + 0.05}{1000} = \frac{0.075}{1000} = 0.018$$

$$\text{eq(HCl)} = \frac{18}{0.05} = 360 \text{ S cm}^2 \text{eq}^{-1}$$

22. (b) The molar conductivity of the dissociated form of crotonic acid is

$$\frac{m(\text{HC})}{m(\text{HCl})} = \frac{m(\text{NaC})}{m(\text{NaCl})}$$

$$\frac{(426 - 383) \text{ cm}^2 \text{mol}^{-1}}{383 \text{ cm}^2 \text{mol}^{-1}} = \frac{m(\text{NaC})}{1 \text{ cm}^2 \text{mol}^{-1}}$$

The molar conductivity of HC $\frac{1000}{m}$


$$\frac{m(\text{HC})}{0.001} = \frac{3.83 \cdot 10^{-5} \text{ cm}^2 \text{mol}^{-1}}{0.001}$$

$$= 38.3 \text{ cm}^2 \text{mol}^{-1}$$

The degree of dissociation,

$$\frac{m(\text{HC})}{m(\text{HC})} = \frac{(38.3 \text{ cm}^2 \text{mol}^{-1})}{(383 \text{ cm}^2 \text{mol}^{-1})} = 0.1$$

$$K_a = \frac{C}{1} \cdot \frac{(10^{-3})(0.1)^2}{1 \cdot 0.1} = 1.1 \cdot 10^{-5}$$



for sparingly soluble salt

$$M = \frac{1000}{2(x_1 + x_2 + x_3)}$$

$$K_{sp} = M^2 = \frac{2.5 \cdot 10^5 x^2}{(x_1 + x_2 + x_3)^2}$$

24. (a) Conductivity of Na_2SO_4 $2.6 \cdot 10^{-4}$

$$m(\text{Na}_2\text{SO}_4) = \frac{1000}{0.001} = 1000$$

$$260 \text{ S cm}^2 \text{mol}^{-1}$$

$$m(\text{SO}_4^{2-}) = \frac{m(\text{Na}_2\text{SO}_4)}{2} = \frac{260}{2} = 130$$

$$160 \text{ S cm}^2 \text{mol}^{-1}$$

Conductivity of CaSO_4 solution

$$7 \cdot 10^{-4} + 2.6 \cdot 10^{-4} = 4.4 \cdot 10^{-4} \text{ S cm}^{-1}$$

$$m(\text{CaSO}_4) = \frac{m(\text{Ca}^{2+})}{2} = \frac{m(\text{SO}_4^{2-})}{2}$$

$$\frac{120}{280} = \frac{160}{280}$$

$$280 \text{ S cm}^2 \text{mol}^{-1}$$

$$\text{Solubility } C = \frac{1000}{280} = 3.57 \text{ M}$$

$$K_{sp} = [\text{Ca}^{2+}][\text{SO}_4^{2-}]_{\text{total}}$$

$$= (0.00157)(0.00157 + 0.001) = 4.0 \cdot 10^{-6} \text{ M}^2$$

25. (b) $K_a = \frac{C}{1} \cdot \frac{1.6 \cdot 10^{-5}}{0.01} = 0.04$

$$\frac{1.6 \cdot 10^{-5}}{0.01} = 0.04$$

$$\frac{m}{m} = \frac{0.04}{380} = 10^{-4}$$

$$15.2 \cdot 10^4 \text{ S cm}^2 \text{mol}^{-1}$$

$$C = \frac{15.2 \cdot 10^4}{10^3} = 15.2 \text{ M}$$

$$(\because 1 \text{ m}^3 = 1000 \text{ litre})$$

$$k = G \cdot G^* \text{ and } G^* = 0.01 \text{ m}^{-1}$$

where $G^* = \frac{l}{A}$

$$G = \frac{1.52 \times 10^{-2}}{0.01} = 1.52 \text{ S}$$

26. (c) Cell X : At anode Cl_2 gas evolved;
At cathode H_2 gas evolved.

Cell Y : At anode O_2 gas evolved;
At cathode Ag deposited.

Cell Z : At anode O_2 gas evolved;
At cathode Cu deposited

$$\text{Eqn. of Ag deposited} = \frac{W}{E} = \frac{21.6}{108} = 0.20$$

Eqn. of O_2 gas evolved; Vol. of O_2 gas evolved 0.2 5.6 1.12 L

Vol. of H_2 evolved Vol. of Cl_2 evolved
0.2 11.2 2.24 L

27. (a) Eq. of H_2 Eq. of O_2 Eq. of $\text{H}_2\text{S}_2\text{O}_8$

$$2 \frac{22.4}{22.4} \quad 4 \frac{8.4}{22.4} \quad 2n$$

$$2 \quad 1.5 \quad 2n \quad n \quad 0.25$$

Level 3

Passage-1

1. (c), 2. (a), 3. (d)

Equivalent of Zn^{2+} produced 0.1 or
moles of $\text{Zn}^{2+} = \frac{0.1}{2} = 0.05$

+ve charge increases in first compartment so due to interaction and maintain electrical neutrality Zn^{2+} move toward II compartment and NO_3^- move towards first compartment. Solution is always electrically neutral so charge of 1 Zn^{2+} is neutralized by 2 NO_3^- .

$[\text{Zn}^{2+}]$ in first compartment

$$1 \frac{0.05}{2} = 1.025 \text{ M}$$

Concentration of NO_3^- in second compartment

$$1 \quad 0.05 \quad 0.95 \text{ M}$$

In third compartment moles of Cu^{2+} reduced

$$\frac{0.05}{2} = 0.025$$

Relatively +ve charge increased so SO_4^{2-} and Na move toward opposite direction to maintain electrical neutrality

$$[\text{SO}_4^{2-}]_{\text{remaining}} = 1 \frac{0.025}{2} = 0.975 \text{ M}$$

Passage-2

1. (b) From equation 1 and 2, $S = \frac{d(G)}{dT}$

$$S = \frac{d(nFE)}{dT} = nF \frac{dE}{dT} \text{ or } \frac{dE}{dT} = \frac{S}{nF}$$

2. (b) From equation 2;

$$H = nF \frac{dE}{dT} = T \frac{nFE_{\text{cell}}}{dT}$$

$$218 \quad 1000 \quad 2 \quad 96500 \quad 300 \quad \frac{dE}{dT} = 2 \quad 96500 \quad 1.015$$

$$\frac{dE}{dT} = 3.81 \times 10^{-4} \text{ VK}^{-1}$$

3. (a) $S = nF \frac{dE}{dT} = 2 \quad 96500 \quad 3.81 \times 10^{-4}$
= 75.53 J/mol-K

Passage-3

2. (b)

$$m \cdot C = \frac{m \cdot M}{1000}$$

$$\frac{200 \quad 0.04}{1000} = 8 \times 10^3 \text{ Scm}^{-1}$$

$$G = \frac{l}{A} = 8 \times 10^{-3} \quad G = \frac{0.50}{2}$$

$$\text{or } G = 0.032 \text{ S; } V = IR \quad \frac{I}{G}$$

$$I = 5 \quad 0.032 \quad 0.16 \text{ A}$$


Passage-4

1. (c) $2 \quad \frac{4.48 \quad 1}{22.4} = \frac{2 \quad 1}{2} \quad \frac{9650}{96,500}$

I = 4 amp.

2. (b) Mass of $\text{H}_2\text{O} = \frac{4.48}{22.4} \times 18 = 3.6 \text{ g}$

3. (b) $2\text{H}_2\text{O}(l) \quad 4\text{H}^+(aq) \quad \text{O}_2(g) \quad 4e^-$
 $2\text{Cu}^{2+} \quad 4e^- \quad 2\text{Cu}(s)$





$$\text{Moles}_{\text{Cu}} = 2 \quad \frac{4 \quad 241.25}{96,500}$$

$$\text{Mole}_{\text{H}} = 2 \quad \text{mole of Cu} = 0.005 \quad 2 \quad 0.01$$

pH = 2

Passage-5



$$K_{sp}(\text{Agx}) \quad x(x-y) \quad 3 \quad 10^{-12} \quad \dots(1)$$

$$K_{sp}(\text{Agy}) \quad y(x-y) \quad 10^{-12} \quad \dots(2)$$

$$K_{(\text{Agx} \rightleftharpoons \text{Agy})} = \frac{K_{\text{Ag}}}{K_x} = \frac{K_y}{K_x} = 0.4 \quad 10^{-6} \quad \dots(3)$$

$$\text{Ag} \quad \frac{1000 \quad K_{\text{Ag}}}{(x-y)} = 60 \quad \dots(4)$$

$$x \quad \frac{1000 \quad K_x}{x} = 90 \quad \dots(5)$$

$$K_y = \frac{1.45 \quad 10^{-7}}{\frac{1000 \quad 1.45 \quad 10^{-7}}{5 \quad 10^{-7}}} = 290$$


$$x-y = 2 \quad 10^{-6} \quad \dots(6)$$

$$\frac{x}{y} = 3 \quad \dots(7)$$

$$x = 1.5 \quad 10^{-6}, y = 5 \quad 10^{-7}, K_{\text{Ag}} = 1.2 \quad 10^{-7}$$

$$K_x = 1.35 \quad 10^{-7} \quad \frac{4}{3} x = 2 \quad 10^{-6}$$

Subjective Problems



Required charge for reduction of 1 mole $\text{C}_6\text{H}_5\text{NO}_2 = 6F$

2. Equivalent mass $\text{O}_2 = \frac{32}{4} = 8$




3. $\frac{0.967}{63.5} \times 2 \times \frac{2}{197} n_f$

4. $\frac{0.18}{\text{Eq. mass}} \times \frac{5 \quad 60 \quad 9.65}{96500}, \text{Eq. mass} = 6$

5. $2 \text{ mole Cl}_2 \times \frac{4F}{F}$

6. $G = 96.5 \quad 10^3 \quad 1 \quad 96500 \quad E$


7. $K_C = 10^{\frac{2(0.34 \quad 0.15)}{0.0591}} = 2.6 \quad 10^6$



$2 \quad F \quad (0.40) - 2 \quad F \quad (0.61) = RT \ln K_f$

$\ln K_f = \frac{2F}{RT} (0.61 - 0.4)$

$\log K_f = \frac{2 \quad F \quad (0.4) - 2F(0.61)}{0.0591} = 7 \quad K_f = 10^7$



$\text{pH} = 7$

11. $1500 \times \frac{1}{2 \quad 10^{-3}} = \frac{l}{A}$

Cell constant $1500 \times 2 \quad 10^{-3} = 3$



% ionisation of $\text{CH}_3\text{COOH} = \frac{0.034}{0.038} \times 100 = 89.47\%$

$\frac{2.72 \quad 10^{-3}}{0.034} \times 100 = 8$

13. $0.12 \times \frac{0.0591}{2} \log \frac{1}{x} = 0.24$

$\log \frac{1}{x} = \frac{0.12 \times 2}{0.06} = 4$

$x = 10^{-4}$



$\text{pH} = 4, [\text{H}^+] = 10^{-4} \quad C$

$\frac{m}{m} = \frac{200}{400} = 0.5;$

$K_a = \frac{(C)}{(1)} \times \frac{10^{-4}(0.5)}{(1)} = 10^{-4}; \text{p}K_a = 4$

15. $E_{\text{Cl}^-/\text{AgCl}/\text{Ag}} = E_{\text{Ag}/\text{Ag}^+} + \frac{0.0591}{1} \log K_{sp}$

$0.209 = 0.80 + \frac{0.0591}{1} \log K_{sp}$

$K_{sp} = 10^{-10}$; Let solubility of AgCl in 0.01 M NaCl solution is x

$10^{-10} = x(x - 0.01)$

$x = 10^{-8}$


Moles of AgCl dissolved in

$10 \text{ L} \quad 10^{-8} \quad 10^{-7}$

DILUTE SOLUTIONS

Vapour Pressure : The pressure exerted by the vapours of liquid which are in equilibrium with it at a given temperature.

- ❖ Vapour pressure is not a surface phenomena.
- ❖ Vapour pressure increases with increase in temperature.



$$\ln \frac{K_{p_2}}{K_{p_1}} = \ln \frac{P_2}{P_1} = -\frac{H_{\text{vap.}}}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right) ;$$

where $H_{\text{vap.}}$ is molar enthalpy of vaporisation (J/mol or cal/mol)

Raoult's Law

(i) Liquid solution containing non volatile solute :

Vapour pressure of liquid solution Mole fraction of solvent in liquid solution.

$$\frac{P_{\text{Solution}}}{P_{\text{Solvent}}} = X_{\text{Solvent}}$$

Where P_{Solvent} Vapour pressure of pure liquid solvent, X_{Solvent} Mole fraction of solvent.

For binary liquid solution ; $X_{\text{Solute}} + X_{\text{Solvent}} = 1$

Relative lowering of vapour pressure $\frac{P - P_{\text{Solution}}}{P} = X_{\text{Solute}}$

(ii) Vapour Pressure of Solution containing volatile solute (B) and volatile solvent (A) :

The equilibrium vapour pressure of a volatile component is linearly proportional to the mole fraction of that component in liquid phase.

A solution contains volatile solute (B) and volatile solvent (A) :

$$\begin{array}{ccc} P_A & P_A X_A & \text{and } P_B & P_B X_B \\ P & P_A X_A & P_B X_B & \end{array}$$

Where :

P_A and P_B are partial vapour pressure of A and B at given temperature.

P_A vapour pressure of pure A. ; P_B vapour pressure of pure B.

X_A mole fraction of A in liquid phase. ; X_B mole fraction of B in liquid phase.

y_A mole fraction of A in vapour phase ; y_B mole fraction of B in vapour phase.

Partial vapour pressure of 'A' mole fraction of 'A' in vapour phase total vapour pressure.

$$P_A = y_A P \quad \text{and} \quad P_B = y_B P$$

$$P_A X_A = y_A P \quad \text{and} \quad P_B X_B = y_B P$$

$$\frac{X_A}{y_A} \frac{P}{P_A} = \frac{X_B}{y_B} \frac{P}{P_B} = 1;$$

$$\frac{y_A}{P_A} = \frac{y_B}{P_B} = \frac{1}{P}$$

$$y_A = \frac{P_A X_A}{P_A X_A + P_B X_B}; \quad y_B = \frac{P_B X_B}{P_A X_A + P_B X_B}$$

Ideal Solutions

The solutions which obey Raoult's Law are called ideal solutions. For ideality :

$F_{\text{Solute}} \dots \dots \text{Solvent}$ $F_{\text{Solute}} \dots \dots \text{Solute}$ (F Intermolecular attractive interactions)

$F_{\text{Solvent}} \dots \dots \text{Solvent}$

$$H_{\text{mix.}} = 0; \quad V_{\text{mix.}} = 0$$

$$S_{\text{mix.}} = 0; \quad G_{\text{mix.}} = 0$$

Examples : Benzene & Toluene, Hexane & Heptane, $\text{C}_2\text{H}_5\text{Br}$ & $\text{C}_2\text{H}_5\text{I}$

Non-ideal solutions : The mixture which do not follow Raoult's law is known as non-ideal solutions.

Types of Real or Non-ideal Solutions

- ❖ Non-ideal solution with positive deviation.
- ❖ Non-ideal solution with negative deviation.

Solution shows positive deviation

$$P_{T, \text{exp}} > (P_A X_A + P_B X_B)$$


$$\begin{array}{l} A \text{ --- } A \\ A \text{ --- } B \\ B \text{ --- } B \end{array}$$

$$H_{\text{mix.}} \quad \text{ve}$$

$$V_{\text{mix.}} \quad \text{ve}$$

$$S_{\text{mix.}} \quad \text{ve}$$

$$G_{\text{mix.}} \quad \text{ve}$$



Solution shows negative deviation

$$P_{T, \text{exp}} < (P_A X_A + P_B X_B)$$


$$\begin{array}{l} A \text{ --- } A \\ A \text{ --- } B \\ B \text{ --- } B \end{array}$$

$$H_{\text{mix.}} \quad \text{ve}$$

$$V_{\text{mix.}} \quad \text{ve}$$

$$S_{\text{mix.}} \quad \text{ve}$$

$$G_{\text{mix.}} \quad \text{ve}$$



Azeotropic Mixture. During distillation, the mole fraction of more volatile component in vapour state is higher than that in liquid state. This makes distillation possible. However, there exist some solutions for particular compositions of which the mole fraction of components in liquid and vapour state is same. Thus, no distillation is possible, a mixture and it is termed as **azeotropic mixture**.

Azeotropes are binary mixtures having the same composition in liquid and vapour phase and boil at a constant temperature.

The solutions which show a large positive deviation from Raoult's law form minimum boiling azeotrope at a specific composition.

The solutions which show a large negative deviation from Raoult's law form maximum boiling azeotrope at a specific composition.

Completely Immiscible Liquids : When they are distilled, they distil in the ratio of their vapour pressure at that temperature.

$$\frac{P}{P_A} = \frac{P_B}{P_A}$$

For vapour of pure liquid we can apply : $P_A V = n_A RT$ and $P_B V = n_B RT$

So,
$$\frac{P_A}{P_B} = \frac{n_A}{n_B}$$

When A and B are distilled mass ratio is given as
$$\frac{w_B}{w_A} = \frac{P_B}{P_A} \frac{M_B}{M_A}$$

Colligative Properties : Colligative Properties depend on the number of solute particles irrespective to their nature.

- ❖ Relative lowering of vapour pressure
- ❖ Elevation of boiling point
- ❖ Depression of freezing point
- ❖ Osmotic pressure

Normal Colligative Properties : When neither association nor dissociation of solute particles take place.

(i) Relative Lowering of Vapour Pressure :
$$\frac{P - P_s}{P} = X_{\text{Solute}}$$

(ii) Elevation of boiling point : $T_b = K_b m$

(iii) Depression of freezing point : $T_f = K_f m$

(iv) Osmotic pressure : $\pi = CRT$

(i) Relative lowering of Vapour Pressure :

$$\frac{P - P_s}{P} = X_{\text{Solute}} = \frac{n}{n + N}$$


Where, n moles of solute, N moles of solvent

$$\frac{P - P_s}{P} = \frac{n}{N}$$

$$\text{Molality (m)} = \frac{P - P_s}{P} \times \frac{1000}{M(\text{in g/mol})}$$

Where, M molar mass of solvent.

Ostwald-Walker Method : Experimental measure of relative lowering in V.P



Loss in mass of solution containers P

Loss in mass of solvent containers $(P - P')$

gain in mass of dehydrating agent P'

$$\frac{P - P'}{P} = \frac{\text{Loss in mass of solvent}}{\text{Gain in mass of dehydrating agent}}$$

(ii) Elevation in Boiling Point :

$$T_b = K_b m$$

where T_b T_b T_b

K_b Boiling point elevation constant
or ebullioscopic constant

$$K_b = \frac{RT_b^2}{1000 L_v}$$

L_v Latent heat of vapourisation per gram

$$K_b = \frac{MRT_b^2}{1000 H_{\text{vap}}}$$

H_{vap} enthalpy of vapourisation per mole

M molar mass of solvent (in g/mol)

T_b Boiling point of solvent

(iii) Depression in Freezing point :

$$T_f = K_f m$$

where T_f T_f T_f

K_f Freezing point depression constant
or cryoscopic constant

$$K_f = \frac{RT_f^2}{1000 L_f}$$

L_f Latent heat of fusion per gram

$$K_f = \frac{MRT_f^2}{1000 H_{\text{fus}}}$$

H_{fus} enthalpy of fusion per mole

M molar mass of solvent (in g/mol)

T_f freezing point of solvent

(iv) Osmotic Pressure () : The hydrostatic pressure built up on the solution which just stops osmosis. Alternatively, it may be defined as the pressure which must be applied to the concentrated solution in order to stop osmosis.

For dilute solutions $\pi = CRT$ $h = \frac{\pi}{\rho g}$

where C is the total molar concentration of all the free species present in the solution, h is the height developed by the column of the concentrated solution and ρ is the density of the solution in the column.

On the basis of osmotic pressure, the solutions can be classified in three classes.

❖ **Isotonic solutions:** Two solutions having same osmotic pressures are called isotonic solutions.

(This implies $C_1 = C_2$ at same temperature).

❖ **Hypertonic and hypotonic solution :** When two solutions are being compared, then the solution with higher osmotic pressure is termed as hypertonic. The solution with lower osmotic pressure is termed as **hypotonic**.

❖ **Abnormal Colligative Properties** : When solute particle associated or dissociated in solvent.

$$i = \text{Vant Hoff factor} = \frac{\text{Actual moles of solute}}{\text{Moles of solute without dissociation or association}} = \frac{\text{Observed or experimental colligative properties}}{\text{Theoretical or calculated colligative properties}}$$

$$(i) \text{ Relative lowering of vapour pressure : } \frac{P - P_s}{P} = \frac{i \cdot n_{\text{solute}}}{n_{\text{solvent}}}$$

$$(ii) \text{ Elevation of boiling point : } T_b = i \cdot K_b m$$

$$(iii) \text{ Depression of freezing point : } T_f = i \cdot K_f m$$

$$(iv) \text{ Osmotic pressure : } \pi = i \cdot CRT$$

$$\text{For dissociation : } i = 1 + \frac{\alpha}{n-1}$$

$$\text{For association : } i = 1 - \frac{\alpha}{n-1}$$

$$\text{Degree of dissociation } \alpha = \frac{i-1}{n-1}$$

$$\text{Degree of association } \alpha = \frac{1-i}{1/n-1}$$

Where n = Total number of particles of solute after dissociation or association.

Henry Law

This law deals with dissolution of gas in liquid. It states that mass of any gas dissolved in any solvent per unit volume is proportional to pressure of gas in equilibrium with liquid.

$$(i) \quad m = k \cdot P$$

Where, m = mass of gas dissolved per unit volume.

k = proportionality constant.

P = pressure of gas.

$$(ii) \quad P = K_H \cdot X_{\text{gas}}$$

Where, P = Pressure of gas.

K_H = Henry's constant

X_{gas} = Mole fraction of gas




Level 1


- The vapour pressure of a given liquid will decrease if :
 - surface area of liquid is decreased
 - the volume of liquid in the container is decreased
 - the volume of the vapour phase is increased
 - the temperature is decreased
- The normal boiling point of water is 373 K. Vapour pressure of water at temperature T is 19 mm Hg. If enthalpy of vaporisation is 40.67 kJ/mol, then temperature T would be (Use : $\log 2 = 0.3$, $R : 8.3 \text{ JK}^{-1} \text{ mol}^{-1}$):
 - 250 K
 - 291.4 K
 - 230 K
 - 290 K
- A sample of liquid H_2O of mass 18.0 g is injected into an evacuated 7.6 L flask maintained at 27.0°C . If vapour pressure of H_2O at 27°C is 24.63 mm Hg, what weight percentage of the water will be vapourised when the system comes to equilibrium? Assume water vapours behaves as an ideal gas. The volume occupied by the liquid water is negligible compared to the volume of the container:
 - 1%
 - 10%
 - 18%
 - 20%
- Raoult's law is obeyed by each constituent of a binary liquid solution when:
 - the forces of attractions between like molecules are greater than those between unlike molecules
 - the forces of attractions between like molecules are smaller than those between unlike molecules
 - the forces of attractions between like molecules are identical with those between unlike molecules
 - the volume occupied by unlike molecules are different
- For a binary ideal liquid solution, the total vapour pressure of the solution is given as:
 - $P_{\text{total}} = P_A^\circ + (P_A^\circ - P_B^\circ) X_B$
 - $P_{\text{total}} = P_B^\circ + (P_A^\circ - P_B^\circ) X_A$
 - $P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ) X_A$
 - $P_{\text{total}} = P_B^\circ + (P_B^\circ - P_A^\circ) X_B$
- For an ideal binary liquid solution with P_A and P_B , which relation between X_A (mole fraction of A in liquid phase) and Y_A (mole fraction of A in vapour phase) is correct ?
 - $Y_A = Y_B$
 - $X_A = X_B$
 - $\frac{Y_A}{Y_B} = \frac{X_A}{X_B}$
 - $\frac{Y_A}{Y_B} = \frac{X_A}{X_B}$
- An ideal solution is formed by mixing two volatile liquids A and B. X_A and X_B are the mole fractions of A and B respectively in the solution and Y_A and Y_B are the mole fractions of A and B respectively in the vapour phase. A plot of $\frac{1}{Y_A}$ along y-axis against $\frac{1}{X_A}$ along x-axis gives a straight line. What is the slope of the straight line ?
 - $\frac{P_B}{P_A}$
 - $\frac{P_A}{P_B}$
 - $P_B - P_A$
 - $P_A - P_B$

8. For a dilute solution, Raoult's law states that :
- (a) the lowering of vapour pressure is equal to the mole fraction of solute
 - (b) the relative lowering of vapour pressure is equal to the mole fraction of solute
 - (c) the relative lowering of vapour pressure is proportional to the amount of solute in solution
 - (d) the vapour pressure of the solution is equal to the mole fraction of solvent
9. The solubility of a specific non-volatile salt is 4 g in 100 g of water at 25°C. If 2.0 g, 4.0 g and 6.0 g of the salt added of 100 g of water at 25°C, in system X, Y and Z. The vapour pressure would be in the order:
- (a) X Y Z (b) X Y Z (c) Z X Y (d) X Y Z
10. The boiling point of C_6H_6 , CH_3OH , $C_6H_5NH_2$ and $C_6H_5NO_2$ are 80°C, 65°C, 184°C and 212°C respectively. Which will show highest vapour pressure at room temperature :
- (a) C_6H_6 (b) CH_3OH (c) $C_6H_5NH_2$ (d) $C_6H_5NO_2$
11. 6.0 g of urea (molecular mass = 60) was dissolved in 9.9 moles of water. If the vapour pressure of pure water is P , the vapour pressure of solution is:
- (a) 0.10 P (b) 1.10 P (c) 0.90 P (d) 0.99 P
12. An ideal solution was found to have a vapour pressure of 80 torr when the mole fraction of a non-volatile solute was 0.2. What would be the vapour pressure of the pure solvent at the same temperature?
- (a) 64 torr (b) 80 torr (c) 100 torr (d) 400 torr
13. The vapour pressure of an aqueous solution of sucrose at 373 K is found to be 750 mm Hg. The molality of the solution at the same temperature will be :
- (a) 0.26 (b) 0.73 (c) 0.74 (d) 0.039
14. Estimate the lowering of vapour pressure due to the solute (glucose) in a 1.0 M aqueous solution at 100°C :
- (a) 10 torr (b) 18 torr (c) 13.45 torr (d) 24 torr
15. Calculate the mass of non-volatile solute having molecular mass 40, which should be dissolved in 57 gm octane to reduce its vapour pressure to 80%:
- (a) 47.2 g (b) 5 g (c) 106.2 g (d) None of these
16. Equal mass of a solute are dissolved in equal mass of two solvents A and B and formed very dilute solution. The relative lowering of vapour pressure for the solution B has twice the relative lowering of vapour pressure for the solution A. If M_A and M_B are the molecular mass of solvents A and B respectively, then:
- (a) $M_A = M_B$ (b) $M_B = 2 M_A$ (c) $M_A = 4 M_B$ (d) $M_A = 2 M_B$
17. An ideal solution has two components A and B. A is more volatile than B, i.e., $P_A > P_B$ and also $P_A > P_{total}$. If X_A and Y_A are mole fractions of components A in liquid and vapour phases, then:
- (a) $X_A > Y_A$ (b) $X_A < Y_A$ (c) $X_A = Y_A$ (d) Data insufficient
18. At 25°C, the vapour pressure of pure liquid A (mol. mass = 40) is 100 torr, while that of pure liquid B is 40 torr, (mol. mass = 80). The vapour pressure at 25°C of a solution containing 20 g of each A and B is:
- (a) 80 torr (b) 59.8 torr (c) 68 torr (d) 48 torr

19. Two liquids A and B form ideal solutions. At 300 K, the vapour pressure of solution containing 1 mole of A and 3 mole of B is 550 mm Hg. At the same temperature, if one more mole of B is added to this solution, the vapour pressure of the solution increases by 10 mm Hg. Determine the vapour pressure of A and B in their pure states (in mm Hg):
 (a) 400, 600 (b) 500, 500 (c) 600, 400 (d) None of these
20. Two liquids A and B have vapour pressure in the ratio $P_A : P_B = 1 : 3$ at a certain temperature. Assume A and B form an ideal solution and the ratio of mole fractions of A to B in the vapour phase is 4 : 3, then the mole fraction of B in the solution at the same temperature is :
 (a) $\frac{1}{5}$ (b) $\frac{2}{3}$ (c) $\frac{4}{5}$ (d) $\frac{1}{4}$
21. Two liquids A and B have P_A and P_B in the ratio of 1 : 3 and the ratio of number of moles of A and B in liquid phase are 1 : 3. Then mole fraction of ' A ' in vapour phase in equilibrium with the solution is equal to:
 (a) 0.1 (b) 0.2 (c) 0.5 (d) 1.0
22. Based on the given diagram, which of the following statements regarding the homogenous solutions of two volatile liquids are correct?



- (1) Plots AD and BC show that Raoult's law is obeyed for the solution in which B is a solvent and A is the solute and as well as for that in which A is solvent and B is solute.
 (2) Plot CD shows that Dalton's law of partial pressures is obeyed by the binary solution of components A and B .
 (3) $EF \parallel EG \parallel EH$; and AC and BD correspond to the vapour pressures of the pure solvents A and B respectively.
 (a) Only 1 (b) 2 and 3 (c) 1 and 3 (d) All
23. Which represents correct difference when non-volatile solute is present in an ideal solution?



- (a) I, II, III (b) I, III (c) II, III (d) I, II


24. Select correct statement :

- (a) Solution has more molecular randomness than a pure solvent. The entropy change between solution and solid is larger than the entropy change between pure solvent and solid
- (b) Heat of fusion of solution and solvent are similar since similar forces of intermolecular forces are involved
- (c) Solution containing sugar freezes at a lower temperature than pure water
- (d) All are correct statements

25. Select correct statement?


- (a) Heats of vaporisation for a pure solvent and for a solution are similar because similar intermolecular forces between solvent molecules must be overcome in both cases
- (b) Entropy change between solution and vapour is smaller than the entropy change between pure solvent and vapour
- (c) Boiling point of the solution is larger than that of the pure solvent
- (d) All are correct statements

26. The vapour pressure curves of the same solute in the same solvent are shown below. The curves are parallel to each other and do not intersect. The concentrations of solutions are in order of:



- (a) $I < II < III$
- (b) $I = II = III$
- (c) $I > II > III$
- (d) $I > III > II$

27. Boiling point composition diagram of the liquid-vapour equilibrium for A and B is shown in the figure. If a binary liquid mixture of A and B is distilled fractionally, which of the following would be correct observation?



- (a) Composition of the still (residue) will approach pure liquid B only
- (b) Composition of the distillate will approach pure A only
- (c) Composition of distillate and residue will approach pure A and B respectively
- (d) Neither of the component can be obtained in pure state

28. The boiling point of an azeotropic mixture of water-ethanol is less than that of both water and ethanol. Then:

- (a) the mixture will show negative deviation from Raoult's law
- (b) the mixture will show positive deviation from Raoult's law
- (c) the mixture will show no deviation from Raoult's law
- (d) this mixture can be considered as true solution

29. Formation of a solution from two components can be considered as :

- (i) Pure solvent separated solvent molecules, H_1
- (ii) Pure solute separated solute molecules, H_2
- (iii) separated solvent and solute molecules solution, H_3

Solution so formed will be ideal if :

- (a) $H_{\text{Soln}} = H_1 + H_2 + H_3$
- (b) $H_{\text{Soln}} = H_1 + H_2 - H_3$
- (c) $H_{\text{Soln}} = H_1 + H_2 - H_3$
- (d) $H_{\text{Soln}} = H_3 + H_1 + H_2$


30. Total vapour pressure of mixture of 1 mol X ($P_X = 150$ torr) and 2 mol Y ($P_Y = 300$ torr) is 240 torr. In this case:

- (a) there is a negative deviation from Raoult's law
- (b) there is a positive deviation from Raoult's law
- (c) there is no deviation from Raoult's law
- (d) can not be decided

31. In a mixture of A and B, components show positive deviation when:

- (a) A-B interaction is stronger than A-A and B-B interaction
- (b) A-B interaction is weaker than A-A and B-B interaction
- (c) $V_{\text{mix}} < 0$, $S_{\text{mix}} < 0$
- (d) $V_{\text{mix}} < 0$, $S_{\text{mix}} < 0$

32. A liquid mixture having composition corresponding to point Z in the figure shown is subjected to distillation at constant pressure. Which of the following statement is correct about the process?



- (a) The composition of distillate differs from the mixture
- (b) The boiling point goes on changing
- (c) The mixture has highest vapour pressure than for any other composition
- (d) Composition of an azeotrope alters on changing the external pressure

33. Which will form maximum boiling azeotrope ?

- (a) C_6H_6 - $\text{C}_6\text{H}_5\text{CH}_3$ solution
- (b) HNO_3 - H_2O solution
- (c) $\text{C}_2\text{H}_5\text{OH}$ - H_2O solution
- (d) *n*-hexane and *n*-heptane

34. Total vapour pressure of mixture of 1 mole of volatile component A ($P_A = 100$ mm Hg) and 3 mole of volatile component B ($P_B = 80$ mm Hg) is 90 mm Hg. For such case:

- (a) there is positive deviation from Raoult's law
- (b) boiling point has been lowered
- (c) force of attraction between A and B is weaker than that between A and A or between B and B
- (d) all the above statements are correct

35. The azeotropic mixture of water (B.P. = 100 °C) and HCl (B.P. = 86°C) boils at about 120°C. During fractional distillation of this mixture, it is possible to obtain :
 (a) pure HCl (b) pure H₂O
 (c) pure H₂O as well as pure HCl (d) neither H₂O nor HCl
36. Azeotropic mixture of water and C₂H₅OH boils at 351 K. By distilling the mixture it is possible to obtain :
 (a) Pure C₂H₅OH only (b) Pure water only
 (c) Neither C₂H₅OH nor water (d) Both water and C₂H₅OH in pure state
37. An azeotropic mixture of two liquids has a boiling point higher than either of them when it:
 (a) shows positive deviation from Raoult's law
 (b) shows negative deviation from Raoult's law
 (c) shows ideal behaviour
 (d) is saturated
38. If two liquids A (P_A = 100 torr) and B (P_B = 200 torr) which are completely immiscible with each other (each one will behave independently of the other) are present in a closed vessel, the total vapour pressure of the system will be:
 (a) less than 100 torr (b) greater than 200 torr
 (c) between 100 to 200 torr (d) 300 torr
39. When a liquid that is immiscible with water was steam distilled at 95.2°C at a total pressure of 748 torr, the distillate contained 1.25 g of the liquid per gram of water. The vapour pressure of water is 648 torr at 95.2°C, what is the molar mass of liquid?
 (a) 7.975 g/mol (b) 166 g/mol (c) 145.8 g/mol (d) None of these
40. Water and chlorobenzene are immiscible liquids. Their mixture boils at 89°C under a reduced pressure of 7.7×10^4 Pa. The vapour pressure of pure water at 89°C is 7×10^4 Pa. Mass per cent of chlorobenzene in the distillate is :
 (a) 50 (b) 60 (c) 78.3 (d) 38.46
41. Which of the following is not a colligative property?
 (a) Vapour pressure (b) Depression in f.pt.
 (c) Elevation in b.pt. (d) Osmotic pressure
42. The degree of dissociation of an electrolyte is α and its van't Hoff factor is i . The number of ions obtained by complete dissociation of 1 molecule of the electrolyte is :
 (a) $\frac{i-1}{\alpha}$ (b) $i-1$ (c) $\frac{i-1}{1-\alpha}$ (d) $\frac{i-1}{1}$
43. One mole of a solute A is dissolved in a given volume of a solvent. The association of the solute take place as follows:

$$nA \rightleftharpoons A_n$$

 If α is the degree of association of A, the van't Hoff factor i is expressed as:
 (a) $i = 1 - \alpha$ (b) $i = 1 - \frac{\alpha}{n}$ (c) $i = \frac{1-\alpha}{1}$ (d) $i = 1$
44. The van't Hoff factor i for an electrolyte which undergoes dissociation and association in solvent are respectively:
 (a) greater than one and less than one (b) less than one and greater than one
 (c) less than one and less than one (d) greater than one and greater than one

45. Which solution has the highest vapour pressure?
(a) 0.02 M NaCl at 50°C (b) 0.03 M sucrose at 15°C
(c) 0.005 M CaCl₂ at 50°C (d) 0.005 M CaCl₂ at 25°C
46. An aqueous solution is 1.00 molal in KI. Which change will cause the vapour pressure of the solution to increase ?
(a) addition of water (b) addition of NaCl
(c) addition of Na₂SO₄ (d) Addition of 1.0 molal KI
47. Four solutions of K₂SO₄ with the concentrations 0.1 m, 0.01 m, 0.001 m and 0.0001 m are available. The maximum value of colligative property corresponds to :
(a) 0.0001 m solution (b) 0.001 m solution (c) 0.01 m solution (d) 0.1 m solution
48. Moles of Na₂SO₄ to be dissolved in 12 mole water to lower its vapour pressure by 10 mm Hg at a temperature at which vapour pressure of pure water is 50 mm is:
(a) 1.5 mole (b) 2 mole (c) 1 mole (d) 3 mole
49. A very diluted saturated solution of a sparingly soluble salt X₃Y₄ has a vapour pressure of 20 mm Hg at temperature T, while pure water exerts a pressure of 20.0126 mm Hg at the same temperature. Calculate molality (m) at temperature T:
(a) 6.3 × 10⁻⁴ (b) 3.5 × 10⁻² (c) 5 × 10⁻³ (d) None of these
50. When 1 mole of a solute is dissolved in 1 kg of H₂O, boiling point of solution was found to be 100.5°C. K_b for H₂O is:
(a) 0.5 (b) 100 (c) 100.5 (d) 95.5
51. Chloroform, CHCl₃, boils at 61.7°C. If the K_b for chloroform is 3.63°C/molal, what is the boiling point of a solution of 15.0 kg of CHCl₃ and 0.616 kg of acenaphthalene, C₁₂H₁₀?
(a) 61.9 (b) 62.0 (c) 52.2 (d) 62.67
52. A compound has the empirical formula C₁₀H₈Fe. A solution of 0.26 g of the compound in 11.2 g of benzene (C₆H₆) boils at 80.26°C. The boiling point of benzene is 80.10°C; the K_b is 2.53°C/molal. What is the molecular formula of the compound?
(a) C₃₀H₂₄Fe₃ (b) C₁₀H₈Fe (c) C₅H₄Fe (d) C₂₀H₁₆Fe₂
53. A solution of 0.640 g of azulene in 100.0 g of benzene boils at 80.23°C. The boiling point of benzene is 80.10°C; and K_b is 2.53°C/molal. What is the molecular mass of azulene?
(a) 108 (b) 99 (c) 125 (d) 134
54. One molal solution of a carboxylic acid in benzene shows the elevation of boiling point of 1.518 K. The degree of association for dimerization of the acid in benzene is (K_b for benzene 2.53 K kg mol⁻¹):
(a) 60% (b) 70% (c) 75% (d) 80%
55. The boiling point elevation constant for toluene is 3.32 K kg mol⁻¹. The normal boiling point of toluene is 110.7°C. The enthalpy of vaporisation of toluene would be nearly:
(a) 17.0 kJ mol⁻¹ (b) 34.0 kJ mol⁻¹ (c) 51.0 kJ mol⁻¹ (d) 68.0 kJ mol⁻¹
56. Which one of the following aqueous solutions will exhibit highest boiling point ?
(a) 0.015 M urea (b) 0.01 M KNO₃ (c) 0.10 M Na₂SO₄ (d) 0.015 M glucose
57. Calculate the percentage degree of dissociation of an electrolyte XY₂ (Normal molar mass 164) in water if the observed molar mass by measuring elevation in boiling point is 65.6:
(a) 75% (b) 25% (c) 65% (d) None of these

58. If the elevation in boiling point of a solution of non-volatile, non-electrolytic and non-associating solute in solvent (K_b x K.kg.mol⁻¹) is y K, then the depression in freezing point of solution of same concentration would be (K_f of the solvent z K.kg.mol⁻¹)
- (a) $\frac{2xz}{y}$ (b) $\frac{yz}{x}$ (c) $\frac{xz}{y}$ (d) $\frac{yz}{2x}$
59. When a solution containing non-volatile solute freezes, which equilibrium would exist?
- (a) solid solvent $\rightleftharpoons$ liquid solvent (b) solid solute $\rightleftharpoons$ liquid solution
(c) solid solute $\rightleftharpoons$ liquid solvent (d) solid solvent $\rightleftharpoons$ liquid solution
60. Bromoform has a normal freezing point of 7.734°C and K_f 14.4°C/m. A solution of 2.60 g of an unknown substance in 100 g of bromoform freezes at 5.43°C. What is the molecular mass of the unknown substance?
- (a) 16.25 (b) 162.5 (c) 100 (d) none of these
61. C₆H₆ freezes at 5.5°C. At what temperature will a solution of 10.44 g of C₄H₁₀ in 200 g of C₆H₆ freeze? K_f (C₆H₆) 5.12°C/m
- (a) 4.608°C (b) 0.892°C (c) 5.5°C (d) none of these
62. How much ethyl alcohol must be added to 1.0 L of water so that solution will not freeze at -4°F? (K_f 1.86°C/m)
- (a) < 20 g (b) < 10.75 g (c) < 494.5 g (d) > 494.5 g
63. The freezing point of a solution of 2.40 g of biphenyl (C₁₂H₁₀) in 75.0 g of benzene (C₆H₆) is 4.40°C. The normal freezing point of benzene is 5.50°C. What is the molal freezing point constant (°C/m) for benzene?
- (a) 5.3 (b) 5.1 (c) 4.6 (d) 4.8
64. A solution containing 1.8 g of a compound (empirical formula CH₂O) in 40 g of water is observed to freeze at -0.465°C. The molecular formula of the compound is (K_f of water 1.86 kg K mol⁻¹) :
- (a) C₂H₄O₂ (b) C₃H₆ (c) C₄H₈O₄ (d) C₆H₁₂O₆
65. Freezing point of the following equilibrium, liquid solvent $\rightleftharpoons$ solid solvent is :
- (a) $\frac{H}{T} \frac{G}{S}$ (b) $\frac{H}{S}$ (c) $\frac{G}{S}$ (d) $\frac{S}{H}$
66. Freezing point of a solution is smaller than freezing point of a solvent. It is due to :
- (a) H of solution and solvent is almost identical since intermolecular forces between solvent molecules are involved
(b) S of solution (between solution and solid) is larger than that of the S of solvent (between solvent and solid)
(c) S of the solution is smaller than that of the solvent
(d) H of the solution is much higher than of solvent but S of solution is smaller than that of the solvent
67. When 36.0 g of a solute having the empirical formula CH₂O is dissolved in 1.20 kg of water, the solution freezes at -0.93°C. What is the molecular formula of the solute? (K_f 1.86°C kg mol⁻¹)
- (a) C₂H₄O (b) C₂H₂O₂ (c) C₂H₄O₃ (d) C₂H₄O₂

68. Calculate the molecular mass of a substance whose 7.0% by mass solution in water freezes at -0.93°C . The cryoscopic constant of water is $1.86^{\circ}\text{C kg mol}^{-1}$:
- (a) 140 g mol^{-1} (b) 150.5 g mol^{-1}
(c) 160 g mol^{-1} (d) 155 g mol^{-1}
69. Camphor is often used in molecular mass determination because
- (a) it is readily available (b) it has a very high cryoscopic constant
(c) it is volatile (d) it is solvent for organic substances
70. For 1 molal solution of each compound minimum freezing point will be assuming complete ionisation in each case:
- (a) $[\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_3$ (b) $[\text{Fe}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2 \cdot \text{H}_2\text{O}$
(c) $[\text{Fe}(\text{H}_2\text{O})_4\text{Cl}_2]\text{Cl} \cdot 2\text{H}_2\text{O}$ (d) $[\text{Fe}(\text{H}_2\text{O})_3\text{Cl}_3] \cdot 3\text{H}_2\text{O}$
71. For 1 molal solution of each compound maximum freezing point will be assuming complete ionisation in each case:
- (a) $[\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_3$ (b) $[\text{Fe}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2 \cdot \text{H}_2\text{O}$
(c) $[\text{Fe}(\text{H}_2\text{O})_4\text{Cl}_2]\text{Cl} \cdot 2\text{H}_2\text{O}$ (d) $[\text{Fe}(\text{H}_2\text{O})_3\text{Cl}_3] \cdot 3\text{H}_2\text{O}$
72. $\text{PtCl}_4 \cdot 6\text{H}_2\text{O}$ can exist as a hydrated complex. Its 1 molal aq. solution has depression in freezing point of 3.72. Assume 100% ionisation and $K_f(\text{H}_2\text{O}) = 1.86^{\circ}\text{C mol}^{-1} \text{ kg}$, then complex is :
- (a) $[\text{Pt}(\text{H}_2\text{O})_6]\text{Cl}_4$ (b) $[\text{Pt}(\text{H}_2\text{O})_4\text{Cl}_2]\text{Cl}_2 \cdot 2\text{H}_2\text{O}$
(c) $[\text{Pt}(\text{H}_2\text{O})_3\text{Cl}_3]\text{Cl} \cdot 3\text{H}_2\text{O}$ (d) $[\text{Pt}(\text{H}_2\text{O})_2\text{Cl}_4] \cdot 4\text{H}_2\text{O}$
73. A complex is represented as $\text{CoCl}_3 \cdot x\text{NH}_3$. Its 0.1 molal solution in water shows $T_f = 0.558 \text{ K}$. K_f for H_2O is $1.86 \text{ K molality}^{-1}$. Assuming 100% ionisation of complex and co-ordination number of Co is six, calculate formula of complex:
- (a) $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$ (b) $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
(c) $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$ (d) none of these
74. The freezing point of equimolal aqueous solutions will be highest for:
- (a) $\text{C}_6\text{H}_5\text{NH}_3\text{Cl}$ (b) $\text{Ca}(\text{NO}_3)_2$ (c) $\text{La}(\text{NO}_3)_3$ (d) $\text{C}_6\text{H}_{12}\text{O}_6$
75. The freezing point of a 4% aqueous solution of 'A' is equal to the freezing point of 10% aqueous solution of 'B'. If the molecular mass of 'A' is 60, then the molecular mass of 'B' will be:
- (a) 160 (b) 90 (c) 45 (d) 180
76. Depression in freezing point of 0.01 molal aqueous HCOOH solution is 0.02046. 1 molal aqueous urea solution freezes at -1.86°C . Assuming molality equal to molarity, pH of HCOOH solution is:
- (a) 2 (b) 3 (c) 4 (d) 5
77. When mercuric iodide is added to the aqueous solution of KI, then the:
- (a) freezing point is raised (b) freezing point is lowered
(c) freezing point does not change (d) boiling point does not change
78. Dimer of acetic acid in liquid benzene is in equilibrium with acetic acid monomer at certain temperature and pressure. If 25% of the dimer molecules are separated out then
- (a) freezing point of the solution reduces (b) average molar mass of solute increases
(c) boiling point of solution increases (d) molar mass of solute decreases





79. The temperature of a city was found to be -9.3°C . A car was used, whose radiator was filled with 5 L of water. What minimum quantity of antifreezing agent ethylene glycol were added to water of radiator in order to use the car for travelling? (K_f of water $1.86\text{ K kg mol}^{-1}$)
- (a) 3200 g (b) 1670 g (c) 1550 g (d) 2100 g
80. The cryoscopic constant of water is $1.86\text{ K kg mol}^{-1}$. A 0.01 molal acetic acid solution produces a depression of 0.0194°C in the freezing point. The degree of dissociation of acetic acid is:
- (a) zero (b) 0.043 (c) 0.43 (d) 1
81. If 1m solution of benzoic acid in benzene has a freezing point depression of 3.84°C . (K_f $5.12^{\circ}\text{C mol}^{-1}\text{ kg}$) and boiling point elevation of 2.53°C (K_b $2.53^{\circ}\text{C mol}^{-1}\text{ kg}$), then select the correct statement/s :
- Statement I : there is dimer formation when undergoing freezing
Statement II : there is no change when undergoing boiling
Statement III : reverse of I and II
Statement IV : dimer formation in freezing and boiling state
- (a) I, II (b) II, III (c) III, I (d) only I
82. In a 0.5 molal solution KCl, KCl is 50% dissociated. The freezing point of solution will be (K_f $1.86\text{ K kg mol}^{-1}$):
- (a) 274.674 K (b) 271.60 K (c) 273 K (d) none of these
83. A 1.0 g sample of $\text{Co}(\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2)_3\text{Cl}_3$ is dissolved in 25.0 g of water and the freezing point of the solution is -0.87°C . How many ions are produced per mole of compound? The K_f of water is $1.86^{\circ}\text{C/molal}$
- (a) 2 (b) 3 (c) 4 (d) 5
84. An aqueous solution contains 3% and 1.8% by mass. urea and glucose respectively. What is the freezing point of solution? (K_f 1.86 C/m)
- (a) -1.172°C (b) -2.27°C (c) -1.5°C (d) none of these
85. Phenol associates in benzene to a certain extent in dimerisation reaction. A solution containing 0.02 kg of phenol in 1.0 kg of benzene has its freezing point depressed 0.69 K . Hence, degree of association of phenol dimerized will be :
- [$K_f(\text{C}_6\text{H}_6)$ $5.12\text{ K kg mol}^{-1}$]
- (a) 0.63 (b) 0.73 (c) 0.83 (d) 0.93
86. Assuming complete ionisation, the solution having maximum freezing point will be:
- (a) 1 M CaF_2 (b) 1.5 M $\text{Al}_2(\text{SO}_4)_3$ (c) 2 M NaCl (d) 1 M AgNO_3
87. In a 0.2 molal aqueous solution of a weak acid HX the degree of dissociation is 0.25. The freezing point of the solution will be nearest to: (K_f $1.86\text{ K kg mol}^{-1}$)
- (a) -0.26°C (b) 0.465°C (c) -0.48°C (d) -0.465°C
88. An aqueous solution of 0.01 M KCl causes the same elevation in boiling point as an aqueous solution of urea. The concentration of urea solution is:
- (a) 0.01 M (b) 0.005 M (c) 0.02 M (d) 0.04 M
89. When some NaCl was dissolved in water, the freezing point depression was numerically equal to twice the molal f.p. depression constant. The relative lowering of vapour pressure of the solution is nearly :
- (a) 0.036 (b) 0.018 (c) 0.0585 (d) 0.072

90. Which one of the following statements is false ?
(a) The correct order of osmotic pressure for 0.01 M aqueous solution of each compound is $\text{BaCl}_2 > \text{KCl} > \text{CH}_3\text{COOH} > \text{sucrose}$
(b) Isotonic solutions are those solutions which have the same osmotic pressure
(c) Raoult's law states that the vapour pressure of a component over a solution is proportional to its mole fraction in liquid state
(d) Two sucrose solutions of same molality prepared in different solvents will have the same freezing point depression
91. 0.1 molal aqueous solution of an electrolyte AB_3 is 90% ionised. The boiling point of the solution at 1 atm is : ($K_{b(\text{H}_2\text{O})} = 0.52 \text{ K kg mol}^{-1}$)
(a) 273.19 K (b) 374.92 K (c) 376.4 K (d) 373.19 K
92. Which of the following aqueous solutions has osmotic pressure nearest to pure solvent?
(a) Na_2SO_4 (b) BaCl_2 (c) $\text{Al}_2(\text{SO}_4)_3$ (d) $\text{C}_{12}\text{H}_{22}\text{O}_{11}$
93. 0.1M NaCl and 0.05M BaCl_2 solutions are separated by a semi-permeable membrane in a container. For this system, choose the correct answer :
(a) There is no movement of any solution across the membrane
(b) Water flows from BaCl_2 solution towards NaCl solution
(c) Water flows from NaCl solution towards BaCl_2 solution
(d) Osmotic pressure of 0.1M NaCl is lower than the osmotic pressure of BaCl_2 (assume complete dissociation)
94. Two aqueous solutions, A and B, are separated by a semi-permeable membrane. The osmotic pressure of solution A immediately begins to decrease. Which of the following statement is true ?
(a) The solvent molecules are moving from the solution of higher osmotic pressure to that of lower osmotic pressure.
(b) The initial osmotic pressure of solution B is greater than that of solution A.
(c) Solvent molecules are moving from solution B into solution A.
(d) Both (a) and (b) are true statements.
95. Which one of the following pairs of solution can we expect to be isotonic at the same temperature?
(a) 0.1 M urea and 0.1 M NaCl (b) 0.1 M urea and 0.2 M MgCl_2
(c) 0.1 M NaCl and 0.1 M Na_2SO_4 (d) 0.1 M $\text{Ca}(\text{NO}_3)_2$ and 0.1 M Na_2SO_4
96. The empirical formula of a non-electrolyte is CH_2O . A solution containing 3 g L^{-1} of the compound exerts the same osmotic pressure as that of 0.05M glucose solution. The molecular formula of the compound is :
(a) CH_2O (b) $\text{C}_2\text{H}_4\text{O}_2$ (c) $\text{C}_4\text{H}_8\text{O}_4$ (d) $\text{C}_3\text{H}_6\text{O}_3$
97. A semipermeable membrane used in the measurement of osmotic pressure of a solution allows the passage of:
(a) solute molecules through it (b) solvent molecules through it
(c) both solvent and solute molecules (d) either solvent or solute
98. In the case of osmosis, solvent molecules move from :
(a) higher vapour pressure to lower vapour pressure
(b) higher concentration to lower concentration

- (c) lower vapour pressure to higher vapour pressure
(d) higher osmotic pressure to lower osmotic pressure
- 99.** The osmotic pressures of equimolar solutions of urea, BaCl_2 and AlCl_3 will be in the order :
(a) $\text{AlCl}_3 > \text{BaCl}_2 > \text{urea}$ (b) $\text{BaCl}_2 > \text{AlCl}_3 > \text{urea}$
(c) $\text{urea} > \text{BaCl}_2 > \text{AlCl}_3$ (d) $\text{BaCl}_2 > \text{urea} > \text{AlCl}_3$
- 100.** Which of the following solutions will have maximum osmotic pressure? Assume 90% dissociation of each salt:
(a) decimolar aluminium sulphate
(b) decimolar barium chloride solution
(c) decimolar sodium sulphate solution
(d) solution of equal volumes of decimolar barium chloride and decimolar sodium sulphate solutions
- 101.** Consider 0.1 M solutions of two solutes X and Y. The solute X behaves as a univalent electrolyte while the solute Y dimerises in solution. Which of the following statements are correct regarding these solutions?
(1) The boiling point of the solution of X will be higher than that of Y
(2) The osmotic pressure of the solution of Y will be lower than that of X
(3) The freezing point of the solution of X will be lower than that of Y
(4) The relative lowering of vapour pressure of both the solutions will be the same
Select the correct answer from the option given below :
(a) 1, 2 and 3 (b) 2, 3 and 4
(c) 1, 2 and 4 (d) 1, 3 and 4
- 102.** If M_{normal} is the normal molecular mass and α is the degree of ionization of $\text{K}_3[\text{Fe}(\text{CN})_6]$, then the abnormal molecular mass of the complex in the solution will be:
(a) $M_{\text{normal}} (1 - 2\alpha)^{-1}$ (b) $M_{\text{normal}} (1 - 3\alpha)^{-1}$
(c) $M_{\text{normal}} (1 - \alpha)^{-1}$ (d) equal to M_{normal}
- 103.** Equal volumes of 0.1 M urea and 0.1 M glucose are mixed. The mixture will have:
(a) lower osmotic pressure (b) same osmotic pressure
(c) higher osmotic pressure (d) none of these
- 104.** A 5% (w/V) solution of cane sugar (molecular mass 342) is isotonic with 1% (w/V) solution of a substance X. The molecular mass of X is:
(a) 34.2 (b) 171.2 (c) 68.4 (d) 136.8
- 105.** Insulin is dissolved in a suitable solvent and the osmotic pressure (π) of solutions of various concentrations (g/cm^3) C is measured at 20°C . The slope of a plot of π against C is found to be 4.65×10^{-3} . The molecular mass of the insulin (g/mol) is:
(a) 3×10^5 (b) 9×10^5
(c) 4.5×10^5 (d) 5.16×10^6
- 106.** An aqueous solution of sucrose ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$) having a concentration of 34.2 gram/litre has an osmotic pressure of 2.38 atmospheres at 17°C . For an aqueous solution of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) to be isotonic with this solution, its concentration should be :
(a) 34.2 gram per litre (b) 17.1 gram per litre
(c) 18.0 gram per litre (d) 36.0 gram per litre


- 107.** Which of the following experimental methods is adopted to determine osmotic pressure?
 (a) Berkley-Hartely's method (b) Beckmann's method
 (c) Landsberger's method (d) Differential method
- 108.** Based upon the technique of reverse osmosis the approximate pressure required to desalinate sea water containing 2.5% (mass/volume) KNO_3 at 27°C will be
 (a) 10.5 atm (b) 21 atm (c) 12.2 atm (d) 6.09 atm
- 109.** A 1% (mass/vol) KCl solution is ionised to the extent of 80%. The osmotic pressure at 27°C of the solution will be:
 (a) 6.95 atm (b) 5.94 atm
 (c) 2.71 atm (d) 3.30 atm
- 110.** Osmotic pressure of blood is 7.40 atm, at 27°C . Number of moles of glucose to be used per litre for an intravenous injection that is to have same osmotic pressure of blood is:
 (a) 0.3 (b) 0.2 (c) 0.1 (d) 0.4
- 111.** The relationship between osmotic pressures (π_1 , π_2 and π_3) at a definite temperature when 1 g glucose, 1 g urea and 1 g sucrose are dissolved in 1 litre of water is (assume $i = 1$ for all):
 (a) $\pi_1 = \pi_2 = \pi_3$ (b) $\pi_3 = \pi_1 = \pi_2$
 (c) $\pi_2 = \pi_1 = \pi_3$ (d) $\pi_2 = \pi_3 = \pi_1$
- 112.** van't Hoff proved that osmotic pressure (π) is a colligative property. For an ideal solution, osmotic pressure (π) is helpful to determine that molecular mass of solute using $M_B = \frac{W_B RT}{\pi V}$

Relation can be expressed by the curve (C = concentration):







- 113.** A solution containing 4.0 g of PVC in 2 litre of dioxane (industrial solvent) was found to have an osmotic pressure 3.0×10^{-4} atm at 27°C . The molar mass of the polymer (g/mol) will be :
 (a) 1.6×10^4 (b) 1.6×10^5 (c) 1.6×10^3 (d) 1.6×10^2
- 114.** The osmotic pressures of 0.010 M solutions of KI and of sucrose ($\text{C}_{12}\text{H}_{22}\text{O}_{11}$) are 0.432 atm and 0.24 atm respectively. The van't Hoff factor for KI is :
 (a) 1.80 (b) 0.80 (c) 1.2 (d) 1.0
- 115.** What is the correct sequence of osmotic pressure of 0.01 M aq. solution of :
 (1) $\text{Al}_2(\text{SO}_4)_3$ (2) Na_3PO_4
 (3) BaCl_2 (4) Glucose
 (a) 4 2 3 1 (b) 3 4 2 1
 (c) 3 4 1 2 (d) 1 2 3 4

- 116.** 1.0 molar solution of the complex salt, $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$, displays an osmotic pressure of $3RT$. 0.5 L of the same solution on treatment with excess of AgNO_3 solution will yield (assume 1) :
 (a) 0.5 mole of AgCl (b) 1.0 mole of AgCl (c) 1.5 mole of AgCl (d) 3.0 mole of AgCl
- 117.** A 0.010 g sample of $\text{Cr}(\text{NH}_3)_4(\text{SO}_4)\text{Cl}$ is dissolved in 25.0 mL of water and the osmotic pressure of the solution is 59.1 torr at 25°C . How many moles of ions are produced per mole of compound?
 (a) 1 (b) 4 (c) 2 (d) 3
- 118.** Which of the following aqueous solutions should have the highest osmotic pressure?
 (a) 0.011 M AlCl_3 at 50°C (b) 0.03 M NaCl at 25°C
 (c) 0.012 M $(\text{NH}_4)_2\text{SO}_4$ at 25°C (d) 0.03 M NaCl at 50°C
- 119.** X_3Y_2 (i 5) when reacted with A_2B_3 (i 5) in aqueous solution gives brown colour. These are separated by a semipermeable membrane AB as shown. Due to osmosis there is:



- (a) brown colour formation in side X (b) brown colour formation in side Y
 (c) formation in both of the sides X and Y (d) no brown colour formation
- 120.** Which of the following curves represents the Henry's law?







- 121.** According to Henry's law, the solubility of a gas in a given volume of liquid increases with increase in:
 (a) temperature (b) pressure
 (c) Both (a) and (b) (d) None of these
- 122.** At 300 K, 40 mL of $\text{O}_3(\text{g})$ dissolves in 100 g of water at 1.0 atm. What mass of ozone dissolved in 400 g of water at a pressure of 4.0 atm at 300 K?
 (a) 0.1 g (b) 1.24 g
 (c) 0.48 g (d) 4.8 g

- 123.** 1 kg of water under a nitrogen pressure of 1 atmosphere dissolves 0.02 gm of nitrogen at 293 K. Calculate Henry's law constant :
- (a) 7.7×10^4 atm (b) 7.7×10^3 atm
(c) 2×10^5 atm (d) 2×10^2 atm
- 124.** According to Henry's law, the partial pressure of gas (P_g) is directly proportional to mole fraction of gas in liquid solution, $P_{\text{gas}} = K_H \cdot X_{\text{gas}}$, where K_H is Henry's constant. Which is incorrect ?
- (a) K_H is characteristic constant for a given gas-solvent system
(b) Higher is the value of K_H , lower is solubility of gas for a given partial pressure of gas
(c) K_H has temperature dependence
(d) K_H decreases with increase of temperature
- 125.** At 760 torr pressure and 20°C temperature, 1 L of water dissolves 0.04 gm of pure oxygen or 0.02 gm of pure nitrogen. Assuming that dry air is composed of 20% oxygen and 80% nitrogen (by volume), the masses (in g/L) of oxygen and nitrogen dissolved by 1 L of water at 20°C exposed to air at a total pressure of 760 torr are respectively :
- (a) 0.008, 0.016 (b) 0.016, 0.008
(c) 0.16, 0.08 (d) 0.04, 0.02




Level 2

- Two beaker A and B present in a closed vessel. Beaker A contains 152.4 g aqueous solution of urea, containing 12 g of urea. Beaker B contains 196.2 g glucose solution, containing 18 g of glucose. Both solutions allowed to attain the equilibrium. Determine mass % of glucose in its solution at equilibrium :
 (a) 6.71 (b) 14.49 (c) 16.94 (d) 20
- Two components A and B form an ideal solution. The mole fractions of A and B in ideal solution are X_A and X_B , while that of in vapour phase, these components have their mole fractions as Y_A and Y_B . Then, the slope and intercept of plot of $\frac{1}{Y_A}$ vs. $\frac{1}{X_A}$ will be :
 (a) $\frac{P_A}{P_B}, \frac{P_B}{P_B} \frac{P_A}{P_B}$ (b) $\frac{P_B}{P_A}, \frac{P_A}{P_A} \frac{P_B}{P_A}$ (c) $\frac{P_B}{P_A}, \frac{P_B}{P_B} \frac{P_A}{P_A}$ (d) $P_A \quad P_B, \frac{P_A}{P_B}$
- At 48°C , the vapour pressure of pure CS_2 is 850 torr. A solution of 2.0 g of sulphur in 100 g of CS_2 has a vapour pressure 844.9 torr. Determine the atomicity of sulphur molecule:
 (a) 1 (b) 2 (c) 4 (d) 8
- An ideal solution contains two volatile liquids A ($P^\circ = 100$ torr) and B ($P^\circ = 200$ torr). If mixture contain 1 mole of A and 4 mole of B then total vapour pressure of the distillate is:
 (a) 150 (b) 180 (c) 188.88 (d) 198.88
- The vapour pressure of two pure liquids A and B , that form an ideal solution are 100 and 900 torr respectively at temperature T . This liquid solution of A and B is composed of 1 mole of A and 1 mole of B . What will be the pressure, when 1 mole of mixture has been vaporized?
 (a) 800 torr (b) 500 torr (c) 300 torr (d) None of these
- Which of the following represents correctly the changes in thermodynamic properties during the formation of 1 mole of an ideal binary solution :



7. A saturated solution of XCl_3 has a vapour pressure 17.20 mm Hg at 20°C , while pure water vapour pressure is 17.25 mm Hg. Solubility product (K_{sp}) of XCl_3 at 20°C is:
 (a) 9.8×10^{-2} (b) 10^{-5} (c) 2.56×10^{-6} (d) 7×10^{-5}
8. A certain non-volatile electrolyte contains 40% carbon, 6.7% hydrogen and 53.3% oxygen. An aqueous solution containing 5% by mass of the solute boils at 100.15°C . Determine the molecular formula of the compound ($K_b = 0.51^\circ\text{C/m}$):
 (a) HCHO (b) CH_3OH (c) $\text{C}_2\text{H}_5\text{OH}$ (d) $\text{C}_6\text{H}_{12}\text{O}_6$
9. A 0.10 M solution of a monoprotic acid ($d = 1.01\text{ g/cm}^3$) is 5% dissociated. What is the freezing point of the solution? The mol. mass of the acid is 300 and $K_f(\text{H}_2\text{O}) = 1.86^\circ\text{C/m}$:
 (a) -0.189°C (b) -0.194°C (c) -0.199°C (d) None of these
10. An aqueous solution boils at 101°C . What is the freezing point of the same solution? (Given: $K_f = 1.86^\circ\text{C/m}$ and $K_b = 0.51^\circ\text{C/m}$)
 (a) 3.647°C (b) -3.647°C (c) -0.364°C (d) None of these
11. An industrial waste water is found to contain 8.2% Na_3PO_4 and 12% MgSO_4 by mass in solution. If % ionisation of Na_3PO_4 and MgSO_4 are 50 and 60 respectively then its normal boiling point is [$K_b(\text{H}_2\text{O}) = 0.50\text{ K kg mol}^{-1}$]:
 (a) 102.3°C (b) 103.35°C (c) 101.785°C (d) None of these
12. Ratio of $\frac{T_b}{K_b}$ of 10 g AB_2 and 14 g A_2B per 100 g of solvent in their respective, solution (AB_2 and A_2B both are non-electrolytes) is 1 mol/kg in both cases. Hence, atomic wt. of A and B are respectively:
 (a) 100, 40 (b) 60, 20 (c) 20, 60 (d) None of these
13. The freezing point of a solution containing 0.2 g of acetic acid in 20.0 g benzene is lowered by 0.45°C . Calculate the degree of dimerization of acetic acid in benzene. K_f for benzene is $5.12\text{ K mol}^{-1}\text{ kg}$:
 (a) 0.527 (b) 0.80 (c) 0.945 (d) None of these
14. If boiling point of an aqueous solution is 100.1°C , what is its freezing point? Given, enthalpy of fusion and vaporisation of water are 80 cal g^{-1} and 540 cal g^{-1} respectively:
 (a) 0.361°C (b) -0.361°C (c) -3.61°C (d) None of these
15. 100 g of $\text{C}_6\text{H}_{12}\text{O}_6$ (aq.) solution has vapour pressure is equal to 40 torr at certain temperature. Vapour pressure of $\text{H}_2\text{O}(l)$ is 40.18 torr at same temperature. If this solution is cooled to -0.93°C , what mass of ice will be separated out? ($K_f = 1.86\text{ K kg mol}^{-1}$)
 (a) 95.5 g (b) 4.5 g (c) 45.5 g (d) 47.8 g
16. 1.0 g of a monobasic acid HA in 100 g water lowers the freezing point by 0.155 K. If 0.75 g, of same acid requires 25 mL of $N/5$ NaOH solution for complete neutralisation then %, degree of ionization of acid is (K_f of $\text{H}_2\text{O} = 1.86\text{ K kg mol}^{-1}$):
 (a) 20% (b) 25% (c) 40% (d) 50%
17. 0.1 M KI and 0.2 M AgNO_3 are mixed in 3 : 1 volume ratio. The depression of freezing point of the resulting solution will be [$K_f(\text{H}_2\text{O}) = 1.86\text{ K kg mol}^{-1}$]:
 (a) 3.72 K (b) 1.86 K (c) 0.93 K (d) 0.279 K

18. If $0.1 \text{ M H}_2\text{SO}_4(\text{aq.})$ solution shows freezing point -0.3906°C then what is the K_{a_2} for H_2SO_4 ?
(Assume $m \approx M$ and $K_{f(\text{H}_2\text{O})} = 1.86 \text{ K kg mol}^{-1}$)
(a) 0.122 (b) 0.0122 (c) 1.11×10^{-3} (d) None of these
19. A living cell contains a solution which is isotonic with 0.2 M glucose solution. What osmotic pressure develops when the cell is placed in 0.05 M BaCl_2 solution at 300 K ?
(a) 1.23 atm (b) 3.69 atm (c) 6.15 atm (d) None of these
20. What is the osmotic pressure of $0.2 \text{ M HX}(\text{aq.})$ solution at 300 K ?
(Given : $K_{a(\text{HX})} = 8 \times 10^{-5}$)
(a) 4.926 atm (b) 0.5024 atm (c) 5.024 atm (d) None of these
21. A solution contain 8 g of a carbohydrate in 100 g of water has a density 1.025 g/mL and an osmotic pressure of 5 atm at 27°C . What is the molar mass of the carbohydrate?
(a) 387 (b) 374 (c) 3740 (d) None of these
22. Study the following figure and choose the correct options. Assuming complete dissociation of electrolyte:




- (a) There will be no net movement of any substance across the membrane
(b) MgCl_2 will flow towards the $\text{Al}_2(\text{SO}_4)_3$ solution
(c) $\text{Al}_2(\text{SO}_4)_3$ will flow towards the MgCl_2 solution
(d) The (osmotic pressure) of 0.1 M MgCl_2 is higher than the of $0.05 \text{ M Al}_2(\text{SO}_4)_3$
23. The total vapour pressure of a 4 mole % solution of NH_3 in water at 293 K is 50.0 torr. The vapour pressure of pure water is 17.0 torr at this temperature. Applying Henry's and Raoult's laws, calculate the total vapour pressure for a 5 mole % solution:
(a) 58.25 torr (b) 33 torr (c) 42.1 torr (d) 52.25 torr
- 24.-25. The vapour pressure of two pure liquids A and B which form an ideal solution are 500 and 800 torr respectively at 300 K . A liquid solution of A and B for which the mole fraction of A is 0.60 is contained in a cylinder closed by a piston on which the pressure can be varied. The solution is slowly vaporized at 300 K by decreasing the applied pressure.
24. The composition of vapour when first bubble formed is:
(a) $y_A = 0.6; y_B = 0.4$ (b) $y_A = 0.48; y_B = 0.52$
(c) $y_A = 0.52; y_B = 0.48$ (d) $y_A = 0.5; y_B = 0.5$
25. What is the composition of last droplet of liquid remaining in equilibrium with vapour?
(a) $x_A = 0.6; x_B = 0.4$ (b) $x_A = 0.5; x_B = 0.5$
(c) $x_A = 0.7; x_B = 0.3$ (d) $x_A = 0.3; x_B = 0.7$

Level 3

Passage 1

Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighted separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and the solvent is kept constant.



- Loss in mass of solvent (w_{II}) will be proportional to:
 - $P - P$
 - $P - P$
 - $\frac{P}{P}$
 - $P - P$
- Gain in mass of anhydrous CaCl_2 is proportional to:
 - P
 - P
 - $P - P$
 - $P - P$
- $\frac{P - P}{P}$ is equal to:
 - $\frac{w_I}{w_I - w_{II}}$
 - $\frac{w_{II}}{w_I - w_{II}}$
 - $\frac{w_I}{w_I - w_{II}}$
 - $\frac{w_{II}}{w_I}$
- Dry air was passed through 9.24 g of solute in 108 g of water and then through pure water. The loss in mass of solution was 3.2 g and that of pure water was 0.08 g. The molecular mass (g/mol) of solute is nearly :
 - 50
 - 62
 - 70
 - 80

Passage 2

A dilute solution contains 'x' moles of solute A in 1 kg of solvent with molal elevation constant K_b . The solute dimerises in the solution according to the following equation. The degree of association is () :



- The van't Hoff factor will be:
 - $i = 1 - \frac{2}{2}$
 - $i = 1 - \frac{2}{2}$
 - $i = 1 - \frac{2}{2}$
 - $i = 1$
- The molecular mass observed will be:
 - greater than actual molecular mass
 - lesser than actual molecular mass
 - equal to the actual molecular mass
 - cannot be predicted by the data given

3. The degree of association is equal to:

- (a) $\frac{(K_b x \quad T_b)}{T_b \cdot 2}$ (b) $\frac{2(K_b x \quad T_b)}{K_b x}$ (c) $2 \frac{2 \quad T_b}{K_b x}$ (d) $\frac{T_b}{2K_b x}$

One or More Answers is/are Correct



- Which of the following statement(s) is/are correct, if intermolecular forces in liquids A, B and C are in the order of A > B > C?
 - B evaporates more readily than A
 - B evaporates more readily than C
 - A evaporates more readily than C
 - all evaporate at same rate at same temperature.
- When non-volatile solute is added to a pure solvent, the:
 - vapour pressure of the solution becomes lower than the vapour pressure of the pure solvent
 - rate of evaporation of solvent is reduced
 - solute does not affect the rate of condensation
 - none of these
- The total vapour pressure of a binary solution is given by

$$P = (100 X_A + 260 X_B) \text{ mm Hg}$$
 where, X_A and X_B are the mole fractions of components A and B. This indicates that the:
 - vapour pressure of solution is less than the pure B component
 - vapour pressure of solution is more than that of pure A component
 - vapour pressure of pure A is 100 mm Hg and that of pure B is 260 mm Hg
 - the vapour pressure of pure A and B are 260 mm Hg and 100 mm Hg respectively
- Which of the following is correct for an ideal solution?
 - $H_{\text{mix}} = 0$ and $V_{\text{mix}} = 0$
 - $V_{\text{mix}} = 0$ and $S_{\text{mix}} = 0$
 - $H_{\text{mix}} = 0$ and $S_{\text{mix}} = 0$
 - $G_{\text{mix}} = 0$ and $S_{\text{mix}} = 0$
- For a non-ideal solution with a negative deviation:
 - H_{mix} ve
 - V_{mix} ve
 - S_{mix} ve
 - G_{mix} ve
- A binary solution of liquids A and B will show positive deviation from Raoult's law if it fulfils the following condition:
 - $P_A < X_A P_A$ and $P_B < X_B P_B$
 - The intermolecular forces of A < B < A, B < B
 - H mixing is positive
 - V mixing is negative
- Which of the following statements is/are correct about acetone and trichloromethane mixture?
 - Mixtures of acetone and trichloromethane shows positive deviation from Raoult's law
 - The forces of attraction acting between molecules of acetone and trichloromethane in a mixture are stronger than those acting between the molecules in pure acetone
 - Pure acetone can be obtained by the careful fractional distillation of any mixture of acetone and trichloromethane
 - When acetone and trichloromethane are mixed, the enthalpy change is negative

8. The azeotropic solutions of two miscible liquids:
(a) can be separated by simple distillation
(b) may show positive or negative deviation from Raoult's law
(c) are supersaturated solution
(d) behave like a single component and boil at a constant temperature
9. For exact determination of molecular mass through colligative properties measurement:
(a) solute must be volatile
(b) solution must be very dilute
(c) solution must be formed by similar nature of substances
(d) solute must not be dissociated or associated
10. For the depression in freezing point experiment, the correct statement(s) is/are:
(a) vapour pressure of pure solvent is more than that of solution
(b) vapour pressure of pure solvent is less than that of solution
(c) only solute molecules solidify at the freezing point
(d) only solvent molecules solidify at the freezing point
11. The cryoscopic constant value depends upon:
(a) the molar mass of the solute in the solution
(b) the molar mass of the solvent in the solution
(c) the enthalpy of fusion of the solvent
(d) the freezing point of the solvent
12. Consider 0.1 M solutions of two solutes X and Y. The solute X behaves as univalent electrolyte, while the solute Y dimerises in solution. Select correct statement(s) regarding these solutions:
(a) The boiling point of solution of 'X' will be higher than that of 'Y'
(b) The osmotic pressure of solution of 'Y' will be lower than that of 'X'
(c) The freezing point of solution of 'X' will be lower than that of 'Y'
(d) The relative lowering of vapour pressure of both the solution will be the same
13. Consider following solutions:
(I) 1 M glucose (aq) (II) 1 M sodium chloride (aq)
(III) 1 M acetic acid in benzene (IV) 1 M ammonium phosphate (aq)
(a) all are isotonic solutions (b) III is hypotonic of I, II, IV
(c) I, II, IV are hypertonic of III (d) IV is hypertonic of I, II, III
14. Which of the following statements is (are) incorrect?
(a) 0.1 M KCl solution will have the same osmotic pressure as 0.1 M glucose solution
(b) 0.1 M KCl solution will have the same boiling point as 0.1 M urea solution
(c) 0.1 m glucose and 0.1 m urea are isotonic
(d) 0.1 m MgCl_2 solution will have less relative lowering of vapour pressure than 0.1 m NaCl
15. Consider following solutions:
0.1 m $\text{C}_6\text{H}_5\text{NH}_3^+\text{Cl}$; 0.1 m KCl; 0.1 m Glucose; 0.1 m $\text{Na}_2\text{C}_2\text{O}_4 \cdot 10\text{H}_2\text{O}$
(a) the solution with highest boiling point is 0.1 m $\text{Na}_2\text{C}_2\text{O}_4 \cdot 10\text{H}_2\text{O}$
(b) the solution with highest freezing point is 0.1 m glucose
(c) 0.1 m $\text{C}_6\text{H}_5\text{NH}_3\text{Cl}$ and 0.1 m NaCl will have the same osmotic pressure
(d) 0.1 m glucose solution will have the lowest osmotic pressure

Match the Column



Column-I and **Column-II** contains four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I may have the matching with the same entries of column-II.

- | Column-I | Column-II |
|--|---|
| (A) <i>n</i> -hexane + <i>n</i> -heptane | (P) Can be separated by fractional distillation |
| (B) Acetone + chloroform | (Q) Maximum boiling azeotrope |
| (C) Chlorobenzene and bromobenzene | (R) Cannot be separated by fractional distillation completely |
| (D) Ethanol + water | (S) Minimum boiling azeotrope |

- | Column-I | Column-II |
|---|---|
| (A) $_1 : 0.1\text{ M}$ glucose; $_2 : 0.1\text{ M}$ urea | (P) $_1$ and $_2$ are isotonic |
| (B) $_1 : 0.1\text{ M}$ NaCl; $_2 : 0.1\text{ M}$ Na ₂ SO ₄ | (Q) No net migration of solvent across the membrane |
| (C) $_1 : 0.1\text{ M}$ NaCl; $_2 : 0.1\text{ M}$ KCl | (R) $_1$ is hypertonic to $_2$ |
| (D) $_1 : 0.1\text{ M}$ CuSO ₄ ; $_2 : 0.1\text{ M}$ sucrose | (S) $_1$ is hypotonic to $_2$ |

- | Column-I | Column-II |
|--|---|
| (A) 0.1 M Al ₂ (SO ₄) ₃ | (P) Solution with highest boiling point |
| (B) 0.1 M AlPO ₄ | (Q) van't Hoff factor is greater than 1 |
| (C) 0.1 M urea | (R) Solution with lowest osmotic pressure |
| (D) 0.1 M MgCl ₂ | (S) Solution with lowest freezing point |

- | Column-I | Column-II |
|---|---------------------------|
| (Solute) | (Van't Hoff factor, i) |
| (A) AlCl ₃ if $_0$ 0.8 | (P) i 3.4 |
| (B) BaCl ₂ if $_0$ 0.9 | (Q) i 2.8 |
| (C) Na ₃ PO ₄ if $_0$ 0.9 | (R) i 3.8 |
| (D) K ₄ [Fe(CN) ₆] if $_0$ 0.7 | (S) i 3.7 |

- | Column-I | Column-II |
|-------------------------------|-------------------------------|
| (A) Elevation of B.P. | (P) Colligative property |
| (B) Osmotic pressure | (Q) Ebullioscopic constant |
| (C) Relative lowering in V.P. | (R) Berkeley-Heartley method |
| (D) Depression of F.P. | (S) Ostwald and Walker method |

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statement carefully and mark the correct answer according to the instructions given below:

- (A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** An increase in surface area increases the rate of evaporation.
STATEMENT-2 : Stronger the intermolecular attraction forces, faster is the rate of evaporation at a given temperature.
2. **STATEMENT-1 :** An ideal solution obeys Raoult's law.
STATEMENT-2 : In an ideal solution, solute-solute as well as solvent-solvent, interactions are similar to solute-solvent interactions.
3. **STATEMENT-1 :** If a liquid solute more volatile than the solvent is added to the solvent, the vapour pressure of the solution is greater than vapour pressure of pure solvent.
STATEMENT-2 : Vapour pressure of solution is equal to vapour pressure of solvent.
4. **STATEMENT-1 :** V_{mix} and S_{mix} for an ideal solution is zero.
STATEMENT-2 : A...B interaction in an ideal solution are same as between A... A and B...B.
5. **STATEMENT-1 :** Elevation in boiling point will be high if the molal elevation constant of the liquid is high.
STATEMENT-2 : Elevation in boiling point is a colligative property.
6. **STATEMENT-1 :** The boiling point of 0.1 M urea solution is less than that of 0.1 M KCl solution.
STATEMENT-2 : Elevation of boiling point is directly proportional to the number of moles of non-volatile solute particles present in the solution.
7. **STATEMENT-1 :** The observed molar mass of acetic acid in benzene is more than the normal molar mass of acetic acid.
STATEMENT-2 : Molecules of acetic acid dimerise in benzene due to hydrogen bonding.
8. **STATEMENT-1 :** Addition of ethylene glycol to water lowers the freezing point of water, therefore, used as antifreeze substance.
STATEMENT-2 : Ethylene glycol is soluble in water.
9. **STATEMENT-1 :** Osmotic pressure is a colligative property.
STATEMENT-2 : Osmotic pressure is developed in a column due to osmosis.
10. **STATEMENT-1 :** Osmosis involves movement of solvent molecules from lower concentration to higher concentration.
STATEMENT-2 : Solutions having the same osmotic pressure are called isotonic solutions.

- 11. STATEMENT-1 :** Isotonic solutions must have the same molar concentration.
STATEMENT-2 : Solutions which have the same osmotic pressure are known as isotonic solution.
- 12. STATEMENT-1 :** Isotonic solutions do not show phenomenon of osmosis.
STATEMENT-2 : Isotonic solutions have same molar concentration at same temperature.
- 13. STATEMENT-1 :** When dried fruits and vegetables are placed in water, they slowly get swollen.
STATEMENT-2 : It happens due to the phenomenon of osmosis.
- 14. STATEMENT-1 :** Reverse osmosis is used to purify sea water.
STATEMENT-2 : Solvent molecules pass from concentrate solution to pure solvent through semipermeable membrane if high pressure () is applied on solution side.
- 15. STATEMENT-1 :** All solutes becomes more soluble in water at higher temperature.
STATEMENT-2 : Solubility of solute depends upon temperature.
- 16. STATEMENT-1 :** Henry's law is always applicable for gases.
STATEMENT-2 : Raoult's law is a special case of Henry's law.
- 17. STATEMENT-1 :** Increasing pressure on pure water decrease its freezing point.
STATEMENT-2 : Density of water is maximum at 273 K.
- 18. STATEMENT-1 :** The molecular mass of acetic acid determined by depression in freezing point method in benzene and water was found to be different.
STATEMENT-2 : Water is polar and benzene is non-polar.
- 19. STATEMENT-1 :** If red blood cells were removed from the body and placed in pure water, pressure inside the cell increases.
STATEMENT-2 : The concentration of the salt content in the cells increases.
- 20. STATEMENT-1 :** Azeotrope is a binary mixture formed by ideal solutions.
STATEMENT-2 : Azeotrope boils with unchanged composition.

Subjective Problems



- The vapour pressure of two pure liquids A and B are 5 and 10 torr respectively. Calculate the total pressure of the solution (in torr) obtained by mixing 2 mole of A and 3 mole of B.
- The vapour pressure of two pure liquids A and B are 50 and 40 torr respectively. If 8 moles of A is mixed with x moles of B, then vapour pressure of solution obtained is 48 torr. What is the value of x.
- The vapour pressure of a liquid solution containing A and B is 99 torr. Calculate mole % of B in vapour phase.
(Given : P_A 100 torr ; P_B 80 torr)
- If 30 g of a solute of molecular mass 154 is dissolved in 250 g of benzene. What will be the elevation in boiling point of the resulting solution ?
(Given : $K_{b(C_6H_6)}$ $2.6 K kg mol^{-1}$)

5. Calculate elevation in boiling point for 2 molal aqueous solution of glucose.
(Given : $K_{b(\text{H}_2\text{O})}$ $0.5 \text{ K kg mol}^{-1}$)
6. Calculate depression of freezing point for 0.56 molal aq. solution of KCl.
(Given : $K_{f(\text{H}_2\text{O})}$ $1.8 \text{ K kg mol}^{-1}$).
7. What is the maximum value of van't Hoff factor for AlCl_3 ?
8. A solution containing 500 g of a protein per litre is isotonic with a solution containing 3.42 g sucrose per litre. The molecular mass of protein is 5×10^x , hence x is.
9. An aqueous solution of urea has a freezing point of -0.515°C . Predict the osmotic pressure (in atm) of the same solution at 37°C .
(Given : $K_{f(\text{H}_2\text{O})}$ $1.86 \text{ K kg mol}^{-1}$)
10. 0.2 M aq. solution of KCl is isotonic with 0.2 M K_2SO_4 at same temperature. What is the van't Hoff factor of K_2SO_4 ?

ANSWERS

Level 1

1. (d)	2. (b)	3. (a)	4. (c)	5. (b)	6. (c)	7. (a)	8. (b)	9. (d)	10. (b)
11. (d)	12. (c)	13. (c)	14. (c)	15. (b)	16. (b)	17. (c)	18. (a)	19. (a)	20. (a)
21. (a)	22. (d)	23. (a)	24. (d)	25. (d)	26. (a)	27. (c)	28. (b)	29. (a)	30. (a)
31. (b)	32. (d)	33. (b)	34. (d)	35. (d)	36. (c)	37. (b)	38. (d)	39. (c)	40. (d)
41. (a)	42. (a)	43. (c)	44. (a)	45. (c)	46. (a)	47. (d)	48. (c)	49. (c)	50. (a)
51. (d)	52. (d)	53. (c)	54. (d)	55. (b)	56. (c)	57. (a)	58. (b)	59. (d)	60. (b)
61. (b)	62. (d)	63. (a)	64. (d)	65. (b)	66. (b)	67. (d)	68. (b)	69. (b)	70. (a)
71. (d)	72. (c)	73. (b)	74. (d)	75. (a)	76. (b)	77. (a)	78. (b)	79. (c)	80. (b)
81. (a)	82. (b)	83. (c)	84. (a)	85. (b)	86. (d)	87. (d)	88. (c)	89. (a)	90. (d)
91. (d)	92. (d)	93. (b)	94. (c)	95. (d)	96. (b)	97. (b)	98. (a)	99. (a)	100. (a)
101. (a)	102. (b)	103. (b)	104. (c)	105. (d)	106. (c)	107. (a)	108. (c)	109. (b)	110. (a)
111. (c)	112. (a)	113. (b)	114. (a)	115. (d)	116. (b)	117. (c)	118. (d)	119. (d)	120. (a)
121. (b)	122. (b)	123. (a)	124. (d)	125. (a)					

Level 2

1. (b)	2. (b)	3. (d)	4. (c)	5. (c)	6. (c)	7. (d)	8. (d)	9. (c)	10. (b)
11. (c)	12. (b)	13. (c)	14. (b)	15. (d)	16. (b)	17. (d)	18. (b)	19. (a)	20. (c)
21. (b)	22. (d)	23. (a)	24. (b)	25. (c)					



Level 3

Passage-1: 1. (a) 2. (b) 3. (b) 4. (b)

Passage-2: 1. (b) 2. (a) 3. (b)

One or More Answers is/are Correct

1. (b,c) 2. (a,b,c) 3. (a,b,c) 4. (a,b,d) 5. (a,b,d) 6. (a,b,c) 7. (b,d) 8. (b,d)
9. (b,d) 10. (a,d) 11. (b,c,d) 12. (a,b,c) 13. (b,c,d) 14. (a,b,d) 15. (a,b,c,d)

Match the Column

- | | | | |
|---------------|---------|---------|--------|
| 1. A P; | B Q, R; | C P; | D R, S |
| 2. A P, Q; | B S; | C P, Q; | D R |
| 3. A P, Q, S; | B Q; | C R; | D Q |
| 4. A P; | B Q; | C S; | D R |
| 5. A P, Q; | B P, R; | C P, S; | D P |

Assertion-Reason Type Questions

- | | | | | | | | | | |
|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1. (C) | 2. (A) | 3. (C) | 4. (D) | 5. (B) | 6. (A) | 7. (A) | 8. (B) | 9. (B) | 10. (B) |
| 11. (D) | 12. (A) | 13. (A) | 14. (B) | 15. (D) | 16. (D) | 17. (C) | 18. (A) | 19. (C) | 20. (D) |

Subjective Problems

- | | | | | | | | | | |
|--------|------|------|------|------|------|------|------|------|-------|
| 1. (8) | 2. 2 | 3. 4 | 4. 2 | 5. 1 | 6. 2 | 7. 4 | 8. 4 | 9. 7 | 10. 2 |
|--------|------|------|------|------|------|------|------|------|-------|

Hints and Solutions

Level 1

1. (d) $x(l) \xrightleftharpoons{Eq^m} x(g)$, $K_p = \frac{P_p \text{ of liq.}}{P^\circ}$ ve T, V_p

2. (b) Given $P_1 = 19 \text{ mm Hg}$, $P_2 = 760 \text{ mm Hg}$;
 $H_{\text{vap.}} = 40670 \text{ J/mol}$

Applying Clausius-Clapeyron's equation

$$\log \frac{P_2}{P_1} = \frac{H_{\text{vap}}}{2.303 R} \left(\frac{T_2}{T_1 T_2} \right)$$

or $\log \frac{760}{19} = \frac{40670}{2.303 \times 8.3} \left(\frac{T_1}{T_1 T_2} \right)$

on solving, we get $T_1 = 291.4 \text{ K}$

3. (a) Initial mass of $\text{H}_2\text{O}(l) = 18 \text{ g}$;

$$n_{\text{H}_2\text{O}}(g) = \frac{\frac{24.63}{760} \times 7.6}{0.0821 \times 300} = 0.01$$

% wt. of H_2O vaporised $= \frac{0.01 \times 18}{18} \times 100 = 1\%$

4. (c) Raoult's law is applicable for ideal solution.

5. (b) $P = P_A^\circ x_A + P_B^\circ x_B = P_B^\circ x_A (P_A^\circ / P_B^\circ + 1)$
 $[\because x_B = 1 - x_A]$

6. (c) $y_A = \frac{P_A^\circ x_A}{P_A^\circ x_A + P_B^\circ x_B} = \frac{x_A}{x_A + x_B \frac{P_B^\circ}{P_A^\circ}}$
 $\frac{y_A}{x_A} = \frac{1}{x_A + x_B \frac{P_B^\circ}{P_A^\circ}} = 1 \quad [\because P_A^\circ = P_B^\circ]$
 $\dots(1)$

Similarly, $y_B = \frac{P_B^\circ x_B}{P_A^\circ x_A + P_B^\circ x_B} = \frac{x_B}{x_B + x_A \frac{P_A^\circ}{P_B^\circ}}$
 $\dots(2)$

$$\frac{y_B}{y_A} = \frac{x_B}{x_A}$$

7. (a) $y_A = \frac{P_A^\circ x_A}{P_A^\circ x_A + P_B^\circ x_B} = \frac{P_A^\circ x_A}{P_B^\circ (P_A^\circ / P_B^\circ + 1) x_A}$
 $\frac{1}{y_A} = \frac{P_B^\circ x_B + P_A^\circ x_A}{P_A^\circ x_A} = 1 + \frac{P_B^\circ}{P_A^\circ} \frac{1}{x_A}$
 $\frac{1}{y_A} = 1 + \frac{P_B^\circ}{P_A^\circ} \frac{1}{x_A}$
 Slope $= \frac{P_B^\circ}{P_A^\circ}$

8. (b) $P_{\text{Solution}} = \frac{P_{\text{Solution}}^\circ X_{\text{Solvent}}}{P} = \frac{P}{X_{\text{Solute}}}$

9. (d) Solution X is unsaturated so v.p. will be more, solution Y and Z are saturated so v.p. of Y = v.p. of Z and 2 gm of solute would be present in form of solid in system Z.

10. (b) Highest vapour pressure for lowest boiling point.

11. (d) $\frac{P}{P} = x_{\text{solute}} \text{ or } \frac{P}{P} = \frac{n}{n + N}$
 $\frac{0.1}{0.1 + 9.9} = 0.01; P = 0.99 P$

12. (c) $80 P (0.8) = P \times 100 \text{ torr}$.

13. (c) Given $P_A = 750 \text{ mm Hg}$
 $\therefore 373 \text{ K}$ is boiling point of water.

Thus, $P_A = 760 \text{ mm Hg}$

$$m = \frac{P}{P} = \frac{1000}{M_{\text{solvent}}} = \frac{10}{750} \times \frac{1000}{18} = 0.74$$

14. (c) Normal boiling point of water is 100°C , hence

$P(\text{H}_2\text{O}) = 760 \text{ torr}$; $M(\text{H}_2\text{O}) = 18 \text{ g/mol}$

$$m = \frac{P}{P} = \frac{1000}{M_{\text{H}_2\text{O}}}$$

$P = 746.5$; $P = 13.45 \text{ torr}$

15. (b) $P_s = \frac{80}{100} P$, $w = ?$,

$M_{\text{solute}} = 40$, $w = 114 \text{ g}$, $M_{\text{solvent}} = 114$

$$\therefore \frac{P}{P_s} \frac{P_s}{M_{\text{solute}}} \frac{w}{M_{\text{solvent}}} \frac{M_{\text{solvent}}}{W}$$

$$\text{or } \frac{P}{(80/100)P} \frac{(80/100)P}{40} \frac{w}{57} = \frac{114}{57}$$

$$16. (b) \frac{P}{P} \frac{P}{n} \frac{w}{N} \frac{5 \text{ g}}{N} \quad (\text{for very dilute solution})$$

$$2 \frac{n}{N_A} \frac{n}{N_B} \frac{M_B}{2M_A}$$

17. (c) We know that

$$Y_A \frac{P_A X_A}{P_{\text{total}}} \text{ or } \frac{Y_A}{X_A} \frac{P_A}{P_{\text{total}}} \\ \therefore P_A \frac{P_A}{P_{\text{total}}} \text{ so } \frac{Y_A}{X_A} = 1 \text{ or } Y_A = X_A$$

$$18. (a) \text{ Moles of A } \frac{20}{40} = 0.5; \\ \text{moles of B } \frac{20}{80} = 0.25 \\ x_A = \frac{0.5}{0.5 + 0.25} = 0.67; \quad x_B = 0.33$$

$$\frac{P_{\text{total}}}{100} \frac{P_A X_A}{0.67} \frac{P_B X_B}{40} \frac{0.33}{80} = 80 \text{ torr}$$

19. (a) Since, $P = X_A P_A + X_B P_B$, we have

$$\frac{1}{1} \frac{P_A}{3} \frac{3}{1} \frac{P_B}{3} = 550 \text{ mm Hg};$$

$$\frac{1}{1} \frac{P_A}{4} \frac{4}{1} \frac{P_B}{4} = 560 \text{ mm Hg}$$

$$\text{That is, } 0.25 P_A + 0.75 P_B = 550 \text{ mm Hg};$$

$$0.20 P_A + 0.8 P_B = 560 \text{ mm Hg}$$

Solving for P_A and P_B , we get;

$$P_A = 400 \text{ mm Hg and } P_B = 600 \text{ mm Hg}$$

$$20. (a) y_A \frac{P_A}{P} \frac{P_A X_A}{P} \text{ and } y_B \frac{P_B X_B}{P};$$


$$\frac{y_A}{y_B} \frac{P_A}{P_B} \frac{x_A}{x_B} = \frac{4}{3} \frac{1}{3} \frac{x_A}{(1-x_A)}$$

$$x_A = \frac{4}{5} \text{ or } x_B = \frac{1}{5}$$

$$21. (a) x_A = \frac{1}{1+3} = \frac{1}{4}; \quad x_B = \frac{3}{4};$$

$$\frac{y_A}{y_B} \frac{P_A}{P_B} \frac{x_A}{x_B} = \frac{1}{3} \frac{1}{3} \frac{1}{3} \\ \frac{1}{9} \text{ or } y_A = \frac{1}{10}$$

22. (d)



Plots AD show vapour pressure of B.

Plots BC show vapour pressure of A.

Plot CD show vapour pressure of liquid solution containing A and B volatile components.

$$\frac{P}{EH} \frac{P_A}{EF} \frac{P_B}{EG}$$

$$23. (a) \therefore (T_b)_{\text{Solvent}} (T_b)_{\text{Solution}}; \\ S_{\text{Solv.}} S_{\text{Solution}}$$

(H_{Solution}) (H_{Solvent})
[Only solvent participate in vaporisation]
[Due to elevation in BP of solution]

$$24. (d) (fS)_{\text{Solution}} (fS)_{\text{Solvent}} \\ \text{[Since solution has more randomness than solvent]}$$

(fH)_{Solution} (fH)_{Solvent}
[Similar forces of intermolecular forces are involved in solution and solvent]

Due to presence of non-volatile solute (Sugar), freezing point of solution decrease.


$$25. (d) (H_{\text{Vap}})_{\text{Solvent}} (H_{\text{Vap}})_{\text{Solution}} \\ \text{[Because similar intermolecular forces are involved in both cases].}$$

$$\therefore (T_b)_{\text{Solvent}} (T_b)_{\text{Solution}} \\ (S_{\text{Vap}})_{\text{Solution}} (S_{\text{Vap}})_{\text{Solvent}}$$

26. (a) Elevation in boiling point concentration of a solution. Thus, the order of concentration of solutions is I < II < III.

27. (c) T_B° T_A° [From diagram] B Residue
A Distillate
28. (b) The mixture will show positive deviation from Raoult's law hence boiling point of an azeotropic mixture of water-ethanol is less than that of both water and ethanol.
29. (a) H_{Solution} H_1 H_2 H_3
30. (a) $P_{\text{Ideal solution}}$ $\frac{1}{3}$ 150 $\frac{2}{3}$ 300 250
 $\therefore P_{\text{Solution}} > P_{\text{Ideal solution}}$; () ve deviation.
31. (b) When A-B interaction is weaker than A-A and B-B interaction, then mixture of A and B show positive deviation.
32. (d) On changing external pressure, composition of azeotrope alters.
33. (b) After mixing HNO_3 and H_2O , intermolecular force of attraction increase.
34. (d) $\therefore$
 $P_{\text{Ideal solution}}$ $\frac{1}{4}$ 100 80 $\frac{3}{4}$ 85 mmHg
 P_{Solution} (90 mm Hg)
Solution show positive deviation from Raoult's law.
35. (d) From an azeotropic mixture the components cannot be separated at boiling point because it is a constant temperature boiling mixture.
36. (c) See solution of Q. 35.
37. (b) Show negative deviation from Raoult's law.
38. (d) $P = P_A^\circ + P_B^\circ = 300$ torr.
39. (c) For two immiscible liquid;
 P_A P_{total} $P_{\text{H}_2\text{O}}$ 748 648 100
 $\frac{W_A}{W_B} = \frac{P_A M_A}{P_B M_B}$; $M_A = \frac{1.25}{1} \cdot \frac{648}{100} = 145.8$
40. (d) $\frac{W_A}{W_B} = \frac{P_A}{P_B} \cdot \frac{M_A}{M_B} = \frac{0.7}{7} \cdot \frac{112.5}{18} = 0.625$
 $\% \frac{W_A}{W_B} = 100 \cdot \frac{0.625}{1.625} = 38.46$
41. (a) Lowering in vapour pressure is a colligative property.
42. (a) $i = 1 - (n-1) \cdot \frac{1}{n}$ so $n = \frac{1}{1-i}$
43. (c) $i = \frac{C(1-\alpha)}{C} = \frac{1-\alpha}{n}$ $i = 1 - \frac{1}{n}$
44. (a) For dissociation $i = 1 + (n-1)\alpha$ $i = 1 + \frac{1}{n}$
For dissociation $i = 1 + \frac{1}{n}$ $i = 1 + \frac{1}{n}$
45. (c) For high v.p. concentration of solute should be low and temperature should be high.
46. (a) On addition of strong electrolyte, vapour pressure decrease. On addition of water, vapour pressure increases due to increase in X_{Solvent} .
47. (d) Colligative property $\propto$ moles of non-volatile solute.
48. (c) $\frac{P}{P} = \frac{10}{50} = \frac{n}{n-12}$ $n = 3$
(Number of ions of solute), moles of $\text{K}_2\text{SO}_4 = 1$
49. (c) $m_i = \frac{P}{P} \cdot \frac{1000}{M_{\text{solvent}}}$;
 $7m = \frac{(20.0126 - 20) \cdot 1000}{20 \cdot 18}$;
 $m = 5 \cdot 10^{-3}$
50. (a) T_b 100.5 100 K_b 1 K_b 0.5
51. (d) $T_b = K_b \cdot m = 3.63 \cdot \frac{0.616/154}{15} = 1000$;
 $T_b = 61.7$ 0.968
62.67°C
52. (d) $T_b = 80.26$, $T_b = 0.16$;
0.16 2.53 $\frac{0.26/M}{11.20} = 1000$; $M = 367$
that is almost molar mass of $\text{C}_{20}\text{H}_{16}\text{Fe}_2$.
53. (c) 80.23 80.10 2.53 $\frac{0.64/x}{100} = 10^3$
 $x = 125$
54. (d) T_b (normal) $K_f m = 2.53 \cdot 1 = 2.53$ K;
 $i = \frac{T_{b(\text{obs})}}{T_{b(\text{nor})}} = \frac{1.518}{2.53} = 0.6$
 $i = 1 - \frac{1}{n}$; $n = 0.8$
55. (b) $H_{\text{vap.}} = \frac{M_{\text{solvent}} \cdot R \cdot T_b^2(\text{solvent})}{1000 \cdot K_b}$;
 $H_{\text{vap.}} = \frac{92 \cdot 8.314 \cdot (383.7)^2}{1000 \cdot 3.32} = 33.91$ kJ/mol $= 34$ kJ/mol

56. (c) B.P. moles of non-volatile solute.



58. (b) $T_b \quad K_b \quad m$

$m = \frac{y}{x} \dots (1)$

$T_f \quad K_f m \dots (2)$

$T_f = \frac{y}{x} z$

59. (d) Liquid solution $\rightleftharpoons$ solid solvent.

60. (b) $T_f \quad 2.304; m = \frac{2.304}{14.4} \quad 0.16;$

$m = \frac{n_{\text{solute}}}{W_{\text{solvent}}} \quad 1000$

$0.16 = \frac{2.6}{M} \cdot \frac{1000}{100}; M = 162.5$

61. (b) $n_{C_4H_{10}} = \frac{10.44}{58} \quad 0.18;$

$m = \frac{0.18}{200} \cdot 1000 \quad 0.9$

$T_f = 5.12 \quad 0.9 \quad 4.608;$

$T_f = 5.5 \quad 4.608 \quad 0.892^\circ\text{C}$

62. (d) $4 \text{ F} \quad 20 \text{ C}; \quad T_f \quad 20 \text{ C}$

$T_f \quad K_f \cdot m \quad m = \frac{20}{1.86} \quad 10.75$

mass of C_2H_5OH required

$10.75 \quad 46 \quad 494.5 \text{ g}$

63. (a) $4.4 \quad 5.5 \quad K_f = \frac{2.4/154}{75} \cdot 10^3 \quad K_f = 5.3$

64. (d) $0.465 \quad 1.86 \quad \frac{1.8/x}{40} \cdot 10^3$

$x = 180 \text{ g/mol} \quad C_6H_{12}O_6$

65. (b) $T_f = \frac{H}{S}$

66. (b) $\therefore \frac{H_f}{S_f} \text{ Solution} = \frac{H_f}{S_f} \text{ Solvent}$

67. (d) $T_f \quad K_f \cdot m \quad 0.93 \quad 1.86 \quad \frac{36}{M} \cdot \frac{1}{1.2},$

$M = 60 \text{ and } 30 \quad n = 60; \quad n = 2$

so compound is $C_2H_4O_2$.

68. (b) $0.93 \quad 1.86 \quad \frac{7/x}{93/1000}$

$x = 150.5 \text{ g mol}^{-1}.$

69. (b) It has a very high cryoscopic constant.

70. (a) Maximum ions minimum T_f .

71. (d) Maximum T_f for minimum ions.

72. (c) $3.72 \quad 1.86 [i m]$

$i = 2 \quad [Pt (H_2O)_3Cl_3]Cl \cdot 3H_2O$

73. (b) $T_f \quad K_f \text{ molality } i;$

$0.558 \quad 1.86 \quad 0.1 \quad i \text{ or } i = 3$

Thus complex is $[CoCl_x \cdot xNH_3] \cdot Cl_{2-x}$. Since co-ordination number of CO is six, thus complex is $[Co(NH_3)_5Cl]Cl_2$.

74. (d) Depression in freezing point, $T_f = iK_f m$. The value of van't Hoff factor (i) is minimum for the glucose, which is a non-electrolyte. Hence, aqueous solution of glucose has highest freezing point.

75. (a) Molality of solution should be same $m_1 = m_2;$

$\frac{4}{60} = \frac{1000}{96} = \frac{10}{M_B} = \frac{1000}{90} \text{ or } M_B = 160$

76. (b) $T_f \quad K_f \cdot m \quad 1.86 \quad K_f = 1$

$K_f(H_2O) = 1.86$

for $HCOOH$; $0.02046 \quad 1.86 \quad 0.01(1 \quad);$

$0.1; \text{ so } [H^+] = 10^{-3}; \text{ pH } = 3$

77. (a) The reaction when KI is added in HgI_2

$HgI_2 + 2KI \rightleftharpoons K_2HgI_4 \rightleftharpoons 2K^+ + HgI_4^{2-}$

Since, number of ions decreased, so freezing point increases.

78. (b) $T_f = \frac{mK_f i}{T_b = \frac{mK_b i}{M_{\text{obs}} = \frac{M}{i}}}$

i , So M_{obs} increases

79. (c) To use car; $T_f \quad T_f \quad T_f \quad 9.3 \text{ K}$

$T_f \quad K_f \cdot m \quad 9.3 \quad 1.86 \quad m \quad m = 5$

number of moles of ethylene glycol added to 5 L of water of radiator $5 \quad 5 \quad 25$

or $25 \quad 62 \quad 1550 \text{ g}$

80. (b) Given $K_1 = 1.86 \text{ K kg mol}^{-1}, m = 0.01,$

$T_f = 0.0194^\circ\text{C}$

$T_f = i \cdot K_f m; \quad \frac{0.0194}{1.86 \cdot 0.01} = i$

or $i = 1.043 \text{ or } 0.043$

81. (a) $3.84 \quad i = 5.12 \quad 1$

$i = 0.75 \quad 1$

$$2.53 \quad i \quad 2.53 \quad 1$$

$$i \quad 1$$

$$82. (b) \quad T_f \quad (1 \quad 0.5) \quad 1.86 \quad 0.5 \quad 1.395$$

$$T_f \quad 273 \quad 1.395 \quad 271.60$$

$$83. (c) \quad T_f \quad K_f \cdot m \cdot i;$$

$$0.87 \quad 1.86 \quad i \quad \frac{1}{345.5} \quad \frac{1000}{25}; \quad i \quad 4$$

$$84. (a) \quad T_f \quad K_f \quad \frac{n_1}{W_{\text{solvent}}} \quad \frac{n_2}{1000};$$

$$T_f \quad 1.86 \quad \frac{\frac{3}{60} \quad \frac{1.8}{180}}{95.2} \quad 1000 \quad 1.172$$

$$T_f \quad T_f \quad 1.172 \quad C; \quad T_f \quad 1.172 \quad C$$

$$85. (b) \quad T_f \quad K_f \cdot m \cdot i \quad i \quad 0.633;$$

$$i \quad 1 \quad \frac{2}{2} \quad 0.73$$

$$86. (d) \text{ Maximum freezing point for minimum ion.}$$

$$87. (d) \quad i \quad 1 \quad 1.25$$

$$T_f \quad K_f \cdot m \cdot i \quad 1.86 \quad 0.2 \quad 1.25$$

$$\text{or } T_f \quad 0.465 \quad \text{or } T_f \quad 0.465 \quad C$$

$$88. (c) \quad 0.01 \quad K_b \quad 2 \quad K_b \quad x \quad 1$$

$$x \quad 0.02 \quad M$$

$$89. (a) \quad \frac{P^\circ}{P^\circ} \quad \frac{P}{N} \quad \frac{in}{2} \quad \frac{m}{10^3} \quad \frac{M_{\text{solution}}}{10^3}$$

$$\frac{2 \quad 1 \quad 18}{1000} \quad \frac{9}{250} \quad 0.036$$

$$T_f \quad k_f(m), \quad \frac{T_f}{K_f} \quad 2 \quad im, \quad m \quad 1, \quad i \quad 2$$

$$90. (d) \text{ Depression in freezing point}$$

$$K_f m \text{ (as } K_f, T_f)$$

K_f depends upon the solvent.

$$91. (d) \quad T \quad 100 \quad 0.1(1 \quad 3 \quad 0.9) \quad 0.52$$

$$T \quad 373.19 \quad K$$

$$92. (d) \quad \text{No. of ions for 1 mole solute.}$$

$$93. (b) \text{ Water (solvent) flows from lower concentration to higher concentration.}$$

$$94. (c) \text{ Solvent molecules are moving from solution B into solution A hence, osmotic pressure of solution A immediately begins to decrease.}$$

$$95. (d) \text{ Solutions having same no. of ions are isotonic}$$

$$\text{Urea (i 1), NaCl (i 2), MgCl}_2 \text{ (i 3), Na}_2\text{SO}_4 \text{ (i 3), Ca(NO}_3)_2 \text{ (i 3).}$$

$$96. (b) \quad \text{Glucose} \quad \text{Non-electrolyte}$$

$$0.05 \quad RT \quad \frac{3}{x} \quad RT$$

$$x \quad 60, \quad n \quad \frac{MFM}{EFM} \quad \frac{60}{30} \quad 2$$

$$M.F. \quad (E.F.)_2, \quad C_2H_4O_2$$

$$97. (b) \text{ A semipermeable membrane allow the passage of only solvent molecules.}$$

$$98. (a) \text{ In osmosis, solvent molecules move from lower concentration (higher vapour pressure) to higher concentration (lower vapour pressure)}$$

$$99. (a) \quad i \quad [\text{Equal concentration}]$$

$$\text{AlCl}_3 (i \quad 4), \text{ Urea (i 1), BaCl}_2 (i \quad 3)$$

$$100. (a) \quad i \quad [\text{Equal concentration}]$$

$$\text{Aluminium sulphate } \text{Al}_2(\text{SO}_4)_3 \text{ (i 5),}$$

$$\text{Barium chloride } \text{BaCl}_2 (i \quad 3),$$

$$\text{Sodium sulphate } \text{Na}_2\text{SO}_4 (i \quad 3).$$

$$101. (a) \text{ For } x (i \quad 2) \text{ and } Y (i \quad 1)$$

$$(T_b)_x \quad (T_b)_y; \quad x \quad y; \quad (T_f)_x \quad (T_f)_y$$

$$102. (b) \quad \text{K}_3[\text{Fe}(\text{CN})_6] \rightleftharpoons 3\text{K}^+ + [\text{Fe}(\text{CN})_6]^{3-}$$

$$\text{At } t \quad 0 \quad 1 \quad 0 \quad 0$$

$$\text{After ionization (1) } \quad 3 \quad ; \quad i \quad 1 \quad 3$$

$$\text{Since, } i \quad \frac{M_{\text{normal}}}{M_{\text{abnormal}}};$$


$$\frac{1 \quad 3}{1} \quad \frac{M_{\text{normal}}}{M_{\text{abnormal}}}$$

$$103. (b) \therefore [\text{Urea}] \quad [\text{Glucose}],$$

Isotonic solutions

$$104. (c) \quad 1 \quad 2; \quad \frac{5}{342} \quad \frac{1000}{100} \quad \frac{1}{X} \quad \frac{1000}{100}$$

$$X \quad 68.4$$



$$105. (d)$$

$$\frac{W}{M} \quad \frac{RT}{V} \quad \text{or} \quad \frac{RT}{C} \quad \frac{RT}{M}$$

$$\text{slope} \quad \frac{RT}{M}$$

$$\text{or } M \quad \frac{0.082 \quad 293}{4.65 \quad 10^3 \quad 10^3}$$

$$5.16 \quad 10^6 \quad g$$

- 106. (c)** $\frac{\text{Sucrose}}{342} RT \times \frac{\text{Glucose}}{180} RT$
 Concentration of glucose $0.1 \frac{180}{18 \text{ g/L}}$
- 107. (a)** Berkley-Hartely's method is adopted to determine osmotic pressure.
- 108. (c)** $P = \frac{2.5}{39+14+48} \frac{1000}{100} \times 2 \times R \times 300$
 12.2 atm.
- 109. (b)** $i \cdot C \cdot RT$
 $1.8 \times \frac{1}{74.5} \times \frac{1}{0.1} \times 24.63$
 5.94
- 110. (a)** $7.4 \times n \times 0.0821 \times 300$
 $n = 0.3$
- 111. (c)** $iCRT$; Where, $i = 1$ for each of the solution
 $\therefore \frac{1}{\text{molecular mass of solute}};$
 $\frac{2}{\text{urea (60)}} \quad \frac{1}{\text{glucose (180)}} \quad \frac{3}{\text{sucrose (342)}}$
- 112. (a)** CRT
 $T [C \text{ const.}]$
 $C [T \text{ const.}]$
- 113. (b)** $M_B = \frac{W_B}{V} \times \frac{R \times T}{4 \times 0.0821 \times 300}$
 1.6×10^5
- 114. (a)** $i = \frac{\text{Number of particles in KI solution}}{\text{Number of particles in sucrose solution}}$
 $\frac{(\text{KI solution})}{(\text{sucrose solution})} = \frac{0.432 \text{ atm}}{0.24 \text{ atm}} \times 1.80$
- 115. (d)** $iCRT$; $i = 5$ for $\text{Al}_2(\text{SO}_4)_3$,
 $i = 4$ for Na_3PO_4 ,
 $i = 3$ for BaCl_2 , $i = 1$ for glucose
 So osmotic pressure of :
 $\text{Glucose} < \text{BaCl}_2 < \text{Na}_3\text{PO}_4 < \text{Al}_2(\text{SO}_4)_3$
- 116. (b)** $iCRT$, $i = 3$ $(n = 1)$;
 For the salt 1 , hence $n = 3$
 $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]\text{Cl}_2 \cdot \text{H}_2\text{O} + 2\text{AgNO}_3$
 $[\text{Cr}(\text{H}_2\text{O})_5\text{Cl}]^{2+} + 2\text{AgCl}(s)$
 $0.5 \text{ L of } 1m \text{ salt} \quad 0.5 \times 2 = 1.0 \text{ mole of AgCl}$
- 117. (c)** $\frac{59.1}{760} \times \frac{0.01}{(52 \times 4 \times 17 \times 96 \times 35.5)} \times \frac{1000}{25} \times 0.0821 \times 298(n)$
 $n = 2$
- 118. (d)** iCT ; iCT is maximum for 0.03 M NaCl at 50°C .
- 119. (d)** Only solvent molecules can pass through SPM so only dilution is possible.
- 120. (a)** Henry's law is $m = K P$; where, m mass of gas absorbed by given volume of the solvent.
 P pressure of gas;
 $\log m \quad \log K \quad \log P$
- 121. (b)** $W_{\text{Gas}} \propto P_{\text{Gas}}$ at constant T .
- 122. (b)** $\therefore n_{\text{O}_3}$ dissolved in 100 g water at 1.0 atm
 $\frac{40 \times 1 \times 10^3}{0.0821 \times 300}$
 n_{O_3} dissolved in 400 g water at 4 atm
 $\frac{4 \times 4 \times 40 \times 10^3}{24.63}$
 $W_{\text{O}_3} = \frac{640 \times 10^3}{24.63} \times 48 \times 1.2 \text{ g}$
- 123. (a)** $P_{\text{N}_2} = (K_H)_{\text{N}_2} X_{\text{Gas}}$
 $1 = (K_H)_{\text{N}_2} \times \frac{0.02/28}{0.02 \times \frac{1000}{28} + \frac{18}{18}}$
 $(K_H)_{\text{N}_2} = 7.7 \times 10^4 \text{ atm}$
- 124. (d)** $P_{\text{Gas}} = \frac{1}{K} n_{\text{Solvent}} \frac{MM_{\text{Gas}}}{X_{\text{Gas}}} \times K_H X_{\text{Gas}}$
 $\therefore K_H = \frac{n_{\text{Solvent}}}{K} \frac{MM_{\text{Gas}}}{X_{\text{Gas}}}$
 As T , K_H .
- 125. (a)** Mass of oxygen $\frac{20}{100} \times 0.04 = 0.008 \text{ g/L}$
 Mass of nitrogen $\frac{80}{100} \times 0.02 = 0.016 \text{ g/L}$

Level 2

- 1. (b)** Mole fraction of urea in its solution

$$\frac{\frac{12}{60}}{\frac{12}{60} + \frac{140.4}{18}} = 0.025$$

Mole fraction of glucose

$$\frac{18}{180} \quad 0.01$$

$$\frac{18}{180} \quad \frac{178.2}{18}$$

$\therefore$ Mole fraction of glucose is less so vapour pressure above the glucose solution will be higher than the pressure above urea solution, so some H_2O molecules will transfer from glucose to urea side in order to make the solutions of equal mole fraction to attain equilibrium, let x moles H_2O transferred

$$\begin{array}{ccc|ccc} 0.2 & & & 0.1 & & \\ \hline 0.2 & 7.8 & x & 0.1 & 9.9 & x \end{array} \quad x \quad 4$$

now mass of glucose solution

$$196.2 \quad 4 \quad 18 \quad 124.2$$

$$\text{wt. \% of glucose} \quad \frac{18}{124.2} \quad 100 \quad 14.49$$

2. (b) $\therefore \quad \frac{P_A}{P_B} = \frac{P_A X_A}{P_B X_B}$
 $(P_A \text{ and } P_B \text{ v.p. of pure A and B})$

$$Y_A = \frac{P_A}{P_A + P_B}$$

$$\frac{P_A X_A}{P_A X_A + P_B (1 - X_A)}$$

$$Y_A = \frac{P_A X_A}{X_A (P_A + P_B) + P_B}$$

$$\frac{1}{Y_A} = \frac{P_A + P_B}{P_A} \quad \frac{P_B}{P_A} \quad \frac{1}{X_A}$$

So, slope is $\frac{P_B}{P_A}$ and intercept $\frac{P_A + P_B}{P_A}$

3. (d) $\frac{P}{P} = \frac{n}{N} \quad \frac{W}{M} \quad \frac{N}{N}$

$$\frac{850}{844.9} = \frac{2}{M} \quad \frac{76}{100}$$

$$M = 252; \quad n = \frac{252}{32} = 8$$

atomicity of sulphur is 8

4. (c) $P = P_A x_A + P_B x_B \quad 100 \quad \frac{1}{5} \quad 200 \quad \frac{4}{5}$
 180 torr

y_A (composition of A vapour phase) $\frac{P_A x_A}{P}$

$$\frac{20}{180} \quad \frac{1}{9}, \quad Y_B = \frac{8}{9}$$

For condensation $y_A = x_A^1; y_B = x_B^1$

$$P_{\text{total}} = P_A x_A^1 + P_B x_B^1$$

$$P_{\text{total}} = 100 \quad \frac{1}{9} \quad \frac{8}{9} \quad 200 \quad 188.88 \text{ torr}$$

5. (c) Let n_B mole of B present in 1 mole of mixture that has been vaporized. Thus, $y_B = \frac{n_B}{1}$

Mole fraction of B in the remaining liquid phase will be $x_B = \frac{1 - n_B}{1}$

$$x_B = \frac{P}{P_B} \quad \frac{P_T}{P_T} \quad \dots (1)$$

$$[\because P = P_T = (P_B + P_T) x_B]$$

and $y_B = \frac{P_B}{P} = \frac{P_B x_B}{P} \quad \dots (2)$

After substitution of values of x_B and y_B in (1) and (2)

we get $1 - n_B = \frac{P}{P_B} \quad \frac{P_T}{P_T} \quad \dots (3)$

and $n_B = \frac{(1 - n_B) P_B}{P} \quad \dots (4)$

or $n_B = \frac{P_B}{P} \quad \frac{P_T}{P_T}$

so $1 = \frac{P_B}{P} \quad \frac{P_T}{P_T} \quad \frac{P}{P_B} \quad \frac{P_T}{P_T}$

$$P = \sqrt{P_B P_T} = \sqrt{100 \times 900}$$

$$300 \text{ torr}$$

6. (c) For an ideal solution $H_{\text{mix}} = 0$ and S_{mix} is always positive so G_{mix} is negative.

7. (d) $m_i = \frac{P}{P} \quad \frac{1000}{M_{\text{solvent}}}$

$$4.m = \frac{(17.25 \times 17.20) \times 1000}{17.2 \times 18}; \quad m = 0.04$$

for dilute solution molality $\sim$ molarity or S ;
 $K_{sp} = 27 S^4 \sim 7 \times 10^{-5}$

8. (d) Mole ratio of C : H : O is 1 : 2 : 1 so empirical formula is CH_2O

$$m = \frac{T_b}{K_b} = \frac{0.15}{0.51} = 0.294;$$

$$0.294 = \frac{50}{M} \quad \frac{1000}{950}; \quad M = 180$$

$(\text{CH}_2\text{O})_n = 180$ or $30 \quad n = 180$ or $n = 6$;
 molecular formula is $\text{C}_6\text{H}_{12}\text{O}_6$.

9. (c) Mass of 1 litre of solution 1010 g;
 mass of solvent 1010 300 0.1 980 g

$$m = \frac{0.1}{0.98} = 0.102;$$

$$T_f = K_f \cdot m \cdot i \quad (1) \quad K_f m$$

$$T_f = 1.05 \cdot 1.86 \cdot 0.102 = 0.199^\circ\text{C};$$

$$T_f = 0 + 0.199 = 0.199^\circ\text{C}$$
10. (b) $m = \frac{T_b}{K_b} = \frac{1}{0.51} = \frac{T_f}{K_f}$

$$\frac{1}{0.51} = \frac{T_f}{1.86}$$

$$T_f = 3.647 \text{ or } T_f = 3.647^\circ\text{C}$$
11. (c) For Na_3PO_4 , $i = 1 + 3 = 4$
 = 2.5; for MgSO_4 , $i = 1 + 1 = 2$
 100 g solution contains 8.2 g Na_3PO_4 and 12 g MgSO_4

$$K_b = \frac{\text{effective no. of moles of } (\text{Na}_3\text{PO}_4 + \text{MgSO}_4)}{\text{mass of solvent (in g)}} \cdot 1000$$

$$T_b = 0.50 \cdot \frac{8.2}{164} + 2.5 \cdot \frac{12}{120} = 1.785^\circ\text{C}$$

$$T_b = 100 + 1.785 = 101.785^\circ\text{C}$$
12. (b) $T_b = K_b \cdot \frac{w_{\text{solute}}}{W_{\text{solvent}}} = \frac{1}{M_{\text{solute}}} \cdot 1000$

$$1 = \frac{1000}{100} \cdot \frac{10}{M_{AB_2}} \Rightarrow M_{AB_2} = 100$$

 similarly $M_{A_2B} = 140$

$$100 = M_A + 2 \cdot M_B$$

 and $140 = 2 \cdot M_A + M_B$

$$M_A = 60 \text{ and } M_B = 20$$
13. (c) Given $w = 0.2 \text{ g}$, $W = 20 \text{ g}$, $T = 0.45^\circ\text{C}$

$$T_f = \frac{1000 \cdot K_f \cdot w}{M \cdot W}$$
 or $0.45 = \frac{1000 \cdot 5.12 \cdot 0.2}{20 \cdot M}$

$$M (\text{observed}) = 113.78$$

 Now for $2\text{CH}_3\text{COOH} \rightleftharpoons (\text{CH}_3\text{COOH})_2$
 Before association 1 0
 After association 1 1/2;
 where α is the degree of association.

$$\frac{M_{\text{normal}}}{M_{\text{observed}}} = \frac{1}{2};$$
 or $\frac{60}{113.78} = \frac{1}{2}; \quad 0.945$

14. (b) Given $H_{\text{fus.}} = 80 \text{ cal g}^{-1}$,
 $H_{\text{vap.}} = 540 \text{ cal g}^{-1}$
 We know, $T_b = K_b \cdot m$
 and $T_f = K_f \cdot m$;
 Also $K = \frac{RT^2}{1000 \cdot H}$

$$\frac{T_b}{T_f} = \frac{K_b}{K_f}$$

$$\frac{RT_b^2}{1000 \cdot H_{\text{vap.}}} = \frac{1000 \cdot H_{\text{fus.}}}{RT_f^2}$$

$$\frac{T_b}{T_f} = \frac{T_b^2}{T_f^2} \cdot \frac{H_{\text{fus.}}}{H_{\text{vap.}}}$$

$$\frac{0.1}{T_f} = \frac{373^2}{273^2} \cdot \frac{80}{540}$$

$$\therefore T_f = 0.361 \text{ so, } T_f = 0.361^\circ\text{C}$$

15. (d) Molality of solution $\frac{P}{P} = \frac{P}{M} = \frac{1000}{M}$

$$\frac{40.18}{40} = \frac{1000}{M} \Rightarrow M = 0.25$$

1000 g water present with 45 g glucose or
 100 g solution has 4.31 g glucose and 95.69 g H_2O . Final molality is 0.5, 1000 g solvent
 contain 90 g glucose or 4.31 g glucose
 present with $\frac{1000}{90} = 4.31$ 47.88 g H_2O

Mass of ice formed = 95.69 – 47.88
 = 47.8 g

16. (b) $T_f = K_f \cdot \frac{w_{\text{acid}}}{M_{\text{acid}} \cdot W_{\text{H}_2\text{O}}} = 1000;$

$M_{\text{acid}} (\text{observed}) = 120$

Normal molecular mass of acid can be
 calculated as


milli-equivalents of acid = milli-equivalents
 of base
 (for HA and NaOH, $N = M$)

$$\frac{0.75}{M_{\text{acid}}} = \frac{1000}{25} \cdot \frac{1}{5}$$

$M_{\text{acid}} (\text{Normal}) = 150$

$$i = \frac{150}{120} = 1.25$$

$$i = 1 + \alpha; \quad 0.25 = \alpha$$





$$[\text{Solute}] = \frac{0.6}{4} M$$

$$T_f = K_f \frac{0.6}{4}$$

$$1.86 \frac{0.6}{4} = 0.279 K$$




first step of H_2SO_4 is strong so



$$C(1) \quad C \quad C \quad C \quad C$$

$$i = \frac{C(1) + C + C + C + C}{C} = 2.1;$$

$$0.1$$



$$K_{a_2} = \frac{C(1) \cdot C}{C(1) \cdot C}$$

$$\frac{1.1 \cdot 0.1 \cdot 0.1}{0.9} = 0.0122$$

19. (a) $(iC_1 - i_2C_2) RT$
 $(1 \cdot 0.2 \cdot 3 - 0.05) 0.0821 \cdot 300$

$$1.23 \text{ atm}$$

20. (c) $\sqrt{\frac{8 \cdot 10^5}{0.2}} = 0.02;$

$$i = (1 + 1.02)$$

$$i \cdot C \cdot R \cdot T$$

$$1.02 \cdot 0.2 \cdot 0.0821 \cdot 300$$

$$5.024 \text{ atm}$$

21. (b) 8 g of carbohydrate is present with 100 g of water in solution
 or 108 g solution contains 8 g of carbohydrate;

$$V_{\text{solution}} = \frac{108}{1.025} \cdot 10^{-3} \text{ litre}$$

$$\frac{n}{V} = \frac{R}{T}$$

$$\frac{8}{M} = \frac{1.025}{108} \cdot 1000 \cdot 0.0821 \cdot 300 \cdot 5;$$

$$M = 374 \text{ g/mol}$$

22. (d) Only solvent molecule can pass through SPM.

Osmotic pressure iC (at constant T).

23. (a) The given data are

$$P_{\text{water}} = 17.0 \text{ torr};$$

$$P_{\text{total}} \text{ (4 mole \% solution)}$$

$$P_{\text{NH}_3} = 50.0 \text{ torr}$$

$$x_{\text{NH}_3} = 0.04 \text{ and } x_{\text{water}} = 0.96$$

Now according to Raoult's law;

$$P_{\text{water}} = x_{\text{water}} P_{\text{water}}^{\circ}$$

$$0.96 \cdot 17.0 \text{ torr} = 16.32 \text{ torr}$$

Now Henry's law constant for ammonia is

$$K_{\text{H}}(\text{NH}_3) = \frac{P_{\text{NH}_3}}{x_{\text{NH}_3}} = \frac{33.68 \text{ torr}}{0.04} = 842 \text{ torr}$$

Hence, for 5 mole % solution, we have

$$P_{\text{NH}_3} = K_{\text{H}}(\text{NH}_3) \cdot x_{\text{NH}_3} = (842 \text{ torr}) (0.05) = 42.1 \text{ torr}$$

$$P_{\text{water}} = P_{\text{water}}^{\circ} \cdot x_{\text{water}}$$

$$(17 \text{ torr}) (0.95) = 16.15 \text{ torr}$$

Thus, P_{total} (5 mole % solution)


$$P_{\text{NH}_3} + P_{\text{water}} = 42.1 + 16.15 = 58.25 \text{ torr}$$

24. (b) $P = P_A x_A + P_B x_B$
 $500 = 0.6 \cdot 800 + 0.4 \cdot 620 \text{ torr}$

$$y_A = \frac{P_A}{P} = \frac{300}{620}$$

$$0.48; \quad y_B = 0.52$$

25. (c) $P_A = 500; \quad P_B = 800 \text{ torr}$



When most of the liquid has vaporized $x_A = 0.6$ (given) would be $y_A = 0.6$

$$y_A = \frac{P_A x_A}{P_A x_A + P_B (1 - x_A)}$$

$$0.6 = \frac{500 x_A}{500 x_A + 800 (1 - x_A)}$$

$$x_A = 0.70; \quad x_B = 0.30$$

Level 3

Passage-1

- (a) Loss of mass of solvent $\rightarrow$ lowering in p ($p < P$).
- (b) Gain in mass of anhydrous $\text{CaCl}_2 \rightarrow P$.
- (b) $\frac{P}{P} = \frac{P}{P} \cdot \frac{w_{II}}{w_I} \cdot \frac{w_{II}}{w_I} \therefore \frac{P}{P} = \frac{w_I}{P} \cdot \frac{w_{II}}{w_I}$
- (b) $\frac{w_{II}}{w_I} = \frac{9.24/x}{9.24/108} = \frac{0.08}{3.28}$
 $x = 62$

Passage-2

- (b) $2A \rightleftharpoons A_2$; $n = 2$
 $i = 1 \quad \frac{1}{n} = \frac{1}{2} \quad 1 \quad \frac{1}{2}$
- (a) $i = \frac{M_T}{M} = 1$
- (b) $T_b = 1 + \frac{1}{2} = 1 + K_b x$
 $\frac{2(K_b x - T_b)}{K_b x}$

Subjective Problems

- $P = \frac{2}{5} \cdot 5 + \frac{3}{5} \cdot 10 = 8 \text{ torr}$
- $48 = \frac{8}{8 - x} \cdot 50 + \frac{x}{x - 8} \cdot 40$
 $x = 2$
- $P = P_A^\circ x_B (P_B^\circ - P_A^\circ)$
 $99 = 100 x_B (20 - \frac{1}{20})$
 $y_B = \frac{80}{99} \cdot \frac{1}{20} = \frac{4}{99}$
 $\% \text{ mole}_B \text{ in vapour phase} = \frac{4}{99} \cdot 100 = 4$
- $T_b = \frac{2.6 \cdot 30 + 1000}{154 + 250} = 2$
- $T_b = 0.5 + 2 = 1$
- $T_f = 2 + 0.56 = 1.8$
- For 100% ionization $i = 4$
- For isotonic solution

Protein	Sucrose	
$\frac{500}{x}$	$\frac{3.42}{342}$	$x = 5 \cdot 10^4$
- $T_f = 0.515 + 1.86 m$
 $m = 0.276$
 $0.276 \cdot 0.0821 \cdot 310 = 7 \text{ atm}$
- | | | |
|-----------------|--------------------------------|-----|
| KCl | K ₂ SO ₄ | |
| $\frac{0.2}{2}$ | $\frac{0.2}{2}$ | x |
| $x = 2$ | | |

SOLID STATE

Types of Solids

S. No.	Crystalline solids	Amorphous solids
1.	They have a definite crystal shape.	They do not have a definite geometrical shape
2.	Long-range order	Short-range order
3.	They have sharp melting points	They do not have sharp melting points.
4.	Enthalpy of fusion is definite	Enthalpy of fusion is not definite
5.	They are hard and rigid solids.	They are comparatively soft and not very rigid.
6.	Possess definite crystal system (cubic, tetragonal, hexagonal, octahedral, etc.)	Crystal system absent ; no regular repeating unit
7.	They are true solid	They are pseudo solid
8.	They split into two pieces when cut with a sharp-edged tool ; newly generated surfaces are plane and smooth.	They cut into two pieces with irregular surfaces when cut with a sharp-edged tool.
9.	They are anisotropic (refractive index and electrical conductivity are different in different directions) (e. g. , NaCl, CsBr, CaF ₂ and ZnS	They are isotropic. (physical properties are same in all directions) (e. g. , Rubber, plastic and glass)

Types of Crystalline Solids

Type of solid	Intermolecular forces	Properties	Examples
Ionic	Ion-ion forces	Brittle, hard high melting	NaCl, KCl, MgCl ₂
Molecular	Dispersion forces/Dipole-Dipole H-bond	Soft, low melting non-conducting	H ₂ O, Br ₂ , CO ₂ , CH ₄
Covalent network	Covalent bonds	Hard: High melting	C-Diamond, SiO ₂
Metallic	Metallic bonds	Variable hardness and melting point conducting	Na, Zn, Cu, Fe

Crystal System

Crystal Systems		Bravais Lattice	Unit Cell Parameters	
			Intercepts	Crystal Angles
1.	Cubic	Primitive, face centered, body centered	$a \ b \ c$	90
2.	Orthorhombic	Primitive, face centered, body centered, end centered	$a \ b \ c$	90
3.	Rhombohedral	Primitive	$a \ b \ c$	90
4.	Monoclinic	Primitive, end centered	$a \ b \ c$	90 , 90
5.	Triclinic	Primitive	$a \ b \ c$	90
6.	Tetragonal	Primitive, body centered	$a \ b \ c$	90
7.	Hexagonal	Primitive	$a \ b \ c$	90 , 120

Some Important Formula

(i) $d = \frac{ZM}{N_0 a^3}$; (ii) Packing fraction $\frac{Z V_{\text{Atom}}}{V_{\text{Unit cell}}}$; (iii) Void fraction $1 - \text{Packing fraction}$;

(iv) C.No. No. of first neighbours
 where d density; M Atomic mass of element;
 N_A Avogadro's number; a Edge length of cube.

Cubic System

S.No.	Property	Simple Cubic Lattice	BCC Lattice	FCC Lattice
1.	Atomic radius (r), a edge length of cube	$r = \frac{a}{2}$	$r = \frac{a\sqrt{3}}{4}$	$r = \frac{a\sqrt{2}}{4}$
2.	No of atoms per unit cell (Z)	$Z = 1$	$Z = 2$	$Z = 4$
3.	Co-ordination No. (C.No.)	6	8	12
4.	Packing fraction (P.F.)	0.52	0.68	0.74

Types of Packing

S.No.	Property	Hexagonal close packing (HCP)	Cubic close packing (CCP)
1.	Co-ordination No. (C.No.)	12	12
2.	No. of atoms per unit cell (z)	z = 6	z = 4
3.	Packing fraction (P.F.)	P.F. = 0.74	P.F. = 0.74
4.	Type of packing	ABAB	ABCABC

Types of Voids

S. No.	Name of Void	$r_{\text{Void}}/r_{\text{Sphere}}$	Co-ordination Number (C.No.)
1.	Triangular void	0.155	3
2.	Tetrahedral void	0.225	4
3.	Octahedral void	0.414	6
4.	Cubic void	0.732	8

Radius Ratio

S.No.	Limiting radius ratio $\frac{r_{\text{cation}}}{r_{\text{anion}}}$	C.No.	Shape
1.	0.155	2	Linear
2.	0.155 — 0.225	3	Planar triangle
3.	0.225 — 0.414	4	Tetrahedral
4.	0.414 — 0.732	6	Octahedral
5.	0.732 — 0.999	8	Like body centred cubic

Types of Ionic Structure


S.No.	Name of Structure	Location of Particle
1.	Rock salt (AB) NaCl	B : ccp lattice ; A : Octahedral void
2.	Zinc blende (ZnS)	S^{2-} : ccp lattice ; Zn^{2+} : Alternate tetrahedral void
3.	CsCl	Cs : Cube centre ; Cl : Corner of cube
4.	Fluorite structure (CaF ₂)	Ca^{2+} : fcc lattice ; F : Tetrahedral void
5.	Antifluorite structure Na ₂ O	O^{2-} : fcc lattice ; Na : Tetrahedral void

Bragg's Equation

$$2d \sin \theta = n\lambda$$

n 1 First order reflection ; n 2 Second order reflection ; θ angle of reflection.
 d distance between planes ; λ wavelength of x-ray

Defects in Crystals



(1) Stoichiometric defects

- ❖ **Vacancy:** When constituent particle missing, then vacancy defects occurs.
- ❖ **Frenkel defect :** The defect in which an ion is displaced from its regular position to an interstitial position creating a vacancy. Such defect is known as Frenkel defects or dislocation defect.
- ❖ **Schottky defect :** Defect in which a pair of one cation and one anion of equal valence is missing from an ionic crystal leading to a pair of vacant sites. Such defect is known as Schottky defect.

(2) Non-Stoichiometric defects :

- ❖ **Metal excess defect :** The defect occurs due to anionic vacancies or due to the presence of extra cations at the interstitial site.
- ❖ **Metal deficiency defect :** The defect occurs due to absence of positive ions from lattice site or extra interstitial negative ions.

Electrical Properties

The solids can be **conductors**, **insulators** or **semiconductors**.

- ❖ Conductors conduct electricity through movement of electrons (metals) or ions (electrolytes).
- ❖ The atomic orbitals of metals form molecular orbitals which are so close in energy that they form bands. If the gap between the filled valence band the next higher unoccupied conduction band overlaps, the electrons can jump and are called conductors. If this gap is more, the electrons can not jump and the substance behaves as an insulator.
- ❖ In semiconductors, this gap is small. For example, silicon and germanium. Their conductivity can be increased by adding impurity (doping) which can be electron rich or electron deficient.
- ❖ When increase in conductivity is due to the electron-rich impurity, it is called n-type semiconductor.
- ❖ When the increase in conductivity is due to electron-deficient impurity, the conductors are called p-type semiconductors.

Magnetic Properties

- ❖ **Paramagnetic** : Substance weakly attracted in magnetic field and magnetized in same direction is known as paramagnetic substance.
- ❖ **Diamagnetic** : Substance weakly repelled in magnetic field is known as diamagnetic substance
- ❖ **Ferromagnetic** : Substance attracted very strongly by magnetic field is known as ferromagnetic substance.
- ❖ **Antiferromagnetic** : Substance in which oppositely located domains cancel out each other's magnetic moment is known as Antiferromagnetic substance.
- ❖ **Ferrimagnetic** : Substance in which magnetic moment of domains are arranged in parallel and antiparallel directions in unequal numbers is known as ferrimagnetic substance.



Level 1

- Which of the following statement is true for ionic solids?
 - Ionic solids are soluble in non-polar solvent
 - Under the electric field cations and anions acquire translatory motion in opposite directions
 - Structural units have strong electrostatic force of attraction
 - Structural units have dipole-dipole interactions
- Which one is called pseudo solid ?
 - CaF_2
 - Glass
 - NaCl
 - All
- Solids which do not show the same physical properties in different directions are called:
 - pseudo solids
 - isotropic solids
 - polymorphic solids
 - anisotropic solids
- Graphite is an example of :
 - ionic solid
 - covalent solid
 - metallic crystal
 - none of these
- Amorphous solids are:
 - isotropic and supercooled liquids
 - anisotropic and supercooled liquids
 - isoenthalpic and superheated liquids
 - isotropic and superheated solids
- Crystals which are good conductor of electricity and heat are:
 - ionic crystals
 - covalent crystals
 - metallic crystals
 - molecular crystals
- The bond length and bond angles in molecules in the solid state are calculated by:
 - X-ray diffraction technique
 - neutrons bombardment
 - protons bombardment
 - none of these
- The unit cell with parameters a , b , c and angles 90° is :
 - cubic
 - triclinic
 - hexagonal
 - tetragonal
- Triclinic crystal has the following unit cell parameters :
 - $a \neq b \neq c$; $\alpha \neq \beta \neq \gamma$
 - $a \neq b \neq c$; $\alpha = \beta = \gamma = 90^\circ$
 - $a \neq b \neq c$; $\alpha = \beta \neq \gamma$
 - $a \neq b \neq c$; $\alpha = \beta = \gamma \neq 90^\circ$
- If all three interfacial angles defining the unit cell, are equal in magnitude, the crystal cannot be:
 - rhombohedral
 - cubic
 - hexagonal
 - tetragonal
- In a hexagonal crystal :
 - $a = b \neq c$; $\alpha = \beta = 120^\circ$; $\gamma = 90^\circ$
 - $a = b \neq c$; $\alpha = \beta = 90^\circ$; $\gamma = 120^\circ$
 - $a = b \neq c$; $\alpha = \beta = \gamma = 90^\circ$
 - $a = b \neq c$; $\alpha = \beta = 120^\circ$; $\gamma = 90^\circ$
- Orthorhombic crystal has the following unit cell parameters :
 - $a \neq b \neq c$; $\alpha = \beta = \gamma = 90^\circ$
 - $a \neq b \neq c$; $\alpha = \beta = 90^\circ$; $\gamma = 120^\circ$
 - $a \neq b \neq c$; $\alpha = \beta = \gamma = 120^\circ$
 - $a \neq b \neq c$; $\alpha = \beta = \gamma = 90^\circ$
- Which of the crystal systems contains the maximum number of Bravais lattices?
 - Cubic
 - Hexagonal
 - Triclinic
 - Orthorhombic


14. The most unsymmetrical and symmetrical systems are, respectively :
(a) Tetragonal, Cubic (b) Triclinic , Cubic
(c) Rhombohedral, Hexagonal (d) Orthorhombic, Cubic
15. The crystal system of a compound with unit cell parameters,
 a 0.328 nm, b 0.328 nm, c 0.527 nm and 90° is :
(a) cubic (b) tetragonal (c) monoclinic (d) rhombohedral
16. In the primitive cubic unit cell, the atoms are present at the:
(a) corners of the unit cell (b) centre of the unit cell
(c) centre of each face of the unit cell (d) one set of faces of the unit cell
17. In the body centered unit cell, the lattice points are present at the:
(a) corners of unit cell only
(b) corners and centre of unit cell
(c) corners and centre of each face of the unit cell
(d) corners and at one set of faces of unit cell
18. In the face centered unit cell, the lattice points are present at the:
(a) corners of unit cell only (b) corners and centre of the unit cell
(c) corners and face centres of the unit cell (d) face centres of the unit cell
19. The effective number of atoms per unit cell in a simple cube, face centred cube and body centred cube are respectively:
(a) 1, 4, 2 (b) 1, 2, 4 (c) 8, 14, 9 (d) 8, 4, 2
20. What would be the effective number of atoms per unit cell in end centered cubic unit cell, if this type of unit cell exist in nature ?
(a) 1 (b) 2 (c) 3 (d) 4
21. In the body centered cubic unit cell and simple cubic unit cell, the radius of atom in terms of edge length (a) of the unit cell is respectively:
(a) $\frac{a}{2}, \frac{a}{2\sqrt{2}}$ (b) $\frac{a}{\sqrt{2}}, \frac{a}{2}$ (c) $\frac{a}{2\sqrt{2}}, \frac{a}{2}$ (d) $\frac{\sqrt{3}a}{4}, \frac{a}{2}$
22. In the face centered cubic unit cell, the radius of atoms in terms of edge length (a) of unit cell is:
(a) $a/2$ (b) $a/\sqrt{2}$ (c) $a/2\sqrt{2}$ (d) $\sqrt{3}a/4$
23. The fraction of volume occupied by atoms in a primitive cubic unit cell is nearly:
(a) 0.48 (b) 0.52 (c) 0.55 (d) 0.68
24. The fraction of volume occupied by atoms in a body centered cubic unit cell is:
(a) 0.32 (b) 0.48 (c) 0.68 (d) 0.74
25. The fraction of volume occupied by atoms in a face centered cubic unit cell is:
(a) 0.32 (b) 0.48 (c) 0.68 (d) 0.74
26. Which of the following crystal lattice has the minimum empty space?
(a) Simple cubic (b) Body centred cubic
(c) Face centred cubic (d) Simple tetragonal
27. Which of the following has the smallest packing efficiency for atoms of a single type?
(a) Body centered cubic (b) Face centered cubic
(c) Simple cubic (d) None of these
28. Polonium crystallizes in a simple cubic structure. The edge of the unit cell is 0.236 nm. What is the radius of the polonium atom:
(a) 0.144 nm (b) 0.156 nm (c) 0.118 nm (d) 0.102 nm

29. Lithium crystallizes as body centered cubic crystals. If the length of the side of unit cell is 350 pm, the atomic radius of lithium is:
(a) 303.1 pm (b) 606.2 pm (c) 151.5 pm (d) 123.7 pm
30. Gold has a face centered cubic lattice with an edge length of the unit cube of 407 pm. The diameter of the gold atom is:
(a) 576.6 pm (b) 287.8 pm (c) 352.5 pm (d) 704.9 pm
31. The neon atoms has a radius of 160 pm. What is the edge of the unit cell of a face centered structure of neon?
(a) 490 pm (b) 320 pm (c) 453 pm (d) 481 pm
32. What are the number of atoms per unit cell and the number of nearest neighbours in a simple cubic structure?
(a) 1, 6 (b) 4, 12 (c) 2, 8 (d) 2, 6
33. What are the number of atoms per unit cell and the number of nearest neighbours in a face centered cubic structure?
(a) 4, 8 (b) 2, 8 (c) 2, 6 (d) 4, 12
34. What are the number of atoms per unit cell and the number of nearest neighbours in a body centered cubic structure?
(a) 4, 12 (b) 1, 6 (c) 2, 8 (d) 2, 5
35. Each edge of a cubic unit cell is 400 pm long. If atomic mass of the element is 120 and its density is 6.25 g/cm^3 , the crystal lattice is : (use $N_A = 6 \times 10^{23}$)
(a) primitive (b) body centered (c) face centered (d) end centered
36. Tungsten has an atomic radius of 0.136 nm. The density of tungsten is 19.4 g/cm^3 . What is the crystal structure of tungsten? (Atomic mass $W = 184$)
(a) Simple cubic (b) Body centered cubic
(c) Face centered cubic (d) None of these
37. The density of argon (face centered cubic cell) is 1.83 g/cm^3 at 20°C . What is the length of an edge a unit cell? (Atomic mass : Ar = 40)
(a) 0.599 nm (b) 0.569 nm (c) 0.525 nm (d) 0.551 nm
38. The density of nickel (face centered cubic cell) is 8.94 g/cm^3 at 20°C . What is the radius of the atom? (Atomic mass : Ni = 59)
(a) 0.124 nm (b) 0.136 nm (c) 0.149 nm (d) 0.110 pm
39. The density of krypton (face centered cubic cell) is 3.19 g/cm^3 . What is the radius of the atom? (Atomic mass : Kr = 84)
(a) 0.198 nm (b) 0.221 nm (c) 0.206 nm (d) 0.225 nm
40. The face centered cubic cell of platinum has an edge length of 0.392 nm. Calculate the density of platinum (g/cm^3) : (Atomic mass : Pt = 195)
(a) 20.9 (b) 20.4 (c) 19.6 (d) 21.5
41. The body centered cubic cell of chromium has an edge length of 0.288 nm. Calculate the density of chromium (g/cm^3) : (Atomic mass : Cr = 52.0)
(a) 6.80 (b) 7.60 (c) 6.60 (d) 7.23
42. An element crystallizes in a face centered cubic lattice and the edge of the unit cell is 0.559 nm. The density is 3.19 g/cm^3 . What is the atomic mass?
(a) 87.6 (b) 79.9 (c) 85.5 (d) 83.9

43. The element crystallizes in a body centered cubic lattice and the edge of the unit cell is 0.351 nm. The density is 0.533 g/cm^3 . What is the atomic mass?
(a) 12.0 (b) 6.94 (c) 9.01 (d) 10.8
44. An element X (At. mass 80 g/mol) has fcc structure. Calculate no. of unit cells in 8 gm of X:
(a) $0.4 N_A$ (b) $0.1 N_A$ (c) $4 N_A$ (d) none of these
45. Molybdenum (At. mass 96 g mol^{-1}) crystallizes as bcc crystal. If density of crystal is 10.3 g/cm^3 , then radius of Mo atom is (use $N_A = 6 \times 10^{23}$):
(a) 111 pm (b) 314 pm (c) 135.96 pm (d) none of these
46. What is the coordination number of an atom for an element crystallizing with a cubic lattice? Calculate the corresponding coordination number for the simple, fcc and bcc lattices:
(a) 12 sc; 12 fcc; 8 bcc (b) 6 sc; 14 fcc; 8 bcc
(c) 8 sc; 12 fcc; 6 bcc (d) 6 sc; 12 fcc; 8 bcc
47. Which of the following layering pattern will have a void fraction of 0.260?
(a) ABCCBAABC (b) ABBAABBA (c) ABCABCABC (d) ABCAABCA
48. The most malleable metals (Cu, Ag, Au) have close-packing of the type :
(a) Hexagonal close-packing
(b) Cubic close-packing
(c) Body-centred cubic packing
(d) Malleability is not related to type of packing
49. The co-ordination number of a metal crystallising in a hexagonal close-packed structure is:
(a) 12 (b) 4 (c) 8 (d) 6
50. If the ratio of coordination no. of A to that of B is $x : y$, then the ratio of no. of atoms of A to that no. of atoms of B in unit cell is:
(a) $x : y$ (b) $y : x$ (c) $x^2 : y$ (d) $y : x^2$
51. The atomic radius of strontium (Sr) is 215 pm and it crystallizes with a cubic closest packing. Edge length of the cube is:
(a) 430 pm (b) 608.2 pm (c) 496.53 pm (d) none of these
52. By X-ray diffraction it is found that nickel (at mass 59 g mol^{-1}), crystallizes with ccp. The edge length of the unit cell is $3.5 \text{ \AA}$. If density of Ni crystal is 9.0 g/cm^3 , then value of Avogadro's number from the data is :
(a) 6.05×10^{23} (b) 6.11×10^{23} (c) 6.02×10^{23} (d) 6.023×10^{23}
53. Stacking of square close packed layers give rise to :
(a) bcc structure (b) fcc structure
(c) Simple cubic structure (d) hcp structure
54. In a hexagonal close packed (hcp) structure of spheres, the fraction of the volume occupied by the sphere is A. In a cubic close packed structure the fraction is B. The relation for A and B is :
(a) $A = B$
(b) $A < B$
(c) $A > B$
(d) A is equal to the fraction in a simple cubic lattice.

55. The unit cell present in ABCABC, closest packing of atoms is:
 (a) hexagonal (b) tetragonal
 (c) face centered cubic (d) primitive cubic
56. The number of atoms present in a hexagonal close-packed unit cell is :
 (a) 4 (b) 6 (c) 8 (d) 12
57. The unit cell present in ABAB, closest packing of atoms is:
 (a) hexagonal (b) tetragonal
 (c) face centered cubic (d) primitive cubic
58. The numbers of tetrahedral and octahedral holes in a hexagonal primitive unit cell are respectively:
 (a) 8, 4 (b) 6, 12 (c) 2, 1 (d) 12, 6
59. Which one of the following schemes of ordering closed packed sheets of equal sized spheres do not generate closest packed lattice?
 (a) ABCABC (b) ABACABAC (c) ABBAABBA (d) ABCBCABCBC
60. In the closest packing of atoms, there are:
 (a) one tetrahedral void and two octahedral voids per atom
 (b) two tetrahedral voids and one octahedral void per atom
 (c) two of each tetrahedral and octahedral voids per atom
 (d) one of each tetrahedral and octahedral void per atom
61. Which of the following figures represents the cross-section of an octahedral site?
- 


(a)



(b)



(c)




(d)
62. In which of the following pairs of structures, tetrahedral as well as octahedral holes are found?
 (a) bcc and fcc (b) hcp and simple cubic
 (c) hcp and ccp (d) bcc and hcp
63. The number of octahedral void in bcc structure is :
 (a) 0 (b) 1 (c) 2 (d) 4
64. An ionic compound is expected to have octahedral structure if r_c/r_a ($r_c = r_a$) lies in the range of:
 (a) 0.414 to 0.732 (b) 0.732 to 0.82
 (c) 0.225 to 0.414 (d) 0.155 to 0.225
65. An ionic compound is expected to have tetrahedral structure if r_c/r_a :
 (a) lies in the range of 0.414 to 0.732 (b) lies in the range of 0.225 to 0.414
 (c) lies in the range of 0.155 to 0.225 (d) is more than 0.732
66. An ionic compound is expected to have body centered type cubic unit cell if r_c/r_a :
 (a) is greater than 0.732 (b) lies in the range of 0.414 to 0.732
 (c) lies in the range of 0.255 to 0.414 (d) lies in the range of 0.155 to 0.225
67. In the closest packing of atoms A (radius : r_a), the radius of atom B that can be fitted into tetrahedral void is:
 (a) $0.155 r_a$ (b) $0.225 r_a$ (c) $0.414 r_a$ (d) $0.732 r_a$

68. In the closest packing of atoms A (radius : r_a), the radius of atom B that can be fitted octahedral void is:
(a) $1.155 r_a$ (b) $0.225 r_a$ (c) $0.414 r_a$ (d) $0.732 r_a$
69. How many nearest neighbours are there in an atom or ion for an octahedral hole of a closed packed structure?
(a) 4 (b) 6 (c) 8 (d) 12
70. How many “nearest” and “next nearest” neighbours, respectively, does potassium have in bcc lattice ?
(a) 8, 8 (b) 8, 6 (c) 6, 8 (d) 6, 6
71. In the closest packing of atoms:
(a) the size of tetrahedral void is greater than that of octahedral void
(b) the size of tetrahedral void is smaller than that of octahedral void
(c) the size of tetrahedral void is equal to that of octahedral void
(d) the size of tetrahedral void may be greater or smaller or equal to that of octahedral void depending upon the size of atoms
72. In the ionic compound AB the ratio $r_A : r_B$ is 0.414. Indicate the correct statement among the following:
(a) Cations form close packing and anions exactly fit into the octahedral voids
(b) Anions form close packing and cations occupy precisely half of the tetrahedral voids
(c) Anions form close packing and cations occupy precisely all the octahedral voids
(d) Anions form close packing and cations fit into the octahedral voids loosely
73. In the unit cell of KCl ($NaCl$ type), Cl^- ions constitute ccp and K^+ ions fall into the octahedral holes. These holes are:
(a) one at the centre and 6 at the centres of the faces
(b) one at the centre and 12 at the centres of the edges
(c) 8 at the centres of 8 small cubes forming the unit cell
(d) none of these
74. Which is incorrect statement ?
(a) In $NaCl$ structure, tetrahedral voids are unoccupied
(b) In ZnS structure, octahedral voids are unoccupied
(c) In CaF_2 structure, all tetrahedral voids are occupied
(d) In Na_2O structure, all tetrahedral voids are unoccupied
75. If the radius of the anion in an ionic solid is 200 pm, what would be the radius of the cation that fits exactly into a cubic hole?
(a) 146.4 pm (b) 82.8 pm (c) 45 pm (d) None of these
76. The $CsCl$ type structure is exhibited by alkali halides only when the radius of the cation is large enough to keep touching its eight nearest neighbour anions. Below what minimum ratio of cation to anion radii (r^+/r^-) this contact is prevented?
(a) 0.225 (b) 0.414
(c) 0.632 (d) 0.732
77. MgO crystallizes in a cubic type crystal system. The ionic radii for Mg^{2+} and O^{2-} are 0.066 and 0.140 nm respectively. One can conclude that the Mg^{2+} ions occupy:
(a) a cubic hole in a simple cubic structure
(b) every tetrahedral hole in a close packed structure

- (c) an octahedral hole in a cubic close packed structure
 (d) every other tetrahedral hole in a close packed structure
- 78.** The unit cell of diamond is made up of:
 (a) 6 carbon atoms, 4 atoms constitute ccp and two atoms occupy half of octahedral voids
 (b) 8 carbon atom, 4 atoms constitute ccp and 4 atoms occupy all the octahedral voids
 (c) 8 carbon atoms, 4 atoms form fcc lattice and 4 atoms occupy half of the tetrahedral voids alternately
 (d) 12 carbon atoms, 4 atoms form fcc lattice and 8 atoms occupy all the tetrahedral holes
- 79.** In diamond, the coordination number of carbon is:
 (a) four and its unit cell has eight carbon atoms
 (b) four and its unit cell has six carbon atoms
 (c) six and its unit cell has four carbon atoms
 (d) four and its unit cell has four carbon atoms
- 80.** Predict coordination number of the cation in crystals of the following compounds:
 1. MgO : r_c 0.65 Å; r_a 1.40 Å, 2. MgS : r_c 0.65 Å; r_a 1.84 Å
 (a) 6, 4 (b) 4, 6 (c) 3, 4 (d) 6, 8
- 81.** In a cubic unit cell, seven of the eight corners are occupied by atoms A and centres of faces are occupied by atoms B . The general formula of the compound is :
 (a) A_7B_6 (b) A_7B_{12} (c) A_7B_{24} (d) $A_{24}B_7$
- 82.** CaS exists in a cubic close packed arrangement of S^{2-} ions in which Ca^{2+} ions occupy $1/2$ of the available tetrahedral holes. How many Ca^{2+} and S^{2-} ions are contained in the unit cell?
 (a) 1, 1 (b) 2, 4 (c) 4, 4 (d) 4, 2
- 83.** In the spinel structure, oxides ions are cubical-closest packed whereas $1/8$ th of tetrahedral voids are occupied by A^{2+} cation and $1/2$ of octahedral voids are occupied by B^{3+} cations. The general formula of the compound having spinel structure is:
 (a) $A_2B_2O_4$ (b) AB_2O_4 (c) $A_2B_4O_2$ (d) $A_4B_2O_2$
- 84.** If the anions (A) form hexagonal closest packing and cations (C) occupy only $2/3$ octahedral voids in it, then the general formula of the compound is:
 (a) CA (b) CA_2 (c) C_2A_3 (d) C_3A_2
- 85.** In a solid, oxide ions are arranged in ccp, cations A occupy $(1/8)^{\text{th}}$ of the tetrahedral voids and cations B occupy $(1/4)^{\text{th}}$ of the octahedral voids. The formula of the compound is:
 (a) ABO_4 (b) AB_2O_3 (c) A_2BO_4 (d) AB_4O_4
- 86.** In a face centered cubic arrangement of A and B atoms whose A atoms are at the corner of the unit cell and B atoms at the face centers. One of the B atoms missing from one of the face in unit cell. The simplest formula of compound is:
 (a) AB_3 (b) A_8B_5 (c) A_2B_5 (d) $AB_{2/5}$
- 87.** An alloy of copper, silver and gold is found to have copper constituting the ccp lattice. If silver atoms occupy the edge centres and gold is present at body centre, the alloy will have the formula:
 (a) $\text{Cu}_4\text{Ag}_2\text{Au}$ (b) $\text{Cu}_4\text{Ag}_4\text{Au}$ (c) $\text{Cu}_4\text{Ag}_3\text{Au}$ (d) CuAgAu

- 88.** Which of the following statements is correct in the rock-salt structure of an ionic compound?
- Co-ordination number of cation is four and anion is six
 - Co-ordination number of cation is six and anion is four
 - Co-ordination number of each cation and anion is four
 - Co-ordination number of each cation and anion is six
- 89.** Which of the following statements is correct in the body centered type cubic structure of an ionic compound?
- Co-ordination number of each cation and anion is two
 - Co-ordination number of each cation and anion is four
 - Co-ordination number of each cation and anion is six
 - Co-ordination number of each cation and anion is eight
- 90.** Which of the following statements is correct in the zinc blende type structure of an ionic compound?
- Co-ordination number of each cation and anion is two
 - Co-ordination number of each cation and anion is four
 - Co-ordination number of each cation and anion is six
 - Co-ordination number of each cation and anion is eight
- 91.** Which of the following expression is correct in case of a sodium chloride unit cell (edge length, a)?
- $r_c \quad r_a \quad a$
 - $r_c \quad r_a \quad a/2$
 - $r_c \quad r_a \quad 2a$
 - $r_c \quad r_a \quad \sqrt{2}a$
- 92.** In an ideal closest rock salt structure (edge length, a), which of the following expressions is correct?
- $r_a \quad \sqrt{2}a$
 - $r_a \quad a/\sqrt{2}$
 - $r_a \quad a/2 \sqrt{2}$
 - $r_a \quad a/4$
- 93.** Which of the following expressions is correct in case of a CsCl unit cell (edge length, a)?
- $r_c \quad r_a \quad a$
 - $r_c \quad r_a \quad a/\sqrt{2}$
 - $r_c \quad r_a \quad \sqrt{3}a/2$
 - $r_c \quad r_a \quad a/2$
- 94.** In NaCl the centres of two nearest like-charged ions are present at a distance of:
- $\frac{1}{2} a\sqrt{2}$
 - $\frac{1}{2} a$
 - $\frac{\sqrt{3}}{2} a$
 - $\frac{1}{\sqrt{2}} 2a$
- 95.** In sodium chloride crystal, the number of next nearest neighbours of each Na^+ ions is:
- 8 Cl^- ions
 - 12 Na^+ ions
 - 12 Cl^- ions
 - 24 Cl^- ions
- 96.** In an ionic compound $\text{A}^+ \text{X}^-$, the radii of A^+ and X^- ions are 1.0 pm and 2.0 pm, respectively. The volume of the unit cell of the crystal AX will be :
- 27 pm^3
 - 64 pm^3
 - 125 pm^3
 - 216 pm^3

97. The coordination number of cation and anion in fluorite CaF_2 and anti-fluorite Na_2O are respectively:
 (a) 8 : 4 and 6 : 3 (b) 6 : 3 and 4 : 4
 (c) 8 : 4 and 4 : 8 (d) 4 : 8 and 8 : 4
98. Select the incorrect statement in a CsCl crystal:
 (a) Cs forms a simple cubic lattice, Cl forms a simple cubic lattice
 (b) Cl occupies body centre of Cs
 (c) Cs occupies body centre of Cl
 (d) It is impossible for Cl to occupy body centre of Cs because the body centre void of Cs is smaller than Cl ion size
99. The radius of a divalent cation A^{2+} is 94 pm and of divalent anion B^{2-} is 146 pm. The compound AB has :
 (a) Rock salt structure (b) Zinc blende structure
 (c) Antifluorite structure (d) Caesium chloride like structure
100. A binary solid (AB) has a rock salt structure. If the edge length is 400 pm, and radius of cation is 80 pm the radius of anion is:
 (a) 100 pm (b) 120 pm
 (c) 250 pm (d) 325 pm
101. For a solid with the structure shown in Fig., the coordination numbers of the points A and B, respectively, are



- (a) 6, 8 (b) 8, 8 (c) 6, 6 (d) 4, 6
102. An ionic compound AB_2 has fluorite type structures. If the radius B is 200 pm, then the ideal radius of A would be:
 (a) 82.8 pm (b) 146.4 pm (c) 40 pm (d) 45 pm
103. In which of the following structures, the anion has maximum coordination number ?
 (a) NaCl (b) ZnS (c) CaF_2 (d) Na_2O
104. CsCl has bcc structure with Cs at the centre and Cl ion at each corner. If r_{Cs} is 1.69 Å and r_{Cl} is 1.81 Å, what is the edge length of the cube?
 (a) 3.50 Å (b) 3.80 Å
 (c) 4.04 Å (d) 4.50 Å
105. CsBr has bcc type structure with edge length 4.3 pm. The shortest inter ionic distance in between Cs and Br is:
 (a) 3.72 pm (b) 1.86 pm (c) 7.44 pm (d) 4.3 pm


- 106.** If the radius of Cl⁻ ion is 181 pm, and the radius of Na⁺ ion is 101 pm then the edge length of unit cell is:
(a) 282 pm (b) 285.71 pm
(c) 512 pm (d) 564 pm
- 107.** Ammonium chloride crystallizes in a body centered cubic lattice with edge length of unit cell equal to 387 pm. If the size of Cl⁻ ion is 181 pm, the size of NH₄⁺ ion would be:
(a) 116 pm (b) 154 pm
(c) 174 pm (d) 206 pm
- 108.** Salt AB has a zinc blende structure. The radius of A²⁺ and B²⁻ ions are 0.7 Å and 1.8 Å respectively. The edge length of AB unit cell is:
(a) 2.5 Å (b) 5.09 Å
(c) 5 Å (d) 5.77 Å
- 109.** Transition metals, when they form interstitial compounds, the non-metals (H, B, C, N) are accommodated in :
(a) voids or holes in cubic-packed structure (b) tetrahedral voids
(c) octahedral voids (d) all of these
- 110.** In diamond, each carbon atom is bonded to four other carbon atoms tetrahedrally. Alternate tetrahedral voids are occupied by carbon atoms. The number of carbon atoms per unit cell is :
(a) 4 (b) 6
(c) 8 (d) 12
- 111.** Which of the following statements for crystals having Schottky defect is not correct?
(a) Schottky defect arises due to the absence of a cation and anion from the position which it is expected to occupy
(b) Schottky defect are more common in ionic compounds with high co-ordination numbers
(c) The density of the crystals having Schottky defect is larger than that of the perfect crystal
(d) The crystal having Schottky defect is electrically neutral as a whole
- 112.** Which is correct statement ?
(a) When temperature increases then number of defects decreases.
(b) Schottky defect occurs when radius of cation is smaller
(c) Frenkel defect occurs when radius of cation is smaller
(d) None of these
- 113.** Which of the following statements for crystals having Frenkel defect is not correct?
(a) Frenkel defects are observed where the difference in sizes of cations and anions is large
(b) The density of crystals having Frenkel defect is lesser than that of a pure perfect crystal
(c) In an ionic crystal having Frenkel defect may also contain Schottky defect
(d) Usually alkali halides do not have Frenkel defect
- 114.** When anion leaves the normal lattice site and electron occupies interstitial sites in its crystal lattice, it is called:
(a) Schottky defect (b) Frenkel defect
(c) Metal excess defect (d) Stoichiometric defect
- 115.** Which of the following defects does KBr show?
(a) Frenkel (b) Schottky
(c) Metal excess (d) Metal deficiency

- 116.** Dopping of AgCl crystals with CdCl_2 results in:
(a) Schottky defect (b) Frenkel defect
(c) Substitutional cation vacancy (d) Formation of F-centres
- 117.** NaCl shows Schottky defects and AgCl Frenkel defects. Their electrical conductivity is due to the:
(a) motion of ions and not the motion of electrons
(b) motion of electrons and not the motion of ions
(c) lower coordination number of NaCl
(d) higher coordination number of AgCl
- 118.** Which one of the following crystals does not exhibit Frenkel defect?
(a) AgBr (b) AgCl
(c) CsCl (d) ZnS
- 119.** Select the incorrect statement:
(a) Stoichiometry of crystal remains unaffected due to Schottky defect
(b) Frenkel defect is usually shown by ionic compounds having low coordination number
(c) F-centres generation is responsible factor for imparting the colour to the crystal
(d) Density of crystal always increases due to substitutional impurity defect
- 120.** In diamond, carbon atom occupy fcc lattice points as well as alternate tetrahedral voids. If edge length of the unit cell is 356 pm, then diameter of carbon atom is:
(a) 77.07 pm (b) 154.14 pm
(c) 251.7 pm (d) 89 pm
- 121.** When NaCl is dopped with 10^{-5} mole % of SrCl_2 , what is the no. of cationic vacancies?
(a) $10^{-5} N_A$ (b) $10^{-7} N_A$
(c) $2 \times 10^{-7} N_A$ (d) None of these
- 122.** The composition of a sample of Wustite is $\text{Fe}_{0.93}\text{O}$. What is the percentage of iron present as Fe^{3+} in total iron?
(a) 15.05% (b) 25%
(c) 35% (d) 45%
- 123.** A certain sample of cuprous sulphide is found to have composition $\text{Cu}_{1.8}\text{S}$, because of incorporation of Cu^{2+} ions in the lattice. What is the mole % of Cu^{2+} in total copper content in this crystal?
(a) 99.8% (b) 11.11%
(c) 88.88% (d) None of these
- 124.** Ferrimagnetism is in :
(a) (b)
(c) (d) none of these
- 125.** Fe_3O_4 is ferrimagnetic at room temperature but at 850 K it becomes :
(a) diamagnetic (b) ferromagnetic
(c) non-magnetic (d) paramagnetic



Level 2


- When heated above 916°C , iron changes its bcc crystalline form to fcc without the change in the radius of atom. The ratio of density of the crystal before heating and after heating is:
(a) 1.069 (b) 0.918 (c) 0.725 (d) 1.231
- $\text{TiAl}(\text{SO}_4)_2 \cdot x\text{H}_2\text{O}$ is bcc with ' a ' 1.22 nm. If the density of the solid is 2.32 g/cc, then the value of x is (Given : $N_A = 6 \times 10^{23}$; at. mass : Ti 204, Al 27, S 32).
(a) 2 (b) 4 (c) 47 (d) 70
- In an atomic bcc lattice what fraction of edge is not covered by atoms?
(a) 0.32 (b) 0.16 (c) 0.134 (d) 0.268
- The packing efficiency of a simple cubic crystal with an interstitial atom exactly fitting at the body center is:
(a) 0.48 (b) 0.52 (c) 0.73 (d) 0.91
- An atomic solid crystallizes in a body centre cubic lattice and the inner surface of the atoms at the adjacent corner are separated by 60.3 pm. If the atomic mass of A is 48, then density of the solid, is nearly:
(a) 2.7 g/cc (b) 5.07 g/cc (c) 3.5 g/cc (d) 1.75 g/cc
- Sodium ($\text{Na} = 23$) crystallizes in bcc arrangement with the interfacial separation between the atoms at the edge 53.6 pm. The density of sodium crystal is:
(a) 2.07 g/cc (b) 2.46 g/cc (c) 1.19 g/cc (d) none of these
- The density of solid argon ($\text{Ar} = 40$ g/mol) is 1.68 g/mL at 40 K. If the argon atom is assumed to be a sphere of radius 1.50×10^{-8} cm, what % of solid Ar is apparently empty space? (use $N_A = 6 \times 10^{23}$)
(a) 35.64 (b) 64.36 (c) 74 (d) none of these
- A body centered cubic lattice is made up of hollow spheres of B . Spheres of solid A are present in hollow spheres of B . Radius of A is half of radius of B . What is the ratio of total volume of spheres of B unoccupied by A in a unit cell and volume of unit cell?
(a) $\frac{7\sqrt{3}}{64}$ (b) $\frac{7\sqrt{3}}{128}$ (c) $\frac{7}{24}$ (d) none of these
- First three nearest neighbour distances for primitive cubic lattice are respectively (edge length of unit cell a):
(a) $a, \sqrt{2}a, \sqrt{3}a$ (b) $\sqrt{3}a, \sqrt{2}a, a$ (c) $a, \sqrt{2}a, 2a$ (d) $a, \sqrt{3}a, 2a$
- First three nearest neighbour distances for body centered cubic lattice are respectively:
(a) $\sqrt{2}a, a, \sqrt{3}a$ (b) $\frac{a}{\sqrt{2}}, a, \sqrt{3}a$ (c) $\frac{\sqrt{3}a}{2}, a, \sqrt{2}a$ (d) $\frac{\sqrt{3}a}{2}, a, \sqrt{3}a$
- Given : The unit cell structure of compound is shown below.



The formula of compound is :

- (a) $A_8B_{12}C_5$ (b) AB_2C_3 (c) $A_2B_2C_5$ (d) ABC_5
12. The density of a pure substance 'A' whose atoms are in cubic close pack arrangement is 1 g/cc. If all the tetrahedral voids are occupied by 'B' atom, what is the density of resulting solid in g/cc. [Atomic mass (A) 30 g/mol and atomic mass (B) 50 g/mol]
 (a) 3.33 (b) 4.33 (c) 2.33 (d) 5.33
13. In a planar tetra-atomic molecule, XY_3 , X is at the centroid of the equilateral triangle formed by the atoms, Y. If the X—Y bond distance is 1 Å, what is the distance between the centres of any two Y atoms?
 (a) $1/\sqrt{3}$ Å (b) $\sqrt{2}$ Å
 (c) $\sqrt{3}$ Å (d) $1/\sqrt{2}$ Å
14. How many unit cells are present in 5.0 gm of crystal AB (formula mass of AB 40) having rock salt type structure? (N_A Avogadro's no.)
 (a) N_A (b) $\frac{N_A}{10}$
 (c) $4 N_A$ (d) none of these
15. The density of CaF_2 (fluorite structure) is 3.18 g/cm^3 . The length of the side of the unit cell is:
 (a) 253 pm (b) 344 pm
 (c) 546 pm (d) 273 pm
16. A crystal of lead (II) sulphide has NaCl structure. In this crystal the shortest distance between a Pb^{2+} ion and S^{2-} ion is 297 pm. What is the volume of unit cell in lead sulphide?
 (a) $209.6 \times 10^{-24} \text{ cm}^3$ (b) $207.8 \times 10^{-23} \text{ cm}^3$
 (c) $22.3 \times 10^{-23} \text{ cm}^3$ (d) $209.8 \times 10^{-23} \text{ cm}^3$
17. CdO has NaCl structures with density 8.27 g/cc. If the ionic radius of O^{2-} is 1.24 Å, determine ionic radius of Cd^{2+} :
 (a) 1.5 Å (b) 1.1 Å
 (c) 1.9 Å (d) 1.5 Å
18. KCl crystallizes in the same type of lattice as does NaCl. Given that r_{Na}/r_{Cl} 0.50 and r_{Na}/r_K 0.70, calculate the ratio of the side of the unit cell for KCl to that for NaCl:
 (a) 1.143 (b) 1.224
 (c) 1.414 (d) 0.875
19. Ferrous oxide has a cubic structure and edge length of the unit cell is 5.0 Å. Assuming the density of ferrous oxide to be 3.84 g/cm^3 , the no. of Fe^{2+} and O^{2-} ions present in each unit cell be : (use N_A 6×10^{23}) :
 (a) 4 Fe^{2+} and 4 O^{2-} (b) 2 Fe^{2+} and 2 O^{2-}
 (c) 1 Fe^{2+} and 1 O^{2-} (d) 3 Fe^{2+} and 4 O^{2-}
20. If an element (at. mass 50) crystallises in fcc lattice, with a 0.50 nm. What is the density of unit cell if it contains 0.25% Schottky defects (use N_A 6×10^{23})?
 (a) 2.0 g/cc (b) 2.66 g/cc
 (c) 3.06 g/cc (d) none of these

21. An element X (atomic mass 24 gm/mol) forms a face centered cubic lattice. If the edge length of the lattice is $4 \times 10^{-8} \text{ cm}$ and the observed density is $2.40 \times 10^3 \text{ kg/m}^3$, then the percentage occupancy of lattice points by element X is : (Use $N_A = 6 \times 10^{23}$) :
- (a) 96 (b) 98
(c) 99.9 (d) none of these
22. In fcc lattice, A, B, C, D atoms are arranged at corner, face center, octahedral void and tetrahedral void respectively, then the body diagonal contains:
- (a) $2A, C, 2D$ (b) $2A, 2B, 2C$
(c) $2A, 2B, D$ (d) $2A, 2D$
23. The distance between an octahedral and tetrahedral void in fcc lattice would be:
- (a) $\sqrt{3}a$ (b) $\frac{\sqrt{3}a}{2}$
(c) $\frac{\sqrt{3}a}{3}$ (d) $\frac{\sqrt{3}a}{4}$
24. A_2B molecules (molar mass 259.8 g/mol) crystallises in a hexagonal lattice as shown in figure. The lattice constants were $a = 5 \text{ \AA}$ and $b = 8 \text{ \AA}$. If density of crystal is 5 g/cm^3 then how many molecules are contained in given unit cell? (use $N_A = 6 \times 10^{23}$)



- (a) 6 (b) 4 (c) 3 (d) 2
25. Graphite has h.c.p. arrangements of carbon atoms and the parallel planes are $3.35 \text{ \AA}$ apart. Determine density of graphite:
- (a) 2.46 g/cc (b) 0.41 g/cc (c) 1 g/cc (d) 1.41 g/cc
26. How many effective Na^+ and Cl^- ions are present respectively in a unit cell of NaCl solid (Rock salt structure) if all ions along line connecting opposite face centres are absent?
- (a) 3, 3 (b) $\frac{7}{2}, 4$ (c) $\frac{7}{2}, \frac{7}{2}$ (d) $4, \frac{7}{2}$
27. A crystal is made of particles X and Y . X forms fcc packing and Y occupies all the octahedral voids. If all the particles along one body diagonal are removed then the formula of the crystal would be:
- (a) X_4Y_3 (b) X_5Y_4 (c) X_4Y_5 (d) none of these

28. Select right expression for determining packing fraction (P.F.) of NaCl unit cell (assume ideal), if ions along an edge diagonal are absent:




(a) P.F. $\frac{\frac{4}{3} (r^3 + r^3)}{16\sqrt{2} r^3}$

(b) P.F. $\frac{\frac{4}{3} + \frac{5}{2} r^3 + 4r^3}{16\sqrt{2} r^3}$


(c) P.F. $\frac{\frac{4}{3} + \frac{5}{2} r^3 + r^3}{16\sqrt{2} r^3}$

(d) P.F. $\frac{\frac{4}{3} + \frac{7}{2} r^3 + r^3}{16\sqrt{2} r^3}$

29. A crystal is made of particles X, Y and Z. X forms fcc packing. Y occupies all the octahedral voids of X and Z occupies all the tetrahedral voids of X. If all the particles along one body diagonal are removed then the formula of the crystal would be:



30. A crystal is made of particles A and B. A forms fcc packing and B occupies all the octahedral voids. If all the particles along the plane as shown in figure are removed, then, the formula of the crystal would be:



31. In the rock salt AB, if C introduced in tetrahedral voids such that no distortion occurs, then formula of resultant compound is :



32. Given length of side of hexagonal unit cell is $\frac{100}{\sqrt{2}}$ pm. The volume of hexagonal unit cell is

(in pm^3) :

(a) 8×10^6

(b) 1.5×10^6

(c) 64×10^6

(d) 36×10^6



Level 3



Passage 1

Packing fraction of a unit cell is defined as the fraction of the total volume of the unit cell occupied by the atom(s).

$$\text{P.F.} = \frac{\text{Volume of the atom(s) present in a unit cell}}{\text{Volume of unit cell}} = \frac{Z \cdot \frac{4}{3} \pi r^3}{a^3}$$

and % of empty space $100 - \text{P.F.} \times 100$
 where Z effective number of atoms in a cube
 r radius of an atom
 a edge length of the cube

- % of empty space in body centered cubic unit cell is nearly:
 (a) 52.36 (b) 68 (c) 32 (d) 26
- Packing fraction in face centered cubic unit cell is:
 (a) 0.7406 (b) 0.6802 (c) 0.5236 (d) None of these



Passage 2

Density of a unit cell is represented as

$$\frac{\text{Effective no. of atom(s)} \cdot \text{Atomic mass / formula mass}}{\text{Volume of a unit cell}} = \frac{Z \cdot M}{N_A \cdot a^3}$$

where, mass of unit cell = mass of effective no. of atom(s) or ion(s).

M At. mass/formula mass
 N_A Avogadro's no. 6.023×10^{23}
 a edge length of unit cell

- Silver crystallizes in a fcc lattice and has a density of 10.6 g/cm^3 . What is the length of an edge of the unit cell?
 (a) 40.7 nm (b) 0.2035 nm (c) 0.101 nm (d) 4.07 nm
- An element crystallizes in a structure having fcc unit cell of an edge 200 pm. Calculate the density, if 100 g of this element contains 12×10^{23} atoms:
 (a) 41.66 g/cm^3 (b) 4.166 g/cm^3 (c) 10.25 g/cm^3 (d) 1.025 g/cm^3
- The density of KBr is 2.75 g/cm^3 . The length of the edge of the unit cell is 654 pm. To which type of cubic crystal, KBr belongs to?
 (a) simple cubic (b) bcc (c) fcc (d) none of these



Passage 3

A spinel is an important class of oxides consisting of two types of metal ions with the oxide ions arranged in ccp layers. The normal spinel has one-eighth of the tetrahedral holes occupied by one type of metal ion and one-half of the octahedral holes occupied by another type of metal ion. Such a spinel is formed by Mg^{2+} , Al^{3+} and O^{2-} . The neutrality of the crystal is being maintained.


- The formula of the spinel is:
 (a) Mg_2AlO_4 (b) MgAl_2O_4 (c) $\text{Mg}_3\text{Al}_2\text{O}_6$ (d) none of these
- Type of hole occupied by Al^{3+} ions is:
 (a) tetrahedral (b) octahedral (c) both (a) and (b) (d) none of these
- Type of hole occupied by Mg^{2+} ions is:
 (a) tetrahedral (b) octahedral (c) both (a) and (b) (d) none of these
- If oxide ion is replaced by $\text{X}^{8/3}$, the number of anionic vacancy per unit cell is:
 (a) 1 (b) 2 (c) 3 (d) $3/4$




Passage 4

Ionic lattice has two major points defects, (1) Schottky (2) Frenkel defects. Schottky defects occurs due to the cation-anion pair's missing from the lattice sites. Frenkel defects occurs when an ion leaves its lattice site and fits into an interstitial space. The neutrality of the crystal is being maintained and we considered all losses from interstitial positions.

- Which defect decreases density of the crystal?
 (a) Frenkel defect (b) Schottky defect (c) Both (a) and (b) (d) None of these
- Structure shown here represents:



- Schottky defect
 - Frenkel defect
 - Metal excess defect
 - None of these
- Structure shown here represents:



- Schottky defect
- Frenkel defect
- Both defects
- None of these



Passage 5

Doping means introduction of small amount of impurities like phosphorus, arsenic or boron into the pure crystal. In pure silicon, there are four valencies used in bonding with other four adjacent silicon atoms. When a silicon crystal is doped with a group-15 element (with five valence electrons) such as P, As, or Bi, the structure of the crystal lattice remains unchanged. Out of the five valence electrons of group-15 doped element, four electrons are used in normal covalent bonding with silicon while fifth electron is delocalised and thus conducts electricity.

Doping a silicon crystal with a group-13 element (with three valence electrons) such as B, Al, Ga or In produces a semiconductor with three electrons in dopant. The place where fourth electron is missing is called an electron vacancy or hole. Such hole can move through the crystal like a positive charge giving rise to conduction of electricity.

- No. of valence electrons in silicon are:
(a) 3 (b) 4 (c) 5 (d) 6
- Silicon that has been doped with group-15 elements is called:
(a) *p*-type semiconductor (b) *n*-type semiconductor
(c) electron vacancy or hole (d) none of these
- Silicon that has been doped with group-13 elements is called:
(a) *p*-type semiconductor (b) *n*-type semiconductor
(c) electron vacancy or hole (d) none of these



Passage 6

Metallic Gold crystallises in fcc lattice and the length of cubic unit cell is 407 pm.
(Given : Atomic mass of Gold = 197, $N_A = 6 \times 10^{23}$)

- The density if it has 0.2% Schottky defect is (in gm/cm^3) :
(a) 4.86 (b) 9.72 (c) 19.48 (d) 19.44
- The distance between next neighbours is :
(a) 407 pm (b) $407\sqrt{2}$ (c) $\frac{407}{\sqrt{2}}$ (d) $\frac{407\sqrt{3}}{2}$



Passage 7

In diamond structure, carbon atoms form fcc lattice and 50% tetrahedral voids are occupied by carbon atoms. Every carbon atom is surrounded tetrahedrally by four carbon atoms with bond length 154 pm. Germanium, silicon and grey tin also crystallise in the same way as diamond. ($N_A = 6 \times 10^{23}$)

- The mass of diamond unit cell is :
(a) 96 amu (b) 96 gm (c) 144 amu (d) 144 gm
- The side length of diamond unit cell is (in pm) :
(a) 154 (b) 1422.63 (c) 711.32 (d) 355.66

3. The density of diamond unit cell is (in gm/cm^3) :

- (a) 28.48 (b) 0.0556 (c) 0.445 (d) 3.56

One or More Answers is/are Correct



- Select the correct statement(s):
 - Co-ordination no. of an atom at a lattice point in simple cubic arrangement is 6
 - Co-ordination no. of an atom at octahedral site is 8
 - Co-ordination no. of an atom at a lattice point in hcp arrangement is 6
 - Co-ordination no. of an atom at octahedral site is 6
- Packing fraction of an identical solid sphere is 74% in:
 - simple cubic structure
 - fcc structure
 - hcp structure
 - bcc structure
- In fcc structure octahedral voids are present at:
 - edge centers
 - face centers
 - body centers
 - corners
- If the radius of Na is 95 pm and that of Cl ions is 181 pm then:
 - co-ordination no. of Na is 6
 - co-ordination no. of Cl is 8
 - length of the unit cell is 552 pm
 - length of the unit cell is 380 pm
- Select the correct statement(s):
 - The co-ordination number of each type of ion in a CsCl crystal is twelve
 - A metal that crystallizes in a bcc structure has a co-ordination number of twelve
 - A unit cell of an ionic crystal shares some of its ions with other unit cells
 - The length of the unit cell in NaCl is 552 pm (given that $r_{\text{Na}} = 85 \text{ pm}$ and $r_{\text{Cl}} = 181 \text{ pm}$)
- The hcp and ccp structure for a given element would be expected to have:
 - the same co-ordination number
 - the same density
 - the same packing fraction
 - all of the above
- Select the correct statement(s) for the rock-salt structure (NaCl) :
 - The tetrahedral voids are smaller than the octahedral sites
 - The octahedral voids are occupied by cations and the tetrahedral sites are empty
 - The radius ratio (r_c/r_a) is 0.225
 - The radius ratio (r_c/r_a) is 0.732
- Select the correct statement(s):
 - The ionic crystal of AgBr may have Schottky defect
 - The unit cell having crystal parameters, $a = b \neq c$, $\alpha = 90^\circ$, $\beta = 120^\circ$ is hexagonal
 - In ionic compounds having Frenkel defect the ratio r_+/r_- is high
 - The co-ordination number of Na ion in NaCl is 6
- Which of the following represents the closest packed arrangement of uniform solid spheres:
 - simple cubic unit cell
 - body centered cubic unit cell
 - face centered cubic unit cell
 - hcp unit cell

- 10.** Select the correct statement(s):
- A cubic system possesses a total of 23 elements of symmetry
 - A cubic system contains centre of symmetry, planes of symmetry as well as axes of symmetry
 - For triclinic system $a \neq b \neq c$ and $\alpha \neq \beta \neq \gamma$
 - The total no. of Bravais space lattice belonging to all the seven crystals are 14
- 11.** Select the correct statement(s):
- Co-ordination no. of Cs⁺ and Cl⁻ are 8, 8 in CsCl crystal
 - If radius ratio (r_c/r_a) < 0.225 then shape of compound must be linear
 - If radius ratio (r_c/r_a) lies between 0.414 to 0.732 then shape of ionic compound may be square planer (Ex. PtCl_4^{2-})
 - If radius ratio is less than 0.155 then shape of compound is linear
- 12.** Select the correct statement(s):
- CsCl changes to NaCl structure on heating
 - NaCl changes to CsCl structure on applying pressure
 - Co-ordination number decreases on applying pressure
 - Co-ordination number increases on heating
- 13.** Select the correct statement(s):
- A NaCl type AB crystal lattice can be interpreted to be made up of two individual fcc type unit lattice of A and B fused together in such a manner that the corner of one unit lattice becomes the edge centre of the other
 - In a fcc unit cell, the body center is an octahedral void
 - In an scc lattice, there can be no octahedral void
 - In an scc lattice, the body center is the octahedral void
- 14.** In a AB unit cell (Rock salt type) assuming A forming fcc:
- The nearest neighbour of A is 6B ion
 - The nearest neighbour of B is 6A ion
 - The second neighbour of A is 12A
 - The packing fraction of AB crystal is $\frac{\sqrt{3}}{8}$
- 15.** Amorphous solids:
- do not have sharp melting points
 - are isotropic
 - have same physical properties in all directions
 - are supercooled liquids
- 16.** Which is/are correct statement about zinc blende structure ?
- The number of first neighbours of S^{2-} is 4
 - The maximum distance between Zn^{2+} is $\frac{a\sqrt{3}}{2}$, where 'a' edge length of unit cell
 - If all tetrahedral voids are occupied by Zn^{2+} then C.N. of S^{2-} is 8.
 - If all tetrahedral voids are occupied by Zn^{2+} then C.N. change from 4 : 4 to 8 : 8.
- 17.** Compound X_2Y have antifluorite structure. Which is/are correct statement ?
- The minimum distance between X is $\frac{a}{2}$, where 'a' edge length of unit cell

- (b) The co-ordination number ratio of x and y is $8 : 4$
 (c) If X removed from alternate tetrahedral void then CN is $4 : 4$.
 (d) If X removed from alternate tetrahedral void then CN is $4 : 8$
- 18.** Which is/are correct statement ?
 (a) Packing fraction in 2D-hcp is 0.785 (b) Packing fraction in AAA is 0.52
 (c) Packing fraction in ABAB is 0.74 (d) Void fraction in ABCABC is 0.26
- 19.** Which is/are correct statement ?
 (a) In simple cubic close packed arrangement no octahedral void is present at edge centre.
 (b) In fcc unit cell octahedral void and tetrahedral void are vacant.
 (c) Packing fraction : simple cubic unit cell < bcc unit cell < fcc unit cell
 (d) Size of void: cubic void > octahedral void > tetrahedral void
- 20.** Given : Radius of A^2 100 pm ; Radius of C 240 pm ; Radius of B^2 300 pm ; Radius of D 480 pm Which is/are correct statement ?
 (a) Coordination number of A^2 in comp AB is 4
 (b) Coordination number of A^2 in comp AB is 6.
 (c) Coordination number of C in comp CD is 6
 (d) Coordination number of C in comp CD is 8.

Match the Column



Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I

- (A) Tetragonal and Hexagonal
 (B) Cubic and Rhombohedral
 (C) Monoclinic and Triclinic
 (D) Cubic and Orthorhombic

Column-II

- (P) are two crystal systems
 (Q)
 (R) $a \neq b \neq c$
 (S) $a = b = c$

2.

Column-I

- (A) If radius ratio

$$x = \frac{r_c}{r_a} = 0.155$$

 (B) If 0.225 $x = 0.414$
 (C) If 0.414 $x = 0.732$
 (D) If 0.732 $x = 1$

Column-II

- (P) Co-ordination no. is 8
 (Q) Co-ordination no. is 4
 (R) Co-ordination no. is 6
 (S) Co-ordination no. is 2

3.

Column-I

(Shape of Compound)

- (A) Linear
 (B) Triangular planar
 (C) Square planar
 (D) Octahedral

Column-II

(Co-ordination No.)

- (P) 6
 (Q) 4
 (R) 2
 (S) 3

4.

Column-I

- (A) Rock salt structure
 (B) Zinc blende structure
 (C) Fluorite structure
 (D) Anti fluorite structure

Column-II

- (P) general formula is AB
 (Q) general formula is AB_3
 (R) general formula is A_2B
 (S) general formula is AB_2

5.

Column-I

- (A) Co-ordination no. of Ca^{2+} and F^- in fluorite structure
 (B) C.No. of Zn^{2+} and S^{2-} in zinc blende structure
 (C) C.No. of Cs^+ and Cl^- in CsCl (bcc type) structure
 (D) C.No. of Li^+ and O^{2-} in antifluorite structure

Column-II

- (P) 8, 4
 (Q) 8, 8
 (R) 4, 8
 (S) 4, 4

6.

Column-I**[Bravais Lattice(s)]**

- (A) Primitive, face centered, body centered, end centered
 (B) Primitive, face centered, body centered
 (C) Primitive, body centered
 (D) Primitive only

Column-II**(Crystal System)**

- (P) Cubic
 (Q) Orthorhombic
 (R) Hexagonal
 (S) Tetragonal

7. Column-I (A) Magnetic moment in a paramagnetic substance (B) Magnetic moment in a ferromagnetic (C) Magnetic moment in an antiferromagnetic (D) Magnetic moment in a ferrimagnetic	Column-II (P)  (Q)  (R)  (S) 
8. Column-I (Structure) (A) Rock salt (NaCl) (B) CsCl (C) Zinc blende (ZnS) (D) Anti fluorite (Na₂O)	Column-II (Edge length of unit cell) (P) $a (r_{\text{Cation}} + r_{\text{Anion}})$ (Q) $a \frac{4}{\sqrt{3}} (r_{\text{Cation}} + r_{\text{Anion}})$ (R) $a \frac{2}{\sqrt{3}} (r_{\text{Cation}} + r_{\text{Anion}})$ (S) $a \cdot 2 (r_{\text{Cation}} + r_{\text{Anion}})$
9. Column-I (Ideal structure) (A) Rock salt (B) Zinc blende (C) Fluorite (D) CsCl	Column-II (Packing fraction) (P) 0.729 (Q) 0.756 (R) 0.793 (S) 0.748
10. Column-I (Structure) (A) Rock salt (B) Zinc blende (C) Fluorite (D) Anti fluorite (Na₂O)	Column-II (Voids occupied) (P) 100% tetrahedral voids occupied by cation (Q) 100% tetrahedral voids occupied by anion (R) 100% octahedral voids occupied by cation (S) 50% tetrahedral voids occupied by cation

11.	Column-I (Void representation)	Column-II (Name of void)
(A)		(P) Tetrahedral void
(B)		(Q) Octahedral void
(C)		(R) Cubic void
(D)		(S) Triangular void

Assertion-Reason Type Questions

Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

- (A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** In any ionic solid $[MX]$ with Schottky defects, the number of positive and negative ions are the same.

STATEMENT-2 : Equal number of cation and anion vacancies are present.

2. **STATEMENT 1 :** Amorphous solids are isotropic.

STATEMENT 2 : Amorphous solids lack a regular three-dimensional arrangement of atoms.

- 3. STATEMENT-1 :** Diamond is a covalent solid.
STATEMENT-2 : The co-ordination number of each carbon atom in diamond is 4.
- 4. STATEMENT-1 :** In NaCl structure, Na⁺ ions occupy octahedral holes and Cl⁻ ions occupy ccp.
STATEMENT-2 : The distance of the nearest neighbours in NaCl structure is $a/2$ where a is the edge length of the cube.
- 5. STATEMENT-1 :** For fluorite structure, the F⁻ ions occupy tetrahedral void and Ca²⁺ ions in ccp
STATEMENT-2 : The radius ratio of fluorite structure is 0.414.
- 6. STATEMENT-1 :** In the body centered cubic type structure of CsCl the arrangement of Cs⁺ is present in central octahedral void.
STATEMENT-2 : In CsCl, the Cs⁺ ions are at body centered position and Cl⁻ are at corners.
- 7. STATEMENT-1 :** Electrical conductivity of semi-conductors increases with increasing temperature.
STATEMENT-2 : With increase in temperature, number of electrons which jumps from the valence bond to the conduction band in semi-conductors increase.
- 8. STATEMENT-1 :** FeO is non-stoichiometric with formula Fe_{0.95}O.
STATEMENT-2 : Some Fe²⁺ ions are replaced by Fe³⁺ as $3\text{Fe}^{2+} \rightarrow 2\text{Fe}^{3+}$ to maintain electrical neutrality.
- 9. STATEMENT-1 :** The number of tetrahedral voids is double the number of octahedral voids.
STATEMENT-2 : The size of the tetrahedral void is half of that of the octahedral void.
- 10. STATEMENT-1 :** Due to Frenkel defect there is no effect on density of a solid.
STATEMENT-2 : Ions shift from lattice sites to interstitial sites in Frenkel defect.
- 11. STATEMENT-1 :** Antiferromagnetic substances become paramagnetic on heating to high temperature.
STATEMENT-2 : Heating results in spins of electrons becoming random.

Subjective Problems



- In seven possible crystal system how many crystal system have _____ ?
- In seven possible crystal system how many crystal system have more than one Bravais lattice ?
- How many effective no. of atoms present in bcc unit cell ?
- In solid X atoms goes to corner of the cube and two alternate face centre. Calculate effective number of atom of X in unit cell ?
- Find the distance (in pm) between the body centered atom and one corner atom in an element ($a = 2.32$ pm)

6. The structure of MgO is similar to NaCl. What is the co-ordination number of Mg ?
7. How many number of formula units in a unit cell of CaF_2 ?
8. What is the co-ordination number of sodium in Na_2O ?
9. Density of Li crystal is 0.53 g/cm^3 . The edge length of Li unit-cell is $3.5 \text{ \AA}$. Find out the number of Li atoms in a unit cell. ($N_A = 6.0 \times 10^{23} \text{ mol}^{-1}$, $M = 6.94 \text{ g mol}^{-1}$)
10. What is the co-ordination number of Cl in CsCl structure ?
11. In cubic system how many atoms arrangement exist in nature ?
12. The ionic radii of A and B are $1.7 \text{ \AA}$ and $1.8 \text{ \AA}$ respectively. Find the co-ordination number of A .
13. If edge fraction unoccupied in ideal anti-fluorite structure is x. Calculate the value of Z. Where
$$Z = \frac{x}{0.097}$$
14. Ionic solid NaA crystallise in rock salt type structure. 2.592 gm of ionic solid salt NaA dissolved in water to make 2 litre solution. The pH of this solution is 8. If distance between cation and anion is 300 pm, calculate density of ionic solid (in gm/cm^3).
(Given : $pK_w = 13$, $pK_a(\text{HA}) = 5$, $N_A = 6 \times 10^{23}$)
15. Calculate the value of $\frac{Z}{10}$. Where
 Z Co-ordination number of 2D-square close packing
 Co-ordination number of 2D-hcp
 Co-ordination number of 3D-square close packing
 Co-ordination number of 3D, ABCABC Packing
 Co-ordination number of 3D, ABAB packing

ANSWERS

Level 1

1. (c)	2. (b)	3. (d)	4. (b)	5. (a)	6. (c)	7. (a)	8. (d)	9. (c)	10. (c)
11. (d)	12. (c)	13. (d)	14. (b)	15. (b)	16. (a)	17. (b)	18. (c)	19. (a)	20. (b)
21. (d)	22. (c)	23. (b)	24. (c)	25. (d)	26. (c)	27. (c)	28. (c)	29. (c)	30. (b)
31. (c)	32. (a)	33. (d)	34. (c)	35. (b)	36. (b)	37. (c)	38. (a)	39. (a)	40. (d)
41. (d)	42. (d)	43. (b)	44. (d)	45. (c)	46. (d)	47. (c)	48. (b)	49. (a)	50. (b)
51. (b)	52. (b)	53. (c)	54. (a)	55. (c)	56. (b)	57. (a)	58. (d)	59. (c)	60. (b)
61. (d)	62. (c)	63. (a)	64. (a)	65. (b)	66. (a)	67. (b)	68. (c)	69. (b)	70. (b)
71. (b)	72. (c)	73. (b)	74. (d)	75. (a)	76. (d)	77. (c)	78. (c)	79. (a)	80. (a)
81. (c)	82. (c)	83. (b)	84. (c)	85. (a)	86. (c)	87. (c)	88. (d)	89. (d)	90. (b)
91. (b)	92. (c)	93. (c)	94. (a)	95. (b)	96. (d)	97. (c)	98. (d)	99. (a)	100. (b)
101. (c)	102. (d)	103. (d)	104. (c)	105. (a)	106. (d)	107. (b)	108. (d)	109. (d)	110. (c)
111. (c)	112. (c)	113. (b)	114. (c)	115. (b)	116. (c)	117. (a)	118. (c)	119. (d)	120. (b)
121. (b)	122. (a)	123. (b)	124. (c)	125. (d)					

Level 2

1. (b)	2. (c)	3. (c)	4. (c)	5. (d)	6. (c)	7. (b)	8. (d)	9. (a)	10. (c)
11. (b)	12. (b)	13. (c)	14. (d)	15. (c)	16. (a)	17. (b)	18. (a)	19. (a)	20. (b)
21. (a)	22. (a)	23. (d)	24. (d)	25. (b)	26. (a)	27. (b)	28. (b)	29. (d)	30. (a)
31. (b)	32. (b)								



Level 3

Passage-1:	1. (c)	2. (a)		
Passage-2:	1. (a)	2. (a)	3. (c)	
Passage-3:	1. (b)	2. (b)	3. (a)	4. (a)
Passage-4:	1. (b)	2. (b)	3. (d)	
Passage-5:	1. (b)	2. (b)	3. (a)	
Passage-6:	1. (d)	2. (a)		
Passage-7:	1. (a)	2. (d)	3. (d)	

One or More Answers is/are correct

1. (a,d) 2. (b,c) 3. (a,c) 4. (a,c) 5. (c) 6. (a,c) 7. (a,b) 8. (a,b,d)
 9. (c,d) 10. (a,b,c,d) 11. (a,c,d) 12. (a,b) 13. (a,b,c) 14. (a,b,c) 15. (a,b,c,d) 16. (a,b,c)
 17. (a,c) 18. (a,b,c,d) 19. (a,b,c,d) 20. (a,c)

Match the Column

- | | | | |
|----------|---------|---------|--------|
| 1. A P; | B P, S; | C P, R; | D P, Q |
| 2. A S; | B Q; | C Q, R; | D P |
| 3. A R; | B S; | C Q; | D P |
| 4. A P; | B P; | C S; | D R |
| 5. A P; | B S; | C Q; | D R |
| 6. A Q; | B P; | C S; | D R |
| 7. A P; | B R; | C Q; | D S |
| 8. A S; | B R; | C Q; | D Q |
| 9. A R; | B S; | C Q; | D P |
| 10. A R; | B S; | C Q; | D P |
| 11. A Q; | B Q; | C P; | D R |

Assertion-Reason Type Questions

1. (A) 2. (A) 3. (B) 4. (B) 5. (C) 6. (D) 7. (A) 8. (A) 9. (C) 10. (A)
 11. (A)

Subjective Problems

1. 5	2. 4	3. 2	4. 2	5. 2	6. 6	7. 4	8. 4	9. 2	10. 8
11. 3	12. 8	13. 3	14. 4	15. 4					

Hints and Solutions

Level 1

13. (d) Orthorhombic contains

Simple, Body centre, Face centre & End centre bravais lattices.

14. (b) Cubic-symmetrical

$$90^\circ, a = b = c$$

Triclinic-unsymmetrical

$$90^\circ, a \neq b \neq c$$

15. (b) Tetragonal $a = b \neq c$, 90°

19. (a) Simple cubic cell "Z" $8 \times \frac{1}{8} = 1$

Face centred cubic unit cell

$$Z = 8 \times \frac{1}{8} + 6 \times \frac{1}{2} = 4$$

20. (b) Effective no. of atoms in End centred unit cell

$$8 \times \frac{1}{8} + 2 \times \frac{1}{2} = 2$$

atoms are present at corners as well as at end face's centre of unit cell.

21. (d) Simple cubic Body centred cubic

$$a = 2R \quad \sqrt{3}a = 4R$$

22. (c) Face centred cubic $\sqrt{2}a = 4R$

23. (b) Packing fraction (simple cubic) $\frac{1 \times \frac{4}{3} R^3}{a^3}$

$$= \frac{1}{6} = 0.52$$

$$a = 2R$$

24. (c) Body centred cubic $\frac{2 \times \frac{4}{3} R^3}{a^3} = \frac{\sqrt{3}}{8}$

$$\sqrt{3}a = 4R$$

25. (d) Face centred cubic $\frac{4 \times \frac{4}{3} R^3}{a^3} = 0.74$

$$\sqrt{2}a = 4R$$

26, 27. (c, c)

Packing efficiency

In simple cubic 0.52

Body centred cubic 0.68

Face centred cubic 0.74

28. (c) $a = 2R$
 $R = \frac{0.236}{2} = 0.118 \text{ nm}$

29. (c) $\sqrt{3}a = 4R$

30. (b) $\sqrt{2}a = 4R$

31. (c) $\sqrt{2}a = 4R$
 $a = 2\sqrt{2}R$

32. (a) $Z = 8 \times \frac{1}{8} = 1$

Nearest neighbour 6

33. (d) $Z = 8 \times \frac{1}{8} + 6 \times \frac{1}{2} = 4$

Nearest neighbour 12

34. (c) $Z = 8 \times \frac{1}{8} + 1 = 2$

Nearest neighbour 8

35. (b) $D = \frac{ZM}{N_A a^3} = 6.25 \times \frac{Z \times 12}{6.02 \times 10^{23} (4 \times 10^{-8})^3}$

$Z = 2$ body centred

36. (b) $D = \frac{ZM}{N_A a^3} \quad Z = 2$

37. (c) $D = \frac{ZM}{N_A a^3} = \frac{4 \times 40}{6.02 \times 10^{23} a^3} = 1.83$

$a = 0.525 \text{ nm}$

38. (a) $D = \frac{ZM}{N_A a^3} \quad Z = 4$

$$\sqrt{2}a = 4R$$

39. (a) Refer Q.no. 38

40. (d) Refer Q.no. 38

41. (d) $D = \frac{ZM}{N_A a^3} \quad Z = 2$

42. (d) Refer Q.no. 38


43. (b) Refer Q.no. 38

44. (d) Mole of 'x' $\frac{8}{80} = 0.1$

No. of atoms $0.1 N_A$

No. of unit cell $\frac{0.1 N_A}{4} = 0.025 N_A$

45. (c) $D = \frac{ZM_w}{N_A a^3}$
 $Z = 2, M_w = 96, D = 10.3 \text{ g/cm}^3$
 $\sqrt{3}a = 4R \frac{\sqrt{3}a}{4}$
46. (d) Coordination no. of
 simple cubic 6
 Face centred cubic 12
 Body centred cubic 8
47. (c) F.C.C. unit cell $ABCABCABC \dots\dots$
 packing efficiency 74 %
50. (b) C.N. of A $x(B)$
 C.N. of B $y(A)$
 Formula $A_x B_y$
51. (b) $D = \frac{ZM}{N_A a^3}$
 $Z = 4, a = 2\sqrt{2}R$
52. (b) $D = \frac{ZM}{N_A a^3}$
 $Z = 4, a = 3.5 \times 10^{-8} \text{ cm}, M_w = 59$
54. (a) Packing efficiency of HCP & CCP is same.
56. (b) $Z = 12 \times \frac{1}{6} = 2 \times \frac{1}{2} = 3 \times 6$
58. (d) No. of octahedral hole in a unit cell
 effective no. of atoms per unit cell
 Z
 No. of tetrahedral holes 2 octahedral holes.
59. (c) If sheet is repeated than packing is poor.
60. (b) Refer Q. no. 58
62. (c) Tetrahedral & octahedral holes are present in hcp & ccp.
63. (a) Refer Q. no. 62
67. (b) $\frac{r_c}{r_a} = 0.225$ tetrahedral holes
68. (c) $\frac{r_c}{r_a} = 0.414$ octahedral holes
69. (b) an octahedral holes is formed by 6 sphere.
70. (b) No. of nearest neighbour 8 (All body centre atom w.r.t. corner atom)
 No. of next nearest neighbour 6 (No. of corner atoms along edge w.r.t. to any corner).
71. (b) Refer Q. no. 67, 68
72. (c) Refer Q. no. 68
73. (b) Octahedral sites in fcc are present at each edge centre as well as at body centre.
74. (d) In Na_2O all tetrahedral void are occupied.
75. (a) $\frac{r_c}{r_a} = 0.732$
 $r_c = 0.732 \times 200 = 146.4 \text{ pm}$
76. (d) for cubic void, $\frac{r_c}{r_a} = 0.732$
77. (c) $\frac{r_c}{r_a} = \frac{0.066}{0.140} = 0.47$
 So, cation (Mg^{2+}) fitted in octahedral holes of O^{2-} are present at fcc lattice point.
78. (c) In diamond C-atoms are present in fcc lattice points as well as in alternate tetrahedral void.
79. (a) Refer Q. no. 78
80. (a) For MgO , $\frac{r_c}{r_a} = \frac{0.65}{1.4} = 0.47$
 Mg^{2+} is in octahedral void so C.N. 6
 For MgS , $\frac{r_c}{r_a} = \frac{0.65}{1.84}$
 Mg^{2+} is in tetrahedral void so C.N. 4
81. (c) No. of A $7 \times \frac{1}{8} = \frac{7}{8}$
 No. of B $6 \times \frac{1}{2} = 3$
 $A_{7/8} B_3$ $A_7 B_{24}$
82. (c) No. of S^{2-} $8 \times \frac{1}{8} = 1$ $6 \times \frac{1}{2} = 3$ 4
 No. of Ca^{2+} $\frac{1}{2} \times 8 = 4$
83. (b) No. of A^{2+} $\frac{1}{8} \times 8 = 1$
 No. of B^{3+} $\frac{1}{2} \times 4 = 2$
 No. of O^{2-} $8 \times \frac{1}{8} = 1$ $6 \times \frac{1}{2} = 3$ 4
 (AB_2O_4)
84. (c) No. of C $6 \times \frac{2}{3} = 4$
 No. of A 6
 (C_2A_3)
85. (a) No. of O^{2-} $8 \times \frac{1}{8} = 1$ $6 \times \frac{1}{2} = 3$ 4

- No. of A $\frac{1}{8}$ 8 1
- No. of B $\frac{1}{4}$ 4 1
86. (c) No. of A 8 $\frac{1}{8}$ 1
- No. of B 5 $\frac{1}{2}$ $\frac{5}{2}$
- A_2B_5
87. (c) No. of Cu 8 $\frac{1}{8}$ 6 $\frac{1}{2}$ 4
- No. of Ag 12 $\frac{1}{4}$ 3
- No. of Au 1
- $Cu_4Ag_3Au_1$
88. (d) In NaCl structure
Cl⁻ ions are present at fcc lattice point
Na⁺ ions are present in all octahedral void.
89. (d) In B.C.C. ionic structure
Cation is present in cubic void
Anions are present at each corner of cubic
90. (b) In ZnS, S^{2-} ions are present in F.C.C. Lattice point
 Zn^{2+} ions are present in alternate tetrahedral void.
91. (b) "edge"  r r $\frac{a}{2}$
92. (c) In ideal arrangement of anions
 $\frac{\sqrt{2}a}{2}$ $\frac{4r_a}{2\sqrt{2}}$
93. (c) r_c r_a $\frac{\sqrt{3}a}{2}$
Anion are at corners and cations are at body centre.
94. (a) Distance b/w two nearest Cl $\frac{a}{\sqrt{2}}$
Distance b/w two nearest Na $\frac{a}{\sqrt{2}}$
95. (b) w.r.t. body centre Na⁺ ion
all 12 Na⁺ ions of edge centre.
96. (d) $\frac{r}{r}$ $\frac{1}{2}$ 0.5 F.C.C. lattice point
Cations are at octahedral void
- $(r_c - r_a) \frac{a}{2} = 3$
- $a = 6$
volume $(6)^3 = 216 \text{ pm}^3$
97. (c) In CaF_2 cations are present at F.C.C. lattice points and anions are at all tetrahedral void, antifluorite structure is reverse arrangement of CaF_2 .
98. (d) Refer Q. no. 89.
99. (a) $\frac{r_c}{r_a} = \frac{94}{146} = 0.64$
100. (b) $r_c - r_a = \frac{a}{2}$
102. (d) $\frac{r_A}{r_B} = 0.225$
 $r_A = 45$
103. (d) In Na_2O O^{2-} are at F.C.C. lattice point
Na⁺ are at all tetrahedral void
C.N. of $\text{O}^{2-} = 8$
104. (c) $r_c - r_a = \frac{\sqrt{3}a}{2}$
 $a = \frac{2}{\sqrt{3}} [1.69 \quad 1.81]$
105. (a) $r_c - r_a = \frac{\sqrt{3}a}{2}$
106. (d) $r_c - r_a = \frac{a}{2}$
107. (b) $r_c - r_a = \frac{\sqrt{3}a}{2}$
108. (d) $r_c - r_a = \frac{\sqrt{3}a}{4}$
110. (c) Refer Q. no. 78
111. (c) Due to missing of ions. Density decreases in Schottky defect.
112. (c) In Frenkel defect ions are shifted in its interstitial site but density remains unchanged.
113. (b) Refer Q.no. 112
115. (b) Alkali metal halide shows schottky defect.
116. (c) Two Ag⁺ ions will be replaced by one Cd^{2+} , so there is one vacancy for each Cd^{2+}
117. (a) In Schottky defect ions are missing from their position and in Frenkel defect ions are shifted in their interstitial site from their lattice point.

118. (c) Refer Q. no. 115.

119. (d) Density of crystal always increases due to substitutional impurity defect.

120. (b) $\frac{\sqrt{3}a}{4} (2R)$

121. (b) No. of vacancy No. of Sr^{2+}
 No of Sr^{2+} in 1 mole NaCl $10^7 N_A$

122. (a) Fe^{3+} Fe^{2+}
 $x\%$ $(100 - x)\%$
 $\frac{200}{93} \cdot \frac{3x}{100} = \frac{2(100 - x)}{100}$
 $\% \text{Fe}^{3+} = 15.05\%$

123. (b) Cu^{2+} Cu
 $x\%$ $(100 - x)\%$
 $\frac{20}{18} \cdot \frac{2x}{100} = \frac{(100 - x)}{100}$
 $x = 11.11$

Level 2

1. (b) $1 \cdot \frac{2 \cdot 56}{4r \sqrt{3}}; 2 \cdot \frac{4 \cdot 56}{(2\sqrt{2}r)^3}$
 $\frac{1}{2} \cdot 0.918$

2. (c) $2.32 \cdot \frac{2 \cdot M}{6 \cdot 10^{23} (1.22)^3 \cdot 10^{-21}}$
 $M = 1264 \quad x = 47$

3. (c) Edge not covered by atom $a = 2r$
 also in bcc, $4r = \sqrt{3}a$
 Therefore, edge not covered
 $a = \frac{\sqrt{3}}{2}a \quad a = \frac{2\sqrt{3}}{2}$

Fraction of edge not covered by atoms
 $\frac{2\sqrt{3}}{2} \cdot 0.134$

4. (c) In a simple cubic crystal, $a = 2r$
 Let the radius of the interstitial atom in simple cubic structure be x , then
 $2(r + x) = \sqrt{3}a = \sqrt{3} \cdot 2r$
 So, $x = \sqrt{3}r - r = 0.732r$
 Packing fraction $\frac{\frac{4}{3}r^3}{(2r)^3} = \frac{4}{3} (0.732r)^3$

$$\frac{\frac{4}{3} r^3 [1 - (0.732)^3]}{8r^3}$$

$$\frac{1}{6} [1 - (0.732)^3] = 0.73$$

5. (d) Given $a = 2r = 60.3$ and for bcc, $4r = \sqrt{3}a$
 $a = \frac{\sqrt{3}}{2}a = 60.3 \quad a = 450 \text{ pm}$

Density () $\frac{2 \cdot 48}{6.023 \cdot 10^{23} (4.5)^3 \cdot 10^{-24}}$
 1.75 g/cc

6. (c) $a = 2r = 53.6 \text{ pm} \quad \dots(i)$

also $4r = \sqrt{3}a$
 $a = \frac{\sqrt{3}}{2}a = 53.6$
 $a = \frac{53.6}{2} \cdot \frac{2}{\sqrt{3}} = 400 \text{ pm}$

Density () $\frac{2 \cdot 23}{6.023 \cdot 10^{23} \cdot 4^3 \cdot 10^{-24}}$
 1.19 g/cc

7. (b) Vol. of all atoms in 1.68 g of Ar
 $\frac{1.68}{40} \cdot 6 \cdot 10^{23} \cdot \frac{4}{3} \cdot \frac{22}{7} (1.50^3 \cdot 10^{-8})$
 0.3564

Vol. of solid Ar 1 cm^3
 % of empty space $(1 - 0.3564) \cdot 100$
 64.36


8. (d) Effective no. of atoms of B present in a unit cell 2
 Total volume of B unoccupied by A in a unit cell

$$2 \cdot \frac{4}{3} (R^3 - r^3) \quad \therefore r = \frac{R}{2}$$

Volume of unit cell a^3
 $\frac{4R^3}{\sqrt{3}} = \frac{64}{3\sqrt{3}} R^3 \quad (\because \sqrt{3}a = 4R)$


Desired ratio $\frac{\frac{7}{3} R^3}{\frac{64}{3\sqrt{3}} R^3} = \frac{7\sqrt{3}}{64}$

9. (a)



- $AB = a$ (nearest)
 $BD = \sqrt{2}a$ (next-nearest)
 $CE = \sqrt{3}a$ (next-next-nearest)

10. (c)



- $AB = \frac{\sqrt{3}a}{2}$ (nearest)
 $AC = a$ (next-nearest)
 $CD = \sqrt{2}a$ (next-next-nearest)

11. (b) For atoms of A $8 \frac{1}{8} 1$
 For atoms of B $8 \frac{1}{4} 2$
 For atoms of C $4 \frac{1}{2} 1 3$

12. (b) Let volume of fcc unit cell V

$$A \frac{4 M_A}{N_A \cdot V}$$


$$B \frac{8 M_B}{N_A \cdot V}$$

$$\frac{A}{B} \frac{M_A}{2M_B} \frac{30}{2 \cdot 50} = 0.3$$

$$B \frac{A}{0.3} = 3.33 \text{ g/cc}$$

$$\text{Total density } A \quad B \quad 4.33 \text{ g/cc}$$

13. (c)



$$\frac{BE}{BC} = \cos 30 = \frac{\sqrt{3}}{2}$$

Distance between the centres of two B atoms

$$2 BE = \sqrt{3} a$$

14. (d) 4 formula units of AB are present in a unit cell

$$\text{no. of unit cell} = \frac{4}{40} \frac{N_A}{4}$$

$$0.025 N_A$$

15. (c) Use $\frac{Z \cdot M}{N_A \cdot a^3}$

$$Z = 4; M = 40 \quad 2 \quad 19$$

16. (a) $a = 2(r_+ + r_-)$
 $2 \cdot 297 \text{ pm}$

$$\text{Volume of unit cell} = a^3 = 209.6 \cdot 10^{24} \text{ cm}^3$$

17. (b) $8.27 \frac{4 \cdot 128}{6.023 \cdot 10^{23} a^3}$

$$a = 46.8 \cdot 10^{-9} \text{ cm} = 4.68 \text{ \AA}$$

$$2[r_{\text{O}^{2-}} + r_{\text{Cd}^{2+}}]$$

18. (a) $\frac{r_{\text{Na}^+}}{r_{\text{Cl}^-}} = \frac{1.1 \text{ \AA}}{1.5}$

$$\text{and } \frac{r_{\text{K}^+}}{r_{\text{Cl}^-}} = \frac{5}{7} \text{ or } \frac{r_{\text{K}^+}}{r_{\text{Cl}^-}} = \frac{12}{7}$$

$$\frac{a_1}{a_2} = \frac{2(r_{\text{K}^+} + r_{\text{Cl}^-})}{2(r_{\text{Na}^+} + r_{\text{Cl}^-})} = \frac{12}{7} \cdot \frac{1}{1.5} = 1.143$$

19. (a) Mass of unit cell $d = V$
 $(5 \cdot 10^{-8})^3 = 3.84 \cdot 6 \cdot 10^{23}$
 288 a.m.u.

Mass of formula unit of FeO $56 \quad 16 \quad 72$
 a.m.u.

$$\text{no. of formula unit} = \frac{288}{72} = 4$$

20. (b) Density of crystals decrease due to Schottky defects effective no. of atoms per unit cell in given lattice

$$4 \cdot 1 \frac{0.25}{100} = 3.99$$

$$50 \cdot 3.99$$

$$6 \cdot 10^{23} (0.50 \cdot 10^{-7})^3$$

$$50 \cdot 3.99$$

$$6 \cdot 125 \cdot 0.1$$

$$2.66 \text{ g/cm}^3$$

21. (a) Theoretical density $\frac{Z \cdot M}{N_A \cdot a^3}$

$$\frac{4 \cdot 24 \cdot 10^3}{6 \cdot 10^{23} \cdot (4 \cdot 10^{-10})^3}$$

$$2.5 \cdot 10^3 \text{ kg/m}^3$$

% occupancy $\frac{\text{Observed density}}{\text{Ideal density}} \cdot 100$

$$\frac{2.4 \cdot 10^3}{2.5 \cdot 10^3} \cdot 100 = 96\%$$

23. (d) Octahedral void present at the centre of cube and tetrahedral void is present at $(1/4)^{\text{th}}$ of the distance along each body diagonal.

$$\frac{\sqrt{3}a}{2} \cdot 2 \text{ distance between octahedral}$$

and tetrahedral void.

24. (d) Vol. of unit cell $a^2 \sin 60^\circ \cdot b$

$$173.2 \cdot 10^{-24} \text{ cm}^3$$

$$a = 5 \text{ \AA}$$

Mass of unit cell

$$173.2 \cdot 10^{-24} \cdot 5 \cdot 6 \cdot 10^{23}$$

$$519.6 \text{ g}$$

No. of molecules present in given unit cell

$$\frac{519.6}{259.8} = 2$$

25. (b) In hcp arrangement total height

$$h = 4 \cdot \sqrt{\frac{2}{3}} \cdot r$$

as per given

$$\frac{h}{2} = 2 \cdot \sqrt{\frac{2}{3}} \cdot r = 3.35 \text{ \AA} = 3.35 \cdot 10^{-8} \text{ cm.}$$

Volume of unit cell $24 \sqrt{2} r^3$

$$24 \sqrt{2} \cdot \left(\sqrt{\frac{3}{2}} \cdot \frac{1}{2}\right)^3 \cdot 3.35^3 \cdot 10^{-8}$$

$$2.93 \cdot 10^{-22} \text{ cm}^3$$

Effective no. of atoms in hexagonal unit cell

$$6$$

density $\frac{6 \cdot 12}{6.023 \cdot 10^{23} \cdot 2.93 \cdot 10^{-22}}$

$$0.41 \text{ g/cc}$$

26. (a) No. of Na $4 \cdot 1 \cdot 3$

No. of Cl $4 \cdot 2 \cdot \frac{1}{2} = 3$

27. (b) Along one body diagonal 2X atoms from 2 corners, one Y particle (at the centre of cube) will be removed.

So, effective no. of X particles in a unit cell

$$4 \cdot 2 \cdot \frac{1}{8} = \frac{15}{4}$$

and effective no. of Y particles in a unit cell

$$4 \cdot 1 \cdot 3$$

$$X : Y = \frac{15}{4} : 3$$

or $5 : 4$

28. (b) Effective no. of Na $4 \cdot 1 \cdot 2 \cdot \frac{1}{4} = \frac{5}{2}$

For Cl $4\sqrt{2}a \cdot 4r$

P.F. $\frac{\text{Vol. of effective no. of cations and anions}}{\text{Vol. of unit cell}}$

29. (d) No. of X particles per unit cell 4

No. of Y particles 4

No. of Z particles 8

Along one body diagonal 2X atoms from 2 corners, one Y particle and 2Z particles will be removed.

So, effective no. of X particles in a unit cell

$$4 \cdot \frac{1}{8} = 2 \cdot \frac{15}{4}$$

Effective no. of Y particles in a unit cell

$$4 \cdot 1 \cdot 3$$

Effective no. of Z particles in a unit cell

$$8 \cdot 2 \cdot 6$$

$$X : Y : Z$$

$$\frac{15}{4} : 3 : 6$$

$$5 : 4 : 8$$

30. (a) Effective no. of A particles are removed

$$4 \cdot \frac{1}{8} = 2 \cdot \frac{1}{2} = \frac{3}{2}$$

Effective no. of A particles present in a unit cell

$$4 \cdot \frac{3}{2} = \frac{5}{2}$$

Effective no. of B particles are removed

$$2 \cdot \frac{1}{4} = 1 \cdot \frac{3}{2}$$

Effective no. of B particles present in a unit cell

$$4 \times \frac{3}{2} \times \frac{5}{2}$$

$$A : B$$

$$\frac{5}{2} : \frac{5}{2} \quad \text{or} \quad 1 : 1$$

Level 3

Passage-1

1. (c) P.F. $\frac{2 \times \frac{4}{3} r^3}{\frac{4r}{\sqrt{3}}} \sim 0.68$

% of empty space 100 0.68 100 32

2. (a) P.F. $\frac{4 \times \frac{4}{3} r^3}{\frac{4r}{\sqrt{2}}} 0.7406$

Passage-2

1. (a) $10.6 \times \frac{4 \times 108}{a^3 \times 6.023 \times 10^{23}}$

$a = 40.7 \text{ nm}$

2. (a) Mass of 12×10^{23} atoms 100 gm

Mass of 6.022×10^{23} atom

$$\frac{100}{12 \times 10^{23}} \times 6.023 \times 10^{23}$$

$$50.18$$

$$\frac{4 \times 50.18}{6.023 \times 10^{23} \times (200 \times 10^{10})^3}$$

$$41.66 \text{ g/cm}^3$$

3. (c) By Hit and Trial : For fcc $Z = 4$ (4 pairs of KBr)

$M = 39 \quad 80 \quad 119$

Passage-3

3. (a) 1 to 3

c.c.p. generated f.c.c., octahedral voids = effective number of atoms in a unit cell, tetrahedral voids = 2 effective no. of atoms in a unit cell.

4. (a) O^{2-} is replaced by $X^{8/3-}$ so formula of spinel is $MgAl_2X_3$ and deficiency by one anion.

Passage-6

1. (d) $D_{\text{Theo}} = \frac{ZM}{N_A a^3}$

$$\frac{4 \times 197}{6 \times 10^{23} \times (407 \times 10^{-10})^3}$$

$$\frac{4 \times 197}{6 \times 10^{23} \times 6.74 \times 10^{-23}}$$

$$19.485 \text{ g/cm}^3$$

2. (a) Distance between next neighbour a

407 pm

Passage-7

1. (a) Effective no. of C-atoms per unit cell 8

mass 12 8 96 amu

2. (d) $\frac{\sqrt{3}a}{4} B.L.C-C$

$D = \frac{4 \times 154}{\sqrt{3}} = 355.66 \text{ pm}$

3. (d) $D = \frac{ZM}{N_A a^3} = \frac{8 \times 12}{6 \times 10^{23} \times (355.66 \times 10^{-10})^3}$

3.56 g/cm^3

One or More Answers is/are Correct

4. (a, c)

$$\frac{r_{\text{Na}}}{r_{\text{Cl}}} = \frac{95}{181} = 0.52$$

Cl ion at F.C.C. lattice point

Na ion are at octahedral void

C.No. of NaCl 6 : 6

$$r_c : r_a = \frac{a}{2}$$

7. (a, b)

Refer Q. no. 4

11. (a, c, d)

CsCl ionic solid is B.C.C.

12. (a, b)

On increasing temperature C.No.

C.No. change 8 : 8 to 6 : 6

On increasing pressure C.No.

14. (a, b, c)

Refer Q. no. 4

16. (a, b, c)

Refer Q. no. 90 (Level-1)

17. (a, c)

Refer Q. no. 97 (Level-1)

18. (a, b, c, d)

(A) 2D-hexagonal packing

$$\text{Packing efficiency} = \frac{R^2 \cdot 6 \cdot \frac{R^2}{3}}{6 \cdot \frac{\sqrt{3}(2R)^3}{4}}$$

$$\text{(B) Simple cubic} \quad \frac{1 \cdot \frac{4}{3} R^3}{a^3}$$

$$\text{H.C.P. (3D)} \quad \frac{6 \cdot \frac{4}{3} R^3}{24\sqrt{2}R^3}$$

$$\text{F.C.C. (3D)} \quad \frac{4 \cdot \frac{4}{3} R^3}{(2\sqrt{2}R)^3}$$

20. (a, c)

$$\frac{r_{A^2}}{r_{B^2}} = \frac{100}{300} = 0.333 \quad \frac{r_C}{r_D} = \frac{240}{480} = 0.5$$

B^2 are at F.C.C. D at F.C.C. lattice point
lattice point

and A^2 ions are C in all octahedral void
alternate

tetrahedral void

C.No. of AB = 4 : 4 C.No. of CD = 6 : 6

Subjective Problems

1. Cubic 90
Tetragonal 90
Orthorhombic 90
Rhombohedral 90
Hexagonal 90
2. Cubic, tetragonal, orthorhombic,
monoclinic
3. $Z = 8 \cdot \frac{1}{8} = 1$
4. No. of x-atoms = $8 \cdot \frac{1}{8} = 1$
5. Distance b/w body centred and corner atom
 $\frac{\sqrt{3}a}{2}$
6. Mg^{2+} is in octahedral void
So its C.No. = 6

7. No. Ca^{2+} per unit cell = 4

No. of F per unit cell = 8

No. of CaF_2 unit per unit cell = 48. Na^+ ions are present in tetrahedral void in
 Na_2O , So C.No. of Na^+ = 4

$$9. D = \frac{ZM}{N_A a^3}$$

 $Z = 2$ 10. Cl^- ions are at corners and Cs^+ at body
centreSo C.No. of Cl^- = 8 Cs^+ 11. Simple cubic
body centred cubic
face centred cubic

$$12. \frac{r_A}{r_B} = \frac{1.7}{1.8} = 0.944$$

B^2 are at corner of cube A^2 at body centre
C.No. of A^2 = 8

13. Fraction of edge unoccupied = $\frac{a - 2R}{a}$

$$a = 2\sqrt{2}R \quad X = \frac{2(\sqrt{2} - 1)}{2\sqrt{2}} = \frac{0.414}{1.414} = 0.293$$

$$Z = \frac{X}{0.097} = \frac{0.293}{0.097} = 3$$

14. NaA is salt of SBWA

$$pH = \frac{pK_w}{2} - \frac{1}{2}(pK_a + \log c)$$

$$8 = 6.5 - \frac{1}{2}(5 + \log c)$$

$$1.5 = 2 - \frac{5}{2} \log c$$

$$\log c = \frac{2}{2} = 1$$

$$c = 0.01 \quad \frac{2.592}{M \cdot 2}$$

$$M = \frac{2.592}{2 \cdot 0.01} = 129.6$$

$$D = \frac{ZM}{N_A a^3} \quad Z = 4$$

$$4 \text{ g/cm}^3 \quad a = 2 \cdot 300 = 600 \text{ pm}$$

$$15. \frac{Z}{10} = \frac{4 \cdot 6 \cdot 6 \cdot 12 \cdot 12 \cdot 40}{40} = 4$$

SURFACE CHEMISTRY

Terms Related to Adsorption and Absorption

- ❖ **Adsorption** : The accumulation of molecular species at the surface rather than bulk of solid or liquid is known as adsorption.
- ❖ **Adsorbate** : The molecular species or substance which accumulates at the surface is known as adsorbate.
- ❖ **Adsorbent** : The surface at which accumulation of adsorbate takes place is known as adsorbent.

Ex : Water vapours (adsorbate) adsorb at the surface of silica gel (adsorbent).

- ❖ **Desorption** : The process of removing an adsorbed substance from adsorbent is known as desorption. When equilibrium is established then rate of adsorption and rate of desorption are equal.
- ❖ **Absorption** : When atoms, molecules or ions enter in bulk phase of solid or liquid is known as absorption.

Ex. Water vapours absorbed by anhydrous CaCl_2 .

- ❖ **Sorption** : When adsorption and absorption takes place simultaneously then phenomena is known as sorption.

Ex. Dyeing of fabric

Characteristics of Adsorption

- ❖ Adsorption is surface phenomena.
- ❖ In adsorption enthalpy decreases.
- ❖ In adsorption entropy decreases.
- ❖ When decrease in free energy takes place then adsorption takes place.

Types of Adsorption

(1) Based on concentration :

- ❖ Positive adsorption : If the concentration of adsorbate is more at the surface as compared to its concentration in the bulk phase then it is called positive adsorption
- ❖ Negative adsorption : If concentration of adsorbate is less at the surface as compared to its concentration in the bulk phase then it is called negative adsorption

(2) Based on forces existing between adsorbate molecule and adsorbent :

- ❖ **Physical adsorption** : If the forces of attraction existing between adsorbate and adsorbent are vander Waal's forces, then adsorption is called physical adsorption.
- ❖ **Chemical adsorption** : If the forces of attraction existing between adsorbate particles and adsorbent are almost of the same strength as chemical bonds then adsorption is called chemical adsorption.

Comparison between physisorption and chemisorption

S.No.	Physisorption	Chemisorption
1.	Low heat of adsorption usually in range of 20 - 40 kJ/mol.	High heat of adsorption in the range of 80 - 240 kJ/mol.
2.	Forces of attraction are vander Waal's forces.	Forces of attraction are chemical bond forces.
3.	It is reversible in nature.	It is irreversible in nature.
4.	It usually takes place at low temp and it decreases with increasing temp.	It takes place at high temperature and it increase with increasing temperature.
5.	More easily liquefiable gases are adsorbed readily.	Gases which can react with the adsorbent show chemisorption.
6.	It forms multimolecular layers.	It forms monomolecular layer.
7.	It does not require any activation energy.	It requires activation energy.
8.	It is not specific.	It is highly specific.
9.	It increases with increase the surface area of adsorbent.	It also increases with increase the surface area of adsorbent.

Factors Affecting Adsorption


- ❖ **Nature of the gas** : Easily liquefiable gases adsorb to greater extent.
- ❖ **Effect of nature of adsorbent** : When adsorbent is activated then extent of adsorption increases.
- ❖ **Specific area of the solid** : Greater the specific area of the solid, greater would be its adsorbent power.
- ❖ **Effect of pressure of the gas** : The adsorption causes a net decrease in pressure as the gas gets adsorbed and thus the increase in pressure favours the process of adsorption.

Adsorption Isotherm

Graph between extent of the adsorption $\frac{x}{m}$ & pressure (P) at a given temperature is called adsorption isotherm. There are two types of adsorption isotherms

(i) Freundlich Adsorption isotherm :

$$\text{Freundlich's Equation } \frac{x}{m} = k P^{1/n}$$




where x = mass of adsorbate adsorbed, m = mass of adsorbent, p = pressure

❖ **At low pressure :** $\frac{x}{m} = k P$

❖ **At intermediate pressure :** $\frac{x}{m} = k P^{1/n}$, where $n > 1$

❖ **At high pressure :** $\frac{x}{m} = \text{constant}$

$$\log \frac{x}{m} = \log k + \frac{1}{n} \log p$$



(ii) Langmuir Adsorption isotherm : According to Langmuir

- ❖ There is adsorption of gas molecules on the surface of the solid.
- ❖ There is desorption of the adsorbed molecules from the surface of the solid.
- ❖ There is formation of unimolecular layer and so it is chemisorption
- ❖ A dynamic equilibrium is attained when rate of adsorption = rate of desorption.

$$\frac{x}{m} = \frac{ap}{1 + bp}$$


Where a & b Langmuir parameters.

Case-I At very high pressure

$$\frac{bp}{1 + bp} \approx 1, \quad \frac{x}{m} = \frac{ap}{bp} = \frac{a}{b} = \text{constant}$$

Case-II At very low pressure


$$\frac{bp}{1 + bp} \approx bp, \quad \frac{x}{m} = ap$$



Adsorption Isobar


Graph between extent of the adsorption $\frac{x}{m}$ & T at a given pressure is called adsorption isobar.

A physical adsorption isobar shows a decrease in x/m as the temperature rises.



Physical adsorption isobar

The isobar of chemisorption shows an increase in the beginning and then decrease as the temperature rises.



Chemical adsorption isobar

Application of Adsorption

- ❖ Activated charcoal is used in gas masks to remove poisonous gases such as CH_4 , CO etc.
- ❖ Animal charcoal is used as decolorizer in manufacture of sugar.
- ❖ Silica gel is used for removing moisture and controlling humidity.
- ❖ Some industrial processes take place in the presence of catalysts which are based upon adsorption.
- ❖ Chromatographic purification of compounds which is based upon adsorption.
- ❖ The ion exchange resins used for removing hardness of water are also based upon adsorption.


Catalyst

1. Catalyst : A substance which influence the rate of reaction is known as catalyst provided it remains unchanged in amount as well as in composition.

- ❖ **Positive catalyst :** The catalyst which increase the rate of chemical reaction is known as positive catalyst. Fe act as positive catalyst in following reaction.



- ❖ **Negative catalyst :** The catalyst which decrease the rate of chemical reaction is known as negative catalyst. $\text{C}_2\text{H}_5\text{OH}$ act as negative catalyst in following reaction.




- ❖ **Auto catalyst :** When one of product act as catalyst then such type of catalyst is known as auto catalyst. Mn^{2+} act as auto catalyst in following reactions



2. Catalyst promoters : Substance which increase the activity of catalyst is known as catalyst promoters. Mo act as catalyst promoters in following reaction.



3. Homogeneous catalytic reaction : If catalyst is present in same phase as reactant then it is called homogenous catalytic reaction.



4. Heterogeneous catalytic reaction : If catalyst is present in different phase as reactant then it is called heterogenous catalytic reaction.




5. Adsorption theory of heterogeneous catalysis : The mechanism of heterogenous catalysis involves the following steps :

- ❖ Adsorption of reactant molecules on the catalyst surface.
- ❖ Occurrence of a chemical reaction through the formation of an intermediate.
- ❖ Desorption of products from the catalyst surface.
- ❖ Diffusion of products away from the catalyst surface.

Enzyme Catalyst

Complex nitrogenous compounds which are produced by living plants and animals are known as enzymes. The characteristic features of enzyme catalyst are :

- ❖ They are highly efficient.
- ❖ They are highly specific in nature.
- ❖ They are highly active under optimum temperature.
- ❖ They are highly active under optimum pH.
- ❖ There is increase in their activity in the presence of activators and coenzymes.
- ❖ They are influenced by inhibitors and poisons.



Mechanism of Enzyme Catalysis

Step 1 : Binding of enzyme to substrate to form an activated complex $\text{E} + \text{S} \rightleftharpoons \text{ES}$

Step 2 : Decomposition of the activated complex to form products $\text{ES} \rightarrow \text{E} + \text{P}$

Colloids

Depending on size of particles mixture are divided into three classes :

- ❖ **True solution :** In true solution size of particles is less than 1 nm.
- ❖ **Suspension :** In suspension size of particles is greater than 1000 nm.
- ❖ **Colloidal solution :** In colloidal solution size of particles lies from 1 nm to 1000 nm.

It consists two phase *i.e.*, dispersed phase and dispersion medium.

Dispersed Phase : It is the component present in small proportion and is just like a solute in a true solution. For example, in the colloidal state of sulphur in water, the former acts as a dispersed phase.

Dispersion Medium : It is normally the component present in excess and is just like a solvent in a solution.

The particles of the dispersed phase are scattered in the dispersion medium in a colloidal system.

Classification of Colloids

1. Based on Physical State of Dispersed Phase & Dispersion Medium :

Dispersed phase	Dispersion medium	Colloidal system	Examples
Gas	Liquid	Foam or froth	Soap sols, lemonade froth, whipped cream.
Gas	Solid	Solid foam/Sol	Pumice stone, styrene, foam, foam rubber.
Liquid	Gas	Aerosols of Liquids	Fog, clouds, fine insecticide sprays.
Liquid	Liquid	Emulsions	Milk
Liquid	Solid	Gels	Cheese, butter, boot polish, table jellies.
Solid	Gas	Aerosols of Solid	Smoke, dust
Solid	Liquid	Sols	Most paint starch dispersed in water, gold sol, muddy water, inks.
Solid	Solid	Solid sols	Ruby glass, some gem stones.

2. Based on Interaction or Affinity of Phases :

(a) Lyophilic Colloids : The colloidal system in which the particle of dispersed phase have great affinity for the dispersion medium, are called lyophilic

(b) Lyophobic colloids : The colloidal system in which the dispersed phase have no affinity for the dispersion medium are called lyophobic.

	Property	Lyophilic sols	Lyophobic sols
1.	Nature	Reversible	Irreversible
2.	Preparation	They are prepared very easily by shaking or warming the substance with dispersion medium.	They are difficult to prepare, Special methods are used.
3.	Stability	They are very stable and are not easily coagulated by electrolytes.	They are generally unstable and get easily coagulated on addition of electrolytes.
4.	Charge	Particles carry no or very little charge depending upon the pH of the medium.	Colloidal particles have characteristic charge (positive or negative)
5.	Viscosity	Viscosity is much higher than that of the medium.	Viscosity is nearly the same as that of the medium
6.	Surface Tension	Surface tension is usually less than that of the medium.	Surface tension is nearly the same as that of the medium.

7.	Migration in electric field	The particles may or may not migrate in an electric field.	The colloidal particles migrate either towards cathode or anode in an electric field.
8.	Solvation	Particles are heavily solvated.	Particles are not solvated.
9.	Visibility	The particles can not be seen under ultra microscope.	The particles though invisible, can be seen under ultra microscope.
10.	Tyndall effect	Less distinct.	More distinct.
11.	Action of electrolyte	Large amount of electrolyte is required to cause coagulation.	Small amount of electrolyte is sufficient to cause coagulation.
12.	Examples	Mostly organic substances e.g., starch, gums, proteins, gelatin etc.	Generally inorganic substance e.g., metal sols, sulphides and oxides sols.

3. Based on Type of Particles of the Dispersed Phase

❖ **Multimolecular colloids** : The multimolecular colloidal particles consists of aggregate of atoms of small molecules with diameters less than 10^{-9} m or 1 nm.

For example, a sol of gold contains particles of various sizes having several atoms.

❖ **Macromolecular colloids** : The macromolecular colloidal particles themselves are large molecules. They have very high molecular mass varying from thousands to millions. These substances are generally polymers.

For example, starch, cellulose and proteins, polyethene, nylon etc.

❖ **Associated colloids** : Certain substances behave as strong electrolytes at low concentration but at higher concentrations these substances exhibit colloidal characteristics due to the formation of aggregated particles. These aggregated particles are called micelles, which are also known as associated colloids.

Kraft Temperature : The formation of micelles takes place only above a particular temperature called Kraft temperature.

Critical Micelle Concentration : Formation of micelle takes place above kraft temperature and above a particular concentration called the critical micelle concentration (CMC).

For soaps, the CMC is about 10^{-4} M to 10^{-3} M.

4. Method of Preparation of Colloids :

❖ **Mechanical Dispersion** : In this method the substance is first finely powdered and a suspension is made by shaking the powdered substance with the dispersion medium. This suspension is then passed through a colloidal mill where suspension is converted into colloids. The colloidal solutions of rubber, ink, paints and varnishes are prepared by this method.

❖ **Electrical Dispersion (Bredig's arc method)** : In this method an electric arc is struck between two metallic rods and kept under the dispersion medium. The particles of the metal break away from the rods and disperse in the liquid. Colloidal solutions of Au, Pt, Ag, Cu and such other metals can be prepared by this method.

❖ **Peptization** : The process in which freshly formed precipitate converts into a sol by the addition of suitable electrolyte is known as peptization.

(a) Added electrolyte is known as peptizing agent.

(b) The peptization action is due to preferential adsorption of common ion.

Ex.	AgI ppt	AgNO ₃ Peptising agent	[AgI] Ag : NO ₃
	AgI ppt	KI Peptising agent	[AgI] I : K ⁺

❖ Chemical methods :

Double decomposition :	As ₂ O ₃	3H ₂ S	As ₂ S ₃ Yellow sol	3H ₂ O
Oxidation :	2H ₂ S	SO ₂	2H ₂ O	3S Sol
Reduction :	2AuCl ₃	3SnCl ₂	3SnCl ₄	2Au Sol
Hydrolysis :	FeCl ₃	3H ₂ O	Fe(OH) ₃ Red sol	3HCl

5. Purification of Colloidal Solution :

- ❖ **Dialysis** : The process of separating the particles of colloid from those of crystalloid, by means of diffusion through a suitable membrane is called dialysis. Its principle is based upon the fact that colloidal particles can not pass through a parchment or cellophane membrane, while the ions of the electrolyte can pass through it.
- ❖ **Electrodialysis** : The ordinary process of dialysis is slow. To increase the process of purification, the dialysis is carried out by applying electric field. This process is called electrodialysis. The artificial kidney machine works on the principle of dialysis.

6. Properties of Colloidal Solution :

(i) Physical Properties :

- ❖ **Heterogeneity** : Colloidal solutions are heterogeneous in nature consisting of two phases viz, the dispersed phase and the dispersion medium.
- ❖ **Filterability** : Colloidal particles readily pass through ordinary filter papers. It is because the size of the pores of the filter paper is larger than that of the colloidal particles.
- ❖ **Non-settling nature** : Colloidal solutions are quite stable as the colloidal particles remain suspended in the dispersion medium indefinitely. Thus there is no effect of gravity on the colloidal particles.
- ❖ **Colour** : The colour of the colloidal solution is not always the same as the colour of the substances in the bulk. The colour of the colloidal solution depends upon the following factors :
 - Size and shape of colloidal particles.
 - Wavelength of the source of light.
 - Method of preparation of the colloidal solution.
 - Nature of the colloidal solution.
 - The way an observer receives the light.
- ❖ **Stability** : Colloidal solutions are quite stable. Only a few solutions of larger particles may settle but very slowly.

(ii) Mechanical Properties :

❖ **Brownian movement** : It involves motion of colloidal particles in zig-zag paths. It is due to the moving molecules of the dispersion medium which are constantly colliding with colloidal particles.

- (i) It is not observed in ordinary suspension.
- (ii) It offers a visible proof of the random kinetic motion of molecules in a liquid.
- (iii) It provides a direct demonstration of ceaseless motion of molecule as postulated by kinetic theory.
- (iv) It provide stability to colloidal sol by not allowing them to settle down.

❖ **Sedimentation** : Under the influence of gravity, colloids tends to settle down very slowly. This phenomena is known as sedimentation. It can be accelerated by ultracentrifugation.

(iii) Optical properties :

❖ **Tyndall Effect** : The scattering of light by colloidal particles is known as Tyndall effect.

The colloidal particles first absorb light and then a part of the absorbed light is scattered from the surface of the colloidal particles. Maximum scattered intensity is being in a plane at right angles to the plane of incident light. The path becomes visible when seen from that direction.

(iv) Electrical properties : Colloidal particles carry an electric charge which is equal and opposite to dispersion medium.

❖ **Electrophoresis** : The movement of colloidal particles towards a particular electrode in an electric field is known as electrophoresis. It provides information about the nature of the electrical charge on the colloidal particle.

❖ **Electro-osmosis** : When electrophoresis of dispersed particles in a colloidal system is prevented by suitable means, it is observed that the dispersion medium itself begins to move in an electric field. This phenomenon is called electro-osmosis.

Electrically charged sols

	Positively charged sols	Negatively charged sols
1.	Ferric hydroxide, aluminium hydroxide	Metals such as Pt, Cu, Au, Ag, Sb ₂ S ₃ , CdS Metals sulphides, <i>e.g.</i> , arsenic sulphide
2.	Basic dyes such as methylene blue	Starch, clay, silicic acid, gum, gelatin, charcoal
3.	Haemoglobin	Acid dyes, such as eosin, congo Red

❖ **Electrical double layer** : The surface of a colloidal particle acquires a positive or a negative charge by selective adsorption of ions carrying +ve or -ve charges respectively. The charged layer attracts counter ions from the medium which forms a second layer. Thus, an electrical double layer is formed on the surface of the particles.

[AgI] I ⁻	:	K ⁺
[AgI] Ag ⁺	:	NO ₃ ⁻
[Fe(OH) ₃] Fe ³⁺	:	3Cl ⁻
Solid	+	-
	+	-
	+	-
	+	-

Selective adsorption of positive ions solid

Electro kinetic potential : The potential difference between the fixed charge layer and diffused layer of opposite charge is called electro kinetic potential or zeta potential.

❖ **Coagulation :** The stability of the colloidal system is due to the existence of charged particles. If the charges are neutralised or destroyed, then the colloidal solution gets precipitated. This is called Coagulation or Flocculation. It can be done by

- adding electrolyte
- mutual action of sols
- persistent dialysis
- By cooling
- By electrophoresis

Coagulation value : The minimum concentration of the electrolyte required in millimoles per litre of solution to cause coagulation or flocculation is called coagulation value.

Hardy Schulze Rule : According to this rule the coagulating power of the active ion increases with the valency of the active ion.

Coagulating power of cations : Al³⁺ Ba²⁺ or Mg²⁺ Na or K .

Coagulating power of anion : [Fe(CN)₆]⁴⁻ PO₄³⁻ SO₄²⁻ Cl

❖ **Isoelectric Point of Colloid :** The hydrogen ion concentration at which the colloidal particles are neither positively charged nor negatively charged (i.e., uncharged) is known as isoelectric point of the colloid.

❖ **Protective colloids :** In presence of lyophilic colloids like starch no coagulation of lyophobic colloids takes place and lyophilic colloids act as protective colloids. Greater protecting power of a lyophilic colloid lesser would be gold number.

Gold Number : It is defined as the minimum amount of the protective colloid in milligrams which must be added to 10 mL solution of standard gold sol, which prevents coagulation on adding one mL of 10% NaCl solution.

❖ **Emulsions :** An emulsion is a colloidal solution of a liquid. It may be defined as a heterogeneous system consisting of more than one immiscible liquids dispersed in one another in the form of droplets

Oil in water type emulsion (O/W) : In this emulsion, oil is the dispersed phase and water is the dispersion medium. It is denoted by O/W or O in W. For example, milk (liquid fat dispersed in water), vanishing cream, etc.

Water in oil type : In this emulsion, water is the dispersed phase and oil is the dispersion medium. It is denoted by W/O or W in O. For example, butter, cod liver oil, cold cream, etc.

Distinction between Two Types of Emulsions

Dye test : It involves the addition of oil soluble dye to the emulsion under experiment. If the emulsion acquires the colour of the dye readily, it is water-in-oil type emulsion and if the emulsion remains colourless, it is oil-in-water type emulsion.

Dilution test : As a general rule, an emulsion can be diluted with the dispersion medium while the addition of the dispersed phase forms a separate layer. Thus, if an emulsion can be diluted with oil, it is water-in-oil type.

Preparation of emulsion (Emulsification) : Emulsification is the process which involves the preparation of emulsion. Generally, an emulsion is prepared by subjecting a mixture of the immiscible liquid to a distinct layers upon standing. This substance which stabilizes the emulsion is called emulsifier or emulsifying agent. The commonly used emulsifying agents are soaps, detergents and lyophilic colloids, casein.

Function of emulsifier : The main function of emulsifier or emulsifying agents is to lower the interfacial tension between oil and water and thus helps the intermixing of two liquids.

Demulsification : The process which involves the breaking of an emulsion into two separate liquid layers is called demulsification.

- ❖ **Gels :** Colloidal system in which liquids are the dispersed phase and solid act as the dispersion medium are called gels. The common examples are : boot polishes, jelly, gum arabic, agar agar, processed cheese and silicic acid.

When the gels are allowed to stand for a long time, they give out small quantities of trapped liquids with accumulate on its surface. This action of gels is known as Synresis or Weeping.

❖ **Uses of Colloids**


- (i) Medicines
- (ii) Dyes
- (iii) Rubber industry
- (iv) Formation of delta
- (v) Purification of water
- (vi) Artificial rain
- (vii) Smoke precipitation
- (viii) Sewage disposal
- (ix) Cleansing action of soap and detergent
- (x) In Photography



Level 1

1. The size of particles in suspension, true solution and colloidal solution varies in the order :
(a) suspension > colloidal > true solution
(b) true solution > suspension > colloidal
(c) suspension > colloidal > true solution
(d) none of these
2. A colloidal system has what size of particles?
(a) 10^{-4} m to 10^{-10} m
(b) 10^{-5} m to 10^{-7} m
(c) 10^{-9} m to 10^{-12} m
(d) 10^{-6} m to 10^{-9} m
3. Which are not purely surface phenomena?
(a) Adsorption, surface tension
(b) Surface tension, viscosity
(c) Adsorption, viscosity
(d) Absorption, viscosity
4. Adsorbed acetic acid on activated carbon is :
(a) adsorber
(b) absorber
(c) adsorbent
(d) adsorbate
5. Adsorption due to strong chemical forces is called :
(a) chemisorption
(b) physisorption
(c) both (a) and (b)
(d) none of these
6. Adsorption of gases on solid surface is exothermic because :
(a) free energy increases
(b) entropy decreases
(c) entropy increases
(d) interaction developed between gas and solid particles
7. The nature of bonding forces in adsorption are :
(a) purely physical such as van der Waals' forces
(b) purely chemical
(c) both chemical and physical are possible
(d) none of these
8. Which one of the following is not applicable to chemisorption?
(a) Heat of adsorption is negative
(b) It takes place at high temperature
(c) It is reversible
(d) It forms mono-molecular layer
9. Which one of the following characteristics is correct for physical adsorption?
(a) It is very specific
(b) Adsorption on solids is irreversible
(c) Adsorption decreases with decrease in temperature
(d) Generally both enthalpy and entropy of adsorption are negative
10. Which of the following statement is not correct?
(a) Physical adsorption is due to van der Waals' forces
(b) Physical adsorption is irreversible

- (c) Chemical adsorption increases with increase in temperature upto certain limit than decreases
- (d) Enthalpy of adsorption ($|\Delta H|$) for a chemical adsorption is greater than that of physical adsorption
11. Which gas will be adsorbed on a solid to greater extent?
- (a) Having non-polar molecule
(b) Having highest critical temperature
(c) Having lowest critical temperature
(d) Having lowest critical pressure
12. Which of the following factors affects the adsorption of a gas on solid?
- (a) Critical temperature (T_c)
(b) Temperature of gas
(c) Pressure of gas
(d) All of these
13. Which gas is adsorbed to maximum amount by activated carbon?
- (a) $H_2(g)$
(b) $He(g)$
(c) $CO(g)$
(d) $CO_2(g)$
14. The volume of gases NH_3 , CO_2 and H_2 adsorbed by one gram of charcoal at 300 K are in order of:
- (a) $H_2 > CO_2 > NH_3$
(b) $NH_3 > H_2 > CO_2$
(c) $NH_3 > CO_2 > H_2$
(d) $CO_2 > NH_3 > H_2$
15. Which of the following is used to adsorb water?
- (a) Silica gel
(b) Calcium acetate
(c) Hair gel
(d) Anhydrous $CaCl_2$
16. Absorption and adsorptions are respectively:
- (a) surface phenomena, bulk phenomena
(b) bulk phenomena, surface phenomena
(c) both are bulk phenomena
(d) both are surface phenomena
17. Adsorption is multilayer in case of:
- (a) physical adsorption
(b) chemisorption
(c) both (a) and (b)
(d) none of these
18. Reversible adsorption is:
- (a) chemical adsorption
(b) physical adsorption
(c) both (a) and (b)
(d) none of these
19. The effect of pressure on adsorption is high if:
- (a) temperature is low
(b) temperature is high
(c) temperature is very high
(d) larger charcoal piece is taken
20. Sorption is the phenomenon:
- (a) reverse of adsorption
(b) reverse of absorption
(c) when adsorption and absorption takes place simultaneously
(d) none of these
21. Which of the following adsorption isotherms represents the adsorption of a gas by a solid involving multilayers of formation? (P_s saturation pressure)



22. A plot of $\log \frac{x}{m}$ against $\log P$ for the adsorption of a gas on a solid gives a straight line with slope equal to:
- (a) $\frac{1}{n}$ (b) n (c) $\log K$ (d) K
23. The heat evolved in chemisorption lies in the range (in kJ/mol) of :
- (a) 80 to 240 (b) 20 to 40
(c) 40 to 80 (d) 20 to 100
24. The heat evolved in physisorption lies in the range (in kJ/mol) of :
- (a) 20–40 (b) 40–100
(c) 100–200 (d) 200–400
25. According to the adsorption theory of catalysis, the speed of the reaction increases because :
- (a) in the process of adsorption, the activation energy of the molecules becomes large
(b) adsorption produces heat which increases the speed of the reaction
(c) adsorption lowers the activation energy of the reaction
(d) adsorption increases the activation energy of the reaction
26. 3.6 gram of oxygen is adsorbed on 1.2 g of metal powder. What volume of oxygen adsorbed per gram of the adsorbent at 1 atm and 273 K?
- (A) 0.19 L g^{-1} (b) 1 L g^{-1}
(C) 2.1 L g^{-1} (d) None of these
27. A catalytic poison renders the catalyst ineffective because:
- (a) It is preferentially adsorbed on the catalyst
(b) It adsorbs the molecules of the reactants
(c) It combines chemically with the catalyst
(d) It combines chemically with one of the reactants
28. The catalyst used in the hydrogenation of oils is:
- (a) Fe (b) Ni (c) Pt (d) V_2O_5
29. The function of zymase is to:
- (a) change starch into sugar
(b) ferment glucose to alcohol and CO_2
(c) change malt sugar into glucose
(d) change starch into malt sugar and dextrin
30. The conversion of maltose to glucose is possible by the enzyme:
- (a) zymase (b) lactase
(c) maltase (d) diastase
31. Shape selective catalysis is a reaction catalysed by:
- (a) enzymes (b) ziegler-Natta Catalyst
(c) zeolites (d) platinum

32. The process which is catalysed by one of the product is called:
(a) acid-base catalysis (b) autocatalysis
(c) negative catalysis (d) homogeneous catalysis
33. An inhibitor is essentially:
(a) a negative catalyst (b) a heterogeneous catalyst
(c) an auto catalyst (d) a homogeneous catalyst
34. A catalyst in the finely divided form is most effective because:
(a) less surface area is available
(b) more active sites are formed
(c) more energy gets stored in the catalyst
(d) none of these
35. Identify the correct statement regarding enzymes:
(a) Enzymes are specific biological catalysts that normally works at high temperature
(b) Enzymes are normally heterogeneous catalysts which decrease the reaction rate
(c) Enzymes are specific biological catalysts with low molar masses
(d) Enzymes are specific biological catalysts that are very specific in nature
36. A liquid leaves no residue when passed through the ultra-filter paper. The liquid is:
(a) a suspension (b) oil (c) a colloidal sol (d) a true solution
37. Crystalloids differ from colloids mainly in respect of :
(a) electrical behaviour (b) particle nature
(c) particle size (d) solubility
38. Surface tension of lyophilic sols is:
(a) lower than water (b) more than water
(c) equal to water (d) none of these
39. Which one of the following is not used for preparing lyophilic sols?
(a) Starch (b) Gum
(c) Gelatin (d) Metal sulphide
40. Which one of the sols acts as protective colloid?
(a) As_2S_3 (b) Gelatin (c) Au (d) $\text{Fe}(\text{OH})_3$
41. Which one of the following is lyophilic colloid?
(a) Pt (b) Gum (c) Fog (d) Blood
42. Small liquid droplets dispersed in another liquid is called:
(a) suspension (b) emulsion
(c) gel (d) true solution
43. Which of the following is an example of associated colloid?
(a) Protein in water (b) Soap in water
(c) Rubber in benzene (d) FeCl_3 in H_2O
44. Among the following, select the properties of lyophilic colloidal sols:
(a) viscosity same as that of the medium
(b) extensive hydration takes place
(c) particles migrate either towards cathode or anode in an electric field
(d) particle can be readily detected under ultramicroscope
45. Fog is an example of colloidal system of:
(a) liquid in a gas (b) gas in a liquid
(c) gas in a solid (d) solid in a liquid

46. Colloidal systems are not classified on the basis of :
(a) molecular size
(b) nature of the particles
(c) surface tension value
(d) interaction between disperse phase and dispersion medium
47. All colloids:
(a) are suspensions of one phase in another
(b) are two-phase systems
(c) contain only water-soluble particles
(d) are true solutions
48. Which of the following is a lyophobic colloid?
(a) Gelatin (b) Sulphur
(c) Starch (d) Gum arabic
49. Which of the following is not a gel?
(a) Cheese (b) Jellies (c) Curd (d) Milk
50. Cleaning action of soap occurs because:
(a) non-polar tails of soap molecules dissolve in grease
(b) oil and grease dissolved into hydrophilic centres of soap micelles and washed away
(c) hydrophilic heads dissolve in grease
(d) grease dissolves in soap solution
51. Arsenous sulphide sol is prepared by passing H_2S through arsenous oxide solution. The charge developed on the particles is due to adsorption of:
(a) H^+ (b) S^{2-} (c) OH^- (d) O^{2-}
52. Bredig's arc method cannot be used for the preparation of colloidal sol of :
(a) copper (b) gold (c) silver (d) sodium
53. As_2S_3 sol is:
(a) positive colloid (b) negative colloid
(c) neutral colloid (d) none of the above
54. Which of the following electrolyte will be most effective in coagulation of negative sol ?
(A) KNO_3 (b) $K_4[Fe(CN)_6]$
(C) Na_3PO_4 (d) $MgCl_2$
55. The minimum amount of an electrolyte required to cause coagulation of a sol is called :
(a) Coagulation value (b) Gold number
(c) Protective value (d) None of these
56. The electrical charge on a colloidal particle is indicated by:
(a) Brownian movement (b) electrophoresis
(c) ultramicroscope (d) molecular sieves
57. Colloidal particles in a sol can be coagulated by :
(a) heating (b) adding an electrolyte
(c) adding oppositely charged sol (d) any of the above methods
58. Peptization involves:
(a) precipitation of colloidal particles
(b) disintegration of colloidal aggregates
(c) purification of colloids
(d) impact of molecules of the dispersion medium on the colloidal particles

59. Hardy-Schulze law states that:
- solution must have higher gold number
 - disperse phase and dispersion medium must be of the same sign
 - micelles coagulate in presence of surfactants
 - the ions carrying more opposite charge to that of sol particle are effective in coagulation
60. Given below are a few electrolytes, indicate which one among them will bring about the coagulation of a gold sol quickest and in the least of concentration?
- NaCl
 - MgSO₄
 - Al₂(SO₄)₃
 - K₄[Fe(CN)₆]
61. The ability of an ion to bring about coagulation of a given colloid depends upon:
- its size
 - the magnitude of its charge only
 - the sign of its charge alone
 - both magnitude and sign of its charge
62. An emulsifying agent is a substance which
- stabilizes the emulsion
 - de-stabilizes the emulsion
 - coagulates the emulsion
 - break the interfacial film between suspended particle and medium
63. Colloidal solution of gold is prepared by:
- colloidal mill
 - double decomposition method
 - Bredig's method
 - peptization
64. The formation of colloid from suspension is called:
- peptisation
 - condensation
 - sedimentation
 - fragmentation
65. The method usually employed for the destruction of a colloidal solution is:
- dialysis
 - addition of electrolytes
 - diffusion through animal membrane
 - condensation
66. Which of the following has minimum flocculation value for positively charged sol?
- Cl⁻
 - SO₄²⁻
 - PO₄³⁻
 - [Fe(CN)₆]⁴⁻
67. Which of the following will have the highest coagulating power for Fe(OH)₃ colloid?
- PO₄³⁻
 - SO₄²⁻
 - Ca²⁺
 - Al³⁺
68. Artificial rain is caused by spraying:
- opposite charged collidal dust over a cloud
 - same charged collidal dust over a cloud
 - both
 - none of these
69. Colloids can be purified by :
- condensation
 - peptization
 - coagulation
 - dialysis


70. At what pH range the rate of enzyme catalyzed reaction is maximum ?
(a) 5 to 7 (b) 6 to 8
(c) 7 to 9 (d) 8 to 10
71. Protective sols are:
(a) lyophilic (b) lyophobic
(c) both (a) and (b) (d) none of these
72. Electro-osmosis is observed when:
(a) dispersion medium particles begins to move in an electric field
(b) dispersed phase begins to move in an electric field
(c) both (a) and (b)
(d) none of the above
73. On adding AgNO_3 solution into KI solution, a negatively charged colloidal sol is obtained when they are in:
(a) 50 mL of 0.1 M AgNO_3 + 50 mL of 0.1 M KI
(b) 50 mL of 0.1 M AgNO_3 + 50 mL of 0.2 M KI
(c) 50 mL of 0.2 M AgNO_3 + 50 mL of 0.1 M KI
(d) None of these
74. A sol is prepared by addition of excess of AgNO_3 solution in KI solution. The charge likely to develop on colloidal particles is:
(a) positive (b) negative
(c) no charge (d) both charges
75. The gold numbers of protective colloids A, B, C and D are 0.04, 0.004, 10 and 40 respectively. The protective powers of A, B, C and D are in the order:
(A) $A > B > C > D$ (b) $B > A > C > D$
(C) $D > C > A > B$ (d) $D > C > B > A$
76. In Brownian motion, the paths of the particles are:
(a) linear (b) curved
(c) zig-zag (d) uncertain
77. Which one of the following is not represented by sols?
(a) Adsorption (b) Tyndall effect
(c) Flocculation (d) Paramagnetism
78. The Tyndall effect associated with colloidal particles is due to:
(a) presence of electrical charges (b) scattering of light
(c) absorption of light (d) reflection of light
79. Blue colour of the sky is due to:
(a) absorption of light by dust particles (b) reflection of light by dust particles
(c) scattering of light by dust particles (d) presence of clouds
80. The apparatus used to coagulate carbon particles from smoke is called:
(a) cottrel smoker (b) cottrell precipitator
(c) cottrell absorber (d) none of these




Level 2

1. Select correct adsorption isobars for chemisorption and physisorption respectively :

(where $\frac{x}{m}$ extent of adsorption, T temperature)




2. Which among the following statements is false?
- Increase of pressure increases the amount of adsorption
 - Increase of temperature may decrease the amount of adsorption
 - Adsorption may be monolayered or multilayered
 - Particle size of the adsorbent will not affect the amount of adsorption
3. Select incorrect statement;
- Lyophilic sols are reversible
 - Lyophilic sols are self stabilized
 - Lyophobic sols are obtained from inorganic materials
 - Lyophobic sols particles are hydrated
4. Which one of the following statements is not correct in respect of lyophilic sols?
- There is a considerable interaction between the dispersed phase and dispersion medium
 - These are quite stable and are not easily coagulated
 - They always carry significant characteristic charge
 - The particles are hydrated
5. Alums purify muddy water by:
- dialysis
 - absorption
 - coagulation
 - ultrafiltration
6. Lyophilic sols are more stable than lyophobic sols because:
- the colloidal particles have positive charge
 - the colloidal particles have negative charge
 - the colloidal particles are solvated
 - there is strong electrostatic repulsion between the colloidal particles
7. Arrange the following electrolytes in the increasing order of coagulation power for the coagulation of As_2S_3 sol:



- (a) $I < II < III < IV$ (b) $I = III < II < IV$
(c) $II < IV < I < II$ (d) $II < III < IV < I$
8. Equal volume each of two sols of AgI, one obtained by adding $AgNO_3$ to slight excess of KI and another obtained by adding KI to slight excess of $AgNO_3$, are mixed together. Then:
(a) the two sols will stabilize each other
(b) the sol particles will acquire more electric charge
(c) the sols will coagulate each other mutually
(d) a true solution will be obtained
9. Which of the following statements is not true about the oil-in-water type emulsion?
(a) Addition of small amount of water, no separate layer of water is formed
(b) Addition of a small amount of oil soluble dye renders the entire emulsion coloured
(c) Addition of oil results in the formation of two layers
(d) Addition of a small amount of an electrolyte increases the conductivity of the emulsion
10. Under the influence of an electric field, the particles in a sol migrate towards cathode. The coagulation of the same sol is studied using NaCl, Na_2SO_4 and Na_3PO_4 solutions. Their coagulating values will be in maximum for:
(a) NaCl (b) Na_2SO_4 (c) Na_3PO_4 (d) same for all
11. Select incorrect statement:
(a) Micelles are associated colloids
(b) The electrical charge on a colloid particle is indicated by electrophoresis
(c) Formation of micelles takes place above Kraft temperature
(d) Formation of micelles takes place below CMC
12. Bredig's arc method involves:
(a) dispersion of metal (b) condensation of metal
(c) dispersion as well as condensation (d) neither dispersion nor condensation
13. A freshly prepared $Fe(OH)_3$ precipitate is peptized by adding $FeCl_3$ solution. The charge on the colloidal particle is due to preferential adsorption of:
(a) Cl^- ions (b) Fe^{3+} ions (c) OH^- ions (d) none of these
14. Select incorrect statement :
(a) Gold sol is multimolecular colloid
(b) Large number of particles of a substance aggregate together and formed multimolecular colloids
(c) Metal sulphides are lyophobic colloids
(d) Sulphur sol is multimolecular colloid and hydrophilic in nature
15. Associated colloids :
(a) raise both the surface tension and viscosity of water
(b) lower both the surface tension and viscosity of water
(c) lower the surface tension and raise the viscosity of water
(d) have greater concentration at the surface layer than the bulk of the solution
16. Below critical micelle concentration (CMC):
(a) the surfactant molecules of ions undergo association to form clusters
(b) the viscosity of solution increases abruptly
(c) substances like grease, fat, etc. dissolve colloiddally
(d) salt behave as normal, strong electrolyte

17. During electro-osmosis of $\text{Fe}(\text{OH})_3$ sol:
- sol particles move towards anode
 - sol particles move towards cathode
 - the dispersion medium moves towards anode
 - the dispersion medium moves towards cathode
18. Select incorrect statement:
- Soap and detergent lower the interfacial surface tension between oil and water
 - Basic principle of peptization is reverse of coagulation
 - Soap and detergent used as emulsifiers
 - Lyophilic sols need stabilizing agent
19. Smoke precipitator works on the principle of :
- centrifugation
 - neutralization of charge on colloids
 - absorption
 - addition of electrolytes
20. What is the correct sequence of the decreasing coagulation value of the following electrolyte for the coagulation of ferric hydroxide sol?
- (I) Na_3PO_4 (II) KCl (III) K_2SO_4 (IV) $\text{K}_4[\text{Fe}(\text{CN})_6]$
- $\text{IV} > \text{I} > \text{III} > \text{II}$
 - $\text{II} > \text{III} > \text{I} > \text{IV}$
 - $\text{I} > \text{II} > \text{III} > \text{IV}$
 - $\text{IV} > \text{III} > \text{II} > \text{I}$
21. Which of the following is true in respect of chemical adsorption?
- H 0, S 0, G 0
 - H 0, S 0, G 0
 - H 0, S 0, G 0
 - H 0, S 0, G 0
22. For the graph below, select correct order of temperature?



- $T_1 \quad T_2 \quad T_3$
 - $T_2 \quad T_3 \quad T_1$
 - $T_3 \quad T_2 \quad T_1$
 - $T_1 \quad T_2 \quad T_3$
23. Although, nitrogen does not adsorb on a surface at room temperature, it adsorbs on the surface at 83 K. Which one of the following statements is correct ?
- At 83 K, there is formation of monolayer
 - At 83 K, nitrogen is adsorbed as atoms
 - At 83 K, nitrogen molecules are held by chemical bonds
 - At 83 K, there is formation of multimolecular layers
24. For the coagulation of 50 mL of ferric hydroxide sol 10 mL of 0.5 M KCl is required. What is the coagulation value of KCl?
- 5
 - 10
 - 100
 - None of these
25. 100 mL of 0.6 M acetic acid is shaken with 2 g activated carbon. The final concentration of the solution after adsorption is 0.5 M. What is the amount of acetic acid adsorbed per gram of carbon?
- 0.6 g
 - 0.3 g
 - 1.2 g
 - None of these

26. A detergent ($\text{C}_{12}\text{H}_{25}\text{SO}_4^-\text{Na}^+$) solution becomes a colloidal sol at a concentration of 10^{-3} M . On an average 10^{13} colloidal particles are present in 1 mm^3 . What is the average number of ions which are contained by one colloidal particle (micelle)? [Given: $N_A = 6 \times 10^{23}$]
(a) 6×10^7 (b) 10 (c) 60 (d) None of these
27. One gram of activated carbon has a surface area of 1000 m^2 . Considering complete coverage as well as monomolecular adsorption, how much ammonia at 1 atm and 273 K would be adsorbed on the surface of $\frac{44}{7} \text{ g}$ carbon if radius of ammonia molecules is 10^{-8} cm .
[Given: $N_A = 6 \times 10^{23}$]
(a) 7.46 L (b) 0.33 L (c) 44.8 L (d) 23.5 L
28. At 1 atm and 273 K the volume of nitrogen gas required to cover a sample of silica gel, assuming Langmuir monolayer adsorption, is found to be $1.30 \text{ cm}^3 \text{ g}^{-1}$ of the gel. The area occupied by a nitrogen molecule is 0.16 nm^2 . What is the surface area per gram of silica gel?
[Given: $N_A = 6 \times 10^{23}$]
(a) $5.568 \text{ m}^2 \text{ g}^{-1}$ (b) $3.48 \text{ m}^2 \text{ g}^{-1}$ (c) $1.6 \text{ m}^2 \text{ g}^{-1}$ (d) None of these
29. 10% sites of catalyst bed have adsorbed by H_2 . On heating H_2 gas is evolved from sites and collected at 0.03 atm and 300 K in a small vessel of 2.46 cm^3 .
No. of sites available is 5.4×10^{16} per cm^2 and surface area is 1000 cm^2 . Find out the no. of surface sites occupied per molecule of H_2 . (Given $N_A = 6 \times 10^{23}$)
(a) 1 (b) 2 (c) 3 (d) None of these
30. A sample of 16 g charcoal was brought into contact with CH_4 gas contained in a vessel of 1 litre at 27°C . The pressure of gas was found to fall from 760 to 608 torr. The density of charcoal sample is 1.6 g/cm^3 . What is the volume of the CH_4 gas adsorbed per gram of the adsorbent at 608 torr and 27°C ?
(a) 125 mL/g (b) 16.25 mL/g (c) 26 mL/g (d) None of these



Level 3



Passage 1


A graph between x/m and the pressure P of the gas at a constant temperature is called adsorption isotherm. Where x is the no. of moles of the adsorbate and m is the mass of the adsorbent. Adsorption isotherms of different shapes have been experimentally observed. According to Freundlich adsorption isotherm.

$$x/m = kP^{1/n}$$

where k and n are constant parameters depending upon the nature of the solid and gas.

1. In the given isotherm select the incorrect statement :


- $\frac{x}{m} \propto P^{1/n}$ along OA
- $\frac{x}{m} \propto P^0$ when point B is reached
- $\frac{x}{m}$ does not increase as rapidly with pressure along BC due to less surface area available for adsorption
- nature of isotherm is different for two gases for same adsorbent



2. Adsorption isotherm of $\log \frac{x}{m}$ and $\log P$ was found of the type :

This is true when :


- $P \rightarrow 0$
- $P \rightarrow \infty$
- $\frac{1}{n} \rightarrow 1$
- $\frac{1}{n} \rightarrow 0$



3. Graph between $\log \frac{x}{m}$ and $\log P$ is a straight line at angle 45° with intercept OA as shown.

Hence, $\frac{x}{m}$ at a pressure of 2 atm is :

- 2
- 4
- 8
- 16



Passage 2

The protective power of the lyophilic colloids is expressed in terms of gold number, a term introduced by Zsigmondy. Gold number is the number of milli-gram of the protective colloid which prevent the coagulation of 10 mL of red gold sol, when 1 mL of a 10 per cent solution of sodium chloride is added to it. Thus, smaller the gold number of lyophilic colloid, the greater is the protective power.

1. On addition of one mL of solution of 10% NaCl to 10 mL of red gold sol in presence of 0.025 g of starch, the coagulation is just prevented. The gold number of starch is:
(a) 0.025 (b) 0.25 (c) 2.5 (d) 25
2. Which of the following statement(s) is/are correct?
(a) Higher the gold number, more protective power of colloid
(b) Lower the gold number, more the protective power
(c) Higher the coagulation value, more the coagulation power
(d) Lower the coagulation value, higher the coagulation power
3. Gold number gives an indication of:
(a) protective nature of colloids
(b) purity of gold in suspension
(c) the charge on a colloidal solution of gold
(d) gram of gold per litre of solution



Passage 3

Coagulation is the process by which the dispersed phase of a colloid is made to aggregate and thereby separate from the continuous phase. The minimum concentration of an electrolyte in milli-moles per litre of the electrolyte solution which is required to cause the coagulation of colloidal sol is called coagulation value.

Therefore higher is the coagulating power of effective ion, smaller will be the coagulation value of the electrolyte.

$$\text{Coagulation value} = \frac{1}{\text{coagulating power}}$$

The coagulation values of different electrolytes are different. This behaviour can be easily understood by Hardy-Schulze rule which states.

“The greater is the valency of the effective ion greater is its coagulating power.”

1. Which one has the highest coagulating power?
(a) K^+ (b) Ca^{2+} (c) Al^{3+} (d) Sn^{4+}
2. As_2S_3 sol is negatively charged. Capacity to precipitate it is highest in:
(a) K_2SO_4 (b) Na_3PO_4 (c) $AlCl_3$ (d) $CaCl_2$
3. The ability of an ion to bring coagulation of a given colloid depends upon:
(a) the sign of its charge (b) magnitude of its charge
(c) both magnitude and sign (d) none of these
4. The coagulation of colloidal particles of the sol can be caused by:
(a) heating
(b) adding electrolyte
(c) adding oppositely charged sol
(d) all of these

**Passage 4**

Emulsions are normally prepared by shaking the two components together vigorously although some kind of emulsifying agent usually has to be added to stabilize the product. This emulsifying agent may be a soap or other surfactant (surface active) species or a lyophilic sol that forms a protective film around the dispersed phase.

Emulsions are broadly classified into two types :

- (i) Oil in water emulsions (O/W) : Oil acts as dispersed phase and water acts as dispersion medium.
- (ii) Water in oil emulsions (W/O) : Water acts as dispersed phase and oil acts as dispersion medium. Dye test, dilution test may be employed for identification of emulsions.

1. Read two statements:

- (1) Milk is an example of oil in water (O/W) type emulsion
- (2) Cold cream is an example of water in oil (W/O) type emulsion
- (a) Only statement 1 is correct
- (b) Only statement 2 is correct
- (c) Both are correct
- (d) None of these

2. Select correct statement:

- (a) Water in oil emulsions are less viscous than the aqueous emulsions
- (b) Electrical conductance of aqueous emulsions is less than that of oil emulsions
- (c) Deemulsification can be done by soap or detergent
- (d) An emulsion can be diluted with H_2O then it is oil in water (O/W) type

One or More Answers is/are Correct

1. Select the correct statement(s) :

- (a) Physical adsorption is specific in nature
- (b) Chemical adsorption is highly specific in nature
- (c) Physical adsorption is due to free valence of atoms
- (d) Chemical adsorption is due to stronger interaction or bond formation

2. Select the correct statement(s):

- (a) Adsorption is a non-spontaneous process
- (b) Surface energy decreases during the process of adsorption
- (c) Adsorption takes place with decrease of entropy
- (d) In general adsorption is exothermic process

3. Select the correct statement(s):

- (a) Physisorption is favoured by low temperature
- (b) Chemisorption is favoured by very high temperature because the process is endothermic
- (c) Chemisorption increases with increase in temperature owing to high activation energy
- (d) Oxygen adsorbed by charcoal can be desorbed by lowering pressure and temperature

4. If adsorption of a gas on a solid is limited to monolayer formation, then which of the following statements are true?
- (a) At low pressures, $\frac{x}{m}$ varies proportionately with p
 - (b) At moderate pressures, $\frac{x}{m}$ varies less than proportionately with p
 - (c) At high pressures, $\frac{x}{m}$ becomes independent of p
 - (d) At high pressures, $\frac{x}{m}$ varies more than proportionately with p
5. Which of the following are colloids?
- (a) Sulphur sol (b) Starch (c) Gold sol (d) Soap solution
6. Which of the following is not ?
- (a) Gelatin sol (b) Silver sol
 - (c) Sulphur sol (d) As_2S_3 sol
7. Which of the following is/are correct for sols?
- (a) Its surface tension is lower than that of water
 - (b) Its viscosity is higher than that of water
 - (c) Its surface tension is higher than that of water
 - (d) Its viscosity is equal to that of water
8. Select the correct statement(s):
- (a) Benzene is dispersed phase in benzosol
 - (b) sols are irreversible and not so stable
 - (c) sol can be produced by double decomposition
 - (d) When a solution of sulphur in alcohol is added in excess of water a sol of alcohol is formed
9. When negatively charged colloid like As_2S_3 sol is added to positively charged $\text{Fe}(\text{OH})_3$ sol in stoichiometric amounts
- (a) both the are precipitated simultaneously
 - (b) this process is called mutual coagulation
 - (c) they becomes positively charged colloid
 - (d) they become negatively charged colloid
10. Colloidal gold can be prepared by:
- (a) Bredig's arc method (b) reduction of AuCl_3
 - (c) hydrolysis (d) peptization
11. The coagulation of sol particles may be brought about by:
- (a) boiling (b) persistent dialysis
 - (c) adding electrolyte (d) adding oppositely charged sol
12. Select the correct statement(s) :
- (a) A sol is prepared by addition of excess of AgNO_3 solution in KI solution. The charge likely to develop on colloidal particle is positive
 - (b) The effects of pressure on physical adsorption is high if temperature is low
 - (c) Ultracentrifugation process is used for preparation of colloids
 - (d) Gold number is the index for extent of gold plating done

13. Colloidal solution can be purified by:
 (a) dialysis (b) electrodialysis (c) electrophoresis (d) ultrafiltration
14. Coagulation of colloids can be achieved by:
 (a) centrifugation (b) adding electrolyte (c) change in pH (d) adding water
15. Which are the properties of colloidal solution ?
 (a) Adsorption (b) Tyndall effect
 (c) Flocculation (d) Depression of freezing point
16. In the aqueous solution of soaps above :
 (a) the cations associate to form the aggregates
 (b) the anions associate to form the clusters of colloidal dimension
 (c) the polar ends of the ions forming the clusters are directed towards water
 (d) the non-polar (hydrocarbon) ends are directed towards water
17. Amongst the following which is/are correct statement about the metal sulphide sols?
 (a) The sol particles are positively charged due to preferential adsorption of metal ions
 (b) The sol particles are negatively charged due to preferential adsorption of sulphide ions
 (c) The cations of added electrolytes are effective in causing the coagulation of the sol
 (d) The sol is due to both the electric charge and hydration of the particles
18. Emulsion can be destroyed by:
 (a) the addition of an emulsifier (b) electrophoresis with a high potential
 (c) freezing (d) all of these
19. Which of the following statement is/are correct for electrophoresis?
 (a) Colloids are uncharged particles and do not migrate towards the electrodes when electric field is applied
 (b) In electrophoresis, sol migrates either to the anode or to the cathode depending on the positively or negatively charged sol
 (c) Electrophoresis is useful for finding the charge on a sol
 (d) All of these
20. Select the false statement(s):
 (a) Brownian motion and Tyndall effect are shown by true solutions
 (b) Sorption process is combinations of adsorption and absorption process
 (c) Law Hardy Schulze is related with coagulation of a sol
 (d) Higher is the gold number greater will be the protective power of a colloid
21. Select correct statement(s)
 (a) colloids are used to protect lyophilic colloids
 (b) Dehydrating agent is used to coagulate sols
 (c) Rubber is obtained by coagulation of latex
 (d) Sometimes, the rainfall occurs when two oppositely charged clouds meet
22. In which of the followings, effect is/are not observed?
 (a) Sugar solution (b) Emulsions
 (c) Urea solution (d) Solution of proteins
23. Which of the following reactions are examples for heterogeneous catalysis?
 (a) $2\text{H}_2\text{O}_2(l) \xrightarrow{\text{MnO}_2(s)} 2\text{H}_2\text{O}(l) + \text{O}_2(g)$
 (b) $2\text{SO}_2(g) + \text{O}_2(g) \xrightarrow{\text{NO}(g)} 2\text{SO}_3(g)$
 (c) $\text{H}_2(g) + \text{C}_2\text{H}_4(g) \xrightarrow{\text{Ni}(s)} \text{C}_2\text{H}_6(g)$
 (d) $\text{CH}_3\text{COOC}_2\text{H}_5(aq) + \text{H}_2\text{O}(l) \xrightarrow{\text{HCl}(aq)} \text{CH}_3\text{COOH}(aq) + \text{C}_2\text{H}_5\text{OH}(aq)$

24. Select correct statement(s)

- (a) The role of a catalyst in a reversible reaction is to allow the equilibrium to be achieved quickly
- (b) Diffusion process is involved in mechanism of heterogeneous catalysis process
- (c) Hydrolysis of cane sugar is catalysed by H
- (d) Promoters enhance the activity of a catalyst

25. Select correct statement(s)

- (a) Blood is a colloidal solution
- (b) Alum is used in water purification
- (c) River water is a colloidal solution of clay
- (d) Colloidal medicines are more effective due to large surface area

Match the Column

Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.**Column-I**

- (A) Chemisorption
- (B) Physical adsorption
- (C) Desorption
- (D) Activation of adsorbent

Column-II

- (P) Exothermic
- (Q) Endothermic
- (R) Removal of adsorbed material
- (S) Highly specific in nature

2.**Column-I**

- (A) Chemisorption
- (B) Physisorption
- (C) Desorption of a solute on liquid surface
- (D) Adsorption of a solute on a liquid surface

Column-II

- (P) Not specific and decreases with temperature
- (Q) Specific and increases with temperature
- (R) Increases the surface tension of the liquid
- (S) Decreases the surface tension of the liquid

3.**Column-I**

- (A) Milk
- (B) Dust
- (C) Cheese
- (D) Froth

Column-II

- (P) Aerosol
- (Q) Emulsion
- (R) Gel
- (S) Foam

4.	Column-I (A) Liquid dispersed in gas (B) Gas dispersed in liquid (C) Liquid dispersed in solid (D) Liquid dispersed in liquid	Column-II (P) Foam (Q) Emulsion (R) Aerosol (S) Gel
5.	Column-I (A) As_2S_3 sol (B) Sulphur sol (C) Starch (D) Soap	Column-II (P) Lyophobic colloid (Q) Macromolecular colloid (R) Multimolecular colloid (S) Associated colloid
6.	Column-I (A) Coagulation (B) Electrophoresis (C) Tyndall effect (D) Brownian movement	Column-II (P) Due to presence of charge (Q) Due to scattering of light (R) Due to neutralization of charge (S) Due to impact of the molecules of the dispersion medium with colloidal particles
7.	Column-I (A) Peptization (B) Ultra centrifugation (C) Electrodialysis (D) Bredig's arc method	Column-II (P) Preparation of sols (Q) Purification of sols (R) Preparation of metal sols (S) Movement of ions across the membrane in presence of electric field

Assertion-Reason Type Questions



Each questions contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statement carefully and mark the correct answer according to the instructions given below:

- (A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- (B) If both the statements are TRUE and STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- (C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- (D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1** : For adsorption G , H , S all have $-ve$ values.
STATEMENT-2 : Adsorption is a exothermic process in which randomness decreases due to force of attraction between adsorbent and adsorbate.
2. **STATEMENT-1** : The extent of adsorption of CO_2 is much more higher than of H_2 .
STATEMENT-2 : $CO_2(g)$ has higher critical temperature and more van der Waals' force of attraction as compared to $H_2(g)$.
3. **STATEMENT-1** : In absorption, the molecules of a substance are uniformly distributed throughout the body of other substance.
STATEMENT-2 : In some cases, both absorption and adsorption takes place simultaneously.
4. **STATEMENT-1** : More heat evolved in physisorption than in chemisorption.
STATEMENT-2 : Molecules of adsorbate and adsorbent are held by van der Waals' forces in physisorption and by chemical bonds in chemisorption.
5. **STATEMENT-1** : Colloidal solution is electrically neutral.
STATEMENT-2 : Due to similar nature of the charge carried by the dispersed phase particles, they repel each other and do not combine to form bigger particles.
6. **STATEMENT-1** : Soap and detergent are macro-molecular colloids.
STATEMENT-2 : Soap and detergent are molecules of large size.
7. **STATEMENT-1** : Micelles are formed by surfactant molecules above the critical micelle concentration (CMC).
STATEMENT-2 : The conductivity of a solution having surfactant molecules decrease sharply at the CMC.
8. **STATEMENT-1** : The micelle formed by sodium stearate in water has COO^- groups at the surface.
STATEMENT-2 : Surface tension of water is reduced by the addition of stearate.
9. **STATEMENT-1** : Protein, starch are lyophilic colloids.
STATEMENT-2 : They have strong interaction with the dispersion medium.
10. **STATEMENT-1** : Colloidal AgI is prepared by adding KI in slight excess to $AgNO_3$ solution, the sol particles migrate toward cathode under electric field.
STATEMENT-2 : Colloidal particles adsorb ions and thus becomes electrically neutral.

- 11. STATEMENT-1 :** When SnO_2 is reacted with NaOH , then its sol particles are attracted towards cathode.
STATEMENT-2 : When SnO_2 is reacted with NaOH , then it gives SnO_3^{2-} which is adsorbed by SnO_2 , so it is negatively charged sol.
- 12. STATEMENT-1 :** For coagulation of positively charged sols, $[\text{Fe}(\text{CN})_6]^{4-}$ ion has higher coagulating power than that of PO_4^{3-} , SO_4^{2-} , Cl^- .
STATEMENT-2 : Because according to Hardy Schulze rule, higher is the valency of ions for the oppositely charged sol particles, better will be the precipitation.
- 13. STATEMENT-1 :** Dispersed phase particles of colloidal solution cannot pass through ultra-filter paper.
STATEMENT-2 : The size of colloidal particles are larger than the size of true solution particles.
- 14. STATEMENT-1 :** When a finely divided active carbon is stirred into a dilute solution of a dye, the intensity of colour in the solution is decreased.
STATEMENT-2 : The dye is adsorbed on the solid surface.
- 15. STATEMENT-1 :** Silica gel is used for drying air.
STATEMENT-2 : Silica gel adsorbs moisture from air.
- 16. STATEMENT-1 :** ZSM-5 is a type of zeolites used as a catalyst in petrochemical industries.
STATEMENT-2 : Zeolites are microporous aluminosilicates three dimensional network silicates in which some silicon atoms are replaced by aluminium atoms.
- 17. STATEMENT-1 :** Lyophilic colloids are called as reversible sols.
STATEMENT-2 : Lyophilic sols are extensively hydrated.
- 18. STATEMENT-1 :** A catalyst is more effective in finely divided form.
STATEMENT-2 : Finely divided form has more surface area.
- 19. STATEMENT-1 :** Sky appears blue in colour.
STATEMENT-2 : Colloidal particles of dust along with water suspended in air scatter blue light.
- 20. STATEMENT-1 :** The conversion of fresh precipitate to colloidal state is called peptization.
STATEMENT-2 : It is caused by addition of common ions.
- 21. STATEMENT-1 :** Colloidal solutions are stable but colloidal particles do not settle down.
STATEMENT-2 : Brownian movement counters the force of gravity act on colloidal particles.
- 22. STATEMENT-1 :** A colloid gets coagulated by addition of an electrolyte.
STATEMENT-2 : Coagulation depends on the valance and sign of the charge of the coagulant ion.
- 23. STATEMENT-1 :** Suspensions are visible to naked eye.
STATEMENT-1 : Size of suspension particles are smaller than collidal particles.
- 24. STATEMENT-1 :** At pH of isoelectric point, the sol particles of amino acids neither move towards anode nor towards cathode.
STATEMENT-2 : Because at the isoelectric point, the concentration of conjugate base and conjugate acid of the Zwitter ions becomes equal and so one's charge is counterbalanced by other.

25. STATEMENT-1 : Gold sol is multimolecular and hydrophobic in nature.

STATEMENT-2 : Gold sol is prepared by Bredig's arc method.

Subjective Problems



1. How many colloidal systems exist in nature ?
2. How many colloidal systems exist in nature in which liquid is dispersed phase ?
3. How many colloidal systems exist in nature in which gas is dispersed phase ?
4. What is the minimum diameter (in nm) of colloidal particles ?
5. For soaps critical micelle concentration (CMC) is 10^{-x} (min.) to 10^{-y} (max.) mol/L. What is the value of x ?

ANSWERS

Level 1

1. (a)	2. (d)	3. (d)	4. (d)	5. (a)	6. (d)	7. (c)	8. (c)	9. (d)	10. (b)
11. (b)	12. (d)	13. (d)	14. (c)	15. (a)	16. (b)	17. (a)	18. (b)	19. (a)	20. (c)
21. (a)	22. (a)	23. (a)	24. (a)	25. (c)	26. (c)	27. (a)	28. (b)	29. (b)	30. (c)
31. (c)	32. (b)	33. (a)	34. (b)	35. (d)	36. (d)	37. (c)	38. (a)	39. (d)	40. (b)
41. (b)	42. (b)	43. (b)	44. (b)	45. (a)	46. (c)	47. (b)	48. (b)	49. (d)	50. (a)
51. (b)	52. (d)	53. (b)	54. (d)	55. (a)	56. (b)	57. (d)	58. (b)	59. (d)	60. (c)
61. (d)	62. (a)	63. (c)	64. (a)	65. (b)	66. (d)	67. (a)	68. (a)	69. (d)	70. (a)
71. (a)	72. (a)	73. (b)	74. (a)	75. (b)	76. (c)	77. (d)	78. (b)	79. (c)	80. (b)

Level 2

1. (c)	2. (d)	3. (d)	4. (c)	5. (c)	6. (c)	7. (b)	8. (c)	9. (b)	10. (a)
11. (d)	12. (c)	13. (b)	14. (d)	15. (c)	16. (d)	17. (c)	18. (d)	19. (b)	20. (b)
21. (b)	22. (c)	23. (d)	24. (c)	25. (b)	26. (c)	27. (a)	28. (a)	29. (c)	30. (b)

Level 3

Passage-1	1. (b)	2. (c)	3. (b)		
Passage-2	1. (d)	2. (b)	3. (a)		
Passage-3	1. (d)	2. (c)	3. (c)	4. (d)	
Passage-4	1. (c)	2. (d)			

One or More Answers is/are Correct

1. (b,d) 2. (b,c,d) 3. (a,c, d) 4. (a,b,c) 5. (a,c) 6. (b,c,d) 7. (a,b) 8. (b,c)
9. (a,b) 10. (a,b) 11. (a,b,c,d) 12. (a,b) 13. (a,b,d) 14. (a,b,c) 15. (a,b,c,d) 16. (b,c)
17. (b,c) 18. (b,c) 19. (c) 20. (a,d) 21. (b,c,d) 22. (a,c) 23. (a,c) 24. (a,b,c,d)
25. (a,b,c,d)

Match the Column

- | | | | |
|------------|---------|---------|--------|
| 1. A P, S; | B P; | C Q, R; | D Q |
| 2. A Q; | B P; | C R; | D S |
| 3. A Q; | B P; | C R; | D S |
| 4. A R; | B P; | C S; | D Q |
| 5. A P, R; | B P, R; | C Q; | D S |
| 6. A R; | B P; | C Q; | D S |
| 7. A P; | B Q; | C Q, S; | D P, R |

Assertion-Reason Type Questions

1. (A) 2. (A) 3. (B) 4. (D) 5. (B) 6. (D) 7. (A) 8. (B) 9. (A) 10. (C)
11. (D) 12. (A) 13. (A) 14. (A) 15. (A) 16. (B) 17. (B) 18. (A) 19. (A) 20. (B)
21. (A) 22. (B) 23. (C) 24. (A) 25. (B)


Subjective Problems

1. 8	2. 3	3. 2	4. 1	5. 4															
------	------	------	------	------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Hints and Solutions


Level 1

1. (a) Solution Particle size
True solution Size < 1 nm
Colloidal solution 1 nm < size < 1000 nm
Suspension Size > 1000 nm
2. (d) 10^{-6} m to 10^{-9} m
3. (d) Absorption is not surface phenomena.
4. (d) Adsorbate adsorb at the surface of adsorbent.
5. (a) Chemical adsorption.
6. (d) Attractive force developed at the surface between adsorbent and adsorbate.
7. (c) Physical adsorption and chemical adsorption.
8. (c) Chemical adsorption is irreversible in nature.
9. (d) H O S O
10. (b) Physical adsorption is reversible in nature.
11. (b) Critical temp. Attractive force
Extent of adsorption
12. (d) Adsorption T_c
Adsorption T
Adsorption P
13. (d) Van der Waal force adsorption
14. (c) Van der Waal force adsorption
15. (a) Silica gel adsorb water.
16. (b) Adsorption : surface phenomena
Absorption : bulk phenomena
17. (a) Physical adsorption is multilayer adsorption.
18. (b) Physical adsorption is reversible adsorption.
19. (a) At low temperature, pressure have greater effect on adsorption.



20. (c) In sorption, adsorption and absorption takes place simultaneously.

21. (a)



$$22. (a) \frac{x}{m} = KP^{\frac{1}{n}}$$

$$\log \frac{x}{m} = \log K + \frac{1}{n} \log P$$

$$\text{Slope} = \frac{1}{n}$$

$$23. (a) H : 80 \text{ to } 240 \text{ kJ mol}^{-1}$$

$$24. (a) H : 20 \text{ to } 40 \text{ kJ mol}^{-1}$$

25. (c) Adsorption lowers the activation energy of reaction because surface energy released during adsorption.

$$26. (c) \text{Mass of } O_2 \text{ per gram of adsorbent} = \frac{3.6}{1.2} = 3$$

No. of moles of O_2 per gram of adsorbent

$$\frac{3}{32}$$

Volume of O_2 per gram of adsorbent

$$\frac{3}{32} \times \frac{0.0821 \times 273}{1}$$


$$= 2.10$$

27. (a) It is preferentially adsorb on the catalyst.
28. (b) Ni used as catalyst in hydrogenation of oil.
29. (b) $C_6H_{12}O_6 \xrightarrow{\text{Zymase}} CO_2 + C_2H_5OH$
30. (c) Maltose $\xrightarrow{\text{Maltase}}$ glucose
31. (c) Zeolites is shape selective catalyst.
32. (b) Autocatalyst
33. (a) Negative catalyst
34. (b) Surface area No of sites
35. (d) Enzyme are specific in nature.
36. (d) True solution.
37. (c) particle size
38. (a) Surface tension of lyophilic solution is lower than water.
39. (d) Metal sulphide are lyophobic solution.


40. (b) Gelatin act as protective colloids.
41. (b) Gum is lyophilic solution.
42. (b) Emulsion : D.P. liquid
D.M. liquid
43. (b) Soap formed associated colloids in water.
44. (b) D.P. have greater affinity towards D.M.
45. (a) fog : liquid in gas.
46. (c) Colloids divided on the basis of molecular size, affinity of D.P. towards D.M. and particle nature.
47. (b) Colloids have D.P. and D.M.
48. (b) Sulphur is lyophobic colloids.
49. (d) Milk is emulsion.
50. (a) Non-polar tails of soap molecules dissolve grease.
51. (b) As_2S_3 H_2S $\text{As}_2\text{S}_3 : \text{S}^{2-} : \text{H}$
Preferential adsorption of S^{2-} takes place. So negative charge developed.
52. (d) Bredig's arc method is used to prepare metal sol.
53. (b) Metal sulphides is negative colloids.
54. (d) Positive charge Coagulating power
 MgCl_2
55. (a) Millimoles of electrolyte required to coagulates one litre colloidal solution is called coagulation value.
56. (b) By electrophoresis, we can determine charge of colloidal solution.
57. (d) Coagulation takes place by heating, adding electrolyte and addition of opposite charge colloids.
58. (b) In peptisation, suspended state converts into colloidal state.
59. (d) Coagulating power charge.
60. (c) Gold sol. have negative charge. So $\text{Al}_2(\text{SO}_4)_3$ is most effective for coagulation.
61. (d) Coagulating power depend upon sign of charge and magnitude of charge.
62. (a) Emulsifying agent stabilise emulsion.
63. (c) Gold solution is prepared by Bredig's arc method.
64. (a) In peptisation, colloid formed from suspension.
65. (b) By addition of electrolyte.
66. (d) For positive charge colloids coagulating power coagulation value $[\text{Fe}(\text{CN})_6]^{4-}$.
67. (a) $\text{Fe}(\text{OH})_3$ is a positive sol. Hence, greater the charge on anion more is the coagulation power.
68. (a) Mutual coagulation
69. (d) Dialysis is a method of purification of colloids.
70. (a) 5 to 7
71. (a) Lyophilic solution are protective colloids.
72. (a) In electro osmosis D.M. move towards opposite charge electrode.
73. (b) AgNO_3 KI AgI(s) KNO_3
5 10^3 10 10^3
0 5 10^3 5 10^3
 AgI(s) I^- $\text{AgI} : \text{I}^-$
Negative charge colloids.
74. (a) AgNO_3 KI AgI KNO_3
excess
 AgI Ag^+ AgI Ag^+
75. (b) Smaller is the 'gold number' of protective colloid, greater is its protective power.
76. (c) Zig-zag path.
77. (d) paramagnetism.
78. (b) Tyndall effect takes place due to scattered of light.
79. (c) Light scattered by dust particles.
80. (b) Cottrell precipitator.

Level 2

1. (c)



Chemical adsorption



Physical adsorption

2. (d) Surface area rate of adsorption
3. (d) In lyophobic sol D.P. have less affinity towards D.M.
4. (c) In lyophilic sol no charge present.

5. (c) Alum, purified to muddy water by coagulation.
6. (c) D.P. particles have great affinity towards D.M. in lyophilic colloids.
7. (b) Charge coagulation power
8. (c) The sols obtained in the two cases will be oppositely charged so coagulate each other.
9. (b) Colour of dye appear in patches in o/w emulsion.
10. (a) The sol particles migrate towards cathode. So they are positively charged. Hence, anions would be effective in coagulation. Greater is the valence of effective ion, smaller will be its coagulating value.
11. (d) Formation of micelles takes place above CMC.
12. (c) Bredig's arc method involve dispersion and condensation.
13. (b) Preferential adsorption of Fe^{3+} takes place. So positive charge develop.
14. (d) Sulphur solution is multimolecular by hydrophobic in nature.
15. (c) Associate colloids decrease surface tension and increase in viscosity.
16. (d) Below CMC, salt behave as normal electrolyte.
17. (c) In electro-osmosis, the sol particles are prevented from migration whereas the dispersion medium migrates in the direction opposite to those of particles. Here, the medium is negatively charged.
18. (d) No stabilising agent required for the preparation of lyophilic sol.
19. (b) Neutralisation of charge.
20. (b) Ferric hydroxide solution have positive charge. So order of coagulation value.
 $\text{KCl} \quad \text{K}_2\text{SO}_4 \quad \text{Na}_3\text{PO}_4 \quad \text{K}_4[\text{Fe}(\text{CN})_6]$
 Charge coagulation value
21. (b) $H \quad 0, \quad S \quad 0, \quad G \quad 0$
22. (c) Temp Extent of adsorption
23. (d) At 83 K physical adsorption takes place effectively so multimolecular layers formed.
24. (c) Coagulation value

$$\frac{\text{Number of millimoles of electrolyte required}}{\text{Volume of colloidal solution (in litre)}}$$

$$\frac{10 \cdot 0.5}{50} \quad 1000 \quad 100$$

25. (b) Mass of acetic acid adsorbed by 2 g charcoal

$$100 \quad 10^{-3} \quad (0.6 - 0.5) \quad 60$$

$$0.6; \frac{x}{m} \quad \frac{0.6}{2} \quad 0.3$$

26. (c) No. of sodium lauryl sulphate molecules ($\text{CH}_3(\text{CH}_2)_{11}\text{SO}_4\text{Na}^+$) in 1 litre solution

$$10^{-3} \quad 6 \quad 10^{23}$$

$$6 \quad 10^{20}$$

$$\text{No. of sodium lauryl sulphate molecules per mm}^3 = 6 \cdot 10^{14}$$

$$\text{No. of colloidal particles per mm}^3 = 10^{13}$$

$$\text{No. of molecules per colloidal particle}$$

$$\frac{6 \cdot 10^{14}}{10^{13}}$$

$$= 60$$

27. (a) Total surface area of carbon $\frac{44}{7} \cdot 10^7 \text{ cm}^2$

$$\text{Surface area of } \text{NH}_3 = r^2$$

$$\frac{22}{7} \cdot 10^{16} \text{ cm}^2$$

$$\text{No. of } \text{NH}_3 \text{ molecules adsorbed} \quad \frac{\frac{44}{7} \cdot 10^7}{\frac{22}{7} \cdot 10^{16}}$$

$$2 \cdot 10^{23}$$

$$\text{Vol. of } \text{NH}_3 \text{ adsorbed at 1 atm 273 K}$$

$$\frac{2 \cdot 10^{23}}{6 \cdot 10^{23}} \cdot 22.4 = 7.46 \text{ L}$$

28. (a) No. of molecules per gram of N_2 in monolayer

$$\frac{6 \cdot 10^{23}}{22,400} \quad 1.30$$

$$= 3.48 \cdot 10^{19}$$

$$\text{Cross-sectional area of a molecule}$$

$$1.6 \cdot 10^{-19} \text{ m}^2$$

$$\text{Area covered by molecules per gram}$$

$$3.48 \cdot 10^{19} \cdot 1.6 \cdot 10^{-19} \quad 5.568 \text{ m}^2$$

$$\text{Surface area} \quad 5.568 \text{ m}^2 \text{ g}^{-1}$$

29. (c) Adsorbed moles of $\text{H}_2 = \frac{0.03 \cdot 2.46 \cdot 10^{-3}}{0.0821 \cdot 300}$

$$\text{No. of absorbed molecules of}$$

$$\text{H}_2 \quad 3 \cdot 10^6 \quad 6 \cdot 10^{23}$$

$$18 \cdot 10^{17}$$

Total no. of surface sites available

$$5.4 \times 10^{16} \times 1000$$

$$5.4 \times 10^{19} \text{ cm}^2$$

No. of surface sites that is occupied by adsorption of

$$\text{H}_2 \times \frac{10}{100} \times 5.4 \times 10^{19}$$

$$5.4 \times 10^{18}$$

No. of surface sites occupied by one molecule

$$\text{H}_2 \times \frac{5.4 \times 10^{18}}{18 \times 10^{17}} = 3$$

30. (b) Final volume of gas, at 608 torr pressure

$$V_2 \times \frac{P_1 V_1}{P_2} = \frac{760 \times 1}{608}$$

$$1.25 \text{ L or } V_2 = 1250 \text{ mL}$$

Volume occupied by gas Volume of vessel
Volume occupied by charcoal

$$1000 \times \frac{16}{1.6} = 990 \text{ mL}$$

Difference of volume is due to adsorption of gas by charcoal.

Volume of gas adsorbed by charcoal

$$1250 - 990$$

$$260 \text{ mL}$$

Volume of the gas adsorbed per gram of charcoal

$$\frac{260}{16} = 16.25 \text{ mL/g at 608 torr and } 27^\circ \text{C.}$$

Level 3

Passage-1

- (b) At point B $\frac{x}{m} = P^{1/n}$
- (c) $\log \frac{x}{m} = \log K + \frac{1}{n} \log P$
- (b) $\frac{x}{m} = K P^{1/n}$
 $\frac{x}{m} = 2 \times 2^1 = 4$

Passage-2

- (d) Gold no. $0.025 \times 10^3 = 25$
- (b) Gold no. protecting power

- (a) Protective nature of colloids.

Passage-3


- (d) Charge coagulating power
- (c) Al^{3+} charge coagulating power
- (c) Coagulation depend upon charge and magnitude.
- (d) Coagulation takes place by heating, adding electrolyte and mutual coagulation.

Passage-4

- (c) Milk : o/w emulsion
Cold cream : w/o emulsion
- (d) o/w emulsion diluted by addition of water.

One or More Answers is/are Correct

- (b, d)
Physical adsorption are not specific in nature due to van der Waal's forces and chemical adsorption are specific in nature and due to chemical bond.
- (b,c,d)
Adsorption is spontaneous process.
Enthalpy of adsorption is -ve
Change in entropy = -ve
- (a, c, d)
Low temperature is favourable for physical adsorption.
Chemical adsorption required activation energy.
- (a, b, c)



- (a, c)
Sulphur and gold are multimolecular colloids.
- (b, c, d)
Gelatin is lyophilic colloids.
- (a, b)
Surface tension is less than water and viscosity is greater than water in case of lyophilic sol.

8. (b, c)
Lyophobic sol are ir-reversible and less stable and they produced by double decomposition.
9. (a, b)
Mutual coagulation.
10. (a, b)
Gold solution prepared by Bredig's arc method and by reduction of AuCl_3 .
11. (a, b, c, d)
Coagulation takes place by boiling,
persistent dialysis
add of electrolyte
mutual coagulation.
12. (a, b)
 AgNO_3 KI AgI KNO_3
 AgI AgNO_3 $\text{AgI} \cdot \text{AgNO}_3$
At low temperature, extent of physical adsorption is greater due to increase in pressure.
13. (a, b, d)
Colloidal solution can be purified by dialysis, electrolysis and ultrafiltration.
14. (a, b, c)
Coagulation occurs due to centrifugation, add of electrolyte and change in pH.
15. (a, b, c, d)
Sol shows Tyndall effect adsorption coagulation sol have colligative properties
16. (b, c)
Above CMC, associate colloids formed.
Non polar part directed towards water.
17. (b, c)
Charge develop due to preferential adsorption of ion addition of electrolyte cause coagulation.
18. (b, c)
Coagulation occurs due to electrophoresis and freezing.
19. (c)
In electrophoresis colloidal particle migrate towards opposite charge electrode.
20. (a, d)
Tyndall effect does not shown by true solution due to small size of particles.
Sorption involve adsorption and absorption coagulating power charge, smaller the gold number greater would be the protecting power.
21. (b, c, d)
Lyophilic colloids used as protecting colloids electrolyte used for coagulation. Rubber is obtained by coagulation of latex. By mutual coagulation rain fall occurs.
22. (a, c)
Due to size of particle
23. (a, c)
 MnO_2 , Ni
24. (a, b, c, d)
In presence of catalyst equilibrium obtained quickly.
Mechanism of hetrogenous catalyst is based on adsorption.
H act as auto catalyst
Catalyst promotor increase the efficiency of catalyst.
25. (a, b, c, d)
Blood is colloidal solution, water purification by alum is based on coagulation. River water is colloidal solution of clay surface area preferential adsorption .

Subjective Problems

- Eight gas colloidal system does not exists.
- Aerosol, Emulsion, Gels.
- Foam, Solid foam.
- Size 1nm
- CMC of soap
 10^{-4} M to 10^{-3} M
 $\times 4$