

Limin Ge Research Catalog

26 result families in 719 manuscripts

October 9, 2026

An overview of the [manuscript collection](#). Results are grouped by subject; entry numbers follow the catalog order and do not indicate a ranking. PDF links follow each summary.

Contents

Real projective geometry	2
Coding theory	2
Discrete geometry	4

Real projective geometry

001. Packings of lines in $\mathbb{R}^{10}$. First values for 7 blank cells ($N = 101, 103, 106, 109, 112, 119$ – 120) in the row $m = 10, n = 1$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 101 \cdot N = 103 \cdot N = 106 \cdot N = 109 \cdot N = 112 \cdot N = 119 \cdot N = 120$

002. Packings of lines in $\mathbb{R}^{11}$. First values for 17 blank cells ($N = 84$ – 100) in the row $m = 11, n = 1$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 84 \cdot N = 85 \cdot N = 86 \cdot N = 87 \cdot N = 88 \cdot N = 89 \cdot N = 90 \cdot N = 91 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100$

003. Packings of lines in $\mathbb{R}^{12}$. First values for 14 blank cells ($N = 95$ – $100, 104$ – $105, 109, 111$ – $112, 116, 119$ – 120) in the row $m = 12, n = 1$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100 \cdot N = 104 \cdot N = 105 \cdot N = 109 \cdot N = 111 \cdot N = 112 \cdot N = 116 \cdot N = 119 \cdot N = 120$

004. Packings of lines in $\mathbb{R}^{13}$. First values for 44 blank cells ($N = 55$ – $89, 92$ – 100) in the row $m = 13, n = 1$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 55 \cdot N = 56 \cdot N = 57 \cdot N = 58 \cdot N = 59 \cdot N = 60 \cdot N = 61 \cdot N = 62 \cdot N = 63 \cdot N = 64 \cdot N = 65 \cdot N = 66 \cdot N = 67 \cdot N = 68 \cdot N = 69 \cdot N = 70 \cdot N = 71 \cdot N = 72 \cdot N = 73 \cdot N = 74 \cdot N = 75 \cdot N = 76 \cdot N = 77 \cdot N = 78 \cdot N = 79 \cdot N = 80 \cdot N = 81 \cdot N = 82 \cdot N = 83 \cdot N = 84 \cdot N = 85 \cdot N = 86 \cdot N = 87 \cdot N = 88 \cdot N = 89 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100$

005. Packings of lines in $\mathbb{R}^{14}$. First values for 42 blank cells ($N = 56$ – $60, 66$ – $100, 105, 120$) in the row $m = 14, n = 1$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 56 \cdot N = 57 \cdot N = 58 \cdot N = 59 \cdot N = 60 \cdot N = 66 \cdot N = 67 \cdot N = 68 \cdot N = 69 \cdot N = 70 \cdot N = 71 \cdot N = 72 \cdot N = 73 \cdot N = 74 \cdot N = 75 \cdot N = 76 \cdot N = 77 \cdot N = 78 \cdot N = 79 \cdot N = 80 \cdot N = 81 \cdot N = 82 \cdot N = 83 \cdot N = 84 \cdot N = 85 \cdot N = 86 \cdot N = 87 \cdot N = 88 \cdot N = 89 \cdot N = 90 \cdot N = 91 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100 \cdot N = 105 \cdot N = 120$

006. Packings of lines in $\mathbb{R}^{15}$. First values for 50 blank cells ($N = 51$ – 100) in the row $m = 15, n = 1$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 51 \cdot N = 52 \cdot N = 53 \cdot N = 54 \cdot N = 55 \cdot N = 56 \cdot N = 57 \cdot N = 58 \cdot N = 59 \cdot N = 60 \cdot N = 61 \cdot N = 62 \cdot N = 63 \cdot N = 64 \cdot N = 65 \cdot N = 66 \cdot N = 67 \cdot N = 68 \cdot N = 69 \cdot N = 70 \cdot N = 71 \cdot N = 72 \cdot N = 73 \cdot N = 74 \cdot N = 75 \cdot N = 76 \cdot N = 77 \cdot N = 78 \cdot N = 79 \cdot N = 80 \cdot N = 81 \cdot N = 82 \cdot N = 83 \cdot N = 84 \cdot N = 85 \cdot N = 86 \cdot N = 87 \cdot N = 88 \cdot N = 89 \cdot N = 90 \cdot N = 91 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100$

007. Packings of lines in $\mathbb{R}^{16}$. First values for 8 blank cells ($N = 51$ – 58) in the row $m = 16, n = 1$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 51 \cdot N = 52 \cdot N = 53 \cdot N = 54 \cdot N = 55 \cdot N = 56 \cdot N = 57 \cdot N = 58$

Coding theory

008. Packings of planes in $\mathbb{R}^8$. First values for 10 blank cells ($N = 102, 104, 106, 108$ – $109, 111, 113, 117, 119$ – 120) in the row $m = 8, n = 2$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 102 \cdot N = 104 \cdot N = 106 \cdot N = 108 \cdot N = 109 \cdot N = 111 \cdot N = 113 \cdot N = 117 \cdot N = 119 \cdot N = 120$

009. Packings of planes in $\mathbb{R}^{11}$. First values for 17 blank cells ($N = 84$ – 100) in the row $m = 11, n = 2$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 84 \cdot N = 85 \cdot N = 86 \cdot N = 87 \cdot N = 88 \cdot N = 89 \cdot N = 90 \cdot N = 91 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100$

010. Packings of planes in $\mathbb{R}^{12}$. First values for 6 blank cells ($N = 95$ – 100) in the row $m = 12, n = 2$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100$

020. Packings of three-spaces in $\mathbb{R}^{16}$. First values for 49 blank cells ($N = 51$ – 62 , 65 – 78 , 81 – 100 , 104 , 107 , 120) in the row $m = 16$, $n = 3$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 51 \cdot N = 52 \cdot N = 53 \cdot N = 54 \cdot N = 55 \cdot N = 56 \cdot N = 57 \cdot N = 58 \cdot N = 59 \cdot N = 60 \cdot N = 61 \cdot N = 62 \cdot N = 65 \cdot N = 66 \cdot$
 $N = 67 \cdot N = 68 \cdot N = 69 \cdot N = 70 \cdot N = 71 \cdot N = 72 \cdot N = 73 \cdot N = 74 \cdot N = 75 \cdot N = 76 \cdot N = 77 \cdot N = 78 \cdot N = 81 \cdot N = 82 \cdot$
 $N = 83 \cdot N = 84 \cdot N = 85 \cdot N = 86 \cdot N = 87 \cdot N = 88 \cdot N = 89 \cdot N = 90 \cdot N = 91 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot N = 95 \cdot N = 96 \cdot$
 $N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100 \cdot N = 104 \cdot N = 107 \cdot N = 120$

021. Packings of four-spaces in $\mathbb{R}^9$. First values for 15 blank cells ($N = 5$ – 19) in the row $m = 9$, $n = 4$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 5 \cdot N = 6 \cdot N = 7 \cdot N = 8 \cdot N = 9 \cdot N = 10 \cdot N = 11 \cdot N = 12 \cdot N = 13 \cdot N = 14 \cdot N = 15 \cdot N = 16 \cdot N = 17 \cdot N = 18 \cdot N = 19$

022. Packings of four-spaces in $\mathbb{R}^{10}$. First values for 15 blank cells ($N = 6$ – 20) in the row $m = 10$, $n = 4$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 6 \cdot N = 7 \cdot N = 8 \cdot N = 9 \cdot N = 10 \cdot N = 11 \cdot N = 12 \cdot N = 13 \cdot N = 14 \cdot N = 15 \cdot N = 16 \cdot N = 17 \cdot N = 18 \cdot N = 19 \cdot$
 $N = 20$

023. Packings of four-spaces in $\mathbb{R}^{11}$. First values for 34 blank cells ($N = 5$ – 20 , 83 – 100) in the row $m = 11$, $n = 4$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 5 \cdot N = 6 \cdot N = 7 \cdot N = 8 \cdot N = 9 \cdot N = 10 \cdot N = 11 \cdot N = 12 \cdot N = 13 \cdot N = 14 \cdot N = 15 \cdot N = 16 \cdot N = 17 \cdot N = 18 \cdot$
 $N = 19 \cdot N = 20 \cdot N = 83 \cdot N = 84 \cdot N = 85 \cdot N = 86 \cdot N = 87 \cdot N = 88 \cdot N = 89 \cdot N = 90 \cdot N = 91 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot$
 $N = 95 \cdot N = 96 \cdot N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100$

024. Packings of four-spaces in $\mathbb{R}^{12}$. First values for 21 blank cells ($N = 5$ – 20 , 94 – 95 , 97 , 99 – 100) in the row $m = 12$, $n = 4$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 5 \cdot N = 6 \cdot N = 7 \cdot N = 8 \cdot N = 9 \cdot N = 10 \cdot N = 11 \cdot N = 12 \cdot N = 13 \cdot N = 14 \cdot N = 15 \cdot N = 16 \cdot N = 17 \cdot N = 18 \cdot$
 $N = 19 \cdot N = 20 \cdot N = 94 \cdot N = 95 \cdot N = 97 \cdot N = 99 \cdot N = 100$

025. Packings of four-spaces in $\mathbb{R}^{13}$. First values for 32 blank cells ($N = 5$ – 27 , 92 – 100) in the row $m = 13$, $n = 4$ of the table of best Grassmannian packings, blank in Sloane's 1997 table and in Cohn's current one. Each value beats every packing obtained by deleting subspaces from a larger known one.

$N = 5 \cdot N = 6 \cdot N = 7 \cdot N = 8 \cdot N = 9 \cdot N = 10 \cdot N = 11 \cdot N = 12 \cdot N = 13 \cdot N = 14 \cdot N = 15 \cdot N = 16 \cdot N = 17 \cdot N = 18 \cdot$
 $N = 19 \cdot N = 20 \cdot N = 21 \cdot N = 22 \cdot N = 23 \cdot N = 24 \cdot N = 25 \cdot N = 26 \cdot N = 27 \cdot N = 92 \cdot N = 93 \cdot N = 94 \cdot N = 95 \cdot N = 96 \cdot$
 $N = 97 \cdot N = 98 \cdot N = 99 \cdot N = 100$

Discrete geometry

026. Spreading points in the plane. First values for $n = 32$ – 50 (19 cells) in Erich Friedman's table of point sets in the plane minimizing the ratio of the largest to the smallest distance, which stops at $n = 31$. Each value is smaller than what deleting a point from the next configuration gives.

$n = 32 \cdot n = 33 \cdot n = 34 \cdot n = 35 \cdot n = 36 \cdot n = 37 \cdot n = 38 \cdot n = 39 \cdot n = 40 \cdot n = 41 \cdot n = 42 \cdot n = 43 \cdot n = 44 \cdot n = 45 \cdot n = 46 \cdot$
 $n = 47 \cdot n = 48 \cdot n = 49 \cdot n = 50$