Variation connects repeated measurements with uncertainty

Contents

1 Describe spread through departures from a centre 2

2 A finite-data description and a population estimate answer different questions 4

3 Estimating the centre changes the variance estimator's expectation 7

4 Standard deviation describes precision on the measurement scale 15

5 Match the spread estimate to the repeated stage of the procedure 19

6 Independent repetition improves the precision of an average 23


1 Describe spread through departures from a centre

An average describes location, while the individual measurements show how consistently the procedure produces that location. The observations in Table 1 and their plot in Fig. 1 establish the finite data set. Each result contributes both to the average and to the observed spread around it.

Table 1. Authored repeated measurement values used to construct deviations and squared deviations.

Series	Values
A	9, 10, 13
B	5, 10, 17

Fig. 1. Repeated observations and their centre, retaining the observed spread around the mean.



A deviation subtracts the observed average from an individual value (Eq. 1). Its sign identifies the side of the centre, and its magnitude gives the distance in the original measurement unit. The signed deviations sum to zero (Eq. 2), an algebraic consequence of centring on the arithmetic mean. Squaring retains information about magnitude when the positive and negative departures balance.

Eq. 1. Deviation of an observed value from its sample mean, in the original measurement unit.

$$d_i = x_i - \bar{x}$$

Symbols
d_i Signed deviation of observation i from its sample mean.
i Observation index, running from 1 to the declared observation count.
x_i i-th observed value
$\bar{x}$ sample arithmetic mean

Eq. 2. The signed deviations from the arithmetic mean sum to zero.

$$\sum_{i=1}^{n} d_i = 0$$

Symbols
d_i Signed deviation of observation i from its sample mean.
i Observation index, running from 1 to the declared observation count.
n number of observations in the sample
$\sum$ Sum over the indexed observations.

The sum of squared deviations in Eq. 3 therefore describes aggregate departure. Dividing it by the number of observations gives the empirical mean squared deviation (Eq. 4), a descriptive summary of this finite data set. Squaring the original measurement unit is part of its meaning. Taking a square root later returns a spread summary to the original scale.

Eq. 3. Sum of squared deviations from the observed mean, with squared units.

$$d_i^2 = (x_i - \bar{x})^2$$
$$SS = \sum_{i=1}^{n} d_i^2$$

Symbols
Square each deviation x_i: observation i; i: index; $\bar{x}$: sample mean; d_i: their signed difference. Superscript 2 denotes squaring.
SS: Sum of Squares n: observation count; Σ: sum from 1 to n. SS adds the squared deviations. Squaring mg/L produces the unit $(\text{mg/L})^2$.

Eq. 4. Empirical mean squared deviation: squared departures averaged over the observed data.

$$\text{MSD} = \frac{SS}{n} = \frac{\sum_{i=1}^{n}(x_i - \bar{x})^2}{n}$$

Symbols
Mean squared deviation MSD: mean squared deviation; SS: sum of squared deviations. x_i: observation i; i: index; $\bar{x}$: sample mean; n: count; Σ: sum from 1 to n.
Squared unit, not absolute distance The deviations are squared before averaging, so MSD has unit $(\text{mg/L})^2$. Division by a count does not change that unit. MSD is not mean absolute distance.

2 A finite-data description and a population estimate answer different questions

Describing the recorded values and estimating variation in a generating population require different reference objects. The empirical quantity in Eq. 5 summarizes the fixed observations. The population quantity in Eq. 6 describes variation under the probability model. To study an estimator, we imagine repeatedly drawing new samples from that model and recalculating the statistic each time.

Eq. 5. Empirical squared-deviation summary of a fixed observed sample.

$$\mathrm{MSD}_n = \frac{1}{n}\sum_{i=1}^{n}(x_i - \bar{x})^2$$

Symbols
MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.
i Observation index, running from 1 to the declared observation count.
n number of observations in the sample
x_i i-th observed value
$\bar{x}$ sample arithmetic mean
$\sum$ Sum over the indexed observations.

Eq. 6. Population variance as an expectation under the specified probability model.

$$\begin{aligned} \mu &= \mathrm{E}[X] \\ \mathrm{Var}(X) &= \mathrm{E}[(X - \mathrm{E}[X])^2] \\ &= \mathrm{E}[(X - \mu)^2] = \sigma^2 \end{aligned}$$

Symbols
E Expectation, the probability-weighted average of a random quantity under repeated sampling.
Var Variance: expected squared deviation of a random quantity from its expectation.
X One random observation from the population, before a specific value is observed.
μ population mean
σ^2 Population variance, the expected squared deviation from the population mean.

Expectation rules express the average behaviour over those repeated samples. Fixed-factor and linearity rules appear in Eq. 7 and Eq. 8. Variance rules additionally account for dependence: Eq. 9 handles fixed transformations, and Eq. 10 retains covariance contributions. Independence is a sufficient condition for the relevant covariance terms to vanish; the assumptions must match the sampling design.

Eq. 7. Expectation rule for fixed terms, with the full authored conventions.

$$\begin{aligned} \mathrm{E}[c] &= c \\ \mathrm{E}[aY + b] &= a\,\mathrm{E}[Y] + b \end{aligned}$$

Symbols
E Expectation, the probability-weighted average of a random quantity under repeated sampling.
Y A random quantity with the moments required by the stated identity.
a Fixed real scale factor in an affine transformation.
b Fixed real additive shift in an affine transformation.
c A scalar placeholder; a fixed constant when pulled outside a sum or expectation.

Eq. 8. Linearity of expectation for the authored random quantities.

$$\mathrm{E}[U + V] = \mathrm{E}[U] + \mathrm{E}[V]$$
$$\mathrm{E}[\textstyle\sum_{i=1}^{n} Y_i] = \textstyle\sum_{i=1}^{n} \mathrm{E}[Y_i]$$

Symbols
E Expectation, the probability-weighted average of a random quantity under repeated sampling.
U An integrable random quantity in an expectation identity.
V A second integrable random quantity.
Y_i The indexed random quantity i in a finite sum.
i Observation index, running from 1 to the declared observation count.
n number of observations in the sample
$\sum$ Sum over the indexed observations.

Eq. 9. Variance under fixed shifts and scaling.

$$\mathrm{Var}(c) = 0$$
$$\mathrm{Var}(aY + b) = a^2 \,\mathrm{Var}(Y)$$

Symbols
Var Variance: expected squared deviation of a random quantity from its expectation.
Y A random quantity with the moments required by the stated identity.
a Fixed real scale factor in an affine transformation.
b Fixed real additive shift in an affine transformation.
c A scalar placeholder; a fixed constant when pulled outside a sum or expectation.

Eq. 10. Variance of combined random quantities, including covariance and its conditions.

$$\begin{aligned}\mathrm{Var}(U + V) = \mathrm{Var}(U) + \mathrm{Var}(V) \\ + 2\,\mathrm{Cov}(U, V)\end{aligned}$$
$$\begin{aligned}\mathrm{Cov}(U, V) = \mathrm{E}[(U - \mathrm{E}[U]) \\ \cdot (V - \mathrm{E}[V])]\end{aligned}$$

Symbols
Cov Covariance of two random quantities; it measures their joint linear variation.
E Expectation, the probability-weighted average of a random quantity under repeated sampling.
U An integrable random quantity in an expectation identity.
V A second integrable random quantity.
Var Variance: expected squared deviation of a random quantity from its expectation.

The random-sample expression in Eq. 11 makes the empirical calculation into an estimator. Its target question is stated in Eq. 12. The repeated-sampling simulation in Fig. 2 illustrates how the resulting estimates vary. Averaging estimates over many simulated samples investigates the estimator's expectation under the chosen model; the spread of those estimates describes a second, sampling-level source of variation.

Eq. 11. Mean squared deviation calculated from a random sample and its random sample mean.

$$\mathrm{MSD}_n = \frac{1}{n}\sum_{i=1}^{n}(X_i - \bar{X})^2$$


Symbols
MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.
X_i Random observation i before its value x_i is observed.
$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.
i Observation index, running from 1 to the declared observation count.
n number of observations in the sample
$\sum$ Sum over the indexed observations.

Eq. 12. Expectation question connecting the random squared-deviation summary with population variance.

$$\mathrm{E}[\mathrm{MSD}_n] \stackrel{?}{=} \sigma^2$$

Symbols
E Expectation, the probability-weighted average of a random quantity under repeated sampling.
MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.
σ^2 Population variance, the expected squared deviation from the population mean.

Fig. 2. Repeated-sampling simulation of the variance calculation under the authored model and parameters.



3 Estimating the centre changes the variance estimator's expectation

The sample mean is calculated from the same observations whose departures are being squared. This shared origin matters: a fitted centre follows the sample and reduces its aggregate squared departure relative to using the population centre. The algebra in Eq. 13 separates those contributions before any expectation is taken.

Eq. 13. Stepwise squared-deviation decomposition, retaining the authored operations, explanations and symbol legend.

Find the source of the MSD shortfall

$$X_i - \bar{X} = (X_i - \mu) - (\bar{X} - \mu)$$

Square, sum and collect the centered terms (Mathematical implication)

Expand each centered difference, distribute the finite sum, and use Σ(a_i−c) = n($\bar{a}$−c). The two cross-term contributions combine with the last squared shift to leave one negative squared shift. The fitted mean removes a nonnegative term. This is an exact algebraic identity for each realization, not an expectation or an unbiasedness claim.

$$\frac{1}{n}\sum_i (a_i - \bar{a})^2 = \frac{1}{n}\sum_i (a_i - c)^2 - (\bar{a} - c)^2$$

Centering at the fitted arithmetic mean removes a nonnegative squared shift. The result is an exact identity, not an average statement. Open the full algebra detail for each expansion and summation operation. The population mean is ordinarily unknown; this identity explains the estimator, rather than giving a formula that requires knowing μ.

$$\boxed{\mathrm{MSD}_n = \frac{1}{n}\sum_{i=1}^{n}(X_i - \mu)^2 - (\bar{X} - \mu)^2}$$

Arithmetic mean of the same n values n is a positive integer, $\bar{a}$ = (1/n)Σa_i, and c is fixed with respect to the index. The identity holds deterministically for every finite realization.

$$\boxed{\mathrm{MSD}_n = \frac{1}{n}\sum_{i=1}^{n}(X_i - \mu)^2 - (\bar{X} - \mu)^2}$$

Symbols MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.

X_i Random observation i before its value x_i is observed.

$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.

i Observation index, running from 1 to the declared observation count.

n number of observations in the sample

$\sum$ Sum over the indexed observations.

μ population mean

Full algebra: why fitting the mean removes a squared shift

Full algebra: why fitting the mean removes a squared shift

$$X_i - \bar{X} = (X_i - \mu) - (\bar{X} - \mu)$$

Square the centered difference and expand (Mathematical implication)

Multiplying a+b by itself gives a^2, the two equal cross-products ab, and b^2. Squaring both sides of an equality is a valid implication, not generally a reversible operation without sign information.

$$(a + b)^2 = a^2 + 2ab + b^2$$

Use a = $X_i - \mu$ and b = $-(\bar{X} - \mu)$ in the binomial identity. Squaring an equality gives a consequence; we do not claim to recover all signs by reversing this step.

$$\begin{aligned}(X_i - \bar{X})^2 = (X_i - \mu)^2 \\ - 2(X_i - \mu)(\bar{X} - \mu) \\ + (\bar{X} - \mu)^2\end{aligned}$$

$$\begin{aligned}(X_i - \bar{X})^2 = (X_i - \mu)^2 \\ - 2(X_i - \mu)(\bar{X} - \mu) \\ + (\bar{X} - \mu)^2\end{aligned}$$

Symbols X_i Random observation i before its value x_i is observed.

$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.

μ population mean

Full algebra: why fitting the mean removes a squared shift

$$\begin{aligned}(X_i - \bar{X})^2 = (X_i - \mu)^2 \\ - 2(X_i - \mu)(\bar{X} - \mu) \\ + (\bar{X} - \mu)^2\end{aligned}$$

Sum, divide by n and factor index-independent terms (Mathematical implication)

Use linearity for each of the three summands and move the index-independent coefficient c outside the middle sum. The last sum is not simplified yet.

$$\begin{aligned}\sum_{i=1}^{n}(a_i + cb_i + q) \\ = \sum_{i=1}^{n} a_i + c\sum_{i=1}^{n} b_i \\ + \sum_{i=1}^{n} q\end{aligned}$$

Distribute the finite sum over the three terms, then divide by the positive n. The same $\bar{X}-\mu$ occurs at every index, so it factors out of the cross term. Its squared last term is added n times; the divisor n cancels that count. This is deterministic within each realization.

$$\begin{aligned}\frac{1}{n}\sum_i (X_i-\bar{X})^2 = \frac{1}{n}\sum_i (X_i-\mu)^2 \\ -\frac{2}{n}(\bar{X}-\mu)\sum_i (X_i-\mu) \\ +(\bar{X}-\mu)^2\end{aligned}$$

Constant with respect to the index c has the same value for every index i within the sum; n is a positive integer.

$$\begin{aligned}\frac{1}{n}\sum_i (X_i-\bar{X})^2 = \frac{1}{n}\sum_i (X_i-\mu)^2 \\ -\frac{2}{n}(\bar{X}-\mu)\sum_i (X_i-\mu) \\ +(\bar{X}-\mu)^2\end{aligned}$$

Symbols X_i Random observation i before its value x_i is observed.

$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.

i Observation index, running from 1 to the declared observation count.

n number of observations in the sample

$\sum$ Sum over the indexed observations.

μ population mean

Full algebra: why fitting the mean removes a squared shift

$$\begin{aligned}\frac{1}{n}\sum_i (X_i-\bar{X})^2 = \frac{1}{n}\sum_i (X_i-\mu)^2 \\ -\frac{2}{n}(\bar{X}-\mu)\sum_i (X_i-\mu) \\ +(\bar{X}-\mu)^2\end{aligned}$$

Insert the sum of population-centered distances (Equivalent transformation)

Split the finite sum over subtraction, count c exactly n times, and substitute Σa_i = n $\bar{a}$. This needs the arithmetic mean of those same values and does not require probability assumptions.

$$\sum_{i=1}^{n}(a_i-c) = n(\bar{a}-c)$$

$\Sigma(\mathrm{X}_i-\mu) = \Sigma\mathrm{X}_i-\mathrm{n}\mu = \mathrm{n}(\bar{X}-\mu)$, because $\bar{X}$ is the mean of these same observations. Substitute that sum into the cross term: $-(2/\mathrm{n})(\bar{X}-\mu)\mathrm{n}(\bar{X}-\mu) = -2(\bar{X}-\mu)^2$.

$$\begin{aligned}\frac{1}{n}\sum_i (X_i-\bar{X})^2 = \frac{1}{n}\sum_i (X_i-\mu)^2 \\ -2(\bar{X}-\mu)^2+(\bar{X}-\mu)^2\end{aligned}$$

Arithmetic mean of the same n values n is a positive integer, $\bar{a} = (1/\mathrm{n})\Sigma \mathrm{a_i}$, and c is fixed with respect to the index. The identity holds deterministically for every finite realization.

$$\frac{1}{n}\sum_i (X_i - \bar{X})^2 = \frac{1}{n}\sum_i (X_i - \mu)^2 - 2(\bar{X} - \mu)^2 + (\bar{X} - \mu)^2$$

Symbols X_i Random observation i before its value x_i is observed.

$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.

i Observation index, running from 1 to the declared observation count.

n number of observations in the sample

$\sum$ Sum over the indexed observations.

μ population mean

Full algebra: why fitting the mean removes a squared shift

$$\frac{1}{n}\sum_i (X_i - \bar{X})^2 = \frac{1}{n}\sum_i (X_i - \mu)^2 - 2(\bar{X} - \mu)^2 + (\bar{X} - \mu)^2$$

Combine the squared shifts and identify the MSD (Equivalent transformation)

Expand each centered difference, distribute the finite sum, and use $\Sigma(\mathrm{a_i}-\mathrm{c}) = \mathrm{n}(\bar{a}-\mathrm{c})$. The two cross-term contributions combine with the last squared shift to leave one negative squared shift. The fitted mean removes a nonnegative term. This is an exact algebraic identity for each realization, not an expectation or an unbiasedness claim.

$$\frac{1}{n}\sum_i (a_i - \bar{a})^2 = \frac{1}{n}\sum_i (a_i - c)^2 - (\bar{a} - c)^2$$

The coefficients of the shared squared shift are -2 and $+1$, leaving -1. The left side is the defining MSD about the fitted sample mean. The result is exact for every sample. It does not by itself prove the estimator's average bias.

$$\boxed{\mathrm{MSD}_n = \frac{1}{n}\sum_{i=1}^{n} (X_i - \mu)^2 - (\bar{X} - \mu)^2}$$

Arithmetic mean of the same n values n is a positive integer, $\bar{a} = (1/\mathrm{n})\Sigma \mathrm{a_i}$, and c is fixed with respect to the index. The identity holds deterministically for every finite realization.

$$\boxed{\mathrm{MSD}_n = \frac{1}{n}\sum_{i=1}^{n} (X_i - \mu)^2 - (\bar{X} - \mu)^2}$$

Symbols MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.

X_i Random observation i before its value x_i is observed.

$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.

i Observation index, running from 1 to the declared observation count.

n number of observations in the sample

$\sum$ Sum over the indexed observations.

μ population mean

Applying the stated sampling assumptions gives the expectation calculation in Eq. 14. The variance of the sample mean (Eq. 15) supplies the term that accounts for estimating the centre. For independent observations with a common finite variance, dividing the squared-deviation sum by the sample size yields an expectation reduced by the factor shown in Eq. 16. This identifies the estimator's average relationship to its target.

Eq. 14. Expectation of the decomposition under the authored sampling assumptions.

Average the MSD identity and recognize variance

$$\begin{aligned}\mathrm{E}[\mathrm{MSD}_n] = \frac{1}{n}\sum_i \mathrm{E}[(X_i - \mu)^2] \\ - \mathrm{E}[(\bar{X} - \mu)^2]\end{aligned}$$

Recognize expected centered squares as variances (Requires statistical assumptions)

For a random variable with finite second moment, variance is the expected squared distance from its expectation.

$$\mathrm{Var}(Y) = \mathrm{E}[(Y - \mathrm{E}[Y])^2]$$

Linearity gives the displayed expectation of the finite squared-distance average; it needs no independence. Each population-centered squared observation has expectation σ^2. The sample mean has expectation μ under the common-mean condition, so its expected squared shift is $\mathrm{Var}(\bar{X})$. The arithmetic average of n copies of σ^2 is σ^2. The complete sample-mean argument is available with the next result.

$$\begin{aligned}\mathrm{E}[(X_i - \mu)^2] &= \sigma^2 \\ \mathrm{E}[(\bar{X} - \mu)^2] &= \mathrm{Var}(\bar{X})\end{aligned}$$

Common mean and finite variance; fixed n $\geq$ 1 For each i = 1, ..., n, E[X_i] = μ and Var(X_i) = $\sigma^2 < \infty$. The positive integer n is fixed. Identical distribution is sufficient but equality of these moments is all that is needed here.

Finite second moment Y has a finite second moment, so its expectation and variance exist.

$$\mathrm{E}[\mathrm{MSD}_n] = \sigma^2 - \mathrm{Var}(\bar{X})$$

Symbols E Expectation, the probability-weighted average of a random quantity under repeated sampling.

MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.

Var Variance: expected squared deviation of a random quantity from its expectation.

X_i Random observation i before its value x_i is observed.

$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.

i Observation index, running from 1 to the declared observation count.

n number of observations in the sample

$\sum$ Sum over the indexed observations.

μ population mean

σ^2 Population variance, the expected squared deviation from the population mean.

Eq. 15. Variance of the sample mean under the stated common-variance sampling model.

$$\boxed{\text{Var}(\bar{X}) = \frac{\sigma^2}{n}}$$

Symbols
Var Variance: expected squared deviation of a random quantity from its expectation.
$\bar{X}$ Random sample mean across repeated samples, as distinct from an observed sample mean $\bar{x}$.
n number of observations in the sample
σ^2 Population variance, the expected squared deviation from the population mean.

Eq. 16. Expectation factor for the squared-deviation estimator divided by sample size.

From the fitted-mean identity to unbiased variance

$$\text{E}[\text{MSD}_n] = \sigma^2 - \frac{\sigma^2}{n}$$

Factor out the population variance (Equivalent transformation)

Write a as na/n, subtract a/n and factor out a. The result holds for every real a and positive integer n. This is algebra, not an additional statistical assumption.

$$a - \frac{a}{n} = \frac{n-1}{n}a$$

In the general factorization use a = σ^2. The fraction left after subtraction is (n−1)/n, strictly less than one for n > 1. Thus the MSD is biased downward when $\sigma^2 > 0$. A particular realized MSD may lie above or below σ^2. Bias is the repeated-sampling expectation, not the direction of every individual error.

$$\text{E}[\text{MSD}_n] = \frac{n-1}{n}\sigma^2$$

Finite observation count n is a positive integer; all indexed equalities hold for i = 1, ..., n.

$$\text{E}[\text{MSD}_n] = \frac{n-1}{n}\sigma^2$$

Symbols E Expectation, the probability-weighted average of a random quantity under repeated sampling.

MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.

n number of observations in the sample

σ^2 Population variance, the expected squared deviation from the population mean.

Bessel's correction compensates for that factor (Eq. 17). For at least two observations under the stated common-variance, uncorrelated sampling conditions, the practical formula in Eq. 18 is unbiased for population variance. The simulation in Fig. 3 compares this expectation behaviour across repeated samples. Unbiasedness describes an average over sampling; a particular sample still supplies an estimate with its own sampling variation.

Eq. 17. Bessel correction derived from the estimator's expectation factor.

Bessel's correction

$$\mathrm{E}[\mathrm{MSD}_n] = \frac{n-1}{n}\sigma^2$$

Multiply by the reciprocal bias factor, for n > 1 (Requires statistical assumptions)

A constant fixed across repeated samples multiplies the expectation. A random offset such as Δ must not itself be pulled outside E; the fixed count n may be.

$$\mathrm{E}[cY] = c\,\mathrm{E}[Y]$$

Define the random statistic S^2 by multiplying MSD_n by the fixed reciprocal n/(n−1). Linearity moves that coefficient outside E and the factors cancel. S^2 is unbiased under the stated assumptions. Its observed value s^2 is not generally exactly σ^2, and the square root is not automatically unbiased.

$$\mathrm{E}[S^2] = \frac{n}{n-1}\,\mathrm{E}[\mathrm{MSD}_n] = \frac{n}{n-1}\frac{n-1}{n}\sigma^2 = \sigma^2$$

Common mean and finite variance; fixed n ≥ 1 For each i = 1, ..., n, E[X_i] = μ and Var(X_i) = $\sigma^2 < \infty$. The positive integer n is fixed. Identical distribution is sufficient but equality of these moments is all that is needed here.

Expectations exist The displayed random quantities have finite expectations. A constant moved outside E is fixed across samples.

Nonzero divisor n > 1, so n−1 is positive and division by n−1 is valid.

Pairwise zero covariance Cov(X_i,X_j) = 0 for distinct indices. Independent finite-variance observations satisfy this condition; independence is sufficient, not necessary. Correlated observations require covariance terms.

$$S^2 := \frac{n}{n-1}\mathrm{MSD}_n$$
$$\boxed{\mathrm{E}[S^2] = \sigma^2}$$

Symbols E Expectation, the probability-weighted average of a random quantity under repeated sampling.

MSD_n Mean squared deviation about the fitted sample mean, with divisor n; a statistic across repeated samples.

S^2 The corrected sample variance statistic across hypothetical samples; lowercase s^2 is its realized value.

n number of observations in the sample


σ^2 Population variance, the expected squared deviation from the population mean.

Eq. 18. Practical sample-variance estimator with denominator sample size minus one and complete authored notation.

$$s^2 = \frac{1}{n-1}\sum_{i=1}^{n}(x_i - \bar{x})^2$$

Symbols
i Observation index, running from 1 to the declared observation count.
n number of observations in the sample
s^2 Sample variance, residual sum of squares divided by n minus one for the usual mean-fitted iid estimator.
x_i i-th observed value
$\bar{x}$ sample arithmetic mean
$\sum$ Sum over the indexed observations.

Fig. 3. Repeated-sampling behaviour of the corrected variance estimator under the authored simulation conditions.



4 Standard deviation describes precision on the measurement scale

Taking the square root of sample variance gives the sample standard deviation (Eq. 19). Its unit matches the original observations, so the interpretation in Eq. 20 can be expressed directly on the measurement scale. The summary table in Table 2 keeps variance and standard deviation connected while distinguishing their units.

Eq. 19. Sample standard deviation as the square root of sample variance, in the original measurement unit.

Concentration → Variance → Standard deviation

$$s_A^2 = \frac{13}{3}\ (\text{mg/L})^2$$

Take the nonnegative square root (Mathematical implication)

Standard deviation is the nonnegative square root of variance. Taking the square root also restores the original physical unit. It is not the mean absolute deviation, and an unbiased variance estimate does not imply an unbiased square root.

$$s = \sqrt{s^2}$$

Series A has sample variance 13/3 $(\text{mg/L})^2$. Take its nonnegative square root: $\sqrt{(13/3)} \approx 2.081666$, reported here as 2.082 mg/L. The concentration unit is no longer squared. This is a root-mean-square measure of residual size with the stated divisor, not an arithmetic mean of absolute distances.

$$s_A = \sqrt{\frac{13}{3}}\ \text{mg/L} \approx 2.082\ \text{mg/L}$$

$$s_A = \sqrt{\frac{13}{3}}\ \text{mg/L} \approx 2.082\ \text{mg/L}$$

Symbols s sample standard deviation

s^2 Sample variance, residual sum of squares divided by n minus one for the usual mean-fitted iid estimator.

$\sqrt{\ }$ Nonnegative square root.

Eq. 20. Authored interpretation of standard deviation on the observed measurement scale.

$$s = \sqrt{\frac{\sum_{i=1}^{n}(x_i - \bar{x})^2}{n-1}}$$

Symbols
i Observation index, running from 1 to the declared observation count.
n number of observations in the sample
s sample standard deviation
x_i i-th observed value
$\bar{x}$ sample arithmetic mean
$\sum$ Sum over the indexed observations.
$\sqrt{\ }$ Nonnegative square root.

Table 2. Location and dispersion summaries with their respective units.

Summary	A	B
Mean (mg/L)	$32/3 \approx 10.667$	$32/3 \approx 10.667$
MSD ((mg/L)^2)	$26/9 \approx 2.889$	$218/9 \approx 24.222$
s^2 ((mg/L)^2)	$13/3 \approx 4.333$	$109/3 \approx 36.333$
s (mg/L)	2.082	6.028

The optional sources in Listing 1 and Listing 2 reproduce the calculations. For quality-control repeats, the observations in Table 3 describe precision under the repeated conditions. The target illustration in Fig. 4 contrasts narrower and wider scatter around a deliberately chosen reference centre. Assessing trueness additionally requires evidence connecting the method's expectation with an appropriate reference value.

Listing 1. Optional R reproduction of the variance calculation.

```
a <- c(9, 10, 13)
b <- c(5, 10, 17)
mean_a <- mean(a)
mean_b <- mean(b)
c(mean_a, mean_b)
```

c(...)

Combines values into a vector; also collects results for output.

mean(x)

Calculates the arithmetic mean of the values in x.

name <- value

Assignment operator: stores a value under a name.

Listing 2. Optional R reproduction of the standard-deviation calculation.

```
a <- c(9, 10, 13)
b <- c(5, 10, 17)
va <- var(a)
vb <- var(b)
sa <- sd(a)
sb <- sd(b)
c(va, vb, sa, sb)
```

c(...)

Combines input values or collects output results into a vector.

var(x)

Calculates sample variance using divisor n−1.


sd(x)

Calculates sample SD, the square root of var(x).

Table 3. Five quality-control repeats per method, with their means and sample standard deviations in mg/L.


Method	Values	Mean	SD
A	9.8, 10.0, 10.1, 9.9, 10.2	10.000	0.158
B	8.7, 10.8, 9.5, 11.2, 9.8	10.000	1.007

Fig. 4. Illustrative target patterns centred by construction: crosses have smaller spread and open circles have larger spread. The coordinates are schematic.



The shifted-data comparison in Fig. 5 clarifies that location and spread are complementary. A common shift changes the mean while preserving distances between observations and their centre. Consequently, a precision statement identifies the measurement conditions and the variation it summarizes; a broader quality statement also addresses calibration, reference values and possible systematic effects.

Fig. 5. A common shift changes the location of the authored observations while retaining their spread.



5 Match the spread estimate to the repeated stage of the procedure

A measurement result can vary because of several stages: collection, preparation and instrument reading. The process diagram in Fig. 6 identifies the authored stages. Repeating a reading of one preparation represents a different experiment from preparing new portions or returning to new sampling sites.

Fig. 6. Measurement-process stages used to identify the scope of replication and variation components.

1. Sampling site

Between-site variation

⟶ 2. collect

2. Independent sample

Between samples within a site

⟶ 3. prepare

3. Sample preparation

Between preparations of a sample

⟶ 4. measure

4. Instrument repeat

Instrumental replicates within a preparation

The component model in Eq. 21 describes how the relevant variance contributions combine under its stated assumptions. Converting the combined variance to standard deviation gives Eq. 22. Variances add for the specified uncorrelated components; standard deviations combine through the corresponding square-root calculation. A design with dependence requires its covariance structure to be included.

Eq. 21. Variance-component combination with the complete authored assumptions and notation.

Combining independent analytical variance contributions

$$\sigma^2_{\text{total}} = \text{Var}(A + B + C)$$

Add the independent component variances (Requires statistical assumptions)

For independent finite-variance contributions, covariance terms vanish. Therefore Var(A+B) = Var(A)+Var(B), and the finite-sum extension gives total variance as the sum of component variances. For correlated sources Var(A+B) also contains 2 Cov(A,B). This law concerns an assumed model, not differences between pooled experimental SDs.

$$\text{Var}(A + B) = \text{Var}(A) + \text{Var}(B)$$
$$\sigma^2_{\text{total}} = \sum_k \sigma^2_k$$

Treat A, B and C as additive random contributions, all expressed in mg/L. Their illustrative variances are 4, 1 and 0.25 $(\text{mg/L})^2$. Independence removes the covariance terms. Add the three variances to obtain 5.25 $(\text{mg/L})^2$. The numbers are model inputs for teaching, not variance components isolated from the lake or QC data.

$$\sigma^2_{\text{total}} = 4 + 1 + 0.25 = 5.25\ (\text{mg/L})^2$$

Illustrative independent additive model Illustrative model, not estimated components: sampling variance 4, preparation variance 1, instrument variance 0.25 $(\text{mg/L})^2$. Assume independent additive contributions with finite variance. Correlated sources require covariance terms.

$$\sigma^2_{\text{total}} = 4 + 1 + 0.25 = 5.25\ (\text{mg/L})^2$$

Symbols A Additive sampling contribution in the illustrative model.

B Additive preparation contribution in the illustrative model.

C Additive instrument contribution in the illustrative model.

Var Variance: expected squared deviation of a random quantity from its expectation.

σ^2_{total} Total variance in $(\text{mg/L})^2$.

Eq. 22. Standard deviation corresponding to the combined component variance.

Combining independent analytical variance contributions

$$\sigma^2_{\text{total}} = 4 + 1 + 0.25 = 5.25\ (\text{mg/L})^2$$

Take one square root after adding (Mathematical implication)

After adding variances in the same squared unit, take one nonnegative square root. Standard deviations are not directly additive. This root-sum-of-squares calculation needs the independence/additivity assumptions used in the preceding step.

$$\sigma_{\text{total}} = \sqrt{\sum_k \sigma_k^2}$$

Take one square root of the total 5.25 $(\text{mg/L})^2$, giving approximately 2.291 mg/L. Adding the individual SDs would give 2 + 1 + 0.5 = 3.5 mg/L, which is not the SD of this independent sum. If contributions were correlated, include covariance terms before taking the square root.

$$\sigma_{\text{total}} = \sqrt{5.25}\ \text{mg/L} \approx 2.291\ \text{mg/L}$$

Illustrative independent additive model Illustrative model, not estimated components: sampling variance 4, preparation variance 1, instrument variance 0.25 $(\text{mg/L})^2$. Assume independent additive contributions with finite variance. Correlated sources require covariance terms.

$$\sigma_{\text{total}} = \sqrt{5.25}\ \text{mg/L} \approx 2.291\ \text{mg/L}$$

Symbols σ_{total} Total standard deviation in mg/L.

σ^2_{total} Total variance in $(\text{mg/L})^2$.


$\sqrt{\ }$ Nonnegative square root.

The replication matrix in Table 4 connects repeated stages with the variation they can reveal. The site comparison in Fig. 7 adds the spatial context. Relative standard deviation (Eq. 23) expresses spread relative to the mean; the authored comparison in Table 5 retains both absolute and relative information. A useful relative comparison requires a meaningful ratio scale and a reference mean sufficiently separated from zero.

Table 4. Replication design matrix connecting repeated procedural stages with the variation represented.

Repeated level	Measurement structure	Readings	Included variation / shared levels
Instrumental repeats	3 injections, one vial	3	Instrument; preparation and field sample shared
Preparation repeats	3 preparations × 3 injections	9	Preparation + instrument; field sample shared
Field samples	3 samples × 3 preparations × 3 injections	27	Within-site sampling + preparation + instrument; site shared
Sampling sites	3 sites × 3 samples × 3 preparations × 3 injections	81	Between-site + all downstream variation; nested readings remain dependent

Fig. 7. Site-level measurement comparison under the authored sampling and replication design.



Eq. 23. Relative standard deviation, including the ratio-scale and denominator conventions.

$$\mathrm{RSD} = \frac{s}{|\bar{x}|}, \qquad \mathrm{RSD}(\%) = 100\,\frac{s}{|\bar{x}|}$$

Symbols
Spread relative to level RSD: relative standard deviation; s: sample SD; $\bar{x}$: sample mean. $|\ |$ takes its absolute value. The mean must be nonzero and sufficiently far from zero.
Ratio or percentage The units cancel. RSD is dimensionless. Multiplying by 100 expresses its numerical percentage; % denotes percent. A ratio of 0.02 is 2%.
Meaningful ratios Use a suitable ratio-scale quantity with a meaningful zero and compatible measurement conditions. A nonzero mean alone is not sufficient.

Table 5. Absolute and relative spread summaries for the authored comparison data.

Quantity	QC A (mg/L)	QC B (g/L)	QC B back in mg/L
Mean	10	0.010	10
Sample SD	0.158114	0.00100747	1.00747
RSD (%)	1.5811	10.0747	10.0747

6 Independent repetition improves the precision of an average

The standard deviation of individual results and the standard deviation of their average refer to different random quantities. Under independent sampling with a common finite variance, the sample-mean variance derived earlier gives the population standard error in Eq. 24. Replacing the population standard deviation by its sample estimate gives Eq. 25.

Eq. 24. Population standard error of an average under the stated independent common-variance sampling model.

$$\mathrm{SE}(\bar{Y}) = \frac{\sigma}{\sqrt{n}}$$

Symbols
SE Standard error: the standard deviation of an estimator's sampling distribution. Here, random uncertainty of the sample mean under the stated sampling assumptions.
$\bar{Y}$ Random sample mean of n measurements Y_i; Y is a local renaming of the earlier measurement variable X, not a new dataset.
n number of observations in the sample
σ population standard deviation

Eq. 25. Estimated standard error using the sample standard deviation.

$$\widehat{\mathrm{SE}}(\bar{Y}) = \frac{s}{\sqrt{n}}$$

Symbols
$\bar{Y}$ Random sample mean of n measurements Y_i; Y is a local renaming of the earlier measurement variable X, not a new dataset.
$\widehat{\mathrm{SE}}$ Estimated standard error, using the observed sample SD s in place of unknown population SD σ.
n number of observations in the sample
s sample standard deviation

The square-root rate in Eq. 26 explains why improving mean precision through repetition has diminishing returns: reducing the standard error by a given factor requires increasing the independent sample size by that factor squared. The worked example in Eq. 27 keeps the authored units and sample size visible. The individual-result spread remains a property of the measurement conditions while the average becomes more stable.

Eq. 26. Square-root sample-size relationship for mean precision under the authored assumptions.

$$\mathrm{SE}_n = \frac{\sigma}{\sqrt{n}} \quad \xrightarrow[n\to\infty]{} 0$$

$$\mathrm{SE}_{4n} = \frac{1}{2}\,\mathrm{SE}_n$$

Symbols
SE Standard error: the standard deviation of an estimator's sampling distribution. Here, random uncertainty of the sample mean under the stated sampling assumptions.
n number of observations in the sample
σ population standard deviation

Eq. 27. Worked standard-error calculation with the authored sample size and measurement units.


$$\widehat{\mathrm{SE}}(\bar{Y}) = \frac{0.1581}{\sqrt{5}} \approx 0.0707 \text{ mg/L}$$

$$\boxed{\bar{y} \pm \widehat{\mathrm{SE}}(\bar{Y}) = (10.00 \pm 0.07) \text{ mg/L}}$$

Symbols
$\bar{Y}$ Random sample mean of n measurements Y_i; Y is a local renaming of the earlier measurement variable X, not a new dataset.
$\widehat{\mathrm{SE}}$ Estimated standard error, using the observed sample SD s in place of unknown population SD σ.
$\bar{y}$ Observed arithmetic mean of the same measurements, using the local Y notation.

The replication comparison in Fig. 8 and optional source in Listing 3 connect this rate with design choices. Shared preparation effects or correlated readings change the effective information gained by repetition. Reporting an uncertainty for an average therefore identifies the repeated unit and its dependence assumptions. The next descriptive step examines distribution shape, because equal centres and spreads can still accompany different patterns of observations.

Fig. 8. Authored replication comparison illustrating the information gained by the specified repeated measurements.



Listing 3. Optional R reproduction of the replication comparison.

```
a <- c(9.8, 10.0, 10.1, 9.9, 10.2)
b <- c(8.7, 10.8, 9.5, 11.2, 9.8)
ma <- mean(a); mb <- mean(b)
sa <- sd(a); sb <- sd(b)
ra <- 100*sa/ma; rb <- 100*sb/mb
inside_a <- a >= 9.5 & a <= 10.5
inside_b <- b >= 9.5 & b <= 10.5
na <- sum(inside_a); nb <- sum(inside_b)
c(ma, mb, sa, sb, ra, rb, na, nb)
```

c(...)

Combines input values or collects output results into a vector.

mean(x)

Calculates the arithmetic mean of x.

sd(x)

Calculates sample SD using the corrected variance.

sum(x)

Adds values; for logical x, counts TRUE entries.

>=, <= and &

Operators: compare to inclusive bounds; & requires both conditions.