
Barycentric Coordinates 1

Barycentric coordinates enable you to evenly interpolate between three values among the verticies of a triangle.



$$U = (u_1, u_2, u_3) \tag{1}$$

$$v_1 = (x_1, y_1) (2)$$

$$v_2 = (x_2, y_2) \tag{3}$$

$$v_3 = (x_3, y_3) \tag{4}$$

$$p = (x_p, y_p) \tag{5}$$

The Barycentric coordinates can be defined in terms of the following relationships:

$$\begin{cases} u_1 + u_2 + u_3 = 1 \\ u_1 x_1 + u_2 x_2 + u_3 x_3 = x_p \\ u_1 y_1 + u_2 y_2 + u_3 y_3 = y_p \end{cases}$$
 (6)

Let's reduce the amount of variables in these equations:

$$u_3 = 1 - u_1 - u_2 \tag{7}$$

$$\begin{cases} u_1(x_1 - x_3) + u_2(x_2 - x_3) &= x_p - x_3 \\ u_1(y_1 - y_3) + u_2(y_2 - y_3) &= y_p - y_3 \end{cases}$$
 (8)

Now we can turn the system of equations into matrix form:

$$T = \begin{bmatrix} x_1 - x_3 & x_2 - x_3 \\ y_1 - y_3 & y_2 - y_3 \end{bmatrix}$$

$$U = \begin{bmatrix} u1 \\ u2 \end{bmatrix}$$

$$R = \begin{bmatrix} x_p - x_3 \\ y_p - y_3 \end{bmatrix}$$

$$(10)$$

$$U = \begin{bmatrix} u1\\u2 \end{bmatrix} \tag{10}$$

$$R = \begin{bmatrix} x_p - x_3 \\ y_p - y_3 \end{bmatrix} \tag{11}$$

$$T \cdot U = R \tag{12}$$

So the solution is

$$U = T^{-1} \cdot R \tag{13}$$

The main effort goes towards finding T^{-1}

$$T^{-1} = \frac{adj(T)}{det(T)} \tag{14}$$

$$det(T) = (x_1 - x_3)(y_2 - y_3) - (x_2 - x_3)(y_1 - y_3)$$
(15)

$$adj(T) = \begin{bmatrix} y_2 - y_3 & x_3 - x_2 \\ y_3 - y_1 & x_1 - x_3 \end{bmatrix}$$

$$\tag{16}$$

$$T^{-1} = \frac{1}{\det(T)} \cdot \begin{bmatrix} y_2 - y_3 & x_3 - x_2 \\ y_3 - y_1 & x_1 - x_3 \end{bmatrix}$$
 (17)

$$T^{-1} \cdot R = \frac{1}{\det(T)} \cdot \begin{bmatrix} (y_2 - y_3)(x_p - x_3) + (x_3 - x_2)(y_p - y_3) \\ (y_3 - y_1)(x_p - x_3) + (x_1 - x_3)(y_p - y_3) \end{bmatrix}$$
(18)

And the final formula you need to find (u_1, u_2, u_3) given the points v_1, v_2, v_3, p is

$$u_{1} = \frac{(y_{2} - y_{3})(x_{p} - x_{3}) + (x_{3} - x_{2})(y_{p} - y_{3})}{(x_{1} - x_{3})(y_{2} - y_{3}) - (x_{2} - x_{3})(y_{1} - y_{3})}$$

$$u_{2} = \frac{(y_{3} - y_{1})(x_{p} - x_{3}) + (x_{1} - x_{3})(y_{p} - y_{3})}{(x_{1} - x_{3})(y_{2} - y_{3}) - (x_{2} - x_{3})(y_{1} - y_{3})}$$

$$(20)$$

$$u_2 = \frac{(y_3 - y_1)(x_p - x_3) + (x_1 - x_3)(y_p - y_3)}{(x_1 - x_3)(y_2 - y_3) - (x_2 - x_3)(y_1 - y_3)}$$
(20)

$$u_3 = 1 - u_2 - u_1 \tag{21}$$