

Extraction of Effective Temperatures from Tsallis Fits to Transverse-Momentum Spectra of Direct Photons and Heavy Quarkonia in High-Energy Collisions

Peng-Cheng Zhang^{1,*}, Hailong Zhu^{1,†} and Fu-Hu Liu^{1,‡}

¹*Institute of Theoretical Physics & College of Physics and Electronic Engineering,
State Key Laboratory of Quantum Optics Technologies and Devices & Collaborative
Innovation Center of Extreme Optics, Shanxi University, Taiyuan 030006, China*

Abstract:

Keywords: Tsallis statistics; effective temperature

PACS numbers:

I. RESULTS AND DISCUSSION

A central feature of the nonprompt direct-photon spectra [1] is that they are not described by a single exponential: the local inverse slope rises with transverse momentum, and the data were accordingly characterized by two separate exponential fits, yielding $T_{\text{eff}} \approx 0.26$ GeV/ c in the range $0.8 < p_T < 1.9$ GeV/ c and $T_{\text{eff}} \approx 0.38$ GeV/ c in $2 < p_T < 4$ GeV/ c . The present analysis shows that this evolving slope is captured naturally, and without piecewise treatment, by a single Tsallis form fitted over the full momentum range.

The equivalence follows directly from the structure of the Tsallis function. For $dN/dp_T \propto p_T [1 + (q - 1)p_T/T]^{q/(1-q)}$, the local inverse slope is not constant but a smooth, monotonically increasing function of p_T ,

$$T_{\text{eff}}(p_T) \equiv - \left(\frac{d \ln(dN/dp_T)}{dp_T} \right)^{-1}, \quad (1)$$

which reduces to the constant T only in the Boltzmann limit $q \rightarrow 1$. A value $q > 1$ therefore encodes precisely the spectral hardening that the two-exponential description represents through a discrete jump in T_{eff} . The single parameter q thus subsumes the two independent slopes of the piecewise fit.

This is borne out quantitatively. Evaluating $T_{\text{eff}}(p_T)$ for the 0%–20% best-fit parameters ($q = 1.084$, $T = 0.128$ GeV) gives $T_{\text{eff}} \approx 0.26$ GeV/ c near $p_T = 1.3$ GeV/ c and $T_{\text{eff}} \approx 0.40$ GeV/ c near $p_T = 3$ GeV/ c , in close agreement with the two inverse slopes extracted independently [1] for the corresponding p_T intervals. The two regimes are recovered not as separate fits but as the low- and high- p_T limits of one continuous curve.

The Tsallis description is therefore both more economical and more physically suggestive. It replaces two phenomenological slopes by a single functional form with one shape parameter q , and it admits a transparent interpretation within superstatistics: a Tsallis spectrum arises from a superposition of Boltzmann emitters with a distribution of inverse temperatures β , with $q - 1 \simeq \text{Var}(\beta)/\langle \beta \rangle^2$. The hardening described piecewise by two slopes is thus reframed as a single spectrum emitted by a system spanning a range of temperatures—a picture consistent with photon emission integrated over the cooling history of the expanding fireball.

* 202312602003@email.sxu.edu.cn

† Correspondence: zhuhl@sxu.edu.cn

‡ Correspondence: fuhuliu@163.com; fuhuliu@sxu.edu.cn




Figure 1a. Nonprompt direct-photon spectra in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV for the four centrality classes, with the Tsallis fits (curves) evaluated at the best-fit (q, T, Nor) values.

-
- [1] N. J. Abdulameer *et al.* (PHENIX Collaboration), “Nonprompt direct-photon production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Physical Review C* 109, no. 4 (2024): 044912, <https://doi.org/10.1103/PhysRevC.109.044912>.




Figure 1d. Tsallis-fit confidence regions in the (q, T) plane for four centrality classes. Closed curves show the 68.3% joint confidence region ($\Delta\chi^2 = 2.30$); filled circles mark the best fits, with bars giving single-parameter 1σ errors ($\Delta\chi^2 = 1$).