



Idea paper: An experimental framework for determining the degree of intraguild predation in a three-species omnivorous food web

Journal:	<i>Ecological Research</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Special Feature
Date Submitted by the Author:	n/a
Complete List of Authors:	Hsu, Gen-Chang; National Taiwan University Department of Life Science,
Manuscript Category:	5: Species interaction, 40: Stable isotope
Manuscript Keywords:	community, experiment, ideas for fundamental questions, intraguild predation, stable isotope analysis
Abstract:	Intraguild predation (IGP) is common in natural and human-managed systems and plays a critical role in food web dynamics. Although previous studies have documented the occurrence of IGP across a wide range of predator taxa, few have qualitatively examined the degree of IGP. Here, I propose an experimental framework combining controlled feeding trials and stable isotope analysis of field samples to determine the degree of IGP in a three-species omnivorous food web. Such an approach can provide a useful tool for studying IGP in a more accurate (controlled feeding trials) and realistic (stable isotope analysis of field samples) fashion. If proven successful, the present framework can be extended to food webs involving more complex interactions (e.g., cannibalism, multiple prey) and further complemented with other approaches to capture a more complete picture of IGP dynamics in the field.

Article category

Special features

Title

Idea paper: An experimental framework for determining the degree of intraguild predation in a three-species omnivorous food web

Author

Gen-Chang Hsu¹

Author's institutional affiliation

¹Department of Life Science, National Taiwan University, Taipei, Taiwan

Correspondence

Gen-Chang Hsu, Department of Life Science, National Taiwan University, Taipei 10617, Taiwan.

Email: genchanghsu@gmail.com

ORCID: <https://orcid.org/0000-0002-6607-4382>

Abstract

Intraguild predation (IGP) is common in natural and human-managed systems and plays a critical role in food web dynamics. Although previous studies have documented the occurrence of IGP across a wide range of predator taxa, few have qualitatively examined the degree of IGP. Here, I propose an experimental framework combining controlled feeding trials and stable isotope analysis of field samples to determine the degree of IGP in a three-species omnivorous food web. Such an approach can provide a useful tool for studying IGP in a more accurate (controlled feeding trials) and realistic (stable isotope analysis of field samples) fashion. If proven successful, the present framework can be extended to food webs involving more complex interactions (e.g., cannibalism, multiple prey) and further complemented with other approaches to capture a more complete picture of IGP dynamics in the field.

Systematic keyword selection

community, experiment, ideas for fundamental questions, intraguild predation, stable isotope analysis

Research question

Intraguild predation (IGP) is common in natural and human-managed ecosystems (Arim & Marquet, 2004; Muller & Brodeur, 2002; Polis & Holt, 1992). Previous studies have documented the occurrence of IGP across various taxa (Polis et al., 1989). However, few have qualitatively examined the intensity/degree of IGP among predators. To address this gap, here I propose an experimental framework combining controlled feeding trials and stable isotope analysis of field samples to determine the degree of IGP in a three-species (top predator, mesopredator, and shared prey) omnivorous food web. The degree of IGP in this study is defined as the relative consumption of mesopredator (compared to shared prey) in the diet of top predator (i.e., a high degree of IGP means that the top predator consumes a high proportion of mesopredator in its diet).

Value

IGP could substantially affect the abundance and distribution of interacting species (Polis et al., 1989), which may have profound ecological and evolutionary consequences for food web dynamics. A better quantitative understanding of IGP can provide insights into the complex predator-predator-prey trophic interactions and may help predict the community structure and stability (Arim & Marquet, 2004; Nakazawa & Yamamura, 2006; Pahl et al., 2020). Furthermore, such understanding can have useful implications for agricultural management, for example, evaluation of the effectiveness of biocontrol agents in pest control programs (Muller & Brodeur, 2002).

Relevant hypothesis

Previous studies have used manipulative experiments (e.g., cage experiments) to assess the intensity of IGP by comparing the differences in the numbers of prey or mesopredator in the presence vs. absence of top predator (Denno et al., 2004; Provost et al., 2005). This approach can reveal the causal relationships between predator-prey interactions, allowing for strong inferences

about IGP. However, the use of enclosures could potentially alter the encounter rates between individuals and thus lead to biased results.

Stable isotopes, particularly nitrogen isotope ratios ($\delta^{15}\text{N}$), have been used to estimate the trophic level of predators in the field (Abd El-Wakeil, 2009; Halaj et al., 2005; Rickers et al., 2006). It is suggested that IGP would increase the $\delta^{15}\text{N}$ of predators and thus their trophic level (Ponsard & Arditi, 2000), but this proposal has not been experimentally verified. In addition, the trophic level of predators in previous studies was often calculated based on assumed trophic enrichment factors (TEFs) (e.g., Klarner et al., 2013; Ponsard & Arditi, 2000; Svanbäck et al., 2015), which can affect the subsequent inferences made about IGP in the field.

Recently, researchers have applied molecular gut content analysis (MGCA) and immunological techniques to reliably detect the presence of certain food items in predators' diet (Gagnon et al., 2011; Hagler, 2006; Mansfield & Hagler, 2016). These advances in technology have allowed researchers to compute the incidence rates (i.e., the percentage of top predator individuals with mesopredator detected in the gut contents) of IGP among predator individuals. Nonetheless, a high incidence of IGP does not necessarily imply a high degree of IGP (Raso et al., 2014). For example, it is possible that a high percentage of individuals in a top predator population feed on other predator species despite on average low consumption in the diet. In this case, the high incidence rates of IGP could be misleading and may result in incorrect inferences of high degree of IGP among predators in the field.

New research idea

In this study, I propose an experimental framework combining controlled feeding trials and stable isotope analysis of field samples to more accurately determine the degree of IGP in a three-species omnivorous food web. Specifically, I predict that the $\delta^{15}\text{N}$ of top predator individuals engaged in IGP will be higher than the $\delta^{15}\text{N}$ of individuals not engaged in IGP. Moreover, the more the top predator consumes the mesopredator in the diet, the higher the $\delta^{15}\text{N}$ of top predator would be, and the increase in $\delta^{15}\text{N}$ (i.e., trophic enrichment) can be used to determine the degree of IGP.

91 **How to tackle the question through the proposed new idea**

92 Consider a three-species omnivorous food web, in which a top predator and a
 93 mesopredator feed on a shared prey, while the top predator also feeds on the mesopredator (Fig.
 94 1a). Two sets of controlled feeding trials will be conducted. In the first trial, the top predator and
 95 the mesopredator will be fed the shared prey for two weeks to allow for the incorporation of
 96 isotopes into the tissues (Gratton & Forbes, 2006) (Fig. 1b). The purpose of the first feeding trial
 97 is to ensure that both predators have reached an isotopic equilibrium state with the shared prey.
 98 In the second trial, the top predator will be fed mixed diets with different proportions of shared
 99 prey and mesopredator individuals (the organisms are from the first feeding trial): (1) shared
 100 prey only, (2) 75% of shared prey + 25% of mesopredator, (3) 50% of shared prey + 50% of
 101 mesopredator, (4) 25% of shared prey + 75% of mesopredator, and (5) mesopredator only (Fig.
 102 1c). The exact numbers of shared prey and mesopredator individuals used in each diet treatment
 103 will be determined based on their field densities. The purpose of the second feeding trial is to
 104 simulate different degrees of IGP under different encounter rates among the organisms.

105 After two weeks of feeding, the $\delta^{15}\text{N}$ of top predator individuals in each diet treatment
 106 will be analyzed, and their TEFs (relative to the $\delta^{15}\text{N}$ of shared prey) are used to construct a
 107 standard IGP curve (Fig. 1d). Finally, field samples of top predator and shared prey individuals
 108 will be collected, with their $\delta^{15}\text{N}$ analyzed to obtain the empirical TEF for the top predator. The
 109 degree of IGP in the field can then be determined by comparing the empirical TEF to the
 110 standard curve (Fig. 1e).

111 The proposed experimental framework combines the strengths of previous approaches to
 112 studying IGP—the controlled feeding trials can yield accurate trophic enrichment factors to
 113 construct a standard curve, whereas the stable isotope analysis of field samples allows for trophic
 114 interactions under natural settings. Therefore, this framework provides a useful tool for
 115 determining the degree of IGP in the field in a more quantitative and realistic fashion.

116 Agricultural systems are ideal for testing the proposed framework. IGP has been
 117 frequently documented among predators in such systems (Rosenheim et al., 1995). In addition,
 118 the species compositions are relatively simple compared to natural systems, which can largely
 119 reduce the potential confounding effects of other species on the trophic interactions among focal

organisms. If proven successful, this framework can be extended to food webs involving more complex interactions (e.g., cannibalism, multiple shared prey) and further complemented with other approaches (e.g., MGCA) to provide a more complete picture of the IGP dynamics in the field. Hopefully, it will provide an important piece of the puzzle in food web ecology.

Motivation

I have been using stable isotope analysis to quantify the diet compositions of generalist arthropod predators in rice agro-ecosystems in Taiwan. In my previous manuscript (published in the journal *Ecosphere*), a few reviewers expressed the concern over whether IGP would affect the diet compositions of predators. In fact, IGP may occur among the predators in our system, yet we were not able to quantify IGP due to the limitations of stable isotope mixing models. This question really puzzled me at that time and haunted my mind for long, which eventually brought me to the idea of using controlled feeding experiments along with stable isotope analysis to determine the degree of IGP in the field. I hope that this study can inspire new ideas and we will be able to develop a more thorough method to solve this question in the future.

Acknowledgements

I would like to thank Jia-Ang (William) Ou for his useful comments and editing work on this manuscript.

140 **References**

- 141 Abd El-Wakeil K. F. (2009). Trophic structure of macro-and meso-invertebrates in Japanese
142 coniferous forest: Carbon and nitrogen stable isotopes analyses. *Biochemical Systematics*
143 *and Ecology*, 37, 317-324. doi: <https://doi.org/10.1016/j.bse.2009.05.008>
- 144 Arim M., & Marquet P. A. (2004). Intraguild predation: a widespread interaction related to
145 species biology. *Ecol Lett*, 7, 557-564. doi: [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2004.00613.x)
146 0248.2004.00613.x
- 147 Denno R. F., Mitter M. S., Langellotto G. A., Gratton C., & Finke D. L. (2004). Interactions
148 between a hunting spider and a web-builder: consequences of intraguild predation and
149 cannibalism for prey suppression. *Ecological entomology*, 29, 566-577. doi:
150 <https://doi.org/10.1111/j.0307-6946.2004.00628.x>
- 151 Gagnon A. E., Heimpel G. E., & Brodeur J. (2011). The Ubiquity of Intraguild Predation among
152 Predatory Arthropods. *PLoS One*, 6, 7. doi: <https://doi.org/10.1371/journal.pone.0028061>
- 153 Gratton C., & Forbes A. E. (2006). Changes in $\delta^{13}\text{C}$ stable isotopes in multiple tissues of insect
154 predators fed isotopically distinct prey. *Oecologia*, 147, 615-624. doi:
155 <https://doi.org/10.1007/s00442-005-0322-y>
- 156 Hagler J. (2006). Development of an immunological technique for identifying multiple predator–
157 prey interactions in a complex arthropod assemblage. *Annals of Applied Biology*, 149,
158 153-165. doi: <https://doi.org/10.1111/j.1744-7348.2006.00076.x>
- 159 Halaj J., Peck R. W., & Niwa C. G. (2005). Trophic structure of a macroarthropod litter food
160 web in managed coniferous forest stands: a stable isotope analysis with $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$.
161 *Pedobiologia*, 49, 109-118. doi: <https://doi.org/10.1016/j.pedobi.2004.09.002>
- 162 Klarner B., Maraun M., & Scheu S. (2013). Trophic diversity and niche partitioning in a species
163 rich predator guild–Natural variations in stable isotope ratios ($^{13}\text{C}/^{12}\text{C}$, $^{15}\text{N}/^{14}\text{N}$) of
164 mesostigmatid mites (Acari, Mesostigmata) from Central European beech forests. *Soil*
165 *Biology and Biochemistry*, 57, 327-333. doi: <https://doi.org/10.1016/j.soilbio.2012.08.013>

- 166 Mansfield S., & Hagler J. R. (2016). Wanted dead or alive: scavenging versus predation by three
167 insect predators. *Food Webs*, 9, 12-17. doi: <https://doi.org/10.1016/j.fooweb.2016.03.003>
- 168 Muller C. B., & Brodeur J. (2002). Intraguild predation in biological control and conservation
169 biology. *Biol Control*, 25, 216-223. doi: [https://doi.org/10.1016/s1049-9644\(02\)00102-0](https://doi.org/10.1016/s1049-9644(02)00102-0)
- 170 Nakazawa T., & Yamamura N. (2006). Community structure and stability analysis for intraguild
171 interactions among host, parasitoid, and predator. *Population Ecology*, 48, 139-149. doi:
172 <https://doi.org/10.1007/s10144-005-0249-5>
- 173 Pahl K. B., Yurkowski D. J., Lees K. J., & Hussey N. E. (2020). Measuring the occurrence and
174 strength of intraguild predation in modern food webs. *Food Webs*, e00165. doi:
175 <https://doi.org/10.1016/j.fooweb.2020.e00165>
- 176 Polis G. A., & Holt R. D. (1992). Intraguild predation: the dynamics of complex trophic
177 interactions. *Trends in ecology & evolution*, 7, 151-154. doi:
178 [https://doi.org/10.1016/0169-5347\(92\)90208-S](https://doi.org/10.1016/0169-5347(92)90208-S)
- 179 Polis G. A., Myers C. A., & Holt R. D. (1989). The ecology and evolution of intraguild predation:
180 potential competitors that eat each other. *Annual review of ecology and systematics*, 20,
181 297-330. doi: <https://doi.org/10.1146/annurev.es.20.110189.001501>
- 182 Ponsard S., & Arditì R. (2000). What can stable isotopes ($\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) tell about the food
183 web of soil macro-invertebrates? *Ecology*, 81, 852-864. doi:
184 [https://doi.org/10.1890/0012-9658\(2000\)081\[0852:WCSINA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[0852:WCSINA]2.0.CO;2)
- 185 Provost C., Coderre D., Lucas E., Chouinard G., & Bostanian N. J. (2005). Impact of intraguild
186 predation and lambda-cyhalothrin on predation efficacy of three acarophagous predators.
187 *Pest Management Science: formerly Pesticide Science*, 61, 532-538. doi:
188 <https://doi.org/10.1002/ps.1027>
- 189 Raso L., Sint D., Mayer R., Plangg S., Recheis T., Brunner S., Kaufmann R., & Traugott M.
190 (2014). Intraguild predation in pioneer predator communities of alpine glacier forelands.
191 *Molecular ecology*, 23, 3744-3754. doi: <https://doi.org/10.1111/mec.12649>

192 Rickers S., Langel R., & Scheu S. (2006). Stable isotope analyses document intraguild predation
193 in wolf spiders (Araneae: Lycosidae) and underline beneficial effects of alternative prey
194 and microhabitat structure on intraguild prey survival. *Oikos*, 114, 471-478. doi:
195 <https://doi.org/10.1111/j.2006.0030-1299.14421.x>

196 Rosenheim J. A., Kaya H. K., Ehler L. E., Marois J. J., & Jaffee B. A. (1995). Intraguild
197 predation among biological-control agents: theory and evidence. *Biol Control*, 5, 303-335.
198 doi: <https://doi.org/10.1006/bcon.1995.1038>

199 Svanbäck R., Quevedo M., Olsson J., & Eklöv P. (2015). Individuals in food webs: the
200 relationships between trophic position, omnivory and among-individual diet variation.
201 *Oecologia*, 178, 103-114. doi: <https://doi.org/10.1007/s00442-014-3203-4>

202

203 **Figure legend**

204 **Figure 1.** Schematic diagram of the proposed experimental framework for determining the
205 degree of intraguild predation in a three-species omnivorous food web, in which a top predator
206 and a mesopredator feed on a shared prey, while the top predator also feeds on the mesopredator
207 (a). In the first feeding trial (b), the top predator and the mesopredator are fed the shared prey for
208 two weeks to ensure that both predators have reached an isotopic equilibrium state with the
209 shared prey. In the second feeding trial (c), the top predator is fed mixed diets with different
210 proportions of shared prey (from the stock) and mesopredator (from the first feeding trial)
211 individuals to simulate different degrees of IGP. (d) The trophic enrichment factors ($\Delta^{15}\text{N}$) of top
212 predator individuals (relative to the shared prey) in each diet treatment are used to construct a
213 standard IGP curve. Note that the curve may not necessarily be linear due to complex isotope
214 routing. (e) Field samples of top predator and shared prey individuals are collected, and their
215 $\delta^{15}\text{N}$ values are analyzed to obtain the empirical $\Delta^{15}\text{N}$ for the top predator. The degree of IGP in
216 the field can then be determined by comparing the empirical $\Delta^{15}\text{N}$ to the standard curve.




Figure 1

253x190mm (300 x 300 DPI)