List of definite integrals

In mathematics, the definite integral:

$$\int_a^b f(x)\,dx$$

is the area of the region in the xy-plane bounded by the graph of f, the x-axis, and the lines x = a and x = b, such that area above the x-axis adds to the total, and that below the x-axis subtracts from the total.

The <u>fundamental theorem of calculus</u> establishes the relationship between indefinite and definite integrals and introduces a technique for evaluating definite integrals.

If the interval is infinite the definite integral is called an improper integral and defined by using appropriate limiting procedures. for example:

$$\int_a^\infty f(x)\,dx = \lim_{b o\infty} \left[\int_a^b f(x)\,dx
ight]$$

A constant, such pi, that may be defined by the integral of an algebraic function over an algebraic domain is known as a period.

The following is a list of the most common definite Integrals. For a list of indefinite integrals see List of indefinite integrals

Contents

Definite integrals involving rational or irrational expressions

Definite integrals involving trigonometric functions

Definite integrals involving exponential functions

Definite integrals involving logarithmic functions

Definite integrals involving hyperbolic functions

Frullani integrals

See also

References

Definite integrals involving rational or irrational expressions

$$\int_{0}^{\infty} rac{dx}{x^{2} + a^{2}} = rac{\pi}{2a}$$
 $\int_{0}^{\infty} rac{x^{m}dx}{x^{n} + a^{n}} = rac{\pi a^{m-n+1}}{n \sin\left(rac{m+1}{n}\pi
ight)} \quad ext{for } 0 < m+1 < n$
 $\int_{0}^{\infty} rac{x^{p-1}dx}{1+x} = rac{\pi}{\sin(p\pi)} \quad ext{for } 0
 $\int_{0}^{\infty} rac{x^{m}dx}{1+2x\cos\beta+x^{2}} = rac{\pi}{\sin(m\pi)} \cdot rac{\sin(m\beta)}{\sin(\beta)}$
 $\int_{0}^{a} rac{dx}{\sqrt{a^{2}-x^{2}}} = rac{\pi}{2}$
 $\int_{0}^{a} \sqrt{a^{2}-x^{2}} dx = rac{\pi a^{2}}{4}$
 $\int_{0}^{a} x^{m}(a^{n}-x^{n})^{p} dx = rac{a^{m+1+np}\Gamma\left(rac{m+1}{n}
ight)\Gamma(p+1)}{n\Gamma\left(rac{m+1}{n}+p+1
ight)}$$

$$\int_0^\infty rac{x^m dx}{\left(x^n+a^n
ight)^r} = rac{(-1)^{r-1}\pi a^{m+1-nr}\Gamma\left(rac{m+1}{n}
ight)}{n\sin\Bigl(rac{m+1}{n}\pi\Bigr)(r-1)!\,\Gamma\left(rac{m+1}{n}-r+1
ight)} \quad ext{for } n(r-2) < m+1 < nr$$

Definite integrals involving trigonometric functions

$$\int_{0}^{\pi} \sin(mx) \sin(nx) dx = \begin{cases} 0 & \text{if } m \neq n \\ \frac{\pi}{2} & \text{if } m = n \end{cases} & \text{for } m, n \text{ positive integers} \\ \frac{\pi}{2} & \text{if } m = n \end{cases} & \text{for } m, n \text{ positive integers} \\ \int_{0}^{\pi} \cos(mx) \cos(nx) dx = \begin{cases} 0 & \text{if } m \neq n \\ \frac{\pi}{2} & \text{if } m = n \end{cases} & \text{for } m, n \text{ positive integers} \end{cases} \\ \int_{0}^{\pi} \sin(mx) \cos(nx) dx = \begin{cases} 0 & \text{if } m + n \text{ even} \\ \frac{2m}{m^2 - n^2} & \text{if } m + n \text{ odd} \end{cases} \\ \int_{0}^{\frac{\pi}{2}} \sin^2(x) dx = \int_{0}^{\frac{\pi}{2}} \cos^{2m}(x) dx = \frac{\pi}{4} \\ \int_{0}^{\frac{\pi}{2}} \sin^{2m}(x) dx = \int_{0}^{\frac{\pi}{2}} \cos^{2m}(x) dx = \frac{1 \times 3 \times 5 \times \dots \times (2m - 1)}{2 \times 4 \times 6 \times \dots \times 2m} \cdot \frac{\pi}{2} & \text{for } m = 1, 2, 3 \dots \end{cases} \\ \int_{0}^{\frac{\pi}{2}} \sin^{2m+1}(x) dx = \int_{0}^{\frac{\pi}{2}} \cos^{2m+1}(x) dx = \frac{2 \times 4 \times 6 \times \dots \times 2m}{1 \times 3 \times 5 \times \dots \times (2m + 1)} & \text{for } m = 1, 2, 3 \dots \end{cases} \\ \int_{0}^{\frac{\pi}{2}} \sin^{2p-1}(x) \cos^{2p-1}(x) dx = \frac{\Gamma(p)\Gamma(q)}{2\Gamma(p+q)} = \frac{1}{2}B(p,q) \\ \int_{0}^{\infty} \frac{\sin(px)}{x} dx = \begin{cases} \frac{\pi}{2} & \text{if } p > 0 \\ 0 & \text{if } p = 0 \text{ (see Dirichlet integral)} \\ -\frac{\pi}{2} & \text{if } 0 < q < p \\ \frac{\pi}{4} & \text{if } p = q > 0 \end{cases} \\ \int_{0}^{\infty} \frac{\sin px \sin qx}{x^2} dx = \begin{cases} \frac{\pi p}{2} & \text{if } 0$$

$$\int_{0}^{\infty} \frac{\cos px - \cos qx}{x^{2}} dx = \frac{\pi(q - p)}{2}$$

$$\int_{0}^{\infty} \frac{\cos mx}{x^{2} + a^{2}} dx = \frac{\pi}{2a} e^{-ma}$$

$$\int_{0}^{\infty} \frac{x \sin mx}{x^{2} + a^{2}} dx = \frac{\pi}{2} e^{-ma}$$

$$\int_{0}^{2\pi} \frac{\sin mx}{a + b \sin x} dx = \frac{\pi}{2a^{2}} (1 - e^{-ma})$$

$$\int_{0}^{2\pi} \frac{dx}{a + b \cos x} = \frac{2\pi}{\sqrt{a^{2} - b^{2}}}$$

$$\int_{0}^{\pi} \frac{dx}{a + b \cos x} = \frac{\cos^{-1} \left(\frac{b}{a}\right)}{\sqrt{a^{2} - b^{2}}}$$

$$\int_{0}^{2\pi} \frac{dx}{(a + b \sin x)^{2}} = \int_{0}^{2\pi} \frac{dx}{(a + b \cos x)^{2}} = \frac{2\pi a}{(a^{2} - b^{2})^{3/2}}$$

$$\int_{0}^{2\pi} \frac{dx}{1 - 2a \cos x + a^{2}} = \frac{2\pi}{1 - a^{2}} \text{ for } 0 < a < 1$$

$$\int_{0}^{\pi} \frac{x \sin x dx}{1 - 2a \cos x + a^{2}} = \frac{\pi}{1 - a^{2}} \text{ for } a^{2} < 1, m = 0, 1, 2, \dots$$

$$\int_{0}^{\infty} \sin ax^{2} dx = \int_{0}^{\infty} \cos ax^{2} = \frac{1}{2} \sqrt{\frac{\pi}{2a}}$$

$$\int_{0}^{\infty} \sin ax^{2} dx = \int_{0}^{\infty} \cos ax^{2} = \frac{1}{2} \sqrt{\frac{\pi}{2a}}$$

$$\int_{0}^{\infty} \sin ax^{n} = \frac{1}{na^{1/n}} \Gamma\left(\frac{1}{n}\right) \sin \frac{\pi}{2n} \text{ for } n > 1$$

$$\int_{0}^{\infty} \cos ax^{n} = \frac{1}{na^{1/n}} \Gamma\left(\frac{1}{n}\right) \cos \frac{\pi}{2n} \text{ for } n > 1$$

$$\int_{0}^{\infty} \frac{\sin x}{x^{p}} dx = \int_{0}^{\infty} \frac{\cos x}{\sqrt{x}} dx = \sqrt{\frac{\pi}{2}}$$

$$\int_{0}^{\infty} \frac{\sin x}{x^{p}} dx = \frac{\pi}{2\Gamma(p) \sin\left(\frac{p\pi}{2}\right)} \text{ for } 0
$$\int_{0}^{\infty} \sin ax^{2} \cos 2bx dx = \frac{1}{2} \sqrt{\frac{\pi}{2a}} \left(\cos \frac{b^{2}}{a} - \sin \frac{b^{2}}{a}\right)$$

$$\int_{0}^{\infty} \sin ax^{2} \cos 2bx dx = \frac{1}{2} \sqrt{\frac{\pi}{2a}} \left(\cos \frac{b^{2}}{a} - \sin \frac{b^{2}}{a}\right)$$$$

Definite integrals involving exponential functions

$$\int_{0}^{\infty} \sqrt{x} \, e^{-s} \, dx = \frac{1}{2} \sqrt{\pi} \, (\text{see also Gamma function})$$

$$\int_{0}^{\infty} e^{-ax} \cos bx \, dx = \frac{a}{a^2 + b^2}$$

$$\int_{0}^{\infty} e^{-ax} \sin bx \, dx = \tan^{-1} \frac{b}{a}$$

$$\int_{0}^{\infty} \frac{e^{-ax} \sin bx}{x} \, dx = \tan^{-1} \frac{b}{a}$$

$$\int_{0}^{\infty} e^{-ax^2} \, dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} \, \text{ for } a > 0 \, (\text{the Gaussian integral})$$

$$\int_{0}^{\infty} e^{-ax^2} \cos bx \, dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} \, e^{\left(\frac{-a^2}{4a}\right)}$$

$$\int_{0}^{\infty} e^{-(ax^2 + bx + c)} \, dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} \, e^{\left(\frac{a^2 - bx}{4a}\right)} \cdot \text{erfc} \, \frac{b}{2\sqrt{a}}, \text{ where } \text{erfc}(p) = \frac{2}{\sqrt{\pi}} \int_{p}^{\infty} e^{-x^2} \, dx$$

$$\int_{-\infty}^{\infty} e^{-(ax^2 + bx + c)} \, dx = \sqrt{\frac{\pi}{a}} \, e^{\left(\frac{p^2 - bx}{4a}\right)} \cdot \text{erfc} \, \frac{b}{2\sqrt{a}}, \text{ where } \text{erfc}(p) = \frac{2}{\sqrt{\pi}} \int_{p}^{\infty} e^{-x^2} \, dx$$

$$\int_{-\infty}^{\infty} e^{-(ax^2 + bx + c)} \, dx = \sqrt{\frac{\pi}{a}} \, e^{\left(\frac{p^2 - bx}{4a}\right)}$$

$$\int_{0}^{\infty} x^n e^{-ax} \, dx = \frac{\Gamma(n+1)}{a^{n+1}}$$

$$\int_{0}^{\infty} x^2 e^{-ax^2} \, dx = \frac{1}{4} \sqrt{\frac{\pi}{a^2}} \, \text{ for } a > 0$$

$$\int_{0}^{\infty} x^2 e^{-ax^2} \, dx = \frac{1}{2a^2} \int_{0}^{\infty} x^2 e^{-ax^2} \, dx = \frac{(2n-1)!!}{2a^2} \sqrt{\frac{\pi}{a^{2n+1}}} \, \text{ for } a > 0, \, n-1, 2, 3 \dots$$

$$\int_{0}^{\infty} x^2 e^{-ax^2} \, dx = \frac{1}{2a^2} \, \text{ for } a > 0$$

$$\int_{0}^{\infty} x^2 e^{-ax^2} \, dx = \frac{n}{2a^2} \int_{0}^{\infty} x^{2n-1} e^{-ax^2} \, dx = \frac{n!}{2a^{n+1}} \, \text{ for } a > 0, \, n = 0, 1, 2 \dots$$

$$\int_{0}^{\infty} x^n e^{-ax^2} \, dx = \frac{n}{2a^2} \int_{0}^{\infty} x^{2n-1} e^{-ax^2} \, dx = \frac{n!}{2a^{n+1}} \, \text{ for } a > 0, \, n = 0, 1, 2 \dots$$

$$\int_{0}^{\infty} x^n e^{-ax^2} \, dx = \frac{1}{2\sqrt{\frac{n}{a}}} e^{-2\sqrt{ab}}$$

$$\int_{0}^{\infty} e^{\left(-ax^2 - \frac{1}{x^2}\right)} \, dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} e^{-2\sqrt{ab}}$$

$$\int_{0}^{\infty} e^{-(-ax^2 - \frac{1}{x^2})} \, dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} e^{-2\sqrt{ab}}$$

$$\int_{0}^{\infty} \frac{x^{n-1}}{e^x - 1} \, dx = \Gamma(n) \zeta(n)$$

$$\int_{0}^{\infty} \frac{x^{n-1}}{e^x - 1} \, dx = \frac{1}{3^2} - \frac{1}{4^2} + \frac{1}{a^2} - \frac{1}{4^2} + \dots = \frac{\pi^2}{12}$$

$$\int_{0}^{\infty} \frac{\sin mx}{e^{2\pi x} - 1} \, dx = \frac{1}{4} \coth \frac{m}{2} - \frac{1}{2m}$$

$$\int_{0}^{\infty} \left(\frac{1}{1 + x} - e^{-x} \right) \, \frac{dx}{x} = \gamma \text{ (where } \gamma \text{ is } \underline{\text{Euler-Mascheroni constant)}}$$

$$\int_{0}^{\infty} \frac{e^{-x^{2}} - e^{-x}}{x} \, dx = \frac{\gamma}{2}$$

$$\int_{0}^{\infty} \left(\frac{1}{e^{x} - 1} - \frac{e^{-x}}{x} \right) \, dx = \gamma$$

$$\int_{0}^{\infty} \frac{e^{-ax} - e^{-bx}}{x \sec px} \, dx = \frac{1}{2} \ln \frac{b^{2} + p^{2}}{a^{2} + p^{2}}$$

$$\int_{0}^{\infty} \frac{e^{-ax} - e^{-bx}}{x \csc px} \, dx = \tan^{-1} \frac{b}{p} - \tan^{-1} \frac{a}{p}$$

$$\int_{0}^{\infty} \frac{e^{-ax} (1 - \cos x)}{x^{2}} \, dx = \cot^{-1} a - \frac{a}{2} \ln \left| \frac{a^{2} + 1}{a^{2}} \right|$$

$$\int_{-\infty}^{\infty} e^{-x^{2}} \, dx = \sqrt{\pi}$$

$$\int_{-\infty}^{\infty} e^{-x^{2}} \, dx = \sqrt{\pi}$$

$$\int_{-\infty}^{\infty} x^{2(n+1)} e^{-\frac{1}{2}x^{2}} \, dx = \frac{(2n+1)!}{2^{n}n!} \sqrt{2\pi} \quad \text{for } n = 0, 1, 2, \dots$$

Definite integrals involving logarithmic functions

$$\int_{0}^{1} x^{m} (\ln x)^{n} dx = \frac{(-1)^{n} n!}{(m+1)^{n+1}} \quad \text{for } m > -1, n = 0, 1, 2, \dots$$

$$\int_{0}^{1} \frac{\ln x}{1+x} dx = -\frac{\pi^{2}}{12}$$

$$\int_{0}^{1} \frac{\ln x}{1-x} dx = -\frac{\pi^{2}}{6}$$

$$\int_{0}^{1} \frac{\ln(1+x)}{x} dx = \frac{\pi^{2}}{12}$$

$$\int_{0}^{1} \frac{\ln(1-x)}{x} dx = -\frac{\pi^{2}}{6}$$

$$\int_{0}^{\infty} \frac{\ln(a^{2} + x^{2})}{b^{2} + x^{2}} dx = \frac{\pi}{b} \ln(a+b) \quad \text{for } a, b > 0$$

$$\int_{0}^{\infty} \frac{\ln x}{x^{2} + a^{2}} dx = \frac{\pi \ln a}{2a} \quad \text{for } a > 0$$

Definite integrals involving hyperbolic functions

$$\int_0^\infty \frac{\sin ax}{\sinh bx} dx = \frac{\pi}{2b} \tanh \frac{a\pi}{2b}$$

$$\int_0^\infty \frac{\cos ax}{\cosh bx} dx = \frac{\pi}{2b} \cdot \frac{1}{\cosh \frac{a\pi}{2b}}$$

$$\int_0^\infty \frac{x}{\sinh ax} dx = \frac{\pi^2}{4a^2}$$

$$\int_{-\infty}^\infty \frac{1}{\cosh x} dx = \pi$$

Frullani integrals

$$\int_0^\infty \frac{f(ax) - f(bx)}{x} \ dx = \left(\lim_{x \to 0} f(x) - \lim_{x \to \infty} f(x)\right) \ln\left(\frac{b}{a}\right) \text{ holds if the integral exists and } f'(x) \text{ is continuous.}$$

See also

- List of integrals
- Indefinite sum
- Gamma function
- List of limits

References

- Spiegel, Murray R.; Lipschutz, Seymour; Liu, John (2009). Mathematical handbook of formulas and tables (3rd ed.). McGraw-Hill. ISBN 978-0071548557.
- Zwillinger, Daniel (2003). CRC standard mathematical tables and formulae (32nd ed.). CRC Press. ISBN 978-143983548-7.
- Abramowitz, Milton; Stegun, Irene Ann, eds. (1983) [June 1964]. Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables. Applied Mathematics Series. 55 (Ninth reprint with additional corrections of tenth original printing with corrections (December 1972); first ed.). Washington D.C.; New York: United States Department of Commerce, National Bureau of Standards; Dover Publications. ISBN 978-0-486-61272-0. LCCN 64-60036 (https://lccn.loc.gov/64-60036). MR 0167642 (https://www.ams.org/mathscinet-getitem?mr=0167642). LCCN 65-12253 (https://lccn.loc.gov/65012253).

Retrieved from "https://en.wikipedia.org/w/index.php?title=List_of_definite_integrals&oldid=862067044"

This page was last edited on 1 October 2018, at 23:21 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the <u>Terms of Use</u> and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.