

# Tutoriat 9 CMI

## Functiile Beta si Gamma

$$\Gamma: (0, \infty) \rightarrow \mathbb{R}, \quad \Gamma(p) = \int_0^{+\infty} x^{p-1} e^{-x} dx, \quad \forall p \in (0, \infty)$$

$$B: (0, \infty) \times (0, \infty), \quad B(p, q) = \int_0^{1-0} x^{p-1} (1-x)^{q-1} dx, \quad \forall p, q \in (0, \infty)$$

$$B(p, q) = 2 \int_0^{\frac{\pi}{2}-0} (\sin x)^{2p-1} (\cos x)^{2q-1} dx$$

$$B(p, q) = \int_0^{+\infty} \frac{x^{p-1}}{(1+x)^{p+q}} dx$$

$$\Gamma(1) = 1$$

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

$$\Gamma(p+1) = p \Gamma(p), \quad \forall p > 0$$

$$\Gamma(n+1) = n!, \quad \forall n \in \mathbb{N}$$

$$\Gamma(p) \cdot \Gamma(1-p) = \frac{\pi}{\sin(p\pi)}, \quad \forall p \in (0, 1)$$

$$\Gamma(p) \cdot \Gamma\left(p + \frac{1}{2}\right) = \frac{\sqrt{\pi}}{2^{2p-1}} \cdot \Gamma(2p), \quad \forall p > 0$$

$$B(p, q) = \frac{\Gamma(p) \cdot \Gamma(q)}{\Gamma(p+q)}, \quad \forall p, q \in (0, \infty)$$

$$B(p, q) = B(q, p), \quad \forall p, q \in (0, \infty)$$

$\Gamma \rightarrow$  generalizarea a notii de factorial

$$\textcircled{+} \quad \Gamma(2p) = \pi^{-\frac{1}{2}} 2^{2p-1} \Gamma(p) \Gamma\left(p + \frac{1}{2}\right)$$

$$\lim_{\substack{x \rightarrow 0 \\ x > 0}} x \Gamma(x) = 1 \quad (! \text{ nu e m\u00e2nginit\u0103})$$

$$\lim_{x \rightarrow \infty} \frac{\Gamma(x+a)}{x^a \Gamma(x)} = 1, \quad \forall a \in (0, \infty)$$

$$\Gamma \begin{cases} \subset C^\infty \\ \text{convex\u0103} \end{cases}$$