

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}, f(x, y) = \begin{cases} \frac{x^2 y^2}{\sqrt{x^4 + y^{10}}}, & x^2 + y^2 \neq 0 \\ 0, & x = y = 0 \end{cases}$$

f functia este continua pe \mathbb{R}^2 (vezi tut 4)

$$\frac{\partial f}{\partial x}(x, y) = \frac{2xy^2 \sqrt{x^4 + y^{10}} - x^2 y^2 \frac{1}{2\sqrt{x^4 + y^{10}}} 4x^3}{x^4 + y^{10}} = 2x \frac{y^2(x^4 + y^{10}) - x^4 y^2}{(x^4 + y^{10}) \sqrt{x^4 + y^{10}}} = 2x \frac{y^{12}}{(x^4 + y^{10}) \sqrt{x^4 + y^{10}}}$$

$$\frac{\partial f}{\partial y}(x, y) = \frac{2x^2 y \sqrt{x^4 + y^{10}} - x^2 y^2 \frac{1}{2\sqrt{x^4 + y^{10}}} 10y^9}{x^4 + y^{10}} = y \frac{2x^2(x^4 + y^{10}) - 5x^2 y^{10}}{(x^4 + y^{10}) \sqrt{x^4 + y^{10}}} = y \frac{2x^6 - 3x^2 y^{10}}{(x^4 + y^{10}) \sqrt{x^4 + y^{10}}}$$

$$e_1 = (1, 0)$$

$$e_2 = (0, 1)$$

$$\lim_{t \rightarrow 0} \frac{f((0, 0) + t e_1) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, 0) + t(1, 0)) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((t, 0)) - f((0, 0))}{t} =$$

$$= \lim_{t \rightarrow 0} \frac{\frac{t^2 \cdot 0^2}{\sqrt{t^4 + 0^{10}}} - 0}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0 \Rightarrow \exists \frac{\partial f}{\partial x}(0, 0) = 0$$

$$\lim_{t \rightarrow 0} \frac{f((0, 0) + t e_2) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, 0) + t(0, 1)) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, t)) - f((0, 0))}{t} =$$

$$= \lim_{t \rightarrow 0} \frac{\frac{0^2 \cdot t^2}{\sqrt{0^4 + t^{10}}} - 0}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0 \Rightarrow \exists \frac{\partial f}{\partial y}(0, 0) = 0$$

$$\exists \frac{\partial f}{\partial x}(x, y), \frac{\partial f}{\partial y}(x, y) \text{ pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$$\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \text{ functii continue pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$$\mathbb{R}^2 \setminus \{(0, 0)\} \text{ multime deschisa}$$

$$\Rightarrow f \text{ diferentiabila pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$$\text{Fie } T: \mathbb{R}^2 \rightarrow \mathbb{R}, T((x, y)) = \frac{\partial f}{\partial x}(0, 0)x + \frac{\partial f}{\partial y}(0, 0)y = 0x + 0y = 0, \forall (x, y) \in \mathbb{R}^2$$

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{|f(x, y) - f(0, 0) - T((x, y) - (0, 0))|}{\|(x, y) - (0, 0)\|} = \lim_{(x, y) \rightarrow (0, 0)} \frac{\left| \frac{x^2 y^2}{\sqrt{x^4 + y^{10}}} - 0 - T((x, y)) \right|}{\|(x, y)\|} =$$

$$= \lim_{(x, y) \rightarrow (0, 0)} \frac{\left| \frac{x^2 y^2}{\sqrt{x^4 + y^{10}}} - 0 \right|}{\sqrt{x^2 + y^2}} = \lim_{(x, y) \rightarrow (0, 0)} \frac{x^2 y^2}{\underbrace{\sqrt{x^4 + y^{10}} \sqrt{x^2 + y^2}}_{g(x, y)}}$$

$$0 \leq g(x, y) = \frac{x^2 y^2}{\sqrt{x^4 + y^{10}} \sqrt{x^2 + y^2}} \leq \frac{x^2 y^2}{\sqrt{x^4} \sqrt{x^2 + y^2}} = \frac{y^2}{\sqrt{x^2 + y^2}} \leq \frac{y^2}{\sqrt{y^2}} = y$$

$$\begin{array}{c} 0 \leq g(x, y) \leq y \\ \swarrow \quad \searrow \\ \quad \downarrow \quad (x, y) \rightarrow (0, 0) \\ \quad \quad \quad 0 \end{array}$$

$$\Rightarrow \lim_{(x, y) \rightarrow (0, 0)} \frac{|f(x, y) - f(0, 0) - T((x, y) - (0, 0))|}{\|(x, y) - (0, 0)\|} = 0 \Rightarrow f \text{ e diferentiabila in } (0, 0)$$

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}, f(x, y) = \begin{cases} \frac{x^5 y^2}{x^8 + y^4}, x^2 + y^2 \neq 0 \\ 0, x = y = 0 \end{cases}$$

$$\lim_{(x,y) \rightarrow (0,0)} f(x, y) = \lim_{(x,y) \rightarrow (0,0)} \frac{x^5 y^2}{x^8 + y^4} \stackrel{?}{=} 0$$

Alegem:

$$y = x \Rightarrow \lim_{x \rightarrow 0} f(x, x) = \frac{x^5 x^2}{x^8 + x^4} = \lim_{x \rightarrow 0} \frac{x^3}{x^4 + 1} = 0$$

$$y = -x \Rightarrow \lim_{x \rightarrow 0} f(x, -x) = \frac{x^5 x^2}{x^8 + x^4} = \lim_{x \rightarrow 0} \frac{x^3}{x^4 + 1} = 0$$

$$y = \sqrt{x} \Rightarrow \lim_{x \rightarrow 0} f(x, \sqrt{x}) = \frac{x^5 x}{x^8 + x^2} = \lim_{x \rightarrow 0} \frac{x^4}{x^6 + 1} = 0$$

Este posibil ca:

$$\lim_{(x,y) \rightarrow (0,0)} f(x, y) = l (= 0) \in \mathbb{R}$$

Evaluam:

$$0 \leq |f(x, y) - l| \leq g(x, y)$$

$(x, y) \rightarrow (0, 0)$

0

$$0 \leq |f(x, y) - l| = \left| \frac{x^5 y^2}{x^8 + y^4} \right| = |x| \frac{x^4 y^2}{x^8 + y^4} \leq |x| \frac{1}{2}$$

$$0 \leq |f(x, y) - l| \leq \frac{|x|}{2}$$

$(x, y) \rightarrow (0, 0)$

0

$$\frac{\partial f}{\partial x}(x, y) = \frac{5x^4 y^2 (x^8 + y^4) - 8x^{12} y^2}{(x^8 + y^4)^2} = x^4 y^2 \frac{5y^4 - 3x^8}{(x^8 + y^4)^2}$$

$$\frac{\partial f}{\partial y}(x, y) = \frac{2x^5 y (x^8 + y^4) - 4x^5 y^5}{(x^8 + y^4)^2} = 2x^5 y \frac{x^8 - y^4}{(x^8 + y^4)^2}$$

$$e_1 = (1, 0)$$

$$e_2 = (0, 1)$$

$$\lim_{t \rightarrow 0} \frac{f((0, 0) + t e_1) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, 0) + t(1, 0)) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((t, 0)) - f((0, 0))}{t} =$$

$$= \lim_{t \rightarrow 0} \frac{\frac{t^5 \cdot 0^2}{\sqrt{t^8 + 0^4}} - 0}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0 \Rightarrow \exists \frac{\partial f}{\partial x}(0, 0) = 0$$

$$\lim_{t \rightarrow 0} \frac{f((0, 0) + t e_2) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, 0) + t(0, 1)) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, t)) - f((0, 0))}{t} =$$

$$= \lim_{t \rightarrow 0} \frac{\frac{0^5 \cdot t^2}{\sqrt{0^8 + t^4}} - 0}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0 \Rightarrow \exists \frac{\partial f}{\partial y}(0, 0) = 0$$

$$\exists \frac{\partial f}{\partial x}(x, y), \frac{\partial f}{\partial y}(x, y) \text{ pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$$\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \text{ functii continue pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$$\mathbb{R}^2 \setminus \{(0, 0)\} \text{ multime deschisa}$$

$$\Rightarrow f \text{ diferentiabila pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$$\text{Fie } T : \mathbb{R}^2 \rightarrow \mathbb{R}, T((x, y)) = \frac{\partial f}{\partial x}(0, 0)x + \frac{\partial f}{\partial y}(0, 0)y = 0x + 0y = 0, \forall (x, y) \in \mathbb{R}^2$$

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{|f(x, y) - f(0, 0) - T((x, y) - (0, 0))|}{\|(x, y) - (0, 0)\|} = \lim_{(x, y) \rightarrow (0, 0)} \frac{\left| \frac{x^5 y^2}{x^8 + y^4} - 0 - T((x, y)) \right|}{\|(x, y)\|} =$$

$$= \lim_{(x, y) \rightarrow (0, 0)} \frac{\left| \frac{x^5 y^2}{x^8 + y^4} \right| - 0}{\sqrt{x^2 + y^2}} = \lim_{(x, y) \rightarrow (0, 0)} \frac{x^5 y^2}{\underbrace{(x^8 + y^4) \sqrt{x^2 + y^2}}_{g(x, y)}}$$

$$y = x^2 \Rightarrow \lim_{x \rightarrow 0} \frac{x^5 x^4}{(x^8 + x^8) \sqrt{x^2 + x^4}} = \lim_{x \rightarrow 0} \frac{x^9}{2x^9 \sqrt{1 + x^2}} = \frac{1}{2} \neq 0$$

$$\Rightarrow \lim_{(x, y) \rightarrow (0, 0)} \frac{|f(x, y) - f(0, 0) - T((x, y) - (0, 0))|}{\|(x, y) - (0, 0)\|} \neq 0 \Rightarrow f \text{ nu e diferentiabila in } (0, 0)$$

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, f(x, y) = \begin{cases} \frac{x^3 y^7}{\sqrt{x^{16} + y^{16}}}, & x^2 + y^2 \neq 0 \\ 0, & x = y = 0 \end{cases}$$

f functia este continua pe \mathbb{R}^2 (vezi tut 4)

$$\frac{\partial f}{\partial x}(x, y) = \frac{3x^2 y^7 \sqrt{x^{16} + y^{16}} - x^3 y^7 \frac{1}{2\sqrt{x^{16} + y^{16}}} 16x^{15}}{x^{16} + y^{16}} = x^2 y^7 \frac{-5x^{16} + 3y^{16}}{(x^{16} + y^{16}) \sqrt{x^{16} + y^{16}}}$$

$$\frac{\partial f}{\partial y}(x, y) = \frac{7x^3 y^6 \sqrt{x^{16} + y^{16}} - x^3 y^7 \frac{1}{2\sqrt{x^{16} + y^{16}}} 16y^{15}}{x^{16} + y^{16}} = x^3 y^6 \frac{-x^{16} + 7y^{16}}{(x^{16} + y^{16}) \sqrt{x^{16} + y^{16}}}$$

$$e_1 = (1, 0)$$

$$e_2 = (0, 1)$$

$$\lim_{t \rightarrow 0} \frac{f((0, 0) + t e_1) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, 0) + t(1, 0)) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((t, 0)) - f((0, 0))}{t} =$$

$$= \lim_{t \rightarrow 0} \frac{\frac{t^3 \cdot 0^7}{\sqrt{t^{16} + 0^{16}}} - 0}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0 \Rightarrow \exists \frac{\partial f}{\partial x}(0, 0) = 0$$

$$\lim_{t \rightarrow 0} \frac{f((0, 0) + t e_2) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, 0) + t(0, 1)) - f((0, 0))}{t} = \lim_{t \rightarrow 0} \frac{f((0, t)) - f((0, 0))}{t} =$$

$$= \lim_{t \rightarrow 0} \frac{\frac{0^3 \cdot t^7}{\sqrt{0^{16} + t^{16}}} - 0}{t} = \lim_{t \rightarrow 0} \frac{0}{t} = 0 \Rightarrow \exists \frac{\partial f}{\partial y}(0, 0) = 0$$

$$\exists \frac{\partial f}{\partial x}(x, y), \frac{\partial f}{\partial y}(x, y) \text{ pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$$\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \text{ functii continue pe } \mathbb{R}^2 \setminus \{(0, 0)\}$$

$\mathbb{R}^2 \setminus \{(0, 0)\}$ multime deschisa

$\Rightarrow f$ diferentiabila pe $\mathbb{R}^2 \setminus \{(0, 0)\}$

$$\text{Fie } T : \mathbb{R}^2 \rightarrow \mathbb{R}, T((x, y)) = \frac{\partial f}{\partial x}(0, 0)x + \frac{\partial f}{\partial y}(0, 0)y = 0x + 0y = 0, \forall (x, y) \in \mathbb{R}^2$$

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{|f(x, y) - f(0, 0) - T((x, y) - (0, 0))|}{\|(x, y) - (0, 0)\|} = \lim_{(x, y) \rightarrow (0, 0)} \frac{\left| \frac{x^3 y^7}{\sqrt{x^{16} + y^{16}}} - 0 - T((x, y)) \right|}{\|(x, y)\|} =$$

$$= \lim_{(x,y) \rightarrow (0,0)} \frac{\left| \frac{x^3 y^7}{\sqrt{x^{16} + y^{16}}} - 0 \right|}{\sqrt{x^2 + y^2}} = \lim_{(x,y) \rightarrow (0,0)} \underbrace{\frac{|x^3 y^7|}{\sqrt{x^{16} + y^{16}} \sqrt{x^2 + y^2}}}_{g(x,y)}$$

$$x^{16} \leq x^{16} + y^{16} \Leftrightarrow \sqrt[16]{x^{16}} \leq \sqrt[16]{x^{16} + y^{16}} \Leftrightarrow |x| \leq \sqrt[16]{x^{16} + y^{16}} \Leftrightarrow |x|^3 \leq (x^{16} + y^{16})^{\frac{3}{16}}$$

Procedăm la fel și pentru $|y|^7 \leq (x^{16} + y^{16})^{\frac{7}{16}}$

$$\Rightarrow |x^3 y^7| \leq (x^{16} + y^{16})^{\frac{3}{16} + \frac{7}{16}} = (x^{16} + y^{16})^{\frac{5}{8}}$$

$$0 \leq g(x, y) = \frac{x^3 y^7}{\sqrt{x^{16} + y^{16}} \sqrt{x^2 + y^2}} \leq \frac{(x^{16} + y^{16})^{\frac{5}{8}}}{\sqrt{x^{16} + y^{16}} \sqrt{x^2 + y^2}} = \frac{(x^{16} + y^{16})^{\frac{5}{8} - \frac{1}{2}}}{\sqrt{x^2 + y^2}} = \frac{\sqrt[8]{x^{16} + y^{16}}}{\sqrt{x^2 + y^2}} \leq$$

$$\leq \frac{\sqrt[8]{(x^2 + y^2)^8}}{\sqrt{x^2 + y^2}} = \frac{x^2 + y^2}{\sqrt{x^2 + y^2}} = \sqrt{x^2 + y^2}$$

$$0 \leq g(x, y) \leq \sqrt{x^2 + y^2}$$

$(x, y) \rightarrow (0, 0)$

0

$$\Rightarrow \lim_{(x,y) \rightarrow (0,0)} \frac{|f(x, y) - f(0, 0) - T((x, y) - (0, 0))|}{\|(x, y) - (0, 0)\|} = 0 \Rightarrow f \text{ e diferentiabila in } (0, 0)$$