

$$\sum_{n=1}^{\infty} \frac{(-1)^n x^n}{n}$$

$$\begin{cases} x_0 = 0 \\ a_n = \frac{(-1)^n}{n}, n \in \mathbb{N}^* \\ a_0 = 0 \end{cases}$$

$$R = \frac{1}{\rho}$$

$$\rho = \overline{\lim} \sqrt[n]{|a_n|}, \text{ sau } \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} \text{ (daca } \exists \text{)}$$

$$\rho = \lim_{n \rightarrow \infty} \frac{\left| \frac{(-1)^{n+1}}{n+1} \right|}{\left| \frac{(-1)^n}{n} \right|} = \lim_{n \rightarrow \infty} \left| \frac{-1}{1 + \frac{1}{n}} \right| = 1$$

$$R = \frac{1}{\rho} = 1$$

$$A \subseteq \mathbb{R}$$

$$(x_0 - R, x_0 + R) \subseteq A \subseteq [x_0 - R, x_0 + R] = \begin{cases} A \subseteq \mathbb{R} \\ (-1, 1) \subseteq \mathbb{R} \subseteq [-1, 1] \end{cases}$$

$$-1 \in A? \quad \sum_{n=1}^{\infty} \frac{(-1)^n (-1)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{2n}}{n} = \sum_{n=1}^{\infty} \frac{1}{n} \Rightarrow \text{serie armonica cu } \alpha \leq 1 \Rightarrow \text{suma este divergenta} \Rightarrow -1 \notin A$$

$$1 \in A? \quad \sum_{n=1}^{\infty} \frac{(-1)^n (1)^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0(1)$$

$$\frac{1}{n+1} \leq \frac{1}{n} (2)$$

$$\Rightarrow \text{conf. crit. lui Leibniz} \Rightarrow \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{ conv.} \Rightarrow 1 \in A$$

$$A = (-1, 1]$$

$$f : A = (-1, 1] \rightarrow \mathbb{R}, f(x) = \sum_{n=1}^{\infty} \frac{(-1)^n x^n}{n}$$

$$f'(x) = \sum_{n=1}^{\infty} \left(\frac{(-1)^n x^n}{n} \right)' = \sum_{n=1}^{\infty} \frac{n(-1)^n x^{n-1}}{n} = - \sum_{n=1}^{\infty} (-1)^{n-1} x^{n-1} \forall x \in (-1, 1]$$

$$f'(x) = - \sum_{n=1}^{\infty} (-1)^{n-1} x^{n-1}, \forall x \in (-1, 1]$$

$$\underline{\underline{n = m + 1}} - \sum_{m=0}^{\infty} (-1)^m x^m = - \frac{1}{1+x} \forall x \in (-1, 1) \Big| \int$$

$$f(x) = - \int \frac{1}{1+x} dx = - \ln(1+x) + C, \forall x \in (-1, 1)$$

$$f(x_0) = a_0 \Leftrightarrow f(0) = 0 \Leftrightarrow - \ln(1+0) + C = 0 \Leftrightarrow C = 0$$

$$f(x) = - \ln(1+x), \forall x \in (-1, 1)$$

$$f \text{ cont în } 1 \Rightarrow f(1) = \lim_{\substack{x \rightarrow 1 \\ x < 1}} f(x)$$

$$f(1) = -\ln 2$$

$$\lim_{\substack{x \rightarrow 1 \\ x < 1}} f(x) = -\ln 2$$

$$\Rightarrow f(x) = -\ln(1+x)$$

Demonstrație pentru două din seriile de puteri speciale:

$$\begin{aligned} S &= \sum_{n=0}^{\infty} x^n \Leftrightarrow S = \lim_{n \rightarrow \infty} 1 + x + x^2 + \dots + x^n \Leftrightarrow xS = \lim_{n \rightarrow \infty} x + x^2 + x^3 + \dots + x^{n+1} \Leftrightarrow \\ \Leftrightarrow xS - S &= \lim_{n \rightarrow \infty} x + x^2 + x^3 + \dots + x^{n+1} - 1 - x - x^2 - \dots - x^n \Leftrightarrow S(x-1) = \lim_{n \rightarrow \infty} x^{n+1} - 1 \Leftrightarrow \\ \Leftrightarrow S &= \lim_{n \rightarrow \infty} \frac{x^{n+1} - 1}{x - 1}, \text{ dar } x \in (-1, 1) \Rightarrow S = \frac{1}{1-x} \end{aligned}$$

$$f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = \sin x$$

$$f(x) = \sin x, \forall x \in \mathbb{R} \Rightarrow f(0) = 0$$

$$f'(x) = \cos x, \forall x \in \mathbb{R} \Rightarrow f'(0) = 1$$

$$f''(x) = -\sin x, \forall x \in \mathbb{R} \Rightarrow f''(0) = 0$$

$$f'''(x) = -\cos x, \forall x \in \mathbb{R} \Rightarrow f'''(0) = -1$$

$$f^{(4)}(x) = \sin x, \forall x \in \mathbb{R} \Rightarrow f^{(4)}(0) = 0$$

.....

conf. Formulei lui Taylor cu restul sub forma lui Lagrange, $\forall x \in \mathbb{R}^* (i.e. x \neq 0)$,

$\exists c$ între 0 și x (i.e. $c \in (0, x)$ sau $c \in (x, 0)$) a.i.

$$f(x) = f(0) + \frac{f'(0)}{1!}(x-0) + \frac{f''(0)}{2!}(x-0)^2 + \dots + \frac{f^{(n)}(0)}{n!}(x-0)^n + \underbrace{\frac{f^{(n+1)}(c)}{(n+1)!}(x-0)^{n+1}}_{R_n(x)}$$

$$R_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!}(x-0)^{n+1}$$

$$|R_n(x)| = \frac{|f^{(n+1)}(c)|}{n+1} |x|^{n+1} \leq \frac{1}{(n+1)!} |x|^{n+1} = \frac{|x|^{n+1}}{(n+1)!}.$$

$$\text{Fie } x_n = \frac{|x|^{n+1}}{(n+1)!}, \forall n \in \mathbb{N}.$$

$$\lim_{n \rightarrow \infty} \frac{x_{n+1}}{x_n} = \lim_{n \rightarrow \infty} \frac{\frac{|x|^{n+2}}{(n+1)!}}{\frac{|x|^{n+1}}{(n+1)!}} = \lim_{n \rightarrow \infty} \frac{|x|}{n+2} = 0 \leq 1.$$

Conform Crit. raportului (de notat ca crit. raportului are doua def, una pt. limita si una pt. conv.)

$$\lim_{n \rightarrow \infty} x_n = 0.$$

Avem

$$\begin{array}{ccc} 0 & \leq & |R_n(x)| \leq x_n \\ & \searrow & \swarrow \\ & 0 & \end{array}$$

$$\text{Deci } \lim_{n \rightarrow \infty} |R_n(x)| = 0; \text{ i.e. } \lim_{n \rightarrow \infty} R_n(x) = 0.$$

$$\text{Prin urmare } f(x) = \sum_{n=0}^{\infty} \frac{f^n(0)}{n!} (x-0)^n = 0 + \frac{1}{1!}x + 0 - \frac{1}{3!}x^3 + 0 + \frac{1}{5!}x^5 + \dots = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1}$$

$$\left. \begin{aligned} f(0) &= \sin 0 = 0 \\ \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} 0^{2n+1} &= 0 \end{aligned} \right| \Rightarrow f(0) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} 0^{2n+1}$$

$$\text{Deci } f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} x^{2n+1}, \forall x \in \mathbb{R}$$

Restul seriilor de puteri ori se rezolvă în stil asemănător cu acestea ori sunt consecințe ale lor

$$\sum_{n=0}^{\infty} (n+2)(n+1)x^n$$

$$\begin{cases} x_0 = 0 \\ a_n = (n+2)(n+1), n \in \mathbb{N}^* \\ a_0 = 0 \end{cases}$$

$$R = \frac{1}{\rho}$$

$$\rho = \overline{\lim} \sqrt[n]{|a_n|}, \text{ sau } \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} (\text{daca } \exists)$$

$$\rho = \lim_{n \rightarrow \infty} \sqrt[n]{(n+2)(n+1)} = 1$$

$$R = \frac{1}{\rho} = 1$$

$$A \subseteq \mathbb{R}$$

$$(x_0 - R, x_0 + R) \subseteq A \subseteq [x_0 - R, x_0 + R] = \begin{cases} A \subseteq \mathbb{R} \\ (-1, 1) \subseteq \mathbb{R} \subseteq [-1, 1] \end{cases}$$

$$-1 \in A? \quad \sum_{n=0}^{\infty} (n+2)(n+1)(-1)^n$$

$$\lim_{n \rightarrow \infty} (n+2)(n+1) = \infty \Rightarrow \text{suma este div} \Rightarrow -1 \notin A$$

$$1 \in A? \quad \sum_{n=0}^{\infty} (n+2)(n+1)$$

$$\lim_{n \rightarrow \infty} (n+2)(n+1) = \infty \Rightarrow \text{suma este div} \Rightarrow 1 \notin A$$

$$A = (-1, 1)$$

$$\begin{aligned}
f : A(= (-1, 1)) &\rightarrow \mathbb{R}, f(x) = \sum_{n=1}^{\infty} (n+2)(n+1)x^n \\
\int f(x) dx &= \int \sum_{n=0}^{\infty} (n+2)(n+1)x^n dx = \sum_{n=0}^{\infty} \int (n+2)(n+1)x^n dx = \sum_{n=1}^{\infty} (n+2)x^{n+1} + C, \forall x \in (-1, 1) \\
\int \int f(x) + C dx &= \int \sum_{n=0}^{\infty} (n+2)x(n+1) + C dx = \sum_{n=0}^{\infty} \int (n+2)x^{n+1} + C dx = \sum_{n=1}^{\infty} x^{n+2} + C, \forall x \in (-1, 1) \\
\int \int f(x) dx dx &= \sum_{n=1}^{\infty} x^{n+2} + C, \forall x \in (-1, 1) \\
\underline{n = m - 2} \sum_{m=3}^{\infty} x^m + C &= \sum_{m=0}^{\infty} x^m + C - x^0 - x^1 - x^2 = \frac{1}{1-x} - 1 - x - x^2 + C, \forall x \in (-1, 1) \Big|'' \\
f(x) &= \left(\frac{1}{1+x} - 1 - x - x^2 + C \right)'' = \left(-\frac{1}{(1+x)^2} - 1 - 2x \right)' = \frac{2}{(1+x)^3} - 2, \forall x \in (-1, 1) \\
f(x) &= \frac{2}{(1+x)^3} - 2, \forall x \in (-1, 1)
\end{aligned}$$

$$\sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} (x+2)^n$$

$$\begin{cases} x_0 = 2 \\ a_n = \frac{(-2)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} \forall n \in \mathbb{N}^* \\ a_0 = 0 \end{cases}$$

$$R = \frac{1}{\rho}$$

$$\rho = \overline{\lim} \sqrt[n]{|a_n|}, \text{ sau } \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} \text{ (daca } \exists \text{)}$$

$$\rho = \lim_{n \rightarrow \infty} \frac{\left| \frac{(-2)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} \right|}{\left| \frac{(-2)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} \right|} = \lim_{n \rightarrow \infty} \left| -2 \sqrt[3]{\frac{n+1}{n+2}} \sqrt[4]{\frac{n+2}{n+3}} \right| = 2$$

$$R = \frac{1}{\rho} = \frac{1}{2}$$

$$A \leq \mathbb{R}$$

$$(x_0 - R, x_0 + R) \leq A \leq [x_0 - R, x_0 + R] = \begin{cases} A \leq \mathbb{R} \\ (\frac{3}{2}, \frac{5}{2}) \leq \mathbb{R} \leq [\frac{3}{2}, \frac{5}{2}] \end{cases}$$

$$\frac{3}{2} \in A? \quad \sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} \left(\frac{7}{2}\right)^n$$

$$\lim_{n \rightarrow \infty} \frac{\left(\frac{7}{2}\right)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} \stackrel{\infty}{=} \infty$$

$$\sqrt[3]{n+1}\sqrt[4]{n+2} \ll \left(\frac{7}{2}\right)^n, \text{ exponen\c{t}iala cre\c{s}te mult mai rapid dec\c{a}t polinomiala}$$

$$\infty \Rightarrow \text{suma este div} \Rightarrow \frac{3}{2} \notin A$$

$$\frac{5}{2}\in A? \quad \sum_{n=1}^{\infty} \frac{(-2)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} \left(\frac{9}{2}\right)^n$$

$$\lim_{n\rightarrow\infty} \frac{\left(\frac{9}{2}\right)^n}{\sqrt[3]{n+1}\sqrt[4]{n+2}} \stackrel{8}{=} \infty$$

$$\sqrt[3]{n+1}\sqrt[4]{n+2} \ll \left(\frac{9}{2}\right)^n, \text{ exponen\c{t}iala cre\c{s}te mult mai rapid dec\c{a}t polinomiala}$$

$$\infty \Rightarrow \text{suma este div} \Rightarrow \frac{5}{2} \notin A$$

$$A = \left(\frac{3}{2}, \frac{5}{2}\right)$$