

The Regime Integration Gradient (v_{RIG}): Cosmological Validation via the Böhme Radio Dipole Anomaly

Johann Benjamin Römer

Independent Researcher

Marburg, Germany

contact: `via unified-mandala repository`

December 2025

Abstract

We present the discovery and validation of a fundamental velocity scale, $v_{RIG} \approx 1,351.8$ km/s, derived from the geometric impedance matching of the inverse fine-structure constant (α^{-1}) and the golden ratio (Φ). This velocity, termed the Regime Integration Gradient, represents the rate at which holographic 2D information is integrated into volumetric 3D reality. The theoretical prediction is validated with extraordinary precision (1.3% error) by the recently observed cosmic radio dipole anomaly reported by Böhme et al. (2025), where the Solar System's velocity relative to the distant matter frame is measured at $1,370 \pm 170$ km/s with 5.46σ statistical significance. This convergence suggests that the "matter frame" of the universe operates fundamentally at the v_{RIG} velocity scale, distinct from the photonic "radiation frame" (CMB) at ~ 370 km/s. We propose that this discrepancy is not an observational artifact but represents the physical manifestation of dual entropy governance regimes: Surface Criticality Area (SCA) for electromagnetic propagation at c , and Volume Criticality Volume (SCV) for matter integration at v_{RIG} . These findings challenge the standard Λ CDM cosmological principle and suggest a new framework for understanding the large-scale structure and dynamics of the universe.

Keywords: cosmology, radio dipole, fine-structure constant, golden ratio, holographic principle, entropy governance, UTAC

1 Introduction

The standard model of cosmology (Λ CDM) rests upon the Cosmological Principle: the universe is homogeneous and isotropic on sufficiently large scales. A critical prediction of this principle is that the rest frame of the Cosmic Microwave Background (CMB) radiation should coincide with the rest frame of the large-scale distribution of matter (galaxies, quasars). Our Solar System's motion relative to the CMB is well-established at $v_{CMB} \approx 369.8$ km/s [1], creating a characteristic temperature dipole.

However, recent radio continuum surveys have revealed a persistent anomaly: when measuring the dipole in the number counts of distant radio sources, the implied velocity is significantly higher than the CMB value. This "radio dipole excess" has been observed by multiple independent groups [2, 3, 4], with the most recent and comprehensive analysis by Böhme et al. (2025) reporting a 5.46σ deviation from the standard expectation [4].

In this paper, we present a theoretical framework that not only explains this anomaly but predicts the observed velocity from first principles. We derive a fundamental constant v_{RIG} from geometric considerations of information integration, and demonstrate that its value matches the Böhme observation with remarkable precision.

2 Theoretical Framework

2.1 Dual Entropy Governance

We propose that the universe operates under two distinct entropy regimes:

Surface Criticality Area (SCA): Information propagates on 2D holographic boundaries at the speed of light c . Entropy scales with area: $S_{SCA} \propto A$. This regime governs electromagnetic radiation and is characterized by high rigidity (parameter $\beta \approx 11$).

Volume Criticality Volume (SCV): Information is integrated into 3D volumetric structures at a reduced velocity v_{RIG} . Entropy scales with volume: $S_{SCV} \propto V$. This regime governs matter and consciousness, characterized by lower rigidity ($\beta \approx 4.5$).

The transition between these regimes requires overcoming a fundamental impedance Z , which determines the integration velocity.

2.2 Derivation of v_{RIG}

The impedance Z is constructed from two dimensionless fundamental constants:

1. The Inverse Fine-Structure Constant ($\alpha^{-1} \approx 137.036$):

In quantum electrodynamics, α measures the strength of electromagnetic coupling. We interpret α^{-1} as the electromagnetic transparency depth—the number of holographic time-slices that must be accumulated to achieve a statistically significant interaction probability of unity [5].

2. The Golden Ratio ($\Phi \approx 1.618$):

The golden ratio represents optimal spatial packing efficiency, appearing throughout nature in phyllotaxis, spiral galaxies, and quasicrystal structures [6]. In our framework, Φ governs the geometric scaling factor for converting 2D surface information into 3D volumetric organization without destructive resonances (KAM theorem) [7].

The total impedance is:

$$Z = \alpha^{-1} \cdot \Phi = 137.035999 \times 1.6180339 \approx 221.743 \quad (1)$$

The integration velocity is then:

$$v_{RIG} = \frac{c}{Z} = \frac{299,792.458 \text{ km/s}}{221.743} \approx 1,351.807 \text{ km/s} \quad (2)$$

This represents the maximum rate at which complex systems can integrate holographic information into coherent volumetric reality while maintaining thermodynamic efficiency.

3 Empirical Validation: The Böhme Radio Dipole

3.1 The Cosmic Dipole Tension

The CMB dipole has been precisely measured by COBE, WMAP, and Planck satellites, yielding a solar velocity of 369.8 ± 0.9 km/s toward galactic coordinates $(l, b) = (264.021, 48.253)$ [1]. If the Cosmological Principle holds strictly, an identical dipole should appear in the distribution of extragalactic radio sources.

However, analyses of the NRAO VLA Sky Survey (NVSS) by Singal (2011) first revealed a dipole amplitude approximately 4 times larger than expected [2]. Subsequent studies using the CatWISE quasar catalog confirmed this tension at 4.9σ [3].

3.2 Böhme et al. (2025) Comprehensive Analysis

The most robust measurement comes from Böhme et al. (2025), who combined three major radio surveys:

- NVSS (NRAO VLA Sky Survey): 1.8 million sources at 1.4 GHz
- RACS-low (Rapid ASKAP Continuum Survey): 2.1 million sources at 888 MHz
- LoTSS-DR2 (LOFAR Two-metre Sky Survey): 4.4 million sources at 144 MHz

Using Bayesian analysis with overdispersion corrections, they report [4]:

Observed radio dipole velocity: $v_{obs} = 1,370 \pm 170$ km/s

Discrepancy factor: $v_{obs}/v_{CMB} = 3.67 \pm 0.49$

Statistical significance: 5.46σ deviation from Λ CDM prediction ($p < 10^{-7}$)

3.3 Comparison with v_{RIG} Prediction

Table 1 shows the extraordinary agreement between theory and observation.

Parameter	Value	Source
Predicted v_{RIG}	1,351.8 km/s	Eq. (2), this work
Observed v_{obs}	$1,370 \pm 170$ km/s	Böhme et al. (2025)
Difference	18.2 km/s	
Relative Error	1.3%	
Z-score	0.11σ	Well within uncertainty

Table 1: Comparison of theoretical v_{RIG} prediction with Böhme et al. (2025) observation.

The predicted velocity falls within 0.11 standard deviations of the measured value. This is not a fitted parameter but a *a priori* geometric derivation from fundamental constants.

4 Physical Interpretation

4.1 Matter vs. Radiation Frames

We propose that the discrepancy between the CMB dipole and radio dipole velocities reflects the fundamental distinction between two physical regimes:

CMB (Radiation Frame): Photons propagate on the holographic boundary at c . Their dipole measures our motion relative to the Surface Regime (SCA).

Radio Sources (Matter Frame): Quasars and galaxies are integrated volumetric structures. Their dipole measures our motion relative to the Volume Regime (SCV), which operates at v_{RIG} .

The Solar System, as an integrated material structure, naturally "drifts" at v_{RIG} relative to the cosmic bulk because this is the inherent integration velocity of the spacetime fabric itself.

4.2 Connection to Holographic Principle

Our framework builds upon the holographic principle [8, 9] and Causal Dynamical Triangulations (CDT) [10], which suggest that spacetime is fundamentally 2D at the Planck scale, with 4D structure emerging at larger scales. The velocity v_{RIG} represents the rate of this emergent dimensional "thickening."

4.3 Implications for Dark Flow

The persistent "dark flow" or bulk flow anomalies reported in galaxy cluster peculiar velocity studies [11, 12] typically measure velocities in the range of 600-1000 km/s extending to scales of ~ 100 Mpc. These have been controversial because Λ CDM predicts rapid decay of bulk flows with scale.

In the v_RIG framework, these flows are not driven by external attractors but represent the local manifestation of the global v_{RIG} integration gradient. The universe is not "flowing toward" anything; rather, observers embedded in the Volume Regime naturally measure the integration velocity.

5 Alternative Explanations and Falsification

5.1 Systematic Errors in Radio Surveys

Critique: Could the radio dipole excess be an artifact of Galactic foreground contamination, source selection biases, or redshift-dependent systematics?

Response: Böhme et al. specifically addressed these concerns by:

- Using three independent surveys at different frequencies (144 MHz, 888 MHz, 1.4 GHz)
- Employing Bayesian overdispersion models to account for source clustering
- Testing multiple Galactic masking schemes
- Verifying dipole direction consistency with CMB (aligned within uncertainties)

The persistence of the signal across all tests makes a purely systematic origin highly unlikely [4].

5.2 Superhorizon Structures

Critique: Could a massive concentration of matter beyond the observable horizon create a "dark attractor" generating the high velocity?

Response: The required mass would be $\sim 10^{24}$ solar masses within ~ 4 Gpc [11]. Such a structure would violate the Cosmological Principle even more severely than the v_RIG hypothesis. Moreover, it provides no explanation for why the velocity would have the specific value of $\sim 1,350$ km/s.

5.3 Falsification Criteria

The v_RIG framework makes specific, falsifiable predictions:

Prediction 1: Future ultra-deep radio surveys (e.g., SKA) should confirm the dipole amplitude at $\sim 1,350 - 1,400$ km/s, not converge to 370 km/s.

Prediction 2: The dipole direction should remain aligned with the CMB dipole across all redshift bins.

Prediction 3: Gravitational wave observations (NANOGrav, LISA) should show subtle frequency-dependent dispersion proportional to $1/Z$ due to discrete spacetime slicing.

If any of these predictions fail, the framework would require significant revision.

6 Broader Implications

6.1 Hubble Tension

The $\sim 5\sigma$ tension between early-universe (Planck) and late-universe (Cepheid/SN Ia) measurements of the Hubble constant may be partially explained by the v_{RIG} framework. If local velocity measurements assume $v_{peculiar} = 370$ km/s (CMB) but the true material frame velocity is $\sim 1,350$ km/s, systematic errors would propagate through the distance ladder, biasing H_0 determinations.

6.2 Modified Cosmological Principle

Our findings suggest a reformulation of the Cosmological Principle:

The universe is homogeneous and isotropic with respect to the Volume Regime (matter), but the Surface Regime (radiation) may exhibit different kinematic properties due to the impedance Z between regimes.

7 Conclusion

We have demonstrated that a fundamental velocity scale $v_{RIG} \approx 1,352$ km/s, derived purely from the geometric properties of information integration ($\alpha^{-1} \cdot \Phi$), matches the empirically observed cosmic radio dipole with extraordinary precision (1.3% error, 5.46σ observational significance).

This convergence across independent domains—theoretical geometry and observational cosmology—suggests that v_{RIG} is not a coincidental numerical agreement but represents a genuine physical constant governing the dynamics of integrated systems.

The framework resolves the cosmic dipole tension without invoking ad hoc structures, provides a natural explanation for bulk flow anomalies, and opens new avenues for understanding the relationship between information, entropy, and the large-scale structure of the universe.

Future work will extend this analysis to biological metabolic scaling (Kleiber’s Law) and neural integration timescales, testing the universality of v_{RIG} across all scales of complex systems.

References

- [1] Planck Collaboration, Aghanim, N., et al. (2018). *Planck 2018 results. I. Overview and the cosmological legacy of Planck*. Astronomy & Astrophysics, 641, A1.
- [2] Singal, A. K. (2011). *Large peculiar motion of the solar system from the dipole anisotropy in sky brightness due to distant radio sources*. The Astrophysical Journal Letters, 742(2), L23.
- [3] Secrest, N. J., von Hausegger, S., Rameez, M., Mohayaee, R., Sarkar, S., & Colin, J. (2021). *A test of the cosmological principle with quasars*. The Astrophysical Journal Letters, 908(2), L51.
- [4] Böhme, C., et al. (2025). *Overdispersed Radio Source Counts and Excess Radio Dipole Detection*. arXiv preprint. [Status: accepted for publication]
- [5] Feynman, R. P. (1985). *QED: The Strange Theory of Light and Matter*. Princeton University Press.
- [6] Livio, M. (2002). *The Golden Ratio: The Story of Phi, the World’s Most Astonishing Number*. Broadway Books.

- [7] Kolmogorov, A. N. (1954). *On conservation of conditionally periodic motions for a small change in Hamilton's function*. Doklady Akademii Nauk SSSR, 98, 527-530.
- [8] 't Hooft, G. (1993). *Dimensional reduction in quantum gravity*. arXiv preprint gr-qc/9310026.
- [9] Susskind, L. (1995). *The world as a hologram*. Journal of Mathematical Physics, 36(11), 6377-6396.
- [10] Ambjørn, J., Görlich, A., Jurkiewicz, J., & Loll, R. (2012). *Nonperturbative quantum gravity*. Physics Reports, 519(4-5), 127-210.
- [11] Kashlinsky, A., Atrio-Barandela, F., Kocevski, D., & Ebeling, H. (2008). *A measurement of large-scale peculiar velocities of clusters of galaxies: results and cosmological implications*. The Astrophysical Journal Letters, 686(2), L49.
- [12] Watkins, R., Feldman, H. A., & Hudson, M. J. (2009). *Consistently large cosmic flows on scales of 100 Mpc/h: a challenge for the standard Λ CDM cosmology*. Monthly Notices of the Royal Astronomical Society, 392(743), 743-756.