Is seismicity in glaciated regions becoming more sensitive to unloading?

Early-stage tests, upper bounds and a preregistered confirmatory design (P133, v0.2.1 draft)

Johann Benjamin Römer*

Independent researcher, Kirchhain, Germany

jbroemer@web.de

Draft of 15 October 2026 (not peer reviewed)

Abstract

Glacial unloading is known to promote faulting over isostatic time scales and to modulate seismicity seasonally. Whether present-day ice loss already makes seismicity in glaciated regions measurably more sensitive is open. We test this with public catalogues (USGS ComCat and five national networks, 1990–2025), GRACE/GRACE-FO mascons, GNSS and simple physical scenarios, and we report upper bounds wherever we find nothing. Globally, the declustered $M \geq 6$ rate shows no increase since 1990 ($-0.31\,\%$/yr, 95 % CI -0.62 to 0.00), and declustered $M \geq 5$ rates in six glaciated boxes did not rise relative to five tectonic controls (pooled ratio 0.95, 0.65–1.39). Southeast Alaska has a robust seasonal modulation (amplitude 0.11, sensitivity $4.0 \pm 1.1\,\%$/kPa), but its growth since 1990 ($+0.035 \pm 0.023$ per decade) is not significant after permutation and multiplicity correction; no group contrast between glaciated and control regions survives (smallest $q = 0.76$). A preregistered test of the time-varying ETAS background (H2.1) was formally positive (difference $+0.405 \pm 0.138$ per decade, $p = 0.0017$), but the effect disappears when the analysis starts in 2003 and is partly explained by network covariates; we read it as a catalogue artefact rather than evidence. No region shows the predefined "masked background rise" and no tidal-phase test is significant. A viscoelastic lag model and tipping scenarios place detectable rate changes decades ahead (earliest about 2046 in the steepest mountain-glacier scenario) and predict a shallow (4–16 %) "deceptive quiet" before acceleration. We therefore treat glacially driven sensitisation as an early-stage hypothesis: not supported, not excluded, with quantified limits. Two confirmatory tests (C1 from November 2026, C2 from January 2027; looks in 2029 and 2036) were frozen and archived before any of their data exist.

Keywords: glacial isostatic adjustment; seasonal seismicity; ETAS; preregistration; tidal triggering; deceptive quiescence; negative results.

Status labels used throughout: [PREREG] = analysis rule fixed in a timestamped, hashed file before the analysis was run; [EXPL] = exploratory, chosen after seeing data. Retrospective tests on 1990–2025 data are *pre-specified but data-touched*; only C1 and C2 are prospective (Section 4).

Contents

1 Introduction **2**

2 Data **3**

*Corresponding author. Address: Marburger Ring 34 B, 35274 Kirchhain-Großseelheim, Germany. Email: jbroemer@web.de (primary); joberoemer@gmail.com (secondary). The author is a certified IT specialist for application development (Fachinformatiker für Anwendungsentwicklung) with a background in web design and works as an independent researcher.

3 Methods **4**
3.1 Catalogue handling, completeness and declustering . . . 4
3.2 Seasonal modulation and load sensitivity (F1–F3) . . . 5
3.3 Criticality indicators . . . 5
3.4 Two-stage ETAS with a time-varying background (H2) . . . 5
3.5 Tidal phase tests (H3) . . . 6
3.6 Viscoelastic lag model (F5) . . . 6
3.7 Tipping scenarios and deceptive-quiet model (A, B) . . . 6
3.8 Quiescence and monitoring indicators . . . 7
3.9 Global baseline [EXPL] . . . 7

4 Preregistration and confirmatory design **7**

5 Results **8**
5.1 Global baseline [EXPL] . . . 8
5.2 Seasonal modulation in Southeast Alaska (F1, F2) . . . 8
5.3 Criticality indicators . . . 9
5.4 ETAS background: H2.1 positive but not robust . . . 9
5.5 Masked background rise: H2.2 . . . 10
5.6 Tides: H3 . . . 11
5.7 Lag model and tipping scenarios . . . 12
5.8 Deceptive quiet before acceleration . . . 13
5.9 Present-day precursors: GRACE and GNSS . . . 14

6 Review process and errata **15**

7 Discussion **16**

8 Limitations **16**

9 Conclusions **17**

References **18**

1 Introduction

Glacial unloading and faulting. Large ice sheets suppress earthquakes while they stand and promote faulting when they melt (Johnston 1987; Hampel et al. 2010). The clearest examples are the end-glacial faults of northern Fennoscandia, which ruptured in large earthquakes shortly after deglaciation (Lagerbäck and Sundh 2008) and still host microseismicity today, for example on the Pärvie fault (Lindblom et al. 2015). On these time scales the controlling process is glacial isostatic adjustment (GIA): unloading changes the Coulomb stress on faults directly through the elastic response and with a delay through viscous mantle flow (Turcotte and Schubert 2002; Hampel et al. 2010).

Present-day ice loss. Today's ice loss is much smaller than the Pleistocene deglaciation but very fast. Southeast Alaska is rising rapidly in response to post-Little-Ice-Age and recent ice loss (Larsen et al. 2004; Larsen et al. 2005), and Rollins et al. (2021) showed that GIA together with inter-earthquake stress transfer promoted the 1958 $M_w \sim 7.8$ Fairweather earthquake and other events in the region. Comparable rapid uplift is documented in Patagonia (Lange et al. 2014), the Amundsen Sea Embayment (Barletta et al. 2018), Greenland (Khan et al. 2016), Iceland (Pagli

et al. 2007) and the European Alps (Mey et al. 2016); low-viscosity models for Alaska and Iceland have recently been updated by Sasgen et al. (2026).

Seasonal loading as a natural experiment. Seasonal changes of snow, ice and water load modulate seismicity in many settings: in the Himalaya (Bettinelli et al. 2008), in California (Johnson et al. 2017; Sirorattanakul and Avouac 2026), in northeastern Japan (Ueda et al. 2024) and in southern Alaska, where Johnson et al. (2020) found a hydrospheric stress sensitivity of a few per cent per kPa with a lag of about three months. At the Grandes Jorasses in the Mont Blanc massif, Simon et al. (2025) documented a recent onset of seasonal microseismicity that they relate to climate change, and for which a hydraulic mechanism (meltwater infiltration and pore pressure) is a natural candidate (cf. Talwani et al. 2007). Seasonal modulation is therefore an observable that responds to load changes on human time scales.

Volcanism. In Iceland, post-glacial unloading increased eruption rates strongly after the last deglaciation (Maclennan et al. 2002), and mid-Holocene climate and ice changes left a detectable imprint on eruption frequency (Swindles et al. 2018). Present-day retreat of Vatnajökull is modelled to increase mantle melt production (Pagli and Sigmundsson 2008; Schmidt et al. 2013). Whether this translates into measurable changes of eruption frequency within decades is unknown.

Motivation and framing. The question addressed here is narrower than "does ice loss cause earthquakes?": *Is the sensitivity of seismicity in glaciated regions to (un)loading changing now, and can that be detected with existing catalogues?* Two framing choices follow from the physics and from the data.

1. *Distribution, not mean.* A change in loading need not raise the average rate. It can change the seasonal modulation, the background (non-triggered) rate, the response to tides or the share of promoted versus shadowed faults (Harris 1998). A shift of the rate distribution may even appear as a temporary quiet. We therefore test several aspects of the rate distribution, not only the mean.

2. *Absence of evidence, with upper bounds.* Catalogues in glaciated regions are short, sparse and affected by network changes. A non-significant result means little unless the effect size that could have been detected is stated. We report minimum detectable effects (MDE) or confidence intervals for every null result.

By *sensitivity* we mean the seismic response per unit change of load or stress, for example the rate change per kPa of seasonal unloading (γ_0 in F2). A larger response caused only by a larger forcing at unchanged sensitivity is a different statement. F2 normalises by the GRACE load and therefore addresses sensitivity; the seasonal amplitude in F1 and C1 can also grow because the seasonal load itself grows, so F1 and C1 alone cannot separate the two.

Because many tests are run on data that have already been examined, the work also includes two preregistered prospective tests (C1, C2) whose rules were frozen and archived before their data exist. This paper describes the state of the project at release v0.2.0 (Römer 2026a) plus an exploratory global baseline added afterwards; it is intended to accompany release v0.2.1.

2 Data

All raw data are public and were retrieved on 2026-10-10; analyses end on 2025-12-31. Raw downloads are not part of the repository but can be regenerated with `glacial-sensitivity fetch`; archived snapshots are included for the global baseline.

Earthquake catalogues. USGS ComCat (U.S. Geological Survey 2026) for global analyses and for boxes without a national catalogue ("ComCat tier"); the Alaska Earthquake Center

Table 1: National-tier regions (`results/regions_meta.csv`). M_c^{ref}: 90 % quantile of yearly M_c 1990–2025. $N_{\geq M_c}$: events $\geq M_c^{ref}$ 1990–2025; N_{decl}: background events after nearest-neighbour declustering.

Region	Role	Catalogue	M_c^{ref}	$N_{\geq M_c}$	N_{decl}
SE Alaska (`SE_Alaska_AK`)	glaciated, primary	AEC	2.1	11 513	5 314
Alps (`Alps_SED`)	glaciated	SED	2.1	2 163	1 335
NZ Southern Alps	glaciated	GeoNet	2.8	12 486	4 405
Iceland, Vatnajökull	glaciated	IMO	2.5	5 866	1 344
Greenland	ice sheet	GEUS	2.7	898	531
Interior Alaska	control (primary)	AEC	2.1	14 023	7 234
N Switzerland	control	SED	2.3	488	316
NZ Wellington	control	GeoNet	2.7	15 415	10 693

contribution to ComCat (`catalog=ak`; Alaska Earthquake Center 2026); the Swiss Seismological Service (Swiss Seismological Service (SED) 2026); GeoNet New Zealand (GeoNet 2026); the Icelandic Meteorological Office SIL catalogue for Vatnajökull (Icelandic Meteorological Office (Veðurstofa Íslands) 2026); and the GEUS Greenland catalogue (Geological Survey of Denmark and Greenland (GEUS) 2026). The relocated Chilean catalogue of Potin et al. (2025) (data: Potin et al. 2024) serves as an additional control. Table 1 lists the national-tier regions used in the main tests. Depth and polygon filters per catalogue are documented in `docs/DATA.md`. Only events typed as earthquakes are kept; icequakes, landslides or tremor that a network did not flag remain in the data (Section 8).

Loads and geodesy. GRACE/GRACE-FO GSFC mascons RL06 v2.0, monthly 2002-04 to 2026-03 (NASA Goddard Space Flight Center 2026; Loomis et al. 2019), converted to surface load (1 cm water equivalent = 0.0981 kPa); ERA5 precipitation, rain and snowfall via the Open-Meteo archive (Open-Meteo 2026; Hersbach et al. 2020); GNSS daily positions (IGS20) and step files of the Nevada Geodetic Laboratory (Nevada Geodetic Laboratory 2026; Blewitt et al. 2018); FDSN station metadata from EarthScope for network histories (EarthScope Consortium 2026).

Volcanism and tides. Smithsonian GVP eruption starts (Global Volcanism Program 2026); body tides from the degree-2 potential with Meeus (1998) ephemerides; ocean loading with SPOTL (Agnew 1997; Agnew 2012) and TPXO 7.2 (Egbert and Erofeeva 2002).

3 Methods

Table 2 gives the status of every analysis. All code is in the open-source Python package of the project (Römer 2026c; Römer 2026d).

3.1 Catalogue handling, completeness and declustering

Yearly completeness M_c is estimated with the maximum-curvature method plus 0.2; the reference M_c^{ref} is the 90 % quantile of the yearly values 1990–2025, so that most years are complete. Robustness checks use the entire-magnitude-range (EMR) estimator and Mc bootstraps [EXPL]. The main declustering is the nearest-neighbour method of Zaliapin and Ben-Zion (2013) (fractal dimension 1.6, $b = 1$, threshold η_0 from a two-component Gaussian mixture of $\log \eta$). Gardner–Knopoff windows (Gardner and Knopoff 1974) in the form of van Stiphout et al. (2012) serve as an alternative [EXPL] and for the global baseline, with ties in magnitude processed in time order. Because detection in glaciated regions may vary seasonally, M_c was also estimated by calendar month (all years pooled; `seasonal_mc_by_month.csv`): it stays at or below M_c^{ref} in every national-tier region except SE Alaska in December (2.2 versus 2.1). A probabilistic detection model (Ogata and Katsura 1993) was not fitted.

Table 2: Analyses and their status. Retrospective tests use 1990–2025 data that had been looked at in part before the rules were fixed.

Code	Analysis	Data	Status
F1, F2, F3	Growth of seasonal modulation; load sensitivity; rain vs load	1990/2002–2025	pre-specified, retrospective (v0.1)
Crit.	Criticality battery (trend, CSD, group tests)	1990–2025	pre-specified, retrospective
F5, A, B	Lag model; tipping scenarios; deceptive quiet	scenarios	model calculations, no test
M1–M5	Monitoring indicators (GRACE, GNSS, modulation, b, quiet)	2002–2026	frozen thresholds; flags, not tests
H2.1, H2.2	ETAS background trend (DiD); masked-rise classification	1995–2025	[PREREG] retrospective (v0.2)
H3.1, H3.2	Body-tide phase (Schuster), b by tidal phase	1995–2025	[PREREG] retrospective (v0.2)
C1	Seasonal modulation, new vs frozen baseline	from 2026-11-01	[PREREG] prospective
C2	ETAS background, glaciated minus control	from 2027-01-01	[PREREG] prospective
—	Global baseline, ice vs control	1975–2026	[EXPL]
—	Robustness variants, network covariates, ocean loading, M4 lag sensitivity	various	[EXPL]

3.2 Seasonal modulation and load sensitivity (F1–F3)

Seasonality is first tested with the Schuster test (Ader and Avouac 2013). F1 fits a Poisson GLM to monthly declustered counts with year fixed effects,

$$\log \mu_{y,m} = \alpha_y + (\beta_0 + \beta_1 \tau) \cos\Big(2\pi \left[\tfrac{m-0.5}{12} - \varphi\right]\Big), \qquad \tau = (t - 2008)/10, \tag{1}$$

so that β_1 is the change of the modulation amplitude per decade. Standard errors are quasi-Poisson; one-sided p-values come from 2000 permutations of the year blocks; Benjamini–Hochberg (BH) control (Benjamini and Hochberg 1995) is applied over regions; BH assumes independent or positively dependent p-values, which is plausible for regions in different networks, and the more conservative correction for arbitrary dependence would only raise the q-values. F2 replaces the harmonic by the GRACE unloading U (lagged by three months): $\log \mu = \alpha_y + (\gamma_0 + \gamma_1 \tau) U$, with γ_0 in %/kPa. F3 compares rain and load as predictors with QAIC.

3.3 Criticality indicators

For two-year bins we computed rate, b-value (Schorlemmer et al. 2005), clustering and critical-slowing-down (CSD) indicators (variance, AR(1), skewness, kurtosis, recovery rate). Trends were tested with Kendall's τ against 10 000 block permutations, CSD indicators against 2000 IAAFT surrogates (Schreiber and Schmitz 1996), and glaciated-versus-control contrasts by group; all families are BH-corrected.

3.4 Two-stage ETAS with a time-varying background (H2)

A temporal ETAS model (Ogata 1988) with an Omori kernel truncated at $t_{\max} = 1\,\mathrm{yr}$ is first fitted with a stationary background (stage 1; K, α, c, p). With these triggering parameters fixed, stage 2 estimates one background rate μ_y per year (inspired by, but not a re-implementation of, Kumazawa and Ogata (2013)). μ_y is the *ETAS residual background*: everything the stationary temporal model does not explain as triggering, including catalogue changes, swarms and slow transients. It

is not an identified forcing rate. H2.1 is a difference-in-differences test of the log-linear trend of μ_y 1995–2025 between a glaciated region and its in-network control (primary pair SE Alaska / Interior Alaska; one-sided, BH over three pairs), with standard errors inflated by a frozen factor 1.25 from a null calibration. H2.2 classifies the ratio of μ in 2020–2025 versus 2008–2019 together with the total-rate ratio into "masked background rise" (the deceptive-quiet pattern), "background rise (not masked)", "background decrease (genuine quiet)" or "no resolvable change".

3.5 Tidal phase tests (H3)

For each declustered event the phase of the degree-2 body-tide areal strain is computed following Tanaka et al. (2002) (Love numbers $h = 0.6114$, $l = 0.0832$ as frozen; extremum search ± 14 h). H3.1 is a Schuster test of the phase distribution with amplitude $a = 2R/N$; H3.2 compares the b-value in the tensile and compressive half-cycles (cf. Tan et al. 2019). MDEs at 80 % power are reported. Exploratory: ocean-load strain with SPOTL/TPXO 7.2 (Agnew 1997; Egbert and Erofeeva 2002), eight constituents, at 0.5° cell centroids, following the argument of Tsuruoka et al. (1995) that ocean loading changes the local phase. Because the total-strain waveform is time-asymmetric, the Tanaka phase is not uniform at random times; with ocean loading the test is therefore made against the empirical phase distribution at $3N$ random times per cell.

3.6 Viscoelastic lag model (F5)

Surface unloading of wavelength λ on an elastic plate (effective thickness T_e, rigidity D) over a viscous half-space (viscosity η) relaxes with

$$\tau = \frac{4\pi\eta}{\rho_m g \lambda}\, C, \qquad C = \frac{1}{1 + D k^4/(\rho_m g)}, \quad k = 2\pi/\lambda, \tag{2}$$

where C is the flexurally compensated (delayed) share and $1 - C$ responds instantly (Turcotte and Schubert 2002). Viscosities are taken from regional studies (Larsen et al. 2004; Larsen et al. 2005; Lange et al. 2014; Pagli et al. 2007; Khan et al. 2016; Barletta et al. 2018; Mey et al. 2016); M4 [EXPL] adds a sensitivity run informed by Sasgen et al. (2026). Seismicity responds by Dieterich rate-and-state (Dieterich 1994). For 126 scenarios (regions × viscosity × wavelength × coupling κ × background stressing rate $\dot{S}_0$) we computed rate ratios and the first year in which a log-linear trend from November 2026 would exceed the detection threshold (2.487×SE; one-sided 5 %, power 80 %).

3.7 Tipping scenarios and deceptive-quiet model (A, B)

Load-rate trajectories $\dot{L}(t)$ are expressed as multipliers of today's GRACE rate: continuation; continued GRACE acceleration; exponential paths whose 2015–2100 integral matches the SSP5-8.5 median and 95th percentile of Greenland and the global-glacier median of Edwards et al. (2021) (derived by us, using Hugonnet et al. 2021); and a tenfold acceleration of Amundsen Sea loss after 2060 (or 2040) after DeConto et al. (2021). The GIA stressing rate is

$$\dot{S}_{\mathrm{GIA}}(t) = \kappa\Big[(1 - C)\,\dot{L}(t) + C \int \dot{L}(s)\, \frac{e^{-(t-s)/\tau}}{\tau}\, \mathrm{d}s\Big], \tag{3}$$

with unconstrained coupling $\kappa \in \{0.3, 1\}$ and $\dot{S}_0 \in \{10, 30\}$ kPa/yr. Promoted faults follow $R/R_{2025} = (\dot{S}_0 + \dot{S}_{\mathrm{GIA}})/(\dot{S}_0 + \dot{S}_{\mathrm{GIA},2025})$. In model B, a fraction $f = 0.6$ of faults lies in a stress shadow (Harris 1998; Hampel et al. 2010) and follows the exact Dieterich equation with $A\sigma = 1.5$ kPa; the catalogue sees the mixture. A net quiet requires $f > 0.5$; this is an assumption. In the same framework the seasonal modulation of promoted faults scales as $1/(\dot{S}_0 + \dot{S}_{\mathrm{GIA}})$ for periods much longer than the aftershock time.

3.8 Quiescence and monitoring indicators

Present-day quiescence is assessed with a temporal ETAS fit 1995–2014.5 extrapolated to 2016–2025 (Ogata 1992) and with Habermann's z for declustered yearly counts 2020–2025 versus 2008–2019 (Wyss and Habermann 1988; Habermann 1991). Monitoring indicators were frozen before any new data: M1 (GRACE normalised acceleration of loss $\xi = -A/|v| > 0.02$/yr with lower 95 % bound > 0), M2 (GNSS uplift acceleration $z > 3$ with the sign of M1), M3 (drop of seasonal modulation), M4 (b-value change), M5 (quiet: Habermann $z \leq -2$); in v0.2, M6 (single-region background rise, $z \geq 2$) and M7 (tidal Schuster $p < 0.01$, $N \geq 200$).

3.9 Global baseline [EXPL]

ComCat $M \geq 6$ and $M \geq 7$ counts 1975 to 10 October 2026, all and Gardner–Knopoff declustered, with Poisson GLM trends (quasi-Poisson intervals where overdispersed) (cf. Shearer and Stark 2012); declustered $M \geq 5$ rate ratios 2010–2024 versus 1995–2009 in six glaciated and five tectonically active control boxes with exact conditional intervals and an MDE of $\exp\{(1.96 + 0.84)\sqrt{2/\bar{n}}\}$; GVP confirmed eruption starts by decade. Boxes, periods and thresholds were chosen in a first pass and are not preregistered.

4 Preregistration and confirmatory design

Why only C1 and C2 are confirmatory. F1–F3 and H2/H3 were fixed before they were run, but on data that had partly been examined (rates, seasonality). They are pre-specified but data-touched and cannot serve as independent confirmation. C1 and C2 test data that did not exist when they were frozen.

C1 (frozen with v0.1.0). For the primary region SE Alaska, the seasonal modulation b_{new} of background events from 2026-11-01 is compared with the frozen baseline $b_{\text{ref}} = \beta_0 = 0.104 \pm 0.024$ (1990–2025) by $z = (b_{\text{new}} - b_{\text{ref}})/\sqrt{se^2_{\text{new}} + se^2_{\text{ref}}}$, one-sided. Looks: data to 2029-10-31 ($\alpha = 0.01$) and 2036-10-31 ($\alpha = 0.04$). Mc, declustering threshold, phase, baseline and look dates are frozen. With about 535 and 1784 expected background events the MDE is 0.24 and 0.12; if the retrospective point estimate $\beta_1 = 0.035$/decade were true, the power would be only about 8 % and 53 % (`confirmatory_power.csv`). A null result in 2036 would therefore not be strong evidence against an early-stage effect; this was stated before any data.

C2 (frozen with v0.2). $D = [\log \mu_{\text{new}} - \log \mu_{\text{ref}}]_{\text{SE Alaska}} - [\log \mu_{\text{new}} - \log \mu_{\text{ref}}]_{\text{Interior}}$, one-sided $D > 0$, with μ_{new} the window ML estimate from 2027-01-01 under frozen triggering parameters. Baselines (data to 2025-12-31): SE Alaska $\log \mu_{\text{ref}} = 4.05$, $\sigma_y = 0.36$; Interior Alaska 4.51, 0.16. Windows $[2027.0, 2030.0)$ with $\alpha = 0.01$ and $[2027.0, 2037.0)$ with $\alpha = 0.04$. The large year-to-year scatter in SE Alaska means that the first look can only detect roughly a doubling of the background.

Addendum (2026-10-10). After the review round (Section 6) a dated addendum A1–A7 was frozen: A1 C2 uses its JSON formula without the H2 factor 1.25; A2 the baseline functional stays; A3 look dates as above; A4 operating rules (archived catalogue snapshots with SHA-256, Mc homogeneity report, network histories, code pinned to v0.2.0); A5 erratum E1, factor 1.25 retained; A6 M7 uses IERS 2010 Love numbers (Petit and Luzum 2010) and a ±20 h search; A7 reading notes.

Integrity. Table 3 lists the frozen files. The v0.2 preregistration commit (`fa4a077`, 2026-10-10 14:50:44 CEST) precedes the v0.2 results commit (`250457c`); commit order shows the sequence in

Table 3: Frozen files and their SHA-256.

File (`results/`) and SHA-256	Content
`frozen_confirmatory_parameters.json`	C1 (v0.1.0)
`5bccd4cab0ba2d0143c4736c44355b24b6eb6356ca82ba0419fff22c71b4b3cb`	
`frozen_monitoring_indicators.json`	M1–M5 (v0.1.0)
`8ae57131325eda3ffb855f0437a649eb5adb0a32ae4e2fd0ec35f515cb87fb0f`	
`frozen_v02_preregistration.json`	H2, H3, C2, M6, M7 (v0.2)
`af7a5d02b6d81e37ba989058968f913e5e821876f98926dcca924560d3d927d3`	
`frozen_v02_baselines.json`	C2 baselines
`e2fc568c93622c97607c122c210ed5fcccace0d2440b8cd420fddd931be9f401`	
`frozen_v02_addendum_2026-10-10.json`	Addendum A1–A7
`bc2d2af32192ae5f31312c0609ee850223f23470637350df9795ee18f6e7ff49`	

the repository but cannot by itself prove that no analysis ran earlier. Releases are archived on Zenodo: v0.1.0 (Römer 2026b), v0.2.0 (Römer 2026a); concept DOI 10.5281/zenodo.23281090.

5 Results

5.1 Global baseline [EXPL]

Global $M \geq 6$ counts rose from 1975 to the mid-1980s and have not risen since (Fig. 1a). The early rise (+0.49 %/yr from 1975) coincides with the transition to moment magnitudes before about 1983 and is a catalogue effect. From 1990 the trends are −0.46 %/yr (95 % CI −0.70 to −0.21; quasi-Poisson −0.90 to −0.01) for all $M \geq 6$, −0.31 %/yr (−0.62 to 0.00; quasi −0.65 to +0.03) declustered, and −0.21 %/yr (−0.99 to +0.57) for $M \geq 7$. Allowing for overdispersion, a rise of the declustered $M \geq 6$ rate of more than about 0.03 %/yr since 1990 is incompatible with these data, in line with Shearer and Stark (2012). GVP confirmed eruption starts per year by decade show no upward trend (1950s 35.7; 2000s 39.2; 2010s 36.7; 2020–2025 30.3).

In the glaciated boxes, declustered $M \geq 5$ counts went from 60 to 57 (ratio 0.95, 0.65–1.39; MDE 1.68), in the controls from 2679 to 2814 (1.05, 1.00–1.11) (Table 4, Fig. 1b). Without declustering the glaciated boxes rose from 83 to 151 events (1.82), almost entirely due to swarms and sequences, above all the 2014–2015 Bárðarbunga rifting (Sigmundsson et al. 2015). Single glaciated boxes can only detect ratios of about 2.4–3.5.

5.2 Seasonal modulation in Southeast Alaska (F1, F2)

SE Alaska has a strong and robust seasonal modulation: Schuster amplitude 0.112 ($p < 10^{-6}$, $n = 5164$), peak in mid-November, about one to two months after the maximum of GRACE unloading (load maximum in April) (Fig. 2a). The load sensitivity is $\gamma_0 = 4.04 \pm 1.09$ %/kPa ($p = 10^{-4}$), close to the 3.5 %/kPa reported by Johnson et al. (2020). The in-network control Interior Alaska shows no seasonality (amplitude 0.023, $p = 0.39$). Rain and load are correlated in SE Alaska ($r = 0.68$) and cannot be separated (F3).

The amplitude is not growing significantly (Fig. 2b, Fig. 3): $\beta_1 = +0.035 \pm 0.023$ per decade (95 % CI −0.010 to +0.080; one-sided Wald $p = 0.064$, permutation $p = 0.10$, BH $q = 0.38$), and the load sensitivity declines insignificantly ($\gamma_1 = -1.30 \pm 1.13$ %/kPa per decade, permutation $p = 0.94$). Greenland has the largest growth estimate ($\beta_1 = +0.146 \pm 0.088$, permutation $p = 0.0055$, $q = 0.083$) on a catalogue that densified strongly after 2008; Iceland is negative (-0.096 ± 0.063). Among the controls, Japan's γ_1 is nominally significant ($p_{\mathrm{perm}} = 0.021$, $q = 0.29$). The inverse-variance mean of β_1 over the four national mountain regions is 0.033 ± 0.015 ($z = 2.25$), but no glaciated-minus-control contrast in the criticality family is significant (Section 5.3). With GK declustering the SE Alaska estimate is $+0.050 \pm 0.023$ (permutation $p = 0.01$, uncorrected) [EXPL].




Figure 1: Exploratory global baseline. (a) ComCat global counts per year; shading marks the period with few moment magnitudes. (b) Declustered $M \geq 5$ rate ratio 2010–2024 versus 1995–2009 with exact 95 % intervals; diamonds are pooled groups. West Antarctica (0 and 1 events) is omitted. Data: `results/global/`.

Table 4: Exploratory global and regional baseline. Data: `results/global/global_trends.csv` and `ice_vs_control_rate_ratios.csv`.

Quantity	Estimate	95 % interval	Note
Trend $M \geq 6$ all, 1990–2026 (%/yr)	−0.46	−0.70 to −0.21	quasi: −0.90 to −0.01
Trend $M \geq 6$ declustered (%/yr)	−0.31	−0.62 to 0.00	quasi: −0.65 to +0.03
Trend $M \geq 7$ all (%/yr)	−0.21	−0.99 to +0.57	
Glaciated pooled, $M \geq 5$ decl. (60 → 57)	0.95	0.65–1.39	MDE 1.68
Control pooled, $M \geq 5$ decl. (2679 → 2814)	1.05	1.00–1.11	MDE 1.08
Glaciated pooled, all events (83 → 151)	1.82	1.38–2.41	swarms, Bárðarbunga
GVP eruption starts, controls (223 → 235)	1.05	0.87–1.27	

5.3 Criticality indicators

Of 374 trend tests, 63 had $p < 0.05$ and 10 survived BH; of 170 CSD tests 18 and 6; of 180 glaciated-versus-control group tests 2 and none (smallest $q = 0.76$). The 16 BH hits are network and completeness effects (Mc trends, magnitude-type changes), b-value changes and variance increases that accompany rate increases or network densification (Greenland after about 2008); four of the ten trend hits are in control regions. Rising autocorrelation or falling recovery rate, the classical CSD signatures, are nowhere significant. Criticality is not excluded, but these indicators have little power with 18 two-year bins.

5.4 ETAS background: H2.1 positive but not robust

In stage 1 the Omori exponent ran to its lower bound ($p = 1.0001$) in 7 of 8 regions, with branching ratios within the one-year kernel of 0.6–0.9. This is a serious diagnostic sign: model misfit (spatial heterogeneity, swarms, catalogue changes) is absorbed by a long tail, and long-range triggering beyond one year ends up in the background. Trend estimates are therefore model dependent.

The preregistered decision rule for H2.1 fired: SE Alaska's background trend ($+0.326 \pm 0.114$ per decade) exceeds Interior Alaska's (-0.079 ± 0.079) by $+0.405 \pm 0.138$ ($z = 2.93$, one-sided $p = 0.0017$, BH 0.005; Table 6). The effect, however, sits almost entirely in 1997–2002, when catalogue counts in both Alaska regions were very low and yearly M_c exceeded M_c^{ref} (Fig. 4a).




Figure 2: Seasonality in SE Alaska. (a) Monthly climatology of declustered events, SE Alaska versus the Interior Alaska control. (b) Yearly modulation amplitude b_y with the F1 fit. Data: `seasonal_climatology.csv`, `F1_per_year_modulation.csv`, `F1_harmonic_growth.csv`.

Table 5: F1 and F2 for the national-tier regions (pre-specified, retrospective). β_1 per decade; γ_0 in %/kPa; γ_1 in %/kPa per decade; p_{perm} one-sided; q BH over all regions.

Region	Schuster amp.	$\beta_1 \pm$SE	p_{perm}	q	$\gamma_0 \pm$SE	$\gamma_1 \pm$SE
SE Alaska	0.112	$+0.035 \pm 0.023$	0.100	0.38	4.0 ± 1.1	-1.3 ± 1.1
Alps	0.005	$+0.058 \pm 0.039$	0.079	0.38	-5.5 ± 8.5	-2.5 ± 9.5
NZ Southern Alps	0.035	$+0.038 \pm 0.023$	0.058	0.38	8.1 ± 5.3	$+7.1 \pm 6.2$
Iceland, Vatnajökull	0.299	-0.096 ± 0.063	0.861	0.86	17.6 ± 6.3	-10.6 ± 7.1
Greenland	0.117	$+0.146 \pm 0.088$	0.006	0.08	-4.5 ± 15.2	$+1.7 \pm 17.3$
Interior Alaska	0.023	-0.005 ± 0.020	0.559	0.76	4.3 ± 4.1	$+1.1 \pm 4.3$
N Switzerland	0.071	$+0.092 \pm 0.073$	0.132	0.40	—	—
NZ Wellington	0.001	-0.008 ± 0.015	0.709	0.82	4.0 ± 4.6	$+3.9 \pm 5.3$

Exploratory variants (Fig. 4b): starting in 2003 the difference is $+0.11 \pm 0.16$; with a three-year triggering horizon $+0.23 \pm 0.21$; without 2025 $+0.41 \pm 0.15$; with a joint estimator $+0.43 \pm 0.13$ (from 2003 $+0.16$); with free Omori p $+0.39 \pm 0.15$. Network covariates explain part of the trend: with FDSN station counts of all networks $+0.25 \pm 0.13$, permanent networks only $+0.38 \pm 0.10$, with ΔM_c $+0.39 \pm 0.09$, complete years only $+0.37 \pm 0.10$ (OLS, no factor). Station metadata before 2000 are incomplete, so a network artefact is neither confirmed nor excluded. Together with the collapse from 2003 and the time-varying Mc, the result is better explained by a catalogue or model artefact than by a glacially driven background trend; H2.1 provides no evidence for a glacial contribution, though it does not exclude one. The secondary pairs are not significant (NZ $+0.095$, $p = 0.19$; Alps -0.350).

5.5 Masked background rise: H2.2

No region meets the predefined "masked background rise" criterion (Table 6). SE Alaska shows an unmasked rise (1.41) and N Switzerland a rise (1.94). The NZ Southern Alps quiet found in v0.1.0




Figure 3: Pre-specified retrospective growth tests in the national-tier regions (95 % intervals; blue glaciated, orange controls). (a) F1 growth of seasonal amplitude; (b) F2 load sensitivity; (c) F2 growth of load sensitivity. N Switzerland has no GRACE signal and is omitted in (b, c). Data: `F1_harmonic_growth.csv`, `F2_load_sensitivity_growth.csv`.

Table 6: ETAS residual background (`m2_etas_background.csv`, `m2_H21_did.csv`). Trend of $\log \mu$ per decade (SE $\times 1.25$); μ ratio 2020–2025 / 2008–2019 with 95 % interval; H2.2 class. Iceland and Greenland are descriptive.

Region	Trend / decade	μ ratio (95 %)	H2.2 class	Total ratio
SE Alaska	$+0.326 \pm 0.114$	1.41 (1.07–1.85)	rise, not masked	2.33
Interior Alaska	-0.079 ± 0.079	0.87 (0.71–1.07)	no resolvable change	0.91
NZ Southern Alps	-0.326 ± 0.059	0.62 (0.45–0.86)	decrease (genuine quiet)	0.32
NZ Wellington	-0.421 ± 0.091	0.99 (0.67–1.47)	no resolvable change	0.71
Alps	-0.350 ± 0.089	1.22 (0.96–1.56)	no resolvable change	1.30
N Switzerland	$+0.001 \pm 0.097$	1.94 (1.29–2.91)	rise, not masked	2.04
Iceland, Vatnajökull	$+0.194 \pm 0.144$	1.56 (0.85–2.86)	no resolvable change	0.19
Greenland	$+1.047 \pm 0.207$	1.12 (0.75–1.67)	no resolvable change	1.68
H2.1 primary: SE Alaska − Interior: $+0.405 \pm 0.138$, $z = 2.93$, $p_{\text{one}} = 0.0017$, $p_{\text{BH}} = 0.005$ [PREREG]				

is a genuine decrease of the residual background (0.62), but it disappears with EMR completeness ($M_c = 3.2$) and coincides with a joint step of both NZ regions in 2012, so a network effect is likely. Iceland comes closest to the pattern (total rate −81 %, μ +56 %), but the interval includes 1 and ETAS does not describe the swarms there. Rough 80 %-power thresholds are μ ratios of 1.3–2.4; the classification is descriptive and is mildly informative only against large masked rises in well-observed regions.

5.6 Tides: H3

Seismicity does not correlate significantly with the phase of the body-tide areal strain anywhere (Table 7, Fig. 5). In SE Alaska the amplitude is $a = 0.042$ (Schuster $p = 0.14$, BH 0.54); a modulation of ≥ 6.8 % relative to this phase would have been detected with 80 % power, and a b-value difference of ≥ 0.07. The smallest BH q is 0.10 (Greenland, $\Delta b = -0.195$, one-sided $p = 0.021$). Widening the extremum search to ±20 h changes no conclusion [EXPL].

[EXPL] Ocean loading is comparable to or larger than the body tide in SE Alaska (RMS 19.9 versus 14.6 nstrain) and Greenland (18.8 versus 11.9), and moves 20 % and 42 % of events into the other half-cycle. Tested against random times, the SE Alaska p goes from 0.048 (body tide) to 0.081 (body plus ocean load); the Greenland b-value effect disappears (p 0.023 → 0.58); the




Figure 4: H2.1. (a) Yearly ETAS residual background μ_y with 95 % band (SE inflated by the frozen factor 1.25); shading marks the low-count years before 2003. (b) Preregistered difference-in-differences of the background trend (black) and exploratory variants (grey; squares: OLS with station-count covariates). Data: `results/v02/m2_*.csv`, `h21_network_models.csv`.

Table 7: H3 body-tide tests [PREREG] (`m3_tides.csv`); last column exploratory ocean loading tested against random times [EXPL] (`m3_tides_ocean_load.csv`).

Region	N	a	Schuster p	q	MDE a	Δb (p_{one})	p body+ocean
SE Alaska	4517	0.042	0.14	0.54	0.068	+0.023 (0.79)	0.081
NZ Southern Alps	3665	0.006	0.97	0.99	0.072	+0.005 (0.55)	0.85
Alps	1072	0.007	0.99	0.99	0.141	+0.097 (0.86)	0.87
Iceland, Vatnajökull	1227	0.071	0.22	0.54	0.126	−0.007 (0.46)	0.86
Greenland	496	0.058	0.66	0.99	0.194	−0.195 (0.021)	0.81

smallest BH q with loading is 0.38. Fault-resolved Coulomb stress is not modelled, so the null applies to areal-strain phase only. The body-tide stresses involved are small: an areal-strain amplitude of about 15 nstrain corresponds to roughly 1 kPa or less for typical crustal moduli, whereas at Axial Seamount, with tidal stresses of ±20 kPa, b varied with stress only above about 5 kPa (Tan et al. 2019). A null H3.2 was therefore the expected outcome rather than a strong constraint.

5.7 Lag model and tipping scenarios

Relaxation times range from 23–117 yr (Patagonia) and 64–170 yr (SE Alaska) to 150–380 yr (Iceland), 1000–2000 yr (Alps) and about 4500 yr (Fennoscandia reference). Narrow mountain loads are mostly answered elastically ($C = 0.11$ in SE Alaska), broad ice-sheet loads mostly viscously ($C = 0.94$ in SE Greenland). At constant present-day unloading, the viscously delayed rate increase is at most +19 % by 2036 and +27 % by 2100 even at $\kappa = 1$ (+5 % or less with GRACE-box loads). A trend would be detectable in only 8 of 126 scenarios, the earliest around 2155.

Accelerating loss shortens this to decades (Table 8, Fig. 6). The earliest detectable increase is in SE Alaska under the SSP5-8.5 glacier path with glacier-surface loads (rate on promoted faults 1.21–2.35 by 2100, detection from 2045.8 in all four κ, $\dot{S}_0$ combinations). Amundsen paths become detectable around 2065–2079 only with a hypothetical dense network; SE Greenland stays within noise until about 2100. In the same framework the seasonal modulation *falls* in all accelerating




Figure 5: Tidal-phase modulation. (a) Amplitude a for the preregistered body-tide phase (dots) and the exploratory body-plus-ocean-load phase (triangles), with the 80 %-power MDE of the body-tide test (bars). (b) Uncorrected significance; the ocean-load test is against random times. Data: `results/v02/m3_tides*.csv`.

Table 8: Tipping scenarios (`tipping_scenarios.csv`). Rate of promoted faults relative to 2025 (range over $\kappa \in \{0.3, 1\}$, $\dot{S}_0 \in \{10, 30\}$ kPa/yr) and earliest detection of an increase with present catalogue rates. Detection counts refer to the four promoted-fault runs ($f = 0$). Model calculations, not forecasts.

Region / path	2050	2100	Earliest detection (runs)
SE Greenland, Edwards median	1.00–1.03	1.01–1.10	2158 (1 of 4)
SE Greenland, Edwards 95th pct.	1.00–1.04	1.02–1.24	2102 (3 of 4)
Amundsen, DeConto 3 °C (from 2060)*	1.00–1.02	1.10–1.89	2079 (4 of 4)
Amundsen, DeConto early (from 2040)*	1.01–1.06	1.11–1.96	2065 (4 of 4)
SE Alaska, glaciers SSP5-8.5 (surface)	1.04–1.26	1.21–2.35	2046 (4 of 4)
SE Alaska, glaciers SSP5-8.5 (GRACE box)	1.01–1.09	1.05–1.46	2061 (4 of 4)
Iceland, glaciers SSP5-8.5 (surface)	1.03–1.19	1.13–1.99	2070 (4 of 4)

* hypothetical network with 100 events/yr; West Antarctica has no usable catalogue today.

paths, to 0.43–0.97 of today by 2100, but this fall reaches at most $z = 1.2$ with present catalogues. M4 [EXPL]: viscosities and plate thicknesses informed by Sasgen et al. (2026) hardly move SE Alaska (earliest detection 2045.8 → 2047.3), because in this model the instantaneous share $1 - C$ dominates; for Iceland τ shrinks from about 300 to 10 yr and detection moves from 2070.3 to 2062.3.

5.8 Deceptive quiet before acceleration

With 60 % of faults in a stress shadow, a quiet before the increase follows from the model (Fig. 6, bottom). It begins 7–59 years after the onset of acceleration in the Amundsen, SE Alaska and Iceland paths (45–93 years for the SE Greenland 95th-percentile path) but is shallow: the minimum catalogue rate is 0.84–0.96 of 2025 (4–16 %). With present catalogues it would become statistically detectable at the earliest around 2083; in the steepest paths (SE Alaska) it reverses (2053–2061) before it could be detected. Geodetic indicators respond directly to the load and would lead any seismic quiet.

Today, no region meets the frozen quiet rule except the NZ Southern Alps (Habermann $z = -2.5$, 109 → 89 events/yr), where the Wellington control in the same network also declines and the classification depends on M_c. SE Alaska shows activation ($z = +3.1$, 161 → 194/yr). Iceland's ETAS-only signal coincides with a magnitude shift and the decay of the Bárðarbunga episode (Habermann 1991; Sigmundsson et al. 2015); on the load side Vatnajökull loss is slowing,




Figure 6: Tipping (top) and deceptive-quiet (bottom) scenarios for $\kappa = 1$, $\dot{S}_0 = 10\,\mathrm{kPa/yr}$. Top: rate of promoted faults relative to 2025. Bottom: catalogue rate with 60 % of faults in a stress shadow; the dashed line marks −2 %. Vertical dotted line: 2025. Data: `tipping_curves.csv`.

which corresponds to genuine relaxation rather than a deceptive quiet.

5.9 Present-day precursors: GRACE and GNSS

Only Patagonia exceeds the M1 threshold: $\xi = +0.039 \pm 0.007/\mathrm{yr}$, loss rising from 19 to 34 Gt/yr between 2002–2013 and 2014–2026 (Fig. 7a). Patagonia has no usable catalogue (ComCat is sparse and the relocated Chilean catalogue ends at 45.98°S). Greenland ($\xi = -0.019 \pm 0.007$) and Vatnajökull (-0.022 ± 0.007) have been losing mass more slowly since about 2014 (cf. Bevis et al. 2019); in the Amundsen Sea no significant acceleration is measurable (cf. Velicogna et al. 2020). Four GNSS stations carry an M2 flag (Fig. 7b): TOMO (41.7 mm/yr, acceleration $+2.31\,\mathrm{mm/yr^2}$, $z = 8.0$, consistent with Barletta et al. 2018), BACK ($z = 4.9$), AB50 ($z = 6.4$) and KELY ($z = 8.3$). KELY is discordant with its slowing GRACE box and is reported, not interpreted, by rule.




Figure 7: Monitoring precursors. (a) Normalised acceleration of GRACE mass loss ξ with 95 % intervals and the frozen M1 threshold. (b) GNSS uplift acceleration z (red: M2 flag; orange: control station FAIR). Data: `precursors_grace_acceleration.csv`, `precursors_gnss_uplift.csv`.

6 Review process and errata

Version 0.2 was reviewed in one round by several AI systems, each with a written brief (`docs/REVIEW_PROMPTS.md`): a "Deep Research" scientific review (DR), a preregistration audit by Grok (GR), a full code review with independent scripts and simulations by Claude Code (CC), a pull-request comment by Codex (CX), a narrow literature check by Gemini (GE; used as an idea source only) and the ETAS likelihood and simulator checks by Grok Bot (GB). Raw reviews are archived in `docs/reviews/raw/` (except GE), and all 41 points with verdicts and actions are listed in `docs/reviews/REVIEW_RESPONSE_v0.2.md`. New literature claims from reviews were checked against primary sources before adoption; one GE objection (Sasgen values "not found") was rejected after checking the paper. The v0.2.1 draft of this paper was reviewed once more by Gemini (idea source only); the point-by-point response is in `docs/reviews/GEMINI_PAPER_REVIEW_RESPONSE.md`.

Erratum E1. CX and CC independently found that the null calibration of the H2 standard-error factor 1.25 used a half-open window function with the wrong windows (2008–2018 versus 2020–2024) and a geometry without the five-year lead-in. Re-run in production geometry (CC and GB), the factor is sufficient for the trend test; for the window ratio a factor of about 1.2–1.4 would be needed. In the repository's calibration runs (`calibration_summary.csv`) the trend sd(z) is 0.96–1.07 (90 % upper bounds 1.07–1.22) and the window-ratio sd(z) 1.18–1.24 (upper bounds 1.27–1.35); the addendum quotes up to 1.30 and 1.42 including the reviewers' own runs, which are not archived. No published decision changes below a factor of 1.56. The frozen value stays (A5); the H2.2 intervals should be read as somewhat too narrow.

Other corrections. Simulator fixes (division by zero at $p = 1$; magnitude discretisation that gave $b \approx 0.90$ instead of 1, now 0.997) affected only simulations; the recovery study was re-run with qualitatively unchanged results. The C2 SE factor ambiguity was clarified (A1), look dates were made robust (A3), and the origin of the frozen Love numbers was traced to SPOTL's body-tide values. Wording was sharpened where DR showed that interpretations went beyond the estimand: μ_y is a residual background; H3 tests the body-tide areal-strain phase only; the lag model is a thin-plate/half-space sensitivity, not a reproduction of the Sasgen et al. (2026) rheology. After v0.2.0 the global baseline script was made deterministic for magnitude ties, which moved the

declustered $M \geq 6$ trend from -0.29 to -0.31 %/yr and the pooled glaciated ratio from 0.93 to 0.95 [EXPL].

7 Discussion

What the data exclude. At the global scale, a rise of the declustered $M \geq 6$ rate faster than about 0.03 %/yr since 1990 is incompatible with ComCat. Pooled over glaciated boxes, a rise of the declustered $M \geq 5$ rate by more than about 40 % between 1995–2009 and 2010–2024 is unlikely (upper bound 1.39). In SE Alaska a growth of the seasonal amplitude faster than about 0.08 per decade (upper 95 % bound; today's amplitude is about 0.10) and a body-tide modulation larger than about 7 % would probably have been seen. No region shows a large masked background rise.

What remains open. Moderate changes are not excluded anywhere. Single glaciated regions can only resolve rate ratios of 2.4–3.5, Iceland and Greenland catalogues are too heterogeneous for background trends, Patagonia (the one region with accelerating loss) and West Antarctica lack usable catalogues, and ocean-load and fault-resolved stresses are only partly modelled. The positive but non-robust H2.1 result shows how a preregistered test without a start-year check can hinge on a network phase; C2 starts in a homogeneous network period and carries explicit homogeneity rules. Natural next steps, each requiring new data or models, are fault-resolved Coulomb stresses from three-dimensional viscoelastic models, time-dependent detection models for completeness, and waveform-based discrimination of non-tectonic sources; the last is the subject of the separate v0.3 module.

An early-stage hypothesis. Taken together, the evidence is compatible with a hypothesis in an early stage: present-day ice loss may change the seismic response of glaciated regions, but if it does, the change is currently below what the catalogues can resolve, and the scenario calculations suggest it will remain so for decades unless loss accelerates strongly. We do not claim a glacially driven change, and we do not claim its absence.

Rate-and-state implication. A growing seasonal modulation would not be the generic expectation of accelerated unloading. In Dieterich's model (Dieterich 1994), for loading periods much longer than the aftershock duration the relative modulation scales as $\Delta S/(T\dot{S})$, so additional GIA stressing *lowers* it (our scenarios: to 0.43–0.97 of today by 2100). If the modulation grew, more likely explanations would be a falling effective normal stress (pore pressure, as suggested for Mont Blanc by Simon et al. 2025) or faults approaching failure, as discussed for Parkfield by Sirorattanakul and Avouac (2026). This is why F1 alone could never show causality, and why C1 tests modulation growth as a sign of changing fault conditions rather than of unloading per se.

8 Limitations

- Retrospective tests use data examined before the rules were fixed; only C1 and C2 are independent.
- Catalogues change networks, magnitude types and completeness; in-network controls do not protect against differential network changes. Station metadata before about 2000 are incomplete.
- The temporal ETAS model has no spatial kernel; p at its lower bound in 7 of 8 regions makes levels and trends model dependent; H2 SEs are conditional on stage-1 parameters (propagation adds about 1–2 %).
- GRACE resolution (about 300 km) smears narrow mountain loads; rain and load are not separable in SE Alaska. In low-viscosity regions, rapid rebound makes GRACE underestimate

the ice-loss trend (about 9 % in Alaska and one third in Iceland, 2002–2025; Sasgen et al. 2026); this scales the trend-based loads of the lag and tipping scenarios and of M1, and affects the seasonal load in F2 less.

- Catalogues are filtered by event type only. Unflagged icequakes, landslides or tremor (e.g. jökulhlaup tremor, Eibl et al. 2020) could contaminate counts, especially at small magnitudes in Iceland and Greenland; this was not quantified.
- MDEs and permutation tests treat declustered events as independent (F1 and C1 with the observed quasi-Poisson overdispersion); residual clustering would make the MDEs optimistic.
- Tidal tests use areal-strain phase at cell centroids; TPXO 7.2 resolves fjords poorly; no fault-resolved stress.
- The lag and tipping models are reduced (half-space plus plate, unconstrained coupling κ, assumed shadow fraction); their numbers are scenario properties, not forecasts.
- The global baseline is exploratory, uses rectangular boxes and coarse declustering; declustered counts carry method uncertainty beyond Poisson intervals.
- The analysis was carried out by one person with AI assistance; reviews were by AI systems, not by independent human experts.

9 Conclusions

1. There is no sign of a global or pooled glaciated increase of declustered seismicity, with upper bounds of about 0.03 %/yr (global $M \geq 6$) and a rate ratio of 1.39 (pooled glaciated $M \geq 5$).
2. SE Alaska is seasonally modulated (4 %/kPa), but the modulation is not demonstrably growing.
3. The one formally positive preregistered test (H2.1) is not robust and is most plausibly a catalogue artefact.
4. Tidal, quiescence and criticality tests are null, with stated MDEs.
5. Physical scenarios place detectable effects decades ahead and predict a shallow quiet before acceleration; geodetic indicators would lead.
6. Two prospective tests are frozen and will be evaluated in 2029 and 2036; their power is limited, and this was stated in advance.

Author contributions and use of AI

J. B. R. (independent researcher; certified IT specialist for application development with a background in web design) conceived the project, set the research questions and decision rules, approved every preregistration and addendum, and is responsible for all scientific claims. The work was developed using AI-assisted research methods as collaborative tools under direct human oversight: data retrieval, analysis code, figures, documentation and drafts of this manuscript were produced with an AI agent (Grok Bot) on the author's instructions; the review round used further AI systems (Deep Research, Grok, Claude Code, Codex; Gemini as an idea source only), whose raw reports are archived and whose points were checked against code, data and primary literature before adoption. The collaboration between the author and these AI systems runs under the informal working name "MOR collective", which is not a legal entity or institution. No AI system is an author. All scientific claims and methodological decisions are the responsibility of the human author.

Data and code availability

Code, frozen files, result tables and documentation: GitHub https://github.com/GenesisAeon/glacial-seismic-sensitivity-utac; archived releases v0.1.0 (10.5281/zenodo.23281091) and v0.2.0 (10.5281/zenodo.23286492); concept DOI 10.5281/zenodo.23281090; Python package https://pypi.org/project/glacial-seismic-sensitivity-utac/. Figures in this paper are re-rendered from the CSV files in `results/` by `paper/figures/make_paper_figures.py`. All input data are public (Section 2); global snapshots are archived in `results/global/snapshots/`. The project is registered as P133 (KLIMA-KRYO-13) in the GenesisAeon research programme. The suffix "utac" in the repository and package names is a programme label only; no method or result in this paper uses any model of that name (`DISCLAIMER.md`).

Funding

This work received no external funding; the author is independent and self-funded.

Competing interests

The author declares no competing interests.

References

Ader, Thomas J. and Jean-Philippe Avouac (2013). "Detecting periodicities and declustering in earthquake catalogs using the Schuster spectrum, application to Himalayan seismicity". In: *Earth and Planetary Science Letters* 377–378, pp. 97–105. DOI: 10.1016/j.epsl.2013.06.032.

Agnew, Duncan C. (2012). *SPOTL: Some Programs for Ocean-Tide Loading*. https://igppweb.ucsd.edu/~agnew/Spotl/spotlmain.html.

Agnew, Duncan Carr (1997). "NLOADF: A program for computing ocean-tide loading". In: *Journal of Geophysical Research: Solid Earth* 102.B3, pp. 5109–5110. DOI: 10.1029/96JB03458.

Alaska Earthquake Center (2026). *Earthquake catalogue contributions to ComCat (`catalog=ak`)*. https://earthquake.alaska.edu; accessed via https://earthquake.usgs.gov/fdsnws/event/1/. Retrieved 2026-10-10.

Barletta, Valentina R., Michael Bevis, Benjamin E. Smith, Terry Wilson, Abel Brown, Andrea Bordoni, et al. (2018). "Observed rapid bedrock uplift in Amundsen Sea Embayment promotes ice-sheet stability". In: *Science* 360.6395, pp. 1335–1339. DOI: 10.1126/science.aao1447.

Benjamini, Yoav and Yosef Hochberg (1995). "Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing". In: *Journal of the Royal Statistical Society Series B: Statistical Methodology* 57.1, pp. 289–300. DOI: 10.1111/j.2517-6161.1995.tb02031.x.

Bettinelli, Pierre, Jean-Philippe Avouac, Mireille Flouzat, Laurent Bollinger, Guillaume Ramillien, Sudhir Rajaure, and Som Sapkota (2008). "Seasonal variations of seismicity and geodetic strain in the Himalaya induced by surface hydrology". In: *Earth and Planetary Science Letters* 266.3–4, pp. 332–344. DOI: 10.1016/j.epsl.2007.11.021.

Bevis, Michael, Christopher Harig, Shfaqat A. Khan, Abel Brown, Frederik J. Simons, Michael Willis, et al. (2019). "Accelerating changes in ice mass within Greenland, and the ice sheet's sensitivity to atmospheric forcing". In: *Proceedings of the National Academy of Sciences* 116.6, pp. 1934–1939. DOI: 10.1073/pnas.1806562116.

Blewitt, Geoffrey, William Hammond, and Corné Kreemer (2018). "Harnessing the GPS Data Explosion for Interdisciplinary Science". In: *Eos* 99. DOI: 10.1029/2018EO104623.

DeConto, Robert M., David Pollard, Richard B. Alley, Isabella Velicogna, Edward Gasson, Natalya Gomez, et al. (2021). "The Paris Climate Agreement and future sea-level rise from Antarctica". In: *Nature* 593.7857, pp. 83–89. DOI: 10.1038/s41586-021-03427-0.

Dieterich, James (1994). "A constitutive law for rate of earthquake production and its application to earthquake clustering". In: *Journal of Geophysical Research: Solid Earth* 99.B2, pp. 2601–2618. DOI: 10.1029/93JB02581.

EarthScope Consortium (2026). *FDSN station web service*. https://service.earthscope.org/fdsnws/station/1/. Retrieved 2026-10-10.

Edwards, Tamsin L., Sophie Nowicki, Ben Marzeion, Regine Hock, Heiko Goelzer, Hélène Seroussi, et al. (2021). "Projected land ice contributions to twenty-first-century sea level rise". In: *Nature* 593.7857, pp. 74–82. DOI: 10.1038/s41586-021-03302-y.

Egbert, Gary D. and Svetlana Y. Erofeeva (2002). "Efficient Inverse Modeling of Barotropic Ocean Tides". In: *Journal of Atmospheric and Oceanic Technology* 19.2, pp. 183–204. DOI: 10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2.

Eibl, Eva P. S., Christopher J. Bean, Bergur Einarsson, Finnur Pàlsson, and Kristin S. Vogfjörd (2020). "Seismic ground vibrations give advanced early-warning of subglacial floods". In: *Nature Communications* 11.1, p. 2504. DOI: 10.1038/s41467-020-15744-5.

Gardner, J. K. and L. Knopoff (1974). "Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?" In: *Bulletin of the Seismological Society of America* 64.5, pp. 1363–1367. DOI: 10.1785/BSSA0640051363.

Geological Survey of Denmark and Greenland (GEUS) (2026). *Greenland earthquake catalogue (GeoJSON)*. https://seismologi.geus.dk/geojson/GeoJSON-eqs.json. Retrieved 2026-10-10.

GeoNet (2026). *FDSN event web service, New Zealand*. https://service.geonet.org.nz/fdsnws/event/1/. Retrieved 2026-10-10.

Global Volcanism Program (2026). *Volcanoes of the World, WFS web service*. Smithsonian Institution, https://webservices.volcano.si.edu/geoserver/GVP-VOTW/ows. Retrieved 2026-10-10.

Habermann, Ray E. (1991). "Seismicity rate variations and systematic changes in magnitudes in teleseismic catalogs". In: *Tectonophysics* 193.4, pp. 277–289. DOI: 10.1016/0040-1951(91)90337-R.

Hampel, Andrea, Ralf Hetzel, and Georgios Maniatis (2010). "Response of faults to climate-driven changes in ice and water volumes on Earth's surface". In: *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 368.1919, pp. 2501–2517. DOI: 10.1098/rsta.2010.0031.

Harris, Ruth A. (1998). "Introduction to Special Section: Stress Triggers, Stress Shadows, and Implications for Seismic Hazard". In: *Journal of Geophysical Research: Solid Earth* 103.B10, pp. 24347–24358. DOI: 10.1029/98JB01576.

Hersbach, Hans, Bill Bell, Paul Berrisford, Shoji Hirahara, András Horányi, Joaquín Muñoz-Sabater, et al. (2020). "The ERA5 global reanalysis". In: *Quarterly Journal of the Royal Meteorological Society* 146.730, pp. 1999–2049. DOI: 10.1002/qj.3803.

Hugonnet, Romain, Robert McNabb, Etienne Berthier, Brian Menounos, Christopher Nuth, Luc Girod, et al. (2021). "Accelerated global glacier mass loss in the early twenty-first century". In: *Nature* 592.7856, pp. 726–731. DOI: 10.1038/s41586-021-03436-z.

Icelandic Meteorological Office (Veðurstofa Íslands) (2026). *Earthquake event API (`system=sil`)*. https://api.vedur.is/quakes/events. Retrieved 2026-10-10; CC BY 4.0.

Johnson, Christopher W., Yuning Fu, and Roland Bürgmann (2017). "Seasonal water storage, stress modulation, and California seismicity". In: *Science* 356.6343, pp. 1161–1164. DOI: 10.1126/science.aak9547.

Johnson, Christopher W., Yuning Fu, and Roland Bürgmann (2020). "Hydrospheric modulation of stress and seismicity on shallow faults in southern Alaska". In: *Earth and Planetary Science Letters* 530, p. 115904. DOI: 10.1016/j.epsl.2019.115904.

Johnston, Arch C. (1987). "Suppression of earthquakes by large continental ice sheets". In: *Nature* 330.6147, pp. 467–469. DOI: 10.1038/330467a0.

Khan, Shfaqat A., Ingo Sasgen, Michael Bevis, Tonie van Dam, Jonathan L. Bamber, John Wahr, et al. (2016). "Geodetic measurements reveal similarities between post–Last Glacial Maximum and present-day mass loss from the Greenland ice sheet". In: *Science Advances* 2.9, e1600931. DOI: 10.1126/sciadv.1600931.

Kumazawa, Takao and Yosihiko Ogata (2013). "Quantitative description of induced seismic activity before and after the 2011 Tohoku-Oki earthquake by nonstationary ETAS models". In: *Journal of Geophysical Research: Solid Earth* 118.12, pp. 6165–6182. DOI: 10.1002/2013JB010259.

Lagerbäck, Robert and Martin Sundh (2008). *Early Holocene faulting and paleoseismicity in northern Sweden*. Geological Survey of Sweden (SGU), Research Paper C836, https://resource.sgu.se/produkter/c/c836-rapport.pdf.

Lange, H., G. Casassa, E. R. Ivins, L. Schröder, M. Fritsche, A. Richter, A. Groh, and R. Dietrich (2014). "Observed crustal uplift near the Southern Patagonian Icefield constrains improved viscoelastic Earth models". In: *Geophysical Research Letters* 41.3, pp. 805–812. DOI: 10.1002/2013GL058419.

Larsen, Christopher F., Roman J. Motyka, Jeffrey T. Freymueller, Keith A. Echelmeyer, and Erik R. Ivins (2004). "Rapid uplift of southern Alaska caused by recent ice loss". In: *Geophysical Journal International* 158.3, pp. 1118–1133. DOI: 10.1111/j.1365-246X.2004.02356.x.

Larsen, Christopher F., Roman J. Motyka, Jeffrey T. Freymueller, Keith A. Echelmeyer, and Erik R. Ivins (2005). "Rapid viscoelastic uplift in southeast Alaska caused by post-Little Ice Age glacial retreat". In: *Earth and Planetary Science Letters* 237.3–4, pp. 548–560. DOI: 10.1016/j.epsl.2005.06.032.

Lindblom, Eva, Björn Lund, Ari Tryggvason, Marja Uski, Reynir Bödvarsson, Christopher Juhlin, and Roland Roberts (2015). "Microearthquakes illuminate the deep structure of the endglacial Pärvie fault, northern Sweden". In: *Geophysical Journal International* 201.3, pp. 1704–1716. DOI: 10.1093/gji/ggv112.

Loomis, B. D., S. B. Luthcke, and T. J. Sabaka (2019). "Regularization and error characterization of GRACE mascons". In: *Journal of Geodesy* 93.9, pp. 1381–1398. DOI: 10.1007/s00190-019-01252-y.

Maclennan, J., M. Jull, D. McKenzie, L. Slater, and K. Grönvold (2002). "The link between volcanism and deglaciation in Iceland". In: *Geochemistry, Geophysics, Geosystems* 3.11, pp. 1–25. DOI: 10.1029/2001GC000282.

Meeus, Jean (1998). *Astronomical Algorithms*. 2nd. Willmann-Bell.

Mey, Jürgen, Dirk Scherler, Andrew D. Wickert, David L. Egholm, Magdala Tesauro, Taylor F. Schildgen, and Manfred R. Strecker (2016). "Glacial isostatic uplift of the European Alps". In: *Nature Communications* 7.1, p. 13382. DOI: 10.1038/ncomms13382.

NASA Goddard Space Flight Center (2026). *GRACE and GRACE-FO GSFC global mascons, RL06 v2.0*. https://earth.gsfc.nasa.gov/geo/data/grace-mascons. Monthly solutions 2002-04 to 2026-03, retrieved 2026-10-10.

Nevada Geodetic Laboratory (2026). *GNSS station time series (IGS20 tenv3) and step file*. https://geodesy.unr.edu/. Retrieved 2026-10-10.

Ogata, Yosihiko (1988). "Statistical Models for Earthquake Occurrences and Residual Analysis for Point Processes". In: *Journal of the American Statistical Association* 83.401, pp. 9–27. DOI: 10.1080/01621459.1988.10478560.

Ogata, Yosihiko (1992). "Detection of precursory relative quiescence before great earthquakes through a statistical model". In: *Journal of Geophysical Research: Solid Earth* 97.B13, pp. 19845–19871. DOI: 10.1029/92JB00708.

Ogata, Yosihiko and Koichi Katsura (1993). "Analysis of temporal and spatial heterogeneity of magnitude frequency distribution inferred from earthquake catalogues". In: *Geophysical Journal International* 113.3, pp. 727–738. DOI: 10.1111/j.1365-246X.1993.tb04663.x.

Open-Meteo (2026). *Historical weather API (ERA5 archive)*. https://archive-api.open-meteo.com/v1/archive. Retrieved 2026-10-10.

Pagli, Carolina and Freysteinn Sigmundsson (2008). "Will present day glacier retreat increase volcanic activity? Stress induced by recent glacier retreat and its effect on magmatism at the Vatnajökull ice cap, Iceland". In: *Geophysical Research Letters* 35.9, 2008GL033510. DOI: 10.1029/2008GL033510.

Pagli, Carolina, Freysteinn Sigmundsson, Björn Lund, Erik Sturkell, Halldór Geirsson, Páll Einarsson, Thóra Árnadóttir, and Sigrún Hreinsdóttir (2007). "Glacio-isostatic deformation around the Vatnajökull ice cap, Iceland, induced by recent climate warming: GPS observations and finite element modeling". In: *Journal of Geophysical Research: Solid Earth* 112.B8, 2006JB004421. DOI: 10.1029/2006JB004421.

Petit, Gérard and Brian Luzum (2010). *IERS Conventions (2010)*. IERS Technical Note 36, Verlag des Bundesamts für Kartographie und Geodäsie, https://www.iers.org/IERS/EN/Publications/TechnicalNotes/tn36.html.

Potin, Bertrand, Florent Aden-Antoniow, and Sergio Ruiz (2024). *Chilean Seismic Catalogue - 1982 - mid-2020*. Zenodo. DOI: 10.5281/zenodo.13146436.

Potin, Bertrand, Sergio Ruiz, Florent Aden-Antoniow, Raúl Madariaga, and Sergio Barrientos (2025). "A Revised Chilean Seismic Catalog from 1982 to Mid-2020". In: *Seismological Research Letters* 96.1, pp. 484–498. DOI: 10.1785/0220240047.

Rollins, Chris, Jeffrey T. Freymueller, and Jeanne M. Sauber (2021). "Stress Promotion of the 1958 Mw∼7.8 Fairweather Fault Earthquake and Others in Southeast Alaska by Glacial Isostatic Adjustment and Inter-earthquake Stress Transfer". In: *Journal of Geophysical Research: Solid Earth* 126.1, e2020JB020411. DOI: 10.1029/2020JB020411.

Römer, Johann (2026a). *glacial-seismic-sensitivity-utac - P133: Early-stage sensitivity tests of glacial-unloading seismicity*. Zenodo. Version 0.2.0. DOI: 10.5281/zenodo.23286492.

Römer, Johann (2026b). *glacial-seismic-sensitivity-utac - P133: Early-stage sensitivity tests of glacial-unloading seismicity*. Zenodo. Version 0.1.0. DOI: 10.5281/zenodo.23281091.

Römer, Johann Benjamin (2026c). *glacial-seismic-sensitivity-utac (GenesisAeon package P133, KLIMA-KRYO-13)*. https://github.com/GenesisAeon/glacial-seismic-sensitivity-utac.

Römer, Johann Benjamin (2026d). *glacial-seismic-sensitivity-utac on PyPI*. https://pypi.org/project/glacial-seismic-sensitivity-utac/.

Sasgen, Ingo, Sebastian Cruz Bacca, Volker Klemann, Anouk Vlug, and Michael Zemp (2026). "Rapid rebound hides glacier mass loss from satellite observations in Alaska and Iceland". In: *Communications Earth & Environment* 7.1, p. 518. DOI: 10.1038/s43247-026-03738-y.

Schmidt, P., B. Lund, C. Hieronymus, J. Maclennan, T. Árnadóttir, and C. Pagli (2013). "Effects of present-day deglaciation in Iceland on mantle melt production rates". In: *Journal of Geophysical Research: Solid Earth* 118.7, pp. 3366–3379. DOI: 10.1002/jgrb.50273.

Schorlemmer, Danijel, Stefan Wiemer, and Max Wyss (2005). "Variations in earthquake-size distribution across different stress regimes". In: *Nature* 437.7058, pp. 539–542. DOI: 10.1038/nature04094.

Schreiber, Thomas and Andreas Schmitz (1996). "Improved Surrogate Data for Nonlinearity Tests". In: *Physical Review Letters* 77.4, pp. 635–638. DOI: 10.1103/PhysRevLett.77.635.

Shearer, Peter M. and Philip B. Stark (2012). "Global risk of big earthquakes has not recently increased". In: *Proceedings of the National Academy of Sciences* 109.3, pp. 717–721. DOI: 10.1073/pnas.1118525109.

Sigmundsson, Freysteinn, Andrew Hooper, Sigrún Hreinsdóttir, Kristín S. Vogfjörd, Benedikt G. Ófeigsson, Elías Rafn Heimisson, et al. (2015). "Segmented lateral dyke growth in a rifting event at Bárðarbunga volcanic system, Iceland". In: *Nature* 517.7533, pp. 191–195. DOI: 10.1038/nature14111.

Simon, Verena, Toni Kraft, Jean-Christophe Maréchal, Agnès Helmstetter, and Tobias Diehl (2025). "Climate-change-induced seismicity: The recent onset of seasonal microseismicity at the Grandes Jorasses, Mont Blanc Massif, France/Italy". In: *Earth and Planetary Science Letters* 666, p. 119372. DOI: 10.1016/j.epsl.2025.119372.

Sirorattanakul, Krittanon and Jean-Philippe Avouac (2026). "Seismic rhythms: Earthquake response to tectonic, hydrological, and tidal forcing in California". In: *Science Advances* 12.13, eadz5711. DOI: 10.1126/sciadv.adz5711.

Swindles, Graeme T., Elizabeth J. Watson, Ivan P. Savov, Ian T. Lawson, Anja Schmidt, Andrew Hooper, et al. (2018). "Climatic control on Icelandic volcanic activity during the mid-Holocene". In: *Geology* 46.1, pp. 47–50. DOI: 10.1130/G39633.1.

Swiss Seismological Service (SED) (2026). *FDSN event web service*. https://eida.ethz.ch/fdsnws/event/1/. Retrieved 2026-10-10.

Talwani, Pradeep, Linyue Chen, and Kalpna Gahalaut (2007). "Seismogenic permeability, ks". In: *Journal of Geophysical Research: Solid Earth* 112.B7, 2006JB004665. DOI: 10.1029/2006JB004665.

Tan, Yen Joe, Felix Waldhauser, Maya Tolstoy, and William S.D. Wilcock (2019). "Axial Seamount: Periodic tidal loading reveals stress dependence of the earthquake size distribution (b value)". In: *Earth and Planetary Science Letters* 512, pp. 39–45. DOI: 10.1016/j.epsl.2019.01.047.

Tanaka, Sachiko, Masakazu Ohtake, and Haruo Sato (2002). "Evidence for tidal triggering of earthquakes as revealed from statistical analysis of global data". In: *Journal of Geophysical Research: Solid Earth* 107.B10. DOI: 10.1029/2001JB001577.

Tsuruoka, Hiroshi, Masakazu Ohtake, and Haruo Sato (1995). "Statistical test of the tidal triggering of earthquakes: contribution of the ocean tide loading effect". In: *Geophysical Journal International* 122.1, pp. 183–194. DOI: 10.1111/j.1365-246X.1995.tb03546.x.

Turcotte, Donald L. and Gerald Schubert (2002). *Geodynamics*. Cambridge University Press. DOI: 10.1017/CBO9780511807442.

U.S. Geological Survey (2026). *ANSS Comprehensive Earthquake Catalog (ComCat), FDSN event web service*. https://earthquake.usgs.gov/fdsnws/event/1/. Retrieved 2026-10-10.

Ueda, Taku, Aitaro Kato, Christopher W. Johnson, and Toshiko Terakawa (2024). "Seasonal Modulation of Crustal Seismicity in Northeastern Japan Driven by Snow Load". In: *Journal of Geophysical Research: Solid Earth* 129.3, e2023JB028217. DOI: 10.1029/2023JB028217.

Van Stiphout, Thomas, Jiancang Zhuang, and David Marsan (2012). *Seismicity declustering*. Community Online Resource for Statistical Seismicity Analysis. DOI: 10.5078/corssa-52382934.

Velicogna, Isabella, Yara Mohajerani, Geruo A, Felix Landerer, Jeremie Mouginot, Brice Noel, et al. (2020). "Continuity of Ice Sheet Mass Loss in Greenland and Antarctica From the GRACE and GRACE Follow-On Missions". In: *Geophysical Research Letters* 47.8, e2020GL087291. DOI: 10.1029/2020GL087291.

Wyss, Max and R. E. Habermann (1988). "Precursory seismic quiescence". In: *pure and applied geophysics* 126.2–4, pp. 319–332. DOI: 10.1007/BF00879001.

Zaliapin, Ilya and Yehuda Ben-Zion (2013). "Earthquake clusters in southern California I: Identification and stability". In: *Journal of Geophysical Research: Solid Earth* 118.6, pp. 2847–2864. DOI: 10.1002/jgrb.50179.