



A modular tool to aggregate results from bioinformatics analyses across many samples into a single report.

Report generated on 2026-09-28, 08:40 NZDT based on data in: /powerplant/workspace/hraijc/Gitrepos/pairtoolsqc/work/9e/f107adbfc2a58d295a0836b6f9cb4e

General Statistics


Sample Name	≥ 30X	Median	Mean Cov.	Read pairs	Unmapped	One-sided	Two-sided	Duplicated	Unique pairs	Cis
Red19_DNase_4M	0.0%	2X	1.9X	3.9 M	3.1%	12.3%	84.6%	0.5%	3.3 M	62.7%
Red19_DNase_7M	0.0%	3X	3.5X	7.4 M	3.1%	12.3%	84.6%	1.0%	6.2 M	62.7%
Red19_FS1H	0.0%	2X	1.9X	4.4 M	3.3%	7.4%	89.3%	7.2%	3.6 M	78.9%
Red19_FS2H	0.0%	3X	3.4X	7.6 M	2.9%	8.0%	89.1%	5.4%	6.3 M	73.3%
Red19_FS3H	0.0%	3X	3.4X	7.3 M	2.7%	9.8%	87.6%	2.1%	6.2 M	65.7%
Red19_DNase_4M.pairsam.stat				3.9 M	3.1%	12.3%	84.6%	0.0%	3.3 M	62.7%
Red19_DNase_7M.pairsam.stat				7.4 M	3.1%	12.3%	84.6%	0.0%	6.2 M	62.7%
Red19_FS1H.pairsam.stat				4.4 M	3.3%	7.4%	89.3%	0.0%	3.9 M	79.2%
Red19_FS2H.pairsam.stat				7.6 M	2.9%	8.0%	89.1%	0.0%	6.7 M	73.5%
Red19_FS3H.pairsam.stat				7.3 M	2.7%	9.8%	87.6%	0.0%	6.4 M	65.8%

Mosdepth

Fast BAM/CRAM depth calculation for WGS, exome, or targeted sequencing. <https://github.com/brentp/mosdepth> DOI: 10.1093/bioinformatics/btx699


Cumulative coverage distribution

Proportion of bases in the reference genome with, at least, a given depth of coverage. Calculated across the entire genome length



Average coverage per contig

Average coverage per contig or chromosome




pairtools

Toolkit for Chromatin Conformation Capture experiments. Handles short-reads paired reference alignments, extracts 3C-specific information, and perform common tasks such as sorting, filtering, and deduplication. <https://github.com/mirnylab/pairtools> DOI: [10.5281/zenodo.1490830](https://doi.org/10.5281/zenodo.1490830); [10.1101/2023.02.13.528389](https://doi.org/10.1101/2023.02.13.528389)

Pairs by alignment status

Number of pairs classified according to their alignment status, including uniquely mapped (UU), unmapped (NN), duplicated (DD), and others.




Pre-filtered pairs by genomic location (overview)

Distribution of pre-filtered pairs (mapping uniquely and rescued) by genomic separation for cis-pairs and trans-pairs.

Skipping the plot as different samples have different distance categories

Pre-filtered pairs as a function of genomic separation (in detail)

Number of interactions (pre-filtered pairs) as a function of genomic separation, log-binned. Click on an individual curve to reveal information for different read pair orientations.



Fraction of read pairs by strand orientation

Number of pre-filtered pairs reported for every type of read pair orientation. Numbers are reported for different ranges of genomic separation and combined.

