



# Day 1

Introduction to Bash scripting  
Decision tree  
Quality filtering WGS data  
Genome assembly  
Assembly evaluation



# WiFi

---

Wifi Name: **UoA-Guest-WiFi**


Username: **mgss2023@wifi.com**

Password: **2P15Pr6Z**

## Shared working doc

---

<https://tinyurl.com/mgss2023doc>




# Welcome!

---

- **Housekeeping**
- **Etherpad for collaborative Q&A/comments**
  - <https://tinyurl.com/mgss2023doc>
- **Overview of attendees**
- **Any questions?**




# Where are we from?




# Do we have data?

---



# Command line experience?

---



# Genomics Aotearoa - Resources

---

## Genomics Aotearoa – GitHub repositories

<https://github.com/GenomicsAotearoa/>

- Metagenomics Summer School material
- RNA seq workshop
- Environmental metagenomics
  - Metagenomic annotation and binning
- Methods and musings
  - Bin cluster refinement
  - Genome assembly ont
  - Metagenomic ont



# Starting each session

1. Log in to the NeSI Jupyter hub via a browser
1. Open the workshop exercise materials on GitHub
1. *Optional: Open a (plain text) text editor for taking notes*





# Bash scripting



# Task: Bash scripting

---

[Go to Github MGSS webpage](#)

## Tasks:

- Introduction to shell
- Introduction to HPC & HPC job




# Metagenomic decision tree(s)




# Our goal: genome recovery

---




# Overview



# Overview

---



# Decision tree

---

- Starts with experimental design
- DNA extraction
- WGS library prep
- Amount of sequencing



Samples/\$\$\$




Many samples/Limited \$\$\$



Few samples/Ample \$\$\$







Samples/\$\$\$

Many samples/Limited \$\$\$

Few samples/Ample \$\$\$


Aim:  
Read analysis

Aim:  
Genome recovery


Libraries:  
Short overlapping PE

Libraries:  
Longer gapped PE inserts

(e.g. 200 bp DNA fragments  
for 2x125 bp reads)




(e.g.  $\geq 550$  bp DNA fragments)



# DNA input

---

- **Very low inputs (e.g. nanograms) for Nextera library prep = enzymatic fragmentation with broad size distributions**



- **High inputs (e.g. 100s ng) for TruSeq = physical fragmentation with defined size selection**



Tends to yield sequences of larger inserts



Samples/\$\$\$

Many samples/Limited \$\$\$

Few samples/Ample \$\$\$


Aim:  
Read analysis

Aim:  
Genome recovery


Libraries:  
Short overlapping PE

Libraries:  
Longer gapped PE inserts

(e.g. 200 bp DNA fragments  
for 2x125 bp reads)



(e.g.  $\geq 550$  bp DNA fragments)



**Samples/\$\$\$**

**Many samples/Limited \$\$\$**

**Few samples/Ample \$\$\$**

**Aim:  
Read analysis**

**Aim:  
Genome recovery**


**Libraries:  
Short overlapping PE**

**Libraries:  
Longer gapped PE inserts**


**Sequencing depth:  
Shallow (<10 Gbp)**

**Sequencing depth:  
Deep (e.g. >=10s Gbp)**


(e.g. 200 bp DNA fragments  
for 2x125 bp reads)



(e.g. >=550 bp DNA fragments)



# Genome recovery per environment




- Fuel cell microbiome
- Peat (boreal)
- Sponge microbiome
- Home microbiome
- Oiled sediment (seafloor)
- Grassland soil




# Estimate sequencing depth

---



# Community structure matters






# Estimate sequencing depth


---

- Estimate generously
- Determine/guesstimate relative abundance of rarest target organism
- Determine/guesstimate the average genome size
- Factor in larger eukaryote genomes
- Decide the minimum desired coverage (e.g. 30x)



# Prokaryotic genome sizes

---




**Fig. 1.** Ranges of bacterial and archaeal genome sizes. Abscissa shows genome size, Mbp; ordinate shows number of genomes; solid line indicates bacterial genomes; dashed line indicates archaeal genomes; A, *C. ruddii* genome; B, *N. equitans* genome; C, minimal size for free-living microorganisms; D, major peak for genome sizes of bacterial and archaeal genomes; E, minor peak for bacterial genomes; F, *Nostoc punctiforme* genome; G, *Sorangium cellulosum* genome; and H, Van Nimwegen limit.

(Smirnov 2010 Molecular Genetics, Microbiology and Virology)



# Estimate sequencing depth

- Estimate generously
- Determine/guesstimate relative abundance of rarest target organism
- Determine/guesstimate the average genome size
- Factor in larger eukaryote genomes
- Decide the minimum desired coverage (e.g. 30x)



e.g., 5% relative abundance = 5% of sequence data

## Mock parameters:

- Bacterial genome 5 Mbp long
- 5% abundance (need 100/5 or 20x)
- 30x coverage

$$5 \text{ Mbp} \times 20 \times 30 = 3,000 \text{ Mbp (or 3 Gbp)}$$



# When you have so many genomes

---

You need a:

- Clear goal
- Question
- Hypothesis to test



# Q&A

**Approaches to metagenomics analyses, e.g.**

- **Short read vs long read sequencing**
- **Assembled genomes vs unbinned reads/contigs**



# Q&A

**Approaches to metagenomics analyses, e.g.**

- **Short read vs long read sequencing**
- **Assembled genomes vs unbinned reads/contigs**



# Mini-project

- Denitrification (Nitrate or nitrite to nitrogen)
- Ammonia oxidation (Ammonia to nitrite or nitrate)
- Anammox (Ammonia and nitrite to nitrogen)
- Sulfur oxidation (SOX pathway, thiosulfate to sulfate)
- Sulfur reduction (DSR pathway, sulfate to sulfide)
- Photosynthetic carbon fixation
- Non-photosynthetic carbon fixation (Reverse TCA or Wood-Ljungdahl)
- Non-polar flagella expression due to a chromosomal deletion
- Plasmid-encoded antibiotic resistance
- Aerobic (versus anaerobic) metabolism



# Quality control/filtering raw reads





# The FastQ data format

---

```
@SEQUENCE_1  
ATCGATCGATCG  
+  
4:<AIIIFIIII  
@SEQUENCE_2  
AATGATCCATG  
+  
IIIIIIIIIIII  
@SEQUENCE_3  
TGTGTGACATG  
+  
BBGBBCIFIII
```

Each sequence is represented by four lines

1. Sequence name
2. Sequence content
3. Spacer line (+, or +Sequence name)
4. Quality information



# The FastQ data format

---

- What does the quality score even mean?
  - It represents the probability of a nucleotide position being incorrectly called

$$Q = -10 \log_{10} p$$

Q	p	Prob. correct
0	1	0
10	0.1	0.9
20	0.01	0.99
30	0.001	0.999
40	0.0001	0.9999

*How each Q value is encoded varies between sequencing platforms*

Generally we work with the **Illumina 1.8+** ([Phred+33](#)) standard



# The FastQ data format

---

- What does the quality score even mean?
  - It represents the probability of a nucleotide position being incorrectly called

$$Q = -10 \log_{10} p$$



Q	p	Prob. correct
0	1	0
10	0.1	0.9
20	0.01	0.99
30	0.001	0.999
40	0.0001	0.9999

*How each Q value is encoded varies between sequencing platforms*

**Generally** we work with the **Illumina 1.8+** ([Phred+33](#)) standard

(33) : !"#%&'()\*+,-./0123456789:;<=>?@ABCDEFGHI



# Quality filtering WGS data

---


- Remove barcode and adapter regions
- Remove low-quality regions of reads
- Identify potential problems that occurred during sequencing
  - Deciphering 'aberrant' metrics in FastQC
    - e.g. Adapter read-through
    - e.g. Rapid drop off in sequence quality



# Quality filtering WGS data

---

- Remove barcode and adapter regions
- Remove low-quality regions of reads
- Identify potential problems that occurred during sequencing
  - Deciphering 'aberrant' metrics in FastQC
    - e.g. Adapter read-through
    - e.g. Rapid drop off in sequence quality



# Task: Quality filtering

---

[Go to Github MGSS webpage](#)

## Tasks:

- **Visualisation with *FastQC***
  - Inspecting FASTQ files
  - Identifying regions of concern
- **Read trimming and adapter removal with Trimmomatic**
  - Removing adapter sequences
  - Removing low-quality regions
- **Diagnosing poor libraries**
- (Optional) Filtering out host DNA



# Common issues with WGS data

---

**Do I need to remove adapters?**



**Yes.**

**I don't know if adapters have been removed or not**



**Check the per-nucleotide distributions  
You will see 100% skews if they remain.**

**What's the lowest Q to allow when trimming?**



**Assembly is a self-correcting process, so  
you can be surprisingly lenient.**


**What if my GC skew is outside of the expected range?**



**FastQC is calibrated to genome data where you expect GC conservation.  
Metagenomes do not adhere to this assumption.**



# Common issues with WGS data





# Common issues with WGS data

---

**Do I need to remove adapters?**



**Yes.**

**I don't know if adapters have been removed or not**



**Check the per-nucleotide distributions  
You will see 100% skews if they remain.**

**What's the lowest Q to allow when trimming?**



**Assembly is a self-correcting process, so  
you can be surprisingly lenient.**

**What if my GC skew is outside of the expected range?**



**FastQC is calibrated to genome data where you expect GC conservation.  
Metagenomes do not adhere to this assumption.**

<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/Help/3%20Analysis%20Modules/>



# Filtering out host DNA

---

Metagenome data derived from microbial communities associated with a host should ideally be filtered to remove any reads originating from host DNA. This may improve the quality and efficiency of downstream data processing

Important for submission to databases e.g. NCBI

- Ethics for human host DNA
- Taonga species in Aotearoa



# Task: Quality filtering

---

[Go to Github MGSS webpage](#)

## Tasks:

- ✓ • **Visualisation with *FastQC***
  - Inspecting FASTQ files
  - Identifying regions of concern
- ✓ • **Read trimming and adapter removal with Trimmomatic**
  - Removing adapter sequences
  - Removing low-quality regions
- ✓ • **Diagnosing poor libraries**
  - (Optional) Filtering out host DNA



# Assembly



# Genome assembly

---

## Overlap-Consensus-Layout (OCL) assembly



# Genome assembly

---

## Overlap-Consensus-Layout (OCL) assembly

TTGAAGAGTT

GGCTCAGATT

TTTGATCATG

AAGAGTTTGA

AACGCTGGCG

GATTGAACGC

CTCAGATTGA

TGAAGAGTTT

ACGCTGGCGC

TCATGGCTCA



# Genome assembly

---

## Overlap-Consensus-Layout (OCL) assembly

```
TTGAAGAGTTTGGCTCAGATTGAACGCTGGCGC
TTGAAGAGTT          GGCTCAGATT  AACGCTGGCG
          TTTGATCATG          GATTGAACGC
      AAGAGTTTGA          CTCAGATTGAACGCTGGCGC
TGAAGAGTTT  TCATGGCTCA
```

The problem for *de novo* assembly?

$$N. comparisons = \frac{(n)(n-1)}{2} = \frac{(10)(10-1)}{2} = 45$$



# Genome assembly

---

## De Bruijn graph assembly

Break reads into shorter *k*-mers

TTGAAGAGTT  
TTGA  
TGAA  
GAAG  
AAGA  
AGAG  
GAGT  
AGTT

TTGA TGAA GAAG AAGA AGAG GAGT AGTT

Number kmers per sequence =  $(L - k) + 1$

$k$  = k-mer length

$L$  = sequence length





# Genome assembly

---

## De Bruijn graph assembly

Identify sequences of shared  $k$ -mers

TTGAAGAGTT

AAGAGTTTGA

AAGA  
AGAG  
GAGT  
AGTT  
GTTT  
TTTG  
TTGA

TTGA TGAA GAAG AAGA AGAG GAGT AGTT GTTT TTTG TTGA




# Genome assembly

---

## De Bruijn graph assembly

Identify sequences of shared  $k$ -mers



# Genome assembly

---

## De Bruijn graph assembly

Problem #1 –  $k$ -mers are short?

TTGAAGAGTTTGTATCATGGCTCAGATTGAACGCTGGCGC  
TTG TTG TTG TGG  
TGA TGA GGC GGC  
GAA GAA  
TCA TCA CGC CGC




# Genome assembly

---

## De Bruijn graph assembly

Problem #1 –  $k$ -mers are short?

TTGAAGAGTTTGATCATGGCTCAGATTGAACGCTGGCGC



# Genome assembly

---

## De Bruijn graph assembly

Problem #2 –  $k$ -mers are long?

TTGAAGAGTT  
TTGAAGAG  
TGAAGAGT  
GAAGAGTT

AAGAGTTTGA  
AAGAGTTT  
AGAGTTTG  
GAGTTTGA

TTGAAGAG TGAAGAGT GAAGAGTT

AAGAGTTT AGAGTTTG GAGTTTGA



# De Bruijn graph assembly

---

## We want a range of *k*-mer sizes

- Short *k*-mers yield higher coverage
- Long *k*-mers assemble longer contigs (jump repeat regions)


## Other considerations for picking *k*-mer sizes

- Size cannot be longer than read length
- Always pick odd *k*-mer sizes
- The more sizes you use, the longer assembly will take

K-mers	N. contigs	Longest contig	N50 >2kbp	L50 >2kbp
21, 33, 55	4,239,806	660,812	6,782	12,906
43, 55, 77, 99, 121	2,519,669	1,022,083	7,990	12,673
21, 43, 55, 77, 99, 121	3,388,682	1,022,083	7,789	13,327



# Which assembler is best?



Outcomes vary by dataset.

Assembly optimization generally requires empirically testing:

- Assemblers
- Parameters



# Which assembler is best?

---

There are three good options

- SPAdes
- MegaHIT
- IDBA-UD





# Which assembler is best?

---

There are three good options

- SPAdes
- MegaHIT
- IDBA-UD

*In conclusion, it can be said that the choice of assembler should depend on the data at hand and on the exact research question asked. Generally, the best assembly is performed by multi k-mer assemblers such as metaSPAdes, Megahit and IDBA-UD. If micro diversity is not a major issue, and the primary research goal is to bin and reconstruct representative bacterial genomes from a given environment, metaSPAdes should clearly be the assembler of choice. This assembler yields the best contig size statistics while capturing a high degree of community diversity, even at high complexity and low read coverage. If micro diversity is however an issue, or if the degree of captured diversity is far more important than contig lengths, then IDBA-UD or Megahit should be preferred.*

Vollmers et al. 2017 (<https://doi.org/10.1371/journal.pone.0169662>)



# Which assembler is best?

---

There are three good options

- SPAdes
- MegaHIT
- IDBA-UD

*In conclusion, it can be said that the choice of assembler should depend on the data at hand and on the exact research question asked. Generally, the best assembly is performed by multi k-mer assemblers such as metaSPAdes, Megahit and IDBA-UD. If micro diversity is not a major issue, and the primary research goal is to bin and reconstruct representative bacterial genomes from a given environment, metaSPAdes should clearly be the assembler of choice. This assembler yields the best contig size statistics while capturing a high degree of community diversity, even at high complexity and low read coverage. If micro diversity is however an issue, or if the degree of captured diversity is far more important than contig lengths, then IDBA-UD or Megahit should be preferred.*

Vollmers et al. 2017 ([https://doi.org/ 10.1371/journal.pone.0169662](https://doi.org/10.1371/journal.pone.0169662))



# What are some key considerations?

---

## Biological

1. What is your hypothesis?
2. What do you want from the data?

## Computational and resource


1. How much data do you have?
2. What are your computational resources?
3. What are your time resources?



# Genome assembly


---

What are some key considerations?



# Too much data?

- Consider testing sub-samples when coverage is very high, e.g. 100s or 1000s
- Example: abundant groundwater genome at 2000x coverage in full dataset
- Empirical testing of subsample sizes identified assembly sweet spot



(Fig. S1, Handley et al., 2014, Environ. Microbiol.)



# Task: Assembly

---

[Go to Github MGSS webpage](#)

## Tasks:

- **Preparing data for assembly (Run IDBA\_UD assembly)**
- **Exploring assembler options**
  - Configure the basic parameters for assembly
- **Submitting jobs to NeSI via slurm**
  - Prepare an assembly job to run under slurm
- **Run SPAdes and IDBA\_UD assembly**
- (Optional) Submitting variant assemblies to NeSI




# **Future considerations and Assembly evaluation**



# Future considerations

---






# Future considerations

---


## Assembly options:

- **Assemble each community separately**
- **Combine reads and assemble all together (co-assembly)**
- **Combine only reads from the same season (mini co-assemblies)**



# Future considerations: rRNA genes

SSU rRNA reference  
guided and iterative  
assembly




(Miller et al., 2011, Genome Biology)



# Future considerations: rRNA genes

---





3b  
Reference-guided SSU rRNA  
reconstruction from WGS data  
(EMIRGE or phyloFlash)

<https://github.com/csmiller/EMIRGE>

<https://hrgv.github.io/phyloFlash/>



# Future considerations: mRNA



Metatranscriptomics

3b  
Reference-guided SSU rRNA  
reconstruction from WGS data  
(EMIRGE or phyloFLASH)

<https://github.com/csmiller/EMIRGE>

<https://hrgv.github.io/phyloFlash/>



# Genome assembly

---

## Contigs vs Scaffolds

Contigs




# Genome assembly

---

## Contigs vs Scaffolds

- Overlapping insert




# Genome assembly

---

## Contigs vs Scaffolds

- Overlapping insert
- Long read sequencing (hybrid assembly)




# Genome assembly

---

## Contigs vs Scaffolds

- Overlapping insert
- Long read sequencing (hybrid assembly)





# Assembly evaluation

---

Parameters to use in evaluation:

- Number of contigs (less is more)
- Total length of contigs (= amount assembled)
- Total length of contigs usable (e.g. >1,000 bp, or at least the length of one bacterial gene)
- Length distribution of contigs
- N50 (minimum contig length at 50% of the total genome length)
- Recovery of particular genomes (determined at later stage)




# Assembly evaluation

---

N50 vs L50

Contigs




Total length



# Assembly evaluation

---

N50 vs L50




# Assembly evaluation

---

## N50 vs L50

*N50* = length of  
middle contig




# Assembly evaluation

---

## N50 vs L50

*N50* = length of  
middle contig



*L50* = count of contigs



# Assembly evaluation

We can then check multiple assembly metrics (e.g. N50/L50) with `BBMap`.

```
[ ]: module load BBMap/38.73 ginkl 2018b
stats.sh in=spades_scaffolds.01.v1.n1000.fna
```

This gives quite a verbose, but useful output:

```
[ ]: A      C      G      T      N      IUPAC  Other  GC      GC_stddev
0.2773  0.2233  0.2223  0.2773  0.0003  0.0000  0.0000  0.4456  0.0100

Main genome scaffold total:      92
Main genome contig total:      111
Main genome scaffold sequence total: 6,454 MB
Main genome contig sequence total: 6,453 MB      0.029% gap
Main genome scaffold N/L50:      14/124,321 KB
Main genome contig N/L50:      10/100,806 KB
Main genome scaffold N/L90:      47/40,787 KB
Main genome contig N/L90:      60/32,398 KB
Max scaffold length:      505,411 KB
Max contig length:      171,572 KB
Number of scaffolds > 50 KB:      41
% main genome in scaffolds > 50 KB: 86.17%

Minimum scaffold length      Number of scaffolds      Number of contigs      Total scaffold length      Total contig length      Scaffold contig coverage
-----
411      92      111      6,454,447      6,452,574      99.97%
1 KB      92      111      6,454,447      6,452,574      99.97%
2.5 KB      82      101      6,441,098      6,439,225      99.97%
5 KB      76      95      6,420,829      6,418,956      99.97%
10 KB      70      89      6,377,701      6,375,828      99.97%
25 KB      59      78      6,192,714      6,190,841      99.97%
50 KB      41      59      5,561,877      5,560,103      99.97%
100 KB      20      35      3,933,974      3,932,493      99.96%
250 KB      4      12      1,608,581      1,599,791      99.95%
500 KB      1      5      505,411      505,016      99.92%
```



# Assembly evaluation

We can then check multiple assembly metrics (e.g. N50/L50) with `BBMap`.

```
[ ]: module load BBMap/38.73 gimkl 2018b
stats.sh in=spades_scaffolds.01.v1.n1000.fna
```

This gives quite a verbose, but useful output:

```
[ ]: A      C      G      T      N      1UPAC  Other  GC      GC_stddev
    0.2773  0.2233  0.2223  0.2773  0.0003  0.0000  0.0000  0.4456  0.0100
```

```
Main genome scaffold total:      92
Main genome contig total:        111
Main genome scaffold sequence total: 6,454 MB
Main genome contig sequence total: 6,453 MB      0.029% gap
Main genome scaffold N/L50:      14/124,321 KB
Main genome contig N/L50:         10/100,806 KB
Main genome scaffold N/L90:      47/40,787 KB
Main genome contig N/L90:         60/32,398 KB
Max scaffold length:              505,411 KB
Max contig length:                171,572 KB
Number of scaffolds > 50 KB:      41
% main genome in scaffolds > 50 KB: 86.17%
```

Minimum scaffold length	Number of scaffolds	Number of contigs	Total scaffold length	Total contig length	Scaffold contig coverage
411	92	111	6,454,447	6,452,574	99.97%
1 KB	92	111	6,454,447	6,452,574	99.97%
2.5 KB	82	101	6,441,098	6,430,225	99.97%
5 KB	76	95	6,420,829	6,418,956	99.97%
10 KB	70	89	6,377,701	6,375,828	99.97%
25 KB	59	78	6,192,714	6,190,841	99.97%
50 KB	41	59	5,561,877	5,560,103	99.97%
100 KB	20	35	3,933,974	3,932,493	99.96%
250 KB	4	12	1,608,581	1,599,791	99.95%
500 KB	1	5	505,411	505,016	99.92%





# Assembly evaluation

We can then check multiple assembly metrics (e.g. N50/L50) with `BBMap`.

```
[ ]: module load BBMap/38.73 ginkl 2018b
stats.sh in=spades scaffolds.01.v1.n1000.fna
```

This gives quite a verbose, but useful output:

```
[ ]: A      C      G      T      N      IUPAC  Other  GC      GC_stddev
0.2773  0.2233  0.2223  0.2773  0.0003  0.0000  0.0000  0.4456  0.0100
```

```
Main genome scaffold total: 92
Main genome contig total: 111
Main genome scaffold sequence total: 6,454 MB
Main genome contig sequence total: 6,453 MB      0.029% gap
Main genome scaffold N/L50: 14/124,321 KB
Main genome contig N/L50: 10/100,806 KB
Main genome scaffold N/L90: 47/40,787 KB
Main genome contig N/L90: 60/32,398 KB
Max scaffold length: 505,411 KB
Max contig length: 171,572 KB
Number of scaffolds > 50 KB: 41
% main genome in scaffolds > 50 KB: 86.17%
```

Minimum scaffold length	Number of scaffolds	Number of contigs	Total scaffold length	Total contig length	Scaffold contig coverage
All	92	111	6,454,447	6,452,574	99.97%
1 KB	92	111	6,454,447	6,452,574	99.97%
2.5 KB	82	101	6,441,098	6,430,225	99.97%
5 KB	76	95	6,420,829	6,418,956	99.97%
10 KB	70	89	6,377,701	6,375,828	99.97%
25 KB	59	78	6,192,714	6,190,841	99.97%
50 KB	41	59	5,561,877	5,560,103	99.97%
100 KB	20	35	3,933,974	3,932,493	99.96%
250 KB	4	12	1,600,581	1,599,791	99.95%
500 KB	1	5	505,411	505,016	99.92%





# Assembly evaluation

We can then check multiple assembly metrics (e.g. N50/L50) with `BBMap`.

```
[ ]: module load BBMap/38.73 ginkl 2018b  
stats.sh in=spades scaffolds.01.v1.n1000.fna
```

This gives quite a verbose, but useful output:

```
[ ]: A      C      G      T      N      IUPAC  Other  GC      GC_stddev  
0.2773  0.2233  0.2223  0.2773  0.0003  0.0000  0.0000  0.4456  0.0100  
  
Main genome scaffold total:      92  
Main genome contig total:      111  
Main genome scaffold sequence total: 6,454 MB  
Main genome contig sequence total: 6,453 MB      0.029% gap  
Main genome scaffold N/L50:      14/124,321 KB  
Main genome contig N/L50:      10/100,806 KB  
Main genome scaffold N/L90:      47/40,787 KB  
Main genome contig N/L90:      60/32,308 KB  
Max scaffold length:      505,411 KB  
Max contig length:      171,572 KB  
Number of scaffolds > 50 KB:      41  
% main genome in scaffolds > 50 KB: 86.17%  
  
Minimum scaffold length      Number of scaffolds      Number of contigs      Total scaffold length      Total contig length      Scaffold contig coverage  
-----  
411      92      111      6,454,447      6,452,574      99.97%  
1 KB      92      111      6,454,447      6,452,574      99.97%  
2.5 KB      82      101      6,441,098      6,430,225      99.97%  
5 KB      76      95      6,420,829      6,418,956      99.97%  
10 KB      70      89      6,377,701      6,375,828      99.97%  
25 KB      50      78      6,192,714      6,190,841      99.97%  
50 KB      41      59      5,561,877      5,560,103      99.97%  
100 KB      20      35      3,933,974      3,932,493      99.96%  
250 KB      4      12      1,600,581      1,599,791      99.95%  
500 KB      1      5      505,411      505,016      99.92%
```



# Assembly evaluation

We can then check multiple assembly metrics (e.g. N50/L50) with `BBMap`.

```
[ ]: module load BBMap/38.73 ginkl 2018b
stats.sh in=spades_scaffolds.01.v1.n1000.fna
```

This gives quite a verbose, but useful output:

```
[ ]: A      C      G      T      N      IUPAC  Other  GC      GC_stddev
0.2773  0.2233  0.2223  0.2773  0.0003  0.0000  0.0000  0.4456  0.0100
```

```
Main genome scaffold total:      92
Main genome contig total:      111
Main genome scaffold sequence total: 6.454 MB
Main genome contig sequence total: 6.453 MB      0.029% gap
Main genome scaffold N/L50:      14/124.321 KB
Main genome contig N/L50:      10/100.806 KB
Main genome scaffold N/L90:      47/40.787 KB
Main genome contig N/L90:      60/32.398 KB
Max scaffold length:      505.411 KB
Max contig length:      171.572 KB
Number of scaffolds > 50 KB:      41
% main genome in scaffolds > 50 KB: 86.17%
```

Minimum scaffold length	Number of scaffolds	Number of contigs	Total scaffold length	Total contig length	Scaffold contig coverage
All	92	111	6,454,447	6,452,574	99.97%
1 KB	92	111	6,454,447	6,452,574	99.97%
2.5 KB	82	101	6,441,098	6,430,225	99.97%
5 KB	76	95	6,420,829	6,418,956	99.97%
10 KB	70	89	6,377,701	6,375,828	99.97%
25 KB	59	78	6,192,714	6,190,841	99.97%
50 KB	41	59	5,561,877	5,560,103	99.97%
100 KB	20	35	3,933,974	3,932,493	99.96%
250 KB	4	12	1,608,581	1,599,791	99.95%
500 KB	1	5	505,411	505,016	99.92%



# Assembly evaluation

We can then check multiple assembly metrics (e.g. N50/L50) with `BBMap`.

```
[ ]: module load BBMap/38.73 gimkl 2018b
stats.sh in=spades_scaffolds.01.v1.n1000.fna
```

This gives quite a verbose, but useful output:

```
[ ]: A      C      G      T      N      IUPAC  Other  GC      GC_stddev
      0.2773  0.2233  0.2223  0.2773  0.0003  0.0000  0.0000  0.4456  0.0100

Main genome scaffold total:      92
Main genome contig total:      111
Main genome scaffold sequence total: 6,454 MB
Main genome contig sequence total: 6,453 MB      0.029% gap
Main genome scaffold N/L50:      14/124,321 KB
Main genome contig N/L50:      10/100,806 KB
Main genome scaffold N/L90:      47/40,767 KB
Main genome contig N/L90:      60/32,398 KB
Max scaffold length:      505,411 KB
Max contig length:      171,572 KB
Number of scaffolds > 50 KB:      41
% main genome in scaffolds > 50 KB: 86.17%
```

Minimum scaffold length	Number of scaffolds	Number of contigs	Total scaffold length	Total contig length	Scaffold contig coverage
All	92	111	6,454,447	6,452,574	99.97%
1 KB	92	111	6,454,447	6,452,574	99.97%
2.5 KB	82	101	6,441,098	6,430,225	99.97%
5 KB	76	95	6,420,829	6,418,956	99.97%
10 KB	70	89	6,377,701	6,375,828	99.97%
25 KB	59	78	6,192,714	6,190,841	99.97%
50 KB	41	59	5,561,877	5,560,103	99.97%
100 KB	20	35	3,933,974	3,932,493	99.96%
250 KB	4	12	1,600,581	1,599,791	99.95%
500 KB	1	5	505,411	505,016	99.92%



# Task: Assembly evaluation

---

[Go to Github MGSS webpage](#)

Tasks:

- Assembly evaluation
- Short contig removal

