

Topic 2-1: Likelihood Construction & Estimation

Univariate Models

EXST 7160
Department of Experimental Statistics
Louisiana State University

July 25, 2021



Content

Discrete IID Random Variables

Multinomial Likelihoods

Continuous IID Random Variables

Mixtures of Discrete and Continuous Components

Proportional Likelihoods

The Empirical Distribution Function as an MLE

Likelihoods from Censored Data



Contents

- 1 General Concept
- 2 Constructing Likelihood Functions
- 3 More on likelihood functions
- 4 Appendix: The connection between discrete and continuous likelihoods



Contents

- 1 General Concept
 - Introduction
- 2 Constructing Likelihood Functions
- 3 More on likelihood functions
- 4 Appendix: The connection between discrete and continuous likelihoods



Statistical Models

- A statistical model is a general functional relation between the unknown parameter(s) and the observed data.



Statistical Models

- A statistical model is a general functional relation between the unknown parameter(s) and the observed data.
- After a statistical model for the observed data has been formulated, the likelihood function of the data is the natural starting point for the inference in many statistical problems.



Statistical Models

- A statistical model is a general functional relation between the unknown parameter(s) and the observed data.
- After a statistical model for the observed data has been formulated, the likelihood function of the data is the natural starting point for the inference in many statistical problems.
- The likelihood function typically leads to essentially automatic methods of inference, including point estimation, interval estimation, and hypothesis testing.



Statistical Models

- A statistical model is a general functional relation between the unknown parameter(s) and the observed data.
- After a statistical model for the observed data has been formulated, the likelihood function of the data is the natural starting point for the inference in many statistical problems.
- The likelihood function typically leads to essentially automatic methods of inference, including point estimation, interval estimation, and hypothesis testing.
- In this topic, we will focus on constructing the likelihood functions from various types of data, including discrete, continuous, mixture of discrete and continuous, and censored data.



Statistical Models

- The likelihood is the joint density of the observed data to be analyzed.



Statistical Models

- The likelihood is the joint density of the observed data to be analyzed.
- Let the random variables Y_1, \dots, Y_n have a joint density function $f(\mathbf{Y} = (Y_1, \dots, Y_n)^T; \boldsymbol{\theta})$ with unknown b density parameters $\boldsymbol{\theta} = (\theta_1, \dots, \theta_b)$. Then, given observed data $\mathbf{Y} = \mathbf{y}$, where $\mathbf{y} \equiv (y_1, \dots, y_n)^T$, the function of $\boldsymbol{\theta}$

$$L(\boldsymbol{\theta}; \mathbf{y}) = f(\mathbf{Y} = \mathbf{y}; \boldsymbol{\theta})$$

is the likelihood function.



Likelihood for iid data

- If the random variables Y_1, \dots, Y_n are independent, then the likelihood function becomes

$$L(\boldsymbol{\theta}; \mathbf{y}) = \prod_{i=1}^n f_i(Y_i = y_i; \boldsymbol{\theta}),$$

where $f_i(Y_i; \boldsymbol{\theta})$ is the density of Y_i .



Likelihood for iid data

- If the random variables Y_1, \dots, Y_n are independent, then the likelihood function becomes

$$L(\theta; \mathbf{y}) = \prod_{i=1}^n f_i(Y_i = y_i; \theta),$$

where $f_i(Y_i; \theta)$ is the density of Y_i .

- If the random variables Y_1, \dots, Y_n are independent and identically distributed (denoted by iid), then the likelihood function becomes

$$L(\theta; \mathbf{y}) = \prod_{i=1}^n f(Y_i = y_i; \theta),$$

where f is the distribution that all Y_1, \dots, Y_n follow.



Solving the Likelihood

- The value of θ that maximizes the likelihood function $L(\theta; \mathbf{y})$ is called the maximum likelihood estimator (MLE) denoted by $\hat{\theta}_{MLE}$.



Solving the Likelihood

- The value of θ that maximizes the likelihood function $L(\theta; \mathbf{y})$ is called the maximum likelihood estimator (MLE) denoted by $\hat{\theta}_{MLE}$.
- In practice, $\hat{\theta}_{MLE}$ is usually calculated by optimizing the log likelihood function $\log(L(\theta; \mathbf{y}))$



Solving the Likelihood

- The value of θ that maximizes the likelihood function $L(\theta; \mathbf{y})$ is called the maximum likelihood estimator (MLE) denoted by $\hat{\theta}_{MLE}$.
- In practice, $\hat{\theta}_{MLE}$ is usually calculated by optimizing the log likelihood function $\log(L(\theta; \mathbf{y}))$
- Note: $\hat{\theta}_{MLE}$ is “optimal” under some regularity conditions (to be discussed in a future topic).



Solving the Likelihood-differentiable likelihoods

- Denote the log likelihood function $\log(L(\boldsymbol{\theta}; \mathbf{y}))$ by $\ell(\boldsymbol{\theta})$, and assume $\ell(\boldsymbol{\theta})$ is differentiable with respect to $\boldsymbol{\theta}$.



Solving the Likelihood-differentiable likelihoods

- Denote the log likelihood function $\log(L(\boldsymbol{\theta}; \mathbf{y}))$ by $\ell(\boldsymbol{\theta})$, and assume $\ell(\boldsymbol{\theta})$ is differentiable with respect to $\boldsymbol{\theta}$.
- The procedure for obtaining the MLE from $\ell(\boldsymbol{\theta})$ is



Solving the Likelihood-differentiable likelihoods

- Denote the log likelihood function $\log(L(\boldsymbol{\theta}; \mathbf{y}))$ by $\ell(\boldsymbol{\theta})$, and assume $\ell(\boldsymbol{\theta})$ is differentiable with respect to $\boldsymbol{\theta}$.
- The procedure for obtaining the MLE from $\ell(\boldsymbol{\theta})$ is
 - 1 Differentiate $\ell(\boldsymbol{\theta})$ with respect to $\boldsymbol{\theta}$ to obtain the likelihood score function $S(\boldsymbol{\theta}) = (\frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_1}, \dots, \frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_b})^T$.



Solving the Likelihood-differentiable likelihoods

- Denote the log likelihood function $\log(L(\boldsymbol{\theta}; \mathbf{y}))$ by $\ell(\boldsymbol{\theta})$, and assume $\ell(\boldsymbol{\theta})$ is differentiable with respect to $\boldsymbol{\theta}$.
- The procedure for obtaining the MLE from $\ell(\boldsymbol{\theta})$ is
 - 1 Differentiate $\ell(\boldsymbol{\theta})$ with respect to $\boldsymbol{\theta}$ to obtain the likelihood score function $S(\boldsymbol{\theta}) = (\frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_1}, \dots, \frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_b})^T$.
 - 2 Find possible MLE candidates by solving the likelihood equations $S(\boldsymbol{\theta}) = \mathbf{0}_{b \times 1}$.



Solving the Likelihood-differentiable likelihoods

- Denote the log likelihood function $\log(L(\boldsymbol{\theta}; \mathbf{y}))$ by $\ell(\boldsymbol{\theta})$, and assume $\ell(\boldsymbol{\theta})$ is differentiable with respect to $\boldsymbol{\theta}$.
- The procedure for obtaining the MLE from $\ell(\boldsymbol{\theta})$ is
 - 1 Differentiate $\ell(\boldsymbol{\theta})$ with respect to $\boldsymbol{\theta}$ to obtain the likelihood score function $S(\boldsymbol{\theta}) = (\frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_1}, \dots, \frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta_b})^T$.
 - 2 Find possible MLE candidates by solving the likelihood equations $S(\boldsymbol{\theta}) = \mathbf{0}_{b \times 1}$.
 - 3 Check if the solution from the last step is the global maximizer of $\ell(\boldsymbol{\theta})$. If it is, then the solution is the MLE of $\boldsymbol{\theta}$.



An R Example



Remark

- A non-differentiable likelihood example can be found in Example 7.2.9 of Casella and Berger (2002).



Remark

- A non-differentiable likelihood example can be found in Example 7.2.9 of Casella and Berger (2002).
- An example for checking if the solution of the score function equation with two unknown parameters is a global maximizer can be found in Example 7.2.12 of Casella and Berger (2002).



Contents

- 1 General Concept
- 2 Constructing Likelihood Functions
- 3 More on likelihood functions
- 4 Appendix: The connection between discrete and continuous likelihoods



Discrete IID Random Variables



Discrete IID Random Variables



Discrete IID Random Variables

-
-
-



Continuous IID Variables



Continuous IID Variables

-
-



Continuous IID Variables

-
-
- References: More examples for constructing the product likelihood associated with iid data can be found in Section 7.2.2 of Casella and Berger (2002).



Multinomial



Multinomial



Multinomial

-
-
- References: Examples for



Multivariate Normal



Multivariate Normal



Multivariate Normal

-
-
-



Mixture of discrete and continuous



Mixture of discrete and continuous



Mixture of discrete and continuous

-
-
-



Multinomial



Multinomial



Multinomial

-
-
- References: Examples for



Contents

- 1 General Concept
- 2 Constructing Likelihood Functions
- 3 More on likelihood functions**
- 4 Appendix: The connection between discrete and continuous likelihoods



Proportional Likelihood



Proportional Likelihood



Proportional Likelihood

-
-
-



Empirical Likelihood



Empirical Likelihood



Empirical Likelihood

-
-
-



Likelihood With Censored data



Likelihood With Censored data



Likelihood With Censored data

-
-
-



Contents

- 1 General Concept
- 2 Constructing Likelihood Functions
- 3 More on likelihood functions
- 4 Appendix: The connection between discrete and continuous likelihoods



A general working definition of the likelihood



A general working definition of the likelihood



A general working definition of the likelihood

-
-
-



Empirical Likelihood



Empirical Likelihood



Empirical Likelihood

-
-
-



Likelihood With Censored data



Likelihood With Censored data



Likelihood With Censored data

-
-
-

