A CERTIFIED SIMPLE, EVEN GROUND STATE OF THE LOCALIZED WEIL QUADRATIC FORM FOR ALL WINDOWS $L \in (0, 1.4035]$

JOHANN ALEXANDER BOTHA

Abstract. Let Q_L be Weil's quadratic form for the Riemann zeta function, restricted to real test functions supported in $[-L, L]$ on the logarithmic scale; it contains the pole terms, the archimedean term and every prime power n with $\log n < 2L$. Connes and van Suijlekom proved that if the lowest eigenvalue of the associated self-adjoint operator is simple with an even eigenfunction ξ, then $\hat{\xi}$ has only real zeros; for the Weil form this "simple-even" property is one of the missing steps in the strategy of Connes, Consani and Moscovici. We give a computer-assisted proof, in Arb ball arithmetic, that the ground state of Q_L is simple and even for every $L \in (0, 1.4035]$. The proof combines min-max monotonicity in L with three pieces: a Lipschitz argument on $(0, 0.05]$, a covering of $[0.05, 1.19]$ by 31 certified windows based on Zhu's frequency-cutoff reduction, and a covering of $(1.19, 1.4035]$ by 8 windows in which second-eigenvalue lower bounds are obtained from Elston's window-compressed prime-norm certifier combined with a rank-one deflation, and upper bounds from Legendre–Galerkin Rayleigh quotients with rigorous quadrature bounds. On $(0, 0.8]$ the statement had been claimed before; the range $(0.8, 1.4035]$ is, to our knowledge, new. As by-products we give rigorous upper bounds for the lowest eigenvalue in both parity sectors up to $L = 2.25$. The extension of the same computation to $L = 1.5$ is in progress.

Contents

1. Introduction 2
2. Setup and normalisation 3
3. Monotonicity and the grid argument 5
4. Main theorem 5
5. Proof 5
6. Results 11
7. Rigorous upper bounds for larger windows 14
8. A numerical observation: prime-power singular points 15
9. Reproducibility 16
10. Limitations 18
Acknowledgements 18
AI-assistance statement 18
References 18

Date: Version 1.1.0, 31 October 2026.

2020 *Mathematics Subject Classification.* 11M26 (primary); 65G20, 47A75, 65F15 (secondary).

Key words and phrases. Weil explicit formula, Weil positivity, quadratic form, ground state, computer-assisted proof, ball arithmetic.

1. Introduction

1.1. **The localized Weil form.** For a real function $f \in L^2(\mathbb{R})$ with support in $[-L, L]$ put $\tilde{f}(x) = f(-x)$, $g = f * \tilde{f}$ and $F(z) = \hat{f}(z) = \int f(x) e^{izx}\, dx$. Weil's explicit-formula functional [26], evaluated at g, gives the quadratic form

$$Q_L(f) = W(g), \qquad W(g) = \hat{g}(\tfrac{i}{2}) + \hat{g}(-\tfrac{i}{2}) - g(0)\log\pi + \frac{1}{2\pi}\int_{\mathbb{R}} \hat{g}(r)\,\operatorname{Re}\psi\big(\tfrac{1}{4} + \tfrac{ir}{2}\big)\, dr - \sum_{n\geq 2} \frac{\Lambda(n)}{\sqrt{n}}\big[g(\log n) + g(-\log n)\big], \tag{1.1}$$

where $\psi = \Gamma'/\Gamma$ and Λ is the von Mangoldt function. Weil's criterion states that the Riemann hypothesis (RH) is equivalent to $Q_L(f) \geq 0$ for all such f and all $L > 0$ [26, 4]. Since g is supported in $[-2L, 2L]$, only prime powers with $\log n < 2L$ contribute; the prime power n enters at $L = \frac{1}{2}\log n$ (Table 1).

Table 1. Prime powers n and the window $L = \frac{1}{2}\log n$ at which they enter the form, for $L \leq 1.5$.

n	2	3	4	5	7	8
$\frac{1}{2}\log n$	0.3466	0.5493	0.6931	0.8047	0.9730	1.0397
n	9	11	13	16	17	19
$\frac{1}{2}\log n$	1.0986	1.1989	1.2825	1.3863	1.4166	1.4722

1.2. **The simple-even hypothesis.** Connes and van Suijlekom [8, Thm. 6.1] prove: if a quadratic form with Schwartz kernel $\tilde{D}(x-y)$, D a real distribution, defines a lower-bounded self-adjoint operator whose lowest spectral value is a *simple, isolated* eigenvalue with *even* eigenfunction ξ, then all zeros of the entire function $\hat{\xi}$ are real. Connes, Consani and Moscovici [7] propose a strategy towards RH based on a spectral realization of the zeros through the eigenfunctions ξ_λ of the Weil form on growing windows; in [7, §8] they list two "missing steps": (i) the smallest eigenvalue of the Weil quadratic form QW_λ is simple with even eigenvector, and (ii) the zeros of $\hat{\xi}_\lambda$ converge to the nontrivial zeros of ζ. The same two steps are listed in [6, §6.6]. The present paper addresses only step (i), and only on the bounded range $L \in (0, 1.4035]$; it says nothing about step (ii) or about large L, and does not prove positivity beyond what the certified lower bounds imply on this range. In the conventions of [8] the operator acts on $L^2([-\frac{L}{2}, \frac{L}{2}])$, so their L is twice ours; in [7] the window is $[\lambda^{-1}, \lambda]$ on the multiplicative scale, i.e. $L = \log\lambda$. Our form agrees with theirs only up to a positive affine rescaling, see Remark 2.4 and §10.

1.3. **Prior work.** We list what we know to have been proved or claimed; status labels follow the sources. Many items are unrefereed 2026 deposits or preprints, some AI-assisted.

- *Small windows.* Suzuki [23, Thm. 1.4] proves for the full form that the lowest eigenvalue is positive, simple and has an even eigenfunction for $L \in (0, a_0)$ with a_0 not explicit (Kato perturbation theory). The mathia notes PL-294 [19] (unrefereed) show that the full form satisfies the Beurling–Deny criterion iff $L \leq \frac{1}{2}\log\rho_{\text{plastic}} \approx 0.1406$, which gives a simple ground state of fixed sign. Andrade [2] (unrefereed) proves a Perron-type theorem for the pole-removed form and states the same threshold for the full kernel in a remark. An analytic argument for $L \leq 9/50$ is reported in [13]; we have not verified it.

- *Zhu* [28] introduced the frequency-cutoff ("one-stroke") reduction with a head/tail block estimate [Zhu, Thm. 1.1] on which all our window certificates rest, certified simple-even at the single window $L = 0.8$ [Zhu, Thm. 6.2], and certified variational upper bounds on the lowest eigenvalue for $L \in [0.5, 2]$ [Zhu, Table 3]. Zhu [§7] retracts an earlier positivity claim at $L = 1.19$.
- *Kramarenko-Byrd* [17] (unrefereed) claims a rigorous proof of simple-even for all $0 < L \le 4/5$ in our half-width convention. We have not checked it. If it is correct, our result on $(0, 0.8]$ is an independent confirmation.
- *Perturbative simplicity* on a neighbourhood of $L = 0.8$ of non-explicit radius: PL-287 [19].
- *Positivity and per-sector lower bounds* (no simplicity or parity statement): jose-paul-b [16] for $L \le \frac{1}{2}\log 7$; Liu [18] at $L = 17/16$ (block-Schur bounded comparison, $Q \ge 2^{-49162}$); Mori [20] at $L = \frac{1}{2}\log 15$; Senke [22] at $L = 77/64$; Elder [10] at $L = 0.8$; and Elston [11], whose certified lowest-eigenvalue lower bounds at $L = 1.19$, 1.3 and 1.495 (e.g. $\lambda \ge 3.501 \cdot 10^{-96}$ at $L = 1.495$) use the certifier we build on in Piece III.
- *Finite matrices.* Andrade [1] and Andrews Jr. [3] certify simple-even for finite Galerkin matrices QW^N; these are not statements about the continuum form.
- *Classical ancestor.* Yoshida [27] combined finitely many pivots with a rigorous high-frequency complement lemma.

1.4. **Contributions.**

(1) A computer-assisted proof that the ground state of Q_L is simple and even for every $L \in (0, 1.4035]$ (Theorem 4.1). The part we believe to be new is $(0.8, 1.4035]$. The piece $(0, 1.19]$ was released earlier as an unrefereed note [5]; this paper reproduces its mathematical content.
(2) A rank-one deflation added to Elston's certifier (Lemma 5.14, Corollary 5.15), which turns a certified lowest-eigenvalue bound into a certified *second* even eigenvalue bound at no extra assembly cost.
(3) Rigorous upper bounds for the lowest even and odd eigenvalues up to $L = 2.25$ from a fast Legendre convolution recurrence with a rigorous Gauss–Legendre bound (§7).
(4) Numerical measurements of the weak singular points of the computed minimisers at $|y| = |L - \log n|$ reported by Eberwein [9, §13] (§8); their location is not new.

Not claimed as new: the range $(0, 0.8]$ [17] and in particular $(0, 0.1406]$ [19]; the window $L = 0.8$ by itself [Zhu, Thm. 6.2]; positivity on this range; certified upper bounds on the lowest eigenvalue for $L \in [0.5, 2]$ [Zhu, Table 3]; the frequency-cutoff reduction and block estimate [Zhu, Thm. 1.1]; the window-compressed prime-norm reduction and its implementation [11]; the Legendre convolution recurrence [14], already used for this problem by Liu [18].

2. Setup and normalisation

2.1. **Frequency side.**

For real f supported in $[-L, L]$ one has $\hat{g}(t) = F(t)F(-t) = |F(t)|^2$ for real t, and by Plancherel $g(0) = \|f\|^2 = \frac{1}{\pi}\int_0^\infty |F|^2$. Hence

$$Q_L(f) = [\text{pole}] + \frac{1}{\pi}\int_0^\infty \Psi_L(t)|F(t)|^2\,dt, \tag{2.1}$$

$$\Psi_L(t) = \operatorname{Re}\psi\big(\tfrac{1}{4} + \tfrac{it}{2}\big) - \log\pi - \sum_{\log n < 2L} \frac{2\Lambda(n)}{\sqrt{n}}\cos(t\log n),$$

with [pole] $= 2F(\frac{i}{2})F(-\frac{i}{2})$. Here $F(\frac{i}{2}) = \int f e^{-x/2}$ is real; for even f, [pole] $= +2F(\frac{i}{2})^2$, and for odd f, [pole] $= -2F(\frac{i}{2})^2$. This normalisation is that of Zhu [Zhu, §6]; it was checked numerically against the zero side of the explicit formula (first 400 zeros).

2.2. **Position side.** Let γ denote Euler's constant. For $g = f * \tilde{f}$,

$$Q_L(f) = 4\int_0^{2L} g(x)\cosh\tfrac{x}{2}\,dx - g(0)\big(\log\pi + \gamma + \log(1-e^{-4L})\big) + \int_0^{2L} \frac{2\,[e^{-2x}g(0) - e^{-x/2}g(x)]}{1-e^{-2x}}\,dx - 2\sum_{\log n<2L}\frac{\Lambda(n)}{\sqrt{n}}\,g(\log n). \tag{2.2}$$

Derivation. For smooth f, insert $\psi(s) = -\gamma + \int_0^\infty \frac{e^{-t}-e^{-st}}{1-e^{-t}}\,dt$ into (1.1) and apply Fourier inversion (Fubini applies since g is smooth). For $x > 2L$ only the $g(0)$ part of the integrand survives and integrates in closed form, giving the term $\log(1-e^{-4L})$; the pole term gives $4\int_0^{2L} g\cosh(x/2)$. For a polynomial f on $[-L,L]$ (needed for the trial functions), g is a polynomial on $[0,2L]$ and Lipschitz on $\mathbb{R}$; apply (2.2) to the mollification $f * \phi_\epsilon$ on the window $L+\epsilon$ (no $\log n$ equals $2L$, and the $\log(1-e^{-4L})$ term absorbs the change of upper limit) and let $\epsilon \to 0$: the frequency side converges by dominated convergence since $\Psi_L|F|^2 = O(t^{-2}\log t)$, and on the position side $g_\epsilon \to g$ uniformly with a uniform Lipschitz constant, which dominates the integrand $\sim (g(0)-g(x))/(2x)$ near $x = 0$.

2.3. **The operator, parity and discreteness.** Let A_L be the self-adjoint operator associated with the closure of Q_L on $C_c^\infty(-L,L)$ in $L^2(-L,L)$. By Lemma 5.4 below, $\Psi_L(t) \geq \log(t/2\pi) - 1/t - \sum_{\log n<2L} 2\Lambda(n)n^{-1/2} \to \infty$, and the pole term is bounded by $C\|f\|^2$; hence the form norm controls $\int \log(2+t)|F|^2$. For functions supported in $[-L,L]$ the set where this quantity and $\|f\|$ are bounded is compact in L^2 (high frequencies are uniformly small, and on compact t-sets the F are equicontinuous by Paley–Wiener). So A_L has compact resolvent, in agreement with [7, Thm. 3.6]. The kernel is real, so we work with real f throughout.

Lemma 2.1 (Parity splitting)**.** *Let $(Rf)(x) = f(-x)$. Then $Q_L(Rf) = Q_L(f)$, and $Q_L(f_e + f_o) = Q_L(f_e) + Q_L(f_o)$ for f_e even and f_o odd. Hence $A_L = A_L^{\mathrm{ev}} \oplus A_L^{\mathrm{odd}}$.*

Proof. Rf has transform $F(-t)$, which leaves $|F(t)|$ (t real) and $F(\frac{i}{2})F(-\frac{i}{2})$ unchanged. For the bilinear form B of Q_L, $B(f_e,f_o) = B(Rf_e,Rf_o) = -B(f_e,f_o)$. □

Definition 2.2. For each parity sector $\sigma \in \{\mathrm{ev},\mathrm{odd}\}$ let $\lambda_1^\sigma(L) \leq \lambda_2^\sigma(L) \leq \cdots$ be the eigenvalues of A_L^σ, counted with multiplicity; by compactness of the resolvent they coincide with the min-max values

$$\lambda_k^\sigma(L) = \inf_{\substack{V\subset\mathcal{D}^\sigma\\ \dim V = k}} \sup_{0\neq f\in V} \frac{Q_L(f)}{\|f\|^2},$$

where $\mathcal{D}^\sigma$ is the form domain of the sector. (Indices start at 1, as in [28, 11]; the earlier note [5] writes $\lambda_0, \lambda_1^{\mathrm{ev}}, \lambda_0^{\mathrm{odd}}$ for our $\lambda_1^{\mathrm{ev}}, \lambda_2^{\mathrm{ev}}, \lambda_1^{\mathrm{odd}}$.)

Corollary 2.3. *The lowest eigenvalue of A_L is simple with even eigenfunction if and only if*

$$\lambda_1^{\mathrm{ev}}(L) < \min\big(\lambda_2^{\mathrm{ev}}(L), \lambda_1^{\mathrm{odd}}(L)\big).$$

Remark 2.4. The property in Corollary 2.3 is invariant under $Q \mapsto aQ + b\|\cdot\|^2$ with $a > 0$. Conventions that differ from ours by a positive factor (e.g. 2π) or a constant shift (e.g. the placement of $\log\pi$) therefore do not affect the theorem.

3. Monotonicity and the grid argument

Lemma 3.1 (Monotonicity in L)**.** *For $0 < L < L'$, every $k \geq 1$ and $\sigma \in \{\mathrm{ev}, \mathrm{odd}\}$, $\lambda_k^\sigma(L') \leq \lambda_k^\sigma(L)$.*

Proof. Extension by zero $\iota : L^2(-L, L) \to L^2(-L', L')$ is an isometry that preserves parity and maps the form domain into the form domain, and $Q_{L'}(\iota f) = Q_L(f)$: the pole and archimedean terms depend only on F, and a prime power with $2L \leq \log n < 2L'$ contributes $g(\log n) = 0$ because g is continuous and supported in $[-2L, 2L]$ with $g(\pm 2L) = 0$. The min-max infimum over the larger family of subspaces of $L^2(-L', L')$ (which contains all ιV) can only be smaller. □

Corollary 3.2 (Step criterion)**.** *Let $L_i < L_{i+1}$, let $U(L_i) \geq \lambda_1^{\mathrm{ev}}(L_i)$, and let $\ell_2(L_{i+1}) \leq \lambda_2^{\mathrm{ev}}(L_{i+1})$ and $\ell^{\mathrm{odd}}(L_{i+1}) \leq \lambda_1^{\mathrm{odd}}(L_{i+1})$. If*

$$\min\big(\ell_2(L_{i+1}), \ell^{\mathrm{odd}}(L_{i+1})\big) > U(L_i), \tag{3.1}$$

then the ground state of A_L is simple and even for every $L \in [L_i, L_{i+1}]$.

Proof. For such L, by Lemma 3.1,

$$\lambda_1^{\mathrm{ev}}(L) \leq \lambda_1^{\mathrm{ev}}(L_i) \leq U(L_i) < \min(\ell_2, \ell^{\mathrm{odd}})(L_{i+1}) \leq \min\big(\lambda_2^{\mathrm{ev}}(L), \lambda_1^{\mathrm{odd}}(L)\big);$$

conclude with Corollary 2.3. □

We call the quotient $\min(\ell_2, \ell^{\mathrm{odd}})(L_{i+1})/U(L_i)$ the *step ratio*. Prime-power entries inside a step are harmless, since Lemma 3.1 holds across them.

4. Main theorem

Theorem 4.1. *For every $L \in (0, 1.4035]$ the lowest eigenvalue of the operator A_L associated with Weil's quadratic form (1.1) on $L^2(-L, L)$ is simple, and its eigenfunction is even. Quantitatively:*

1. *on $(0, 0.05]$: $\lambda_2^{\mathrm{ev}} - \lambda_1^{\mathrm{ev}} \geq 0.4172$ and $\lambda_1^{\mathrm{odd}} - \lambda_1^{\mathrm{ev}} \geq 0.6624$;*
2. *on each of the 30 grid steps $[L_i, L_{i+1}] \subset [0.05, 1.19]$ of Table 2: step ratio ≥ 1.288;*
3. *on each of the 8 grid steps $[L_i, L_{i+1}] \subset [1.19, 1.4035]$ of Table 4: step ratio ≥ 1.131.*

Remark 4.2 (Status of release 1.1.0). Items (1) and (2) are the certificate of record of [5], re-checked by the gluing script (§9). Item (3) rests on the 16 lower-bound certificates and 8 upper bounds of Table 4, all complete. The goal of this work is $L \in (0, 1.5]$: the extension to 1.5 is in progress, with the even sector at 1.4255 and both sectors at 1.468, 1.4885 and 1.5 still running. The later step $[1.4255, 1.447]$ already passes (Table 4), but it is not part of the theorem, because the step $[1.4035, 1.4255]$ before it is not yet complete. At the single window $L = 1.19$ the Piece III machinery also confirms simple-even with a second, independent code path (§6.2). Piece III beyond $L = 1.19$ has not been audited (§10).

The numbers in (1)–(3) are outward-rounded lower ends. The proof is computer-assisted and uses Arb ball arithmetic [15, 24]; it is given in §5.

5. Proof

5.1. **Piece I: small windows $L \in (0, 0.05]$.** Here $L < \frac{1}{2}\log 2$, so no prime powers occur. Put $f(x) = L^{-1/2} f_1(x/L)$ with $\|f_1\|_{L^2(-1,1)} = 1$ and $F_1 = \hat{f_1}$. The substitution $t = \tau/L$ gives

$$Q_L(f) = \log \tfrac{1}{L} + R_L(f_1), \qquad R_L(f_1) = P_L(f_1) + \frac{1}{\pi}\int_0^\infty m_L(\tau)|F_1(\tau)|^2\, d\tau,$$

with $m_L(\tau) = \operatorname{Re}\psi(\frac{1}{4} + \frac{i\tau}{2L}) - \log\pi + \log L$, $P_L(f_1) = 2L\big(\int_{-1}^{1} f_1 e^{-Ls/2} ds\big)^2$ for even f_1 and $P_L(f_1) = -2L\big(\int_{-1}^{1} f_1 \sinh(Ls/2) ds\big)^2$ for odd f_1. The shift $\log(1/L)$ is common to all f_1, so the gaps of Q_L equal those of R_L, sector by sector, and R_L acts on the fixed space $L^2(-1,1)$. With $u = \tau/(2L)$,

$$\partial_L m_L = \tfrac{1}{L} h(u), \qquad h(u) = 1 - \operatorname{Re}\big(iu\,\psi'(\tfrac{1}{4} + iu)\big).$$

Lemma 5.1. *For $u > 0$, $|h(u)| \le 0.8125/u^2$.*

Proof. From $\psi'(w) = \sum_{n\ge 0}(n+w)^{-2}$ one gets $h(u) = 1 - \sum_{n\ge 0}\varphi(n+\frac{1}{4})$ with $\varphi(a) = 2au^2/(a^2+u^2)^2$ and $\int_0^\infty \varphi = 1$. The sum is the midpoint rule for $\int_{-1/4}^{\infty}\varphi$ with unit cells. The midpoint Peano kernel is bounded by $1/8$ and $\int_{\mathbb{R}} |\varphi''| = 6/u^2$, so the quadrature error is at most $\frac{3}{4}u^{-2}$; and $|\int_{-1/4}^{0}\varphi| \le \int_{-1/4}^{0} 2|a|u^{-2}da = \frac{1}{16}u^{-2}$. □

We bound $\int_0^\infty |h|$ by an upper Riemann sum on $[0,40]$ with cells of width $1/200$, enclosing $\sup|h|$ on each cell by evaluating h on a ball covering the cell, plus $0.8125/40$ from Lemma 5.1. At 768 bits this gives $\int|h| \le 0.593979$. Since $|F_1|^2 \le (\int|f_1|)^2 \le 2$, the multiplier part of $\frac{d}{dL}R_L(f_1)$ is at most $\frac{4}{\pi}\int|h|$. For the even pole term, with $J = \int f_1 e^{-Ls/2}$, $\frac{d}{dL}(2LJ^2) = 2J^2 + 4LJJ'$ and Cauchy–Schwarz gives $J^2 \le 2\sinh L/L$, $|J'| \le \frac{1}{2}(\frac{2}{3}e^L)^{1/2}$; for the odd pole term, with $K = \int f_1\sinh(Ls/2)$, $K^2 \le 2\sinh^2(L/2)$ and $|K'| \le \frac{1}{2}\cosh(L/2)\sqrt{2/3}$. All bounds increase in L; evaluated at $L_0 = 0.05$ they give, uniformly for $L \le L_0$,

$$C_{\mathrm{ev}} = \tfrac{4}{\pi}\int|h| + 2J^2 + 4L_0|J||J'| \le 4.876362, \qquad C_{\mathrm{odd}} \le 0.761666.$$

The forms R_L share the core $C_c^\infty(-1,1)$ and $|R_L(f_1) - R_{L'}(f_1)| \le C_\sigma|L-L'|\|f_1\|^2$ in sector σ; by min-max every sector eigenvalue of R_L is C_σ-Lipschitz in L. With the certified values $\ell_1, U, \ell_2, \ell^{\mathrm{odd}}$ at $L_0 = 0.05$ (Table 2), for $L \le L_0$,

$$\lambda_2^{\mathrm{ev}}(L) - \lambda_1^{\mathrm{ev}}(L) \ge \ell_2 - U - 2C_{\mathrm{ev}}L_0 \ge 0.4172,$$
$$\lambda_1^{\mathrm{odd}}(L) - \lambda_1^{\mathrm{ev}}(L) \ge \ell^{\mathrm{odd}} - U - (C_{\mathrm{ev}} + C_{\mathrm{odd}})L_0 \ge 0.6624.$$

Both are positive, which proves item (1) of Theorem 4.1. □

Remark 5.2. At 53 bits the same procedure gives the smaller enclosures $\int|h| \le 0.592773$, $C_{\mathrm{ev}} \le 4.87484$, $C_{\mathrm{odd}} \le 0.760130$; at 128 bits $\int|h| \le 0.593583$. All are upper bounds under Arb's containment guarantee and we use the largest. Why the higher-precision upper end is larger is not explained (§10).

5.2. **Per-window certificates on $[0.05, 1.19]$.** Fix L and a parity. We use the orthonormal Legendre basis $e_n(x) = c_n P_n(x/L)$, $c_n = \sqrt{(2n+1)/(2L)}$, with $n = 2m$ (even) or $n = 2m+1$ (odd). Up to a unit factor common to the sector, $\hat{e}_n(t) = c_n\, 2L\,(-1)^m j_n(Lt)$ with j_n the spherical Bessel function, and the pole vector is $p_m = \sqrt{2}\,c_n\,2L\,i_n(L/2)$, so that [pole] $= \pm(p\cdot a)^2$ for $f = \sum a_m e_{n(m)}$.

Lemma 5.3 (Binet bound)**.** *For $\operatorname{Re} w > 0$, $\big|\psi(w) - \log w + \frac{1}{2w}\big| \le \dfrac{1}{12|w|\operatorname{Re} w}$.*

Proof. Binet's formula reads $\psi(w) = \log w - \frac{1}{2w} - \int_0^\infty b(x)e^{-wx}dx$ with $b(x) = \frac{1}{e^x-1} - \frac{1}{x} + \frac{1}{2}$, $b(0) = 0$. With $y = x/2$, $b'(x) = \frac{1}{4}(y^{-2} - \sinh^{-2} y) \ge 0$. For $y < \sqrt{3}$ the inequality $b' \le \frac{1}{12}$ reduces to $\sinh^2 y \le 3y^2/(3-y^2)$, which holds coefficientwise $(2^{2k-1}/(2k)! \le 3^{1-k}$, with equality at $k = 1, 2$ and term ratio $4/((2k+1)(2k+2)) \le \frac{1}{3}$); for $y \ge \sqrt{3}$ it is trivial. Integrating by parts, $\int be^{-wx} = w^{-1}\int b'e^{-wx}$, of modulus at most $1/(12|w|\operatorname{Re} w)$. □

Lemma 5.4 (Envelope; [Zhu, Lemma 3.1]). *For $w = \frac{1}{4} + \frac{it}{2}$ and $t \geq 3/2$, $\operatorname{Re}\psi(w) \geq \log\frac{t}{2} - \frac{2}{3t} - \frac{1}{2t^2} \geq \log\frac{t}{2} - \frac{1}{t}$.*

Proof. Lemma 5.3 with $|w| \geq t/2$ and $\operatorname{Re}(1/(2w)) \leq 1/(2t^2)$; the second inequality is equivalent to $t \geq 3/2$. □

Let $A_L^{\sup} = \sum_{\log n < 2L} 2\Lambda(n)/\sqrt{n}$. The prime comb is $\geq -A_L^{\sup}$ and the envelope increases in t, so for $t \geq T^\sharp$

$$\Psi_L(t) \geq \beta := \log\frac{T^\sharp}{2\pi} - \frac{1}{T^\sharp} - A_L^{\sup}. \tag{5.1}$$

Lemma 5.5 (Frequency reduction; [Zhu, Thm. 1.1]). *Assume $\Psi_L \geq \beta > 0$ on $[T^\sharp, \infty)$ and let $\mathcal{R}(f) = [\text{pole}] + \frac{1}{\pi}\int_0^{T^\sharp}(\Psi_L - \beta)|F|^2 dt + \beta\|f\|^2$. Then $Q_L \geq \mathcal{R}$ as forms, hence $\lambda_k^\sigma(Q_L) \geq \lambda_k^\sigma(\mathcal{R})$ for all k and σ.*

Proof. Write $\Psi_L = (\Psi_L - \beta) + \beta$, use $\frac{1}{\pi}\int_0^\infty |F|^2 = \|f\|^2$, drop the nonnegative integral over $[T^\sharp, \infty)$ and apply min-max. □

Lemma 5.6 (Monotonicity of $\operatorname{Re}\psi$). *For $s > 0$, $\operatorname{Re}\psi(s+iy) = \psi(s) + \sum_{n\geq 0}\frac{y^2}{(n+s)((n+s)^2+y^2)}$ is increasing in $|y|$. Consequently $|\Psi_L - \beta| \leq M_\psi := \max(|\operatorname{Re}\psi(\frac{1}{4})|, |\operatorname{Re}\psi(\frac{1}{4} + \frac{iT^\sharp}{2})|) + \log\pi + A_L^{\sup} + \beta$ on $[0, T^\sharp]$.*

Proof. Take real parts in $\psi(w) = -\gamma + \sum_n\big(\frac{1}{n+1} - \frac{1}{n+w}\big)$. □

Lemma 5.7 (Bessel bounds). $|j_n(x)| \leq \dfrac{x^n}{(2n+1)!!}$ *and* $0 \leq i_n(x) \leq \dfrac{x^n}{(2n+1)!!}e^{x^2/2}$ *for* $x \geq 0$.

Proof. Use Poisson's integral and the beta integral,

$$j_n(x) = \frac{x^n}{2^{n+1}n!}\int_{-1}^{1}\cos(xs)(1-s^2)^n\,ds, \qquad \int_{-1}^{1}(1-s^2)^n\,ds = \frac{2^{2n+1}(n!)^2}{(2n+1)!};$$

for i_n replace cos by cosh $\leq e^{x^2/2}$. □

Lemma 5.8 (Gauss–Legendre and Bernstein; [25, Thm. 19.3]). *If g is analytic in the Bernstein ellipse E_ρ with $|g| \leq M$ there, then the n-point Gauss–Legendre rule on $[-1,1]$ has error at most $\frac{64}{15}M\rho^{-2n+2}/(\rho^2-1)$. A polynomial of degree d bounded by M on $[-1,1]$ is bounded by $M\rho^d$ on E_ρ.*

The cited theorem is stated with ρ^{-2n} for the $(n+1)$-point rule; the form above is the same statement for n points.

Remark 5.9 (Correction of a Gauss–Legendre constant). The original runs of the window certificates applied ρ^{-2n} to the n-point rule, so quadrature balls were too small by ρ^2 (about 4.33 for the head panels, 9 for the Rayleigh outer rule). The error was found in internal review and fixed in the code. The original certificate was first corrected by a Weyl shift of every lower bound by $\delta = N(\rho^2-1)q_{\text{old}}$ ($\delta \leq 6.4\cdot 10^{-35}$ for $L \leq 0.8$, $\leq 1.6\cdot 10^{-33}$ for $0.84 \leq L \leq 0.97$, $\leq 3.2\cdot 10^{-32}$ for $0.98 \leq L \leq 1$) and fresh upper bounds; that correction assumed unchanged window parameters between code revisions. It has since been superseded by a clean end-to-end rerun with the corrected code, which is the certificate of record used here and does not depend on the correction.

Head matrix. We compute $C_{jk} = \frac{1}{\pi}\int_0^{T^\sharp}(\Psi_L - \beta)\hat{e}_j\hat{e}_k\,dt$ by 64-point Gauss–Legendre on each unit panel in Arb. The ellipse has half-height $B = 2/5$, $\rho = 2B + \sqrt{4B^2+1}$. For complex t the continuation of $\operatorname{Re}\psi(\frac{1}{4} + \frac{it}{2})$ is $\frac{1}{2}(\psi(w_+) + \psi(w_-))$, $w_\pm = \frac{1}{4} \pm \frac{it}{2}$; on the strip $\operatorname{Re} w_\pm \geq \frac{1}{4} - \frac{B}{2} = 0.05$, and Lemma 5.3 bounds $\psi(w_\pm)$. On the strip $|\hat{e}_j| \leq \sqrt{2L}e^{LB}$ and the comb is bounded by $\sum 2\Lambda(n)n^{-1/2}\cosh(B\log n)$. The per-entry error is $T^\sharp \cdot \frac{32}{15}M\rho^{-126}/(\rho^2-1)$ and is added as a ball. The head is $A = C \pm pp^T + \beta I +$ (quadrature ball), with $+$ in the even and $-$ in the odd sector.

Tails. For tail orders $n \geq n_0$ put $X = LT^\sharp$; then $|\hat{e}_n| \leq \sigma_n := c_n 2LX^n/(2n+1)!!$ on $[0, T^\sharp]$ and $\sigma_{n+2}/\sigma_n \leq r = \sqrt{\frac{2n_0+5}{2n_0+1}}\frac{X^2}{(2n_0+3)(2n_0+5)} < 1$ (asserted), so $\sum\sigma^2 \leq \sigma_{n_0}^2/(1-r^2)$. The tail of the pole vector satisfies $\|p_{\text{tail}}\| \leq p_{\text{tail}} := \sqrt{2}c_{n_0}2Ly^{n_0}/(2n_0+1)!! \cdot e^{y^2/2}/\sqrt{1-y^2}$, $y = L/2$. Put

$$e_D = \frac{M_\psi}{\pi}T^\sharp\sum\sigma^2 + [\text{odd}]\,p_{\text{tail}}^2, \qquad e_B = \frac{M_\psi}{\pi}T^\sharp\sqrt{2L}\sqrt{N}\Big(\sum\sigma^2\Big)^{1/2} + 2\|p_{\text{head}}\|p_{\text{tail}}.$$

Then the tail block satisfies $D \geq (\beta - e_D)I$ (in the even sector the tail pole term $+pp^T \geq 0$ is dropped) and the coupling block satisfies $\|X\| \leq e_B$ by Hilbert–Schmidt bounds.

Lemma 5.10 (Block bound; [Zhu, Thm. 1.1, §6.2])**.** *Let* $\mathcal{R} = \begin{pmatrix} A & X \\ X^T & D \end{pmatrix}$ *with* $D \geq (\beta - e_D)I$ *and* $\|X\| \leq e_B$*. Then* $\lambda_k(\mathcal{R}) \geq \min(\lambda_k(A), \beta - e_D) - e_B$ *for every* k*.*

Proof. $\left\|\begin{pmatrix} 0 & X \\ X^T & 0 \end{pmatrix}\right\| = \|X\|$, so $\mathcal{R} \geq \operatorname{diag}(A, D) - e_B$. The k-th min-max value of $\operatorname{diag}(A, D)$ is at least $\min(\lambda_k(A), \lambda_1(D))$; conclude with Weyl's inequality. □

Verified Cholesky. To show $A - \mu I \succeq 0$ for a ball matrix A, set $d = \mu/100$, compute an approximate Cholesky factor R of $\operatorname{mid}(A) - (\mu + d)I$ in midpoint arithmetic, compute $E = A - (\mu+d)I - RR^T$ rigorously in ball arithmetic, and check $\|E\|_F < d$; then $A - \mu I = RR^T + E + dI \succeq 0$ for every matrix in the ball (cf. [21]). Eigenvalue estimates from inverse iteration on midpoints are used only to choose μ and the deflation vector and carry no rigour. With Lemmas 5.5 and 5.10 this gives $\lambda_1^{\text{ev}} \geq \min(\mu_1, \beta - e_D) - e_B$, $\lambda_1^{\text{odd}} \geq \min(\mu^{\text{odd}}, \beta - e_D^{\text{odd}}) - e_B^{\text{odd}}$, and $\lambda_2^{\text{ev}} \geq \min(\mu_2, \beta - e_D) - e_B$ where μ_2 is certified for $A + vv^T$ (Lemma 5.14 below).

Parameters. The cutoff is $T^\sharp = \max(200, 5\lceil 80/(5L)\rceil, 5\lceil 2\pi e^{A_L^{\sup}+1/2}/5\rceil)$ and the head has $N = \lceil eLT^\sharp/4\rceil + 42$ modes per sector. For $L > 1$ ("sharp cutoff") we use instead $T^\sharp = 2450$ ($L = 1.03$), 4000 ($L = 1.06, 1.09$), 5550 ($L = 1.12$–1.19), the exact dyadic floor $\beta = 3/16$, 112 Gauss nodes per unit panel and $N = \lceil eLT^\sharp/4\rceil + 80$. Lemma 5.4 is applied above the crude cutoff T_2 (asserting $\log(T_2/2\pi) - 1/T_2 - A_L^{\sup} \geq 3/16$), and $\Psi_L > \beta$ on $[T^\sharp, T_2]$ is certified by Arb enclosures of Ψ_L on cells of width 0.01 with bisection (minimum width 10^{-9}). Together $\Psi_L \geq \beta$ on $[T^\sharp, \infty)$, which is all Lemma 5.5 needs. This device is a refinement inside Zhu's reduction; we make no novelty claim for it. Validation at $L = 1$: the sharp path gives $4.978 \cdot 10^{-30} \leq \lambda_1^{\text{ev}} \leq 9.406 \cdot 10^{-30}$, $\lambda_2^{\text{ev}} \geq 1.808 \cdot 10^{-23}$, $\lambda_1^{\text{odd}} \geq 1.255 \cdot 10^{-26}$, overlapping the crude-cutoff values of Table 2.

Spherical Bessel values for $L \geq 0.98$. Orders $n \leq n_1 = \lfloor x\rfloor$ use upward recurrence in ball arithmetic from Arb seeds every 400 orders; orders $n > n_1$ use the ratios $r_k = j_k/j_{k-1}$ from the backward continued fraction $r_k = x/(2k + 1 - xr_{k+1})$, started from $[0, 1]$ and anchored at j_{n_1}.

Lemma 5.11 (Ratio enclosure)**.** *For $x > 0$ and $k \geq \lfloor x\rfloor + 1$, every true ratio r_k lies in $(0, 1]$; hence the interval iteration started from $[0, 1]$ encloses it.*

Proof. If $2k+1 \geq 2x$, the map $\tau \mapsto x/(2k+1-x\tau)$ sends $[0,1]$ into $[x/(2k+1), x/(2k+1-x)] \subset [0,1]$. By downward induction from large K (where $j_K(x) > 0$ and $r_K \in [0,1]$): if $2k+1 \geq 2x$ and $r_{k+1} \in [0,1]$, then $j_{k-1}/j_k = (2k+1)/x - r_{k+1} \geq 1$, so $r_k \in (0,1]$ and $j_{k-1} > 0$. The condition holds for all $k \geq \lfloor x \rfloor + 1$. □

The enclosures need not be sharp at the top orders unless the start index is well above $n_{\max}$ [12]; sharpness only affects the size of the radii.

Upper bound U. The trial function is $f = \sum_{m<m^*} v_m e_{2m}$, a polynomial on $[-L, L]$, with v an approximate ground vector (exact dyadic entries). In $s = x/L$, $g(Ls) = L\int_{-1}^{1-s} f(y)f(y+s)dy$ is computed exactly by $(D+1)$-point Gauss–Legendre ($D = \deg f$); the integral in (2.2) by 450-point Gauss–Legendre on $[0,2]$ with error bounded by Lemma 5.8 ($\rho = 3$, exponent ρ^{-898}, ellipse maximum enclosed by 2048 ball arcs with $|1 - e^{-2Lz}| > 0$ asserted on each arc; the zeros $i\pi k/L$, $k \neq 0$, lie outside the ellipse); the other terms of (2.2) exactly. Then $U \geq Q_L(f)/\|f\|^2 \geq \lambda_1^{\mathrm{ev}}(L)$. A polynomial on $[-L, L]$ has $|\hat{f}|^2 = O(t^{-2})$ and lies in the form domain $\{f : \int \log(2+t)|\hat{f}|^2 < \infty\}$, and in the closure of $C_c^\infty(-L, L)$ there, since the weight is weaker than any H^s, $s < \frac{1}{2}$.

5.3. **Piece II: the grid on** $[0.05, 1.19]$**.** The grid is

$$0.05,\ 0.1,\ 0.2,\ 0.3,\ 0.35,\ 0.4,\ 0.45,\ 0.5,\ 0.55,\ 0.6,\ 0.64,\ 0.68,\ 0.72,\ 0.76,\ 0.8,\ 0.84,$$
$$0.87,\ 0.9,\ 0.93,\ 0.96,\ 0.97,\ 0.98,\ 0.99,\ 1.0,\ 1.03,\ 1.06,\ 1.09,\ 1.12,\ 1.15,\ 1.18,\ 1.19.$$

For each of the 30 steps the criterion (3.1) holds (Table 2); the smallest ratios are 1.288 on $[0.05, 0.1]$, 1.320 on $[0.1, 0.2]$ and, past $L = 1$, 2.148 on $[1.15, 1.18]$. By Corollary 3.2 this proves item (2) of Theorem 4.1. □

Remark 5.12 (Comparisons and dyadic windows). (a) The gluing script compares balls with Python's built-in `min`, which is not a rigorous minimum when balls overlap; here the two lower bounds are disjoint at every window (relative separation ≥ 1.021, radii $< 5 \cdot 10^{-20}$), and the step ratios in Table 2 were recomputed with explicit endpoint comparisons. (b) Each window is certified at the dyadic number $\tilde{L}_i \geq L_i$ with $\tilde{L}_i < L_i + 2^{-p}$, $p \geq 768$, for all four bounds of the window; the covering is then $(0, \tilde{L}_1] \cup \bigcup_i [\tilde{L}_i, \tilde{L}_{i+1}] \supset (0, 1.19]$.

5.4. **Piece III: the grid on** $(1.19, 1.4035]$**.** Beyond $L \approx 1.2$ the route of §5.2 is impractical: the cutoff $T^\sharp \gtrsim 2\pi e^{A_L^{\sup}}$ forced by the pointwise bound (5.1) is about $2 \cdot 10^5$ at $L = 1.3$ and $5 \cdot 10^6$ at $L = 1.5$, and Zhu [Lemma 3.2] shows that $A_L^{\sup}$ is optimal for any pointwise bound. Elston [11] replaces the pointwise bound by an operator bound for the prime comb on a slightly widened window. We state his reduction as a cited proposition; we use his code unchanged except for the deflation option described below.

Notation. Let $a = L$ be the window, $\delta > 0$, $b = a + \delta$, $\beta_K > 0$, and $T^\sharp > \beta_K/\delta$ with $\Omega := T^\sharp - \beta_K/\delta$ and $\Omega\delta \geq 1$. Let $W(u) = I_0(\beta_K\sqrt{1 - u^2/\delta^2})/I_0(\beta_K)$ on $[-\delta, \delta]$ (Kaiser window, 0 outside), $\kappa(u) = \frac{\sin \Omega u}{\pi u} W(u)$ and $m = 1 - \hat{\kappa}$. Let $H(t) = \operatorname{Re}\psi(\frac{1}{4} + \frac{it}{2}) - \log\pi$, $P(t) = \sum_{\log n < 2a} \frac{2\Lambda(n)}{\sqrt{n}} \cos(t \log n)$, so $\Psi_a = H - P$, and $A = \sum_{\log n < 2a} 2\Lambda(n)/\sqrt{n}$. Let $C_b = \sum_{\log n < 2a} \Lambda(n) n^{-1/2}(\tau_{\log n} + \tau_{-\log n})$, compressed to $L^2(-b, b)$, where τ_s is translation by s.

Proposition 5.13 (Elston [11])**.** *Assume*

(H1) $\bar{\mu} \geq \sup\{\langle h, C_b h\rangle/\|h\|^2 : 0 \neq h \in L^2(-b, b)\}$;
(H2) $\varepsilon \geq \sup_{t \geq T^\sharp} |\hat{\kappa}(t)|$ *(in* [11]*,* $\varepsilon = 2/(\pi\beta_K I_0(\beta_K))$*);*
(H3) $0 < \beta' \leq \log \frac{T^\sharp}{2\pi} - \frac{1}{T^\sharp} - \bar{\mu} - (2\varepsilon + \varepsilon^2)(A + \bar{\mu})$.

Put $\Phi = \Psi_a + m^2(P - \bar{\mu})$ *and*

$$R'(f) = [\text{pole}] + \frac{1}{\pi}\int_0^{T^\sharp} (\Phi - \beta')|F|^2\,dt + \beta'\|f\|^2.$$

Then:

(a) $Q_a \geq R'$ *as forms on each parity sector.*

(b) *Let* A_N *be the matrix of* R' *on the first* N *Legendre modes of the sector, and let* $\varepsilon_D, \varepsilon_B$ *be the tail constants of Zhu's reduction computed with* $G \geq \sup_{[0,T^\sharp]} |\Phi - \beta'|$ *(Bessel tail bounds as in Lemma 5.7;* ε_D *by a Gershgorin bound on the tail rows,* ε_B *by a Schur test). Then the tail block of* R' *is* $\succeq (\beta' - \varepsilon_D)I$ *and the coupling block has norm* $\leq \varepsilon_B$.

(c) *The certifier encloses every entry of* A_N *in a ball (Gauss–Legendre nodes and weights,* Ψ *via the complex digamma function,* m *by rigorous integration of the Kaiser main lobe and a mean-value bound for the side lobes,* j_n *by recurrence and ratio enclosure), adds a spectral quadrature debit* c_{err} *from a Bernstein-ellipse bound on panels of width* $1/4$*, and certifies a lower bound* $\lambda_1(A_N) \geq \mu' - \rho_{\mathrm{res}} - N r_{\max} - c_{\mathrm{err}}$*, where* μ' *is a dyadic shift,* ρ_{res} *the* ∞*-norm of an exactly computed integer Cholesky residual divided by the scale* 2^{2P} *(plus* $N2^{-2P}$ *for the floors), and* $r_{\max}$ *the largest entry radius.*

Proof of (a), following [11]. Put $h = f - f * \kappa$; then $\operatorname{supp} h \subset [-b, b]$ and $\hat{h} = mF$. By (H1), $\frac{1}{\pi}\int_0^\infty Pm^2|F|^2 = \langle h, C_b h\rangle \leq \bar{\mu}\|h\|^2 = \frac{1}{\pi}\int_0^\infty \bar{\mu}m^2|F|^2$, so $\frac{1}{\pi}\int m^2(\bar{\mu} - P)|F|^2 \geq 0$ and $Q_a(f) \geq [\text{pole}] + \frac{1}{\pi}\int_0^\infty \Phi|F|^2$. For $t \geq T^\sharp$, $\Phi = H - \bar{\mu} - (1-m^2)(P + \bar{\mu})$ with $|1 - m^2| \leq 2\varepsilon + \varepsilon^2$ by (H2), so Lemma 5.4 and (H3) give $\Phi \geq \beta'$; dropping $\frac{1}{\pi}\int_{T^\sharp}^\infty (\Phi - \beta')|F|^2 \geq 0$ gives $Q_a \geq R'$. □

Elston states (b)–(c) for the lowest eigenvalue; in [11], $\bar{\mu}$ is certified by a weighted Schur test on 4000–8000 cells with a Collatz–Wielandt weight, and the finite certificate is an exact integer Gram residual in the spirit of [21]. An independent audit of the code at $L = 1.19$ (documented in [11]) found and fixed two implementation issues; the code used here is the post-audit version (upstream commit `3cc4cb7b`).

Lemma 5.14 (Rank-one deflation)**.** *Let* A *be a real symmetric* $N \times N$ *matrix and* $w \in \mathbb{R}^N$. *Then* $\lambda_2(A) \geq \lambda_1(A + ww^T)$. *In particular, if* $A + ww^T - \mu I \succeq 0$ *then* A *has at most one eigenvalue below* μ.

Proof. By Courant–Fischer, $\lambda_2(A) = \max_{y \neq 0} \min\{x^T A x : \|x\| = 1,\, x \perp y\} \geq \min\{x^T A x : \|x\| = 1,\, x \perp w\} = \min\{x^T(A + ww^T)x : \|x\| = 1,\, x \perp w\} \geq \lambda_1(A + ww^T)$. (For $w = 0$ the claim is trivial.) □

Corollary 5.15 (Second even eigenvalue)**.** *Under the hypotheses of Proposition 5.13, in the even sector and for any* $w \in \mathbb{R}^N$,

$$\lambda_2^{\mathrm{ev}}(L) \;\geq\; \lambda_2(R'^{\mathrm{ev}}) \;\geq\; \min\bigl(\lambda_1(A_N + ww^T),\; \beta' - \varepsilon_D\bigr) - \varepsilon_B.$$

Proof. By Lemma 5.14, $\lambda_2(A_N) \geq \lambda_1(A_N + ww^T)$. Lemma 5.10 with $k = 2$ (Weyl's inequality applied to the off-diagonal blocks, with the bounds of Proposition 5.13(b)) gives $\lambda_2(R') \geq \min(\lambda_2(A_N), \beta' - \varepsilon_D) - \varepsilon_B$. Since $Q_L \geq R'$ as forms (Proposition 5.13(a)), min-max gives $\lambda_2^{\mathrm{ev}}(L) \geq \lambda_2(R'^{\mathrm{ev}})$. □

Implementation of the deflation (option `--deflate`; our only change to [11]). After the ball matrix A_N is assembled, a vector v approximating the unit lowest eigenvector of $\mathrm{mid}(A_N)$ is computed by 12 steps of inverse iteration; its entries are rounded to 20 significant decimal digits and converted to Arb balls. The matrix $A_N + vv^T$ is formed in ball arithmetic, so it encloses $A_N + ww^T$ for the exact midpoint vector w of v, and the rounding of v enters the

largest entry radius $r_{\max}$ exactly like the Gram radii. The certifier then runs unchanged on $A_N + vv^T$, and Proposition 5.13(c) yields a certified lower bound for $\lambda_1(A_N + ww^T)$. The choice of v affects only sharpness, not rigour.

One form, two codes. The step test (3.1) compares lower bounds from Elston's certifier with upper bounds from our code, so both must refer to the same form with the same norm. Both implement (1.1) in Zhu's normalisation ($\|f\|^2 = \frac{1}{\pi}\int_0^\infty |F|^2$ for real f, pole term $+2F(\frac{i}{2})^2$ in the even and $-2|F(\frac{i}{2})|^2$ in the odd sector). As a consistency check, at $L = 1.19$ Elston's certified even lower bound $6.8131 \cdot 10^{-48}$, our Piece II bracket $[7.8 \cdot 10^{-48}, 1.312 \cdot 10^{-47}]$ and the Legendre–Galerkin value $8.676 \cdot 10^{-48}$ of §7 are mutually compatible; in the odd sector Elston's certified lower bound $3.9197 \cdot 10^{-44}$ lies below our rigorous Galerkin upper bound $4.8461 \cdot 10^{-44}$ ($K = 256$). The proof does not rely on these numerical agreements; it relies only on both codes evaluating (1.1).

Upper bounds $U(L_i)$. At every grid point L_i, including $L_i = 1.19$, $U(L_i)$ is the outward-rounded upper endpoint of a Rayleigh quotient $v^T Q_K v / v^T v$ of the even Legendre–Galerkin matrix Q_K ($K = 640$ even modes, 3000 bits) with an exact dyadic vector v, computed in ball arithmetic with the rigorous quadrature bound of §7 (Proposition 7.1). At $L = 1.19$ this gives $U(1.19) \le 8.712 \cdot 10^{-48}$, sharper than the $1.312 \cdot 10^{-47}$ of Piece II (Table 2). The sharper value is needed: with $1.312 \cdot 10^{-47}$ the first step ratio would be 0.93. Both are valid upper bounds, and the gluing script uses the smaller one at each window.

Grid. The planned grid to $L = 1.5$ is

$$1.19,\ 1.2205,\ 1.25,\ 1.2785,\ 1.3055,\ 1.3315,\ 1.3565,$$
$$1.3805,\ 1.4035,\ 1.4255,\ 1.447,\ 1.468,\ 1.4885,\ 1.5.$$

It was chosen from non-rigorous Galerkin estimates so that $0.95 \min(\lambda_2^{\mathrm{ev}}, \lambda_1^{\mathrm{odd}})(L_{i+1}) \ge 1.25 \cdot 1.03 \cdot \lambda_1^{\mathrm{ev}}(L_i)$. In this release the windows $1.2205, \dots, 1.4035$ are complete in both sectors, as are the odd sector at 1.4255 and both sectors at 1.447; the other runs are in progress. The parameters actually used are in Table 3 and in the field `args` of every certificate file. They follow a design rule (`code/gap/params.py`) that plays no role in the proof: $\Omega = T^\sharp - \beta_K/\delta$ is interpolated log-linearly from Elston's runs at $L = 1.19, 1.3, 1.495$ and inflated by 3%; $N \approx 0.8L(\Omega + 64/\delta) + 20$; the number of Gauss points per panel is raised so that the quadrature debit is at most 10^{-3} times the target eigenvalue (computed in the certifier's plan mode); and β_K is the smallest value for which N times the side-lobe entry radius stays below 10^{-6} times the target eigenvalue. The last rule was added after a first pass with $\beta_K = 64$ at every window gave a negative odd bound at 1.2785; the runs from that pass that gave useful bounds (1.2205, 1.25, and the even sector at 1.2785) are kept. Grid values are passed to the certifier as decimal strings, converted to Arb balls that contain the exact decimal; the Schur window b is rounded up, which is admissible because the supremum in (H1) is monotone in b.

Conclusion of Piece III. Each of the 8 steps from 1.19 to 1.4035 satisfies (3.1) (Table 4); the smallest ratio is 1.131, on $[1.2205, 1.25]$, and the odd sector binds at every window. By Corollary 3.2 this proves item (3) of Theorem 4.1. □

The step $[1.4255, 1.447]$ also satisfies (3.1) (ratio ≥ 1.379), so the ground state is simple and even on that interval too. We report it as a data point and not as part of Theorem 4.1, since $[1.4035, 1.4255]$ is not yet covered. Nothing is claimed for $L \in (1.4035, 1.4255)$ or $L > 1.447$.

6. Results

6.1. **Pieces I and II.** Table 2 lists the certificate of record on $[0.05, 1.19]$. Lower bounds are rounded down and upper bounds up, to four significant digits; the step ratio in row L_{i+1}

is $\min(\ell_2, \ell^{\mathrm{odd}})(L_{i+1})/U(L_i)$, lower end floored. The odd sector binds at every window from $L = 0.1$ on. Piece I on this certificate: $\lambda_2^{\mathrm{ev}} - \lambda_1^{\mathrm{ev}} \geq 0.4172$ and $\lambda_1^{\mathrm{odd}} - \lambda_1^{\mathrm{ev}} \geq 0.6624$ on $(0, 0.05]$.

TABLE 2. Certificate of record on $[0.05, 1.19]$. $T^\sharp$ and N are the cutoff and the head size per sector. At $L = 1$ the crude cutoff is used (the sharp-path validation is in §5.2).

L	$T^\sharp$	N	$\lambda_1^{\mathrm{ev}} \geq$	$\lambda_1^{\mathrm{ev}} \leq U$	$\lambda_2^{\mathrm{ev}} \geq$	$\lambda_1^{\mathrm{odd}} \geq$	step ratio $\geq$
0.05	1600	97	0.9351	0.9827	1.887	1.926	(Piece I)
0.1	800	97	0.4348	0.4572	1.387	1.266	1.288
0.2	400	97	0.09014	0.09499	1.046	0.6037	1.320
0.3	270	98	7.143e-3	7.587e-3	0.7437	0.2110	2.222
0.35	230	97	1.110e-3	1.204e-3	0.6153	0.06086	8.022
0.4	200	97	1.662e-4	1.841e-4	0.2866	0.01361	11.31
0.45	200	104	1.465e-5	1.647e-5	0.08888	2.205e-3	11.98
0.5	200	110	8.482e-7	9.506e-7	0.01650	1.752e-4	10.63
0.55	200	117	4.424e-8	6.066e-8	1.234e-3	1.213e-5	12.76
0.6	200	124	1.321e-9	1.813e-9	8.661e-5	4.921e-7	8.114
0.64	200	129	7.330e-11	1.036e-10	5.914e-6	3.008e-8	16.59
0.68	200	135	1.984e-12	2.832e-12	3.870e-7	1.180e-9	11.39
0.72	200	140	5.422e-14	1.207e-13	1.013e-8	3.167e-11	11.18
0.76	200	146	7.533e-16	1.802e-15	3.574e-10	7.456e-13	6.181
0.8	200	151	9.787e-18	3.681e-17	4.895e-12	1.086e-14	6.031
0.84	830	516	1.134e-19	1.757e-19	8.754e-14	1.322e-16	3.593
0.87	830	533	2.636e-21	4.329e-21	2.568e-15	3.892e-18	22.15
0.9	830	550	3.401e-23	5.545e-23	5.866e-17	5.819e-20	13.44
0.93	830	567	4.303e-25	7.594e-25	8.697e-19	9.536e-22	17.20
0.96	830	584	4.510e-27	7.555e-27	9.996e-21	8.439e-24	11.11
0.97	830	590	8.936e-28	1.613e-27	2.109e-21	1.929e-24	255.3
0.98	3610	2447	1.682e-28	2.165e-28	5.262e-22	4.471e-25	277.2
0.99	3610	2471	2.782e-29	3.552e-29	1.101e-22	7.684e-26	354.9
1.0	3610	2496	4.531e-30	6.601e-30	1.649e-23	1.141e-26	321.5
1.03	2450	1795	2.096e-32	4.110e-32	1.055e-25	6.604e-29	10.00
1.06	4000	2962	7.275e-35	1.249e-34	4.435e-28	2.691e-31	6.548
1.09	4000	3043	1.306e-37	2.162e-37	1.034e-30	4.711e-34	3.774
1.12	5550	4305	1.903e-40	3.001e-40	2.049e-33	8.502e-37	3.933
1.15	5550	4418	1.752e-43	2.799e-43	2.501e-36	8.572e-40	2.857
1.18	5550	4531	9.792e-47	1.636e-46	2.088e-39	6.011e-43	2.148
1.19	5550	4569	7.823e-48	1.312e-47	1.439e-40	4.329e-44	264.7

At $L = 0.8$ our certified values are consistent with Zhu's [Thm. 6.2]: $8.9 \cdot 10^{-18} \leq \lambda_1^{\mathrm{ev}} \leq 2.523 \cdot 10^{-16}$, $\lambda_2^{\mathrm{ev}} \geq 2.085 \cdot 10^{-12}$, $8.206 \cdot 10^{-15} \leq \lambda_1^{\mathrm{odd}} \leq 2.347 \cdot 10^{-14}$. Zhu's certified upper bound $\lambda_1^{\mathrm{ev}}(0.8) \leq 2.27 \cdot 10^{-17}$ [Table 3] is sharper than our $3.681 \cdot 10^{-17}$.

6.2. **Piece III.** *Test at* $L = 1.19$. With Elston's published parameters at $L = 1.19$ ($\delta = 0.3$, $\beta_K = 64$, $T^\sharp = 700$, $N = 680$, 56 Gauss points on panels of width $1/4$) the certifier, run by us, gives

$$\lambda_1^{\mathrm{ev}}(1.19) \geq 6.8131164696015 \cdot 10^{-48}, \qquad \lambda_1^{\mathrm{odd}}(1.19) \geq 3.91966861609542239 \cdot 10^{-44},$$

reproducing Elston's published $6.8131164696007 \cdot 10^{-48}$ (even, 12 digits) and his odd bound to 17 digits; the shift μ_s depends on the thread count through the numerical eigenvalue estimate, and both values are valid lower bounds. With `--deflate`,

$$\lambda_2^{\mathrm{ev}}(1.19) \geq 1.2336395547 \cdot 10^{-40}$$

(non-rigorous Galerkin value $1.23364079 \cdot 10^{-40}$). With $U(1.19) \le 1.312 \cdot 10^{-47}$ from Table 2, $\min(\lambda_2^{\mathrm{ev}}, \lambda_1^{\mathrm{odd}})(1.19)/U(1.19) \ge 2987$: simple-even at $L = 1.19$ is confirmed by a second, independent code path (Elston's reduction instead of Zhu's pointwise cutoff, a different Gram assembly and a different Cholesky certificate).

TABLE 3. Piece III windows and certifier parameters, from the field `args` of each certificate file. Common to all runs: panel width 1/4, Schur test on 4000 cells with 1500 weight iterations. "GL" is the number of Gauss–Legendre points per panel, "bits" the working precision, "time" the wall-clock seconds (field `seconds`) of the λ_2^{ev} (deflated) and λ_1^{odd} runs, with 15–52 worker processes. At 1.2785 the two sectors used different parameters. New prime powers relative to the previous row are listed. Runs not listed are in progress.

L_i	δ	β_K	$T^\sharp$	N	GL	bits	time (s), λ_2^{ev} / λ_1^{odd}	new n
1.19	0.3	64	700	680	56	400	600–720 each	(test; $n \le 9$)
1.2205	0.3	64	850	849	56	400	1009 / 714	11
1.25	0.3	64	990	1010	56	400	1481 / 1034	
1.2785, even	0.3	64	1150	1196	57	400	2418 / —	
1.2785, odd	0.3	74	1130	1133	63	400	— / 1000	
1.3055	0.3	78	1290	1315	65	400	3301 / 2466	13
1.3315	0.2	84	1570	1585	70	400	5673 / 4384	
1.3565	0.2	89	1730	1757	74	600	11116 / 8545	
1.3805	0.2	93	1890	1943	78	600	11639 / 7998	
1.4035	0.2	98	2070	2143	81	600	14363 / 9784	16
1.4255	0.2	102	2240	2356	85	600	running / 13217	17
1.447	0.2	107	2440	2587	89	600	24322 / 20970	

TABLE 4. Piece III certificate. $U(L_i) \ge \lambda_1^{\mathrm{ev}}(L_i)$ is the $K = 640$ Galerkin bound; ℓ^{odd} and ℓ_2 are certified lower bounds for λ_1^{odd} and λ_2^{ev}; the step ratio in row L_{i+1} is $\min(\ell_2, \ell^{\mathrm{odd}})(L_{i+1})/U(L_i)$. Lower bounds are rounded down and upper bounds up, to four significant digits, and ratios are floored. The lower bounds in row 1.19 are the test of §6.2 and enter no step. Below the rule: additional certified data points beyond 1.4035, not part of Theorem 4.1 (the λ_2^{ev} run at 1.4255 is in progress).

L_i	$U(L_i)$	$\lambda_1^{\mathrm{odd}} \ge$	$\lambda_2^{\mathrm{ev}} \ge$	step ratio $\ge$
1.19	8.712e-48	3.91966861609542239e-44	1.2336395547e-40	(Piece II: 264.7)
1.2205	2.336e-51	1.224e-47	4.200e-44	1.405
1.25	4.458e-55	2.642e-51	1.162e-47	1.131
1.2785	8.317e-59	5.889e-55	2.648e-51	1.321
1.3055	1.209e-62	9.466e-59	4.806e-55	1.138
1.3315	1.693e-66	1.706e-62	9.901e-59	1.412
1.3565	1.962e-70	2.187e-66	1.435e-62	1.292
1.3805	2.251e-74	3.032e-70	2.247e-66	1.546
1.4035	2.493e-78	3.492e-74	2.991e-70	1.552
1.4255	2.371e-82	3.564e-78	(running)	—
1.447	2.107e-86	3.269e-82	3.420e-78	1.379

7. Rigorous upper bounds for larger windows

This section gives *upper bounds only*; it makes no lower-bound, positivity or simplicity statement.

Galerkin matrix. Let $e_k(y) = \sqrt{(2k+1)/(2L)}P_k(y/L)$ with $k = 2m+p$, $m < K$, $p \in \{0,1\}$ the parity. With $r = x/L \in [0,2]$, the symmetrised autocorrelation of e_a, e_b is an L-independent polynomial in r: up to the normalisation $\sqrt{(2a+1)(2b+1)}/2$ it equals $C_{ab}(r) = \int_{-1}^{1-r} P_a(t)P_b(t+r)\,dt$ (for $a+b$ even $C_{ab} = C_{ba}$). By (2.2) the Galerkin matrix is $Q_K = \big(\int_0^2 C_{ab}\,d\nu_L\big)_{a,b}$ plus a multiple of the identity, where ν_L is the measure with smooth density (pole and archimedean weights) minus the point masses $2\Lambda(n)n^{-1/2}$ at $r = \log n/L$. The entries follow from the Legendre moments $\int P_j(r-1)\,d\nu_L$ by the convolution recurrence [14]

$$C_{a+1,b} = C_{a-1,b} + \frac{2a+1}{2b+1}\big(C_{a,b-1} - C_{a,b+1}\big),$$

with exact seeds C_{0b}, C_{1b} expressed through $\int_u^1 P_k$. The recurrence is run only on $b \ge a$, where it is forward stable; the cost is $O(K^2)$ instead of $O(K^4)$ for direct quadrature.

Proposition 7.1 (Rigorous moments)**.** *The smooth moments are computed with M_q-point Gauss–Legendre in r, and the truncation error of each is bounded by Lemma 5.8: $|P_j| \le \rho^j$ on E_ρ, the maxima of the weights $4L\cosh(Lr/2)$, $2Lr\,e^{-Lr/2}/(1-e^{-2Lr})$ and of the diagonal weight are enclosed by covering the ellipse with 2000 complex balls, and the poles $r = i\pi m/L$, $m \ne 0$, are checked to lie outside E_ρ. The Legendre values at the nodes are computed by the Chebyshev recurrence on exact midpoints with an a priori rounding bound and a Markov bound for the node radii, followed by the nonnegative Chebyshev-to-Legendre map. Hence every entry of Q_K is enclosed in a ball, and for every exact vector v the upper endpoint of v^TQ_Kv/v^Tv is an upper bound for $\lambda_1^\sigma(L)$ in the sector of v.*

The last statement holds because v defines a polynomial trial function in the form domain (§5.2). The vector v comes from inverse iteration on midpoints and carries no rigour; the reported value is outward-rounded.

Table 5. Rigorous upper bounds for the lowest eigenvalue in each sector. K = number of Legendre modes per sector; "—" = no rigorous run. The Piece III windows are in Table 4 ($K = 640$, 3000 bits).

L	$\lambda_1^{\mathrm{ev}}(L) \le$	$\lambda_1^{\mathrm{odd}}(L) \le$	K, bits (even; odd)
1.5	3.85105826960019e-97	8.21780562022764e-93	384, 2400; 512, 4000
1.75	—	2.06808416452444e-162	—; 512, 4000
2.0	1.84681871820091e-283	3.23821041502563e-278	1024, 6000; 1024, 6000
2.25	9.03204367909380e-476	4.64181809939520e-470	2048, 8000; 1792, 8000

At $L = 2$ our even bound is below Zhu's certified $3.19 \cdot 10^{-283}$ [Table 3]; certified upper bounds for $L \le 2$ are not new. The window $L = 2.25$ lies outside the range of [28]. Non-rigorous Galerkin values continue to decrease slowly with K (at $L = 2.25$: $9.894 \cdot 10^{-476}$, $9.337 \cdot 10^{-476}$, $9.032 \cdot 10^{-476}$ for $K = 1536, 1792, 2048$); we do not have a bound on the omitted-mode tail, so these upper bounds may be far from sharp in relative terms.

Remark 7.2 (Quadrature exponent). The $L = 2$ even run was made before the exponent of Lemma 5.8 was corrected from ρ^{-2n} to ρ^{-2n+2} in this code path ($\rho \le 3$ there). The quadrature contributions to the moment radii are below 10^{-1999} and the reported radius of the Rayleigh

quotient is $4.5 \cdot 10^{-1656}$; scaling the quadrature radii by $\rho^2 \le 9$ cannot reach the slack of the 15-digit outward rounding (order 10^{-297}). A re-run with the corrected exponent has not been made. The bounds U of Table 4 were computed on 30 October 2026 with `fr_sweep.py` as shipped in `code/upper/`: the exported value is the upper end of the Rayleigh-quotient ball raised by a relative 10^{-13} before rounding to 15 digits, while every ball radius is below 10^{-600} times the value, so a factor $\rho^2 \le 9$ on the quadrature part of the radius cannot reach that margin, whichever exponent was in use. The working directory was not under version control, so the revision of `fast_recurrence.py` used for each run is not recorded. Re-running $U(1.19)$ with the shipped code on 31 October 2026 reproduced every exported digit and the radius (`data/piece3/rerun_U1.19.jsonl`). For the other rows of Table 5 (28–30 October; by-products that enter no proof) we have not confirmed which exponent was used.

8. A numerical observation: prime-power singular points

Everything in this section is numerical and non-rigorous. It concerns finite-dimensional approximants of the even minimiser; whether the limiting eigenfunction has the same features is not established. That near-null eigenvectors have weak singularities at $\log n$ and $L - \log n$, and that meshes should be cut there, is due to Eberwein [9, §13], whose spectral-element code already does so; the mechanism is the one familiar from differential-difference equations. Only the measurements below are ours. Note also that second-generation points such as $L - \log 6$ exist, so the singular set is not limited to $|L - \log n|$ with n a prime power.

At $L = 1.19$ the Legendre coefficients of the lowest Galerkin eigenvector ($K = 384$) fall exponentially to about 10^{-25} by degree 160 and then only like $(\text{degree})^{-2.1}$. A periodogram of the high-degree coefficients locates singular points at $|y| \approx 0.088$–0.105, 0.183–0.198, 0.406, 0.510, 0.747, 0.884, 1.003, which match $|L - \log n| = 0.091$, 0.196, 0.419, 0.497, 0.756, 0.889, 1.007 for $n = 3, 4, 5, 2, 7, 8, 9$; there is no signal at $|L - \log 6| = 0.602$. A heuristic reason: in the eigen-equation the shifted copies $f(y \pm \log n)$ (with $f = 0$ outside $[-L, L]$) switch on or off where $y \pm \log n$ crosses $\mp L$, and only prime powers carry weight ($\Lambda(6) = 0$).

To quantify this we use a split-window basis (piecewise Legendre, discontinuous at $\pm(L - \log n)$; its x-quadrature error is checked by node doubling but not bounded) and record the one-sided jumps of the computed eigenvector at the breakpoints, normalised by $\Lambda(n)n^{-1/2}f(L)$. Table 6 gives the absolute values at $L = 1.19$ for four resolutions. The signs are negative for $n = 2, 3$ (where $L - \log n > 0$) and positive otherwise, as the heuristic predicts.

Table 6. $|\text{jump}|/(\Lambda(n)n^{-1/2}f(L))$ at $y = L - \log n$ (or $\log n - L$), $L = 1.19$, even sector, split-window basis with base allocation 24; "CV" is the coefficient of variation over the seven prime powers.

N_{tot}	$n = 3$	4	5	2	7	8	9	mean	CV
600	0.119	0.134	0.112	0.117	0.112	0.141	0.120	0.122	0.085
800	0.112	0.125	0.104	0.117	0.107	0.120	0.110	0.113	0.061
1200	0.107	0.109	0.103	0.107	0.104	0.119	0.118	0.110	0.054
1600	0.100	0.108	0.099	0.106	0.097	0.112	0.107	0.104	0.051

At $L = 1.19$ the normalised jumps of all seven prime powers are thus close to a common value (CV about 5% at the two highest resolutions), although raw weights $\Lambda(n)n^{-1/2}$ range from 0.25 to 0.74; the common value drifts down with resolution (roughly like $1/(\ln(1/h) + b)$ in the mesh size h), so it is not a constant: the computed jumps are finite-resolution images of a logarithmically softened singularity of a continuous function, and they tend to 0. At $L = 1.5$

the value-jump ratios are again close to each other, but the split-window mesh there omits some cuts (including second-generation points close to the $n = 8, 16$ cuts), and we make no per-breakpoint claim. Prior work: the location of the singular points and the cut mesh are Eberwein's [9, §13]; jumps of a trial vector at $L - \log n$ appear in Mori's work [20], and the $\pm \log n$ shift structure in [23, eq. (2.11)].

9. Reproducibility

9.1. **Software and hardware.** Python 3.12 (3.12.13 for Pieces I–II) with python-flint 0.9.0, which bundles FLINT/Arb 3.6.0 [24, 15]; the C tools `vchol` (verified Cholesky) and `veig` (estimates only) of Piece II are built with gcc 15.2.0 against FLINT 3.4. Pieces I–II ran on a 20-thread laptop and a cluster of three 64-core nodes (FLINT versions on the cluster not recorded; a SHA-256 check of the code files on every host was in force for the certificate of record). Piece III and §7 ran on three 64-vCPU virtual machines. Elston's certifier is used at upstream commit `3cc4cb7be6a659f1463ae4e2e99ad7ae5e6646f5` with the `--deflate` option added (`code/gap/elston/`).

Run times. Piece II certificate of record: clean rerun for $L \le 1$ about 5.8 h in three stages; extension 1.03–1.09: 2859 s, 11022 s, 11760 s; continuation 1.12–1.19 about 28.5 h in total (head assembly about 6.5–7 h per window on the cluster). Piece III test at $L = 1.19$: 10–12 min per run with 18 workers. Piece III certificates: per run as in Table 3; the 16 runs behind Theorem 4.1 took about 814 worker-process hours (seconds times workers), all 19 shipped runs about 1177. Upper bounds U ($K = 640$, 3000 bits): 626–822 s per window. Rigorous upper bounds of Table 5: $L = 2$, $K = 1024$: 3166 s (even), 10971 s (odd); $L = 2.25$: 63923 s (even), 77675 s (odd); $L = 1.5$, 1.75 odd: 1278 s, 1293 s; $L = 1.5$ even ($K = 384$): 421 s.

9.2. **Commands.**

- Piece I and Piece II gluing: `python code/theorem.py --check` (seconds; ends with "THEOREM CERTIFIED for all L in (0, 1.19]"). Windows: `python code/theorem.py` with the environment settings recorded in the release.
- Piece III gluing: `python code/gap/check_steps.py` (seconds; reads `data/piece3/`; ends with "CERTIFIED: …simple-even for all L in (0, 1.4035]", compare `data/piece3/check.log`).
- Piece III lower bounds, in `code/gap/elston/`, with the window options `<w>` from the field `args` of the certificate file (e.g. `--a 1.2205 --delta 0.3 --beta-k 64 --tsharp 850 --nmodes 849 --gl 56 --prec 400`):

```
C="--panels-per-unit 4 --schur-k 4000 --schur-iters 1500"
python grid_certificate_rigorous.py <w> $C --sector odd --workers n --json <out>
python grid_certificate_rigorous.py <w> $C --sector even --deflate --workers n \
       --json <out>
```

- Upper bounds, in `code/upper/` (exports the outward-rounded field `lam0_upper`):

```
THREADS=n python fr_sweep.py <out.jsonl> "640" "1.19,1.2205,..." 3000
```

9.3. **Verification and audits.** All reviews so far were carried out by AI agents directed by the author; no independent human referee has checked the proofs or the code.

- *Pieces I–II.* Three independent review rounds of the earlier note produced, among others: the Gauss–Legendre exponent correction (Remark 5.9); reconciliation of the Piece I constants across precisions; outward rounding of all quoted bounds; a downward-induction proof of the Bessel ratio enclosure; the definition of the continuation of $\operatorname{Re}\psi$ off the real axis; documentation of Python `min` on balls; a mollification argument for (2.2); a self-contained discreteness proof; a SHA-256 check of remote code; and

TABLE 7. Provenance of the numbers in this paper. Paths under `highprec/` are in the author's working tree and are not part of the repository; all files the theorem depends on are.

Item	File(s)
Table 2, Piece I constants	`data/theorem_bounds.json`; `code/theorem.py`; logs `data/logs/check.log`, `run_rerun.log`, `run_extend.log` (release of [5])
Remark 5.9	`data/history/theorem_bounds_corrected.json`, `data/logs/run_gl_correction.log`
Piece III parameters (Table 3)	field `args` of each file in `data/piece3/`; design rule `code/gap/params.py`
Piece III certifier	`code/gap/elston/grid_certificate_rigorous.py`, `UPSTREAM_COMMIT`, `LICENSE`
$L = 1.19$ test (§6.2)	run logs on the compute host, not shipped (the test enters no step of the proof)
Table 4 lower bounds	`data/piece3/g_<L>_l2.json` (λ_2^{ev}), `g_<L>_odd.json` (λ_1^{odd}); field `certified_lower_bound`
Table 4 upper bounds	`data/piece3/upper640.jsonl`, field `lam0_upper`; code `code/upper/`
Piece III gluing	`code/gap/check_steps.py`, log `data/piece3/check.log`, checksums `data/piece3/SHA256SUMS`
Table 5, $L = 2$ even	`highprec/data/rig_L2.jsonl`, `rig_L2.log`
Table 5, other rows	`highprec/data/rig_L225.jsonl` ($L = 2.25$ even), `highprec/data/odd.jsonl` (odd sector), `highprec/gap/out_all/upper.jsonl` ($L = 1.5$ even)
Non-rigorous Galerkin values (§7)	`highprec/data/fr_L225*.jsonl`, `fr_sweep*.jsonl`
§8	`highprec/data/tail_probe.json`, `kink_119_*.json`, `kink_15_*.json`

corrected credit to Zhu for the reduction. Every finding was checked against code, logs and data before it was applied. A clean end-to-end rerun with a single code revision (empty cache, tools rebuilt, remote code hashed) then reproduced all lower bounds for $L \leq 1$ to about three significant digits. The sharp-cutoff path and the windows 1.03–1.19 have not been through a review round.

- *Upper-bound code.* An external review of the high-precision notes (six findings, all accepted) led to per-code-path rigour labels, outward rounding of exported upper endpoints, and withdrawal of an unsupported tail claim; an earlier review found and fixed an inaccurate ball inverse, a missing residual check and a missing quadrature bound.
- *Piece III.* Elston's certifier was audited independently at $L = 1.19$ (two issues fixed; documented in [11]); the post-audit code at larger L and our `--deflate` addition have not been audited. The step test (`code/gap/check_steps.py`, about 50 lines) compares explicit Arb endpoints. The reproduction of Elston's $L = 1.19$ bounds to 12 and 17 digits is a consistency check of the installation, not an audit.

Invitation to review. This work has not been peer-reviewed. Code, certificate files and this source are public at https://github.com/GenusCore/weil-ground-state-simple-even; the file `REVIEWING.md` there lists the least-checked parts and concrete checks ordered by effort, from re-running the gluing script on the shipped certificates (minutes) to re-running a single window (hours). Independent review by anyone is welcome; confirmed errors will be corrected publicly and reviewers credited.

10. Limitations

(1) *Range.* The theorem is certified on $(0, 1.4035]$. The extension to 1.5 is in progress (Remark 4.2); the passing step $[1.4255, 1.447]$ is a data point only.
(2) *Piece III is unaudited beyond $L = 1.19$.* Elston's certifier was audited only at $L = 1.19$; the runs at larger L, the parameter changes (β_K, $T^\sharp$, N, Gauss points) and our `--deflate` addition have not been reviewed. The step ratios of Piece III are small (1.131–1.552), so the theorem on $(1.19, 1.4035]$ depends on every bound being correct to within about 13%.
(3) *Computer assistance.* Correctness depends on Arb/FLINT, on our code for Pieces I–II and §7, on Elston's code for Piece III, and on the lemmas above.
(4) *Normalisation.* Our Q_L follows Zhu and was validated against the zero side of the explicit formula. It has not been checked line by line against QW_λ of [8, 7]. Remark 2.4 covers positive affine changes, not a discrepancy in the relative weights of the pole, archimedean and prime terms.
(5) *Self-adjoint realisation.* We use the closure of Q_L on $C_c^\infty(-L, L)$; trial polynomials not vanishing at $\pm L$ are in its form domain by the argument in §5.2. A different realisation would need re-examination.
(6) *Unexplained precision dependence* of the upper end of $\int |h|$ in Piece I (largest value used).
(7) *Comparisons on balls.* Python `min`/`max` on Arb balls occur in the gluing and in some intermediate steps of both certifiers; every current use is separated by many orders of magnitude more than the radii.
(8) *Sharp-cutoff and triangular assembly code* for $1.03 \le L \le 1.19$ in Piece II has not been reviewed; $L = 1.19$ is, however, re-confirmed by the Piece III code path.
(9) *Upper bounds of §7* are upper bounds only; their distance from the true eigenvalue is not controlled (Remark 7.2 for the $L = 2$ quadrature exponent).
(10) *Section 8* is numerical; the split-window x-quadrature is not bounded.
(11) *Scope.* Nothing is claimed for $L > 1.4035$, about the convergence step (ii) of [7], or about RH.

Acknowledgements

The Piece III lower bounds use Zac Elston's certifier *weil-windows* [11] (MIT licence), which we ran with one added option; the reduction it implements and the published $L = 1.19$ certificates are his. The window certificates of Pieces I–II rest on Xuefeng Zhu's frequency-cutoff reduction and block estimate [28]. The survey of lower-bound methods behind Piece III drew on the work of Mori [20] and of Liu [18], whose position-side use of the Legendre convolution recurrence preceded ours.

AI-assistance statement

Computations, code and drafting were carried out with substantial assistance from an AI system (Anthropic's Claude). All results were checked by independent recomputation, and the author takes responsibility for the content.

References

1. Andrade, *Certified simple-even ground states for finite Galerkin matrices qw^n*, 2026, Zenodo record 20887028. Description, not the original title. Unrefereed.
2. ———, *Perron-type theorem for the pole-removed Weil form*, 2026, Zenodo record 20682834. Description, not the original title. Unrefereed.

3. R. Andrews Jr., *Independent reproduction and convergence analysis of the CCM zeta spectral triple*, 2026, Zenodo record 22715669, v2.5. Unrefereed.
4. Enrico Bombieri, *Remarks on Weil's quadratic functional in the theory of prime numbers. I*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **11** (2000), no. 3, 183–233.
5. Johann Alexander Botha, *A certified simple, even ground state of the localized Weil quadratic form for all windows $l \in (0, 1.19]$*, 2026, Computer-assisted note, release of 2026-10-26, repository GenusCore/weil-ground-state-simple-even. Unrefereed.
6. Alain Connes, *The Riemann Hypothesis: past, present and a letter through time*, 2026, arXiv:2602.04022.
7. Alain Connes, Caterina Consani, and Henri Moscovici, *Zeta spectral triples*, 2025, arXiv:2511.22755.
8. Alain Connes and Walter D. van Suijlekom, *Quadratic forms, real zeros and echoes of the spectral action*, 2025, arXiv:2511.23257.
9. Logan Eberwein, *The truncated Weil quadratic form: inertia, visibility, structural measurements, and no-go theorems for the Riemann Hypothesis*, 2026, Zenodo, doi: 10.5281/zenodo.23177438 (6 October 2026); code `https://github.com/loganeberwein/Logan-Eberwein-RH-Weil-Window`. Unrefereed.
10. Elder, *Per-sector lower bounds for the localized Weil form*, 2026, Zenodo record 23113186. Description, not the original title. Unrefereed.
11. Zac Elston, *weil-windows*, 2026, Software and documentation, MIT licence; Zenodo, doi: 10.5281/zenodo.23174403; `https://github.com/zacharyelston/weil-windows`. Unrefereed.
12. Walter Gautschi, *Computational aspects of three-term recurrence relations*, SIAM Rev. **9** (1967).
13. grandchallenge / MATHSOLVE, *Analytic argument for windows $l \leq 9/50$*, 2026, Not read in full; reported only. Unrefereed.
14. Nicholas Hale and Alex Townsend, *An algorithm for the convolution of Legendre series*, SIAM J. Sci. Comput. **36** (2014), no. 3, A1207–A1220.
15. Fredrik Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), no. 8, 1281–1292.
16. jose-paul-b, *weil-positivity-certificates*, 2026, GitHub repository. Unrefereed.
17. Kramarenko-Byrd, *Simple even ground state of the localized Weil form for $0 < l \leq 4/5$*, 2026, Zenodo record 23007100, v1 of 2026-09-27, `https://zenodo.org/records/23007100`. Title paraphrased from the abstract. Unrefereed.
18. Liu, *certified-weil-positivity*, 2026, Manuscript and code, `https://github.com/luciferyu666/certified-weil-positivity`; submitted to Math. Comp. according to the repository. Unrefereed.
19. mathia project, *Project notes PL-282, PL-287, PL-294*, 2026, GitHub repository murillo128/mathia. Unrefereed.
20. Mori, *prolate-weil-bounds, paper VIII (positivity at $l = \frac{1}{2}\log 15$)*, 2026, Zenodo, doi: 10.5281/zenodo.23281236; `https://github.com/moriryota/prolate-weil-bounds`. Unrefereed.
21. Siegfried M. Rump, *Verification methods: rigorous results using floating-point arithmetic*, Acta Numer. **19** (2010), 287–449.
22. A. Senke, *Exact finite Gram closure for Weil positivity on the fixed window $(-77/64, 77/64)$*, 2026, Zenodo record 22871384, v0.3. Unrefereed.
23. Masatoshi Suzuki, *Weil's quadratic form via the screw function*, 2026, arXiv:2606.09096.
24. The FLINT team, *FLINT: Fast Library for Number Theory*, Version 3.6.0 (bundled with python-flint 0.9.0) and 3.4; `https://flintlib.org`.
25. Lloyd N. Trefethen, *Approximation theory and approximation practice*, SIAM, Philadelphia, 2013.
26. André Weil, *Sur les "formules explicites" de la théorie des nombres premiers*, Comm. Sém. Math. Univ. Lund, tome supplémentaire, 1952, pp. 252–265.
27. Hiroyuki Yoshida, *On Hermitian forms attached to zeta functions*, Zeta functions in geometry (Tokyo, 1990), Adv. Stud. Pure Math., vol. 21, Kinokuniya, Tokyo, 1992, pp. 281–325.
28. Xuefeng Zhu, *Weil positivity in compact windows: a finite reduction, certified two-sided bounds, and a Landau–Widom decay law*, 2026, arXiv:2608.24827v2.

GenusCore
Email address: `info@genuscore.com`