

CSCI 447: Project 1

George Engel

*Department of Engineering
Montana State University
Bozeman, MT 59715, USA*

GEOENGEL.Z@GMAIL.COM

Troy Oster

*Department of Engineering
Montana State University
Bozeman, MT 59715, USA*

TOSTER1011@GMAIL.COM

Dana Parker

*Department of Engineering
Montana State University
Bozeman, MT 59715, USA*

DANAHARMONPARKER@GMAIL.COM

Henry Soule

*Department of Engineering
Montana State University
Bozeman, MT 59715, USA*

HSOULE427@GMAIL.COM

Editor: Engel et al.

Abstract

A brief, one paragraph abstract summarizing the results of the experiments

Keywords: Keywords, Go, Here

1. Introduction

Problem statement, including hypothesis

2. Problem Statement

3. Algorithm

Per the instructions of the assignment, we implemented the naive Bayes algorithm on the five provided datasets. Given some example $x \in X$, where X is our dataset, naive Bayes predicts the correct class c of x by computing the probabilities of each possible classification for x . For class c , the probability is denoted as follows: $P(c|a_1, a_2, \dots, a_d)$ where a_k denotes one of d attribute values in x . To compute this probability for each class $c \in C$ where C is the set of all possible classifications, the probabilities of each attribute value are computed from the training set.

Acknowledgments

Acknowledgements go here.

Appendix A.

In this appendix we prove the following theorem from Section 6.2:

Theorem *Let u, v, w be discrete variables such that v, w do not co-occur with u (i.e., $u \neq 0 \Rightarrow v = w = 0$ in a given dataset \mathcal{D}). Let N_{v0}, N_{w0} be the number of data points for which $v = 0, w = 0$ respectively, and let I_{uv}, I_{uw} be the respective empirical mutual information values based on the sample \mathcal{D} . Then*

$$N_{v0} > N_{w0} \Rightarrow I_{uv} \leq I_{uw}$$

with equality only if u is identically 0. ■

Proof. We use the notation:

$$P_v(i) = \frac{N_v^i}{N}, \quad i \neq 0; \quad P_{v0} \equiv P_v(0) = 1 - \sum_{i \neq 0} P_v(i).$$

These values represent the (empirical) probabilities of v taking value $i \neq 0$ and 0 respectively. Entropies will be denoted by H . We aim to show that $\frac{\partial I_{uv}}{\partial P_{v0}} < 0 \dots$

Remainder omitted in this sample. See <http://www.jmlr.org/papers/> for full paper.