

What factors influence the occurrence and abundance of midstorey *Acacia* in Mountain Ash forests?

DAVID LINDENMAYER,^{*1,2}  WADE BLANCHARD,² LACHLAN MCBURNEY,^{1,2}
KITA ASHMAN,^{1,2} ELLE BOWD^{1,2}  AND DAVID BLAIR 

¹*Threatened Species Recovery Hub, National Environmental Science Program, Fenner School of Environment and Society, The Australian National University, Canberra, ACT, Australia; and* ²*Fenner School of Environment and Society, The Australian National University, Canberra, ACT 2601, Australia (Email: david.lindenmayer@anu.edu.au)*

Abstract The midstorey is a critical structural component of many forests globally. Using statistical models, we quantified the influence of two sets of variables on the percentage cover and basal area of two dominant *Acacia* spp. (Montane Wattle [*Acacia frigescens*] and Silver Wattle [*Acacia dealbata*]) in the midstorey of Mountain Ash (*Eucalyptus regnans*) forests in mainland south-eastern Australia. Specifically, we focused on the influence of (1) the age of the overstorey eucalypts (corresponding to the time since the last stand-replacing disturbance), and (2) environmental drivers (aspect, topographic wetness index, slope, elevation). We found evidence for generally non-linear relationships between stand age and the percentage cover and the basal area of both Silver Wattle and Montane Wattle. Silver Wattle had the highest values for percentage cover, and Montane Wattle the lowest, in stands regenerating from fire in 2009. The basal area of Silver Wattle was highest in stands that regenerated after the 2009 wildfires and after disturbance that occurred between 1960 and 1990s. For Montane Wattle, basal area was lowest in stands that regenerated in 2009 but values did not differ among stands of other ages. Both *Acacia* species were a midstorey component in old-growth Mountain Ash forest. No environmental covariates influenced the percentage cover of Montane Wattle or Silver Wattle. However, our model for the basal area of Montane Wattle contained evidence of a positive relationship with topographic wetness. The general paucity of environmental drivers in most of the models we constructed is likely due to the fact that both tree species occur well beyond our study region. Hence, the set of environmental conditions modelled may not be limiting the percentage cover or basal area of these midstorey tree species. Disturbance appears to be the key driver of dynamics of Montane Wattle and Silver Wattle in Mountain Ash forests.

Key words: ash-type eucalypt forest, disturbance, logging, south-eastern Australia, succession, wattle, wildfire.

INTRODUCTION

Many forests globally support multiple layers of vegetation, including a midstorey and an understorey of trees, shrubs and other plant species (e.g. tree ferns) (Franklin *et al.* 2002; Perry *et al.* 2008; Brock *et al.* 2020). These layers are a critically important component of stand structural complexity (*sensu* Lindenmayer & Franklin 2002) and have many key ecological roles. For example, the midstorey contributes to vertical heterogeneity (Brokaw & Lent 1999) which, in turn, creates more niches for more species in groups such as birds (MacArthur 1964; Recher 1969) and bats (Brown *et al.* 1997). Indeed, the midstorey is a key foraging substrate for many

kinds of animals (Morrison *et al.* 2006) ranging from mammals and birds (Smith 1984; Whelan & Maina 2005) to invertebrates (Woinarski & Cullen 1984; Schultz 2002). Other key roles of midstorey plants include providing sites for animals to nest (Beruldsen 2003), acting as nitrogen fixers for other plants such as overstorey trees (May & Attiwill 2003; Chaer *et al.* 2011), and providing a substrate for the establishment of other plants such as epiphytes and bryophytes (Pharo *et al.* 2013).

In forests subject to stand-replacing disturbances, overstorey trees may be killed, but midstorey and understorey plants may persist and recover through resprouting (Bradstock *et al.* 2012; Blair *et al.* 2016; White & Vesk 2019). This may mean that some elements of the midstorey and understorey may be of markedly different age to that of the dominant trees in the overstorey (Mueck *et al.* 1996; Fedrigo *et al.* 2019). In other cases, the overstorey trees may survive a disturbance (see McCarthy & Lindenmayer 1998) with some midstorey tree species killed but

*Corresponding author:

Accepted for publication December 2020.

†Deceased.

Permission to reproduce material from other sources: Not applicable.

then regenerating from seeds in the soil seed bank (Blair *et al.* 2016; Bowd *et al.* 2018). These include *Acacia* spp. trees which can occur as a key midstorey element in many eucalypt forests in Australia (Specht & Specht 1999; Boland *et al.* 2006), as well as other forests around the world (New 1984). *Acacia* spp. are an iconic component of Australia's flora with over 1000 species occurring on the continent (Burrows *et al.* 2018). They form long-lived, persistent soil seed banks that increase annually and which can be viable for multiple decades (Strydom *et al.* 2017; Burrows *et al.* 2018). The size of the *Acacia* spp. seed bank in the soil can increase with standing stem diameter, which when triggered by fire or other forms of soil heating (>60°C), can produce prolific germination of some species (Passos *et al.* 2017; Strydom *et al.* 2017). These reproductive strategies and traits allow *Acacia* spp. trees to persist in ecosystems for long periods (Passos *et al.* 2017; Strydom *et al.* 2017). However, *Acacia* spp. tree populations may be limited by pollinator availability (Cunningham 2000), recruitment issues, small seed set and senescing stands in small, fragmented populations (Broadhurst & Young 2006).

Given the importance of midstorey vegetation, including that comprising *Acacia* spp. trees, it is important to quantify its response to the following: (1) temporal drivers such as disturbances (e.g. stand-replacing fire and logging), and (2) spatial drivers like environmental conditions (e.g. slope, aspect, elevation and topography (Lindenmayer *et al.* 2000b; Huggett & Cheeseman 2002). In this study, we quantified the factors influencing the occurrence and abundance of *Acacia* spp. in the Mountain Ash (*Eucalyptus regnans*) forests of mainland south-eastern Australia.

Mountain Ash forests are wet-sclerophyll forests, dominated by the tall, obligate-seeder tree species, Mountain Ash. The forests support a diverse and well-developed midstorey consisting of *Acacia* spp. trees and other midstorey and understorey trees, broad leaved shrubs, tree ferns and a mesic ground layer rich in fern and herb species (Mueck 1990; Blair *et al.* 2016; Bowd *et al.* 2018). Prominent midstorey tree species include a range of *Acacia* spp. such as Silver Wattle (*Acacia dealbata*), Montane Wattle (*Acacia frutescens*) (sometimes also called Forest Wattle or Frosted Wattle), Mountain Hickory Wattle (*Acacia obliquinervia*) and Blackwood (*Acacia melanoxylon*) (Lindenmayer *et al.* 1994). Other midstorey tree species include the cool temperate rainforest trees, Myrtle Beech (*Nothofagus cunninghamii*) and Southern Sassafras (*Atherosperma moschatum*) that often occur in the cooler, wetter and more sheltered parts of the landscape (Lindenmayer *et al.* 2000b). Soils in these forests consist predominantly of well-drained, acidic, deep dermosols derived from

granitic rock (Bowd *et al.* 2019). In dominant stands, soil pH ranges from 3.4 to 5.2, nitrate and phosphorus content averages approximately 23 mg kg⁻¹ and 50 mg kg⁻¹, respectively, and organic carbon content typically exceeds 5% (Bowd *et al.* 2019).

We sought to identify relationships between overstorey stand age (which is a function of the time elapsed since the last stand-replacing disturbance [typically logging or fire]), environmental factors, and the percentage cover and basal area of *Acacia* spp. trees. We focused on *Acacia* spp. trees because they are key components of the midstorey in Mountain Ash forests (Ashton 1981; van der Meer & Dignan 2007; Serong & Lill 2008). *Acacia* spp. trees contribute to the known habitat requirements of the critically endangered Leadbeater's Possum (*Gymnobelideus leadbeateri*) (Lindenmayer *et al.* 1991b) and provide food and movement pathways not only for this species (Smith 1984; Lindenmayer *et al.* 1994) but also for other species of arboreal marsupials (Seebeck *et al.* 1984) and birds (Loyn 1985; Lindenmayer *et al.* 2009). *Acacia* spp. trees in Mountain Ash forests also contribute to carbon stocks (Keith *et al.* 2009; Fedrigo *et al.* 2019) and form symbiotic relationships with nitrogen-fixing microorganisms (May & Attiwill 2003).

Past studies (e.g. see Adams & Attiwill 1984; Blair *et al.* 2016; Trouvé *et al.* 2019) in Mountain Ash forests have examined the occurrence of some *Acacia* spp. trees in eucalypt forests. However, there has been limited quantification of the factors influencing both percentage cover and basal area of *Acacia* spp. in Mountain Ash forests. Here, we sought to close this knowledge gap by testing two broad hypotheses at the outset of this investigation.

Stand age hypothesis

We hypothesised that the percentage cover and basal area of midstorey *Acacia* spp. trees would exhibit a non-linear relationship with the age of overstorey trees. Mountain Ash forests are subject to stand-replacing disturbances in which the overstorey trees, and the majority of midstorey and understorey plants (including *Acacia* spp. trees), are killed (Ashton 1981; Simkin & Baker 2008). This triggers the germination of *Acacia* spp., which form a dominant component of early successional forest regrowth (Blair *et al.* 2016; Kasel *et al.* 2017; Bowd *et al.* 2018). Based on such prior knowledge of the biology of *Acacia* spp. trees, we therefore anticipated that values for percentage cover and basal area would be low soon after disturbance, then increase 20–80 years after disturbance, before declining again as *Acacia* spp. trees senesce and die (Adams & Attiwill 1984; Trouvé *et al.* 2019).

Environmental drivers hypothesis

We hypothesised that the percentage cover and basal area of midstorey *Acacia* spp. trees would be greatest in areas with the highest available moisture as reflected by broad environmental attributes such as topographic wetness, slope and aspect. That is, percentage cover and basal area values would be highest in areas with high values for topographic wetness, on flat terrain and on sheltered aspects. We hypothesised this response because of well-known relationships between topography and moisture availability (Huggett & Cheeseman 2002), coupled with moisture effects on the balance of overstorey and midstorey vegetation cover (Specht & Morgan 1983; Crombie 1992). That is, after controlling for factors like stand age (and associated structural attributes like tree height and stocking rate), we hypothesised that percentage cover and basal area of *Acacia* spp. trees would be highest where there is greater site productivity (Larson *et al.* 2008) including available moisture (Crombie 1992; Yamaura *et al.* 2020).

In addition to testing the two above hypotheses, given that the percentage cover and basal area of *Acacia* spp. trees were measured on the same sites, we sought to quantify statistical relationships between these two attributes of stand structure. Values for percentage cover may be similar for stands with quite different measures for basal area such as those with a high density of stems versus those with a low density of stems (Curtis *et al.* 2019). Nevertheless, at the outset of this study, we hypothesised there would be high levels of correlation between percentage cover and basal area. Thus, both measures also would exhibit similar responses to stand age and environmental drivers. We also explored patterns of co-occurrence among *Acacia* spp. trees and hypothesised that where values of percentage cover and basal area were high for a given species, they would be low for other species. Finally, we tested whether stand age or environment factors influenced the patterns of co-occurrence of *Acacia* tree species that were observed.

METHODS

Study area

We focused our study on the Mountain Ash forests of the Central Highlands of Victoria. The region lies about 120 km north-east of the city of Melbourne and covers approximately 1/2 degree of latitude and one degree of longitude (378200–378550S and 1458300–1460200E) (Fig. 1). The region experiences mild, humid winters with occasional periods of snow. Summers are generally cool. There is approximately 140 000 ha of Mountain Ash forest in our study area, and this tree species typically occurs at altitudes between 400 and 900 m (Costermans 1994; Boland *et al.* 2006).

We conducted surveys of midstorey *Acacia* spp. trees on 156 long-term monitoring sites in Mountain Ash forests. These sites measured 1 ha in size, and they covered a wide range of environmental conditions including stand age, slope, aspect, elevation and topographic position (Appendix S1).

Vegetation surveys

We visually estimated the projective (per cent) foliage cover (%) of *Acacia* spp. in six 10 × 10 m quadrats at 20 m increments along a 110 m transect in each site. Specifically, we made these estimates from the centre of each quadrat proportionate to the quadrat area (e.g. 1 m² of foliage cover = 1% of cover) in the summer of 2019–2020.

In 2017, at each site, we measured the diameter at breast height (DBH) of all live *Acacia frigescens*, *Acacia dealbata* and eucalypt stems greater than 2 m in height across three 10 m × 10 m plots, 10 m, 50 m and 90 m along a central 100 m transect. We counted the number of stems of each *Acacia* spp. and all eucalypts. We grouped the results into eleven DBH size categories: 0–5 cm, 5–10 cm, 10–20 cm, 20–30 cm, 30–40 cm, 40–60 cm, 60–80 cm, 80–100 cm, 100–140 cm, 140–180 cm and 180 cm+. For each stem, we computed the basal area at breast height using the DBH in metres via $BA = \pi(DBH/2)^2$ (the mid-point of each diameter interval was used in this calculation and 200 cm used for the highest category). We then summed the individual basal areas at breast height across all the stems of a particular species and then converted a measure per hectare by multiplying the total by 10 000 m²/300 m². We summed all eucalypt species at a site to get a total measure of eucalypt basal area. To account for the potential influence of overstorey eucalypts on *Acacia* spp. (Trouvé *et al.* 2019), we included the basal area and percentage cover of all eucalypts in all respective statistical models.

Environmental and other variables

We calculated values for a suite of covariates at each site for subsequent use in constructing statistical modelling. We assigned the age of the forest at each site to one of five age-classes: 1 = old growth dominated by trees that germinated before 1900 (7 sites), 2 = 1939 regrowth (dominated by trees that regenerated as a result of the 1939 wildfires) (75 sites), 3 = 1960–1990s regrowth (i.e. trees that regenerated between 1960 and 1990) (15 sites), 4 = sites were regenerated after the 2009 wildfire (31 sites) and 5 = mixed-aged forest (in which there were two or more distinct age cohorts of trees in the stand) (28 sites). Our age-class classification was based on the dominant age cohort of living overstorey trees in a stand (>85%). We also have made extensive measurements of the diameter of trees and using tree diameter–age relationship developed by Ashton (1976) to confirm stand age for any given site. We note that the vast majority of the mixed-aged stands supported an old-growth component with a number of individual large old living trees.

We interrogated a 20 m resolution Digital Elevation Model to extract data on slope, northerly aspect (aspect




Fig. 1. Location of the field survey sites where vegetation measurements of *Acacia* spp. were completed.

angles between 67.5 and 292.5 were assigned to a non-northerly aspect, and the remainder were assigned northerly aspect) and elevation for the centroid of each site. We also calculated values for a topographic wetness index (TWI) (Moore & Hutchinson 1991) for each site. TWI is a measure of relative position in the landscape and thus potential water distribution. Calculation of TWI requires a Digital Elevation Model (DEM) that has hydrological integrity; we used the ANUDEM algorithm (Hutchinson 2011) to generate a DEM of our study region at a grid resolution of 20 m. For each cell, the size of the catchment that flows to it was divided by its width, adjusted geometrically by the aspect of inflow direction. This 'specific catchment' was then divided by the cell's local slope. Lower values indicate ridges and upper slopes that have little to no contributing catchment, with values increasing for lower slopes, valley bottoms and drainage lines. We present the range of values for key covariates in Appendix S2.

Statistical analyses

Models of percentage cover and basal area

We modelled the percentage cover of each *Acacia* species using a Bayesian hurdle-beta regression model (Ospina & Ferrari 2012). We calculated the mean percent cover across

the six plots on each site and used the proportion data as the beta distribution is restricted to lie in the open interval (0,1); and the hurdle portion of the model was used to accommodate absences on a plot of a given species. In the literature, these models are typically called zero-inflated beta models. However, the name zero-inflated beta is a misnomer, as zeros and ones are not part of the support of the beta distribution. To avoid confusion, we do not use the standard nomenclature and refer to these models as hurdle-beta models. Specifically, the model we employed was the following:

$$PC_i \sim HB(\mu_i, \Phi)$$

$$\mu_i = \text{Intercept} + EPC_i + \text{StandAge}_i + \text{Slope}_i \\ + \text{NortherlyAspect}_i + \text{TWI}_i + \text{Elevation}_i$$

$$HU_i = \text{Intercept} + EPC_i + \text{StandAge}_i + \text{Slope}_i \\ + \text{NortherlyAspect}_i + \text{TWI}_i + \text{Elevation}_i$$

where PC_i is the average percentage cover on the i th site, HB refers to the hurdle-beta model, μ_i is the mean of the hurdle-beta distribution, Φ is the precision parameter of the beta distribution, EPC_i is the eucalypt percent cover, Stand Age (a categorical variable with levels: old growth, 1939, 1960–1990s, 2009, mixed age), slope, northerly aspect, TWI and elevation are covariates, and HU_i is the hurdle

component of the model. Both model components use the logistic link function. As the likelihood function for hurdle models factor in two independent components, we performed model selection for each component of the model independently. We fit each of the 32 possible combinations of the five covariates for each component of the zero-inflated beta regression model and chose the simplest model within two Widely Applicable Information Criterion (WAIC) units (see Gelman *et al.* 2014; Vehtari *et al.* 2016) of the best fitting model for each component. We note that eucalypt percent cover was included in all models.

We modelled site-level basal area data using a Bayesian hurdle-gamma regression model with an identical set of predictor variables as employed for the analysis of site-level average percentage cover described above. We employed a gamma distribution for basal area, as it is a positive variable, which has a gamma distribution. Specifically, the model was as follows:

$$BA_i \sim \text{HG}(\mu_i, \theta)$$

$$\mu_i = \text{Intercept} + \log(\text{EBA}_i) + \text{StandAge}_i + \text{Slope}_i \\ + \text{NortherlyAspect}_i + \text{TWI}_i + \text{Elevation}_i$$

$$\text{HU}_i = \text{Intercept} + \log(\text{EBA}_i) + \text{StandAge}_i + \text{Slope}_i \\ + \text{NortherlyAspect}_i + \text{TWI}_i + \text{Elevation}_i$$

where the covariates are the same as described previously except $\log(\text{EBA}_i)$ is the basal area of all eucalypts, HG is the hurdle gamma, and θ is the shape parameter of the gamma distribution. The logistic link function was used for the hurdle component of the model, while the gamma mean component of the model was modelled with a log link function. We employed the same model selection strategy as for the hurdle-beta regressions described above.

The models for percentage cover and basal area were fit with using the brms (Bayesian regression models using Stan) package (Buerkner 2017) in R version 3.6.1 (R Core Team 2019). We used the default priors in brms for all model parameters. Specifically, we used student-t priors with three degrees of freedom with zero mean and scale parameter ten for the regression parameters and a half student-t with three degrees of freedom with zero mean and scale parameter 10 for the standard deviation of the site random effect. We ran four Markov chains for 2000 iterations, discarding the first 1000 iterations as warm-up leaving 4000 posterior samples for inference. We assessed convergence using the Gelman–Rubin statistic, \hat{R} (Gelman & Rubin 1992). Note, in all cases, the \hat{R} was less than 1.01 indicating adequate mixing of the chains. We present posterior medians and 95% credible intervals for model parameters. We present R^2 values for each model for each species and also provide some basic residual diagnostics for each species.

Relationships between percentage cover and basal area

For each *Acacia* species, we used Spearman's correlation coefficients to investigate the associations between percentage cover and basal area.

Co-occurrence patterns between Silver Wattle and Montane Wattle

We defined co-occurrence as a binary variable, indicating whether or not pairs of species occurred together (1 = if species A and B occur together, 0 = if not) (see Appendix S2, Appendix S13). We modelled each of the binary variables on the previously mentioned covariates using a Bayesian logistic regression to determine whether or not co-occurrence was associated with the covariates. We used an identical model selection strategy as previously described.

RESULTS

We recorded six species of *Acacia* as midstorey components on our 156 survey sites in Mountain Ash forests. These species were as follows: Silver Wattle, Dwarf Silver Wattle (*Acacia nanodealbata*), Mountain Hickory Wattle, Montane Wattle, Blackwood and Prickly Moses (*Acacia verticillata*). Of these, only two species, Montane Wattle and Silver Wattle, were recorded at a sufficient number of sites (>30) to enable subsequent analyses and the construction of robust statistical models.

Silver Wattle occurred on 94 of our 156 sites surveyed in 2019 with percentage cover values ranging from 1 to 37% on the sites where it occurred. Using basal area, we gathered in 2017, Silver Wattle occurred on 83 sites, and when it did occur, values for basal area ranged from 0.02 to 29.52 m² ha⁻¹. Montane Wattle occurred on 64 of our 156 sites surveyed in 2019 with values for percentage cover ranging from 1.7 to 63.3% on sites where it occurred. Using basal area data gathered in 2017, Montane Wattle occurred on 57 sites and when it did occur, values for basal area ranged from 0.02 to 30.79 m² ha⁻¹. We developed separate models for percentage cover and basal area for both species.

Percentage cover of *Acacia*

The results of model selection for the hurdle and Beta component of the model for each species are given in Appendix S3. For the presence of Silver Wattle, we found a stand age effect but no eucalypt percentage cover or environmental covariate effects (Fig. 2a, Appendix S4, Appendix S14). After conditioning on the presence of Silver Wattle at a site, further analyses of percentage cover data revealed no environmental covariate effects but a stand age effect in which values were highest in stands that regenerated after the 2009 fire relative to other forest age-classes (Appendix S4, Appendix S14).

The presence of Montane Wattle was lowest in forest that germinated in 2009 and highest in forest that




Fig. 2. Panel a: Percentage cover of Montane Wattle and Silver Wattle in stands of Mountain Ash forest of different age in the Central Highlands of Victoria. Panel b: Basal area of Montane Wattle and Silver Wattle. The age categories were as follows: 1 = old growth (dominated by trees that regenerated before 1900), 2 = 1939 regrowth (dominated by trees that regenerated as a result of the 1939 wildfires), 3 = 1960–1990s regrowth (dominated by trees that regenerated between 1960 and 1990), 4 = 2009 regrowth (dominated by trees that regenerated from 2009 wildfire) and 5 = mixed-aged forest (in which there were two or more distinct age cohorts of trees in the stand). The percentage cover is a combination of the proportion of zeros (obtained from the hurdle part of the model) and the beta (conditional abundance) part of the model, and basal area is a combination of the proportion of zeros and gamma (conditional abundance) part of the model. The model estimates are plotted as 95% credible intervals (closed circles), and the individual data points are plotted as open circles. The number of zeros, for each species and measurement, is indicated at the bottom of each plot, and the overall number of plots in each group is also indicated in the labels for each age-class.

regenerated between 1960 and 1990 (Fig. 2a, Appendix S5, Appendix S15). No environmental factors, nor the percentage cover of eucalypts influenced the presence of Montane Wattle at a site (Appendix S5). After conditioning on the presence of Montane Wattle at a site, further analyses of percentage cover revealed a stand age effect but again no response to any environmental covariates (Appendix S5, Appendix S15).

Basal area of *Acacia*

The results of the model selection for the hurdle and Gamma component of the model for the basal area of each species of *Acacia* are given in Appendix S6. There was a stand age effect, but no effect of

eucalypt basal area or environmental covariate effects in the hurdle component of the model for the basal area of Silver Wattle (Appendix S7). After conditioning on the presence of Silver Wattle at a site, we found evidence of a stand age effect on basal area; values were highest in stands that regenerated after the 2009 wildfires and after disturbance that occurred between 1960 and the 1990s (Appendix S7, Appendix S16). There also was evidence of a eucalypt basal area effect in which values for the basal area of Silver Wattle were lower with increasing basal area of eucalypts (Appendix S7).

Our model for the basal area of Montane Wattle contained evidence of a stand age effect for the presence of the species (Fig. 2, Appendix S8, Appendix S17) with the lowest values being in forests burnt in 2009. There were no environmental covariate effects,

nor a eucalypt basal area effect for this (hurdle) component of the model (Appendix S8). After conditioning on the presence of Montane Wattle at a site, we found no stand age effects or basal area effects or environmental effects.

Relationships between percentage cover and basal area

We found a high level of correlation, as measured by Spearman's correlation coefficient, between percentage cover and basal area for both Silver Wattle (0.66) (Fig. 3a) and Montane Wattle (0.70) (Fig. 3b).

Co-occurrence patterns between Silver Wattle and Montane Wattle

Table 1 gives the breakdown of the presence/absence co-occurrence pattern of Silver Wattle and Montane Wattle for percentage cover and basal area, and Fig. 4 shows the results graphically. Model selection results for the Bayesian logistic regression for percentage cover (Appendices S9 and S10) revealed no

association with any of our covariates. However, there was evidence of a stand age effect for basal area co-occurrence of the two *Acacia* spp. (Appendices S9 and S11).

Diagnostics

We present residual diagnostic plots (residuals versus fitted values and response versus fitted values) for the percentage cover and basal area analysis in Appendix S18. We give Bayesian R^2 measures for percentage cover, basal area and co-occurrence in Appendix S12.

DISCUSSION

Midstorey vegetation is a critical component of many forests globally and plays a wide range of important ecological roles in ecosystem dynamics and biodiversity conservation (Perry *et al.* 2008). A key finding from our work was that time since disturbance (as reflected by the age of the overstorey eucalypt trees) was an important factor influencing the percentage




Fig. 3. Scatter plots of percentage cover versus basal area for a. Montane Wattle and b. Silver Wattle. The Spearman correlation coefficients are 0.70 and 0.66, respectively, for Montane Wattle and Silver Wattle.

Table 1. Cross-tabulation showing the number of sites for the co-occurrence of Silver Wattle and Montane Wattle as reflected by percentage cover and basal area

| Attribute | Silver Wattle | Montane Wattle | | Total |
|------------------|---------------|----------------|---------|-------|
| | | Absent | Present | |
| Percentage cover | Absent | 34 | 28 | 62 |
| | Present | 58 | 36 | 84 |
| | Total | 92 | 64 | 156 |
| Basal area | Absent | 53 | 24 | 77 |
| | Present | 49 | 30 | 79 |
| | Total | 102 | 54 | 156 |

cover and the basal area of both Silver Wattle and Montane Wattle, although there were marked interspecific differences in responses (Fig. 2). Previous studies in Mountain Ash forests have found a decline in the prevalence of *Acacia* spp. with an increase in the age of the eucalypt overstorey (e.g. Forrester *et al.* 2011; Trouvé *et al.* 2019). However, in this study and in the case of Montane Wattle, the lowest percentage cover values were for stands that regenerated after the 2009 fire (Fig. 2), but there were limited differences in percentage cover among other age-classes (except for mixed-aged forests relative to

stands that regenerated after the 1939 wildfire). Thus, there was a generally non-linear percentage cover–stand age relationship that was broadly consistent with what we hypothesised would occur at the outset of our study. The highest values for the percentage cover of Silver Wattle were for stands regenerating after the 2009 fire with markedly lower levels of cover in other age-classes (Fig. 2). The stand age relationships that we identified for basal area were broadly congruent with those recorded for percentage cover, again with marked interspecific differences in responses between Silver Wattle and Montane Wattle (Fig. 2). Such similar responses for percentage cover and basal area were consistent with the high levels of correlation between these two measures (as discussed further below).

An interesting observation from our datasets was that, similar to Montane Wattle, Silver Wattle was a midstorey component in old-growth Mountain Ash forest (Fig. 2). This is consistent with earlier studies that revealed that *Acacia* spp. trees characterise the midstorey of old-growth montane ash forests (Lindenmayer *et al.* 2000a). An allied investigation (see Lindenmayer *et al.* 1999) showed that very few old-growth stands supported strictly one age of overstorey trees, but rather there was strong empirical




Fig. 4. Scatter plots of Silver Wattle versus Montane Wattle for percentage cover (panel a) and basal area (panel b).

evidence of the presence of at least two (and sometimes more) age-classes (see also Fedrigo *et al.* 2019). It is possible that partial stand-replacing disturbances or site-specific environmental changes (increases in light from gaps in the understorey and canopy) may have triggered the germination and subsequent growth of *Acacia* spp. midstorey in stands of old-growth forest (Lindenmayer *et al.* 2000a; Fedrigo *et al.* 2019). Notably, detailed dendrochronology of a limited number of large old Mountain Ash trees provides evidence of multiple fires which have not killed these trees (Banks 1993). This is consistent with both the presence of fire scars on many living trees, and the occurrence of multi-aged stands in montane ash forests in the Central Highlands of Victoria (Lindenmayer *et al.* 1991a; McCarthy & Lindenmayer 1998). Furthermore, while measures like percentage cover and basal area of *Acacia* spp. may exhibit a general decline with stand age, seed stores increase (Passos *et al.* 2017; Strydom *et al.* 2017), which in the event of high-severity fire can result in a high abundance of *Acacia* spp. that exceeds that of younger forests (Bowd *et al.*, unpublished data, 2020).

As outlined above, our analyses contained evidence of interspecific differences in percentage cover and basal area relationships with stand age. We have not explored the underlying mechanisms for such interspecific differences in responses. However, they may be associated with interspecific competition or, alternatively, differences in life-history attributes such as the regeneration niche and/or subsequent growth patterns following disturbance. Further studies, specifically targeted at quantifying between-species competition, would be required to provide additional insights into the differences in stand age relationships for percentage cover and basal area that we have identified.

The influence of environmental factors on the midstorey

Unexpectedly, we found no evidence of effects of environmental factors such as slope, aspect and topographic wetness on the percentage cover of neither Montane Wattle nor Silver Wattle. This suggests their influence is limited relative to the impacts of disturbance. In contrast, we found a positive relationship between the basal area of Montane Wattle and topographic wetness. Thus, there was some support for our second postulate that *Acacia* spp. midstorey would be best developed in the wettest parts of Mountain Ash landscapes. The findings of this part of our study of *Acacia* spp. were broadly similar to those of previous work on other midstorey components of Mountain Ash forests (see also Kasel *et al.* 2017) such as Myrtle Beech. That study revealed that

strongly influenced by environmental attributes (e.g. slope Lindenmayer *et al.* 2000b).

The reasons for general paucity of environmental effects in most of the models we developed for percentage cover and basal remain unclear. Both Montane Wattle and Silver Wattle species have somewhat broader distributions (that encompass a wider range of environments) than the Central Highlands region where we completed this study. Hence, the relatively restricted set of environmental conditions that we modelled are a subset of the overall environmental domains occupied by these species, and may therefore not be limiting for either species.

Relationships between percentage cover and basal area and co-occurrence patterns

We found broadly similar responses to stand age and environmental drivers for percentage cover and basal area for both species. That is, the shapes of the response curves for stand age for both measures were broadly congruent (compare Figs 2 and 3). This result was perhaps not surprising given the high level of correlation between these measures, particularly for Montane Wattle.

Given the broad similarity of responses of percentage cover and basal area to stand age, the findings reported here will be generally relatable to past work such as that on animal responses to the basal area of *Acacia* (Lindenmayer *et al.* 1991b). However, because there were differences between percentage cover and basal area in responses to environmental factors, there are benefits in measuring both attributes of stand structure.

Co-occurrence patterns

While we have not explicitly sought to complete a formal statistical comparison of the post-disturbance trajectories of Montane Wattle and Silver Wattle, it is important to note that sites with high values of percentage cover for one species often were characterised by an absence of the other species (Fig. 4). This result may, in part, be an outcome of examining compositional data where the total amount of cover has a maximum (capped) value (100%). However, our findings also may reflect interspecific competition or physiological differences which can influence germination and dispersal success (Brown *et al.* 2008; Forrester *et al.* 2011). Notably, no factors influenced the co-occurrence of the two species based on percentage cover data. By contrast, our analysis of basal area data indicated there was a stand age effect, with Silver Wattle and Montane Wattle less likely to co-occur in forests regenerating after the 2009 wildfires.

The reasons for the limited co-occurrence of both species (as reflected by basal area values) in young post-fire forests remain unclear, but this effect may be related to the general rarity of Montane Wattle in forests of this age.

Caveats

We modelled the factors influencing the percentage cover and basal area of two key species of *Acacia* in the understorey of Mountain Ash forests. We recognise the factors in addition to the ones we analysed may have had an important effect. These include soil type. Further studies in which detailed soil data are gathered would be a useful adjunct to the work reported here. In addition, our study did not include early post-regeneration stands, such as those that were between one and nine years old. We recognise that there would be value in completing additional field surveys to capture information on such early successional forests. Finally, we did not explore the effects of interactions between variables such as stand age and environmental attributes such as topographic wetness. This was because stand age was the sole main effect for the vast majority of models that we constructed. Moreover, a larger dataset than we had available to us would be required to build robust models comprising interactions among a suite of variables.

CONCLUDING COMMENTS

We provide insights into the temporal and spatial dynamics of two dominant *Acacia* tree spp. in the Mountain Ash forests of mainland Australia highlighting the critical role of disturbance, and subsequent stand age on their levels of cover and basal area. As *Acacia* spp. trees have important functional roles in forests (e.g. as a foraging substrate, for habitat connectivity, and nitrogen fixation), our findings provide important insights for forest management, including understanding where and when suitable amounts of this kind of vegetation will occur. Indeed, our findings suggest that the dynamics of a given dominant midstorey tree species cannot be well understood in the absence of insights into the dynamics of other relatively common midstorey species in the ecosystem.

ACKNOWLEDGEMENTS

This study was supported by grants from the Australian Government (National Environmental Science Program Threatened Species Recovery Hub), the Victorian Government (Department of Environment,

Land, Water and Planning; and Parks Victoria) and private donors. The authors have no conflict of interest to declare. This work is dedicated to the memory of friend and colleague, Dr David Blair. He was working on key aspects of midstorey dynamics at the time of his very premature death. We thank Dr Chris Taylor for providing updated information on the environmental attributes of our field sites. Claire Shepherd assisted with manuscript preparation.

AUTHOR CONTRIBUTION

David B Lindenmayer: Conceptualization (equal); Funding acquisition (lead); Investigation (equal); Methodology (equal); Resources (lead); Supervision (lead); Writing-original draft (lead); Writing-review & editing (lead). **Wade Blanchard:** Formal analysis (lead); Investigation (equal); Methodology (equal); Writing-review & editing (lead). **Lachlan McBurney:** Data curation (equal); Investigation (equal); Writing-review & editing (equal). **Kita Ashman:** Writing-review & editing (equal). **Elle J Bowd:** Investigation (equal); Writing-review & editing (equal). **David Blair:** Conceptualization (lead); Data curation (equal); Methodology (equal).

Data Availability Statement

The data are available in the Dryad Data Repository. Citation: Lindenmayer, David et al. (2020), What factors influence the extent of midstorey development in Mountain Ash forests? Dryad, Dataset, <https://doi.org/10.5061/dryad.zw3r2285m>

REFERENCES

- Adams M. H. & Attiwill P. M. (1984) The role of *Acacia* spp. in nutrient balance and cycling in regenerating *Eucalyptus regnans* F. Muell forests 1. Temporal changes in biomass and nutrient content. *Aust. J. Bot.* **32**, 205–15.
- Ashton D. H. (1976) The development of even-aged stands of *Eucalyptus regnans* F. Muell. in central Victoria. *Aust. J. Bot.* **24**, 397–414.
- Ashton D. H. (1981) Tall Open Forests. In: *Australian Vegetation*. (ed R. H. Groves) pp. 121–51. Cambridge University Press, Cambridge.
- Banks J. C. (1993) *Tree-ring analysis of two Mountain Ash trees Eucalyptus regnans* F. Muell from the Watts and O'Shannassy Catchments, Central Highlands, Victoria. A Report to the Central Highlands Old Growth Forest Project. August 1993. Department of Conservation and Natural Resources, Melbourne, Australia.
- Beruldsen G. R. (2003) *Australian Birds: Their Nests and Eggs*. G & E Beruldsen, Kenmore Hills.
- Blair D., McBurney L., Blanchard W., Banks S. & Lindenmayer D. B. (2016) Disturbance gradient shows

- logging affects plant functional groups more than fire. *Ecol. Appl.* **26**, 2280–301.
- Boland D. J., Brooker M. I., Chippendale G. M. *et al.* (2006) *Forest Trees of Australia*. CSIRO Publishing, Melbourne.
- Bowd E. J., Banks S. C., Strong C. L. & Lindenmayer D. B. (2019) Long-term impacts of wildfire and logging on forest soils. *Nat. Geosci.* **12**, 113–8.
- Bowd E. J., Lindenmayer D. B., Banks S. C. & Blair D. P. (2018) Logging and fire regimes alter plant communities. *Ecol. Appl.* **28**, 826–41.
- Bradstock R. A., Gill A. M. & Williams R. J. (2012) *Flammable Australia: Fire Regimes, Biodiversity and Ecosystems in a Changing World*. CSIRO Publishing, Melbourne.
- Broadhurst L. M. & Young A. G. (2006) Reproductive constraints for the long-term persistence of fragmented *Acacia dealbata* (Mimosaceae) populations in southeast Australia. *Biol. Conserv.* **133**, 512–36.
- Brock J. M., Morales N. S., Burns B. R. & Perry G. L. W. (2020) The hare, tortoise and crocodile revisited: Tree fern facilitation of conifer persistence and angiosperm growth in simulated forests. *J. Ecol.* **108**, 969–81.
- Brokaw N. V. & Lent R. A. (1999) Vertical structure. In: *Managing Biodiversity in Forest Ecosystems*. (ed M. Hunter) pp. 373–99. Cambridge University Press, Cambridge.
- Brown G. W., Nelson J. L. & Cherry K. A. (1997) The influence of habitat structure on insectivorous bat activity in montane ash forests of the Central Highlands of Victoria. *Aust. Forestry* **60**, 138–46.
- Brown J., Enright N. J. & Miller B. P. (2008) Seed production and germination in two rare and three common co-occurring *Acacia* species from south-east Australia. *Austral Ecol.* **28**, 271–80.
- Buerkner P.-C. (2017) brms: An R Package for Bayesian Multilevel Models using Stan. *J. Stat. Softw.* **80**, 1–28.
- Burrows G. E., Alden R. & Robinson W. A. (2018) The lens in focus – lens structure in seeds of 51 Australian *Acacia* species and its implications for imbibition and germination. *Aust. J. Bot.* **66**, 398–413.
- Chaer G. M., Resende A. S., Campello E. F. C., de Faria S. M. & Boddey R. M. (2011) Nitrogen-fixing legume tree species for the reclamation of severely degraded lands in Brazil. *Tree Physiol.* **31**, 139–49.
- Costermans L. (1994) *Native trees and shrubs of south-eastern Australia*, 2nd edn. Rigby, Sydney.
- Crombie D. S. (1992) Root depth, leaf area and daytime water relations of Jarrah (*Eucalyptus marginata*) forest overstorey and understorey during summer drought. *Aust. J. Bot.* **40**, 113–22.
- Cunningham S. A. (2000) Depressed pollination in habitat fragments causes low fruit set. *Proc. R. Soc. B* **267**, 1149–52.
- Curtis E. J., Gorrod E. J., Ellis M. V. & Chisholm L. A. (2019) A spatio-temporal analysis of canopy dynamics and intra-stand competition in a riparian forest, south-eastern Australia. *Forest Ecol. Manage.* **432**, 189–99.
- Fedrico M., Stewart S., Kasel S., Levchenko V., Trouvé R. & Nitschke C. (2019) Radiocarbon dating informs tree fern population dynamics and disturbance history of temperate forests in southeast Australia. *Radiocarbon* **61**, 445–60.
- Forrester D. I., Vanclay J. K. & Forrester R. I. (2011) The balance between facilitation and competition in mixtures of *Eucalyptus* and *Acacia* changes as stands develop. *Oecologia* **166**, 265–72.
- Franklin J. F., Spies T. A., van Pelt R. *et al.* (2002) Disturbances and the structural development of natural forest ecosystems with silvicultural implications, using Douglas-fir forests as an example. *Forest Ecol. Manage.* **155**, 399–423.
- Gelman A., Hwang J. & Vehtari A. (2014) Understanding predictive information criteria for Bayesian models. *Stat. Comput.* **24**, 997–1016.
- Gelman A. & Rubin D. B. (1992) Inference from iterative simulation using multiple sequences. *Stat. Sci.* **7**, 457–511.
- Huggett R. & Cheeseman J. (2002) *Topography and the Environment*. Prentice Hall, London.
- Hutchinson M. F. (2011) *ANUDEM Version 5.3*. The Australian National University, Canberra. Available at: <https://fennergchool.anu.edu.au/research/products/anudem>.
- Kasel S., Bennett L., Fedrico M. & Nitschke C. (2017) Environmental heterogeneity promotes floristic turnover in temperate forests of south-eastern Australia more than dispersal limitation and disturbance. *Landscape Ecol.* **32**, 1613–29.
- Keith H., Mackey B. G. & Lindenmayer D. B. (2009) Re-evaluation of forest biomass carbon stocks and lessons from the world's most carbon-dense forests. *Proc. Natl. Acad. Sci. USA* **106**, 11635–40.
- Larson A. J., Lutz J. A., Gersonde R. F., Franklin J. F. & Hietpas F. F. (2008) Potential site productivity influences the rate of forest structural development. *Ecol. Appl.* **18**, 899–910.
- Lindenmayer D. B., Boyle S., Burgman M. A., McDonald D. & Tomkins B. (1994) The sugar and nitrogen content of the gums of *Acacia* species in the Mountain Ash and Alpine Ash forests of central Victoria and its potential implications for exudivorous arboreal marsupials. *Austral. J. Ecol.* **19**, 169–77.
- Lindenmayer D. B., Cunningham R. B., Donnelly C. F. & Franklin J. F. (2000a) Structural features of old growth Australian montane ash forests. *For. Ecol. Manage.* **134**, 189–204.
- Lindenmayer D. B., Cunningham R. B., Nix H. A., Tanton M. T. & Smith A. P. (1991a) Predicting the abundance of hollow-bearing trees in montane ash forests of southeastern Australia. *Aust. J. Ecol.* **16**, 91–8.
- Lindenmayer D. B., Cunningham R. B., Tanton M. T., Nix H. A. & Smith A. P. (1991b) The conservation of arboreal marsupials in the montane ash forests of the Central Highlands of Victoria, south-eastern Australia. III. The habitat requirements of Leadbeaters Possum *Gymnobelideus leadbeateri* and models of the diversity and abundance of arboreal marsupials. *Biol. Conserv.* **56**, 295–315.
- Lindenmayer D. B. & Franklin J. F. (2002) *Conserving Forest Biodiversity: A Comprehensive Multiscaled Approach*. Island Press, Washington DC.
- Lindenmayer D. B., Mackey B. G., Cunningham R. B. *et al.* (2000b) Factors affecting the presence of the cool temperate rain forest tree myrtle beech (*Nothofagus cunninghamii*) in southern Australia: integrating climatic, terrain and disturbance predictors of distribution patterns. *J. Biogeogr.* **27**, 1001–9.
- Lindenmayer D. B., Mackey B. G., Mullen I. C. *et al.* (1999) Factors affecting stand structure in forests – are there climatic and topographic determinants? *Forest. Ecol. Manage.* **123**, 55–63.
- Lindenmayer D. B., Wood J., Michael D. *et al.* (2009) Are gullies best for biodiversity? An empirical examination of Australian wet forest types. *Forest Ecol. Manage.* **258**, 169–77.
- Loyn R. H. (1985) Bird populations in successional forests of Mountain Ash *Eucalyptus regnans* in Central Victoria. *Emu* **85**, 213–30.

- MacArthur R. H. (1964) Environmental factors affecting bird species diversity. *Am. Nat.* **98**, 387–97.
- May B. M. & Attiwill P. M. (2003) Nitrogen-fixation by *Acacia dealbata* and changes in soil properties 5 years after mechanical disturbance or slash-burning following timber harvest. *Forest Ecol. Manage.* **181**, 339–55.
- McCarthy M. A. & Lindenmayer D. B. (1998) Multi-aged mountain ash forest, wildlife conservation and timber harvesting. *Forest Ecol. Manage.* **104**, 43–56.
- Moore I. D. & Hutchinson M. F. (1991) Spatial extension of hydrologic process modelling. In: *International Hydrology and Water Resources Symposium 1991: Challenges for sustainable development*. Perth, WA. Institution of Engineers, Australia. pp. 419–27.
- Morrison M. L., Marcot B. G. & Mannan R. W. (2006) *Wildlife-Habitat Relationships. Concepts and Applications*. Island Press, Washington, D.C.
- Mueck S. G. (1990) *The Floristic Composition of Mountain Ash and Alpine Ash Forests in Victoria: Silvicultural Systems Project No.4*. Department of Conservation and Environment, Victoria.
- Mueck S. G., Ough K. & Banks J. C. (1996) How old are wet forest understories? *Aust. J. Ecol.* **21**, 345–8.
- New T. R. (1984) *A Biology of the Acacias*. Oxford University Press, Oxford.
- Ospina R. & Ferrari S. L. (2012) A general class of zero-or-one-inflated beta regression models. *Comput. Stat. Data An.* **56**, 1609–23.
- Passos I., Marchante H., Pinho R. & Marchante E. (2017) What we don't seed: The role of long-lived seed banks as hidden legacies of invasive plants. *Plant Ecol.* **218**, 1313–24.
- Perry D. A., Oren R. & Hart S. C. (2008) *Forest Ecosystems*. John Hopkins University Press, Baltimore.
- Pharo E. J., Meagher D. A. & Lindenmayer D. B. (2013) Bryophyte persistence following major fire in eucalypt forest of southern Australia. *Forest Ecol. Manage.* **296**, 24–32.
- R Core Team (2019) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Recher H. F. (1969) Bird species diversity and habitat diversity in Australia and North America. *Am. Nat.* **103**, 75–121.
- Schultz C. B. (2002) Restoring resources for an endangered butterfly. *J. Appl. Ecol.* **38**, 1007–18.
- Seebeck J. H., Warneke R. M. & Baxter B. J. (1984) Diet of the Bobuck, *Trichosurus caninus* Ogilby (Marsupialia: Phalangeridae) in a mountain forest in Victoria. In: *Possums and Gliders*. (eds A. P. Smith & I. D. Hume) pp. 145–54. Surrey Beatty and Sons, Chipping Norton, Sydney.
- Serong M. & Lill A. (2008) The timing and nature of floristic and structural changes during secondary succession in wet forests. *Aust. J. Bot.* **56**, 220–31.
- Simkin R. & Baker P. J. (2008) Disturbance history and stand dynamics in tall open forest and riparian rainforest in the Central Highlands of Victoria. *Austral Ecol.* **33**, 747–60.
- Smith A. P. (1984) Diet of Leadbeater's Possum *Gymnobelideus leadbeateri* (Marsupialia). *Aust. Wildl. Res.* **11**, 265–73.
- Specht R. L. & Morgan D. G. (1983) The balance between the foliage projective covers of overstorey and understorey in Australian vegetation. *Aust. J. Ecol.* **6**, 193–202.
- Specht R. L. & Specht A. (1999) *Australian Plant Communities. Dynamics of Structure, Growth and Biodiversity*. Oxford University Press, Melbourne.
- Strydom M., Veldtman R., Ngwenya M. Z. & Esler K. J. (2017) Invasive Australian *Acacia* seed banks: Size and relationship with stem diameter in the presence of gall-forming biological control agents. *PLoS One* **12**, e0181763.
- Trouvé R., Nitschke C. R., Andrieux L., Willersdorf T., Robinson A. P. & Baker P. J. (2019) Competition drives the decline of a dominant midstorey tree species. Habitat implications for an endangered marsupial. *Forest Ecol. Manage.* **447**, 26–34.
- van der Meer P. J. & Dignan P. (2007) Regeneration after 8 years in artificial canopy gaps in Mountain Ash (*Eucalyptus regnans* F. Muell.) forest in south-eastern Australia. *Forest Ecol. Manage.* **244**, 102–11.
- Vehtari A., Gelman A. & Gabry J. (2016) Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Stat. Comput.* arXiv preprint arXiv:1507.04544.
- Whelan C. J. & Maina G. G. (2005) Effects of season, understorey vegetation density, habitat edge and tree diameter on patch-use by bark-foraging birds. *Funct. Ecol.* **19**, 529–36.
- White D. J. & Vesk P. A. (2019) Fire and legacy effects of logging on understorey assemblages in wet-sclerophyll forests. *Aust. J. Bot.* **67**, 341–57.
- Woinarski J. C. & Cullen J. M. (1984) Distribution of invertebrates on foliage in forests in south-eastern Australia. *Aust. J. Ecol.* **9**, 207–32.
- Yamaura Y., Lindenmayer D. B., Yamada Y. *et al.* (2020) A spatially-explicit empirical model of structural development processes in natural forests based on climate and topography. *Conserv. Biol.* **34**, 194–206.

SUPPORTING INFORMATION

Additional supporting information may/can be found online in the supporting information tab for this article.

Appendix S1. Descriptive statistics for site-level covariates including the mean, median, minimum and maximum values for key covariates.

Appendix S2. Descriptive statistics for the occurrence of the three most commonly occurring *Acacia* tree species as measured by percentage cover and basal area.

Appendix S3. Model Selection results for the Bayesian hurdle beta regression models for Silver Wattle and Montane Wattle.

Appendix S4. Silver Wattle hurdle-beta posterior model summaries for percentage cover (converted to proportions for modelling).

Appendix S5. Montane Wattle hurdle-beta posterior model summaries for percentage cover (converted to proportions for modelling).

Appendix S6. Model Selection results for the Bayesian hurdle gamma regression models for Silver Wattle and Montane Wattle.

Appendix S7. Silver Wattle hurdle-gamma posterior model summaries for basal area.

Appendix S8. Montane Wattle hurdle-gamma posterior model summaries for basal area.

Appendix S9. Model Selection results for the Bayesian logistic regression models for the co-occurrence of Silver Wattle and Montane Wattle.

Appendix S10. Best fitting Bayesian logistic regression model for the co-occurrence of Silver Wattle and Montane Wattle when assessed via percentage cover.

Appendix S11. Best fitting Bayesian logistic regression model for the co-occurrence of Silver Wattle and Montane Wattle as measured by basal area.

Appendix S12. Bayesian R^2 (in percent) for each of the best fitting models for Silver Wattle and Montane Wattle for percentage cover and basal area.

Appendix S13. Scatter plots of percentage cover and basal area of Montane and Silver Wattle and all *Eucalyptus* species by stand age.

Appendix S14. Estimates and comparisons with 95% credible intervals of the effects of stand age on

the probability of absence, the conditional percentage cover and the unconditional percentage cover of Silver Wattle.

Appendix S15. Estimates and comparisons with 95% credible intervals of the effects of stand age on the probability of absence, the conditional percentage cover and the unconditional percentage cover of Montane Wattle.

Appendix S16. Estimates and comparisons with 95% credible intervals of the effects of stand age on the probability of absence, the conditional basal area and the unconditional basal area of Silver Wattle.

Appendix S17. Estimates and comparisons with 95% credible intervals of the effects of stand age on the probability of absence, the conditional basal area and the unconditional basal area of Montane Wattle.

Appendix S18. Diagnostic plots for Silver Wattle and Montane Wattle for percentage cover and basal area.