

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/323300859>

Populations of Commercial Molluscs within a Highly Biodiverse Marine Protected Area of the Northern Alboran Sea (W Mediterranean): Preferential Habitats, Seasonal Dynamics and Impo...

Article in *Thalassas: An International Journal of Marine Sciences* · October 2018

DOI: 10.1007/s41208-018-0070-5

CITATIONS

6

READS

364

5 authors, including:



Javier Urra

Instituto Español de Oceanografía

123 PUBLICATIONS 1,102 CITATIONS

[SEE PROFILE](#)



Jose L. Rueda

Instituto Español de Oceanografía

208 PUBLICATIONS 2,191 CITATIONS

[SEE PROFILE](#)



Pablo Marina

Instituto Español de Oceanografía (IEO-CSIC)

67 PUBLICATIONS 515 CITATIONS

[SEE PROFILE](#)



Antit BEN REJEB Mouna

University of Tunis El Manar

24 PUBLICATIONS 191 CITATIONS

[SEE PROFILE](#)



Populations of Commercial Molluscs within a Highly Biodiverse Marine Protected Area of the Northern Alboran Sea (W Mediterranean): Preferential Habitats, Seasonal Dynamics and Importance for Artisanal Fisheries

Javier Urrea^{1,2} · José Luis Rueda¹ · Pablo Marina² · Mouna Antit³ · Carmen Salas²

Received: 27 September 2017 /

© Springer International Publishing AG, part of Springer Nature 2018

Abstract

Seasonal dynamics and preferential sedimentary habitats of commercial molluscs have been studied in soft bottoms within the limits of the Special Area of Conservation (SAC) “Calahonda” (S Spain). This SAC harbours a high diversity of invertebrate communities, and it is located in a very touristy area of the littoral of Malaga and within the limits of one of the most productive shellfish fishing areas of the NW Alboran Sea. Here, the artisanal fishery targeting molluscs is a locally important activity because some species (e.g. *Chamelea gallina*, *Callista chione*, *Donax trunculus*) are highly demanded by locals and tourists and therefore reaching high commercial value. More than 200 molluscan species inhabit the soft bottoms of the SAC “Calahonda”, from which eight species collected in this study have economic value. The bivalve *C. gallina* was the most abundant and frequent commercial species, mainly at shallow fine and medium sand bottoms, followed by *C. chione* at coarse sand bottoms. Some species displayed higher abundance values of large or small size individuals at different depths, and most of them showed maximum abundances in spring and summer. Many invertebrate species associated with these soft bottoms are potentially very sensitive to dredging activities, especially echinoderms, decapods and cnidarians. Further research regarding the potential impact of fisheries on these assemblages would be desirable, especially considering the high biodiversity existing within this SAC and the increase of tourists in the last years.

Keywords Alboran Sea · Biodiversity · Commercial molluscs · Conservation

Introduction

Molluscs are within the top-three faunistic groups collected in marine fishing areas worldwide and represent ca. 10.4% (above 9 million tons) of the total world fishery landings (FAO 2012). In this line, molluscs are one of the most exploited marine resources in the Mediterranean Sea

(Fischer et al. 1987), being most of the bivalve species the same ones than those recorded in the classical works of Greek antiquity by Aristotle, Xenocrates and Athenaeus, among others (Poutiers 1987; Voultsiadou et al. 2010). At the same time, molluscs are within the top dominant faunistic groups in several marine habitats, acting as an important component in trophic webs as they represent one of the main primary consumers of plant productivity and nutrient recyclers from the water column (Snelgrove 1998; Mazzella et al. 1992). Besides, molluscs are of interest for studies describing biodiversity patterns in marine ecosystems and changes in community composition and structure (Chemello et al. 2000; Rueda et al. 2009; Urrea et al. 2017a).

The Alboran Sea highlights within the Mediterranean context as it harbours a high biodiversity promoted by its biogeographic location, oceanography and bottom heterogeneity (García Raso et al. 2010; Robles 2010; Templado 2011). This has promoted an important fishing activity with a high

✉ Javier Urrea
javier.urra.recuero@gmail.com

¹ Centro Oceanográfico de Málaga, Instituto Español de Oceanografía, Puerto Pesquero s/n, 29640 Fuengirola, Málaga, Spain

² Departamento de Biología Animal, Universidad de Málaga, Campus de Teatinos s/n, 29071 Málaga, Spain

³ Faculté des Sciences de Tunis, UR11ES12 Biologie de la Reproduction et du Développement Animal, Université de Tunis El Manar, 2092 Tunis, Tunisie

diversity of fishing gears, techniques and target species (Camiñas et al. 2004; Robles 2010). An important part of the fishing fleet of the northern Alboran Sea is involved in artisanal fisheries, characterized by operating with small vessels in shallow fishing grounds close to the coast and in one day fishing trips using small vessels (Camiñas et al. 2004). Artisanal fisheries target many different species alternating gears and techniques, both spatially and temporally depending on market prices, close seasons and seasonality of resources, in order to optimize catches and maximize benefits (Camiñas et al. 2004). Molluscs represent a very important resource for these fisheries in the Alboran Sea, being targeted by the mechanical dredging fleet that involves 247 vessels in the northern basin (MAGRAMA 2012). Among the target species, the smooth clam *Callista chione* (Linnaeus, 1758), the striped venus clam *Chamelea gallina* (Linnaeus, 1758) and the wedge clam *Donax trunculus* Linnaeus, 1758 represent some of the most important commercial species collected by the this fleet (Tirado 2011), with ca. 192,000 $\text{k}\cdot\text{y}^{-1}$ collected for *C. chione* (mean annual value: 412,609 €) and more than 20,000 $\text{k}\cdot\text{y}^{-1}$ collected for *C. gallina* (mean annual value: 165,254 €) and *D. trunculus* (mean annual value: 221,982 €), respectively, in Mediterranean fishing grounds of Andalusia (<http://www.juntadeandalucia.es>). These species present important populations between Estepona and Vélez-Málaga coastal areas (Málaga province), together with those of other molluscan commercial species, clustering the activity of a great part of the fleet in this area.

One of the main marine molluscs fishing areas of the northern Alboran Sea is “Cala del Moral” (code AND31), where the capture of the smooth clam, the striped venus clam, the wedge clam, the banded carpet-shell clam *Tapes rhomboides* (Pennant, 1777), the warty venus clam *Venus verrucosa* Linnaeus, 1758, the tuberculate cockle *Acanthocardia tuberculata* (Linnaeus, 1758), the king scallop *Pecten maximus* (Linnaeus, 1758), the banded murex *Hexaplex trunculus* (Linnaeus, 1758) and the purple dye murex *Bolinus brandaris* (Linnaeus, 1758) among other species, is allowed by artisanal fishery (BOE 2009). This area also harbours the Special Area of Conservation (SAC) “Calahonda” (code: ES6170030; part of the EU Natura 2000 network), a biodiversity hot-spot within the Alboran and Mediterranean contexts (García Raso et al. 2010; Urrea et al. 2017a). Despite widely concerns in the last decades over the environmental impact of different fishing gears (e.g. dredges), which can result in a final loss of biodiversity (Hutchings 1990; Jones 1992; Thrush et al. 1998), the exploitation of commercial molluscs has been coexisting with this high biodiversity in the SAC “Calahonda” for many years, showing a sustainable use of marine resources by artisanal fishermen. Nevertheless, the littoral of Málaga attracts hundreds of thousands of tourists every year, with tourist arrivals

increasingly, and molluscs constitute an important element of the local gastronomy. This high demand of seafood in such touristy coasts could affect mollusc stocks.

Regarding commercial species, previous works in the Alboran Sea have focussed on selectivity with different trawling techniques (Baro et al. 2004a, b; Baro and Muñoz de los Reyes 2007), stock assessments of commercial fishes (Pérez Gil et al. 2004; Quetglas et al. 2004) and the growth and reproduction of different commercial molluscs (Tirado and Salas 1998; Tirado et al. 2002, b, 2003; Rodríguez de la Rúa et al. 2003). Nevertheless, there is scarce information regarding the spatial patterns of commercial mollusc species that are generally exploited by artisanal fisheries in the southern Iberian Peninsula (Chicharro et al. 2002; Rufino et al. 2010). This work aims to identify preferential sedimentary habitats of commercial molluscs within the SAC “Calahonda”, located within “Cala del Moral” fishing area. We also analyzed some information regarding the biota occurring in these fishing grounds, and discuss the importance of the establishment of the SAC for the artisanal fisheries.

Material and Methods

Study Site

The SAC “Calahonda” is located between Punta de Calaburras (36°30.4'N 04°38.3'W) and the marina of Cabopino (36°29'N 04°44.3'W) (S Spain), close to the Strait of Gibraltar (Fig. 1a). The northwestern coast of the Alboran Sea, where this SAC is located, is highlighted by its high biological productivity, promoted by the presence of almost permanent nutrient upwellings of deep Mediterranean waters (Sarhan et al. 2000). Main habitats within the SAC “Calahonda” include fragmented beds of the endemic Mediterranean seagrass *Posidonia oceanica* (Linnaeus) Delile, and of the subtropical seagrass *Cymodocea nodosa* (Ucria) Ascherson, both of them occurring amongst shallow rocky outcrops down to ca. 5 m depth; soft bottoms that include those with fine, medium and coarse sand as well as with bioclasts, which extend beyond the patchy seagrass beds; and finally a rocky outcrop at 15–20 m depth harbouring a highly diverse coralligenous community (García Raso et al. 2010; Urrea et al. 2012) (Fig. 1b). The soft bottoms harbour biodiverse molluscan assemblages with more than 200 species, including well sorted fine sands assemblages at 5 m depth (Sables Fins Bien Calibrás or SFBC), muddy bioclastic sands assemblages between 15 and 25 m depth (Détritiques Envasés or DE) and coastal bioclastic sands assemblages at 25 m depth (Détritiques Côtiers or DC) (Urrea et al. 2011), according to the




Fig. 1 Location of the Special Area of Conservation (SAC) “Calahonda” in southern Spain and sampling stations. The spatial distribution of different soft bottoms according to their granulometric composition within

the limits of the SAC is indicated with different colours (see more details in Urrea et al. 2011b)

classification of benthic communities defined for the Mediterranean by Pérès and Picard (1964).

Sample Collection and Laboratory Procedures

Faunistic samples were collected seasonally in summer (September 04), autumn (November 04), winter (February 05) and spring (May 05) with a small rock dredge (42 cm width, 22 cm height, mesh size: 4 mm) at three different depths (5, 15, 25 m) along four transects oriented perpendicularly to the coastline (Fig. 1a). A total of 117 samples were collected on the soft bottoms considered in this study. A detailed description of the sampling method is presented by Urrea et al. (2017b). Faunistic samples were processed in the laboratory, with live molluscs extracted from the sediment, identified and counted. The characterization of the sediment was done from sediment samples collected in each sampling station following the classification of Buchanan (1984) (see Urrea et al. 2011 for more details).

Data Analysis

The temporal variability of the eight commercial molluscan species identified from the faunistic samples was studied through the analyses of seasonal dynamics for the abundance values. Individuals of the bivalves *Acanthocardia tuberculata*, *Callista chione*, *Pecten*

maximus, *Venus verrucosa* and the gastropod *Hexaplex trunculus* with a shell length lower than 10 mm were considered small individuals, and those with a shell length higher than 10 mm were considered large individuals. On the other hand, individuals of *Chamelea gallina*, *Tapes rhomboides* and *Donax trunculus* with a shell length of less than 5 mm were considered small individuals, and those with a shell length higher than 5 mm were considered large individuals. One-factor ANOVA (Analysis of Variance) analyses were carried out for testing statistical differences in seasonal abundance values. Analyses to test the normality (Kolmogorov-Smirnov) and to verify the homogeneity of variances (Levene) were done prior to ANOVA analyses. A post-hoc Tukey test ($p < 0.05$) was used for posterior multiple comparisons. These statistical procedures were performed using the software SPSS.

A canonical correspondence analysis (CCA) was carried out in order to elucidate the relationships between the commercial species studied and their environment. This method extracts synthetic environmental gradients from the ecological data-sets, providing a general framework for statistical testing of the effects of sediment characteristics on the distribution of the species (ter Braak 1986). The statistical significance of the effect of each variable was tested by a Monte Carlo permutation test. This multivariate analysis was executed using the software CANOCO.

Results

Commercial Molluscs

Acanthocardia tuberculata was represented by 222 individuals. About ca. 69% of them were collected in fine and medium sand bottoms at 5 m depth, followed by ca. 18% in coarse sand bottoms (T2-15 m) and ca. 13% in muddy bioclastic sand bottoms (T3-15 m, T4-15 m) (Fig. 2; Table 1). Most of the collected individuals (ca. 77%) displayed a shell length higher than 10 mm (large individuals), and included adults with commercial size (45 mm). A similar number of small individuals (shell length < 10 mm) were found at 5 and 15 m depth, whereas large individuals were quantitatively more abundant at 5 m depth (Table 1). Mean abundance (N) values of *A. tuberculata* at 5 m depth, where this species was among the top-10 dominant (2.10% of total abundance) and frequent (66.6% Frequency of occurrence in total samples) species, displayed significant seasonal changes (one-factor ANOVA, $F = 5.359$, $p < 0.05$), with the highest N value for large and small *A. tuberculata* individuals observed in September (summer) and, on the other hand, the absence of individuals in May (spring) (Fig. 3).

A total of 219 individuals of *Callista chione* were collected. Although this species was present in most of the sampling stations, many of them were found at T2-15 m (ca. 48%) and at T1-5 m (ca. 24%), where it was one of the top-10 dominant (ca. 2.5%) and most frequent species (up to 100% F at T2-15 m) (Fig. 2; Table 1). Small individuals were more abundant at coarse sand bottoms (T2-15 m; 53% of small individuals) and large ones at shallow fine-medium sand bottoms (70% of large individuals), with maximum N values in November (autumn) at both depths. The mean N values, clustering the data from T2-15 m and T1-5 m, displayed significant seasonal changes (one-factor ANOVA, $F = 9.303$, $p < 0.05$), with maximum abundances in November and minimum ones in September for both large and small individuals (Fig. 3).

Chamelea gallina was mainly found inhabiting well sorted fine sands at T1-5 m (ca. 66%) and T3-5 m (ca. 23%) (Fig. 2; Table 1). This bivalve was present in every sample collected in this type of soft bottoms (100% F), where it displayed dominance values ranging between ca. 22% in spring and ca. 6% in winter, being one of the top-5 dominant species. Mean N values displayed a significant seasonal trend (one-factor ANOVA, $F = 6.229$, $p < 0.05$) with maxima in warm months (September and May) and minima in cold months (November and February), and the highest number of large and small individuals observed in September and May, respectively (Fig. 3).

Tapes rhomboides was found in deep sampling stations (15 and 25 m depth) (Fig. 2). This species was mainly represented by small individuals (ca. 98% from a total of 145 ind. collected) (Table 1). The highest N value was found in muddy bioclastic sand bottoms at T4-25 m (ca. 69%), where it displayed a dominance value of ca. 2.6%, followed by coastal

bioclastic sands at T2-25 m (21%) (Fig. 2). Most individuals found in these and other sampling stations were collected in May (ca. 95% from the total collected) (one-factor ANOVA, $F = 40.87$, $p < 0.05$) (Fig. 3).

Other commercial species were minoritarian and represented by less than 20 individuals such as in the case of *Donax trunculus*, only collected at the shallow stations T1-5 m (4 large ind. and 3 small ind.) and T2-5 m (1 large ind. and 9 small ind.) (one-factor ANOVA, $p > 0.05$) (Figs. 2 and 3), and *Hexaplex trunculus*, collected at the deep stations T2-15 m (6 large ind. and 2 small ind.), T2-25 m (6 large ind.) and T4-25 m (1 large ind.) (one-factor ANOVA, $p > 0.05$) (Figs. 2 and 3). Finally, *Pecten maximus* and *Venus verrucosa* were represented by four large individuals (two individuals collected in September and November, respectively) and one large individual (November) collected at 25 m depth, respectively (Fig. 2, Table 1; not shown in Fig. 3).

The CCA selected depth and percentage of bioclasts, for the first axis, and medium and very fine sand, for the second axis, as the variables explaining most of the species distribution, as detailed in Fig. 4. Thus, a gradient with depth can be observed from the left, with those species that inhabit soft bottoms between 5 and 15 m depth with low size granulometric fractions (very fine-medium sand) such as *C. gallina* and *D. trunculus*, to the right of axis I, with those ones that prefer deeper bottoms (15–25 m depth) with a higher content of coarse sand and bioclasts such as *H. trunculus* and *T. rhomboides* (Fig. 4). The Monte Carlo test was significant for the first axis ($p < 0.05$).

Associated Biota

Within the SAC “Calahonda”, rich faunistic assemblages have been found associated with the soft bottoms where commercial molluscs are collected. Regarding the well sorted fine sand bottoms where *D. trunculus* and *C. gallina* inhabit, the assemblage is represented by decapods including the dominant *Diogenes pugilator* (Roux, 1829) (79% of all decapods collected in the same samples) and *Philocheilus trispinosus* (Hailstone in Hailstone and Westwood, 1835), echinoids such as *Echinocardium mediterraneum* (Forbes, 1844) and ophiuroids such as *Amphipholis squamata* (Delle Chiaje, 1828). Detritic sandy bottoms with population of *H. trunculus*, *C. chione* and *P. maximus* are home of the hermit-crabs *Anapagurus hyndmanni* (Bell, 1846) and *Paguristes eremita* (Linnaeus, 1767), the echinoids *Genocidaris maculata* A. Agassiz, 1869 and *Echinocyamus pusillus* (O.F. Müller, 1776), the ophiuroids *Ophiopsila aranea* Forbes, 1843 and *Ophiothrix quinque maculata* (Delle Chiaje, 1828), as well as of individuals of the gorgoniid anthozoans *Eunicella gazella*

Fig. 2 Spatial distribution of commercial molluscs identified within the Special Area of Conservation “Calahonda” and total abundance of specimens collected in each sampling station











Table 1 Commercial molluscan species identified within the Special Area of Conservation (SAC) “Calahonda”, with quantitative data regarding the abundance of large and small individuals in fine and medium sand bottoms (Sables Fins Bien Calibrés or SFBC, according to Pérès and

Picard 1964), mixed coarse sand bottoms (MX), muddy bioclastic sand bottoms (Détritiques Envasés or DE, according to Pérès and Picard 1964) and coastal bioclastic sand bottoms (Détritiques Côtiers or DC, according to Pérès and Picard 1964)

	SFBC			MX	DE				DC		Total
	T1-5 m	T2-5 m	T3-5 m		T3-15 m	T4-15 m	T3-25 m	T4-25 m	T1-25 m	T2-25 m	
<i>Acanthocardia tuberculata</i>											0
Small indiv.		3	17	17	6	9					52
Large indiv.	63	23	48	22	11	3					170
Total	63	26	65	39	17	12					222
<i>Callista chione</i>											0
Small indiv.	26	17	10	94	6	8		3	9	6	179
Large indiv.	27		1	12							40
Total	53	17	11	106	6	8		3	9	6	219
<i>Chamelea gallina</i>											0
Small indiv.	161	6	30	6		7	1	1		1	213
Large indiv.	418	43	174	17		4				3	659
Total	579	49	204	23		11	1	1		4	872
<i>Tapes rhomboides</i>											0
Small indiv.				3		1	5	100	3	29	141
Large indiv.				2						2	4
Total				5		1	5	100	3	31	145
<i>Donax trunculus</i>											0
Small indiv.	1										1
Large indiv.	6	10									16
Total	7	10									17
<i>Hexaplex trunculus</i>											0
Small indiv.				2							2
Large indiv.				6				1		6	13
Total				8				1		6	15
<i>Pecten maximus</i>											0
Small indiv.											0
Large indiv.								3		1	4
Total								3		1	4
<i>Venus verrucosa</i>											0
Small indiv.										1	1
Large indiv.											0
Total										1	1


Studer, 1878 and *Leptogorgia sarmentosa* (Esper, 1789) that are found attached to small hard substrates such as rocks. Finally, detritic muddy bottoms harbour the echinoid *E. cordatum* (Pennant, 1777), the asteroid *Astropecten irregularis* (Pennant, 1777), dense groups of the epizoanthid anthozoan *Epizoanthus incrustatus* (Düben and Koren, 1847) with some rare architectonic gastropods that feed on them including *Pseudotorinia architae* (O. G. Costa, 1841), *Basisulcata lepida* (Bayer, 1942) and *Heliculus fallaciosus* (Tiberi, 1872), abundant hermit-crabs such as *Anapagurus alboranensis* García-Gómez, 1994 and *Anapagurus petiti*

Dehancé and Forest, 1962, and the pennatulid anthozoan *Pteroeides spinosum* (Ellis, 1764), among other taxa.

Discussion

The marine area of Calahonda was officially declared as a Site of Community Importance (SCI, code ES6170030) in 2006 (EU official journal 2006, September 21st), to protect some of the westernmost populations of the endangered Mediterranean seagrass *Posidonia oceanica*. Further studies focussed on the

Fig. 3 Seasonal dynamics of the commercial molluscs identified within the Special Area of Conservation “Calahonda”. White bars: small individuals (shell length < 10 mm for *A. tuberculata*, *C. chione* and *H. trunculus*, and < 5 mm for *C. gallina*, *T. rhomboides* and *D. trunculus*); grey bars: large individuals (shell length > 10 mm for *A. tuberculata*, *C. chione* and *H. trunculus*, and > 5 mm for *C. gallina*, *T. rhomboides* and *D. trunculus*). Mean + SE. Letters above error bars display the results of post-hoc test; different letters distinguish significantly different means at $p < 0.05$. Those species with non-significant seasonal differences in their abundance values are indicated with *



invertebrate communities inhabiting the different soft and hard bottoms around these seagrass meadows pointed out a high biodiversity that led to an increase of its extension in




Fig. 4 Graphical representation of the commercial molluscan species distribution with respect to the first two axes of the canonical correspondence analysis (CCA). The summary of the results of the CCA is also presented. % bioclats, percentage of bioclats; OM, organic matter

2009 (García Raso et al. 2010). The SCI “Calahonda” was confirmed as a Special Area of Conservation (SAC) in 2015 (DECRETO 369/2015, Andalusian Government official journal 2015, August 7th). Among this richness and regarding molluscs, approximately one third of the 1200 species coexisting in southern Spain can be found within the SAC (Gofas et al. 2011; Urrea et al. 2017a), which makes this coastal area of the northwestern Alboran Sea one of the molluscan hot-spots of European waters identified so far. Regarding molluscs with commercial value, eight species were collected in this study, including *Donax trunculus*, *Chamelea gallina*, *Callista chione*, *Acanthocardia tuberculata*, *Tapes rhomboides*, *Pecten maximus*, *Venus verrucosa* and *Hexaplex trunculus*. Besides, other commercial molluscs are found with the limits of the SAC such as the Mediterranean mussel *Mytilus galloprovincialis* Lamarck, 1819, with juveniles and adults settling on different hard substrates, the solid surf clam *Spisula solida* (Linnaeus, 1758) that presents small populations (<10 ind. collected) in the area, as well as different species of razor clams (*Ensis* spp.) and cephalopods (*Octopus vulgaris* Cuvier, 1797). In this context, and considering the presence of small-scale artisanal fisheries targeting molluscs of economic interest in the area, it is necessary to know the preferential habitat of each one of these species, together with the time period when recruitment events take place (Tirado et al. 2002a), as this information has important implications in terms of management and sustainable use of living resources, as well as for biodiversity conservation purposes.

A depth-related pattern associated with the granulometric composition of the seabed has been observed for the spatial

distribution of the commercial molluscan species found in the SAC “Calahonda”. The most abundant and frequent commercial species found in the study area was *C. gallina*, which has been mainly found inhabiting well sorted fine sand bottoms at 5 m depth (Pérès and Picard 1964; Gofas et al. 2011). In the other hand, the highly demanded *D. trunculus* was poorly represented in samples because it presents the bulk of its population at shallower depths (0–3 m), being usually replaced by populations of other Donacidae such as *D. venustus* Poli, 1795 from 5 m down to 10 m depth (Tirado and Salas 1999; Gaspar et al. 2002), as in the case presented here. Large individuals of *C. chione* and *A. tuberculata* were frequently collected at 5 m depth, whereas smaller individuals were collected at greater depths, being this trend more acute for *C. chione*. This could be due to the fishing activity in the area, as the artisanal fleet targets *C. chione* in bottoms deeper than 10 m, capturing every individual above minimum legal size. On the other hand, *A. tuberculata* displayed quantitatively larger individuals at shallower bottoms, but it showed a uniform distribution pattern of small individuals at 5 and 15 m depth. A true depth segregation pattern between size classes have previously been documented in other European coastal waters, especially concerning *Donax* species, and related to biological factors such as intraspecific competition among juveniles and adults (Ansell and Lagardère 1980; Tirado et al. 2002a; Gaspar et al. 2002). The only commercial gastropod identified from samples was *H. trunculus* and it was collected at deep stations, especially on rocky and coarse sand bottoms at 15 m depth, probably related to its trophic behaviour (i.e. carnivorous) and prey availability, including barnacles, bivalves and other gastropods usually inhabiting these bottoms. Minoritarian commercial bivalves included *T. rhomboides*, *P. maximus* and *V. verrucosa*, which were collected on deep bioclastic bottoms. The habitat preference for the abovementioned species is in line with data available in the literature and regarding European waters (Pérès and Picard 1964; Gofas et al. 2011). The spatial distribution pattern observed for these species is promoted by (1) depth related environmental factors such as granulometric characteristics of sediments and food supply, with a significant increasing trend of clay and mud fractions and of percentage of organic matter with depth (García Raso et al. 2010), and by (2) their different adaptive responses to the environment, with intense current, wave action and sediment instability in high energy environments (i.e. shallow bottoms) (Morin et al. 1985; Brown and McLachlan 1990; Koulouri et al. 2006).

Maximum abundances for most of the studied species were observed in spring and summer months, coinciding with the season of highest shellfish demand by markets. Several studies focussed on reproductive biology and population dynamics of commercial molluscan species in European coastal areas, especially in the Mediterranean Sea, have previously reported maximum abundances in

warm months, coinciding with important spawning periods (i.e. most intense release of gametes) and hatching juveniles (Tirado et al. 2002a; Rodríguez de la Rúa et al. 2003; Lahbib et al. 2011; Tirado et al. 2017). In the case of *A. tuberculata*, it is collected between autumn and winter only when the canning industry demands it. This practice can be considered a resting period for other exploited species as most artisanal fishing vessels targeting molluscs focus their activity on the tuberculate cockle during a period of two to three months of intensive fishing, being likely related to the absence of individuals collected in spring in the study area. In the case presented here, the environmental conditions registered in the northwestern Alboran Sea are characterized by mild seawater temperature and highly nutrient upwellings (Sarhan et al. 2000), which would provide good conditions for long reproductive cycles of commercial species such as molluscs (e.g. *Callista chione*, Tirado et al. 2002b, 2003). Nevertheless, a detailed environmental monitoring should be carried out in this and other areas of interest for artisanal fisheries along the northwestern Alboran Sea, as 1) sea surface temperature values have increased in the western Mediterranean Sea in the last years, including the Alboran basin (Vargas-Yáñez et al. 2010), and 2) an intensification of the decrease of the Ekman transport has been documented for the area (Vargas-Yáñez et al. 2010). The combination of these two processes could have negative effects in the food web structure and productivity, as the decrease of the intensity of upwellings will reduce the contribution of nutrients to the photic zone, with consequences in the primary production and in the rest of the trophic chain, which could be strengthened by warming (O'Connor et al. 2009).

An important issue to be considered, given that the SAC “Calahonda” is within the limits of the fishing area “Cala del Moral”, is the potential impact that dredging can have on the benthic fauna inhabiting the soft bottoms where these commercial species are collected. Bottom dredging and trawling can remove or damage a large part of the macro-epibenthic fauna as the fishing gear is towed across the seafloor (Hutchings 1990; Jennings and Kaiser 1998; Thrush et al. 1998). This can affect biomass, production and species richness of benthic invertebrate communities at a large scale, reducing the abundance of benthic food for commercial fish, and can lead to a loss of habitat heterogeneity and substrate complexity, as well as changes in community composition and size structure (Thrush et al. 1998; Kaiser and de Groot 2000; Queirós et al. 2006). The effects of trawling are more acute in sensitive environments such as detritic soft bottoms and especially hard bottoms, where large, long-lived and high ecologically value species (e.g. cnidarians) inhabit (Hiddink et al. 2006; Queirós et al. 2006).

Benthic communities associated with sedimentary environments within the SAC “Calahonda” seem to have a

relative good conservation status, as most of the species characterizing these communities (e.g. see Pérès and Picard 1964) have been found in the collected samples, besides a high species richness regarding different faunistic groups (García Muñoz et al. 2008; García Raso et al. 2010; Urrea et al. 2011). In this line, Urrea et al. (2017b) reported that only the 15% of the discarded individuals in the wedge clam mechanized dredging fisheries of the northern Alboran Sea, including the study area, displayed any type of damage. Decapods and specially echinoderms were reported by these authors to be vulnerable to dredging, as their bodies are fragile and easily fragmented by fishing gears, besides their autotomize capacity when subjected to stress (Bergmann and Moore 2001a, b). Urrea et al. (2017b) also highlighted that the target species (*Donax trunculus*) displayed a high proportion of undamaged individuals to fishing impacts, representing an important factor for the maintenance of populations of this commercial species on the fishing grounds. Moreover, faunistic assemblages inhabiting shallow and dynamic environments have showed a greater capacity to recover from fishing disturbances (Jones 1992; Currie and Parry 1996; Kaiser et al. 1998), and in the case presented here faunistic assemblages could also benefit from the high local productivity of the northern Alboran Sea (Sarhan et al. 2000). Furthermore, discards of dredging fisheries targeting molluscs in the northern Alboran Sea are released on suitable sedimentary habitats just after on-deck sorting, which would support the survival of discarded individuals. Nevertheless, an exhaustive program should be carried out to monitor 1) the conservation of species protected under different national and international frameworks (e.g. *Pinna nobilis*, *Charonia lampas*) and that have been observed within the limits of the SAC, 2) the different sedimentary habitats (habitat 1110 of the Natura 2000 Habitat Directive), vegetated meadows (e.g. *Posidonia oceanica* meadows, habitat 1120) and infralittoral rocky bottoms with coralligenous communities (habitat 1170) and the associated fauna existing within the SAC “Calahonda”, and 3) the amount of discards generated by the artisanal fleet, as well as the damage caused to invertebrate species, both commercial and non-target ones, by mechanical dredges, in order to maintain the good environmental status of this area.

Acknowledgements We would like to express our sincere gratitude to the captain and crew of the vessel AMA 8 for their cooperation during the sampling campaign; to the colleagues of our department mainly to Dr. Serge Gofas and Dr. J. Enrique García Raso for their help at different stages of this research; and to Terence Edwards for English revision of this manuscript. This work was supported by the ‘Consejería de Medio Ambiente de

la Junta de Andalucía’ (contract with the University of Málaga; reference 807/46.2284).

Compliance with Ethical Standards

Conflict of Interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- Ansell AD, Lagardère F (1980) Observations on the biology of *Donax trunculus* and *D. vittatus* at Ile d'Oléron (French Atlantic coast). *Mar Biol* 57(4):287–300
- Baro J, de los Reyes Muñoz I (2007) Comparación de los rendimientos pesqueros y la selectividad del arte de arrastre empleando mallas cuadradas y rómbicas en el copo, Informe Técnico del Instituto Español de Oceanografía, Centro Oceanográfico de Málaga, 23 pp
- Baro J, García T, del Árbol J, Garrucho S (2004a) Informe del seguimiento científico sobre una acción piloto de selectividad (RAI-AP-51/2004) con artes de arrastre en aguas del litoral de Málaga. Instituto Español de Oceanografía, Centro Oceanográfico de Málaga, 52 pp
- Baro J, Meléndez MJ, del Árbol J (2004b) Informe del seguimiento científico sobre una acción piloto de selectividad (RAI-AP-24/2003) con artes de arrastre en aguas del litoral de Málaga. Instituto Español de Oceanografía, Centro Oceanográfico de Málaga, 45 pp
- Bergmann M, Moore PG (2001a) Mortality of *Asterias rubens* y *Ophiura ophiura* discarded in the Nephrops fishery of the Clyde Sea area, Scotland. *ICES J Mar Sci* 58:531–542
- Bergmann M, Moore PG (2001b) Survival of decapods crustaceans discarded in the Nephrops fishery of the Clyde Sea area, Scotland. *ICES J Mar Sci* 58:163–171
- BOE (2009). Orden ARM/1995/2009, de 6 de julio, por la que se hacen públicas las nuevas relaciones de zonas de producción de moluscos y otros invertebrados marinos en el litoral español. Número 177, Sección III, pp. 63072. <http://www.boe.es/boe/dias/2009/07/23/pdfs/BOE-A-2009-12274.pdf>
- Brown AC, McLachlan A (1990) Ecology of sandy shores. Elsevier Publishers, Amsterdam, 328 pp
- Buchanan JB (1984) Sediment analysis. In: NA Holme ADMI (ed) Methods for the study of marine benthos. Blackwell, Oxford, pp 41–65
- Camiñas JA, Baro J, Abad R (2004) La pesca en el Mediterráneo andaluz, Servicio de publicaciones de la Fundación. Unicaja, Málaga, 270 pp
- Chemello R, Scotti G, Milazzo M (2000) L'uso della malacofauna marina in conservazione della natura. *Bollettino Malacologico* 36(1–4):49–60
- Chicharo L, Chicharo MA, Gaspar M, Alves F, Regala J (2002) Ecological characterization of dredged and recently dredge-free areas of a bivalve fishing ground, off South Portugal. *J Mar Biol Assoc U K* 82:41–50
- Currie DR, Parry GD (1996) Effects of scallop dredging on a soft sediment community: a large-scale experimental study. *Mar Ecol Prog Ser* 134:131–150
- FAO (2012) FAO 2010 yearbook, Fishery and Aquaculture statistics. <http://www.fao.org/fishery/publications/yearbooks/en>
- Fischer W, Bauchot ML, Schneider M (1987) Fiches FAO d'identification des espèces pour les besoins de la pêche, Méditerranée et mer Noire, Vol. I, Végétaux et Invertébrés, CEE, FAO Rome, 760 pp
- García Muñoz JE, Manjón-Cabeza ME, García Raso JE (2008) Decapod crustacean assemblages from littoral bottoms of the Alborán Sea (Spain, west Mediterranean Sea): spatial and temporal variability. *Sci Mar* 72(3):437–449

- García Raso JE, Gofas S, Salas Casanova C, Manjón-Cabeza ME, Uría J, García Muñoz JE (2010). El mar más rico de Europa: Biodiversidad del litoral occidental de Málaga entre Calaburras y Calahonda, Consejería de Medio Ambiente, Sevilla, 138 pp
- Gaspar MB, Chicharro LM, Vasconcelos P, García A, Santos AR, Monteiro CC (2002) Depth segregation phenomenon in *Donax trunculus* (Bivalvia: Donacidae) populations of the Algarve coast (southern Portugal). *Sci Mar* 66(2):111–121
- Gofas S, Moreno D, Salas C (2011) Moluscos marinos de Andalucía, Servicio de Publicaciones e Intercambio Científico, Universidad de Málaga, Málaga, 798 pp
- Hiddink JG, Jennings S, Kaiser MJ, Queirós AM, Duplisea DE, Piet GJ (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Can J Fish Aquat Sci* 63:721–736
- Hutchings P (1990) Review of the effects of trawling on macrobenthic Epifaunal communities. *Aust J Mar Freshwat Res* 41:111–120
- Jennings S, Kaiser MJ (1998) The effects of fishing on marine ecosystems. *Adv Mar Biol* 34:201–352
- Jones JB (1992) Environmental impact of trawling on the seabed: a review. *N Z J Mar Freshw Res* 26:59–67
- Kaiser MJ, de Groot SJ (2000) The effects of fishing on non-target species and habitats: biological. Conservation and Socio-economics Issues, Blackwell Science, Oxford, 399 pp
- Kaiser MJ, Edwards DB, Armstrong PJ, Radford K, Lough NEL, Flat RP, Jones HD (1998) Changes in megafauna benthic communities in different habitats after trawling disturbance. *ICES J Mar Sci* 55(3): 353–361
- Koulouri P, Dounas C, Arvanitidis C, Koutsoubas D, Eleftheriou A (2006) Molluscan diversity along a Mediterranean soft bottom sublittoral ecotone. *Sci Mar* 70:573–583
- Lahbib Y, Abidli S, Trigui El Menif N (2011) Reproductive activity in the commercially exploited mediterranean muricid *Hexaplex trunculus* collected from Boughrara lagoon (southern Tunisia). *Russ J Mar Biol* 37(6):501–508
- MAGRAMA (2012) Estrategia Marina Demarcación Marina del Estrecho y Alborán. Parte 1: Marco general evaluación inicial y buen estado ambiental. Ministerio de Agricultura, Alimentación y Medio Ambiente, Secretaría General Técnica, Centro de Publicaciones, 189 pp + Anexes
- Mazzella L, Buia MC, Gambi MC, Lorenti M, Russo GF, Scipione MB, Zupo V (1992) Plant-animal trophic relationship in the *Posidonia oceanica* ecosystem of the Mediterranean Sea: a review. In: DM John, SJ Hawkins, JH Price, eds, Plant-animal interactions in the marine benthos, systematics association special volume no. 46. Clarendon press, Oxford, pp. 165–187
- Morin JG, Kastendiek JE, Harrington A, Davis N (1985) Organization and patterns of interactions in a subtidal sand community on an exposed coast. *Mar Ecol Prog Ser* 27:163–185
- O'Connor MI, Piehler MF, Leech DM, Anton A, Bruno JF (2009) Warming and resource availability shift food web structure and metabolism. *PLoS Biol* 7(8):e1000178
- Pérès JM, Picard J (1964) Nouveau manuel de bionomie benthique de la Méditerranée. Recueil des Travaux de la Station Marine d'Endoume 31:1–137
- Pérez Gil JL, Fernández A, Baro J, González M, Massutí E, Quetglas A, Quintanilla L (2004) Stocks assessment of hake (*Merluccius merluccius*) from the trawl fishery off the geographical sub-area 01 (northern Alboran), report of the sixth session of the sub-Committee for Stock Assessment, GFCM-SAC, 17 pp
- Poutiers JM (1987) Bivalves. In: Fiches FAO d'identification des espèces pour les besoins de la pêche, Méditerranée et mer Noire, Végétaux et Invertébrés, FAO. Rome:370–512
- Queirós AM, Hiddink JG, Kaiser MJ, Hinz H (2006) Effects of chronic bottom trawling disturbance on benthic biomass, production and size spectra in different habitats. *J Exp Mar Biol Ecol* 335:91–103
- Quetglas A, Massutí E, del Árbol J, Baro J, Fernández A, González M, Pérez-Gil JL, Quintanilla L (2004) Stock assessment of red mullet (*Mullus barbatus*) from the trawl fishery off the geographical sub-area 01 (northern Alboran). Report of the Sixth Session of the Sub-Committee for Stock Assessment, GFCM-SAC, 17 pp
- Robles R (2010) Conservación y desarrollo sostenible del mar de Alborán, Gland. Suiza y Málaga, España, UICN, 112 pp
- Rodríguez de la Rúa A, Prado MA, Bruzón MA (2003). Estudio del ciclo reproductor de *Chamelea gallina* (L., 1758) (Mollusca: Bivalvia) en tres poblaciones del litoral andaluz, Boletín del Instituto Español de Oceanografía, 19(1–4): 57–63
- Rueda JL, Marina P, Uría J, Salas C (2009) Changes in the composition and structure of a molluscan assemblage due to eelgrass loss in southern Spain (Alboran Sea). *J Mar Biol Assoc U K* 89(7):1319–1330
- Rufino MM, Gaspar M, Pereira A, Maynou F, Monteiro CC (2010) Ecology of megabenthic bivalve communities from sandy beaches on the south coast of Portugal. *Sci Mar* 74(1):163–178
- Sarhan T, García Lafuente J, Vargas JM, Plaza F (2000) Upwelling mechanisms in the northwestern Alboran Sea. *J Mar Syst* 23:317–331
- Snelgrove PVR (1998) The biodiversity of macrofaunal organisms in marine sediments. *Biodivers Conserv* 7:1123–1132
- Templado J (2011) La diversidad marina en España. Memorias de la Real Sociedad Española de Historia Natural 9:343–362
- ter Braak CJF (1986) Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology* 67:1167–1179
- Thrush SF, Hewitt JE, Cummings VJ, Dayton PK, Cryer M, Turner SJ, GA Funnell BRG, Milburn CJ, Wilkinson MR (1998) Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecol Appl* 8(3):866–879
- Tirado C (2011) La pesca de moluscos en Andalucía. In: Gofas S, Moreno D, Salas C (eds) Moluscos marinos de Andalucía, Servicio de Publicaciones e Intercambio Científico, Universidad de Málaga, pp 31–37
- Tirado C, Salas C (1998) Reproduction and fecundity of *Donax trunculus* in the littoral of Málaga (S Spain). *J Shellfish Res* 17(1):169–176
- Tirado C, Salas C (1999) Reproduction of *Donax venustus* Poli 1795, *Donax semistriatus* Poli 1795 and intermediate morphotypes (Bivalvia: Donacidae) in the littoral of Málaga (southern Spain). *Mar Ecol* 20(2):111–130
- Tirado C, de la Rúa R, Franch A, Bruzón Gallego MA, López Linares JI, Salas Casanova C, Márquez Pascual I (2002a) La reproducción de bivalvos y gasterópodos de interés pesquero en Andalucía. Consejería de Agricultura y Pesca, Sevilla, 129 pp
- Tirado C, Salas C, López JI (2002b) Reproduction of *Callista chione* L., 1758 (Bivalvia: Veneridae) in the littoral of Málaga (southern Spain). *J Shellfish Res* 21(2):643–648
- Tirado C, de la Rúa R, Franch A, Bruzón Gallego MA, López Linares JI, Salas Casanova C, Márquez Pascual I (2003) La reproducción del pulpo (*Octopus vulgaris*) y el choco (*Sepia officinalis*) en la costa andaluza. Consejería de Agricultura y Pesca, Sevilla, 159 pp
- Tirado C, Marina P, Uría J, Antit M, Salas C (2017) Reproduction and population structure of *Acanthocardia tuberculata* (Linnaeus, 1758) (Bivalvia: Cardiidae) in southern Spain: implications for stock management. *J Shellfish Res* 36(1):61–68
- Uría J, Gofas S, Rueda JL, Marina P (2011) Molluscan assemblages in littoral soft bottoms of the Alboran Sea (western Mediterranean Sea). *Mar Biol Res* 7:27–42
- Uría J, Rueda JL, Gofas S, Marina P, Salas C (2012) A species-rich molluscan assemblage in a coralligenous bottom of the Alboran Sea (southwestern Mediterranean): intra-annual changes and ecological considerations. *J Mar Biol Assoc U K* 92(4):665–677
- Uría J, Gofas S, Rueda JL, Marina P, Mateo-Ramírez Á, Antit M, Salas C (2017a) Biodiversity and biogeographical patterns of molluscan assemblages in vegetated and unvegetated habitats in the northern Alboran Sea (W Mediterranean Sea). *Mar Biodivers* 47:187–201

- Urra J, García T, Gallardo-Roldán H, León E, Lozano M, Baro J, Rueda JL (2017b) Discard analysis and damage assessment in the wedge clam mechanized dredging fisheries of the northern Alboran Sea (W Mediterranean Sea). *Fish Res* 187:58–67
- Vargas-Yáñez M, García Martínez MC, Moya F, Tel E, Parrilla G, Plaza F, Lavín A, García MJ (2010) Cambio Climático en el Mediterráneo español. Instituto Español de Oceanografía, 176 pp
- Voultsiadou E, Koutsoubas D, Achparaki M (2010) Bivalve mollusc exploitation in Mediterranean coastal communities: an historical approach. *Journal of Biological Research-Thessaloniki* 13:35–45