

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/257549688>

# Severe reduction in genetic variation in a montane isolate: The endangered Mount Graham red squirrel (*Tamiasciurus hudsonicus grahamensis*)

Article in *Conservation Genetics* · December 2013

DOI: 10.1007/s10592-013-0511-x

CITATIONS

23

READS

160

3 authors:



**Robert R. Fitak**

University of Central Florida

61 PUBLICATIONS 1,034 CITATIONS

SEE PROFILE



**John Koprowski**

University of Wyoming

261 PUBLICATIONS 3,819 CITATIONS

SEE PROFILE



**Melanie Culver**

The University of Arizona

200 PUBLICATIONS 7,699 CITATIONS

SEE PROFILE

Severe reduction in genetic variation in a montane isolate: the endangered Mount  
Graham red squirrel (*Tamiasciurus hudsonicus grahamensis*)

Robert R. Fitak<sup>1</sup>, John L. Koprowski<sup>1,2</sup> and Melanie Culver<sup>1,2,3</sup>

<sup>1</sup>Graduate Interdisciplinary Program in Genetics, University of Arizona, Tucson,  
AZ, 85721, USA

<sup>2</sup>School of Natural Resources and the Environment, 1311 East Fourth Street,  
University of Arizona, Tucson, AZ, 85721, USA

<sup>3</sup>U. S. Geological Survey Arizona Cooperative Fish and Wildlife Research Unit,  
School of Natural Resources and the Environment, University of Arizona, Tucson,  
AZ, 85721, USA

**The Mount Graham red squirrel (*Tamiasciurus hudsonicus grahamensis*; MGRS) is endemic to the Pinaleno Mountains of Arizona at the southernmost extent of the species' range. The MGRS was listed as federally endangered in 1987, and is currently at high risk of extinction due to declining population size and increasing threats. Here we present a genetic assessment of the MGRS using eight nuclear DNA microsatellite markers and a 472 bp fragment of the mitochondrial cytochrome b gene. We analyzed 34 MGRS individuals and an additional 66 red squirrels from the nearby White Mountains, Arizona (*T. h. mogollonensis*). Both nuclear and mitochondrial DNA analyses revealed an extreme reduction in measures of genetic diversity relative to conspecifics from the White Mountains, suggesting that the MGRS has either experienced multiple bottlenecks, or a single long-term bottleneck. Additionally, we found a high degree of relatedness (mean =  $0.75 \pm 0.18$ ) between individual MGRS. Our study implies that the MGRS may lack the genetic variation required to respond to a changing environment. This is especially important considering this region of the southwest United States is expected to experience profound effects from global climate change. The reduced genetic variability together with the high relatedness coefficients should be taken into account when constructing a captive population to minimize loss of the remaining genetic variation.**

Keywords: endangered species, conservation, effective population size,  
inbreeding, Arizona

Corresponding Author:  
Robert R. Fitak  
Biosciences East Room 317  
1311 East Fourth Street  
Tucson, AZ 85721, USA  
Tel: 520-626-1636  
Fax: 520-621-8801  
Email: rfitak@email.arizona.edu

## 1    **Introduction**

2    Analysis of the factors that contribute to extinction is central to the conservation and  
3    management of endangered and threatened populations (Lande 1998). These populations  
4    become increasingly vulnerable to demographic stochasticity, environmental variation,  
5    random genetic effects, and extreme events like natural catastrophes (Schaffer 1981).  
6    Genetic effects are important because endangered species have small and/or declining  
7    population sizes, which result in an exponential loss of genetic diversity and an increased  
8    frequency of inbreeding (Frankham 2005). The loss of genetic diversity may jeopardize  
9    the ability for a species to respond to environmental changes (Frankel 1970; 1974), and  
10    inbreeding can reduce fitness through increased expression of deleterious traits  
11    (inbreeding depression; see Hedrick and Kalinowski 2000 for a review). Therefore, it is  
12    important for managers to evaluate endangered species from a genetic perspective to  
13    facilitate recovery and preserve remaining evolutionary potential.

14  
15    North American red squirrels (*Tamiasciurus hudsonicus*) are small (<300 g), diurnal  
16    squirrels found in boreal, mixed conifer, and deciduous forests throughout much of North  
17    America. Their distribution includes the northeastern United States, west through  
18    Canada and Alaska, and south along the Rocky Mountains to Arizona and New Mexico  
19    (Hall 1981). At the southernmost extent of the species' range exists an isolated  
20    population of a subspecies, the Mount Graham red squirrel (*T. h. grahamensis*; MGRS).  
21    The MGRS is restricted to high-elevation coniferous forests of the Pinaleño Mountains of  
22    southeastern Arizona (Hall 1981; Brown 1984). Although Allen (1894) originally  
23    described the MGRS as a unique subspecies due to pelage differences and geographic

isolation, both Hall (1981) and Hoffmeister (1986) acknowledged that morphology alone does not strongly differentiate it from its nearest conspecific, *T. h. mogollonensis*. The MGRS can be distinguished through vocalizations from *T. h. lychnuchus* of New Mexico, and its most common vocalization is different from that of *T. h. mogollonensis* (Yamamoto et al. 2001). The MGRS can also be differentiated from other red squirrels using mitochondrial restriction fragment length polymorphisms (RFLP; Riddle et al. 1992) and protein electrophoresis (Sullivan and Yates 1995). Unfortunately, the two most recent phylogenetic works on North American red squirrels did not include the MGRS (Arbogast et al. 2001; Wilson et al. 2005).

In 1987, the MGRS was listed as an endangered species because its range and habitat had been reduced, and remaining habitat was threatened by anthropogenic factors, fire, insect outbreaks, and possible competition with the introduced Abert's squirrel (U.S. Fish and Wildlife Service 1987; 1993; reviewed by Sanderson and Koprowski 2009). Historical accounts suggested that the MGRS was abundant (Mearns 1907; Hoffmeister 1986), but by the mid 1900's the population declined and was believed to be extirpated (Minckley 1968). Recent survey information indicates that the population has remained below 300 individuals since Fall of 2001 (U.S. Fish and Wildlife Service 2011). In response to the declining population size and increasing threats, a thorough genetic assessment employing both nuclear and mitochondrial DNA markers is necessary to develop a conservation strategy for this species.

In this study we used nuclear (microsatellites) and mitochondrial DNA markers to assess

the level and distribution of genetic diversity in the MGRS. To better evaluate these measures, we included a population of its geographically nearest conspecific, *T. h. mogollonensis*, for comparison. Additionally, managers are investigating the possibility of establishing a captive population of the MGRS (U.S. Fish and Wildlife Service 2011). A genetic analysis is necessary to identify the appropriate individuals to remove, to pair for breeding, and to release while acknowledging any population substructure and minimizing genetic loss (Lacy 1994; Ballou et al. 1995; 2010). We discuss our results in the context of establishing a captive breeding program in addition to the implications for conservation and recovery of the MGRS.

## **Materials and Methods**

### *Sample collection and DNA extraction*

We analyzed 100 red squirrel samples collected by the University of Arizona's MGRS Monitoring Program from September 2002 through November 2006. The squirrels were collected from the Pinaleno Mountains (*T. h. grahamensis*; MGRS; n = 34) and from two sites in the White Mountains (*T. h. mogollonensis*; WMRS; n = 66) of Arizona (Fig. 1). We selected either a small piece of ear tissue or 10-20 plucked hairs from live or deceased specimens and stored them at -20 °C. The use of >10 plucked hairs has provided reliable genotype estimates for microsatellites (Goossens et al. 1998). Live specimens were ear-tagged with unique numbered monel steel tags (model 1005-1, National Band & Tag Co., Newport, KY, USA) prior to release to avoid resampling of individuals. We extracted DNA using the Qiagen DNeasy Tissue Kit (Qiagen, Inc., Valencia, CA, USA) following the manufacturer's protocol.

*Microsatellite amplification and analysis*

We amplified eight microsatellite loci developed for *T. hudsonicus* and reported by Gunn et al. (2005) for each individual (Th08, Th14, Th21, Th23, Th25, Th33, Th41, and Th49). Fluorescent labeling was performed using the method detailed by Schuelke (2000). All PCR reactions contained 2 µl DNA template, 1.5 mM MgCl<sub>2</sub>, 0.2 mM each dNTP, 1.0 µM fluorescently labeled M13 primer (6-FAM, NED, or VIC), 0.1 µM forward primer, 1.0 µM reverse primer, 1 unit of AmpliTaq Gold<sup>®</sup> polymerase (Applied Biosystems, Foster City, CA, USA), 1X PCR buffer, and water in a total volume of 10 µl. Thermocycler conditions consisted of an initial denaturation at 94 °C for 2 min, 5 cycles (94 °C for 1 min, 64 °C for 30 s, 72 °C for 30 s), and 30 cycles (94 °C for 1 min, 55 °C for 30 s, 72 °C for 30 s). All fragment lengths were resolved on the same ABI3730 DNA Analyzer at the University of Arizona Genetics Core (UAGC; <http://uagc.arl.arizona.edu>) and alleles were scored using GENOTYPER v2.1 (Applied Biosystems). We subsequently re-amplified all ambiguous samples until a genotype could be determined or we did not assign a genotype.

We classified raw allele sizes into bins using FLEXIBIN v2 (Amos et al. 2007). We checked each locus in each population for scoring errors due to stutter, allele dropout, and null alleles using MICROCHECKER v2.2.3 (van Oosterhout et al. 2004). We performed exact tests for significant deviations from Hardy-Weinberg equilibrium (HWE) and for linkage disequilibrium (LD) between loci using a Markov chain method with a dememorization of 10,000 steps and 100 batches of 10,000 iterations each in GENEPOP

v4 (Rousset 2008). We adjusted the nominal  $p$ -value for multiple comparisons with a Bonferroni correction. We calculated general indices of genetic diversity, including number of alleles, allelic richness, and observed and expected heterozygosities using FSTAT v2.9.3.2 (Goudet 2001). We examined levels of inbreeding by computing the excess or deficiency in heterozygosity in individuals relative to the subpopulation ( $F_{IS}$ ). We tested  $F_{IS}$  computations for a statistical difference from zero using 1,000 permutations in FSTAT.

We used a Bayesian clustering algorithm implemented in STRUCTURE v2.3.2.1 (Pritchard et al. 2000; Falush et al. 2003; Hubisz et al. 2009) to determine the number of clusters (populations) without any *a priori* knowledge of population substructure. The estimated number of populations ( $K$ ) is given as the value  $K$  where the probability of the data,  $\Pr(X | K)$ , reaches a plateau. We tested for  $K = 1-8$  with ten iterations of each possible  $K$ . We assumed admixture and correlated allele frequencies and ran each analysis for 100,000 burn-in generations and 500,000 generations thereafter. For our most probable number of populations,  $K$ , we combined the ten replicates using the full-search algorithm in the software CLUMPP v1.1.2 (Jakobsson and Rosenberg 2007) and plotted the results using the software DISTRUCT v1.1 (Rosenberg 2004). We calculated effective population sizes using an approximate Bayesian method implemented in ONESAMP (Tallmon et al. 2008) using 50,000 iterations and the four loci that were polymorphic in all populations (Th08, Th14, Th23, Th41). Confidence intervals were assessed using the posterior distribution. We tested for presence of a population bottleneck using the software BOTTLENECK v1.2.02 (Cornuet and Luikart 1996; Piry et

al. 1999) implementing a two-phased model of mutation and assessed significance using a Wilcoxon's ranked test. We also used the mode-shift indicator in BOTTLENECK as qualitative evidence for a recent (within a few dozen generations) bottleneck event (Luikart et al. 1998). Pairwise relatedness was estimated in COANCESTRY v1.0.0 (Wang 2010) using the method of Lynch and Ritland (1999) and 1,000 bootstrap replications to examine differences between populations.

### *Mitochondrial DNA amplification and analysis*

We amplified a 472 bp fragment of the mitochondrial cytochrome b gene (cytb) in each individual using primers mcb398 and mcb869 (Verma and Singh 2003). The PCR reaction mix consisted of 3.0  $\mu$ l DNA template, 0.1  $\mu$ M forward and reverse primer, 0.2 mM each dNTP, 1.0 mM  $MgCl_2$ , 0.05 % BSA, 1X PCR buffer, 0.5 units *Taq* polymerase (Qiagen), and water to a final volume of 20  $\mu$ l. Thermocycling conditions included an initial denaturation step at 95 °C for 10 min, followed by 35 cycles (95 °C for 45 s, 52 °C for 1 min, 72 °C for 2 min) and a final extension step at 72 °C for 10 min. All PCR products were purified using a QIAquick PCR Purification Kit (Qiagen) and sequenced in both forward and reverse directions on an ABI3730 DNA Analyzer at the UAGC. All sequences were assembled, had primer bases removed, and were visually inspected for errors in SEQUENCHER 4.9 (Gene Codes Corp., Ann Arbor, MI, USA).

We used the software ARLEQUIN v3.5 (Excoffier and Lischer 2010) to organize the mitochondrial DNA (mtDNA) sequences into haplotypes and to calculate haplotype diversity ( $h$ ), nucleotide diversity ( $\pi$ ). We inferred the phylogenetic relationships



between cytb haplotypes using statistical parsimony in TCS v1.21 (Clement et al. 2000) with a probability of parsimony cutoff value of 0.95. Sequences from each of the recovered haplotypes were deposited into Genbank (Accession numbers KC306932-KC306944).

## Results

### *Microsatellite analyses*

Eight microsatellite loci were amplified and scored for 100 individuals (MGRS = 34, WMRS = 66). All eight loci were polymorphic in WMRS, whereas only four loci (50%, Thu21, Thu25, Thu33, and Thu49) were polymorphic in MGRS. Results from MICROCHECKER indicated that no scoring errors, allele dropout, or null alleles were present. After Bonferroni correction, no loci in either population deviated significantly from HWE, and no significant LD was detected.

We calculated several indices of genetic variability and compared them between the populations (Table 1). Genetic diversity as indicated by the number of alleles, allelic richness, and observed heterozygosity was consistently four to six times greater in WMRS relative to MGRS and significant in all cases using a Welch's t-test ( $p < 0.001$ ). We found a majority of alleles (61.5%) in MGRS to occur at high frequency ( $> 0.8$ ), whereas low frequency alleles were characteristic of WMRS (Fig. 2).

We used a Bayesian inference method of population clustering implemented in STRUCTURE to quantify an individual's proportion of membership in each of K

populations. The  $\text{Pr}(X | K)$  reached a maximum when two populations were assumed. One population consisted entirely of MGRS individuals and the other of WMRS individuals (Fig. 3). Three WMRS individuals, however, did share a moderate amount of ancestry ( $37 \pm 0.002\%$ ,  $19 \pm 0.001\%$ , and  $30 \pm 0.001\%$ ) with MGRS. Examination of additional values of  $K > 2$  revealed no evidence of substructure within MGRS (data not shown).

Mean estimates of effective population size were 10.4 (95% CI = 6.0 - 18.4) in MGRS and 48.8 (95% CI = 37.3 - 63.7) in WMRS. We were unable to detect a population bottleneck in both MGRS and WMRS (Wilcoxon's ranked test:  $p = 1.0$  and 0.098, respectively) although the allele frequency distribution in MGRS did have a shifted mode (Fig. 2). The average pairwise relatedness among MGRS individuals was  $0.75 \pm 0.18$ , significantly greater ( $p < 0.001$ ) than that observed in WMRS (mean =  $-0.01 \pm 0.34$ ; Fig. 4).

#### *Mitochondrial DNA analyses*

After removing primer bases, the final cytb amplicon length was 421 bp. Mitochondrial cytb sequence data also indicated similar reductions in measures of genetic variation for MGRS relative to WMRS. Only a single haplotype existed in MGRS ( $h = 0$ ), whereas we found 10 haplotypes in WMRS ( $h = 0.849 \pm 0.019$ ). Nucleotide diversity ( $\pi$ ) per locus was zero in MGRS and  $3.186 \pm 1.668$  in WMRS. The single MGRS haplotype was not shared with WMRS. The relationships between haplotypes using statistical parsimony indicated the MGRS haplotype (haplotype A) was at least two mutational

steps different from the nearest WMRS haplotypes (haplotypes F, H, and D; Fig. 5), but within the range of differences observed between WMRS haplotypes (maximum of seven steps between haplotypes B and E; Fig. 5)

## **Discussion**

The MGRS has experienced both recent population declines and an increase in threats that may severely impact its long-term viability (U.S. Fish and Wildlife Service 1987; 1993, reviewed by Sanderson and Koprowski 2009). These forces may also have profound effects on the amount and distribution of genetic diversity (Koprowski and Steidl 2009). In this study, we analyzed both nuclear and mitochondrial DNA markers to assess the genetic status of the MGRS.

### *Genetic variability*

For WMRS, measures of genetic variation are consistent with that reported in other North American red squirrels using the same loci (Gunn et al. 2005; Beatty et al. 2011; Kiesow et al. 2012), and heterozygosity is similar to the mean reported for sciurids (0.62, Garner et al. 2005). However, for MGRS, we uncovered a remarkable reduction of both nuclear and mitochondrial genetic diversity. We observed only four polymorphic nuclear loci, similar to Sullivan and Yates (1995) who reported no variation in the 26 allozymes examined. For mitochondrial DNA we found only a single haplotype in MGRS, consistent with the lack of variation in mitochondrial RFLPs reported by Riddle et al. (1992).

Such a drastic decrease in genetic variation is likely the result of many factors. For instance, populations at the periphery of a species' range, like the MGRS, often have lower genetic variation than the core population (Cassel and Tammaru 2003; Schwartz et al. 2003; Eckert et al. 2008; Huang et al. 2009). Additionally, a founder's effect resulting from early Holocene colonization of coniferous mountaintops in southeast Arizona and southwest New Mexico followed by local extinctions with the exception of those in the Pinaleno Mountains (Harris 1990; Sullivan and Yates 1995) may have reduced genetic variation. Similar histories have been suggested for other more northerly portions of the range (Wilson et al. 2005; Chavez et al. 2011). Finally, a dynamic demographic history that includes multiple population bottlenecks, and long-term small population size reduces genetic diversity. We were unable to detect a historical bottleneck from the genetic data, despite probable bottleneck events that include a devastating fire in 1685 that destroyed much of the forest in the Pinaleno Mountains (Grissino-Mayer et al. 1995) and a near extinction in the mid 1900's (Minckley 1968). We did detect a recent population bottleneck, possibly the result of a series of fires and insect outbreaks from 1996 – 2002 that reduced census population estimates by more than half (U.S. Fish and Wildlife Service 2011).

Another striking result is the high pairwise estimates of relatedness between MGRS individuals. We chose the method of Lynch and Ritland (1999), a robust estimator that outperforms many other methods in unstructured populations and can be calculated for biallelic loci (Lynch and Ritland 1999; Olliehoek et al. 2006; Toro et al. 2003). Despite HWE and evidence for random mating ( $F_{IS} \sim 0$ ), a MGRS is, on average, related to a high

degree (mean =  $0.75 \pm 0.18$ ) to any other member of the population. This suggests that mating in the MGRS is random, but, due to their low levels of variation and small population size, individuals often encounter mates that are genetically similar.

#### *Population structure*

Analyses of population structure using nuclear DNA markers revealed a strict pattern of allele frequency differentiation between MGRS and its nearest conspecific. A few WMRS individuals maintained some shared ancestry with MGRS, possibly due to a residual effect of their common origin during the late Pleistocene (Harris 1990; Sullivan and Yates 1995), or the result of immigration and subsequent gene flow from MGRS into WMRS. We found no evidence of gene flow from WMRS into MGRS. Within MGRS, we did not detect the presence of subpopulations. This suggests that MGRS has maintained at least some level of gene flow across different regions of the Pinaleño Mountains, but more extensive sampling is required to substantiate this result.

We also assessed genetic structure using mitochondrial DNA, because matrilinear patterns of genetic structure may be different than that reported from nuclear loci if sex-biased dispersal exists (see Handley and Perrin 2007 for a review). Additionally, hypervariable loci, *i.e.* microsatellites, may overestimate differentiation after large reductions in population size (Hedrick 1999). We found a single cytb haplotype in MGRS that was not shared with WMRS. This demonstrates a similar degree of isolation and subsequent genetic drift as observed in the microsatellite results.

*Conservation implications*

The maintenance of genetic diversity is an important factor in conservation because it is directly correlated with fitness and evolutionary potential (Allendorf and Ryman 2002; Reed and Frankham 2003). In the MGRS, evidence of reduced fitness measured by decreased litter size and survivorship relative to conspecifics has been reported (Rushton et al 2006; Munroe et al. 2009; Zugmeyer and Koprowski 2009). Additionally, many ecosystems in the southwestern U.S., similar to the Pinaleño Mountains, are particularly vulnerable to climate change and will likely experience increased drought, insect outbreaks, and fires (Archer and Predick 2008; Allen et al. 2010). These factors render the MGRS at a high risk of extinction, and management strategies should be designed to mitigate this risk.

Management strategies for a taxon with a single population are limited and include: increasing the population size, establishing additional populations to avoid catastrophes to the single extant population, maximizing the reproductive rate, and insulating from environmental change (Frankham et al. 2002). Our results do not provide evidence that the MGRS is a distinct, phylogenetic unit but rather a population with extensive allele frequency differentiation from its nearest conspecific, likely the result of isolation and extensive genetic drift. Therefore, increasing genetic diversity through the translocation of individuals from WMRS remains a possibility. Such introductions have been successful in other isolated and/or inbred populations (Westemeier et al. 1998; Madsen et al. 1999; Johnson et al. 2010; Olson et al. 2012). However, isolated and rare populations, like the MGRS, may contain a suite of locally adaptive variation that can be negatively

impacted through outbreeding with maladaptive individuals (Lande and Shannon 1996; Tufto 2001). Although behaviors such as vocalizations (Yamamoto et al. 2001), home range size (Koprowski et al. 2008), caching (Angell 2009) and nest use (Leonard and Koprowski 2009) differ between Mt. Graham and other subspecies, we do not know if these represent adaptations or simply behavioral plasticity. Nonetheless, future genetic work using additional nuclear loci, such as a genome-wide set of single nucleotide polymorphisms and gene sequences, will be necessary to investigate the taxonomic status of the MGRS.

Current management actions include a proposal to establish a captive population from 16 founders (U.S. Fish and Wildlife Service 2010, 2011). The removal and pairing of individuals to establish a captive population should consider the high coefficients of relatedness we observed to minimize the loss of remaining genetic variation. Finally, because populations can adapt to captivity in as little as one generation (Christie et al. 2012), we believe that preservation of tissue (*i.e.* gametes, embryos, other tissue) in genome resource banks is prudent. The preservation of this material can extend the generation interval and provide genomic material for future studies (Johnston and Lacy 1995).

## *Conclusions*

Our results demonstrated that the MGRS persists despite incredibly low levels of genetic variation. However, due to demographic, environmental, and genetic concerns, the long-term persistence of the MGRS is in jeopardy. Although some taxa have maintained small

population sizes and/or high levels of inbreeding for long periods of time (see Craig 1994 for examples), many such taxa are extinct. Therefore, informed management actions for the MGRS that incorporate genetics in addition to demographics, natural history, and the environment will be most effective. Finally, studies employing genome-wide markers or sequencing may be useful for identifying important adaptive and detrimental variation that can be managed accordingly (Kohn et al. 2006; Ouborg et al. 2010).

### **Acknowledgments**

We thank A.Naidu, A. Ochoa, J. Leonard, and two anonymous reviewers for their helpful comments on this manuscript. We would also like to thank the individuals who contributed to collecting tissue samples, especially K. Munroe and V. Greer, and T. Dee, D. Sotelo and G. Reida for help with DNA extractions. This research was supported through an Arizona Game and Fish Heritage Program grant to MC and JLK and funds from the USDA Forest Service and the University of Arizona to JLK. RRF was supported by a Science Foundation Arizona fellowship and an NSF-IGERT fellowship in comparative genomics. Mention of specific products does not constitute endorsement by the U.S. Geological Survey.

### **References**

1. Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N, Vennetier M, Kitzeberger T, Rigling A, Breshears DD, Hogg EH, Gonzalez P, Fensham R, Zhang Z, Castro J, Demidova N, Lim JH, Allard G, Running SW, Semerci A, Cobb N (2010) A



- 323 global overview of drought and heat-induced tree mortality reveals emerging climate  
324 change risks for forests. *For Ecol Manage* 259 (4):660-684.  
325 doi:10.1016/j.foreco.2009.09.001
- 326 2. Allen JA (1894) Descriptions of ten new North American mammals, and remarks on  
327 others. *Bull Am Mus Nat Hist* 6:320-321
- 328 3. Allendorf FW, Ryman N (2002) The role of genetics in population viability analysis. In:  
329 Beissinger SR, McCullough DR (eds) *Population viability analysis*. University of  
330 Chicago Press, Chicago, pp 50-85
- 331 4. Amos W, Hoffman JI, Frodsham A, Zhang L, Best S, Hill AVS (2007) Automated  
332 binning of microsatellite alleles: problems and solutions. *Mol Ecol Notes* 7 (1):10-14.  
333 doi:10.1111/j.1471-8286.2006.01560.x
- 334 5. Angell DK (2009) Removal rates and fate of two cone species collected by red squirrels.  
335 In: Sanderson HR, Koprowski JL (eds) *The last refuge of the Mt. Graham red squirrel:  
336 ecology of endangerment*. University of Arizona Press, Tucson, pp 230-243
- 337 6. Arbogast BS, Browne RA, Weigl PD (2001) Evolutionary genetics and Pleistocene  
338 biogeography of North American tree squirrels (*Tamiasciurus*). *J Mammal* 82 (2):302-  
339 319
- 340 7. Archer SR, Predick KI (2008) Climate change and the ecosystems of the southwestern  
341 United States. *Rangelands* 30:23-28. doi: [http://dx.doi.org/10.2111/1551-  
342 501X\(2008\)30\[23:CCAEOT\]2.0.CO;2](http://dx.doi.org/10.2111/1551-501X(2008)30[23:CCAEOT]2.0.CO;2)
- 343 8. Ballou JD, Lacy RC (1995) Identifying genetically important individuals for management  
344 of genetic variation in pedigreed populations. In: Ballou JD, Gilpin ME, Foose TJ (eds)  
345 *Population management for survival and recovery: analytical methods and strategies in*

- 346 small population conservation. Methods and cases in conservation science. Columbia  
347 University Press, New York, pp 76-111
- 348 9. Ballou JD, Lees C, Faust LJ (2010) Demographic and genetic management of captive  
349 populations. In: Kleiman DG, Thompson KV, Baer CK (eds) Wild mammals in captivity:  
350 principles and techniques for zoo management. University of Chicago Press, Chicago, pp  
351 219-252
- 352 10. Beatty WS, Liu Z, Latch EK, Nelson TA (2011) Genetic assessment of the red squirrel in  
353 Illinois: immigrants or exotics? J Wildl Manage 75 (5):1236-1242.  
354 doi:10.1002/jwmg.156
- 355 11. Brown DE (1984) Arizona's tree squirrels. Arizona Game and Fish Department, Phoenix
- 356 12. Cassel A, Tammaru T (2003) Allozyme variability in central, peripheral and isolated  
357 populations of the scarce heath (*Coenonympha hero* : Lepidoptera, Nymphalidae):  
358 implications for conservation. Conserv Genet 4 (1):83-93. doi:10.1023/a:1021884832122
- 359 13. Chavez AS, Saltzberg CJ, Kenagy GJ (2011) Genetic and phenotypic variation across a  
360 hybrid zone between ecologically divergent tree squirrels (*Tamiasciurus*). Mol Ecol 20  
361 (16):3350-3366. doi: 10.1111/j.1365-294X.2011.05184.x
- 362 14. Christie MR, Marine ML, French RA, Blouin MS (2012) Genetic adaptation to captivity  
363 can occur in a single generation. Proc Natl Acad Sci USA 109 (1):238-242.  
364 doi:10.1073/pnas.1111073109
- 365 15. Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene  
366 genealogies. Mol Ecol 9 (10):1657-1659. doi:10.1046/j.1365-294x.2000.01020.x
- 367 16. Cornuet JM, Luikart G (1996) Description and power analysis of two tests for detecting  
368 recent population bottlenecks from allele frequency data. Genetics 144 (4):2001-2014

17. Craig JL (1994) Meta-populations: is management as flexible as nature? In: Olney PJS, Mace GM, Feistner ATC (eds) Creative conservation: interactive management of wild and captive animals. Chapman and Hall, London, pp 50-66
18. Eckert CG, Samis KE, Loughheed SC (2008) Genetic variation across species' geographical ranges: the central-marginal hypothesis and beyond. *Mol Ecol* 17 (5):1170-1188. doi:10.1111/j.1365-294X.2007.03659.x
19. Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* 10 (3):564-567. doi:10.1111/j.1755-0998.2010.02847.x
20. Falush D, Stephens M, Pritchard JK (2003) Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* 164 (4):1567-1587
21. Frankel OH (1970) Variation, the essence of life. *Proc Linn Soc NSW* 95:158-169
22. Frankel OH (1974) Genetic conservation - our evolutionary responsibility. *Genetics* 78 (1):53-65
23. Frankham R (2005) Genetics and extinction. *Biol Conserv* 126 (2):131-140. doi:10.1016/j.biocon.2005.05.002
24. Frankham R, Ballou JD, Briscoe DA (2002) Introduction to conservation genetics. Cambridge University Press, Cambridge
25. Garner A, Rachlow JL, Hicks JF (2005) Patterns of genetic diversity and its loss in mammalian populations. *Conserv Biol* 19 (4):1215-1221. doi:10.1111/j.1523-1739.2005.00105.x
26. Goossens B, Waits LP, Taberlet P (1998) Plucked hair samples as a source of DNA:

392 reliability of dinucleotide microsatellite genotyping. *Mol Ecol* 7 (9):1237-1241.  
 393 doi:10.1046/j.1365-294x.1998.00407.x

394 27. Goudet J (2001) FSTAT, a program to estimate and test gene diversities and fixation  
 395 indices (version 2.9.3). <http://www.unil.ch/popgen/softwares/fstat.htm>. Accessed October  
 396 2009

397 28. Grissino-Mayer HD, Baisan CH, Swetnam TW (1995) Fire history in the Pinaleno  
 398 Mountains of southeastern Arizona: effects of human-related disturbances. *Biodiversity*  
 399 and management of the Madrean Archipelago: the sky islands of southwestern United  
 400 States and northwestern Mexico. U.S. Forest Service General Technical Report, RM-  
 401 GTR-264:399-407

402 29. Gunn MR, Dawson DA, Leviston A, Hartnup K, Davis CS, Strobeck C, Slate JON,  
 403 Coltman DW (2005) Isolation of 18 polymorphic microsatellite loci from the North  
 404 American red squirrel, *Tamiasciurus hudsonicus* (Sciuridae, Rodentia), and their cross-  
 405 utility in other species. *Mol Ecol Notes* 5 (3):650-653. doi:10.1111/j.1471-  
 406 8286.2005.01022.x

407 30. Hall ER (1981) The mammals of North America. vol 1, Second edn. John Wiley and  
 408 Sons, New York

409 31. Handley LJJ, Perrin N (2007) Advances in our understanding of mammalian sex-biased  
 410 dispersal. *Mol Ecol* 16 (8):1559-1578. doi:10.1111/j.1365-294X.2006.03152.x

411 32. Harris AH (1990) Fossil evidence bearing on southwestern mammalian biogeography. *J*  
 412 *Mammal* 71 (2):219-229. doi:10.2307/1382170

413 33. Hedrick PW (1999) Perspective: highly variable loci and their interpretation in evolution  
 414 and conservation. *Evolution* 53 (2):313-318

34. Hedrick PW, Kalinowski ST (2000) Inbreeding depression in conservation biology. *Annu Rev Ecol Syst* 31:139-162. doi:10.1146/annurev.ecolsys.31.1.139
35. Hoffmeister DF (1986) *Mammals of Arizona*. University of Arizona Press, Tucson
36. Huang ZH, Liu NF, Chen YK, Xiao YA (2009) Genetic diversity in peripheral and central populations of rusty-necklaced partridge (*Alectoris magna*) based on mitochondrial and microsatellite DNA. *Acta Zool Acad Sci Hung* 55 (2):187-197
37. Hubisz MJ, Falush D, Stephens M, Pritchard JK (2009) Inferring weak population structure with the assistance of sample group information. *Mol Ecol Resour* 9 (5):1322-1332. doi:10.1111/j.1755-0998.2009.02591.x
38. Jakobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* 23 (14):1801-1806. doi:10.1093/bioinformatics/btm233
39. Johnson WE, Onorato DP, Roelke ME, Land ED, Cunningham M, Belden RC, McBride R, Jansen D, Lotz M, Shindle D, Howard J, Wildt DE, Penfold LM, Hostetler JA, Oli MK, O'Brien SJ (2010) Genetic restoration of the Florida panther. *Science* 329 (5999):1641-1645. doi:10.1126/science.1192891
40. Johnston LA, Lacy RC (1995) Genome resource banking for species conservation - selection of sperm donors. *Cryobiology* 32 (1):68-77. doi:10.1006/cryo.1995.1006
41. Kohn MH, Murphy WJ, Ostrander EA, Wayne RK (2006) Genomics and conservation genetics. *Trends Ecol Evol* 21 (11):629-637. doi:10.1016/j.tree.2006.08.001
42. Kiesow AM, Monroe EM, Britten HB (2012) Genetic structure of the arboreal squirrels (*Glaucomys sabrinus* and *Tamiasciurus hudsonicus*) in the North American Black Hills. *Can J Zoolog* 90 (9):1191-1200. doi: 10.1139/z2012-087

43. Koprowski JL, King SRB, Merrick MJ (2008) Expanded home ranges in a peripheral population: space use by endangered Mt. Graham red squirrels. *Endang Species Res* 4 (1-2):227-232. doi: 10.3354/esr00026
44. Koprowski JL and Steidl RJ (2009) Consequences of small populations and their impacts on Mt. Graham red squirrels. In: Sanderson HR, Koprowski JL (eds) *The last refuge of the Mt. Graham red squirrel: ecology of endangerment*. University of Arizona Press, Tucson, pp 142-152
45. Lacy RC (1994) Managing genetic diversity in captive populations of animals. In: Bowles ML, Whelan CJ (eds) *Restoration and recovery of endangered plants and animals*. Cambridge University Press, Cambridge, pp 63-89
46. Lande R, Shannon S (1996) The role of genetic variation in adaptation and population persistence in a changing environment. *Evolution* 50 (1):434-437. doi:10.2307/2410812
47. Lande R (1998) Anthropogenic, ecological and genetic factors in extinction and conservation. *Res Popul Ecol* 40 (3):259-269. doi:10.1007/bf02763457
48. Leonard KM, Koprowski JL (2009) A comparison of habitat use and demography of red squirrels at the southern edge of their range. *Amer Midl Nat* 162 (1):125-138. doi: <http://dx.doi.org/10.1674/0003-0031-162.1.125>
49. Luikart G, Allendorf FW, Cornuet JM, Sherwin WB (1998) Distortion of allele frequency distributions provides a test for recent population bottlenecks. *J Hered* 89 (3):238-247. doi:10.1093/jhered/89.3.238
50. Lynch M, Ritland K (1999) Estimation of pairwise relatedness with molecular markers. *Genetics* 152 (4):1753-1766
51. Madsen T, Shine R, Olsson M, Wittzell H (1999) *Conservation biology - Restoration of*

- an inbred adder population. *Nature* 402 (6757):34-35. doi:10.1038/46941
52. Mearns EA (1907) Mammals of the Mexican boundary of the United States: a descriptive catalogue of the species of mammals occurring in that region; with a general summary of the natural history, and a list of trees. *B US Natl Mus* 56 pt1:1-530
53. Minckley WL (1968) Possible extirpations of the spruce squirrel from the Pinaleno (Graham) Mountains, southcentral Arizona. *J Ariz Acad Sci* 5:110
54. Munroe KE, Koprowski JL, Greer VL (2009) Reproductive ecology and home range size of squirrels: do Mt. Graham red squirrels fit the pattern? In: Sanderson HR, Koprowski JL (eds) *The last refuge of the Mt. Graham red squirrel: ecology of endangerment*. University of Arizona Press, Tucson, pp 287-298
55. Oliehoek PA, Windig JJ, van Arendonk JAM, Bijma P (2006) Estimating relatedness between individuals in general populations with a focus on their use in conservation programs. *Genetics* 173 (1):483-496. doi:10.1534/genetics.105.049940
56. Olson ZH, Whittaker DG, Rhodes OE (2012) Evaluation of experimental genetic management in reintroduced bighorn sheep. *Ecol Evol* 2 (2):429-443. doi:10.1002/ece3.97
57. Ouborg NJ, Angeloni F, Vergeer P (2010) An essay on the necessity and feasibility of conservation genomics. *Conserv Genet* 11 (2):643-653. doi:10.1007/s10592-009-0016-9
58. Piry S, Luikart G, Cornuet JM (1999) BOTTLENECK: a computer program for detecting recent reductions in the effective population size using allele frequency data. *J Hered* 90 (4):502-503. doi:10.1093/jhered/90.4.502
59. Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155 (2):945-959

60. Reed DH, Frankham R (2003) Correlation between fitness and genetic diversity. *Conserv Biol* 17 (1):230-237
61. Riddle BR, Yates TL, Lee Jr. TE (1992) Molecular divergence and variation in the endangered Mount Graham red squirrel (*Tamiasciurus hudsonicus grahamensis*). Final report submitted to the Mount Graham Red Squirrel Study Committee. University of Arizona, Tucson
62. Rosenberg NA (2004) DISTRUCT: a program for the graphical display of population structure. *Mol Ecol Notes* 4 (1):137-138. doi:10.1046/j.1471-8286.2003.00566.x
63. Rousset F (2008) GENEPOP'007: a complete re-implementation of the genepop software for Windows and Linux. *Mol Ecol Resour* 8 (1):103-106. doi:10.1111/j.1471-8286.2007.01931.x
64. Rushton SP, Wood DJA, Lurz PWW, Koprowski JL (2006) Modelling the population dynamics of the Mt. Graham red squirrel: can we predict its future in a changing environment with multiple threats? *Biol Conserv* 131 (1):121-131. doi:10.1016/j.biocon.2006.02.010
65. Sanderson HR and Koprowski JL (2009) The last refuge of the Mt. Graham red squirrel: ecology of endangerment. University of Arizona Press, Tucson.
66. Schuelke M (2000) An economic method for the fluorescent labeling of PCR fragments. *Nat Methods* 18 (2):233-234. doi:10.1038/72708
67. Schwartz MK, Mills LS, Ortega Y, Ruggiero LF, Allendorf FW (2003) Landscape location affects genetic variation of Canada lynx (*Lynx canadensis*). *Mol Ecol* 12 (7):1807-1816. doi:10.1046/j.1365-294X.2003.01878.x
68. Shaffer ML (1981) Minimum population sizes for species conservation. *Bioscience* 31



507 (2):131-134

508 69. Sullivan RM, Yates TL (1995) Population genetics and conservation biology of relic  
509 populations of red squirrels. In: Istock CA, Hoffman RS (eds) Storm over a mountain  
510 island: conservation biology and the Mt. Graham affair. University of Arizona Press,  
511 Tucson

512 70. Tallmon DA, Koyuk A, Luikart G, Beaumont MA (2008) ONeSAMP: a program to  
513 estimate effective population size using approximate Bayesian computation. Mol Ecol  
514 Resour 8 (2):299-301. doi:10.1111/j.1471-8286.2007.01997.x

515 71. Toro MA, Barragan C, Ovilo C (2003) Estimation of genetic variability of the founder  
516 population in a conservation scheme using microsatellites. Anim Genet 34 (3):226-228.  
517 doi:10.1046/j.1365-2052.2003.00988.x

518 72. Tufto J (2001) Effects of releasing maladapted individuals: A demographic-evolutionary  
519 model. Am Nat 158 (4):331-340. doi:10.1086/321987

520 73. U. S. Fish and Wildlife Service (1987) Endangered and threatened wildlife and plants;  
521 determination of endangered status for the Mount Graham red squirrel. Fed Regist 52  
522 (106):20994-20999

523 74. U. S. Fish and Wildlife Service (1993) Mount Graham red squirrel recovery plan. U.S.  
524 Fish and Wildlife Service, Albuquerque

525 75. U. S. Fish and Wildlife Service (2010) Draft environmental assessment - establishment of  
526 a captive breeding program for the endangered Mount Graham red squirrel. U.S. Fish and  
527 Wildlife Service, Phoenix

528 76. U. S. Fish and Wildlife Service (2011) Draft Mount Graham red squirrel recovery plan,  
529 first revision (*Tamiasciurus hudsonicus grahamensis*). U.S. Fish and Wildlife Service,

Albuquerque

77. Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Notes* 4 (3):535-538. doi:10.1111/j.1471-8286.2004.00684.x
78. Verma SK, Singh L (2003) Novel universal primers establish identity of an enormous number of animal species for forensic application. *Mol Ecol Notes* 3 (1):28-31. doi:10.1046/j.1471-8286.2003.0340.x
79. Wang J (2011) COANCESTRY: a program for simulating, estimating and analysing relatedness and inbreeding coefficients. *Mol Ecol Resour* 11 (1):141-145. doi:10.1111/j.1755-0998.2010.02885.x
80. Westemeier RL, Brawn JD, Simpson SA, Esker TL, Jansen RW, Walk JW, Kershner EL, Bouzat JL, Paige KN (1998) Tracking the long-term decline and recovery of an isolated population. *Science* 282 (5394):1695-1698. doi:10.1126/science.282.5394.1695
81. Wilson GM, Van Den Bussche RA, McBee K, Johnson LA, Jones CA (2005) Intraspecific phylogeography of red squirrels (*Tamiasciurus hudsonicus*) in the central Rocky Mountain region of North America. *Genetica* 125 (2-3):141-154. doi:10.1007/s10709-005-5154-5
82. Yamamoto O, Moore B, Brand L (2001) Variation in the bark call of the red squirrel (*Tamiasciurus hudsonicus*). *West N Am Nat* 61 (4):395-402
83. Zugmeyer CA, Koprowski JL (2009) Severely insect-damaged forest: a temporary trap for red squirrels? *Forest Ecol Manag* 257 (2):464-470. doi:10.1016/j.foreco.2008.09.020

## Tables

MGRS						
Locus	n	A	A <sub>R</sub>	H <sub>O</sub>	H <sub>E</sub>	F <sub>IS</sub> *
Th8	34	3	2.941	0.265	0.329	0.173
Th14	33	2	2	0.242	0.216	-0.123
Th21	34	1	1	0.000	0.000	N/A
Th23	34	2	2	0.206	0.234	0.118
Th25	32	1	1	0.000	0.000	N/A
Th33	34	1	1	0.000	0.000	N/A
Th41	34	2	2	0.147	0.138	-0.065
Th49	32	1	1	0.000	0.000	N/A
Mean	33.375	1.625	1.618	0.108	0.115	0.052
WMRS						
Locus	n	A	A <sub>R</sub>	H <sub>O</sub>	H <sub>E</sub>	F <sub>IS</sub> *
Th8	66	13	12.484	0.924	0.903	-0.023
Th14	66	14	12.757	0.848	0.901	0.059
Th21	66	5	4.485	0.621	0.618	-0.006
Th23	66	6	5.899	0.682	0.745	0.085
Th25	66	4	3.861	0.182	0.224	0.189
Th33	66	5	4.737	0.697	0.730	0.045
Th41	64	7	6.871	0.844	0.816	-0.034
Th49	63	6	5.996	0.698	0.719	0.029
Mean	65.375	7.500	7.136	0.687	0.707	0.028

**Table 1** Summary of microsatellite data analysis for red squirrels (*Tamiasciurus hudsonicus*) in the White (WMRS: *T. h. mogollonensis*) and the Pinaleno (MGRS: *T. h. grahamensis*) Mountains, Arizona. The columns indicate number of successful genotypes (n), number of alleles (A), allelic richness (A<sub>R</sub>), observed heterozygosity (H<sub>O</sub>), expected heterozygosity (H<sub>E</sub>), and inbreeding coefficient (F<sub>IS</sub>)

\*F<sub>IS</sub> values did not differ significantly from zero after 1000 permutations

## Figure Captions

**Fig. 1** Map of the sampling area. The light grey-shaded region indicates the distribution of *T. h. mogollonensis*, and the dark grey-shaded region represents the Pinaleño Mountains and the restricted range of *T. h. grahamensis*. The open circles represent the two sampling locations for the WMRS. (Map adapted from [http://en.wikipedia.org/wiki/File:USA\\_Arizona\\_location\\_map.svg](http://en.wikipedia.org/wiki/File:USA_Arizona_location_map.svg) under the Creative Commons Attribution-Share Alike 3.0 Unported license)

**Fig. 2** Allele frequency distribution for alleles at eight microsatellite loci. The black bars indicate alleles in MGRS (*T. h. grahamensis*) – shifted mode – and the white bars indicate alleles in WMRS – non-shifted mode

**Fig. 3** Results of the STRUCTURE analysis for red squirrels (*Tamiasciurus hudsonicus*) in the Pinaleño (MGRS: *T. h. grahamensis*) and White (WMRS: *T. h. mogollonensis*) Mountains, Arizona. Each vertical bar represents a different individual, while the color of each vertical bar represents the proportion of ancestry from each of  $K = 2$  populations (indicated by the light grey and dark grey colors)

**Fig. 4** Frequency histogram of the pairwise relatedness values for each population calculated using the method of Lynch and Ritland (1999) for red squirrels (*Tamiasciurus hudsonicus*) in the Pinaleño (MGRS; black line; *T. h. grahamensis*) and White (WMRS; grey-dashed line; *T. h. mogollonensis*) Mountains, Arizona

**Fig. 5** A maximum parsimony network indicating the phylogenetic relationships between the recovered cytb haplotypes for red squirrels (*Tamiasciurus hudsonicus*) in the Pinaleño (MGRS; dark grey; *T. h. grahamensis*) and White (WMRS; light grey; *T. h. mogollonensis*) Mountains, Arizona. Each circle represents a haplotype (A – K), with size proportional to the number of haplotypes sampled. Lines indicate a single mutational step, and small black circles reflect hypothetical haplotypes not recovered in the study

Figure 1  
[Click here to download Figure: Fig1.eps](#)




Figure 2  
[Click here to download Figure: Fig2.eps](#)




Figure 3  
[Click here to download Figure: Fig3.eps](#)




Figure 4  
[Click here to download Figure: Fig4.eps](#)






Figure 5  
[Click here to download Figure: Fig5.eps](#)

