

1-1-2022

Freshwater lamprey and fishes of Iran; a revised and updated annotated checklist-2022

SOHEIL EAGDERI

ATTA MOULUDI-SALEH

HAMID RIZA ESMAELI

GOLNAZ SAYYADZADEH

MANONCHEHR NASRI

Follow this and additional works at: <https://journals.tubitak.gov.tr/zoology>



Part of the [Zooology Commons](#)

Recommended Citation

EAGDERI, SOHEIL; MOULUDI-SALEH, ATTA; ESMAELI, HAMID RIZA; SAYYADZADEH, GOLNAZ; and NASRI, MANONCHEHR (2022) "Freshwater lamprey and fishes of Iran; a revised and updated annotated checklist-2022," *Turkish Journal of Zoology*: Vol. 46: No. 6, Article 6. <https://doi.org/10.55730/1300-0179.3104>

Available at: <https://journals.tubitak.gov.tr/zoology/vol46/iss6/6>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Freshwater lamprey and fishes of Iran; a revised and updated annotated checklist-2022

Soheil EAGDERI^{1*}, Atta MOULUDI-SALEH¹, Hamid Reza ESMAEILI²,
Golnaz SAYYADZADEH², Manoochehr NASRI³

¹Department of Fisheries, Faculty of Natural Resources, University of Tehran, Karaj, Iran

²Ichthyology and Molecular Systematics Laboratory, Department of Biology, College of Sciences, Shiraz University, Shiraz, Iran

³Department of Environmental and Fisheries Sciences and Engineering, Faculty of Agriculture and Natural Resources, Lorestan University, Khorramabad, Iran

Received: 08.06.2022

Accepted/Published Online: 25.10.2022

Final Version: 10.11.2022

Abstract: The current status of the inland waters ichthyofauna of Iran is revised and updated. A total of 292 fish species belong to 3 classes, 24 orders, 36 families and 106 genera inhabit the inland waters of Iran. Among these, 29 species (9.9%) are exotic and 102 species (35%) are endemic. Orders with the largest numbers of species in the ichthyofauna of Iran are the Cypriniformes (182 species) followed by Gobiiformes (30 species), Cyprinodontiformes (15 species), Siluriformes (12 species), Clupeiformes (10 species), Acipenseriformes and Cichliformes (7 species) and Mugiliformes (6 species). At the family level, the Cyprinidae has the greatest number of species (74 species; 25.34% of the total species), followed by the Nemacheilidae (47 species), Leuciscidae (42 species), Gobiidae (30 species), Aphaniidae (11 species), Clupeidae (10 species), Acipenseridae and Cichlidae and Sisoridae (7 species in each) and Cobitidae and Mugilidae (6 species).

Key words: Freshwater ichthyofauna, Iran, endemism, introduced

1. Introduction

Iran possesses a rich freshwater fish fauna in terms of diversity and endemism, and its ichthyofauna is characterized by having elements from Palearctic, Oriental, and Ethiopian ecoregions with exotics from the Nearctic and Neotropical origins (Esmaeili et al., 2010a, b, 2013a, b, 2014a, b, 2017a, 2018). The remarkable Iranian ichthyofauna has attracted the interest of foreign naturalists and scientists for a long time and later awareness regarding Iranian fish taxonomy increased among Iranian researchers at the end of the 20th century. During the last two decades, scientists significantly increased their knowledge of Iranian freshwater fish diversity (Esmaeili et al., 2018).

The first checklist of Iranian freshwater fishes was published by Coad (1995) listing 150 species in 25 families, 14 orders, and 3 classes found in the 19 drainage basins. Later Coad (1998) listed 155 species in 67 genera, 24 families, 15 orders, and 3 classes. Subsequently, Esmaeili et al. (2010a) provided a list of the freshwater fishes of Iran and confirmed the presence of 202 species in 104 genera, 28 families, 17 orders, and 3 classes. They also reported 23 species whose presence in Iranian waters needed confirmation by specimens. Later, Jouladeh-Roudbar et al.

(2015a) listed 257 species in 106 genera, 29 families, 18 orders, and 3 classes. According to them, the most diverse order is the Cypriniformes with 162 species or 63.04% of the fauna, followed by Perciformes (32 species, 12.45%), Cyprinodontiformes (17 species, 6.61%) and Clupeiformes (11 species, 4.28%). Keivany et al. (2016) reported a total of 163 species occurring in the freshwater of Iran, excluding the Caspian Sea, mostly based on their own collections. In another publication, Esmaeili et al. (2017a) listed 288 species in 107 genera, 28 families, 22 orders, and 3 classes reported from different Iranian basins. However, according to them, the presence of 23 species in Iranian waters needs confirmation by specimens. Esmaeili et al. (2018) provided a new list of inland/freshwater fishes of Iran reporting 297 species in 109 genera, 30 families, 24 orders, and 3 classes (including 23 unconfirmed species). In the last published checklist, Jouladeh-Roudbar et al. (2020), a list of 274 recognized species in 100 genera, 33 families, 20 orders, and 3 classes was given.

Although there are quite a number of publications on freshwater fish taxonomy, the data set for freshwater fish assemblages are still poor. In this paper, we present an updated checklist of the freshwater fishes of Iran based on previous checklists (Esmaeili et al., 2018), and new species

* Correspondence: soheil.eagderi@ut.ac.ir

records. Nonetheless, the continuing description of new species of Iran freshwater fishes makes it clear that the present checklist underestimates, perhaps to a significant degree, the actual richness of that freshwater ichthyofauna. The present study aims to represent the currently known diversity of the freshwater fishes of Iran, including only confirmed species.

2. Material and methods

The species in the following list are compiled from two different sources of information. The data were cross-checked and supplemented by information from recent publications dealing with species present in the inland waters of Iran and/or describing new species of fishes from Iran. The previously published checklists are taken as a baseline (Esmaili et al., 2018). In addition, information on fish distribution in FishBase was evaluated. Additional fish records published during the last decade were added, and some questionable species appearing in previous checklists were evaluated and either verified or excluded from the present list. We follow the family classification of Nelson et al. (2016), with orders and families arranged systematically, but genera and species alphabetically within each family.

In the present checklist, we include only those fishes which spend some parts of their life or their entire life history in freshwater systems. Some species are excluded from the checklist because (1) they were erroneously identified, (2) taxa were synonymized, and (3) they were probably not occurring in inland waters of Iran and need confirmation by the specimen. Such questionable species are excluded from the list unless their occurrence can be proven.

3. Results

The present checklist includes 292 species inhabiting freshwater systems of Iran, belonging to 3 classes, 24 orders, 36 families, and 106 genera. Among them, 102 species (35%) are considered as endemic and 29 species (9.9%) are exotic (Figure 1). The dominant order in the fish fauna is Cypriniformes (182 species) followed by Gobiiformes (30 species), Cyprinodontiformes (15 species), Siluriformes (12 species), Clupeiformes (10 species), Acipenseriformes and Cichliformes (7 species in each) and Mugiliformes (6 species) (Figure 2). At the family level, the Cyprinidae has the greatest number of species (74 species; 25.34% of the total species), followed by the Nemacheilidae (47 species; 16.1% of the total species), Leuciscidae (42 species; 14.38% of the total species), Gobiidae (30 species; 10.27% of the total species), Aphaniidae (11 species; 3.77% of the total species), Clupeidae (10 species; 3.42% of the total species), Acipenseridae, Cichlidae and Sisoridae (7 species; 2.4% of the total species in each) and Cobitidae and Mugilidae




Figure 1. Percentage of the total occurrence of endemic, exotic, and native fishes of Iranian inland waters.

(6 species; 2.05% of the total species in each) (Figure 3). The highest number of endemic (36 species) belong to the Tigris basin and exotic (17 species) to Tigris and Caspian Sea basins, and the Caspian Sea basin has the highest number of native species (83 species) (Tables 1, 2, and Figure 4).

Species name (Author) [Occurrence]-[IUCN], English Name/ Distribution (Reference/s).

Kingdom Animalia

Phylum Chordata

Subphylum Craniata

Infraphylum Vertebrata

Superclass Petromyzontomorpha

Class Petromyzontida

Order Petromyzontiformes

Family Petromyzontidae

Genus *Caspiomyzon* Berg, 1906

1. *Caspiomyzon wagneri* (Kessler, 1870) [N]-[NT], Caspian lampreys/Caspian Sea Basin (Eagderi et al., 2017d; Namdarian Rad et al., 2017).

Class Chondrichthyes

Order Carcharhiniformes

Family Carcharhinidae

Genus *Carcharhinus* Blainville, 1816

2. *Carcharhinus leucas* (Valenciennes 1839) [N]-[NT], Bull shark/Tigris, Persian Gulf Basin (Esmaili et al., 2018).

Superclass Gnathostomata

Grade Teleostomi

Class Osteichthyes

Subclass Actinopterygii

Infraclass Chondrostei




Figure 2. Number of fish species in different orders.




Figure 3. Percentage of fish species in different families.

Order Acipenseriformes

Family Acipenseridae

Genus *Acipenser* Linnaeus, 1758

3. *Acipenser baerii* Brandt, 1869 [I]-[EN], Siberian Sturgeon/Caspian Sea Basin (Hasanalipour et al., 2013). This species was imported for aquaculture purposes, but it can be found in some natural habitats e.g., Aras River, which probably escaped from fish farms.

4. *Acipenser gueldenstaedtii* Brandt & Ratzeburg, 1833 [N]-[CR], Russian sturgeon/Caspian Sea Basin (Esmaili et al. 2018).

5. *Acipenser nudiiventris* Lovetsky, 1828 [N]-[CR], Fringebarbel sturgeon/Caspian Sea Basin (Esmaili et al., 2018).

6. *Acipenser persicus* Borodin, 1897 [N]-[CR], Persian sturgeon/Caspian Sea Basin (Eagderi et al., 2017a; Esmaili et al., 2018).

7. *Acipenser ruthenus* Linnaeus, 1758 [N]-[VU], Sterlet/Caspian Sea Basin. Comment: Reported from the middle and South Caspian Sea by Naseka and Bogutskaya (2009) but not confirmed by specimens for Iran. This species is transported to the Shahid Rajaei Fish Aquaculture Center at Sari, Mazandaran, and some other fish farms in Guilan and Golestan provinces for aquaculture purposes.

8. *Acipenser stellatus* Pallas, 1771 [N]-[CR], Stellate sturgeon/Caspian Sea Basin (Esmaili et al., 2018).

Table 1. Occurrence percentage of endemic, exotic and native fish species separately for each basin.

Basins	Total	Native		Endemic		Exotic	
		#	%	#	%	#	%
1: Caspian Sea	110	83	75.5	10	9.1	17	15.5
2: Dasht-e Kavir	18	4	21.1	6	33.33	8	44.4
3: Dasht-e Lut	9	5	55.6	1	11.1	3	33.3
4: Esfahan	14	2	14.3	6	42.9	6	42.9
5: Mashkid	18	11	61.1	2	11.1	5	27.8
6: Jaz Murian	14	6	42.9	2	14.3	6	42.9
7: Kor River	22	4	18.2	11	50.0	7	31.8
8: Lake Maharlu	14	3	21.4	4	28.6	7	50.0
9: Lake Urmia	26	11	42.3	7	26.9	8	30.8
10: Namak Lake	23	3	12.5	9	39.1	11	47.8
11: Sirjan	5	1	20.0	1	20.0	3	60.0
12: Sistan	21	11	52.4	3	14.3	7	33.3
13: Hari River	23	14	60.9	0	0	9	39.1
14: Kerman-Nain	6	1	16.7	2	33.3	3	50.0
15: Makran	24	16	66.7	3	12.5	5	20.8
16: Hormuz	35	16	45.7	15	42.9	4	11.4
17: Persis	41	19	45.2	14	34.1	8	19.5
18: Tigris River	107	55	50.9	36	33.6	17	15.9
19: Zohreh River	11	5	45.5	3	27.3	3	27.3
TOTAL	292	161	55.1	102	35	29	9.9

Genus *Huso* Brandt & Ratzeburg, 1833

9. *Huso huso* (Linnaeus, 1758) [N]-[CR], Beluga/Caspian Sea Basin (Asgari et al., 2014; Esmaeili et al., 2018).

Infraclass Holostei**Division Ginglymodi****Order Lepisosteiformes****Family Lepisosteidae****Genus *Atractosteus* Rafinesque, 1820**

10. *Atractosteus spatula* (Lacepède, 1803) [I]-[NE], Alligator gar/Tigris, Persian Gulf basin (Esmaeili et al. 2017b, 2018).

Remark: It has been reported from the southern reaches of the Arvand River/Shatt al-Arab River at Om-al-Rasas Island, Basrah, Iraq during an ichthyologic survey in September 2016 (Mutlak et al., 2017) and from Marivan (Zarivar) Lake, a Tigris River tributary of Iran (Esmaeili et al., 2017b). The aquarium trade pathway is the suspected factor for the presence of this species in the freshwater area of Iran and Iraq. Monitoring of inland waters is recommended.

Neopterygii**Infraclass Holostei****Division Teleostei****Subdivision Teleostei****Supercohort Teleostei****Cohort Elopomorpha****Order Anguilliformes****Family Anguillidae****Genus *Anguilla* Schrank, 1798**

11. *Anguilla anguilla* (Linnaeus, 1758) [I]-[CR], European eel/Introduced to the Caspian Sea basin (Esmaeili et al., 2018).

Clupeocephala**Cohort Otocephala****Superorder Clupeomorpha****Order Clupeiformes****Family Clupeidae**

Kottelat and Freyhof (2007) listed *Alosa volgensis* (Berg, 1913) from Iranian waters but we cannot confirm the presence of this species in Iran. The distribution range of this species is limited to the northern Caspian Sea. *Alosa curensis* (Suvorov, 1907) is poorly known and described from Kyzylagach Bay of Azerbaijan. We could not confirm the presence of this species in Iran.

Table 2. List of species to be excluded and their exclusion criterion (EQ) from the freshwater checklist of Iran (species are listed in alphabetical order).

Species	EQ	Explanation
<i>Alburnoides coadi</i> Mousavi-Sabet, Vatandoust & Doadrio, 2015	2	Junior synonym of <i>Alburnoides namak</i> (Jouladeh-Roudbar and Eagderi 2017).
<i>Alburnoides idignensis</i> Bogutskaya & Coad, 2009	2	Junior synonym of <i>Alburnoides nicolausi</i> (Jouladeh-Roudbar and Eagderi 2019).
<i>Alburnoides parhami</i> Mousavi-Sabet, Vatandoust & Doadrio, 2015	2	Junior synonym of <i>Alburnoides holciki</i> (Jouladeh-Roudbar and Eagderi 2019).
<i>Alburnus zagrosensis</i> (Coad, 2009)	2	Junior synonym of <i>Alburnus sellal</i> (Eagderi et al., 2019b).
<i>Alosa curensis</i> (Suvorov, 1907)	3	Need confirmation by specimen.
<i>Alosa volgensis</i> (Berg, 1913)	3	Need confirmation by specimen.
<i>Aphanius arakensis</i> Teimori, Esmaeili, Gholami, Zarei & Reichenbacher, 2012	3	Junior synonym of <i>Esmacilius sophiae</i> (Heckel, 1849) (Freyhof and Yoğurtçuoğlu 2020).
<i>Aphanius farsicus</i> Teimori, Esmaeili & Reichenbacher, 2011	3	Junior synonym of <i>Esmacilius persicus</i> (Jenkins 1910) (Freyhof and Yoğurtçuoğlu 2020).
<i>Aphanius arakensis</i> Teimori, Esmaeili, Gholami, Zarei & Reichenbacher, 2012	3	Junior synonym of <i>Esmacilius sophiae</i> (Heckel, 1849) (Freyhof and Yoğurtçuoğlu 2020).
<i>Aphanius kavirensis</i> Esmaeili, Teimori, Gholami & Reichenbacher, 2014	3	Junior synonym of <i>Esmacilius sophiae</i> (Heckel, 1849) (Freyhof and Yoğurtçuoğlu 2020).
<i>Aphanius pluristriatus</i> (Jenkins, 1910)	3	Junior synonym of <i>Esmacilius sophiae</i> (Heckel, 1849) (Freyhof and Yoğurtçuoğlu 2020).
<i>Babka gymnotrachelus</i> (Kessler, 1857)	3	Need confirmation by specimen.
<i>Babka macrophthalma</i> (Kessler, 1877)	3	Need confirmation by specimen.
<i>Benthophiloides brauneri</i> Beling and Iljin, 1927	3	Need confirmation by specimen.
<i>Benthophiloides turcomanus</i> (Iljin, 1941)	3	Need confirmation by specimen.
<i>Benthophilus casachicus</i> Ragimov, 1978	3	Need confirmation by specimen.
<i>Benthophilus grimmi</i> Kessler, 1877	3	Need confirmation by specimen.
<i>Benthophilus kessleri</i> Berg, 1927	3	Need confirmation by specimen.
<i>Benthophilus leptocephalus</i> Kessler, 1877	3	Need confirmation by specimen.
<i>Benthophilus leptorhynchus</i> Kessler, 1877	3	Need confirmation by specimen.
<i>Benthophilus mahmudbejovi</i> Ragimov, 1976	3	Need confirmation by specimen.
<i>Benthophilus ragimovi</i> Boldyrev & Bogutskaya, 2004	3	Need confirmation by specimen.
<i>Benthophilus spinosus</i> Kessler, 1877	3	Need confirmation by the specimen.
<i>Benthophilus svetovidovi</i> Pinchuk & Ragimov, 1979	3	Need confirmation by the specimen.
<i>Capoeta kaput</i> Levin, Prokofiev & Roubenyan, 2019	3	Need confirmation by the specimen (Kuljanishvili et al., 2020).
<i>Capoeta sevangi</i> De Filippi, 1865	2	Junior synonym of <i>C. capoeta</i> (Kuljanishvili et al., 2019).
<i>Garra variabilis</i>	3	Need confirmation by the specimen.
<i>Hyracanogobius bergi</i> Iljin, 1928	3	Need confirmation by the specimen.
<i>Mesogobius nigronotatus</i> (Kessler, 1877)	3	Need confirmation by the specimen.
<i>Oncorhynchus keta</i> (Walbaum, 1792)	3	Need confirmation by the specimen.
<i>Oxynoemacheilus arygyrogramma</i> (Heckel, 1877)	1	Misidentified.
<i>Oxynoemacheilus freyhofi</i> Jouladeh-Roudbar, Eagderi & Hosseinpour, 2016	2	Junior synonym of <i>Oxynoemacheilus euphraticus</i> (Bănărescu & Nalbant 1964) (Freyhof 2016b)
<i>Oxynoemacheilus hanae</i> Freyhof & Abdullah, 2017	3	Need confirmation by specimen.
<i>Oxynoemacheilus lenkoranensis</i> (Abdurakhamanov, 1962)	2	Junior synonym of <i>O. bergianus</i> (Freyhof et al., 2022).
<i>Oxynoemacheilus veyselorum</i> Çiçek, Eagderi & Sungur, 2018	3	Need confirmation by specimen.
<i>Paracobitis longicauda</i> (Kessler, 1872)	3	Need confirmation by specimen.
<i>Paraschistura pasatigris</i> (Coad & Nalbant, 2005)	2	Junior synonym of <i>Paraschistura ilamensis</i> Vatandoust & Eagderi 2015 (Eagderi et al., 2019d).
<i>Paracobitis vignai</i> Nalbant & Bianco, 1998	2	Synonym of <i>Paracobitis rhadinaea</i> (Regan, 1906) (Sayyadzadeh et al., 2019a).
<i>Pimephales promelas</i> Rafinesque, 1820	3	Need confirmation by specimen.
<i>Ponticola platystris</i> (Pallas, 1814)	3	Need confirmation by specimen.
<i>Rutilus rutilus</i> (Linnaeus, 1758)	3	Need confirmation by specimen (Kuljanishvili et al., 2020).
<i>Salvelinus fontinalis</i> (Mitchill, 1814)	3	Need confirmation by specimen.
<i>Sander volgensis</i> (Gmelin, 1789)	3	Need confirmation by specimen.




Figure 4. Occurrence percentage of endemic, exotic, and native fish species separately for each basin.

Genus *Alosa* Linck, 1790

12. *Alosa braschnikowi* (Borodin, 1904) [N]-[NE], Caspian marine shad-Brazhnikov's shad/ Caspian Sea Basin (Esmaili et al., 2018; Saffari et al., 2021).

13. *Alosa caspia* (Eichwald, 1838) [N]-[LC], Caspian shad/Caspian Sea Basin (Esmaili et al., 2018).

14. *Alosa kessleri* (Grimm, 1887) [N]-[LC], Caspian anadromous shad-blackback-black-spined herring/ Caspian Sea Basin (Esmaili et al., 2018).

15. *Alosa saposchnikowii* (Grimm, 1885) [N]-[NE], Saposhnikov shad/Caspian Sea Basin (Esmaili et al., 2018).

16. *Alosa sphaerocephala* (Berg, 1913) [N]-[LC], Agrakhan shad/Caspian Sea Basin (Esmaili et al., 2018).

Genus *Clupeonella* Kessler, 1877

17. *Clupeonella caspia* Svetovidov, 1941 [N]-[LC], Caspian tyulka/Caspian Sea Basin (Esmaili et al., 2018; Amiri et al., 2018).

Remark: *Clupeonella cultriventris* is widespread along the coasts of the Black and Caspian Seas, entering the lower reaches of rivers. Based on unpublished data (see Kuljanishvili et al., 2020) including molecular and morphological studies they failed to distinguish the Caspian population (*C. caspia*) from the Black Sea population (*C. cultriventris*). Hoestlandt (1991) followed by Kottelat and Freyhof (2007) separate these two tyulkas largely based on the length of paired fins, a character that is found to be very much overlapping in the materials examined by Kuljanishvili et al. (2020). Therefore, they treated *C. caspia* as a synonym of *C. cultriventris*. More data using different populations from the southern Caspian and the Black seas is needed to accept this synonymy.

18. *Clupeonella cultriventris* (Nordman, 1840) [N]-[LC], Caspian tyulka/Caspian Sea Basin (Amiri et al., 2017a,b; Esmaili et al., 2018).

19. *Clupeonella engrauliformis* (Borodin, 1904) [N]-[NE], Caspian tyulka/Caspian Sea Basin (Esmaili et al., 2018).

20. *Clupeonella grimmi* Kessler, 1877 [N]-[NE], Southern Caspian sprat, Bigeye kilka/Caspian Sea Basin (Esmaili et al., 2018).

Genus *Tenualosa* Fowler, 1934

21. *Tenualosa ilisha* (Hamilton, 1822) [N]-[LC], Hilsa/Tigris and Persis; possibly Hormuz (Esmaili et al., 2018).

Order Gonorynchiformes

Family Chanidae

Genus *Chanos* Lacepède, 1803

22. *Chanos chanos* (Forsskal, 1775) [N]-[LC], Milkfish/Tigris, Persis, Hormuz and Makran (Esmaili et al. 2018).

Superorder Ostariophysi

Series Otophysi

Subseries Cypriniphysi

Order Cypriniformes

SUBORDER Cyprinoidae

Family Acheilognathidae

Genus *Rhodeus* Agassiz, 1832

23. *Rhodeus caspius* Esmaili, Sayyadzadeh, Japoshvili, Eagderi, Abbasi & Mousavi-Sabet, 2020 [E]-[NE], Caspian bitterling/The Caspian Sea basin, introduced to the Urmia Lake and Tigris basins (Esmaili et al., 2020a).

Family Cyprinidae**Genus Arabibarbus Borkenhagen, 2014**

24. *Arabibarbus grypus* (Heckel, 1843) [N]-[VU], Shabout/Tigris, Persis and Hormuz (Esmaili et al., 2018; Soleimanian et al., 2021).

Genus Bangana Hamilton, 1822

25. *Bangana dero* (Hamilton, 1822) [N]-[LC], Kalabans/Mashkid River basin (Esmaili et al., 2013a).

Genus Barbus Cuvier, 1816

26. *Barbus cyri* De Filippi, 1865 [N]-[NE], Kura barbel/Caspian Sea basin (Jalili et al., 2017; Khaefi et al., 2017).

27. *Barbus karunensis* Khaefi, Esmaili, Geiger & Eagderi, 2017 [E]-[NE], Karun barbel/Karun River drainage (Khaefi et al., 2017; Eagderi et al., 2017).

28. *Barbus lacerta* Heckel, 1843 [N]-[LC], Tigris barbel/Tigris (Persian Gulf basin) (Khaefi et al., 2017; Esmaili et al., 2018).

29. *Barbus miliaris* De Filippi, 1863 [N]-[NE], Namak barbel/Namak Lake and Kavir basins (Khaefi et al., 2017; Esmaili et al., 2018).

30. *Barbus urmianus* Eagderi, Nikmehr, Çiçek, Esmaili, Vatandoust & Mousavi-Sabet, 2019 [E]-[NE], Urmia barbel/Urmia Lake basin (Eagderi et al., 2019a).

Genus Capoeta Valenciennes, 1842

31. *Capoeta aculeata* (Valenciennes, 1844) [N]-[NE], Common large-scale scraper/Namak Lake and Kavir basins (Zareian et al., 2018).

32. *Capoeta alborzensis* Jouladeh-Roudbar, Eagderi, Ghanavi & Doadrio, 2016 [E]-[NE], Alborz large scale scraper /Namak and Kavir basins (Jouladeh-Roudbar et al., 2017a).

33. *Capoeta birunii* Zareian & Esmaili, 2017 [E]-[NE], Esfahan small-scale scraper/Zayandehrud basin (Zareian and Esmaili, 2017).

34. *Capoeta buhsei* Kessler, 1877 [E]-[LC], Namak scraper/Namak Lake basin (Zareian and Esmaili, 2017; Ahmddzadeh et al., 2019).

35. *Capoeta capoeta* (Güldenstädt, 1773) [N]-[LC], Caucasian scraper/Caspian Sea basin (Esmaili et al., 2017a; Zareian et al., 2018).

36. *Capoeta coadi* Alwan, Zareian & Esmaili, 2016 [E]-[NE], Karun scraper, Coad's scraper/Karun River drainage (Tigris, Persian Gulf basin) (Alwan et al., 2016).

37. *Capoeta damascina* (Valenciennes, 1842) [N]-[LC], Mesopotamian barb/Tigris (Persian Gulf basin) (Razavi Pour et al., 2014).

38. *Capoeta ferdowsii* Jouladeh-Roudbar, Eagderi, Murillo-Ramos, Ghanavi & Doadrio, 2017 [E]-[NE], Zohreh scraper, Ferdowsi scraper/Zohreh River drainage (Persian Gulf basin) (Jouladeh-Roudbar et al., 2017a).

39. *Capoeta fusca* Nikol'skii, 1897 [N]-[NE], Desert scraper/Hari River, Kavir, Bedjestan, Tedzhen River, Sistan and Lut basins (Zareian et al., 2018).

40. *Capoeta gracilis* (Keyserling, 1861) [E]-[NE], Esfahan large scale scraper/Zayandehrud (Esfahan) basin (Zareian et al., 2018).

41. *Capoeta heratensis* (Keyserling, 1861) [N]-[NE], Hari scraper/Hari-rud River basin (Esmaili et al., 2018; Zareian et al., 2018).

42. *Capoeta macrolepis* (Heckel, 1847) [E]-[NE], Kor scraper/Kor and Tigris (Esmaili et al., 2018; Zareian et al., 2018).

43. *Capoeta pyragyi* Jouladeh-Roudbar, Eagderi, Murillo-Ramos, Ghanavi & Doadrio, 2017 [E]-[NE], Sezar scraper/Tigris River drainage (Persian Gulf basin) (Jouladeh-Roudbar et al., 2017a).

44. *Capoeta raghazensis* Eagderi & Mousavi-Sabet, 2021 [E]-[NE], Raghaz scraper/Raghaz Canyon (Hormuz basin) (Eagderi and Mousavi-Sabet, 2021).

45. *Capoeta razii* Jouladeh-Roudbar, Eagderi, Ghanavi & Doadrio, 2017 [E]-[NE], Caspian scraper/Caspian Sea and Kavir basins (Jouladeh-Roudbar et al., 2017b).

46. *Capoeta saadii* (Heckel, 1847) [E]-[NE], Saadi scraper/Kor, Persis, Hormuz, Kerman and Maharlou Lake (Zareian and Esmaili, 2017).

47. *Capoeta shajariani* Jouladeh-Roudbar, Eagderi, Murillo-Ramos, Ghanavi & Doadrio, 2017 [E]-[NE], Shajarian scraper/Tigris River (Persian Gulf basin) (Jouladeh-Roudbar et al., 2017a).

48. *Capoeta umbla* (Heckel, 1843) [N]-[LC], Tigris scraper/Tigris (Persian Gulf basin) (Esmaili et al., 2016b).

Genus Carasobarbus Karaman, 1971

49. *Carasobarbus kosswigi* (Ladiges, 1960) [N]-[VU], Kiss-lip himri/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

50. *Carasobarbus luteus* (Heckel, 1843) [N]-[LC], Mesopotamian himri/Tigris, Persis, Hormuz, Kor and Maharlou lake (Esmaili et al., 2017a, 2018).

51. *Carasobarbus sublimus* (Coad & Najafpour, 1997) [N]-[NE], Persian himri/Tigris and Zohreh (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

Genus Carassius Jarocki, 1822

52. *Carassius auratus* (Linnaeus, 1758) [I]-[LC], Gold fish/widespread in many parts of Iran (Esmaili et al., 2017a, 2018).

53. *Carassius gibelio* (Bloch, 1782) [I]-[NE], Gibel carp/widespread in many parts of Iran (Esmaili et al., 2017a, 2018; Eagderi et al., 2020a).

54. *Carassius langsdorfii* Temminck & Schlegel, 1846 [I]-[NE], Ginbuna/Lar National Park (Caspian Sea Basin) (Khosravi et al., 2022).

Genus Cyprinion Heckel, 1843

55. *Cyprinion kais* Heckel, 1843 [N]-[LC], Smallmouth lotak/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

56. *Cyprinion macrostomum* Heckel, 1843 [N]-[LC], Largemouth lotak/Tigris and probably Persis (Persian Gulf basin) (Nasri et al., 2013; Esmaili et al., 2017a, 2018).

57. *Cyprinion microphthalmum* (Day, 1880) [N]-[LC], Smalleye lotak/different rivers and Qanats in southeastern Iran (Nasri et al., 2018).

58. *Cyprinion milesi* (Day, 1880) [N]-[NE], Eastern lotak/Makran basin (Sarbaz River) (Nasri et al., 2016).

59. *Cyprinion tenuiradius* Heckel, 1847 [E]-[NE], Qarah Aqaj lotak/Persis (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

60. *Cyprinion watsoni* (Day, 1872) [N]-[LC], Indus lotak/might be limited to Pakistan waters. Need further study. (Nasri et al., 2018).

Genus *Cyprinus* Linnaeus, 1758

61. *Cyprinus carpio* Linnaeus, 1758 [N]-[VU], Common carp/native populations in the Caspian Sea basin; also introduced there and elsewhere in Iran (Esmaili et al., 2017a, 2018).

Genus *Garra* Hamilton, 1822

62. *Garra amirhosseini* Esmaili, Sayyadzadeh, Coad & Eagderi, 2016 [E]-[NE], Hot spring garra, Amirhossein's garra/Tigris (Esmaili et al., 2016c).

63. *Garra elegans* (Günther, 1868) [N]-[LC], Elegant garra/-the lower Little Zab River, the Sirvan River, and the lower Tigris (Freyhof, 2016a).

64. *Garra gymnothorax* Berg, 1949 [E]-[NE], Chest scaleless garra/Tigris (Persian Gulf basin) (Esmaili et al., 2016c).

65. *Garra hormuzensis* Zamani-Faradonbe, Zhang & Keivany, 2021 [E]-[NE], Hormuz garra/Kol River drainage (Zamani-Faradonbe et al., 2021a).

66. *Garra lorestanensis* Mousavi-Sabet & Eagderi, 2016 [E]-[NE], Blind cave garra/ subterranean waters in the Tigris drainage (Persian Gulf basin) (Mousavi-Sabet and Eagderi, 2016a).

67. *Garra meymehensis* Zamani-Faradonbe, Keivany, Dorafshan & Zhang, 2021 [E]-[NE], Meymeh garra/Meymeh River (Tigris basin) (Zamani-Faradonbe et al., 2021b).

68. *Garra mondica* Sayyadzadeh, Esmaili & Freyhof, 2015 [E]-[NE], Mond garra/Mond River drainage, Persis (Persian Gulf basin) (Sayyadzadeh et al., 2015a).

69. *Garra nudiventris* (Berg, 1905) [E]-[NE], Lut garra/Kalat-e-Baba Qanat, Lut basin (Esmaili et al., 2016c).

70. *Garra persica* Berg, 1914 [E]-[NE], Persian garra/Hormuz, Makran and Hamun-e Jaz Murian basins (Esmaili et al., 2016d).

71. *Garra roseae* Mousavi-Sabet, Saemi-Komsari, Doadrio & Eagderi, 2019 [E]-[NE], Rose's garra/Tang-e-Sarhe stream, Makran basin (Mousavi-Sabet et al., 2019a).

72. *Garra rossica* (Nikol'skii, 1900) [N]-[NE], Hari garra/Hari, Bedjestan, Sistan, Lut, Hamun-e Jaz Murian, Mashkid and Makran basin (Esmaili et al., 2016d).

73. *Garra rufa* (Heckel, 1843) [N]-[LC], Red garra/Tigris and Persis (Persian Gulf basin) and Maharlu Lake basin (Esmaili et al., 2016c).

74. *Garra tashanensis* Mousavi-Sabet, Vatandoust, Fatemi & Eagderi, 2016 [E]-[NE], Tashan blind cave garra/ subterranean waters in the Tigris drainage (Persian Gulf basin) (Mousavi-Sabet et al., 2016a).

75. *Garra tiam* Zamani-Faradonbe, Keivany, Dorafshan & Zhang, 2021 [E]-[NE], Tiam garra/Karun River drainage (Tigris basin) (Zamani-Faradonbe et al., 2021b).

76. *Garra typhlops* (Bruun & Kaiser, 1944) [E]-[VU], Discless blind cave garra/subterranean waters in the Tigris drainage (Persian Gulf basin) (Esmaili et al., 2016c).

Genus *Labeo* Cuvier, 1816

77. *Labeo rohita* (Hamilton, 1822) [I]-[LC], Roho labeo/Tigris drainage (Persian Gulf basin) (Eagderi et al., 2019c).

Genus *Luciobarbus* Heckel, 1843

78. *Luciobarbus barbulus* (Heckel, 1847) [N]-[NE], Qarah Aqaj barbel/Helleh, Mond, Zohreh and Tigris (Persian Gulf basin) (Khaefi et al., 2017b).

79. *Luciobarbus brachycephalus* (Kessler, 1872) [N]-[VU], Aral barbel/Caspian Sea basin (Esmaili et al., 2017a, 2018).

80. *Luciobarbus capito* (Güldenstädt, 1773) [N]-[VU], Bulatamai barbel/Caspian Sea basin (Eagderi et al., 2013; Esmaili et al., 2018).

81. *Luciobarbus conocephalus* (Kessler, 1872) [N]-[NE], Hari barbel/Hari basin (Eagderi et al., 2021).

82. *Luciobarbus esocinus* Heckel, 1843 [N]-[VU], Pike barbel/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

83. *Luciobarbus kersin* (Heckel, 1843) [N]-[DD], Kersin barbel/Tigris (Persian Gulf basin) (Khaefi et al., 2018).

84. *Luciobarbus mursa* (Güldenstädt, 1773) [N]-[LC], Mursa/Caspian Sea and Urmia Lake basins (Esmaili et al., 2017a, 2018).

85. *Luciobarbus subquincunciatus* (Günther, 1868) [N]-[CR], Leopard barbel/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

86. *Luciobarbus xanthopterus* Heckel, 1843 [N]-[VU], Gattan/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

Genus *Mesopotamichthys* Karaman, 1971

87. *Mesopotamichthys sharpeyi* (Günther, 1874) [N]-[VU], Binni/Tigris and Persis (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

Genus *Paracapoeta* Turan, Kaya, Aksu & Bektaş, 2022

88. *Paracapoeta trutta* (Heckel, 1843) [N]-[LC], Longspine scraper /Tigris and Zohreh (Persian Gulf basin) (Esmaili et al., 2018).

89. *Paracapoeta anamisensis* (Zareian, Esmaili & Freyhof, 2016) [E]-[NE], Minab scraper/Makran (Zareian et al., 2016).

90. *Paracapoeta mandica* (Bianco & Bănărescu, 1982) [E]-[NE], Mond scraper/Persis basin (Zareian et al., 2016).

Genus *Schizocypris* Regan, 1914

91. *Schizocypris altidorsalis* Bianco & Bănărescu, 1982 [N]-[LC], Gorgak/Sistan basin (Esmaili et al., 2017a, 2018).

Genus *Schizopygopsis* Steindachner, 1866

92. *Schizopygopsis stolickai* Steindachner, 1866 [N]-[NE], False Osman/Sistan basin (Esmaili et al., 2017a, 2018).

Genus *Schizothorax* Heckel, 1838

93. *Schizothorax intermedius* McClelland & Griffith, 1842 [N]-[NE], Common marinka/Sistan basin (Esmaili et al., 2017a, 2018).

94. *Schizothorax pelzami* Kessler, 1870 [N]-[LC], Transcasian Marinka/Hari River and Kavir basins (Esmaili et al., 2018; Mouludi-Saleh et al., 2020a).

95. *Schizothorax zarudnyi* (Nikol'skii, 1897) [N]-[NE], Sistan Marinka/Sistan basin (Esmaili et al., 2017a, 2018).

Genus *Tariqilabeo* Mirza & Saboohi, 1990

96. *Tariqilabeo adiscus* (Annandale, 1919) [N]-[NE], Sistan Latia/Sistan basin (Sayyadzadeh et al., 2015b).

97. *Tariqilabeo diplochilus* (Heckel, 1838) [N]-[NE], Kashmir Latia/Mashkid and Makran basins (Sayyadzadeh et al., 2015b).

Family Danionidae

Genus *Barilius* Hamilton, 1822

98. *Barilius mesopotamicus* Berg, 1932 [N]-[LC], Mesopotamian minnow/Tigris and Persis (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

Genus *Cabdio* Hamilton, 1822

99. *Cabdio morar* (Hamilton, 1822) [N]-[LC], Morar/Makran and Mashkid basin (Esmaili et al., 2015; Radkhah et al., 2022a).

Family Gobionidae

Genus *Gobio* Cuvier, 1816

100. *Gobio nigrescens* (Keyserling, 1861) [N]-[NE], Hari gudgeon/Hari basin (Mousavi-Sabet et al., 2016b).

Genus *Pseudorasbora* Bleeker, 1859

101. *Pseudorasbora parva* (Temminck & Schlegel, 1846) [I]-[LC], Topmouth gudgeon/Introduced to the Caspian Sea, Namak Lake, Hari River, Sistan, Maharlu, Urmia, Persis and Tigris River drainage and probably elsewhere (Ganjali et al., 2021).

Genus *Romanogobio* Bănărescu, 1961

102. *Romanogobio macropterus* (Kamensky, 1901) [N]-[LC], South Caucasian gudgeon/Caspian Sea basin (Esmaili et al., 2017a, 2018).

103. *Romanogobio persus* (Günther, 1899) [E]-[NE], Persian gudgeon/Urmia Lake basin (Esmaili et al., 2017a, 2018).

Family Leuciscidae

Genus *Abramis* Cuvier, 1816

104. *Abramis brama* (Linnaeus, 1758) [N]-[LC], Common bream/the Caspian Sea basin and introduced to the Urmia Lake basin (Sahraeian et al., 2016).

Genus *Acanthobrama* Heckel, 1843

105. *Acanthobrama marmid* Heckel, 1843 [N]-[LC], Mesopotamian bream/Tigris (Persian Gulf basin) (Esmaili et al., 2018; Abbasi Ranjbar et al., 2018).

106. *Acanthobrama microlepis* (De Filippi, 1863) [N]-[LC], Blackbrow bleak/Caspian Sea basin (Esmaili et al., 2018; Abbasi Ranjbar et al., 2018).

107. *Acanthobrama persidis* (Coad, 1981) [E]-[NE], Persian bleak, Kor bleak/Kor, Persis, Maharlu Lake, and Hormuz basins (Teimori et al., 2015).

108. *Acanthobrama urmianus* (Günther, 1899) [E]-[DD], Urmia bream/Urmia Lake basin (Esmaili et al., 2018; Abbasi Ranjbar et al., 2018).

Genus *Alburnoides* Jettles, 1861

109. *Alburnoides damghani* Jouladeh-Roudbar, Eagderi, Esmaili, Coad & Bogutskaya, 2016 [E]-[NE], Damghan spiralin, Damghan riffle minnow/Kavir basin (Jouladeh-Roudbar et al., 2016).

110. *Alburnoides eichwaldii* (De Filippi, 1863) [N]-[LC], South western Caspian spiralin, Eichward's riffle minnow/Caspian Sea basin (Esmaili et al., 2017a, 2018).

111. *Alburnoides holciki* Coad & Bogutskaya, 2012 [N]-[NE], Hari spiralin, Holcik's riffle minnow/Hari and the Caspian Sea/Atrak River basins (Jouladeh-Roudbar et al., 2020).

112. *Alburnoides namaki* Bogutskaya & Coad, 2009 [E]-[NE], Namak spiralin, Namak riffle minnow/Namak Lake basin (Esmaili et al., 2017a, 2018).

113. *Alburnoides nicolausi* Bogutskaya & Coad, 2009 [E]-[NE], Seimareh spiralin, Karkheh spiralin, Nicholas' riffle minnow/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

114. *Alburnoides petrubanarescui* Bogutskaya & Coad, 2009 [E]-[NE], Urmia spiralin, Banarescu's riffle minnow/Urmia Lake basin (Esmaili et al., 2017a, 2018).

115. *Alburnoides qanati* Coad & Bogutskaya, 2009 [E]-[NE], Kor spiralin, Qanat spiralin/Kor and Sirjan basins (Esmaili et al., 2017a, 2018).

116. *Alburnoides samiii* Mousavi-Sabet, Vatandoust & Doadrio, 2015 [E]-[NE], Sefidrud spiralin, Samii riffle minnow/Caspian Sea basin (Mousavi-Sabet et al., 2015a).

117. *Alburnoides tabarestanensis* Mousavi-Sabet, Anvarifar & Azizi, 2015 [E]-[NE], Tajan spiralin/Caspian Sea basin (Mousavi-Sabet et al., 2015b).

Genus *Alburnus* Rafinesque, 1820

118. *Alburnus atropatenae* Berg, 1925 [E]-[NE], Urmia bleak, Urmia shemaya/Urmia Lake basin (Esmaili et al., 2017a, 2018; Mouludi-Saleh et al., 2022a).

119. *Alburnus caeruleus* Heckel, 1843 [N]-[LC], Black spotted bleak/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

120. *Alburnus chalcoides* (Güldenstadt, 1772) [N]-[LC], Caspian shemaya/Caspian Sea basin (Esmaili et al., 2017a, 2018).

121. *Alburnus doriae* De Filippi, 1865 [E]-[NE], Doria bleak/Esfahan, Namak, Tigris (Persian Gulf) basins (Mohammadian-Kalat et al., 2017).

122. *Alburnus filippii* Kessler, 1877 [N]-[LC], Kura bleak/Caspian Sea basin (Esmaili et al., 2017a, 2018).

123. *Alburnus hohenackeri* Kessler, 1877 [N]-[LC], Transcaucasian bleak/native in the Caspian Sea basin and translocated to other basins in Iran (Zareian et al., 2013).

124. *Alburnus sellal* Heckel, 1843 [N]-[LC], Mesopotamian bleak/Tigris, Zohreh, Persis and Hormuz basins (Mohammadian-Kalat et al., 2017; Eagderi et al., 2019b).

125. *Alburnus taeniatus* Kessler, 1874 [N]-[NE], Striped bystranka/Hari basin (Jouladeh-Roudbar et al., 2016).

Genus *Ballerus* Heckel, 1843

126. *Ballerus sapa* (Pallas, 1814) [N]-[LC], White-eye bream/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Genus *Blicca* Heckel, 1843

127. *Blicca bjoerkna* (Linnaeus, 1758) [N]-[LC], Silver bream/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Genus *Chondrostoma* Agassiz, 1832

128. *Chondrostoma cyri* Kessler, 1877 [N]-[LC], Southern Caspian nase/Caspian Sea basin (Esmaili et al., 2017a, 2018).

129. *Chondrostoma esmaeilii* Eagderi, Jouladeh-Roudbar, Birecikligil, Çiçek & Coad, 2017 [E]-[NE], Tigris nase/Tigris (Persian Gulf basin) (Eagderi et al., 2017b).

130. *Chondrostoma orientale* Bianco & Bănărescu, 1982 [E]-[NE], Kor nase/Kor basin (Esmaili et al., 2017a, 2018).

131. *Chondrostoma regium* (Heckel, 1843) [N]-[LC], Mesopotamian nase/Tigris, Zohreh (Persian Gulf), and Esfahan basins (Dastanpoor et al., 2021).

Genus *Leucaspis* Heckel & Kner, 1858

132. *Leucaspis delineatus* (Heckel, 1843) [N]-[LC], Moderlieschen/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Genus *Leuciscus* Cuvier, 1816

133. *Leuciscus aspius* (Linnaeus, 1758) [N]-[LC], Asp, European asp/Caspian Sea basin (Esmaili et al., 2017, 2018; Abbasi et al., 2021a).

134. *Leuciscus vorax* (Heckel, 1843) [N]-[LC], Mesopotamian asp/Tigris (Persian Gulf basin) (Esmaili et al., 2017, 2018).

135. *Leuciscus latus* (Keyserling, 1861) [N]-[DD], Transcasian dace/Transcasian chub/ Hari basin (Eagderi et al., 2021b)

Genus *Pelecus* Agassiz, 1835

136. *Pelecus cultratus* (Linnaeus, 1758) [N]-[LC], Ziege/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Genus *Petroleuciscus* Bogutskaya, 2002

137. *Petroleuciscus ulanus* (Günther, 1899) [E]-[NE], Urmia chub/Urmia Lake basin (Esmaili et al., 2017a, 2018; Abbasi et al., 2021; Mouludi-Saleh et al., 2022).

Genus *Rutilus* Rafinesque, 1820

138. *Rutilus frisii* (Nordmann, 1840) – [N]-[NE], Kutum/Caspian Sea basin (Kuljanishvili et al., 2020).

139. *Rutilus lacustris* (Pallas, 1814) [N]-[NE], Roach, Vobla/Caspian Sea basin (Levin et al., 2016; Kuljanishvili et al., 2020).

Genus *Scardinius* Bonaparte, 1837

140. *Scardinius erythrophthalmus* (Linnaeus, 1758) [N]-[LC], Rudd, redeye, redfin, pearl roach/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Genus *Squalius* Bonaparte, 1837

141. *Squalius berak* Heckel, 1843 [N]-[LC], Mesopotamian chub/Tigris (Esmaili et al., 2016a; Khaefi et al., 2016).

142. *Squalius lepidus* Heckel, 1843 [N]-[LC], Mesopotamian pike chub/Tigris (Esmaili et al. 2017a, 2018).

143. *Squalius namak* Khaefi, Esmaili, Sayyadzadeh, Geiger & Freyhof, 2016 [E]-[NE], Namak Lake chub/ Namak Lake basin (Khaefi et al., 2016).

144. *Squalius turcicus* De Filippi, 1865 [N]-[LC], Transcaucasian chub/Urmia Lake and southern Caspian Sea basins (Mouludi-Saleh et al., 2020b).

Genus *Vimba* Fitzinger, 1873

145. *Vimba persa* (Pallas, 1814) [N]-[LC], Persian vimba, Caspian vimba/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Family Tincidae

Genus *Tinca* Cuvier, 1816

146. *Tinca tinca* (Linnaeus, 1758) [N]-[LC], Tench/ Caspian Sea basin (Esmaili et al., 2017a, 2018).

Family Xenocyprinidae

Genus *Ctenopharyngodon* Steindachner, 1866

147. *Ctenopharyngodon idella* (Valenciennes, 1844) [I]-[NE], Grass carp/Introduced to the Caspian Sea, Tigris River, Kor River, Maharlou Lake, and Sistan basins;

elsewhere in reservoirs throughout Iran (Esmaili et al., 2017a, 2018).

Genus *Hemiculter* Bleeker, 1859

148. *Hemiculter leucisculus* (Basilevsky, 1855) [I]-[LC], Sharpbelly/introduced to the Caspian Sea basin; probably elsewhere in Iran including Urmia Lake and Tigris River basins (Radkhah & Eagderi, 2021).

Genus *Hypophthalmichthys* Bleeker, 1859

149. *Hypophthalmichthys molitrix* (Valenciennes, 1844) [I]-[NT], Silver carp/introduced to Caspian reservoirs and throughout Iran (Esmaili et al., 2017a, 2018).

150. *Hypophthalmichthys nobilis* (Richardson, 1845) [I]-[DD], Bighead carp/introduced to Caspian reservoirs and throughout Iran (Esmaili et al., 2017a, 2018).

Genus *Mylopharyngodon* Peters, 1881

151. *Mylopharyngodon piceus* (Richardson, 1846) [I]-[DD], Black carp/introduced to the Caspian Sea basin (Esmaili et al., 2017a, 2018).

Superfamily Cobitoidea

Family Cobitidae

Genus *Cobitis* Linnaeus, 1758

152. *Cobitis avicennae* Mousavi-Sabet, Vatandoust, Esmaili, Geiger & Freyhof, 2015 [E]-[NE], Avicenna spined loach/Tigris (Persian Gulf basin) (Mousavi-Sabet et al., 2015c). Its occurrence has been confirmed in the Sirvan River by Mouludi-Saleh et al. (2022b).

153. *Cobitis faridpaki* Mousavi-Sabet, Vasil'eva, Vatandoust & Vasil'ev, 2011 [E]-[NE], Faridpak's spine loach or Siahrud spined loach/Caspian Sea basin (Mousavi-Sabet et al., 2011).

154. *Cobitis linea* (Heckel, 1847) [E]-[NE], Persepolis or Kor spined loach/Kor basin (Esmaili et al., 2017a, 2018).

155. *Cobitis saniae* Eagderi, Jouladeh-Roudbar, Jalili, Sayyadzadeh & Esmaili, 2017 [N]-[NE], Sania's spined loach/Caspian Sea basin (Eagderi et al., 2017c).

Genus *Sabanejewia* Vladykov, 1929

156. *Sabanejewia aurata* (De Filippi, 1863) [N]-[LC], Golden spined loach/Caspian Sea basin (Sayyadzadeh et al., 2018a; Çiçek et al., 2022).

157. *Sabanejewia caspia* (Eichwald, 1838) [N]-[NE], Caspian spined loach/Caspian Sea basin (Sayyadzadeh et al., 2018a).

Family Nemacheilidae

Genus *Eidinemacheilus* Hashemzadeh Segherloo, Ghaedrahmati & Freyhof, 2016

158. *Eidinemacheilus smithi* (Greenwood, 1976) [E]-[VU], Zagros blind crested loach/subterranean waters in the Tigris drainage (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

Genus *Oxynoemacheilus* Bănăraescu & Nalbant, 1967

159. *Oxynoemacheilus bergi* (Gratzianov, 1907) [N]-[LC], Berg Loach/Caspian Sea basin (Mohammadi et al., 2018).

160. *Oxynoemacheilus bergianus* (Derjavin, 1934) [N]-[LC], Safid River stone loach/Caspian Sea, Urmia and Namak lake basins (Mouludi-Saleh and Eagderi, 2021a).

161. *Oxynoemacheilus brandtii* (Kessler, 1877) [N]-[LC], Kura loach/Caspian Sea and Urmia lake basins (Sayyadzadeh et al., 2017).

162. *Oxynoemacheilus chomanicus* Kamangar, Prokofiev, Ghaderi & Nalbant, 2014 [N]-[NE], Choman stone loach/Tigris (Persian Gulf basin) (Kamangar et al., 2014).

163. *Oxynoemacheilus elsae* Eagderi, Jalili & Çiçek, 2018 [N]-[NE], Elsa stone loach/ Urmia Lake basin (Eagderi et al., 2018a).

164. *Oxynoemacheilus euphraticus* (Banareescu & Nalbant, 1964) [N]-[NE], Euphrates stone loach/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018). Its occurrence has been confirmed in the Sirvan River by Mouludi-Saleh et al. (2022b).

165. *Oxynoemacheilus frenatus* (Heckel, 1843) [N]-[LC], Banded Tigris loach/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

166. *Oxynoemacheilus karunensis* Freyhof, 2016 [E]-[NE], Karun stone loach/Tigris (Persian Gulf basin) (Freyhof, 2016b; Sayyadzadeh & Esmaili, 2020).

167. *Oxynoemacheilus kiabii* Goltzarianpour, Abdoli & Freyhof, 2011 [E]-[NE], Kiabi stone loach/Tigris (Persian Gulf basin) (Goltzarianpour et al., 2011).

168. *Oxynoemacheilus kurdistanicus* Kamangar, Prokofiev, Ghaderi & Nalbant, 2014 [N]-[NE], Kurdistan stone loach/Tigris (Persian Gulf basin) (Kamangar et al., 2014).

169. *Oxynoemacheilus longipinnis* (Coad & Nalbant, 2005) [E]-[NE], Ilam stone loach/Tigris (Persian Gulf basin) (Sayyadzadeh et al., 2017).

170. *Oxynoemacheilus marunensis* Sayyadzadeh & Esmaili, 2020 [E]-[NE], Marun stone loach/Tigris (Persian Gulf basin) (Sayyadzadeh & Esmaili, 2020).

171. *Oxynoemacheilus parvinae* Sayyadzadeh, Eagderi & Esmaili, 2016 [E]-[NE], Parvin stone loach/Tigris (Persian Gulf basin) (Sayyadzadeh et al., 2016).

172. *Oxynoemacheilus persa* (Heckel, 1847) [E]-[NE], Persian stone loach/Kor River, Mond River and Maharlu Lake basins (Sayyadzadeh et al., 2018b).

173. *Oxynoemacheilus tongiorgii* (Nalbant & Bianco, 1998) [E]-[DD], Tongiorgi stone loach//Kor basin (Esmaili et al., 2017a, 2018).

174. *Oxynoemacheilus zagrosensis* Kamangar, Prokofiev, Ghaderi & Nalbant, 2014 [N]-[NE], Zagros

stone loach/Tigris (Persian Gulf basin) (Kamangar et al., 2014).

175. *Oxynoemacheilus zarzianus* Freyhof & Geiger, 2017 [N]-[NE], Zarzian stone loach/Sirvan River (Tigris) (Eagderi et al., 2022).

Genus *Paracobitis* Bleeker, 1863

176. *Paracobitis abrishamchianorum* Mousavi-Sabet, Vatandoust, Geiger & Freyhof, 2019 [E]-[NE], Abrishamchi crested loach/Caspian Sea basin (Mousavi-Sabet et al., 2019).

177. *Paracobitis atrakensis* Esmaeili, Mousavi-Sabet, Sayyadzadeh, Vatandoust & Freyhof, 2014 [E]-[NE], Atrak crested loach/Caspian Sea and Kavir basins (Esmaeili et al., 2014d).

178. *Paracobitis basharensis* Freyhof, Esmaeili, Sayyadzadeh & Geiger, 2014 [E]-[NE], Bashar crested loach/Tigris (Persian Gulf basin) (Freyhof et al., 2014).

179. *Paracobitis hircanica* Mousavi-Sabet, Sayyadzadeh, Esmaeili, Eagderi, Patimar & Freyhof, 2015 [E]-[NE], Eastern crested loach/Caspian Sea basin (Mousavi-Sabet et al., 2015d).

180. *Paracobitis malapterura* (Valenciennes, 1846) [E]-[NE], Namak Lake crested loach/Namak Lake and Kavir basins (Freyhof et al., 2014).

181. *Paracobitis molavii* Freyhof, Esmaeili, Sayyadzadeh & Geiger, 2014 [N]-[NE], Molavi's crested loach/Tigris (Persian Gulf basin) (Freyhof et al., 2014).

182. *Paracobitis persa* Freyhof, Esmaeili, Sayyadzadeh & Geiger, 2014 [E]-[NE], Persian crested loach/Kor basin (Freyhof et al., 2014).

183. *Paracobitis rhadinaea* (Regan, 1906) [N]-[NE], Sistan crested loach/Sistan basin (Sayyadzadeh et al., 2019a).

Genus *Paraschistura* Prokofiev, 2009

184. *Paraschistura abdoli* Freyhof, Sayyadzadeh, Esmaeili & Geiger, 2015 [E]-[NE], Abdoli's loach/Kol, Hamun-e Jaz Murian and Sirjan drainage basins (Freyhof et al., 2015).

185. *Paraschistura alta* (Nalbant & Bianco, 1998) [N]-[NE], Helmand loach/Sistan basin (Jouladeh-Roudbar et al., 2015b).

186. *Paraschistura aredvii* Freyhof, Sayyadzadeh, Esmaeili & Geiger, 2015 [E]-[NE], Anahita loach/Zohreh River drainage (Persian Gulf basin) (Freyhof et al., 2015).

187. *Paraschistura bampurensis* (Nikol'skii, 1900) [N]-[NE], Bampur loach/Mashkid, Makran and Hamun-e Jaz Murian basins (Freyhof et al., 2015).

188. *Paraschistura cristata* (Berg, 1898) [N]-[NE], Turkmenian crested loach/Hari basin (Freyhof et al., 2015).

189. *Paraschistura delvarii* Mousavi-Sabet & Eagderi, 2015 [E]-[NE], Delvari's loach/Mond River (Persis) (Persian Gulf basin) (Mousavi-Sabet and Eagderi, 2015).

190. *Paraschistura hormuzensis* Freyhof, Sayyadzadeh, Esmaeili & Geiger, 2015 [E]-[NE], Hormuz loach/Makran basin (Freyhof et al., 2015).

191. *Paraschistura ilamensis* Vatandoust & Eagderi, 2015 [E]-[NE], Ilam loach/Tigris (Persian Gulf basin) (Vatandoust and Eagderi, 2015).

192. *Paraschistura kermanensis* Sayyadzadeh, Teimori & Esmaeili, 2019 [E]-[NE], Kerman loach/Kerman-Naein basin (Sayyadzadeh et al., 2019b).

193. *Paraschistura kessleri* (Günther, 1889) [N]-[NE], Kessler loach/Sistan and Mashkid basins (Freyhof et al., 2015).

194. *Paraschistura makranensis* Eagderi, Mousavi-Sabet & Freyhof, 2019 [E]-[NE], Makran loach/Makran basin (Eagderi et al. 2019d).

195. *Paraschistura naumanni* Freyhof, Sayyadzadeh, Esmaeili & Geiger, 2015 [E]-[NE], Naumann loach/Maharlu Lake, Persis and Hormuz basins (Freyhof et al., 2015).

196. *Paraschistura nielsenii* (Nalbant & Bianco, 1998) [E]-[NE], Nielsen's loach/Helleh and Mond River drainages (Persis) (Freyhof et al., 2015).

197. *Paraschistura susiani* Freyhof, Sayyadzadeh, Esmaeili & Geiger, 2015 [E]-[NE], Susian loach, Susa loach/Tigris (Persian Gulf basin) (Freyhof et al., 2015).

198. *Paraschistura turcmenica* (Berg, 1932) [N]-[NE], Turkmen Loach/Bedjestan, Hari River and Kavir basins (Freyhof et al., 2015).

199. *Paraschistura turcomana* (Nikolskii, 1947) [N]-[NE], Turkmen Loach/Hari basin (Mousavi Sabet et al., 2015e).

Genus *Sasanidus* Freyhof, Geiger, Golzarianpour & Patimar, 2016

200. *Sasanidus kermanshahensis* (Bănărescu & Nalbant, 1966) [E]-[NE], Kermanshah stone loach/Tigris (Persian Gulf basin) (Freyhof et al., 2016).

Genus *Turcinoemacheilus* Bănărescu & Nalbant, 1964

201. *Turcinoemacheilus bahaii* Esmaeili, Sayyadzadeh, Özuluğ, Geiger & Freyhof, 2014 [E]-[NE], Bahaii dwarf loach/Zayandeh River (Esfahan) basin (Esmaeili et al., 2014b).

202. *Turcinoemacheilus hafezi* Golzarianpour, Abdoli, Patimar & Freyhof, 2013 [E]-[NE], Hafez dwarf loach/Tigris (Persian Gulf basin) (Esmaeili et al., 2014b).

203. *Turcinoemacheilus kosswigi* Banarescu & Nalbant, 1964 [N]-[LC], Zagros dwarf loach/Tigris (Persian Gulf basin) (Nikmehr et al., 2019, 2020).

204. *Turcinoemacheilus saadii* Esmaeili, Sayyadzadeh, Özuluğ, Geiger & Freyhof, 2014 [E]-[NE], Saadi dwarf loach/Tigris (Persian Gulf basin) (Esmaeili et al., 2014b).

Subseries Characiphysi**Order Characiformes****Suborder Characoidea****Superfamily Erythrinioidea****Family Serrasalminae****Genus *Piaractus* Eigenmann, 1903**

205. *Piaractus brachipomus* (Cuvier, 1818) [I]-[NE], Pirapitinga/Tigris (Persian Gulf basin) (Esmaili et al., 2017b).

Subseries Siluriphysi**Order Siluriformes****Family Bagridae****Genus *Mystus* Scopoli, 1777**

206. *Mystus pelusius* (Solander, 1794) [N]-[LC], Zugzug Catfish, Tigris mystus/Tigris (Persian Gulf) and Kol River (Hormuz) basins (Esmaili et al., 2017a, 2018).

207. *Mystus cyrusi* Esmaili, Sayyadzadeh, Zarei, Eagderi & Mousavi-Sabet, 2022 [E]-[NE], Cyrus catfish/Helleh drainage (Persis) (Esmaili et al. 2022)

Family Siluridae**Genus *Silurus* Linnaeus, 1758**

208. *Silurus glanis* Linnaeus, 1758 [N]-[LC], Wels catfish/Caspian Sea and Urmia Lake basins (Esmaili et al., 2017a, 2018).

209. *Silurus triostegus* Heckel, 1843 [N]-[LC], Mesopotamian catfish/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

Family Sisoridae**Genus *Glyptothorax* Blyth, 1860**

210. *Glyptothorax galaxias* Mosavi-Sabet, Eagderi, Vatandoust & Freyhof, 2021 [E]-[NE], Glaxias catfish/Karun River drainage (Tigris) (Mosavi-Sabet et al., 2021).

211. *Glyptothorax kurdistanicus* (Berg, 1931) [N]-[DD], Kordestan catfish/Tigris (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

212. *Glyptothorax silviae* Coad, 1981 [E]-[NE], Southern catfish/Tigris and Persis (Persian Gulf basin) (Esmaili et al., 2017a, 2018).

213. *Glyptothorax alidaei* Mosavi-Sabet, Eagderi, Vatandoust & Freyhof, 2021 [E]-[NE], Alidaei's catfish/Karkheh drainage (Tigris) (Mosavi-Sabet et al., 2021).

214. *Glyptothorax hosseinpanahii* Mosavi-Sabet, Eagderi, Vatandoust & Freyhof, 2021 [E]-[NE], Hosseinpanahii catfish/Zohreh River drainage (Persian Gulf basin) (Mosavi-Sabet et al., 2021).

215. *Glyptothorax pallens* Mosavi-Sabet, Eagderi, Vatandoust & Freyhof, 2021 [E]-[NE], Pallens sucking catfish/Sirvan River drainage (Tigris) (Mosavi-Sabet et al., 2021).

216. *Glyptothorax shapuri* Mosavi-Sabet, Eagderi, Vatandoust & Freyhof, 2021 [E]-[NE], Shapuri catfish/Helleh drainage (Persis) (Mosavi-Sabet et al., 2021).

Family Heteropneustidae**Genus *Heteropneustes* Bloch, 1794**

217. *Heteropneustes fossilis* (Bloch, 1794) [I]-[LC], Stinging catfish/introduced to the Tigris River drainages (Persian Gulf basin) (Esmaili et al., 2018).

Superorder Protacanthopterygii**Order Salmoniformes****Family Salmonidae****Genus *Coregonus* Linnaeus, 1758**

218. *Coregonus lavaretus* (Linnaeus, 1758) [I]-[VU], European whitefish/Namak Lake basin (Esmaili et al., 2017, 2018).

Genus *Oncorhynchus* Suckley, 1861

219. *Oncorhynchus mykiss* (Walbaum, 1792) [I]-[NE], Rainbow trout/introduced to the Tigris, Caspian Sea, Urmia Lake, Namak Lake, Kavir, Esfahan and Kor River basins, and widely farmed (Esmaili et al., 2017a, 2018).

Genus *Salmo* Linnaeus, 1758

220. *Salmo caspius* Kessler, 1877 [N]-[NE], Caspian trout/Caspian Sea basin (Esmaili et al., 2017a, 2018).

221. *Salmo trutta* Linnaeus, 1758 [I]-[LC], Brown trout/Caspian Sea, Urmia lake and Namak basins (Esmaili et al., 2017a, 2018).

Genus *Stenodus* Richardson, 1836

222. *Stenodus leucichthys* (Güldenstädt, 1772) [N]-[EW], Inconnu, Sheefish/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Order Esociformes**Family Esocidae****Genus *Esox* Linnaeus, 1758**

223. *Esox lucius* Linnaeus, 1758 [N]-[LC], Northern pike/Caspian Sea basin; introduced in some lakes and reservoirs of Iran (Esmaili et al., 2017a, 2018).

Order Gadiformes**Family Gadidae****Genus *Lota* Oken, 1817**

224. *Lota lota* (Linnaeus, 1758) [N]-[LC], Burbot/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Series Percomorpha**Subseries Gobiida****Order Gobiiformes****Family Gobiidae****Genus *Anatirostrum* Iljin, 1930**

225. *Anatirostrum profundorum* (Berg, 1927) [N]-[NE], Duckbill goby, Duckbill pugolovka/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Genus *Benthophilus* Eichwald, 1831

226. *Benthophilus abdurahmanovi* Ragimov, 1978 [N]-[NE], Abdurakhmanov's tadpole goby/Caspian Sea basin (Abbasi, 2017).

227. *Benthophilus baeri* Kessler, 1877 [N]-[NE], Baer pugolovka/Caspian Sea basin (Abbasi, 2017).

- 228. *Benthophilus ctenolepidus*** Kessler, 1877 [N]-[NE], Transparent tadpole goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018).
- 229. *Benthophilus granulosus*** Kessler, 1877 [N]-[LC], Granular tadpole goby, Granular pugolovka/Caspian Sea basin (Abbasi, 2017).
- 230. *Benthophilus leobergius*** Berg, 1949 [N]-[LC], Caspian stellate tadpole goby /Caspian Sea basin (Abbasi, 2017; Zarei et al., 2021a).
- 231. *Benthophilus macrocephalus*** (Pallas, 1787) [N]-[LC], Caspian tadpole goby, bighead tadpole goby/Caspian Sea basin (Abbasi, 2017).
- 232. *Benthophilus pinchuki*** Ragimov, 1982 [N]-[NE], Pinchuk tadpole goby/Caspian Sea basin (Abbasi, 2017).
- Genus *Boleophthalmus* Valenciennes, 1837**
- 233. *Boleophthalmus dussumieri*** Valenciennes, 1837 [N]-[NE], Dussumier's mudskipper/Tigris, Persis, Hormuz and Makran basins (Esmaeili et al., 2017a, 2018).
- Genus *Glossogobius* Gill, 1859**
- 234. *Glossogobius giuris*** (Hamilton, 1822) [N]-[LC], Tang Goby/Hormuz and Makran basins (Esmaeili et al., 2017a, 2018).
- Genus *Knipowitschia* Iljin, 1927**
- 235. *Knipowitschia caucasica*** (Berg, 1916) [N]-[LC], Caucasian dwarf goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018).
- 236. *Knipowitschia iljini*** Berg, 1931 [N]-[NE], Iljin's dwarf goby/Caspian Sea basin (Esmaeili et al. 2017, 2018).
- 237. *Knipowitschia longicaudata*** (Kessler, 1877) [N]-[LC], Longtail dwarf goby/Caspian Sea basin (Abbasi, 2017).
- Genus *Mesogobius* Bleeker, 1874**
- 238. *Mesogobius nonultimus*** (Iljin, 1936) [N]-[NE], Caspian toad goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018).
- Genus *Neogobius* Iljin, 1927**
- 239. *Neogobius bathybius*** (Kessler, 1877) [N]-[NE], Deepwater goby/Caspian Sea basin (Zarei et al., 2021).
- 240. *Neogobius caspius*** (Eichwald, 1831) [N]-[NE], Caspian goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018; Nikmehr et al., 2021).
- 241. *Neogobius melanostomus*** (Pallas, 1814) [N]-[LC], Round goby, black spotted goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018).
- 242. *Neogobius pallasii*** (Berg, 1916) [N]-[LC], Caspian sand goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018; Nikmehr et al., 2021).
- Genus *Periophthalmus* Bloch & Schneider, 1801**
- 243. *Periophthalmus waltoni*** Koumans, 1941 [N]-[LC], Walton's mudskipper/Tigris, Persis, Hormuz, and Makran basins (Esmaeili et al., 2017a, 2018).
- Genus *Ponticola* Iljin, 1927**
- 244. *Ponticola cyrius*** (Kessler, 1874) [N]-[LC], Kura River goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018).
- 245. *Ponticola goebelii*** (Kessler, 1874) [N]-[NE], Caspian ratan or rotan goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018).
- 246. *Ponticola gorlap*** (Iljin, 1949) [N]-[LC], Caspian bighead goby/Caspian Sea basin (Esmaeili et al., 2017, 2018).
- 247. *Ponticola hircaniaensis*** Zarei, Esmaeili, Kovačić, Schliewen & Abbasi, 2022 [E]-[NE], Hircaniaen bighead goby/Caspian Sea basin (Zarei et al., 2022).
- 248. *Ponticola iranica*** Vasil'eva, Mousavi-Sabet & Vasil'ev, 2015 [E]-[NE], Persian goby/Caspian Sea basin (Vasil'eva et al., 2015).
- 249. *Ponticola patimari*** Eagderi, Nikmehr & Poorbagher, 2020 [E]-[NE], Patimar goby/Caspian Sea basin (Eagderi et al., 2020c).
- 250. *Ponticola ratan*** (Nordmann, 1840) [N]-[NE], Ratan goby/Caspian Sea basin (Abbasi, 2017).
- 251. *Ponticola syrman*** (Nordmann, 1840) [N]-[LC], Syrman goby/Caspian Sea basin (Esmaeili et al., 2017a, 2018).
- Genus *Proterorhinus* Smitt, 1900**
- 252. *Proterorhinus nasalis*** (De Filippi, 1863) [N]-[LC], Eastern tubenose goby/Caspian Sea basin (Zare et al., 2021a).
- Genus *Rhinogobius* Gill, 1859**
- 253. *Rhinogobius lindbergi*** Berg 1933 [I]-[NE], Amur goby/Caspian Sea, Urmia Lake, Hari, Persian Gulf basins (Eagderi and Moradi, 2017; Eagderi et al., 2018b).
- Genus *Scartelaos* Swainson, 1839**
- 254. *Scartelaos tenuis*** (Day, 1876) [N]-[LC], Indian Ocean slender mudskipper/Tigris, Persis, Hormuz, and Makran basins (Esmaeili et al., 2017a, 2018).
- Subseries Ovalentaria**
- Order Mugiliformes**
- Family Mugilidae**
- Genus *Chelon* Artedi, 1793**
- 255. *Chelon auratus*** (Risso, 1810) [I]-[LC], Golden grey mullet/Caspian Sea basin (Durand and Borsa, 2015).
- 256. *Chelon saliens*** (Risso, 1810) [I]-[LC], Leaping mullet/Caspian Sea basin (Durand and Borsa, 2015).
- Genus *Ellochelon* Whitley, 1930**
- 257. *Ellochelon vaigiensis*** (Quoy & Gaimard, 1825) [N]-[LC], Squaretail mullet/Tigris; possibly other coastal rivers in the Persian Gulf (Esmaeili et al., 2017a, 2018).
- Genus *Mugil* Linnaeus, 1758**
- 258. *Mugil cephalus*** Linnaeus, 1758 [N]-[LC], Flathead mullet/The Caspian Sea (Exotic), Tigris and Makran basins; possibly other coastal rivers in the Persian Gulf (Esmaeili et al., 2017, 2018).
- Genus *Planiliza* Whitley, 1945**
- 259. *Planiliza abu*** (Heckel, 1843) [N]-[LC], Abu mullet/Tigris, Persis, and Hormuz; possibly introduced in the Lake Maharlou basin (Durand and Borsa, 2015; Mouludi-Saleh et al., 2021).

260. *Planiliza subviridis* (Valenciennes, 1836) [N]-[NE], Greenback mullet/Tigris and Persis (Persian Gulf basin) (Durand and Borsa, 2015).

Order Cichliformes

Family Cichlidae

Genus *Amatitlania* Schmitter-Soto, 2007

261. *Amatitlania nigrofasciata* (Günther, 1867) [I]-[NE], Convict cichlid/Namak and Hormuz basins (Esmaeili et al., 2013; Mousavi-Sabet and Eagderi, 2016b).

Genus *Coptodon* Gervais, 1848

262. *Coptodon zillii* (Gervais, 1848) [I]-[NE], Redbelly tilapia/Tigris and Persis (Persian Gulf basin) (Teimori et al., 2017; Radkhah and Eagderi, 2022).

Genus *Iranocichla* Coad, 1982

263. *Iranocichla hormuzensis* Coad, 1982 [E]-[NE], Hormuz cichlid/Mehran River (Hormuz) (Esmaeili et al., 2016d; Schwarzer et al., 2016).

264. *Iranocichla persa* Esmaeili, Sayyadzadeh & Seehausen, 2016 [E]-[NE], Persis cichlid/Shur, Hasanlangi and Minab River drainages flowing into the Persian Gulf at the Strait of Hormuz (Esmaeili et al., 2016d).

265. *Iranocichla* sp. [E]-[NE], Hormuz (Kol River drainages, see Schwarzer et al., 2016).

Genus *Oreochromis* Günther, 1889

266. *Oreochromis aureus* (Steindachner, 1864) [I]-[LC], Blue tilapia/Tigris (Persian Gulf basin) (Radkhah and Eagderi, 2022).

267. *Oreochromis niloticus* (Linnaeus, 1758) [I]-[NE], Nile tilapia/Persis basin (Rafiee et al., 2017).

Infraseries Atherinomorpha

Order Atheriniformes

Suborder Atherinoidei

Family Atherinidae

Subfamily Atherininae

Genus *Atherina* Linnaeus, 1758

268. *Atherina caspia* Eichwald, 1831 [N]-[NE], Caspian silverside/Caspian Sea basin (Esmaeili et al., 2017a, 2018).

Order Cyprinodontiformes

Superfamily Cyprinodontoidea

Family Aphaniidae

Genus *Aphaniops* Hoedeman, 1951

269. *Aphaniops furcatus* (Teimori, Esmaeili, Erpenbeck & Reichenbacher, 2014) [E]-[NE], Scaleless tooth-carp/Hormuz and Makran basins (Esmaeili et al., 2020b).

270. *Aphaniops ginaonis* (Holly, 1929) [E]-[NE], Geno (Genow) tooth-carp/Hormuz basin (Esmaeili et al., 2020b).

271. *Aphaniops hormuzensis* (Teimori, Esmaeili, Hamidan & Reichenbacher, 2018) [E]-[NE], Hormuz tooth-carp/Mehran River drainage (Esmaeili et al., 2020b; Mouludi-Saleh et al., 2020c; Teimori et al., 2020).

272. *Aphaniops stoliczkanus* (Day, 1872) [N]-[NE], Eastern tooth-carp, Indian tooth-carp/Tigris, Persis, Makran and Mashkid (Esmaeili et al., 2020b).

Genus *Esmaeilius* Freyhof & Yoğurtçuoğlu 2020

273. *Esmaeilius darabensis* (Esmaeili, Teimori, Gholami & Reichenbacher, 2014) [E]-[NE], Darab tooth-carp/Hormuz basin (Esmaeili et al., 2014e).

274. *Esmaeilius persicus* (Jenkins, 1910) [E]-[NE], Persian tooth-carp/ Maharlou Lake basin (Teimori et al., 2011).

275. *Esmaeilius isfahanensis* (Hrbek, Keivany & Coad, 2007) [E]-[NE], Esfahan tooth-carp/Zayandeh Rud (Esfahan) basin (Hrbek et al., 2007).

276. *Esmaeilius shirini* (Gholami, Esmaeili, Erpenbeck & Reichenbacher, 2014) [E]-[NE], Shirin tooth-carp/endemic to the Kor River basin but has been translocated to the Helleh River drainage (Persis) (Gholami et al., 2014; Freyhof & Yoğurtçuoğlu, 2020).

277. *Esmaeilius sophiae* (Heckel, 1849) [E]-[NE], Kor tooth-carp/endemic to the Kor River basin but has been translocated to the Persis and Tigris (Esmaeili et al., 2020b; Freyhof & Yoğurtçuoğlu, 2020).

278. *Esmaeilius vladykovi* (Coad, 1988) [E]-[NE], Zagros tooth-carp/Tigris (Esmaeili et al., 2020b; Freyhof & Yoğurtçuoğlu, 2020).

Genus *Paraphanius* Esmaeili, Teimori, Zarei & Sayyadzadeh, 2020

279. *Paraphanius mento* (Heckel, 1843) [N]-[LC], Iridescent tooth-carp/Tigris (Persian Gulf basin) (Esmaeili et al., 2020b). There is only one record of this species. No recent record.

Superfamily Poecilioidea

Family Poeciliidae

Genus *Gambusia* Poey, 1854

280. *Gambusia holbrooki* Girard, 1859 [I]-[LC], Eastern mosquitofish/introduced and widespread into Iran to provide biological control for mosquitoes (Radkhah et al., 2022b).

Genus *Poecilia* Bloch & Schneider, 1801

281. *Poecilia latipinna* (Lesueur, 1821) [I]-[LC], Sailfin molly/Esfahan and Tigris (Persian Gulf basin) (Moshayedi et al., 2015).

282. *Poecilia reticulata* Peters, 1859 [I]-[NE], Guppy/Namak basin (Mousavi-Sabet et al., 2014).

Genus *Xiphophorus* Heckel, 1848

283. *Xiphophorus hellerii* Heckel, 1848 [I]-[LC], Green swordtail/Namak Lake and Persis (Esmaeili et al., 2010b; Eagderi et al., 2015).

Order Synbranchiformes

Suborder Mastacembeloidei

Family Mastacembelidae

Genus *Mastacembelus* Scopoli, 1777

284. *Mastacembelus mastacembelus* (Banks & Solander, 1794) [N]-[LC], Mesopotamian spiny eel/Tigris,

Kor and Persis basins. Its presence from the Kor River should be confirmed by specimen (Gholamhosseini et al., 2022).

Order Anabantiformes

Family Channidae

Genus *Channa* Scopoli, 1777

285. *Channa gachua* (Hamilton, 1822) [N]-[LC], Dwarf snakehead/Hamun-e Jaz Murian, Makran and Mashkid basins (Mouludi-Saleh et al., 2019).

Order Syngnathiformes

Suborder Syngnathoidei

Superfamily Syngnathoidea

Family Syngnathidae

Genus *Syngnathus* Linnaeus, 1758

286. *Syngnathus caspius* Eichwald, 1831 [N]-[LC], Caspian pipefish/Caspian Sea basin (Zarei et al., 2021b).

Order Perciformes

Suborder Percoidei

Superfamily Percoidea

Family Percidae

Genus *Perca* Linnaeus, 1758

287. *Perca fluviatilis* Linnaeus, 1758 [N]-[LC], European perch/Caspian Sea basin (Abbasi et al., 2021b).

Genus *Sander* Oken, 1817

288. *Sander lucioperca* (Linnaeus, 1758) [N]-[LC], Pike perch/Caspian Sea basin; introduced to lakes and reservoirs throughout Iran (Esmaili et al., 2017a, 2018).

289. *Sander marinus* (Cuvier, 1828) [N]-[DD], Estuarine perch/Caspian Sea basin (Esmaili et al., 2017, 2018).

Order Scorpaeniformes

Suborder Gasterosteoidae

Family Gasterosteidae

Genus *Gasterosteus* Linnaeus, 1758

290. *Gasterosteus aculeatus* Linnaeus, 1758 [I]-[LC], Three-spined stickleback/Caspian Sea, Kavir and Hari basins (Mouludi-Saleh and Eagderi, 2021).

Genus *Pungitius* Coste, 1848

291. *Pungitius platygaster* (Kessler, 1859) [N]-[LC], Ukrainian stickleback/Caspian Sea basin (Esmaili et al., 2017a, 2018).

Order Spariformes

Family Sparidae

Genus *Acanthopagrus* Peters, 1855

292. *Acanthopagrus arabicus* Iwatsuki, 2013 [N]-[LC], Arabian yellowfin seabream/ /Tigris (Persian Gulf), Hormuz and Persis basins (Esmaili et al., 2017a, 2018).

4. Discussion

Based on the present checklist, we document the presence of 292 fish species in 19 inland water basins of Iran. The confirmed freshwater lampreys and freshwater fishes of Iran comprise 106 genera, 36 families, 24 orders, and 3 classes. In addition, 43 species are excluded from the checklist. This reveals species richness and a high degree of endemism in the Iranian ichthyofauna. The endemic fish species comprise 35% (102 species) in Iran.

4.1. Alien species

A total of 29 i.e. *Acipenser baerii*, *Amatitlania nigrofasciata*, *Anguilla anguilla*, *Atractosteus spatula*, *Carassius auratus*, *C. gibelio*, *C. langsdorfii*, *Chelon saliens*, *Chelon auratus*, *Coptodon zillii*, *Coregonus lavaretus*, *Ctenopharyngodon idella*, *Gambusia holbrooki*, *Gasterosteus aculeatus*, *Hemiculter leuciscus*, *Heteropneustes fossilis*, *Hypophthalmichthys molitrix*, *H. nobilis*, *Labeo rohita*, *Mylopharyngodon piceus*, *Oncorhynchus mykiss*, *Oreochromis aureus*, *O. niloticus*, *Poecilia latipinna*, *P. reticulata*, *Pseudorasbora parva*, *Piaractus brachipomus*, *Rhinogobius lindbergi* and *Xiphophorus hellerii* have been introduced deliberately or accidentally into Iranian inland waters.

4.2. Excluded species

According to three exclusion criteria, 43 recorded fish species in Iranian inland waters are misidentified or erroneously recorded (Table 2). Based on available literature, we present 14 synonyms, 1 erroneously listed species that whose occurrence is geographically impossible and 28 species that need confirmation by the specimen (Table 2). However, we keep this list open for later addenda to insert based on further documentation, which may confirm their occurrence by specimens. The citation of doubtful records without verification had caused a great number of errors (Kottelat and Freyhof 2007); therefore, we rechecked and verified the previous lists based on our expedition during the past 20 years and examination of available materials to us. Furthermore, the first records in scientific journals were carefully read and verified. Therefore, further research and taxonomic revisions are crucial to solve these taxonomic errors that we listed in Table 2.

Acknowledgments

This work was financially supported by the University of Tehran.

References

- Abbasi K, Mouludi-Saleh A, Eagderi S, Sarpanah A (2021b). Morphological comparison of Eurasian perch (*Perca fluviatilis* L., 1758) in the Anzali and Chamkhaleh wetlands, Guilan Province. *Journal of Aquatic Ecology* 10 (4): 28-35.
- Abbasi Ranjbar K, Mouludi-Saleh A, Eagderi S, Sarpanah A (2018). Distinguishing Meristic and Morphometric Traits in Three Species of the Genus *Acanthopagrus* from Iranian Inland Waters. *Taxonomy and Biosystematics* 10 (36): 49-58.

- Abbasi K. (2017). Fishes of Guilan. Farhang-e Ilia publication. 206 p.
- Abbasi K, Mouludi-Saleh A, Eagderi S (2021a). Morphological diversity of the Caspian Asp, *Leuciscus aspius*, in the South Caspian Sea basin (Osteichthyes: Cyprinidae). *Zoology in the Middle East* 67 (1): 25-31. doi:10.1080/09397140.2021.1879439
- Abbasi K, Mouludi-Saleh A, Eagderi S (2021c). The variability of morphometric and meristic characteristics of Urmia chub, *Petroleuciscus ulanus* (Günther, 1899) in the Mahabdi-Chai and Godar-Chai rivers, Lake Urmia basin. *Limnology and Freshwater Biology* 2021 (4): 1156-1159. doi:10.31951/2658-3518-2021-A-4-1156
- Abdoli A (2000). The Inland Water Fishes of Iran. Iranian Museum of Nature and Wildlife, Tehran. 378 p.
- Ahmdzadeh M, Poorbagher H, Eagderi S (2019). Calculating the habitat suitability index of Siahmahi (*Capoeta buhsei*, Kessler 1877) using the kernel smoothing in the Jajrood River, Namak basin of Iran. *Aquaculture Sciences* 6 (2): 99-108.
- Alwan NH, Zareian H, Esmaeili HR (2016). *Capoeta coadi*, a new species of cyprinid fish from the Karun River drainage, Iran based on morphological and molecular evidences (Teleostei, Cyprinidae). *ZooKeys* 572: 155. doi:10.3879/zookeys.572.377
- Amiri K, Shabanipour N, Eagderi S (2017a). Predict the potential fishing grounds for Kilka (*Clupeonella* spp.) fishes in southern part of the Caspian Sea using maximum entropy models and remotely sensed satellite data. *Iranian Journal of Ichthyology* 4 (3): 290-298. doi:10.22034/iji.v4i3.238
- Amiri K, Shabanipour N, Eagderi S (2017b). Using kriging and co-kriging to predict distributional areas of Kilka species (*Clupeonella* spp.) in the southern Caspian Sea. *International Journal of Aquatic Biology* 5 (2): 108-113. doi:10.22034/ijab.v5i2.309
- Amiri K, Shabanipour N, Eagderi S (2018). Forecasting the catch of kilka species (*Clupeonella* spp.) using Time Series SARIMA models in the Southern Caspian Sea. *Caspian Journal of Environmental Sciences* 16 (4): 349-358. doi:10.22124/cjes.2018.3203
- Asgari R, Rafiee G, Poorbagher H, Agh N, Zadeh HE (2013). Body shape changes during the early development of the Beluga (*Huso huso*). *International Journal of Aquatic Biology* 1 (1): 1-5. doi.org/10.22034/ijab.v1i1.9
- Çiçek E, Eagderi S, Sungur S (2018). *Oxynoemacheilus veyseli*, a new nemacheilid species from the upper Aras River drainage of Turkey (Teleostei: Nemacheilidae). *Iranian Journal of Ichthyology* 5 (3): 232-242. doi:10.22034/iji.v5i3.302
- Çiçek E, Eagderi S, Secer B, Sungur S (2022). Taxonomic status of the genus *Sabanejewia* (Cobitidae) from Kura-Aras river system (Turkey). *Transylvanian Review of Systematical and Ecological Research* 24 (1). doi:10.2478/trser-2022-0006
- Coad BW (1995). Freshwater Fishes of Iran. *Acta Scientiarum Naturalium Academiae Scientiarum Bohemicae*, Brno 29 (1): 1-64.
- Coad BW (1998). Systematic biodiversity in the freshwater fishes of Iran. *Italian Journal of Zoology* 65 (S1): 101-108. https://doi.org/10.1080/11250009809386802
- Dastanpoor N, Eagderi S, Farahmand H, Mousavi-Sabet H (2021). Morphological Variations and Diagnostic Characteristics of *Chondrostoma regium* Populations in Iranian Inland Waters. *Taxonomy and Biosystematics* 13 (46): 79-92.
- Durand JD, Borsa P (2015). Mitochondrial phylogeny of grey mullets (Acanthopterygii: Mugilidae) suggests high proportion of cryptic species. *Comptes Rendus Biologies* 338 (4): 266-277. doi:10.1016/j.crv.2015.01.007
- Eagderi S, Moradi M (2017). Range extension of the lake goby *Rhinogobius similis* Gill, 1859 (Teleost: Gobiidae) to Urmia Lake basin in northwestern Iran. *Biharean Biologist* 11 (2): 123-125.
- Eagderi S, Mouludi-Saleh A, Ghaderi E, Freyhof J (2022). First record of *Oxynoemacheilus zarzianus* Freyhof & Geiger, 2017 from Iran (Teleostei: Nemacheilidae). *Iranian Journal of Ichthyology* 9 (1): 11-15. doi.org/10.22034/iji.v9i1.829
- Eagderi S, Roudbar AJ, Birecikligil SS, Cicek E, Coad BW (2017b). *Chondrostoma esmaeili*, a new cyprinid species from the Tigris river drainage in Iran (Teleostei: Cyprinidae). *Vertebrate Zoology* 67 (2): 125-132.
- Eagderi S, Jalili P, Çiçek E (2018a). *Oxynoemacheilus elsaе*, a new species from the Urmia Lake basin of Iran (Teleostei: Nemacheilidae). *FishTaxa* 3 (2): 453-459.
- Eagderi S, Jouladeh-Roudbar A, Jalili P, Sayyadzadeh G, Esmaeili HR (2017c). Taxonomic statue of the genus *Cobitis* Linnaeus, 1758 (Teleostei: Cobitidae) in the southern Caspian Sea basin, Iran with description of a new species. *FishTaxa* 2 (1): 48-61.
- Eagderi S, Mojazi Amiri B, Poorbagher H, Nasrollah Pourmoghadam M, Nemati Mobin N (2017d). Effects of cadmium on morphological structure of sperm in *Caspiomyzon wagneri* (Kessler, 1870) (Petromyzontiformes: Petromyzontidae). *Caspian Journal of Environmental Sciences* 15 (2): 125-134.
- Eagderi S, Mojazi AB, Adriaens D (2013). Description of the ovarian follicle maturation of the migratory adult female bulatmai barbel (*Luciobarbus capito*, Guldenstadt 1772) in captivity. *Iranian Journal of Fisheries Sciences* 12 (3): 550-560.
- Eagderi S, Mouludi-Saleh A, Cicek E (2020b). Length-weight relationship of ten species of Leuciscinae sub-family (Cyprinidae) from Iranian inland waters. *International Aquatic Research* 12 (2): 133-136. doi: 10.22034/iar(20).2020.1891648.1004
- Eagderi S, Mouludi-Saleh A, Nazlabadi SA (2019c). First record of the rohu, *Labeo rohita* (Hamilton, 1822) (Cyprinidae) from Karun River, Tigris River drainage, Iran. *FishTaxa* 4 (1): 18-20.
- Eagderi S, Mouludi-Saleh A, Ahmadi S, Javadzadeh N (2020a). Phenotypic plasticity of the body shape in Prussian carp (*Carassius gibelio*), in response to lentic and lotic habitats using geometric morphometric technique. *Iranian Scientific Fisheries Journal* 29 (1): 49-58. doi: 20.1001.1.10261354.1399.29.1.5.3
- Eagderi S, Mouludi-Saleh A, Hosseini V, Mousavi-Sabet H (2021). Documentation of the Turkestan barbel, *Luciobarbus conocephalus* (Kessler, 1872) in the Iranian part of Hari River basin (Teleostei: Cyprinidae: Barbinae). *Iranian Journal of Ichthyology* 8 (1): 67-75. doi: 10.22034/iji.v8i1.615
- Eagderi S, Mousavi-Sabet H, Freyhof J (2019d). *Paraschistura*

- makranensis*, a new loach from the Jegin River drainage in southern Iran with comments on *P. ilamensis* and *P. pasatigris* (Teleostei: Nemacheilidae). Zootaxa 4668 (2): 258-270. doi: 10.11646/zootaxa.4668.2.6
- Eagderi S, Nasri, M, Çiçek E (2018b). First record of the Amur goby *Rhinogobius lindbergi* Berg 1933 (Gobiidae) from the Tigris River drainage, Iran. International Journal of Aquatic Biology 6 (4): 202-207. doi: 10.22034/ijab.v6i4.528
- Eagderi S, Nikmehr, N, Freyhof J (2019b). *Alburnus zagrosensis*, a junior synonym of *Alburnus sellal* (Teleostei: Leuciscidae). Zootaxa 4652 (2): 367-374. doi: 10.11646/zootaxa.4652.2.9
- Eagderi S, Nikmehr N, Poorbagher H (2020c). *Ponticola patimari* sp. nov. (Gobiiformes: Gobiidae) from the southern Caspian Sea basin, Iran. FishTaxa 17: 22-31.
- Eagderi S, Nikmehr N, Çiçek E, Esmaeili HR, Vatandoust S, et al. (2019a). *Barbus urmianus* a new species from Urmia Lake basin, Iran (Teleostei: Cyprinidae). International Journal of Aquatic Biology 7 (4): 239-244. doi: 10.22034/ijab.v7i4.725
- Eagderi S, Mousavi-Sabet H (2021). *Capoeta raghazensis*, a new species of algae-scraping cyprinid from the Raghaz Canyon in Hormuz basin, southern Iran (Teleostei: Cyprinidae). FishTaxa 22: 37-44.
- Eagderi S, Poorbagher H, Moshayedi F, Hosseini SV (2017a). Morphological development and allometric growth patterns of *Acipenser persicus* Borodin, 1897 (Actinopterygii, Acipenseridae) during early development. International Journal of Aquatic Biology 5 (3): 201-207. doi: 10.22034/ijab.v5i3.312
- Eagderi S, Poorbagher H, Parsazade F, Mousavi-Sabet H (2015). Effects of rearing temperature on the body shape of swordtail (*Xiphophorus hellerii*) during the early development using geometric morphometric. Poeciliid Research 5 (1): 24-30.
- Esmaeili HR, Khaefi R, Zammanian Nejad R (2016a). Historical review on the taxonomy of *Squalius berak* Heckel, 1843 (Teleostei: Cyprinidae). FishTaxa 1 (3): 118-126.
- Esmaeili HR, Khajehpanah A, Mehraban H, Elmi A, Malekzahi H et al. (2015). Fishes of the Mashkid and Makran basins of Iran: an updated checklist and ichthyogeography. Iranian Journal of Ichthyology 2 (2): 113-132. doi: 10.22034/iji.v2i2.36
- Esmaeili HR, Masoudi M, Amini Chermahini M, Esmaeili AH, Zarei F et al. (2017b). Invasion of the Neotropical and Nearctic fishes to Iran. FishTaxa 2 (3): 126-133.
- Esmaeili HR, Teimori A, Owfi F, Abbasi K, Coad BW (2014a). Alien and invasive freshwater fish species in Iran: Diversity, environmental impacts and management. Iranian Journal of Ichthyology 1 (2): 62-72. doi: 10.22034/iji.v1i2.4
- Esmaeili HR, Zareian H, Eagderi S, Alwan N (2016b). Review on the taxonomy of Tigris scraper, *Capoeta umbla* (Heckel, 1843) and its confirmation record from the Iranian part of Tigris River, Persian Gulf basin (Teleostei: Cyprinidae). FishTaxa 1: 35-44.
- Esmaeili HR, Coad BW, Gholamifard A, Nazari N, Teimori A (2010a). Annotated checklist of the freshwater fishes of Iran. Zoosystematica Rossica 19: 361-386.
- Esmaeili HR, Coad BW, Mehraban HR, Masoudi M, Khaefi R et al. (2014b). An updated checklist of fishes of the Caspian Sea basin of Iran with a note on their zoogeography. Iranian Journal of Ichthyology 1 (3): 152-184. doi: 10.22034/iji.v1i3.18
- Esmaeili HR, Gholamifard A, Sayyadzadeh G, Parsi B, Mirghiyasi S, Ghasemian S (2013b). New record of the convict cichlid, *Amatitlania nigrofasciata* (Günther, 1867), from the Middle East (Actinopterygii: Cichlidae). Aqua, International Journal of Ichthyology 19: 225-229.
- Esmaeili HR, Gholamifard A, Teimori A, Bagh-Bani S, Coad BW (2010b). *Xiphophorus hellerii* Heckel, 1848 (Cyprinodontiformes, Poeciliidae), a newly introduced fish recorded from natural freshwaters of Iran. Journal of Applied Ichthyology 26: 937-938. doi: 10.1111/j.1439-0426.2010.01515.x
- Esmaeili HR, Malekzahi H, Pazira A, Freyhof J (2013a). First record of the Kalabans, *Bangana dero* (Hamilton, 1822), from Iran (Teleostei: Cyprinidae). Zoology in the Middle East 59 (1): 89-91. doi: 10.1080/09397140.2013.795074
- Esmaeili HR, Mehraban H, Abbasi K, Keivany Y, Coad BW (2017a). Review and updated checklist of freshwater fishes of Iran: taxonomy, distribution and conservation status. Iranian Journal of Ichthyology 4 (Suppl. 1): 1-114. doi: 10.22034/iji.v4iSuppl.%201.220
- Esmaeili HR, Sayyadzadeh G, Eagderi S, Abbasi K (2018). Checklist of freshwater fishes of Iran. FishTaxa 3 (3): 1-95.
- Esmaeili HR, Sayyadzadeh G, Zarei F, Eagderi S, Mousavi-Sabet H (2022). *Mystus cyrusi*, a new species of bagrid catfish (Teleostei: Bagridae) from Middle East. Zootaxa 5099 (3): 325-343. doi: 10.11646/zootaxa.5099.3.2
- Esmaeili HR, Sayyadzadeh G, Seehausen O (2016d). *Iranocichla persa*, a new cichlid species from southern Iran (Teleostei, Cichlidae). ZooKeys 636: 141. doi:10.3897/zookeys.636.10571
- Esmaeili HR, Sayyadzadeh G, Japoshvili B, Eagderi S, Abbasi K, et al. (2020a). *Rhodeus caspius*, a new bitterling from Iran (Teleostei: Cypriniformes: Acheilognathidae). Zootaxa 4851 (2): 319-337. doi: 10.11646/zootaxa.4851.2.6
- Esmaeili HR, Teimori A, Zarei F, Sayyadzadeh G (2020b). DNA barcoding and species delimitation of the Old-World tooth-carps, family *Aphaniidae* Hoedeman, 1949 (Teleostei: Cyprinodontiformes). PLoS One 15: e0231717. doi: 10.1371/journal.pone.0231717
- Esmaeili HR, Sayyadzadeh G, Özuluğ M, Geiger M, Freyhof J (2014b). Three new species of *Turcinoemacheilus* from Iran and Turkey (teleostei: Nemacheilidae). Ichthyological Exploration of Freshwaters 24 (3): 257-273.
- Esmaeili HR, Teimori A, Gholami Z, Reichenbacher B (2014e). Two new species of the tooth-carp *Aphanius* (Teleostei: Cyprinodontidae) and the evolutionary history of the Iranian inland and inland-related *Aphanius* species. Zootaxa 3786 (3): 246-268. doi: 10.11646/zootaxa.3786.3.2
- Esmaeili HR, Sayyadzadeh G, Coad BW, Eagderi S (2016c). Review of the genus *Garra* Hamilton, 1822 in Iran with description of a new species: a morpho-molecular approach (Teleostei: Cyprinidae). Iranian Journal of Ichthyology 3 (2): 82-121.

- Esmaili HR, Mousavi-Sabet H, Sayyadzadeh G, Vatandoust S, Freyhof J (2014d). *Paracobitis atrakensis*, a new species of crested loach from northeastern Iran (Teleostei: Nemacheilidae). *Ichthyological Exploration of Freshwaters* 25 (3): 237-242.
- Freyhof J (2016a). Redescription of *Garra elegans* (Günther, 1868), a poorly known species from the Tigris River drainage (Teleostei: Cyprinidae). *Zootaxa* 4173 (5): 496-500. doi: 10.11646/zootaxa.4173.5.7
- Freyhof J, Geiger MF, Golzarianpour K, Patimar R (2016). *Sasanidus*, a new generic name for *Noemacheilus kermanshahensis* Bănărescu and Nalbant, with discussion of *Ilamnemacheilus* and *Schistura* (Teleostei; Nemacheilidae). *Zootaxa* 4107 (1): 65-80. doi: 10.11646/zootaxa.4107.1.3
- Freyhof J (2016b). *Oxynoemacheilus karunensis*, a new species from the Persian Gulf basin Teleostei: Nemacheilidae. *Zootaxa* 4175 (1): 94-100. doi: 10.11646/zootaxa.4175.1.9
- Freyhof J, Esmaili, HR, Sayyadzadeh G, Geiger M (2014). Review of the crested loaches of the genus *Paracobitis* from Iran and Iraq with the description of four new species (Teleostei: Nemacheilidae). *Ichthyological Exploration of Freshwaters* 25 (1): 11-38.
- Freyhof J, Sayyadzadeh G, Esmaili HR, Geiger M (2015). Review of the genus *Paraschistura* from Iran with description of six new species (Teleostei: Nemacheilidae). *Ichthyological Exploration of Freshwaters* 26 (1): 1-48.
- Freyhof J, Kaya C, Geiger Mf (2022). A practical approach to revise the *Oxynoemacheilus bergianus* species group (Teleostei: Nemacheilidae). *Zootaxa* 5128 (2): 151-194. <https://doi.org/10.11646/zootaxa.5128.2.1>
- Freyhof J, Yoğurtcuoğlu B (2020). A proposal for a new generic structure of the killifish family Aphaniidae, with the description of *Aphaniops teimorii* (Teleostei: Cyprinodontiformes). *Zootaxa* 4810 (3): 421-451. doi.org/10.11646/zootaxa.4810.3.2
- Ganjali Z, Esmaili HR, Zarei F, Sayyadzadeh G, Eagderi S et al. (2021). West Asian colonisation of topmouth gudgeon, *Pseudorasbora parva* (Teleostei: Gobionidae): Genetic admixture at the crossroad of Europe and East Asia. *Freshwater Biology* 66 (4): 699-715. doi: 10.1111/fwb.13671
- Gholami Z, Esmaili HR, Erpenbeck D, Reichenbacher B (2014). Phylogenetic analysis of *Aphanius* from the endorheic Kor River Basin in the Zagros Mountains, South-western Iran (Teleostei: Cyprinodontiformes: Cyprinodontidae). *Journal of Zoological Systematics and Evolutionary Research* 52 (2): 130-141. doi: 10.1111/jzs.12052
- Gholamhosseini A, Razbanian M, Esmaili HR, Eagderi S (2022). Molecular systematics and morphological variation in the Mesopotamian spiny eel *Mastacembelus mastacembelus* (Teleostei: Mastacembelidae). *The European Zoological Journal* 89 (1): 546-555. doi:10.1080/24750263.2022.2057604
- Golzarianpour K, Abdoli A, Freyhof J (2011). *Oxynoemacheilus kiabii*, a new loach from Karkheh River drainage, Iran (Teleostei: Nemacheilidae). *Ichthyological Exploration of Freshwaters* 22 (3): 201.
- Hasanalipour A, Eagderi S, Poorbagher H, Bahmani M (2013). Effects of stocking density on blood cortisol, glucose and cholesterol levels of immature Siberian sturgeon (*Acipenser baerii* Brandt, 1869). *Turkish Journal of Fisheries and Aquatic Sciences* 13 (1). doi: 10.4194/1303-2712-v13_1_04
- Hoestlandt H (1991). The freshwater fishes of Europe, Clupeidae, Anguillidae. Wiesbaden: AULA.
- Hrbek T, Keivany Y, Coad BW (2006). New species of *Aphanius* (Teleostei, Cyprinodontidae) from Isfahan Province of Iran and a reanalysis of other Iranian species. *Copeia* 2006 (2): 244-255. doi: 10.1643/0045-8511(2006)6[244:NSOATC]2.0.CO;2
- Jalili P, Eagderi S, Nikmehr N, Keivany Y (2015). Descriptive osteology of *Barbus cyri* (Teleostei: Cyprinidae) from southern Caspian Sea basin. *Iranian Journal of Ichthyology* 2 (2): 105-112.
- Jouladeh-Roudbar A, Vatandoust S, Eagderi S, Jafari-Kenari S, Mousavi-Sabet H (2015a). Freshwater fishes of Iran; an updated checklist. *Aquaculture, Aquarium, Conservation & Legislation Bioflux* 8(6): 855-909.
- Jouladeh-Roudbar A, Eagderi S (2017). Study on phylogenetic of *Alburnoides namaki* Bogutskaya and Coad, 2009 using COI gene. *Modern Genetic Journal* 11 (4): 531-538.
- Jouladeh-Roudbar A, Eagderi S. (2019). A re-evaluation of taxonomic status of *Alburnoides parhami* using morphological and genetic data. *Modern Genetic Journal* 14 (2): 101-110.
- Jouladeh-Roudbar A, Eagderi S, Vatandoust S (2015b). First record of *Paraschistura alta* (Nalbant and Bianco, 1998) from Eastern Iran and providing its COI barcode region sequences (Teleostei: Nemacheilidae). *Iranian Journal of Ichthyology* 2 (4): 235-243.
- Jouladeh-Roudbar A, Eagderi S, Esmaili HR, Coad BW, Bogutskaya N (2016). A molecular approach to the genus *Alburnoides* using COI sequences data set and the description of a new species, *A. damghani*, from the Damghan River system (the Dasht-e Kavir Basin, Iran) (Actinopterygii, Cyprinidae). *Zookeys* (579): 157. doi:10.3897/zookeys.579.7665
- Jouladeh-Roudbar A, Eagderi S, Murillo-Ramos L, Ghanavi, HR, Doadrio I (2017a). Three new species of algae-scraping cyprinid from Tigris River drainage in Iran (Teleostei: Cyprinidae). *FishTaxa* 2 (3): 134-155.
- Jouladeh-Roudbar A, Ghanavi HR, Doadrio I (2020). Ichthyofauna from Iranian freshwater: Annotated checklist, diagnosis, taxonomy, distribution and conservation assessment. *Zoological Studies* 59. doi: 10.6620/ZS.2020.59-21
- Jouladeh-Roudbar A, Eagderi S, Ghanavi HR, Doadrio I (2017b). A new species of the genus *Capoeta* Valenciennes, 1842 from the Caspian Sea basin in Iran (Teleostei, Cyprinidae). *ZooKeys* (682): 137-155. doi:10.3897/zookeys.682.12670
- Kamangar BB, Prokofiev AM, Ghaderi E, Nalbant TT (2014). Stone loaches of Choman River system, Kurdistan, Iran (Teleostei: Cypriniformes: Nemacheilidae). *Zootaxa* 3755 (1): 033-061. doi: 10.11646/zootaxa.3755.1.2
- Keivany Y, Nasri M, Abbasi K, Abdoli A (2016). Atlas of inland water fishes of Iran. Iran Department of Environment, Tehran, Iran. 218 p.

- Khaefi R, Esmaeili HR, Geiger MF, Eagderi S (2017a). Taxonomic review of the cryptic *Barbus lacerta* species group with description of a new species (Teleostei: Cyprinidae). *FishTaxa*, 2 (2): 90-115.
- Khaefi R, Esmaeili HR, Sayyadzadeh G, Geiger MF, Freyhof J. (2016). *Squalius namak*, a new chub from Lake Namak basin in Iran (Teleostei: Cyprinidae). *Zootaxa* 4169 (1): 145-159. doi: 10.11646/zootaxa.4169.1.7
- Khaefi R, Teimori A, Esmaeili HR (2017b). Phylogenetic relationships and taxonomy of *Luciobarbus barbulus* (Heckel, 1847) (Teleostei: Cyprinidae). *Journal of Ichthyology* 57 (6): 835-845. doi: 10.1134/S0032945217060078
- Khaefi R, Esmaeili HR, Chermahini MA (2018). Natural Hybridization of *Luciobarbus barbulus* x *Luciobarbus kersin* and *Luciobarbus barbulus* x *Luciobarbus xanthopterus* in the Persian Gulf Basin. *Turkish Journal of Fisheries and Aquatic Sciences* 18 (12): 1399-1407. doi: 10.4194/1303-2712-v18_12_08
- Khosravi M, Abdoli A, Tajbakhsh F, Ahmadzadeh F, Nemati H, et al. (2022). An Effort toward Species Delimitation in the Genus *Carassius* (Cyprinidae) using Morphology and the Related Challenges: A Case Study from Inland Waters of Iran. *Journal of Ichthyology* 62: 185-194. doi: 10.1134/S0032945222020096
- Kottelat M, Freyhof J (2007). *Handbook of European Freshwater Fishes*. Kottelat, Cornol, Switzerland and Freyhof. Berlin. Germany. 646 p.
- Kuljanishvili T, Eptashvili G, Freyhof J, Japoshvili B, Kalous L et al. (2020). Checklist of the freshwater fishes of Armenia, Azerbaijan and Georgia. *Journal of Applied Ichthyology* 36 (4): 501-514. doi: 10.1111/jai.14038
- Levin BA, Simonov EP, Ermakov OA, Levina MA, Interesova EA et al. (2016). Phylogeny and phylogeography of the roaches, genus *Rutilus* (Cyprinidae), at the Eastern part of its range as inferred from mtDNA analysis. *Hydrobiologia* 788 (1): 33-46. doi: 10.1007/s10750-016-2984-3
- Kuljanishvili T, Eptashvili G, Freyhof J, Japoshvili B, Kalous L, Levin B, et al. (2020). Checklist of the freshwater fishes of Armenia, Azerbaijan and Georgia. *Journal of Applied Ichthyology* 36 (4): 501-514. doi.org/10.1111/jai.14038
- Mohammadi S, Eagderi S, Poorbagher H (2018). Comparison of morphological characteristics of *Oxyneomacheilus bergi* (Gratsianov, 1907) populations in Aras river drainage. *Protection of Iranian Endemic Freshwater Fishes Conference*. Karaj, Iran. 19 December 2018. pp: 268-274.
- Mohammadian-Kalat T, Esmaeili HR, Aliabadian M, Freyhof J (2017). Re-description of *Alburnus doriae*, with comments on the taxonomic status of *A. amirkabiri*, *A. mossulensis*, *A. sellal* and *Petroleuciscus esfahani* (Teleostei: Cyprinidae). *Zootaxa* 4323 (4): 487-502. doi: 10.11646/zootaxa.4323.4.3
- Moshayedi F, Eagderi S, Jalili P, Mousavi-Sabet H (2015). Allometric growth pattern and morphological development of sailfin molly *Poecilia latipinna* (Cyprinodontiformes, Poeciliidae) during early development. *Poeciliid Research* 5 (1): 1-7.
- Mouludi-Saleh A, Eagderi S, Abbasi K, Nasri M (2022a). Validation of two sympatric fish species of Urmia chub, *Petroleuciscus ulanus* and Urmia bleak, *Alburnus atropatenae*, based on morphologic characters in Mahabad-Chai River. *Nova Biologica Reperta* 8 (4): 289-296.
- Mouludi-Saleh A, Eagderi S, Poorbagher H (2022b). Ichthyofauna of the Iranian part of the Sirvan River drainage with the first record of *Cobitis avicenna* and *Oxyneomacheilus euphraticus*. *International Journal of Aquatic Biology* 10 (3): 242-253. doi: 10.22034/ijab.v10i3.1639
- Mouludi-Saleh A, Eagderi S, Latif-Nejad S, Nasri M (2020a). The morphological study of transcaspian marinka (*Schizothorax pelzami*) in Harirud and Dasht-e Kavir basins using the geometric morphometric technique. *Nova Biologica Reperta* 7 (2): 185-191.
- Mouludi-Saleh A, Eagderi S (2021a). Morphological variations of *Oxyneomacheilus bergianus* (Derzhavin, 1934) in two inland water basins of Iran using geometric morphometric method. *Journal of Applied Ichthyological Research* 8 (4):71-76.
- Mouludi-Saleh A, Eagderi S (2021b). Habitat-associated morphological divergence of *Gasterosteus aculeatus* in the Southern Caspian Sea Basin. *Iranian Journal of Science and Technology, Transactions A: Science* 45 (1), 121-125. doi: 10.1007/s40995-020-01005-z
- Mouludi-Saleh A, Eagderi S, Cicek E, Sungur S (2020b). Morphological variation of Transcaucasian chub, *Squalius turcicus* in southern Caspian Sea basin using geometric morphometric technique. *Biologia* 75 (10): 1585-1590. doi: 10.2478/s11756-019-00409-6
- Mouludi-Saleh A, Eagderi S, Nahavandi R (2020c). The Application of the geometric morphometric technique in evaluating sexual dimorphism in the body shape of *Aphaniops Hormozensis* from Mehran and Shur Rivers (Hormozgan Basin). *Taxonomy and Biosystematics* 12 (43): 35-44.
- Mouludi-Saleh A, Eagderi S, Poorbagher H, Kazemzadeh S (2019). The effect of body shape type on differentiability of traditional and geometric morphometric methods: A case study of *Channa gachua* (Hamilton, 1822). *European Journal of Biology* 78 (2): 165-168. doi: 10.26650/EurJBiol.2019.0011
- Mouludi-Saleh A, Eagderi S, Poorbagher H, Shojaei D, Nasri M (2021). Phenotypic adaptation patterns in Abu Mullet, *Planiliza abu* using traditional and geometric morphometric methods in rivers of the Persian Gulf basin. *Fisheries Science and Technology* 10 (2): 141-150.
- Mousavi-Sabet H, Azimi H, Eagderi S, Bozorgi S, Mahallatipour B (2014). Growth and morphological development of guppy *Poecilia reticulata* (Cyprinodontiformes, Poeciliidae) larvae. *Poeciliid Research* 4 (1): 24-30.
- Mousavi-Sabet H, Eagderi S (2016b). First record of the convict cichlid, *Amatitlania nigrofasciata* (Günther, 1867) (Teleostei: Cichlidae) from the Namak Lake basin, Iran. *Iranian Journal of Ichthyology* 3 (1): 25-30. doi: 10.7508/iji.2016.01.003
- Mousavi-Sabet H, Eagderi S (2016a). *Garra lorestanensis*, a new cave fish from the Tigris River drainage with remarks on the subterranean fishes in Iran (Teleostei: Cyprinidae). *FishTaxa* 1 (1): 45-54.

- Mousavi-Sabet H, Vasil'eva ED, Vatandoust S, Vasil'ev VP (2011). *Cobitis faridpaki* sp. nova - a new spined loach species (Cobitidae) from the Southern Caspian Sea Basin (Iran). Journal of Ichthyology 51 (10): 925-931. doi: 10.1134/S003294521100055
- Mousavi-Sabet H, Ganjbakhsh B, Geiger MF, Freyhof J (2016b). Redescription of *Gobio nigrescens* from the Hari River drainage (Teleostei: Cyprinidae). Zootaxa 4114 (1): 71-80. doi: 11646/zootaxa.4114.1.4
- Mousavi-Sabet H, Vatandoust S, Esmaili HR, Geiger MF, Freyhof J (2015c). *Cobitis avicennae*, a new species of spined loach from the Tigris River drainage (Teleostei: Cobitidae). Zootaxa 3914 (5): 558-568. doi: 10.11646/zootaxa.3914.5.4
- Mousavi-Sabet H, Saemi-Komsari M, Doadrio I, Freyhof J (2019a). *Garra roseae*, a new species from the Makran region in southern Iran (Teleostei: Cyprinidae). Zootaxa 4671 (2): 223-239. doi: 10.11646/zootaxa.4671.2.3
- Mousavi-Sabet H, Vatandoust S, Azizi F (2015b). *Alburnoides tabarestanensis*, a new species of riffle minnow from the southern Caspian Sea basin in Iran (Actinopterygii: Cyprinidae). Aqua, International Journal of Ichthyology 21 (3-15): 144.
- Mousavi-Sabet H, Vatandoust S, Doadrio I (2015a). Review of the genus *Alburnoides* Jeitteles, 1861 (Actinopterygii, Cyprinidae) from Iran with description of three new species from the Caspian Sea and Kavir basins. Caspian Journal of Environmental Sciences 13 (4): 293-331.
- Mousavi-Sabet H, Vatandoust S, Fatemi Y, Eagderi S (2016a). Tashan Cave a new cave fish locality for Iran; and *Garra tashanensis*, a new blind species from the Tigris River drainage (Teleostei: Cyprinidae). FishTaxa 1 (3): 133-148.
- Mousavi-Sabet H, Vatandoust S, Geiger MF, Freyhof J (2019). *Paracobitis abrishamchiani*, a new crested loach from the southern Caspian Sea basin (Teleostei: Nemacheilidae). Zootaxa, 4545 (3): 375-388. doi: 10.11646/zootaxa.4545.3.3
- Mousavi-Sabet H, Sayyadzadeh G, Esmaili HR, Eagderi S, Patimar R et al. (2015d). *Paracobitis hircanica*, a new crested loach from the southern Caspian Sea basin (Teleostei: Nemacheilidae). Ichthyological Exploration of Freshwaters 25 (4): 339-346.
- Mousavi-Sabet H, Eagderi S (2015). *Paraschistura delvarii* spec. nov. - a new species of stone loach from the Persian Gulf basin, southern Iran (Teleostei: Nemacheilidae). Vertebrate Zoology 65 (3): 297-303.
- Mousavi-Sabet H, Vatandoust S, Roudbar AJ, Eagderi S (2015e). Taxonomic status of the genus *Paraschistura* (Teleostei: Nemacheilidae) in the Hari river basin, with re-validation of *P. turcomana*. Journal of Applied Biological Sciences 9 (3): 01-05.
- Mousavi-Sabet H, Eagderi S, Vatandoust S, Freyhof J (2021). Five new species of the sisorid catfish genus *Glyptothorax* from Iran (Teleostei: Sisoridae). Zootaxa 5067 (4): 451-484. doi: 10.11646/zootaxa.5067.4.1
- Mutlak F, Jawad L, Al-Faisal A (2017). *Atractosteus spatula* (Actinopterygii: Lepisosteiformes: Lepisosteidae): A deliberate aquarium trade introduction incidence in the Shatt al-Arab River, Basrah, Iraq. Acta Ichthyologica et Piscatoria 47 (2): 205-207. doi: 10.3750/AIEP/02143
- Namdariyanrad A, Mojazi Amiri B, Eagderi S, Benam S, Abedi M (2017). A comparative histological study of osmotic regulator organs in matured and pre-mature *Caspiomyzon wagneri* migrating to Shiroud River. Journal of Animal Research (Iranian Journal of Biology) 30 (2): 237-243.
- Naseka AM, Bogutskaya NG (2009). Fishes of the Caspian Sea: zoogeography and updated check-list. Zoosystematica Rossica 18 (2): 295-317.
- Nasri M, Eagderi S, Keivany Y, Farahmand H, Dorafshan S et al. (2018). Morphological diversity of *Cyprinion* Heckel, 1843 species (Teleostei: Cyprinidae) in Iran. Iranian Journal of Ichthyology 5 (2): 96-108. doi: 10.22034/iji.v5i2.265
- Nasri M, Eagderi S, Farahmand H (2016). Descriptive and comparative osteology of Bighead Lotak, *Cyprinion milesi* (Cyprinidae: Cypriniformes) from southeastern Iran. Vertebrate Zoology 66 (3): 251-260.
- Nasri M, Eagderi S, Farahmand H, Hashemzade-Segharloo I (2013). Body shape comparison of *Cyprinion macrostomum* (Heckel, 1843) and *Cyprinion watsoni* (Day, 1872) using geometric morphometric method. International Journal of Aquatic Biology 1 (5): 240-244. doi: 10.22034/ijab.v1i5.154
- Nelson JS, Grande TC, Wilson MV (2016). Fishes of the World. John Wiley & Sons.
- Nikmehr N, Eagderi S, Poorbagher H, Abbasi K (2021). Length-weight relationship and condition factor of three endemic fish species, *Ponticola bathybius*, *Neogobius caspius* and *Neogobius pallasii* (Perciformes: Gobiidae) from the Southern Caspian Sea basin, Iran. Ege Journal of Fisheries and Aquatic Sciences, 38 (4), 523-525. doi: 10.12714/egejfas.38.4.14
- Nikmehr N, Eagderi S, Poorbagher H, Farahmand H (2020). Taxonomic status of the *Turcinoemacheilus* populations (Nemachilidae) of the Gaveh River using the COI gene. Wetland Ecology 12 (2): 49-58.
- Nikmehr N, Eagderi S, Poorbagher H, Farahmand H (2019). Phylogenetic relationships of the genus *Turcinoemacheilus* in Iran using osteological characteristics. Taxonomy and Biosystematics 11 (40): 47-58.
- Radkhah AR, Eagderi S (2021). A brief review of the geographic ranges and ecological effects of three major invasive cyprinid species in Iran. Journal of Fisheries 9 (3): 93301-93301. doi: 10.17017/j.fish.305
- Radkhah AR, Eagderi S (2022). Ecological consequences of tilapia species on fish biodiversity of Iran and challenges arising from their introduction. Iranian Journal of Ichthyology 8 (4): 342-350. doi: 10.22034/iji.v8i4.776
- Radkhah AR, Eagderi S, Mouludi-Saleh A. (2022a). Morphological variation, length-weight relationship and condition factor of *Cabdio morar* (Hamilton, 1822) populations in southeast of Iran. Journal of Wildlife and Biodiversity 6 (1). doi: 10.5281/zenodo.6498906
- Radkhah AR, Eagderi S, Poorbagher H, Nowferesti H (2022b). Sexual dimorphism of Eastern mosquitofish (*Gambusia holbrooki* Girard, 1859) in Gamasiab River, Kermanshah Province. Nova Biologica Reperta 9 (1): 50-60.

- Rafiee G, Jouladeh, RA, Eagderi S (2017). Review on the Iranian members of the family Cichlidae (Actinopterygii) with first record of Nile Tilapia, *Oreochromis niloticus*. Journal of Fisheries 70 (1): 1-10.
- Razavi Pour P, Eagderi S, Poorbagher H (2014). Study of osteological characteristics of Tuini fish (*Capoeta damascina* Valenciennes, 1842) from Tigris basin. Journal of Applied Ichthyological Research 2 (3): 1-16.
- Soleimanian S, Amini Chermahini M, Eagderi S, Savari A (2021). Body shape changes of Shirbot, *Arabibarbus grypus* (Heckel, 1843) during early development using geometric morphometric method. Aquaculture Sciences 8 (2): 58-64.
- Saffari S, Poorbagher H, Eagderi S, Farahmand H (2021). Identifying the species of the genus *Alosa* using the scales, wavelet and outline analyses. Journal of Fisheries 74 (2): 315-324.
- Sahraeian MR, Eagderi S, Zibae A, Rafiei GR, Khomeirani R (2016). Allometric growth patterns and morphological development of the Common bream (*Abramis brama*) during early development under cultural condition. Journal of Animal Physiology and Development 2 (33): 13-22.
- Sayyadzadeh G, Esmaeili HR, Abbasi K, Coad BW (2015b). Re-validation of *Gonorhynchus adiscus* and *G. diplochilus* (Teleostei: Cyprinidae) using morphological and molecular data. Zoology in the Middle East 61 (4): 349-361. doi: 10.1080/09397140.2015.1095517
- Sayyadzadeh G, Esmaeili HR, Eagderi S, Jouladeh-Roudbar A, Masoudi M et al. (2017). Re-description of *Oxynoemacheilus longipinnis* from the Persian Gulf basin (Teleostei: Nemacheilidae). Zoology in the Middle East 63 (3): 228-238. doi: 10.1080/09397140.2017.1349243
- Sayyadzadeh G, Esmaeili HR, Freyhof J (2015a). *Garra mondica*, a new species from the Mond River drainage with remarks on the genus *Garra* from the Persian Gulf basin in Iran (Teleostei: Cyprinidae). Zootaxa 4048 (1): 75-89. doi: 10.11646/zootaxa.4048.1.4
- Sayyadzadeh, G, Esmaeili HR (2020). *Oxynoemacheilus marunensis*, a new loach species from the Persian Gulf basin with remarks on *O. frenatus* (Teleostei: Nemacheilidae 4885 (2): 189-206. doi: 10.11646/zootaxa.4885.2.2
- Sayyadzadeh G, Eagderi S, Esmaeili HR (2016). A new loach of the genus *Oxynoemacheilus* from the Tigris River drainage and its phylogenetic relationships among the nemacheilid fishes (Teleostei: Nemacheilidae) in the Middle East based on mtDNA COI sequences. Iranian Journal of Ichthyology 3 (4): 236-250. doi: 10.22034/iji.v3i4.205
- Sayyadzadeh G, Esmaeili HR, Eagderi S (2019a). Taxonomic status of the loaches *Paracobitis vignai* and *P. rhadinaea* (Teleostei: Nemacheilidae) in Iran. Zoology in the Middle East 65 (3): 221-229. doi: 10.1080/09397140.2019.1586143
- Sayyadzadeh G, Teimori A, Esmaeili HR (2019b). *Paraschistura kermanensis*, a new stone loach species from southeastern Iran (Teleostei: Nemacheilidae). Zootaxa 4638 (4): 571-583. doi: 10.11646/zootaxa.4638.4.7
- Schwarzer J, Shabani N, Esmaeili HR, Mwaiko S, Seehausen O (2016). Allopatric speciation in the desert: diversification of cichlids at their geographical and ecological range limit in Iran. Hydrobiologia 791 (1): 1-15. doi: 10.1007/s10750-016-2976-3
- Teimori A, Esmaeili HR, Sayyadzadeh G, Zarei N, Gholamhosseini A (2015). Molecular systematics and distribution review of the endemic cyprinid species, Persian chub, *Acanthobrama persidis* (Coad, 1981) in Southern Iran (Teleostei: Cyprinidae). Molecular Biology Research Communications 4 (4): 189-206.
- Teimori A, Esmaeili HR (2020). Axial skeleton morphology of the Western Palearctic aphaniid fishes (Teleostei: Cyprinodontiformes; Family: Aphanidiidae). Acta Zoologica 103 (3): 282-296. doi:10.1111/azo.12370
- Teimori A, Esmaeili HR, Reichenbacher B (2011). *Aphanius farsicus*, a replacement name for *A. persicus* (Jenkins, 1910) (Teleostei, Cyprinodontidae). Zootaxa 3096 (1): 53-58. doi: 10.11646/zootaxa.3096.1.5
- Teimori A, Esmaeili HR, Gholami Z, Zarei N, Reichenbacher B (2012). *Aphanius arakensis*, a new species of tooth-carp (Actinopterygii, Cyprinodontidae) from the endorheic Namak Lake basin in Iran. ZooKeys 215: 55-76. doi:10.3897/zookeys.215.1731
- Teimori A, Motamedi M, Hesni MA (2017). Translocation and new geographical distribution of the invasive Redbelly Tilapia, *Coptodon zillii* (Gervais, 1848) (Teleostei: Cichlidae) in southern Iran. Check List 13: 1. doi: 10.15560/13.1.2051
- Teimori A, Esmaeili HR, Hamidan N, Reichenbacher B (2020). Comment on "A proposal for a new generic structure of the killifish family Aphanidiidae, with the description of *Aphaniops teimorii* (Teleostei: Cyprinodontiformes)" by Jörg Freyhof & Baran Yoğurtçuoğlu, Zootaxa (July 2020) 4810 (3): 421-451. FishTaxa 17: 15-16
- Turan D, Kaya C, Aksu İ, Bektaş Y (2022). Paracapoeta, a new genus of the Cyprinidae from Mesopotamia, Cilicia and Levant (Teleostei, Cypriniformes). Zoosystematics and Evolution 98 (2): 201-212.
- Vasil'eva ED, Mousavi-Sabet H, Vasil'ev VP (2015). *Ponticola iranica* sp. nov. (Actinopterygii: Perciformes: Gobiidae) from the Caspian Sea basin. Acta Ichthyologica et Piscatoria 45 (2): 189-197. doi: 10.3750/AIP2015.45.2.09
- Vatandoust S, Eagderi S (2015). *Paraschistura ilamensis*, a new species of loach from the Tigris River drainage (Teleostei: Nemacheilidae). International Journal of Aquatic Biology 3 (3): 177-182. doi: 10.22034/ijab.v3i3.95
- Zamani-Faradonbe M, Keivany Y, Dorafshan S, Zhang E (2021b). Two new species of *Garra* (Teleostei: Cyprinidae) from western Iran. Ichthyological Exploration of Freshwaters 30 (1): 249-270.
- Zamani-Faradonbe M, Zhang E, Keivany Y (2021a). *Garra hormuzensis*, a new species from the upper Kol River drainage in the Persian Gulf basin (Teleostei: Cyprinidae). Zootaxa 5052 (3): 380-394. doi: 10.11646/zootaxa.5052.3.4
- Zarei F, Esmaeili HR, Abbasi K, Sayyadzadeh G, Eagderi S et al. (2021b). Genealogical concordance, comparative species delimitation, and the specific status of the Caspian pipefish *Syngnathus caspius* (Teleostei: Syngnathidae). Marine Ecology 42 (1): e12624. doi: 10.1111/maec.12624

- Zarei F, Esmaili HR, Schliwen UK, Abbasi K, Sayyadzadeh G (2021). Mitochondrial phylogeny, diversity, and ichthyogeography of gobies (Teleostei: Gobiidae) from the oldest and deepest Caspian sub-basin and tracing source and spread pattern of an introduced *Rhinogobius* species at the tricontinental crossroad. *Hydrobiologia* 848 (6): 1267-1293. doi: 10.1007/s10750-021-04521-0
- Zarei F, Esmaili HR, Schliwen UK, Abbasi K, Sayyadzadeh G (2021). Mitochondrial phylogeny, diversity, and ichthyogeography of gobies (Teleostei: Gobiidae) from the oldest and deepest Caspian sub-basin and tracing source and spread pattern of an introduced *Rhinogobius* species at the tricontinental crossroad. *Hydrobiologia* 848 (6): 1267-1293. doi: 10.1007/s10750-021-04521-0
- Zarei F, Esmaili HR, Kovačić M, Schliwen UK, Abbasi K (2022). *Ponticola hircaniaensis* sp. nov., a new and critically endangered gobiid species (Teleostei: Gobiidae) from the southern Caspian Sea basin. *Zootaxa* 5154 (4): 401-430. doi: 10.11646/zootaxa.5154.4.1
- Zareian H, Esmaili HR (2017). Mitochondrial phylogeny and taxonomic status of the *Capoeta damascina* species group (Actinopterygii: Cyprinidae) in Iran with description of a new species. *Iranian Journal of Ichthyology* 4: 231-269. doi: 10.22034/iji.v4i3.239
- Zareian H, Esmaili HR, Gholamhosseini A, Japoshvili B, Özüluğ M et al. (2018). Diversity, mitochondrial phylogeny, and ichthyogeography of the *Capoeta capoeta* complex (Teleostei: Cyprinidae). *Hydrobiologia* 806 (1): 363-409. doi: 10.1007/s10750-017-3375-0
- Zareian H, Esmaili HR, Freyhof J (2016). *Capoeta anamisensis*, a new species from the Minab and Hasan Langhi River drainages in Iran (Teleostei: Cyprinidae). *Zootaxa* 4083 (1): 126-142. doi: 10.11646/zootaxa.4083.1.7
- Zareian H, Esmaili HR, Gholamhosseini A, Sayyadzadeh G (2013). New records and geographical distribution of *Alburnus hohenackeri* Kessler, 1870 (Teleostei: Cyprinidae) in Iran. *Check List* 9: 829-831. doi:10.155660/9.4.829