

Tropical seamounts as stepping-stones for coral reef fishes: range extensions and new regional distributions from mesophotic ecosystems in the Coral Sea, Australia

Gemma F Galbraith (✉ gemma.galbraith@jcu.edu.au)

James Cook University <https://orcid.org/0000-0001-9888-2918>

Benjamin J Cresswell

James Cook University <https://orcid.org/0000-0003-0191-8828>

Eva C McClure

James Cook University <https://orcid.org/0000-0002-3073-1264>

Andrew S Hoey

James Cook University <https://orcid.org/0000-0002-4261-5594>

Research Article

Keywords: seamounts, coral reef fishes, Coral Sea, mesophotic coral ecosystems, biodiversity, ROVs, BRUVs

Posted Date: September 15th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3342963/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations:

The authors have no competing interests to declare that are relevant to the content of this article

Tropical seamounts as stepping-stones for coral reef fishes: range extensions and new regional distributions from mesophotic ecosystems in the Coral Sea, Australia

Galbraith, G.F.^{*1,2} Cresswell, B.C.^{1,2}, McClure, E.C.^{1,2} and Hoey, A.S.^{1,2}

*Corresponding author gemma.galbraith@jcu.edu.au

¹Marine Biology and Aquaculture, College of Science and Engineering, and

²ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland 4811, Australia

Author ORCID ID: G.F.G 0000-0001-9888-2918; B.J.C 0000-0003-0191-8828; E.C.M 0000-0002-3073-1264; A.S.H 0000-0002-4261-5594

Key words: seamounts, coral reef fishes, Coral Sea, mesophotic coral ecosystems, biodiversity, ROVs, BRUVs

Abstract

Seamounts and remote oceanic islands provide valuable natural laboratories in which to study patterns and processes in marine biodiversity. A central hypothesis arising from studies of these systems is the ecological function of seamounts as stepping stones for dispersal and population connectivity. Evidence of this mechanism exists for a range of taxa, including coral reef fishes, but is still lacking from many tropical seamounts in remote regions. In this study, we used remotely operated vehicles and baited remote underwater video systems to survey fish and benthic communities between 1 – 100m on seamounts in the Coral Sea Marine Park (CSMP), Australia. We found evidence to support the stepping stone model of ecological connectivity from new observations of sixteen coral reef fishes which have previously not been recorded by quantitative surveys in the region. The widespread distribution of many of these species throughout the full latitudinal extent of the CSMP suggests that there is greater connectivity between mesophotic habitats in the Coral Sea and surrounding biogeographic regions than previously known. We also found a wide variety of mesophotic habitats and recorded significant depth range extensions for seventy-eight fishes in these habitats. This further highlights the potential role of increased habitat area and heterogeneity in a stepping stone effect throughout the region. Four of the fish occurrence records represent significant range extensions into the Coral Sea from adjacent biogeographic regions and thirteen fishes recorded by this study in the CSMP are not known from the neighbouring Great Barrier Reef, despite its close proximity. Although the Coral Sea remains relatively understudied, these findings suggest that larger-scale models of marine biogeography are relevant to communities in the region, particularly at mesophotic depths. Given the extent and the spatial arrangement of seamounts in the Coral Sea, our findings emphasise that the region is an important link between the centre of marine biodiversity in

the Coral Triangle and the Southwest Pacific. Greater mesophotic sampling effort and genetic studies are necessary to understand the nature of connectivity and, to establish the role of regional seamount chains, like the Coral Sea reefs, in broader marine biogeographic processes.

Introduction

The Coral Sea, in the southwest Pacific Ocean, is the second largest tropical marginal sea on earth and is characterised by a complex bathymetry and diverse seascape of distinct marine habitats (Ceccarelli et al. 2013; McKinnon et al. 2014). These habitats include the deep sea, submerged banks, canyons, island chains and large oceanic coral reef systems atop of seamounts rising from deep waters (up to 3000m) (Davies et al. 1989; Bridge et al. 2019). Ecologically, seamounts have been variously considered as either stepping stones for dispersal that can promote regional connectivity (Hubbs 1959; Rowden et al. 2010; Mazzei et al. 2021) and conversely, as isolated islands that can give rise to unique ecological communities (Richer de Forges et al. 2000; Hobbs et al. 2008, 2012; McClain et al. 2009). Generalities for these contrasting hypotheses remain unclear and it is broadly agreed that the ecological function of seamounts in marine connectivity depends on multiple factors including geomorphology, depth, spatial isolation, oceanographic processes and taxa-specific dispersal capabilities (McClain 2007; Clark and Bowden 2015; Miller and Gunasekera 2017; Pinheiro et al. 2017). Despite wide recognition as important marine habitats, in both tropical and temperate oceans, seamounts remain one of the least explored and studied marine biomes on earth (Clark et al. 2010b; Yesson et al. 2011). Consequently, many regional seamount chains require considerably greater sampling effort to establish patterns of biodiversity and

the mechanisms driving ecological connectivity with wider biogeographic regions (Rogers 2018).

Most seamount reef systems in the Coral Sea occur in Australia's Exclusive Economic Zone (EEZ) and are managed as the Coral Sea Marine Park (CSMP). Together with the French Natural Park of the Coral Sea (Le Parc Naturel de la Mer de Corail), the Coral Sea possesses the largest combined protected area in the world (Director of National Parks 2018). In Australia's CSMP, over 30 individual reef systems are spread across 22 degrees of latitude on the Queensland and Marion Plateaux and constitute ~24,000km² of shallow water (< 30m) emergent coral reef habitat (Bridge et al. 2019). The CSMP is bordered by major global marine biodiversity and productivity hotspots; Australia's Great Barrier Reef (GBR) to the west, the Coral Triangle (specifically, Papua New Guinea and the Solomon Islands) to the north, Vanuatu and New Caledonia to the east and the Tasman Sea to the south (Figure 1). It is therefore not surprising that the CSMP supports a high diversity of reef fish (>600 species) and high abundance and biomass of sharks and other large predatory fishes (Ceccarelli et al. 2013; Stuart-Smith et al. 2013; Hoey et al. 2022). Shallow water reef habitats in the CSMP have also been shown to support unique coral and reef fish communities that are distinct from those of the adjacent GBR to the west and share more similarities with those found in New Caledonia to the east (Ceccarelli et al. 2013; Hoey et al. 2020). Oceanographic processes and historic environmental conditions explain a significant proportion of the evolutionary processes driving genetic connectivity and biodiversity patterns within Coral Sea populations and between surrounding regions (Ceccarelli et al. 2013; Kessler and Cravatte 2013; Payet et al. 2022). However, the extent and spatial arrangement of coral reef habitat on seamounts in the Coral Sea is also likely a key component of ecological connectivity in this region and with the wider Central Pacific Ocean.




Fig.1 a) Location of the Coral Sea relative to major marine biodiversity regions adapted from dissimilarity analysis between species occurrence data by Kulbicki et al. (2013) and b) Bathymetry of the Coral Sea from the same region bounded in red as in Figure 1a (data from Beaman 2010). Most seamounts in the Coral Sea Marine Park can be seen in a chain on the Queensland plateau, offshore to the east from the Great Barrier Reef. The Southwest Pacific region in the Coral Sea extends east to New Caledonia and north to the boarder of the Coral Triangle and Papua New Guinea. The northern seamounts of the Coral Sea extend to the boarder with the Central Indo-Pacific region, west through the Torres Strait. Arrows indicate conceptual borders and connectivity with adjacent biogeographical regions.

Although recent large-scale monitoring efforts (e.g. Hoey et al. 2020, 2022) and baseline surveys (e.g. Ayling and Ayling 1985; Oxley et al. 2004; Edgar et al 2015) have established quantitative ecological data for shallow water coral reefs in the CSMP, the remote nature of reefs in the Coral Sea mean they remain poorly documented compared to those in the surrounding GBR or the Coral Triangle (Ceccarelli 2010). Additionally, restrictive institutional and occupational diving regulations in Australia have led to a paucity of coral reef research conducted in depths below 20m (Eyal et al. 2021). Mesophotic Coral Ecosystems (MCEs) are defined as light-dependent coral reef communities in depths of 30–150 m (Loya et al. 2016). The exceptionally clear oligotrophic waters of the Coral Sea allow light penetration to considerable depths and several exploratory studies have confirmed the presence of MCEs in the region to depths of up to 125m (Sarano and Pichon 1988; Bongaerts et al. 2011; Muir et al. 2015; Englebert et al. 2015, 2017). At these depths, the complex bathymetries of the Coral Sea seamounts are also highly variable, both among and within individual reef systems (Harris et al. 2003; Beaman 2010). Some rise as vertical walls to the surface (e.g. Bougainville and Osprey reefs) while others have less abrupt slopes and many possess near-horizontal areas along flanks and shelves (e.g. Holmes Reefs and East Diamond Islet). On more moderate slopes, where hard substrate is present and light availability is optimal, mesophotic depths can support high percentage cover of photosynthetic habitat-forming taxa (Pérez-Rosales et al. 2022). Hard and soft corals, macroalgae, sponges and large benthic foraminifera are all important constituents of MCEs and in turn provide habitats for other marine organisms (Slattery and Lesser 2012; Lesser et al. 2018). This said, the extent of MCEs or other important benthic habitats between 0 -150m have not been quantified or confirmed at most Coral Sea reefs. Further, the few exploratory studies of MCEs in the CSMP are restricted to a select number of reefs and have all focused on benthic organisms, specifically scleractinian corals. To date, there have been no quantitative surveys

of fish communities in the CSMP, or the Coral Sea more broadly, at depths below 20m. The ecology and biodiversity of mesophotic coral reef fish communities in the Coral Sea is therefore scarcely known.

Two major global marine biogeographical regions meet in the Coral Sea; the Southwest Pacific and the Central Indo-Pacific, the latter of which includes the Coral Triangle (Kulbicki et al. 2013) (Figure 1b). There has been increasing recognition that “peripheral habitats”, typically isolated archipelagos, marginal seas and seamounts in regions surrounding major biodiversity centres, can also export biodiversity and connect biogeographical regions, rather than function only as isolated population sinks (Bowen et al. 2013; Simon et al. 2022). Over the past 10 - 20 years, updated species checklists from locations in the Southwest Pacific have listed occurrence records for several Indo-Pacific fishes not previously reported from these areas (Randall et al. 2003; Fricke et al. 2011b, a). Additionally, aquarist collections from one location in the central Coral Sea Marine Park (Holmes Reefs) have also reported deep-water fishes typically known only from either the Central Indo-Pacific or Central Pacific regions (Fenton Walsh, pers. com). Large-scale connectivity patterns are not well established in the Coral Sea but these observations suggest greater latitudinal connectivity between the Central Indo-Pacific and Western Pacific region through the Coral Sea than may currently be known. This may be particularly true for mesophotic species which are not recorded in shallow water reef surveys but utilise deeper reef habitats.

In this study, we used Remotely Operated Vehicles (ROVs) and Baited Remote Underwater Video systems (BRUVs) to survey fish and benthic communities in the Coral Sea Marine Park at depths between 1 – 100m. Here we present the first occurrence records of sixteen species from quantitative ecological surveys of coral reef fishes from seamounts in the Coral

Sea. We also compare fish species richness between shallow water monitoring surveys and ROV/BRUV surveys and present a range of mesophotic habitats found in the CSMP during this study. We discuss these observations in the context of regional connectivity, biogeography and the importance of seamounts in tropical coral reef seascapes.

Methods

Four separate voyages in the Coral Sea Marine Park were undertaken in 2021, 2022 and 2023. Remotely Operated Vehicles (ROVs) and Baited Remote Underwater Video systems (BRUVs) were used to survey fish communities and benthic habitats at 17 reefs and between 1 – 100m (Figure 2). Single camera BRUVS were deployed for 1 hour following the standard operating procedures outlined in Langlois et al. (2020). ROVs (BlueRobotics BlueRov2) were fitted with a forward-facing stereo-video system (SVS) to enable length estimates to be made. SVS cameras (Paralenz + or GoPro Hero 8 systems) were calibrated prior to surveys using the software CAL and associated calibration method (SeaGIS Pty Australia). ROV transects, each 30 x 5m, were conducted parallel to the reef contour using a timed swim method (ROV speed 0.2m/s for 2 minutes 30 seconds) at a constant depth (+/- 2m). For each ROV deployment, two transects were conducted within each 10m depth band, starting with the deepest transects and working upwards to the shallows. Sufficient horizontal and vertical separation was attained between transects and between depth bands by known speed and time of the ROV. Three GoPro Hero8 cameras inside deep-rated T-housings were mounted facing outwards left and right and downwards on the ROV. These cameras were set to the timelapse photo function (1 photo every 10 seconds), capturing an image of the benthos every ~2m (15 photos per transect). For fish community data, videos were interrogated in EventMeasure (SeaGIS, Pty Australia) and each individual fish entering the frame (BRUV) or transect field-

of-view (ROV) identified to the lowest possible taxonomic resolution. For ROV stereo-video footage, length estimates were also made of individual fishes in each transect (fork length) using the software EventMeasure Stereo (SeaGIS, Pty Australia). All major habitats at each reef were surveyed (outer reef, lagoon, back reef and reef passes between the lagoons and leeward outer reef), with BRUVs mostly conducted in lagoons and inner reef areas due to the steep sides of many Coral Sea seamount reefs.

We compared species records from shallow underwater visual census monitoring surveys conducted by divers on SCUBA (1 - 10m depth) at the same reefs during the same voyages (Hoey et al. 2021, 2022; Galbraith et al. 2022), as well as fish species records from previous surveys of the same reefs conducted by the Reef Life Survey Foundation (Edgar and Stuart-Smith 2014). Occurrence records and locations for species only recorded by BRUV and ROV surveys were obtained from the Ocean Biodiversity Information System (OBIS 2023a) and Global Biodiversity Information Facility (GBIF.org 2023a) and were cross-referenced with other online databases; Atlas of living Australia (ALA 2023), Fishes of Australia (Bray and Gomon 2023), Reef Life Survey (RLS 2023), fishbase (Froese and Pauly, 2023), CSIRO Codes for Australian Aquatic Biota (Rees et al. 2023) and the Australian Faunal Directory (ABRS 2020) as well as taxonomic experts. Known depth records for all fishes recorded were also extracted from fishbase using the *rfishbase* package (Boettiger et al. 2012) and compared to depths at which they were observed by ROV and BRUV surveys. To illustrate notable range extensions for three species, previous extant range extents were calculated as Extent of Occurrence (EOO) based on records obtained from the aforementioned databases and plotted in R (R Core Development Team 2023) using the packages *ggmaps* (Kahle and Wickham 2013) and *maps* (Becker et al. 2022).




Fig.2 Map of the Coral Sea, East Coast Australia and survey sites from this study. All sites in the CSMP surveyed by BRUV, ROV and shallow diver surveys are labelled by text. Locations with yellow stars indicate reefs with new locality observations for coral reef fishes. The Holmes Reefs, where several of these species have been previously noted by aquarium fish collectors, are marked by a red circle. Reefs marked with a black circle indicate locations where no new observations of reef fishes were recorded.

Results

A total of 274 ROV transects and 108 BRUV drops were analysed and cumulatively recorded 360 species of fishes from 41 families (Online Resource 1). Compared to available data from shallow (<10m) underwater visual census surveys, 50 additional species were recorded by ROV and BRUV at depths between 1 – 100m. Thirteen of these 50 species were previously only known from a single location in the central Coral Sea through observations and/or collections by aquarium fish collectors at the Holmes Reefs (F.Walsh pers. Com). Outside of these collections, these are the first observations of these species from quantitative fish community surveys in the Coral Sea Marine Park, and broader Coral Sea region (Table 1). A further three species have single records from Boot (*Anampses melanurus*, Atlas of Living Australia 2023), Frederick (*Mulloidichthys pfluegeri*, Atlas of Living Australia 2023) and Osprey (*Valenciennea helsdingeni*, Australian Faunal Directory 2023) reefs respectively but were recorded in our surveys at 11 other reefs (Table 1). These records increase their range extent in the region by between 4 -12 degrees of latitude. Of the 360 fish species recorded by BRUV and ROV surveys in this study, 158 (44%) were observed at depths below their reported maximum known depth as listed on the fishbase database (Online Resource 2). 34 of these depth record extensions are for species observed at depths greater than double their previously reported maximum depth.

- 1 **Table 1** New occurrence records of 16 fishes from mesophotic depths on seamount reefs in the Coral Sea. Previous range extents and
- 2 distributions are based on georeferenced occurrence data from online databases (see methods) and personal communication with taxonomic
- 3 experts and aquarium trade collectors. Reported depth ranges were obtained from fishbase and depth of observations from this study presented
- 4 for comparison.

Species	Family	Known range	New Coral Sea records	Reported depth range (fishbase)	Observed depth (This study)
<i>Hoplostethus randalli</i> ¹ (Allen, Erdmann & Hamilton, 2010)	Malacanthidae	Indo West Pacific: Indonesia, Philippines, Palau, Yap Is. and the Solomon Islands.	Lihou, East Diamond, Boot, Ashmore Reefs	30 -85m	70m – 85m
<i>Cephalopholis polleni</i> ¹ (Bleeker 1868)	Serranidae	Indian Ocean and Western Pacific: Comoros Islands to the Line Islands and French Polynesia. It has not been recorded from continental Australia or the major islands of Indonesia. In Australia it has been recorded from Christmas Island and Cocos (Keeling) Island.	Osprey Reef	30 -120m	91m
<i>Pseudanthias flavicauda</i> ¹ , (Randall and Pyle 2001)	Serranidae	Western-central Pacific: New Caledonia, Fiji, Tonga, Vanuatu and Tahiti.	Osprey, Lihou and Bougainville Reefs	30 – 61m	92m
<i>Liopropoma</i> 'sp. Yellow tail' ¹	Serranidae	Currently undescribed but known to have a wide distribution in the Western-Pacific (Tea, pers com).	Osprey Reef	Na	70-94m
<i>Bodianus paraleucosticticus</i> ¹ (Gomon 2006)	Labridae	Western Pacific: Cook Islands and Papua New Guinea; New Caledonia.	Osprey and Lihou Reefs	25 -115m	70 – 90m
<i>Hoplostethus marcosi</i> ¹	Malacanthidae	Western Pacific Ocean; Indonesia, Philippines,	Lihou, East	18 – 80m	60 – 85m

(Burgess 1978)		Palau, Papua New Guinea and the Solomon Islands. One report from Great Barrier Reef shelf break with Coral Sea (Sih et al. 2017).	Diamond, Ashmore Reefs		
<i>Abalistes filamentosus</i> ¹ (Matsuura & Yoshino, 2004)	Balistidae	Indo-West Pacific: Southern Japan to northern and north-western Australia, and New Caledonia.	Chilcott Reef	60-180 m	65 – 85m
<i>Pogonoperca punctata</i> ¹ (Valenciennes 1830)	Serranidae	Indo-Pacific: Comoros to the Line, Marquesan and Society islands, north to southern Japan, south to New Caledonia. Indian Ocean: southern Natal, South Africa and the Australian territories of Christmas Island and Cocos (Keeling) Islands. In Australia from South of Evans Shoal, Northern Territory.	East Diamond Islet	10 – 216m	53m
<i>Valenciennea helsdingeni</i> ² (Bleeker 1858)	Gobiidae	Indo-West Pacific: southern Red Sea and East Africa to Indonesia, north to southern Japan, south to the Great Barrier Reef. Solomon Islands. Known from Osprey Reef in the Coral Sea.	East Diamond Islet	1 - 45m	52m
<i>Xanthichthys auromarginatus</i> ¹ (Bennett 1832)	Balistidae	Indo Pacific: From Mauritius east to northern Australia to Hawaii and Society Islands. North to Southern Japan. Northern Great Barrier Reef (Escape Reef), in Christmas and Cocos (Keeling) Islands and Lord Howe Island.	Willis Island, Osprey, Bougainville, Mellish, Ashmore, Boot and Wreck Reefs	8 – 140m	40 – 95m
<i>Anampses melanurus</i> ³ (Bleeker 1857)	Labridae	Pacific Ocean: Indonesia to the Marquesas and Society Islands, north to Ryukyu Islands, south to Scott Reef Western Australia. Range extends to Easter Island.	Ashmore, East Diamond, Osprey, Holmes, Wreck Reefs	15 -40m	72m
<i>Pyronotanthias cf aurulentus</i> ¹ (Randall and McCosker 1982)	Serranidae	Eastern Central Pacific: Line Islands	Osprey and Bougainville Reefs	? – 52m	92m
<i>Genicanthus bellus</i> ¹	Pomacanthidae	Western pacific; Tahiti, Guam, Palau, Tonga,	Osprey and	25 – 100m	66 – 82m

(Randall 1975)		the Cook Islands, the Marshall Islands, the Philippines, southern Japan and southern Indonesia. Indian Ocean of the Cocos (Keeling) Islands and Christmas Island.	Bougainville Reefs		
<i>Pycnocrhromis leucura</i> ¹ (Gilbert 1905)	Pomacentridae	Indo-west-central Pacific, including: Madagascar, Mascarenes, Réunion, Mauritius, Andamans, Indonesia, Hawaiian Islands, Marquesas Islands and Gambier Islands, southern Japan and Ryukyu Islands, eastern Indonesia, New Caledonia.	Osprey Reef	20 -120m	84m
<i>Cirrhilabrus roseafascia</i> ¹ (Randall and Lubbock 1982)	Labridae	Western Central Pacific: New Caledonia, Philippines, Fiji, Palau, Samoa and Vanuatu. Australia: Myrmidon region of the GBR (Sih et al 2017).	Osprey Reef, East Diamond Islet	30 – 155m	94m
<i>Mulloidichthys pfluegeri</i> ⁴ (Steindachner 1900)	Mullidae	Indo-West Pacific: Reunion, Hawaiian, Marquesan and Society Islands. Eastern Indonesia, north to Southern Japan and south to Tonga. Known from the GBR in Australia.	Chilcott, Flinders, Herald, Holmes, Kenn, Lihou, Willis, Wreck Reefs	30 -110m	40 – 67m

5 Superscript numbers next to species names indicate observations from three other single CSMP locations ¹Holmes Reefs (F.Walsh Pers.com)

6 ²Osprey Reef (Australian Faunal Directory 2023), ³Boot Reef (Atlas of Living Australia 2023), ⁴Frederick Reef (Atlas of Living Australia 2023).

Four species recorded by ROV in the CSMP represent notable range extensions based on previous global occurrence records. *Hoplolatilus randalli* (Allen et al. 2010), a relatively newly described species of tile fish (family Malcanthidae), is currently known only from Indonesia, the Philippines, Palau, Yap and the Solomon Islands (Froese and Pauly 2023). A total of eight individuals were recorded at reefs spanning the northern and central CSMP (Ashmore and Lihou Reefs and East Diamond Islet), all at depths below 70m (Table 1, Figure 3a). We mostly observed *H. randalli* in pairs beside large mounds of rubble, apparently built by the fish over their burrows. The observations from this study are the southernmost occurrence records for the species and expand the known extent of occurrence for *H. randalli* by almost 10 degrees of latitude. *Cephalopholis polleni* (Bleeker 1868) was previously only known in Australian waters from the Cocos (Keeling) and Christmas Islands in the Indian Ocean. Elsewhere, *C. polleni*, (family Serranidae), occurs at scattered localities on oceanic islands across the Indian Ocean and wider Indo-Pacific (Bray 2023). In the Coral Sea, this study recorded one individual *C. polleni* at 97m under a ledge at Osprey Reef by ROV survey (Fig 3b) and it has also been collected at Holmes Reef (Fenton Walsh pers. com). These Coral Sea records extend the southern range of *C. polleni* in the Southwest Pacific by 6 degrees of latitude. *Pseudanthias flavicauda* (Randall and Pyle 2001) was recorded by ROV survey at Osprey and Bougainville reefs in the northern Coral Sea. *P. flavicauda* (family Serranidae) is known from the Central and South West Pacific (Bray 2022; Froese and Pauly 2023), and recently from Tonga (Fricke et al 2011) and New Caledonia (Fricke and Kulbicki 2007). We observed abundant schools of *P. flavicauda* between 80 -100m at Osprey and Bougainville Reefs and although also collected from the Holmes Reefs (Fenton Walsh pers. com), the observations from this study are the most western records for this species and the most northern extent in the Coral Sea (Figure 3c). *Bodianus paraleucosticticus* (Gomon 2006) was found in ROV surveys at Lihou and Osprey Reefs at depths between 70 – 90m.

Together with collections from Holmes Reefs, these new observations of *B. paraleucosticticus* (family Labridae) extend the previously reported distribution west from New Caledonia and south from Papua New Guinea into the Coral Sea.




Fig.3 Current extent of occurrence plotted as colored hulls for a) *Hoplolatilus randalli* and b) *Cephalopholis polleni* c) *Pseudanthias flavicauda* and d) *Bodianus paraleucosticticus*.

Occurrence data were obtained from OBIS (2023b,c,d,e) and GBIF (2023b,c,d,e). New observations of each species from the Coral Sea by this study are represented by yellow stars.

In addition to range extensions, we found twelve other species, previously only known from a single location in the Coral Sea, at multiple other reefs in ROV and BRUV surveys. None of these species have previously been recorded in shallow water community surveys in the Coral Sea Marine Park and most are deep-water specialists occurring below depths of 50m (Figure 4). Three of these species have been recorded on the GBR but were previously not confirmed to be widely distributed throughout the Coral Sea.




Fig.4 Twelve species of coral reef fish recorded by ROV and BRUV surveys in the Coral Sea Marine Park (CSMP) between depths of 50 to 100 m. All are previously known from one reef location in the CSMP but are reported here from multiple other reefs spanning the full latitudinal extent of the CSMP. a) *Pogonoperca punctata* b) *Pycnochromis leucura* c) *Valenciennaea helsdingeni* d) *Abalistes filamentosus* e) *Liopropoma* “sp. yellow tail” f) *Xanthichthys auromarginatus* g) *Mulloidichthys pfluegeri* h) *Anampses melanurus* i) *Cirrhilabrus roseafascia* j) *Genicanthus bellus* k) *Hoplolatilus marcosi* and l) *Pyronotanthias cf. aurulentus*.

The presence of mesophotic coral ecosystems was confirmed at all 17 of the reefs surveyed by ROV and BRUV. Several sites possessed remarkable hard coral cover (preliminary estimates ~70 -85%) at depths between 50 – 100m (Figure 5a and d). Multiple other non-coral dominated mesophotic habitats were also found including *Cycloclypeus* beds (large benthic foraminifera) (Figure 5b), octocoral dominated walls (Figure 5c), seagrass (Figure 5e) and extensive *Halimeda* meadows (Figure 5f).




Fig.5 Varied mesophotic coral ecosystems and other mesophotic habitats in the Coral Sea a) areas of high coral cover, 77m, Lihou Reef b) *Oxychellinus orientalis* and *Cirrhilabrus bathyphilus* on a dense bed of *Cycloclypeus* at 82m, East Diamond Islet c) *Oxycheilinus orientalis* on a steep wall with soft corals and gorgonians, 94m, Osprey Reef d) High abundance of fishes with high and complex coral cover, 57m Bougainville Reef e) seagrass (*Halophila gradients*) at 43m East Diamond Islet and f) Dense *Halimeda* meadows at 67m, Lihou Reef.

Discussion

The new occurrence records presented here span a considerable latitudinal gradient and provide evidence of more widespread distributions for multiple fishes at mesophotic depths in the region than previously known. These observations support the stepping stone model of ecological connectivity for seamounts within the Coral Sea and between certain neighbouring biogeographic regions. The occurrence of several fishes in the Coral Sea which were previously only known from either the Indo-Pacific region or South and Western Central Pacific also aligns with other ecological hypotheses explaining larger-scale patterns in marine biodiversity. “The biodiversity feedback” theory proposes that peripheral regions actively contribute to the export of taxa and lineages back to biodiversity hotspots, rather than only acting as sink populations (Bowen et al. 2013). Range expansions into the Coral Sea from both the Central Indo-Pacific (e.g. *H. randalii*) and the western Central Pacific (e.g. *P. flavicauda*) suggest multiple directions of connectivity between these biogeographical regions through the Coral Sea. The proximity of the Coral Sea to the global centre of marine biodiversity, The Coral Triangle in the Indo-Pacific, mean reef habitats in the Coral Sea may represent a particularly important link in larger-scale patterns of reef fish biodiversity, rather than isolated populations with limited distribution (Hobbs et al. 2009; Budd and Pandolfi 2010). Though often speculated, prior to this study there has been scant empirical evidence to support the role of the Coral Sea seamounts as stepping stones for reef fish populations (but see Van Herwerden et al. 2009), or in contributing to biodiversity feedback between regions.

The stepping stone model of dispersal has been shown to contribute to the biodiversity feedback process for reef fish assemblages in other regional seamount chains (Pinheiro et al. 2015, 2018; Mazzei et al. 2021). From studies in the Southwest Atlantic, coastal populations

closer to the Brazilian continental shelf represent areas of higher biodiversity, and genetic connectivity exists in both directions between these populations and the most offshore seamounts (Simon et al. 2022). Interestingly, of the sixteen new occurrence records found in this study from the Coral Sea Marine Park, thirteen are not known from the neighbouring Great Barrier Reef. At least nine of these species are deep water-specialists, typically only known from depths greater than 30m and up to 120m. We include *H. marcosi* in these deep water-specialists which, prior to these multiple new Coral Sea observations, is only reported in Australia from a single individual at 100m at the GBR shelf break (Sih et al. 2017) and has not been recorded from the reefs of the GBR itself. Although a lack of observations on the GBR for these thirteen species may reflect low sampling from mesophotic depths, the shallow geomorphology of the GBR shelf (30 - 50m, Hopley 2006) likely restricts the establishment of populations of the deep water specialist, or mesophotic, species reported here. Indeed, the occurrence of these mesophotic species throughout the full latitudinal extent of the CSMP, but not on the GBR, suggests that connectivity for these species is greater latitudinally along the seamount chain, where deep-water habitat is available between the Coral Triangle and island chains of the Southwest Pacific. Connectivity patterns in the Coral Sea remain unclear but for shallow water studies both genetic analyses (Planes et al. 2001; Payet et al. 2022) and dispersal-driven connectivity models (Tremblay et al. 2008) suggest that connectivity between the Coral Sea and GBR is generally weak. The barriers to connectivity between these regions have not been fully established, but the spatial separation of these two regions by deep open water and the lack of mesophotic habitat on the GBR would certainly contribute to a dispersal barrier for mesophotic species from the Coral Sea seamounts to the shallow GBR shelf. Further studies utilising genetic sampling of mesophotic fishes with extended ranges throughout the CSMP and from neighbouring regions is required to test

aspects of the biodiversity feedback hypothesis and quantify the nature and direction of meta-population connectivity throughout deep reefs of the region.

A second mechanism, particularly relevant to patterns of mesophotic fish diversity, is “The Habitat Persistence Hypothesis” (HPH) (Copus et al. 2022). This theory posits that during periods of lower relative sea level, deeper habitats can persist, particularly on complex bathymetries where there are horizontal and low aspect areas, while shallow water habitats are dried out and communities here are lost. Multiple biogeographical hypotheses of global patterns and processes in reef fish diversity are supported to varying degrees by often overlapping empirical evidence (Mora et al. 2003; Gaither and Rocha 2013; Bowen et al. 2013; Cowman et al. 2017). Yet, most of this evidence is derived from shallow water coral reefs (<30m) which are estimated to represent only 20% of global coral reef habitat (Pyle and Copus 2019). These data therefore must be considered incomplete regarding both species inventories and the extent of available habitat. Indeed, unlike shallow water reef fishes, diversity for mesophotic fishes does not appear to attenuate with distance from The Coral Triangle (Pyle 2000, 2005; Pyle and Copus 2019) and this mismatch suggests that there are further mechanisms shaping reef fish diversity than have currently been considered (Pinheiro et al. 2023). Again, although the lack of mesophotic sampling effort in the region must be acknowledged, the absence of fishes which are reported in this study from the adjacent GBR aligns with several mechanisms proposed by the HPH. During periods of lower relative sea level, species richness may have been retained in deeper habitats of the Coral Sea, and, in these persisting habitats evolutionary processes would continue amongst populations, potentially driving higher rates of speciation. Certainly, taxonomic revisions and redefined species complexes within some reef fish genera demonstrate high levels of speciation across broad distributions which span island chains, oceanic islands and seamounts in the Western

and Indo-Pacific. These include the *Pomacentrus philippinus* group, which in the Coral Sea is represented by *P. imitator* but in the adjacent GBR is *P. magniseptus* (Allen et al. 2017), the goby genus *Nemateleotris* (Tea and Larson 2023) and multiple *Pseudanthias* species (Anderson 2018, 2022; Gill 2022) many of which are deep-water specialists. The compilation and comparison of updated species inventories is required to test the applicability of the Habitat Persistence Hypothesis in the Coral Sea, as well as genetic sampling to establish patterns of connectivity between mesophotic populations in adjacent regions. The complex bathymetries of the Coral Sea seamounts do however, constitute a significant system that aligns with many of the mechanisms proposed by the HPH and is a promising region in which to test these concepts further.

Total habitat area and the arrangement of a variety of habitat types are also fundamental components of species-area-isolation relationships which drive biodiversity patterns (MacArthur and Wilson 1967; Connor and McCoy 1979; Fahrig 2013; Hanski 2015). The presence of multiple deep-water habitats at individual Coral Sea reefs highlights that there is considerably greater habitat area and habitat heterogeneity within the Coral Sea than known from shallow reefs alone. This is significant given that most measures of ecological isolation comprise some measure of patch size combined with distance from nearest neighbouring habitat and the properties of the surrounding matrix (Moilanen and Nieminen 2002; Prugh et al. 2008). Compared to continental scales (thousands of kilometres), seamounts in the Coral Sea are separated by relatively small distances (< 450km maximum distance). Isolation can both positively and negatively affect biodiversity, either via demographic effects (Hanski et al. 2013; Fahrig 2013; Jones et al. 2020) or distance from anthropogenic influences (Demartini et al. 2008; Williams et al. 2011; Bennett et al. 2018). In the context of this study the relatively close proximity of many reefs throughout the seamount chain, in combination

with increased habitat area and heterogeneity at mesophotic depths, may represent an optimal level of isolation between populations. This in turn would facilitate dispersal and enhanced connectivity for some taxa through the stepping stone model (Baum et al. 2004; Saura et al. 2014). The relationships between increased habitat area and reduced isolation, together with levels of habitat heterogeneity and quality also drive biodiversity through other ecological dynamics (Gratwicke and Speight 2005; Szangolies et al. 2022). For example, *Halimeda* spp. meadows, seagrass and other macroalgae beds are known to provide valuable nursery habitats for reef fish settlement and recruitment (Sambrook et al. 2019; Tang et al. 2020; Sievers et al. 2020). Although this function has not been extensively tested in MCEs, *Halimeda* beds are known to support diverse mesophotic fish communities (Langston and Spalding 2017; Spalding et al. 2019) and we found these habitats on deep outer reef slopes and in lagoons of Coral Sea reefs. Given their isolation from other coastal nursery habitats, these habitats may be particularly important for the early life stages of fishes and invertebrates, and thereby the replenishment and maintenance of populations. Similarly, areas of high coral cover at mesophotic depths increase total habitat area and resource availability for coral associated and dependent fishes. Although resources at range margins, including depth, can be of lower quality and affect the physiological condition of some reef fishes (Munday 2001; Srinivasan 2003; Hoey et al. 2007), others including highly specialised obligate coral feeding butterflyfishes have been shown to access equal or greater resources from deeper reefs without impacting their fitness (Macdonald et al. 2018; MacDonald et al. 2019). Further, other studies of energetic trade-offs associated with marginal coral reef habitats have found that deep reefs support robust subpopulations through demographic and reproductive plasticity (Goldstein et al. 2016). Biological traits including habitat preference, diet and dispersal ability therefore also strongly influence connectivity differentially between species and ontogenetic stages (Hixon and Jones 2005; Goldstein et al. 2017). In the context of this

study, this is evident from the function of seamounts as stepping stones for deep-sea invertebrates, where mismatches between expected and observed dispersal patterns can be explained by a combination of environmental parameters and biological traits (Miller and Gunasekera 2017). Finally, although there is limited connectivity between shallow and mesophotic ecological assemblages in many regions for some taxa (Morais and Santos 2018; Bongaerts and Smith 2019; Stefanoudis et al. 2019), the depth extensions for 158 species in the Coral Sea reported by this study suggest that boundaries between shallow and deep assemblages may be shifted deeper here than community breaks known from other regions (Lesser et al. 2019). Although MCEs clearly warrant conservation actions and scientific investigation independent from shallow-water reefs (Bridge et al. 2013; Rocha et al. 2018), it is recognised that the degree of MCE species overlap with shallow-water assemblages can vary considerably by taxon and location (Laverick et al. 2018). This necessitates further ecological and environmental sampling from MCEs in remote geographic regions, like the Coral Sea, and in understudied habitats, like seamounts.

Increased mesophotic surveys will likely continue to increase diversity records for the region and comprehensive fish species checklists for the Coral Sea Marine Park, and Coral Sea region more broadly, will undoubtedly continue to expand on the observations presented by this study. Recent large scale survey efforts in the region have collected the most detailed bathymetry data for these reefs to date (Carroll et al. 2021; Beaman et al. 2022; Brooke and Schmitt Ocean Institute 2022) and have substantially expanded our understanding of deep-sea habitats of the Coral Sea. Unfortunately, despite the evident value of deep-sea exploration, technical constraints on large ship-based ROVs mean that such work often only focuses on mesophotic habitats for short periods of time. Our findings highlight the utility of small, affordable ROVs as an effective tool for conducting mesophotic surveys in remote regions

where technical diving is often not feasible. Although beyond the scope of this study, trait-based analysis of the fish community data collected by ROV surveys would be an informative line of further investigation to establish how ecological characteristics (e.g. body size, depth range, habitat preference, dispersal ability) contribute to habitat use and geographical ranges throughout the Coral Sea. Further, despite significant advances in understanding connectivity and recruitment patterns in tropical reef fishes (Jones et al. 1999; Mora et al. 2003; Mora 2004; Planes et al. 2009; Jones 2015; Almany et al. 2017) ecological studies from seamounts are typically focused on cold-water, true deep-sea taxa (Rowden et al. 2005, 2010; Pitcher et al. 2007; Clark et al. 2010a; Rogers 2018). Coral reef communities from tropical seamounts, including those inhabiting MCEs, are therefore underrepresented in an already understudied global marine habitat but are known to support abundant and diverse reef fish communities (Letessier et al. 2019; Galbraith et al. 2021; Leitner et al. 2021). As biodiversity hotspots and important patch habitats for connectivity, seamounts should more widely be considered as conservation priorities in coral reef seascapes (McCook et al. 2009; Santos and Morato 2010; Riva and Fahrig 2022). Increased mesophotic community surveys in the region and in other tropical seamount chains will contribute to baseline knowledge of reef fish community structure and species distributions in these habitats. Further observational studies should also be supported by genetic sampling to establish the nature and direction of meta-population connectivity throughout tropical seamount chains and within adjacent biogeographical regions.

The findings from this study support the stepping stone model of dispersal for seamounts in a tropical seascape. The new records of sixteen fishes recorded throughout the Coral Sea Marine Park confirm that the geographic range of many tropical reef fishes are more widespread than currently reported. Sampling deficiencies in the region and at mesophotic

depths are clearly a significant reason behind new species observations. Nevertheless, these new records from the Coral Sea contribute to evidence that tropical seamounts are important habitats for reef fish dispersal and connectivity between the Indo-Pacific, Coral Triangle and western Pacific region. The discovery of high coral-cover mesophotic reefs, combined with other diverse deep-water benthic communities, demonstrates the potential of mesophotic habitats of the Coral Sea to provide valuable corridors for dispersal for coral reef fishes. The spatial arrangement of the Coral Sea seamounts along the boundary of major biogeographical regions also suggests that these tropical seamounts function in processes driving large scale marine biodiversity patterns. These include but are likely not limited to biodiversity feedback between peripheral regions and centres of marine biodiversity and, differences in the distribution and diversity of mesophotic fishes driven by habitat persistence through periods of sea-level change. These paradigms may operate across other regional tropical seamount chains and suggests these habitats have an important role in the maintenance and regulation of global biodiversity patterns for coral reef taxa.

Acknowledgements

We are grateful to JH Choat, GP Jones and C MacDonald for thoughtful discussion of and comments on this manuscript. Thanks also to D Bray, A Hay, M McNeil, J Pogonoski, YK Tea and F Walsh for assistance with cross referencing species distributions and identification. ROV and BRUV surveys were logistically supported by the skippers of the MV Iron Joy, Rob Benn, and MV Argo, Jeff Farnham, and their crew. Photo of *B. paraleucosticticus* in Fig. 3d courtesy of Richard Bajol/Alain Daoulas.

Compliance with Ethical Standards

The authors have no competing interests to declare that are relevant to the content of this article. All authors contributed to the study conception and design. Material preparation and data collection were performed GFG and BJC. Analysis was performed by GFG, BJC and ECM. The first draft of the manuscript was written by GFG and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding

Funding for all surveys was provided by the Australian Government's Our Marine Parks Grants (Rounds 2 and 3) to ASH, GFG and ECM (grant numbers – 4-FISKTNX and 4-HAY3RAP respectively) and the ARC Centre of Excellence for Coral Reef Studies (ASH). Additional ROV surveys in July 2021 were facilitated by Parks Australia with operational support from Queensland Department of Environment and Science as part of the Coral Sea Island Health voyage.

Ethics and permitting statement

This research was conducted under permission from the Director of National Parks (PA2020-00092-3) and the James Cook University Animal Ethics Committee (approval A2721).

Data availability statement

All species recorded by ROV and BRUV in this study are provided as Online Resource 1. A list of depth extensions for fishes compared to fishbase records is provided in Online Resource 2. Occurrence data used to plot Figure 3 were downloaded from the OBIS database (OBIS 2023a,b,c,d,e), GBIF database (GBIF 2023a,b,c,d,e) and are provided as a csv file at [10.5281/zenodo.7844209](https://doi.org/10.5281/zenodo.7844209).

References

- ALA (2023) Atlas of Living Australia website at <http://www.ala.org.au>. Accessed 4 March 2023
- ABRS 2020. Australian Faunal Directory. Australian Biological Resources Study, Canberra. Viewed 21 March 2023, <https://biodiversity.org.au/afd/home>
- Allen G, Museum WA, Erdmann M V, et al (2017) Descriptions of four new species of damselfishes (Pomacentridae) in the *Pomacentrus philippinus* complex from the tropical western Pacific Ocean. <https://doi.org/10.5281/zenodo.317395>
- Allen GR, Erdmann M V, Hamilton AM (2010) *Hoplolatilus randalli*, a new species of sand tilefish (Pisces : Malacanthidae) from the tropical western Pacific with comments on the validity of *H. luteus*. *Aqua, Int J Ichthyol* 16:171–186
- Almany GR, Planes S, Thorrold SR, et al (2017) Larval fish dispersal in a coral-reef seascape. *Nat Ecol Evol* 1:148. <https://doi.org/10.1038/s41559-017-0148>
- Anderson WD (2022) Additions and emendations to the annotated checklist of anthiadine fishes (Percoidei: Serranidae). 5195:567–578
- Anderson WD (2018) Annotated checklist of anthiadine fishes (Percoidei: Serranidae) WILLIAM. *Zootaxa* 4475:1–62. <https://doi.org/https://doi.org/10.11646/zootaxa.4475.1.1>
- Atlas of Living Australia (2023) *Anampses melanurus* occurrence records <https://doi.org/10.26197/ala.6aff80e8-d157-4549-857b-6950c4949ca5>, accessed 21/03/2023
- Atlas of Living Australia (2023) *Mulloidichthys pfluegeri* occurrence records <https://doi.org/10.26197/ala.0e0763bf-259f-45e7-84d1-ec8ae391355b>, accessed 21/03/2023
- Australian Faunal Directory (2023) *Valenciennesa helsdingenii* occurrence records https://biodiversity.org.au/afd/taxa/Valenciennesa_helsdingenii, accessed 21/03/2023
- Ayling A, Ayling A (1985) Report on a preliminary survey of the Lihou and Coringa/Herald Nature Reserves.
- Baum KA, Haynes KJ, Dilleuth FP, Cronin JT (2004) The matrix enhances the effectiveness of corridors and stepping stones. *Ecology* 85:2671–2676. <https://doi.org/10.1890/04-0500>
- Beaman R. (2010) Project 3DGBR: A high-resolution depth model for the Great Barrier Reef and Coral Sea. Marine and Tropical Sciences Research Facility (MTSRF) Project 2.5i.1a Final Report,
- Beaman RJ, Picard K, Miller A (2022) RV Falkor surveys in Australia 2020-2021. In: Maschke J (ed) *Hydrospatial 2021 Conference*, 16-18 Feb 2022. Australasian Hydrographic Society,. Cairns, Australia
- Becker R., Wilks R., Brownrigg R, et al (2022) maps: Draw Geographical Maps
- Bennett E. (1832) Observations on a collection of fishes from Mauritius, presented by Mr Telfair, with characters of new genera and species, remaining portion. *Proc Comm Sci Corresp Zool Soc London* 1:165–169
- Bennett S, Halford AR, Choat JH, et al (2018) Geography and island geomorphology shape fish

- assemblage structure on isolated coral reef systems. *Ecol Evol* 8:6242–6252.
<https://doi.org/10.1002/ece3.4136>
- Bleeker P (1868) Description de deux espèces inédites d'*Epinephelus* rapportées de l'île de la Réunion par M.M. Pollen et van Dam. *Versl en Meded der K Akad van Wet (Afdeeling Natuurjunde)* 2:336–341
- Bleeker P (1858) Bijdrage tot de kennis der vischfauna van den Goram-Archipel. *Natuurwetenschappelijk Tijdschr voor Ned Indië* 15:97–218
- Bleeker P (1857) Achtste bijdrage tot de kennis der vischfauna van Amboina. *Acta Soc Regiae Sci Indo-Neêrlandicae* 2:1–102
- Bray, D.J. & Gomon, M.F. (2023) Fishes of Australia. Museums Victoria and OzFishNet, accessed 21 March 2023, <http://fishesofaustralia.net.au/>
- Boettiger C, Lang DT, Wainwright PC (2012) Rfishbase: Exploring, manipulating and visualizing FishBase data from R. *J Fish Biol* 81:2030–2039. <https://doi.org/10.1111/j.1095-8649.2012.03464.x>
- Bongaerts P, Bridge TCL, Kline DI, et al (2011) Mesophotic coral ecosystems on the walls of Coral Sea atolls. *Coral Reefs* 30:335–335. <https://doi.org/10.1007/s00338-011-0725-7>
- Bongaerts P, Smith TB (2019) Beyond the “Deep Reef Refuge” Hypothesis: A Conceptual Framework to Characterize Persistence at Depth. In: *Coral Reefs of the World*. Springer, Cham, pp 881–895
- Bowen BW, Rocha LA, Toonen RJ, Karl SA (2013) The origins of tropical marine biodiversity. *Trends Ecol Evol* 28:359–366. <https://doi.org/10.1016/J.TREE.2013.01.018>
- Bray DJ (2023) *Cephalopholis polleni* in Fishes of Australia,. <https://fishesofaustralia.net.au/home/species/5007>. Accessed 25 Mar 2023
- Bray DJ (2022) *Pseudanthias flavicauda* in Fishes of Australia. <https://fishesofaustralia.net.au/home/species/4998>. Accessed 27 Mar 2023
- Bridge TC, Hughes TP, Guinotte JM, Bongaerts P (2013) Call to protect all coral reefs. *Nat Clim Chang* 3:528–530. <https://doi.org/10.1038/nclimate1879>
- Bridge TCL, Beaman RJ, Bongaerts P, et al (2019) The Great Barrier Reef and Coral Sea. In: Loya Y, Puglise KA, Bridge TCL (eds) *Mesophotic Coral Ecosystems*. Springer International Publishing, Cham, pp 351–367
- Brooke B, Schmitt Ocean Institute (2022) Schmidt Ocean Institute Expedition Report: Seamounts, Canyons and Reefs of The Coral Sea.
- Budd AF, Pandolfi JM (2010) Evolutionary novelty is concentrated at the edge of coral species distributions. *Science* (80-) 328:1558–1562
- Burgess WE (1978) Two new species of tilefishes (family Branchiostegidae) from the western Pacific. *Trop Fish Hobbyist* 26:43–47
- Carroll A, Brooke B, Nichol S, et al (2021) Seamounts, Reefs and Canyons of Australia's Coral Sea and Great Barrier Reef Marine Parks: A Comprehensive Deep Sea Study. In: *ICRS 2021 - 14th International Coral Reef Symposium*. p Abstract no.: ICRS2021-2115.

- Ceccarelli DM (2010) Research and Monitoring in Australia's Coral Sea: A Review. Report for DSEWPac
- Ceccarelli DM, David McKinnon A, Andréfouët S, et al (2013) The Coral Sea: Physical Environment, Ecosystem Status and Biodiversity Assets. In: *Advances in Marine Biology*. pp 213–290
- Clark M., Bowden D. (2015) Seamount biodiversity: high variability both within and between seamounts in the Ross Sea region of Antarctica. *Hydrobiologia* 761:161–180. <https://doi.org/10.1007/s10750-015-2327-9>
- Clark MR, Rowden AA, Schlacher T, et al (2010a) The Ecology of Seamounts: Structure, Function, and Human Impacts. *Annu Rev Mar Sci* 2:253–78. <https://doi.org/10.1146/annurev-marine-120308-081109>
- Clark MR, Schlacher TA, Rowden AA, et al (2010b) Science Priorities for Seamounts: Research Links to Conservation and Management. *PLoS One* 7:e29232. <https://doi.org/10.1371/journal.pone.0029232>
- Connor EF, McCoy ED (1979) The Statistics and Biology of the Species-Area Relationship. *Am Nat* 113:791–833
- Copus JM, Pyle RL, Bowen BW, et al (2022) The Habitat Persistence Hypothesis: a new perspective on the distribution of coral-reef organisms. *Front Biogeogr* 14:. <https://doi.org/10.21425/F5FBG57427>
- Cowman PF, Parravicini V, Kulbicki M, Floeter SR (2017) The biogeography of tropical reef fishes: endemism and provinciality through time. *Biol Rev* 92:2112–2130. <https://doi.org/10.1111/brv.12323>
- Davies P, Symonds P, Feary D, Pigram C (1989) The evolution of carbonate platforms of northeast Australia. In: Crevello P, Wilson J, Sarg J, Read J (eds) *Controls on Carbonate Platform and Basin Development*. SEPM Special Publications, Tulsa, pp 233–258.
- Demartini EE, Friedlander AM, Sandin SA, Sala E (2008) Differences in fish-assemblage structure between fished and unfished atolls in the northern Line Islands, central Pacific. 365:199–215. <https://doi.org/10.3354/meps07501>
- Director of National Parks (2018) Coral Sea Marine Park Management Plan. Canberra
- Edgar G, Stuart-Smith R (2014) Reef Life Survey (RLS): Global reef fish dataset
- Edgar GJ, Ceccarelli DM, Stuart-Smith RD, (2015) Reef Life Survey Assessment of Coral Reef Biodiversity in the Coral Sea. Report for the Department of the Environment. The Reef Life Survey Foundation Inc. and Institute of Marine and Antarctic Studies
- Englebert N, Bongaerts P, Muir P, et al (2015) Deepest zooxanthellate corals of the Great Barrier Reef and Coral Sea. *Mar Biodivers* 45:1–2. <https://doi.org/10.1007/s12526-014-0221-8>
- Englebert N, Bongaerts P, Muir PR, et al (2017) Lower Mesophotic Coral Communities (60- 125 m Depth) of the Northern Great Barrier Reef and Coral Sea. 1–16. <https://doi.org/10.1371/journal.pone.0170336>
- Eyal G, Laverick JH, Bongaerts P, et al (2021) Mesophotic Coral Ecosystems of the Great Barrier Reef Are Understudied and Underexplored. *Front Mar Sci* 8:. <https://doi.org/10.3389/fmars.2021.622856>

- Fahrig L (2013) Rethinking patch size and isolation effects: the habitat amount hypothesis. *J Biogeogr* 40:1649–1663. <https://doi.org/10.1111/jbi.12130>
- Fricke R, Earle JL, Pyle RL, Séret B (2011a) Checklist of the fishes. In: Bouchet P, Le Guyader H, Pascal O (eds) *The natural history of Santo - Patrimoines naturels*. Paris (MNHN, PNI), Marseille (PNI), pp 343–409
- Fricke R, Kulbicki M (2007) Checklist of the shore fishes of New Caledonia. In: Ayri C., Richer de Forges B (eds) *Compendium of marine species from New Caledonia*. 2nd edition, 2nd Editio. pp 357–401, pls. 15/1 and 15/2.
- Fricke R, Kulbicki M, Wantiez L (2011b) Checklist of the fishes of New Caledonia, and their distribution in the Southwest Pacific Ocean (Pisces). *Stuttgarter Beitrage Zur Naturkunde, Ser A* 341–463
- Froese R, Pauly D (2023) FishBase. In: *World Wide Web Electron. Publ.* www.fishbase.org
- Gaither MR, Rocha LA (2013) Origins of species richness in the Indo-Malay-Philippine biodiversity hotspot: Evidence for the centre of overlap hypothesis. *J Biogeogr* 40:1638–1648. <https://doi.org/10.1111/jbi.12126>
- Galbraith G, McClure E, Barnett A, et al (2022) Diving into the Deep: the Unique Deep Habitats of the Coral Sea Marine Park. Report prepared for Parks Australia. [10.13140/RG.2.2.10516.68487](https://doi.org/10.13140/RG.2.2.10516.68487)
- Galbraith GF, Cresswell BJ, McCormick MI, et al (2021) High diversity, abundance and distinct fish assemblages on submerged coral reef pinnacles compared to shallow emergent reefs. *Coral Reefs* 40:335–354. <https://doi.org/10.1007/s00338-020-02044-z>
- GBIF.org (2023a), GBIF Home Page. Available from: <https://www.gbif.org>
- GBIF.org (2023b) GBIF Occurrence Download <https://doi.org/10.15468/dl.uyvhj9> [Accessed 1 March 2023]
- GBIF.org (2023c) GBIF Occurrence Download <https://doi.org/10.15468/dl.f7ak26> [Accessed 1 March 2023]
- GBIF.org (2023d) GBIF Occurrence Download <https://doi.org/10.15468/dl.qspxgx> [Accessed 1 March 2023]
- GBIF.org (2023e) GBIF Occurrence Download <https://doi.org/10.15468/dl.zjpsfj> [Accessed 1 March 2023]
- Gilbert C. (1905) II. The deep-sea fishes. In: Jordan D., Evermann B. (eds) *The aquatic resources of the Hawaiian Islands*. Bulletin of the United States Fish Commission : 23(2). pp 575–713 figs 230–276 pls 66–101
- Gill AC (2022) Revised definitions of the anthiadine fish genera *Mirolabrichthys* Herre and *Nemanthias* Smith, with description of a new genus (Teleostei: Serranidae). *Zootaxa* 5092:41–66. <https://doi.org/10.11646/zootaxa.5092.1.2>
- Goldstein ED, D'Alessandro EK, Sponaugle S (2017) Fitness consequences of habitat variability, trophic position, and energy allocation across the depth distribution of a coral-reef fish. *Coral Reefs* 36:957–968. <https://doi.org/10.1007/s00338-017-1587-4>
- Goldstein ED, D'alessandro EK, Sponaugle S (2016) Demographic and reproductive plasticity across

- the depth distribution of a coral reef fish. *Nat Publ Gr*. <https://doi.org/10.1038/srep34077>
- Gomon MF (2006) A revision of the labrid fish genus *Bodianus* with descriptions of eight new species. *Rec Aust Museum* 58:1–133. <https://doi.org/10.3853/j.0812-7387.30.2006.1460>
- Gratwicke B, Speight MR (2005) The relationship between fish species richness, abundance and habitat complexity in a range of shallow tropical marine habitats. *J Fish Biol* 66:650–667. <https://doi.org/10.1111/j.0022-1112.2005.00629.x>
- Hanski I (2015) Habitat fragmentation and species richness. *J Biogeogr* 42:989–993. <https://doi.org/10.1111/jbi.12478>
- Hanski I, Zurita GA, Bellocq MI, Rybicki J (2013) Species-fragmented area relationship. *Proc Natl Acad Sci U S A* 110:12715–12720. https://doi.org/10.1073/PNAS.1311491110/SUPPL_FILE/PNAS.201311491SI.PDF
- Harris P, Heap A, Passlow V, et al (2003) Geomorphic Features of the Continental Margin of Australia GEOSCIENCE AUSTRALIA
- Hixon MA, Jones GP (2005) Competition, predation, and density-dependent mortality in demersal marine fishes. *Ecology* 86:2847–2859. <https://doi.org/10.1890/04-1455>
- Hobbs J-PA, Jones GP, Munday PL, et al (2012) Biogeography and the structure of coral reef fish communities on isolated islands. *J Biogeogr* 39:130–139. <https://doi.org/10.1111/j.1365-2699.2011.02576.x>
- Hobbs JPA, Choat JH, Robbins WD, et al (2008) Unique fish assemblages at world's southernmost oceanic coral reefs, Elizabeth and Middleton Reefs, Tasman Sea, Australia. *Coral Reefs* 27:15. <https://doi.org/10.1007/S00338-007-0301-3/FIGURES/1>
- Hobbs JPA, Frisch AJ, Allen GR, Van Herwerden L (2009) Marine hybrid hotspot at Indo-Pacific biogeographic border. *Biol Lett* 5:258–261. <https://doi.org/10.1098/rsbl.2008.0561>
- Hoey A, Harrison H, McClure E, et al (2021) Coral Sea Marine Park Coral Reef Health Survey 2021.
- Hoey A, McClure E, Burn D, et al (2022) Coral Sea Marine Park Coral Reef Health Survey 2022. Report prepared for Parks Australia.
- Hoey AS, Harrison H., Pratchett M. (2020) Coral Reef Health in the Coral Sea Marine Park – Surveys 2018-2020
- Hoey J, McCormick MI, Hoey AS (2007) Influence of depth on sex-specific energy allocation patterns in a tropical reef fish. *Coral Reefs* 26:603–613. <https://doi.org/10.1007/s00338-007-0246-6>
- Hopley D (2006) Coral reef growth on the shelf margin of the Great Barrier Reef with special reference to the Pompey complex. *J Coast Res* 22:. <https://doi.org/doi:10.2112/05A-0012.1>
- Hubbs CL (1959) Initial Discoveries of Fish Faunas on Seamounts and Offshore Banks in the Eastern Pacific!
- Jones G., Milicich MJ, Emslie MJ, Lunow C (1999) Self-recruitment in a coral reef fish population. *Nature* 402:802–804
- Jones GP (2015) Mission impossible: unlocking the secrets of coral reef fish dispersal. In: Mora C (ed) *Ecology of Fishes on Coral Reefs*. Cambridge University Press, Cambridge, pp 16–27

- Jones GP, Barone G, Sambrook K, Bonin MC (2020) Isolation promotes abundance and species richness of fishes recruiting to coral reef patches. *Mar Biol* 167:167. <https://doi.org/10.1007/s00227-020-03772-0>
- Kahle D, Wickham H (2013) ggmap: Spatial Visualization with ggplot2. *R J* 5:144–161
- Kessler WS, Cravatte S (2013) Mean circulation of the Coral Sea. *J Geophys Res Ocean* 118:6385–6410. <https://doi.org/10.1002/2013JC009117>
- Kulbicki M, Parravicini V, Bellwood DR, et al (2013) Global Biogeography of Reef Fishes: A Hierarchical Quantitative Delineation of Regions. *PLoS One* 8:81847. <https://doi.org/10.1371/journal.pone.0081847>
- Langlois T, Goetze J, Bond T, et al (2020) A field and video annotation guide for baited remote underwater stereo-video surveys of demersal fish assemblages. *Methods Ecol Evol* 11:1401–1409. <https://doi.org/10.1111/2041-210X.13470>
- Langston RC, Spalding HL (2017) A survey of fishes associated with Hawaiian deep-water *Halimeda* *kanaloana* (Bryopsidales: Halimedaceae) and *Avrainvillea* sp. (Bryopsidales: Udoteaceae) meadows. *PeerJ* 2017:. <https://doi.org/10.7717/peerj.3307>
- Laverick JH, Piango S, Andradi-Brown DA, et al (2018) To what extent do mesophotic coral ecosystems and shallow reefs share species of conservation interest? A systematic review. *Environ Evid* 7:1–13. <https://doi.org/10.1186/s13750-018-0127-1>
- Leitner A, Friedrich T, Kelley C, et al (2021) Biogeophysical influence of large-scale bathymetric habitat types on mesophotic and upper bathyal demersal fish assemblages: a Hawaiian case study. *Mar Ecol Prog Ser* 659:219–236. <https://doi.org/10.3354/meps13581>
- Lesser MP, Slattery M, Laverick JH, et al (2019) Global community breaks at 60 m on mesophotic coral reefs. *Glob Ecol Biogeogr* 00:1–14. <https://doi.org/10.1111/geb.12940>
- Lesser MP, Slattery M, Mobley CD (2018) Biodiversity and Functional Ecology of Mesophotic Coral Reefs. *Annu Rev Ecol Evol Syst* 49:49–71. <https://doi.org/10.1146/annurev-ecolsys-110617>
- Letessier TB, Mouillot D, Bouchet PJ, et al (2019) Remote reefs and seamounts are the last refuges for marine predators across the Indo-Pacific. *PLOS Biol* 17:e3000366. <https://doi.org/10.1371/journal.pbio.3000366>
- Loya Y, Eyal G, Treibitz T, et al (2016) Theme section on mesophotic coral ecosystems: advances in knowledge and future perspectives. *Coral Reefs* 35:1–9. <https://doi.org/10.1007/s00338-016-1410-7>
- MacArthur RH, Wilson EO (1967) *The theory of island biogeography*. Princeton University Press, New Jersey
- MacDonald C, Bridge TCL, McMahon KW, Jones GP (2019) Alternative functional strategies and altered carbon pathways facilitate broad depth ranges in coral-obligate reef fishes. *Funct Ecol* 33:1962–1972. <https://doi.org/10.1111/1365-2435.13400>
- Macdonald C, Jones GP, Bridge T (2018) Marginal sinks or potential refuges? Costs and benefits for coral-obligate reef fishes at deep range margins. *Proc R Soc B Biol Sci* 285:20181545. <https://doi.org/10.1098/rspb.2018.1545>
- Mazzei EF, Pinheiro HT, Simon T, et al (2021) Mechanisms of dispersal and establishment drive a

- stepping stone community assembly on seamounts and oceanic islands. *Mar Biol* 168:1–11. <https://doi.org/10.1007/S00227-021-03919-7/FIGURES/4>
- McClain CR (2007) Seamounts : identity crisis or split personality ? *J Biogeogr* (J 34:2001–2008. <https://doi.org/10.1111/j.1365-2699.2007.01783.x>
- McClain CR, Lundsten L, Ream M, et al (2009) Endemicity, Biogeography, Composition, and Community Structure On a Northeast Pacific Seamount. *PLoS One* 4:e4141. <https://doi.org/10.1371/journal.pone.0004141>
- McCook LJ, Almany GR, Berumen ML, et al (2009) Management under uncertainty: Guide-lines for incorporating connectivity into the protection of coral reefs. *Coral Reefs* 28:353–366. <https://doi.org/10.1007/s00338-008-0463-7>
- McKinnon AD, Williams A, Young J, et al (2014) Tropical Marginal Seas: Priority Regions for Managing Marine Biodiversity and Ecosystem Function. *Ann Rev Mar Sci* 6:415–437. <https://doi.org/10.1146/annurev-marine-010213-135042>
- Miller KJ, Gunasekera RM (2017) A comparison of genetic connectivity in two deep sea corals to examine whether seamounts are isolated islands or stepping stones for dispersal. *Sci Rep* 7:46103. <https://doi.org/10.1038/srep46103>
- Moilanen A, Nieminen M (2002) Simple connectivity measures in spatial ecology. *Ecology* 83:1131–1145. [https://doi.org/10.1890/0012-9658\(2002\)083\[1131:SCMISE\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[1131:SCMISE]2.0.CO;2)
- Mora C (2004) Importance of dispersal in coral reef fishes. 784:156
- Mora C, Chittaro PM, Sale PF, et al (2003) Patterns and processes in reef fish diversity. *Nature* 421:933–936. <https://doi.org/10.1038/nature01393>
- Morais J, Santos BA (2018) Limited potential of deep reefs to serve as refuges for tropical Southwestern Atlantic corals. *Ecosphere* 9:. <https://doi.org/10.1002/ecs2.2281>
- Muir P, Wallace C, Bridge TCL, Bongaerts P (2015) Diverse Staghorn Coral Fauna on the Mesophotic Reefs of North-East Australia. *PLoS One* 10:e0117933. <https://doi.org/10.1371/journal.pone.0117933>
- Munday PL (2001) Fitness consequences of habitat use and competition among coral-dwelling fishes. *Oecologia* 128:585–593. <https://doi.org/10.1007/s004420100690>
- OBIS (2023a) Ocean Biodiversity Information System. Intergovernmental Oceanographic Commission of UNESCO. www.obis.org.
- OBIS (2023b) Distribution records of *Hoplostethus medius* (Allen, Erdmann & Hamilton, 2010) [Dataset] (Available: Ocean Biodiversity Information System. Intergovernmental Oceanographic Commission of UNESCO. www.obis.org. Accessed: 21 March 2023)
- OBIS (2023c) Distribution records of *Cephalopholis polleni* (Bleeker 1868) [Dataset] (Available: Ocean Biodiversity Information System. Intergovernmental Oceanographic Commission of UNESCO. www.obis.org. Accessed: 21 March 2023)
- OBIS (2023d) Distribution records of *Pseudanthias flavicauda* (Randall and Pyle 2001) [Dataset] (Available: Ocean Biodiversity Information System. Intergovernmental Oceanographic Commission of UNESCO. www.obis.org. Accessed: 21 March 2023)

- OBIS (2023e) Distribution records of *Bodianus paraleucosticticus* (Gomon 2006) [Dataset] (Available: Ocean Biodiversity Information System. Intergovernmental Oceanographic Commission of UNESCO. www.obis.org. Accessed: 21 March 2023)
- Oxley W, Ayling A, Cheal A, Osborne K (2004) Marine surveys undertaken in the Elizabeth and Middleton Reefs Marine National Nature Reserve, December 2003
- Payet SD, Pratchett MS, Saenz-Agudelo P, et al (2022) Demographic histories shape population genomics of the common coral grouper (*Plectropomus leopardus*). *Evol Appl* 15:1221–1235. <https://doi.org/10.1111/EVA.13450>
- Pérez-Rosales G, Hernández-Agreda A, Bongaerts P, et al (2022) Mesophotic depths hide high coral cover communities in French Polynesia. *Sci Total Environ* 844:157049. <https://doi.org/10.1016/J.SCITOTENV.2022.157049>
- Pinheiro HT, Bernardi G, Simon T, et al (2017) Island biogeography of marine organisms. *Nature* 549:82–85. <https://doi.org/10.1038/nature23680>
- Pinheiro HT, MacDonald C, Quimbayo JP, et al (2023) Assembly rules of coral reef fish communities along the depth gradient. *Curr Biol* 1–10. <https://doi.org/10.1016/j.cub.2023.02.040>
- Pinheiro HT, Mazzei E, Moura RL, et al (2015) Fish Biodiversity of the Vitória-Trindade Seamount Chain, Southwestern Atlantic: An Updated Database. *PLoS One* 10:e0118180. <https://doi.org/10.1371/journal.pone.0118180>
- Pinheiro HT, Rocha LA, Macieira RM, et al (2018) South-western Atlantic reef fishes: Zoogeographical patterns and ecological drivers reveal a secondary biodiversity centre in the Atlantic Ocean. *Divers Distrib* 24:951–965. <https://doi.org/10.1111/ddi.12729>
- Pitcher TJ, Morato T, Hart PJB, et al (2007) *Seamounts: Ecology, Fisheries & Conservation*. Blackwell Publishing Ltd
- Planes S, Doherty PJ, Bernardi G (2001) Strong genetic divergence among populations of a marine fish with limited dispersal, *Acanthochromis polyacanthus*, within the Great Barrier Reef and the Coral Sea. *Evolution* (N Y) 55:2263–2273. <https://doi.org/10.1111/j.0014-3820.2001.tb00741.x>
- Planes S, Jones GP, Thorrold SR (2009) Larval dispersal connects fish populations in a network of marine protected areas. *Proc Natl Acad Sci* 106:5693–5697. <https://doi.org/10.1073/pnas.0808007106>
- Prugh LR, Hodges KE, Sinclair ARE, Brashares JS (2008) Effect of habitat area and isolation on fragmented animal populations. *Proc Natl Acad Sci U S A* 105:20770–20775. <https://doi.org/10.1073/pnas.0806080105>
- Pyle LR (2000) Assessing undiscovered fish biodiversity on deep coral reefs.pdf. *Mar Technol Soc J* 34:82–91. <https://doi.org/doi.org/10.4031/MTSJ.34.4.11>
- Pyle RL (2005) Recent Discoveries of New Fishes Inhabiting Deep Pacific Coral Reefs, with Biogeographic Implications. In: 7th Indo-Pacific Fish Conference (IPFC7). Taipei, Taiwan.
- Pyle RL, Copus JM (2019) Mesophotic Coral Ecosystems: Introduction and Overview. *Coral Reefs of the World* 12:3–27. https://doi.org/10.1007/978-3-319-92735-0_1
- R Core Development Team (2023) *R: A language and environment for statistical computing*

- Randall J, Williams J, Smith D, et al (2003) Checklist of the shore and epipelagic fishes of Tonga. Washington DC
- Randall JE (1975) A Revision of the Indo-Pacific Angelfish Genus *Genicanthus*, with Descriptions of Three New Species. *Bull Mar Sci* 25:393–421
- Randall JE, Lubbock R (1982) Three new labrid fishes of the genus *Cirrhitilabrus* from the southwestern Pacific. *Occas Pap Bernice P Bish Museum* 25:12pp.
- Randall JE, McCosker JE (1982) Two new serranid fishes of the genus *Anthias* from the central Pacific. *J Aquaric* 2:59–69
- Randall JE, Pyle RL (2001) Four new serranid fishes of the anthiine genus *Pseudanthias* from the South Pacific. *Raffles Bull Zool* 49:19–34
- Rees, A.J.J., Yearsley, G.K., Gowlett-Holmes, K. and Pogonoski, J. (2023) Codes for Australian Aquatic Biota (on-line version).CSIRO National Collections and Marine Infrastructure (formerly CSIRO Marine and Atmospheric Research), World Wide Web electronic publication, 1999 onwards. Available at: <https://www.marine.csiro.au/data/caab/>.
- RLS (2023) Reef Life Survey website <https://reeflifesurvey.com/survey-data>. Accessed 25 March 2023
- Richer de Forges B, Koslow JA, Poore GCB (2000) Diversity and endemism of the benthic seamount megafauna in the southwest Pacific. *Nature* 405:944–947
- Riva F, Fahrig L (2022) The disproportionately high value of small patches for biodiversity conservation. *Conserv Lett* 15:1–7. <https://doi.org/10.1111/conl.12881>
- Rocha LA, Pinheiro HT, Shepherd B, et al (2018) Mesophotic coral ecosystems are threatened and ecologically distinct from shallow water reefs. *Science* 361:281–284. <https://doi.org/10.1126/science.aag1614>
- Rogers AD (2018) The Biology of Seamounts: 25 Years on. *Adv Mar Biol* 79:137–224. <https://doi.org/10.1016/BS.AMB.2018.06.001>
- Rowden AA, Clark MR, Wright IC (2005) Physical characterisation and a biologically focused classification of ‘seamounts’ in the New Zealand region. *New Zeal J Mar Freshw Res* 39:. <https://doi.org/10.1080/00288330.2005.9517374>
- Rowden AA, Dower JF, Schlacher TA, et al (2010) Paradigms in seamount ecology: fact, fiction and future. *Mar Ecol* 31:226–241. <https://doi.org/10.1111/j.1439-0485.2010.00400.x>
- Sambrook K, Hoey AS, Andréfouët S, et al (2019) Beyond the reef: The widespread use of non-reef habitats by coral reef fishes. *Fish Fish* 20:903–920. <https://doi.org/10.1111/faf.12383>
- Santos RS, Morato T (2010) Conservation and utilization of biodiversity in seamounts Ricardo Serrão Santos & Telmo. In: UN Conference on Ecosystems and People – Biodiversity for development – The road to 2010 and beyond. Norway, pp 162–169
- Sarano F, Pichon M (1988) Morphology and ecology of the deep fore reef slope at Osprey Reef (Coral Sea). In: Choat J. (ed) *Proceedings of the 6th International Coral Reef Symposium*. Townsville, Australia.
- Saura S, Bodin Ö, Fortin MJ (2014) Stepping stones are crucial for species’ long-distance dispersal

- and range expansion through habitat networks. *J Appl Ecol* 51:171–182.
<https://doi.org/10.1111/1365-2664.12179>
- SeaGis Pty Australia CAL, EventMeasure, TransectMeasure
- Sievers KT, McClure EC, Abesamis RA, Russ GR (2020) Non-reef habitats in a tropical seascape affect density and biomass of fishes on coral reefs. *Ecol Evol* 10:13673.
<https://doi.org/10.1002/ECE3.6940>
- Sih TL, Cappo M, Kingsford M (2017) Deep-reef fish assemblages of the Great Barrier Reef shelf-break (Australia). *Sci Rep* 7:10886. <https://doi.org/10.1038/s41598-017-11452-1>
- Simon T, Pinheiro HT, Santos S, et al (2022) Comparative phylogeography of reef fishes indicates seamounts as stepping stones for dispersal and diversification. *Coral Reefs* 41:551–561.
<https://doi.org/10.1007/S00338-021-02178-8/FIGURES/2>
- Slattery M, Lesser MP (2012) Mesophotic coral reefs: a global model of community structure and function. 9–13
- Spalding HL, Amado-Filho GM, Bahia RG, et al (2019) Macroalgae. Springer, Cham, pp 507–536
- Srinivasan M (2003) Depth distributions of coral reef fishes: the influence of microhabitat structure, settlement, and post-settlement processes. *Oecologia* 137:76–84. <https://doi.org/10.1007/s00442-003-1320-6>
- Stefanoudis P V., Rivers M, Smith SR, et al (2019) Low connectivity between shallow, mesophotic and rariphotic zone benthos. *R Soc Open Sci* 6:190958. <https://doi.org/10.1098/rsos.190958>
- Steindachner F (1900) Fische aus dem Stillen Ocean. Ergebnisse einer Reise nach dem Pacific (Schauinsland, 1896-1897). *nzeiger der Kais Akad der Wissenschaften Math Klasse, Wien* 37:174–178
- Stuart-Smith R, Crawford T, Cooper A, et al (2013) Coral Sea marine biodiversity.
- Szangolies L, Rohwäder MS, Jeltsch F (2022) Single large AND several small habitat patches: A community perspective on their importance for biodiversity. *Basic Appl Ecol* 65:16–27.
<https://doi.org/10.1016/j.baae.2022.09.004>
- Tang S, Graba-Landry A, Hoey AS (2020) Density and height of Sargassum influence rabbitfish (f. Siganidae) settlement on inshore reef flats of the Great Barrier Reef. *Coral Reefs* 39:467–473.
<https://doi.org/10.1007/s00338-020-01908-8>
- Tea YK, Larson HK (2023) Synopsis of the ptereleotrine goby genus *Nemateleotris*, with description of a new species from the western and central Pacific Ocean (Teleostei: Gobiidae). *Raffles Bull Zool* 71:248–266. <https://doi.org/10.26107/RBZ-2023-0019>
- Valenciennes A (1830) in Cuvier G & Valenciennes, *A Histoire naturelle des poissons*. Tome Sixième. Livre sixième. Partie I. Des Sparoïdes; Partie II. Des Ménides. v. 6: i-xxiv + 6 pp. + 1-559, Pls. 141-169. [Valenciennes is author of pp. 1-425, 493-559; Cuvier 426-491. i-x
- Van Herwerden L, Howard Choat J, Newman SJ, et al (2009) Complex patterns of population structure and recruitment of *Plectropomus leopardus* (Pisces: Epinephelidae) in the Indo-West Pacific: Implications for fisheries management. *Mar Biol* 156:1595–1607.
<https://doi.org/10.1007/s00227-009-1195-0>

- Williams ID, Richards BL, Sandin SA, et al (2011) Differences in Reef Fish Assemblages between Populated and Remote Reefs Spanning Multiple Archipelagos Across the Central and Western Pacific. *J Mar Biol* 2011:1–14. <https://doi.org/10.1155/2011/826234>
- Yesson C, Clark MR, Taylor ML, Rogers AD (2011) The global distribution of seamounts based on 30 arc seconds bathymetry data. *Deep Res Part I Oceanogr Res Pap* 58:442–453. <https://doi.org/10.1016/j.dsr.2011.02.004>

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [OnlineResource1.docx](#)
- [OnlineResource2.docx](#)