

## Abscisic acid and reactive oxygen species were involved in slightly acidic electrolyzed water-promoted seed germination in watermelon

Xue Wu<sup>1</sup>, Cuinan Wu<sup>1</sup>, Zhonghua Bian<sup>3</sup>, Zhangying Ye<sup>4</sup>, Lili Meng<sup>1</sup>, Liru Xia<sup>1,2</sup>, Encai Bao<sup>1,2,\*</sup>, Kai Cao<sup>1,2,\*</sup>

<sup>1</sup> The Agriculture Ministry Key Laboratory of Agricultural Engineering in the Middle and Lower Reaches of Yangtze River, Institute of Agricultural Facilities and Equipment, Jiangsu Academy of Agricultural Sciences, Nanjing 210014, Jiangsu, China

<sup>2</sup> School of Agricultural Engineering, Jiangsu University, Zhenjiang 212013, Jiangsu, China

<sup>3</sup> Institute of Urban Agriculture, Chinese Academy of Agricultural Sciences, Chengdu 610299, Sichuan, China

<sup>4</sup> School of Biosystems Engineering and Food Science, Zhejiang University, Hangzhou 310058, Zhejiang, China



### ARTICLE INFO

#### Keyword:

Slightly Acidic Electrolyzed Water  
Seed Germination  
Abscisic Acid  
Reactive Oxygen Species

### ABSTRACT

Seed germination is a vital process in seedling stage. The delayed seed germination and irregular emergence frequently occur in watermelon. It is critical to exploit technologies for promoting seed germination. This study aims to explore the effects of slightly acidic electrolyzed water (SAEW) on the seed germination of watermelon. The results showed that SAEW significantly improved the germination indexes and caused a mass of changes to nutrient substances and enzymatic activities in watermelon seeds. In addition, SAEW treatment decreased abscisic acid (ABA) contents, while increased gibberellin (GA) contents of watermelon seeds by regulating the gene expressions of metabolic related enzymes. Thereinto, SAEW remarkably upregulated the transcriptional level of *CYP707A1* and *CYP707A2* involved in ABA catabolism, which inhibited the accumulation of ABA. Moreover, SAEW could inhibit reactive oxygen species (ROS) accumulation in seeds and improve the seed germination which was identical to the effect of DMTU (dimethyl thiourea, ROS inhibitor) on the seed germination. In addition, the expressions of *RbohC* and *RbohD* involved in ROS accumulation were notably depressed by SAEW. This study demonstrated that exogenous SAEW could inhibit the accumulation of ABA content and decrease the accumulation of ROS, which finally improves the germination ability of watermelon seeds.

### 1. Introduction

Seed germination is one of the most critical phases during seedling establishment, which converts dormant seeds into active seedlings (Baskin et al., 2019; Hubbard et al., 2012). There are a number of factors controlling seed germination including seed vigor, temperature and humidity of germinating environment (Malcolm et al., 2003; Huehne and Bhinija, 2012; Dahlquist-Willard et al., 2016; Jiao et al., 2016; Yu et al., 2016; Guan et al., 2020). Phytohormones include auxin, ABA (abscisic acid), ethylene, GA (gibberellin), CTK (cytokinin), BR (brassinosteroid), and so forth, which could regulate seed dormancy and germination in plants (Miransari and Smith, 2014). It has long been known that GA could break seed dormancy and promote germination (Kucera et al., 2005; Wang et al., 2011). However, ABA is the only hormone known to maintain seed dormancy and delay seed germination (Cutler et al., 2010; Hubbard et al., 2010). Furthermore, more recent

researches have shown that reactive oxygen species (ROS) played roles in seed germination (Lin et al., 2013; Ortiz-Espin et al., 2017; Luo et al., 2021). It was revealed that ROS played important roles in the radicle elongation and endosperm weakening as a non-enzymatic player during seed germination (Yang et al., 2020).

Watermelon (*Citrullus lanatus*) is a highly popular fruit crop with a great economic value and widely cultivated in China, Turkey and Iran with its annual production of over 119 million tons in 2017 (FAOSTAT, 2017). A mass of studies about watermelon have focused on the improvement of nutritional quality, saccharinity and survival rate of grafting as well as the tolerance to biotic and abiotic stresses (Dabirian et al., 2017; Li et al., 2017; Filippou et al., 2019; Jawad et al., 2020; Wei et al., 2020; Zhang et al., 2020). However, there is a scarce study on the seed germination of watermelon. Due to the specific structure and reduced viability of watermelon seeds stored in the natural condition, the delayed seed germination and irregular emergence frequently occur,

\* Corresponding authors: (E Bao), Tel: +86-25-84390180, (K Cao), Tel: +86-25-84390180

E-mail addresses: baoencai1990@163.com (E. Bao), kcao@jaas.ac.cn (K. Cao).

especially for triploid varieties of watermelon. Triploid watermelon seeds are smaller than diploid varieties in size, which causes a limited number of reserves to support germination and seedling growth of triploid watermelon (Pratibha et al., 2020). Prompt and uniform seed germination is crucial for taking full advantage of production resources in commercial agriculture (Chen and Arora, 2013). Therefore, improving the germination rate and emergence quality of watermelon is important, which is beneficial to the growth of later watermelon seedlings and bringing watermelon for the market ahead of time. Up to now, there are few reports on the positive impacts of exogenous substances to the germination of watermelon in natural conditions.

Slightly acidic electrolyzed water (SAEW) is a novel disinfectant generated by the electrolysis of dilute sodium chloride or hydrochloric acid solutions or both in an electrolytic cell without a separating membrane. SAEW, colourless, odourless, and harmless to humans, has been directly used on food surfaces to extend shelf-life in Japan and America (Jung et al., 2017; Wang et al., 2017; Ye et al., 2017; Huang et al., 2021). Meanwhile, SAEW is regarded as an environmental friendly measure for broad-spectrum microbial decontamination in recent years (Ye et al., 2017; Wang et al., 2018). In this research, the effects of SAEW on the germination in watermelon seeds were investigated. We evaluated the effects of SAEW on germination rate, germination energy, germination index and vitality index of watermelon seed. Furthermore, analyses of plant hormone content were used to identify the significant phytohormone responding to SAEW treatment in watermelon seeds. The pharmacology experiment was conducted to preliminarily establish the connection between SAEW-induced germination and ROS. These results may help us to understand the germination mechanisms of watermelon seed, as well as to further reveal the physiological regulatory functions not merely the sterilization of SAEW in plants.

## 2. Materials and methods

### 2.1. Preparation of slightly acidic electrolyzed water (SAEW)

SAEW was prepared by electrolysis of a 6% HCl solution using a generator (HD-240L, Fuqiang-Want Sanitary Accessories Ltd, Shanghai, China) with a non-membrane electrolytic chamber at a voltage of 220 V. SAEW was prepared on the day of the experiments. After that, the pH and the available chlorine concentration of were respectively determined by a pH meter (FE-28, METTLER TOLEDO, Shanghai, China) and chlorine concentration test paper (Newstar paper, Q/HSSC 202-2016, Hangzhou, China).

### 2.2. Plant material and treatments

Seeds of watermelon (*Citrullus lanatus* L. cv. 8424), a kind of triploid hybrid variety, were chosen as the research objective. Watermelon seeds were purchased from 'Hefeng Seed Industry Co., Ltd of China'. The hundred-grain weight and seed germination on the label of seeds were 5.20 g and 85% respectively. Firstly, seeds were respectively soaked in deionized water (Control, Con) or different concentrations of SAEW (20, 40, 60, 80 ppm) at 25°C for 5 h. After that, the effect of SAEW on seed germination was performed by cultivating seeds at 25°C on cotton gauze soaked with deionized water or different concentrations of SAEW in 90 mm plastic petri dish (30 seeds per dish) at 25°C all night in an incubator. The germination indices were measured at indicated time (24, 48, 72, 96 and 120 h). The germination energy was calculated by a formula: GE (%) = M/N × 100. GE is the germination energy (%), parameter M is the number of germinating seeds in five days (120 h) with different treated concentrations of SAEW, and N is the total number of seeds from the germination test. Germination rate (%), GR = A/B × 100. Parameter A is the number of germinating seeds, and B is the total number of seeds from the germination test. Germination index (GI) =  $\sum(x/y)$ , parameter x is the number of germinating seeds in x day, and y is the corresponding days. Finally, Vitality index (VI) = GI × L, parameter GI is the

germination index, and L is the total length of radicle in x day.

In the time-course experiment of germination, watermelon seeds were respectively soaked in deionized water (Con) or 60 ppm SAEW for 1, 2, 5 and 10 h. After that, the time effect of SAEW on seed germination was performed by cultivating seeds at 25°C on cotton gauze soaked with deionized water (Con) or 60 ppm SAEW in 90 mm plastic petri dish (30 seeds per dish) at 25°C all night in an incubator. The germination indices were measured at indicated time (24, 48, 72, 96 and 120 h).

### 2.3. Assay of antioxidant enzyme activity

Fresh seeds of watermelon (0.5 g) with different treatments were homogenized with 10 mL precooled sodium phosphate buffer (50 mM, pH 7.8). After that, the mixed liquor was centrifuged with 12,000 rpm at 4°C for 20 min. The collected supernatant solution was used for enzyme activity assays.

Superoxide dismutase (SOD) and peroxidase (POD) activities were analyzed with the methods described by Han et al. (2007) with some changes. One unit of SOD (U) represented the amount of enzyme required to inhibit 50% nitro blue tetrazolium (NBT) reduction which was measured at 560 nm. U/g FW was represented the activity of SOD. The oxidation rate of guaiacol in presence of H<sub>2</sub>O<sub>2</sub> at 470 nm for 3 min was used to perform the POD activity of watermelon seeds. The absorbance was kept track for every minute and expressed as U/min·g·FW. The activity of catalase (CAT) was determined by Huang et al. (2006) with measuring the reduction of H<sub>2</sub>O<sub>2</sub> at 240 nm for 30 s.

### 2.4. Quantification of soluble sugar, starch, protein and fat

Fresh samples of the shelled watermelon seeds (1 g) were ground in a mortar with 2.5 mL of distilled water. The homogenates were centrifuged at 13,000 rpm for 30 min. Then, the supernatant was used to analyze the contents of soluble sugar. The soluble sugar content was determined using the sulfuric acid anthrone method with a spectrophotometer (UV-5200 spectrophotometer, Shanghai Metash Instruments Co., Ltd, Shanghai, China) at a wavelength of 630 nm (Morris, 1948).

For the analysis of starch content, frozen shelled watermelon seeds were thoroughly ground with liquid nitrogen. The obtained residues were washed with 80% ethanol and filtered again and then dissolved in 20 mL hot distilled water. The mixture was incubated in boiling water bath for 15 min. After incubation, the ice-cooled perchloric acid was added to the mixture and continued ice-cooled for 15 min. After that, the mixture was filtered and the residues were dissolved in 10 mL of distilled water and then incubated in a hot water bath at 100°C for 15 min. The resulting solutions were extracted with 2 mL of perchloric acid (4.6 mol·L<sup>-1</sup>) for 15 min and then filtered. The filtrates were collected and washed with distilled water three times. The resulting solution was adjusted to 100 mL with distilled water to analyze starch content (Lv et al., 2021).

The shelled watermelon seeds samples (0.05 g FW) were ground in a mortar with liquid nitrogen, and the powder transferred with 3 mL of a phosphate buffered solution (pH 7.0) into centrifuge tubes. After 15 min centrifugation at 13,000 g (4°C), 0.1 mL of the supernatant was combined with 5 mL of Coomassie brilliant blue G-250 solution. Two minutes later, the soluble protein content (mg·g<sup>-1</sup> FW) was determined at a wavelength of 595 nm (Su et al., 2017).

The fat content was evaluated according to the modified method of Mohammed et al., 2020. The shelled watermelon seeds samples were homogenized in a solvent mixture of chloroform/ methanol (2:1, v/v) and the mixture was centrifuged at 5,000 g for 20 min. The residues were dissolved in 2 mL of toluene/ethanol mixture (4/1, v/v). The extracted lipids were concentrated by a rotary evaporator and extracted fat was weighed in vials to calculate the fat content.

## 2.5. Determination of phytohormone contents

Seeds were respectively soaked in deionized water (Con) and 60 ppm SAEW at 25°C for 5 h. After that, the effect of SAEW on seed germination was performed by cultivating seeds at 25°C on cotton gauze soaked with deionized water or different concentrations of SAEW in 90 mm plastic petri dish (30 seeds per dish) at 25°C all night in an incubator. The seeds were sampled at indicated time (0, 24, 48 and 72 h). Endogenous bioactive IAA, ABA, GA<sub>1</sub>, GA<sub>3</sub>, GA<sub>4</sub>, GA<sub>7</sub> and Zeatin concentrations in watermelon seeds of the investigated cultivars were measured using HPLC-MS/MS following the protocols described in Pérez-Jiménez et al. (2014) and Camilo et al. (2019), respectively.

## 2.6. Localization of hydrogen peroxide ( $H_2O_2$ ) and superoxide anion ( $O^{2-}$ ) by 3,3'-diaminobezidine (DAB) and nitroblue tetrazolium (NBT)

On account of the high oil and fatty acid concentrations of the seeds, ROS staining in the seeds is difficult. Thus, radicles (germinated for 72 h) were used for further ROS staining processes.

DAB was dissolved in sterilized water (pH=3.8) and produced a solution with a final concentration of 1 mg•ml<sup>-1</sup> according to Luo et al., 2021. Seeds were soaked in deionized water (Con) and 60 ppm SAEW (SAEW) for 5 h. Then, radicles were incubated in DAB-staining solution for 12 h. After that, the stained radicles were washed with 75% ethanol and photographed.

Histochemical localization of superoxide radicals by NBT staining in watermelon was performed according to Zhang et al., (2014). Seeds were soaked in deionized water (Con) and 60 ppm SAEW (SAEW) for 5 h. Then, radicles of seed were incubated 1 mM NBT in 50 mM phosphatic buffer solution (PBS, pH 7.5) at 37°C for 30-60 min. After that, the stained radicles were washed with 75% ethanol and photographed.

## 2.7. Determination of hydrogen peroxide ( $H_2O_2$ ) and superoxide anion ( $O^{2-}$ ) concentrations in plants

$H_2O_2$  content was detected as described in Yu et al., (2019). Seeds were soaked in deionized water (Con) and 60 ppm SAEW (SAEW) for 5 h. Then, 0.1 g radicles were ground in liquid nitrogen and extracted with 1 ml cooled acetone. After having an ice bath for 5 minutes, the samples were centrifuged for 10 min at 10,000 rpm at 4°C. Then, 1.5 ml liquid supernatant was mixed with 0.1 ml 5% (w/v) titanium sulfate and 0.2 ml ammonia. The miscible liquid was centrifuged for 10 min at 10,000 rpm at 4°C. After that, the precipitate was washed with the cooled acetone until it became colorless. 3ml of 2M  $H_2SO_4$  was used to dissolve the precipitate, and then immediately measured the value of  $A_{415}$ . The  $H_2O_2$  content was calculated through a standard curve produced from a series of  $H_2O_2$  standards at known concentrations.

$O^{2-}$  content in this study was quantified according to Luo et al., 2021. Seeds were soaked in deionized water (Con) and 60 ppm SAEW (SAEW) for 5 h. Then, 0.1 g radicles were ground in liquid nitrogen, and the superoxide anion content was measured by manufacturer's instructions of superoxide anion test kit (BC1290; Solarbio Corp.).

## 2.8. RNA extraction and real time quantitative PCR (qRT-PCR)

Watermelon seeds were respectively soaked in deionized water (Con) or 60 ppm SAEW for 5 h. After that, the seed were cultivated at 25°C on cotton gauze soaked with deionized water (Con) or 60 ppm SAEW in 90 mm plastic petri dish (30 seeds per dish) at 25°C all night in an incubator. Then, the seed were sampled at indicated germination time (0, 24, 48 and 72 h). The RNA extraction and qRT-PCR were performed followed the report of Wu et al., (2019). RNA purity was verified based on the ratio (>1.9) of 260/280 nm absorbance using a spectrophotometer. The primer sequences of different genes were designed by Primer Premier 5 software and listed in Table S1. Three independent biological experiments were performed with three replications.

## 2.9. Statistical analysis

Statistical analyses were performed through SPSS statistical software package (version 11.0). And the differences among treatments were analyzed by one-way analysis of variance (ANOVA) combined with Duncan's multiple range test, with P < 0.05 as the threshold. Before the ANOVA, all the data passed the normality test (P > 0.05).

## 3. Results

### 3.1. Effect of slightly acidic electrolyzed water (SAEW) on the germination of watermelon seeds


To compare the effects of SAEW on the seed germination, the watermelon seeds were respectively soaked at 20, 40, 60 and 80 ppm SAEW for 5 h (Fig. 1). Besides, the seeds immersed in deionized water were used as control (Con). There were no differences in the germination rate among various treatments until the germination time reached 48 h (Fig. S1A). As shown in the embedded histogram of Fig. S1A, the germination rate was gradually increased by elevated concentrations of SAEW at 48 h. Thereinto, the germination rate was reached 93.33% when the seeds were treated with 60 ppm SAEW for 48 h. However, the germination rate was only reached 16.67% in Con. In addition, the germination energy, germination index and vitality index of watermelon seeds performed the similar tendency with germination rate, which were gradually increased by SAEW (Fig.S1B-D). Thereinto, the germination energy, germination index and vitality index were respectively increased by 48.28%, 47.49% and 69.35% with 60 ppm SAEW-treatment compared with the control. All the results revealed that 60 ppm SAEW could significantly promote the germination of watermelon seeds.

After the verification of SAEW-treated concentration, the time effect of 60 ppm SAEW on the seed germination was measured by the germination rate, the germination energy, the germination index and the vitality index. As shown in Fig.S2A, the germination rate was increased with the extension of germination time. Thereinto, the significant differences were mainly performed at 48 and 72 h. It was shown that the germination rate, germination energy, germination index and vitality index peaked with 60 ppm SAEW treatment for 5 h. When the treated time was prolonged to 10 h, the germination rate and germination energy were reduced by 60 ppm SAEW (Fig.S2 A and B). Above all, 5 h was the optimum time for SAEW to promote the germination of watermelon seeds.

### 3.2. Effect of slightly acidic electrolyzed water (SAEW) on the concentrations of nutrient substances and activities of enzymes

The process of germination could cause a mass of changes to nutrient substances in seeds. As shown in Fig.S2A, the content of soluble sugar was increased as the time prolonged and peaked at 48 h. Moreover, the soluble sugar content in the seeds treated with SAEW was significantly higher than it in control at 48, 72 and 96 h. Thereinto, the content of soluble sugar at 48 h was increased by 48.87% with SAEW treatment. On the contrary, the changing trends of starch and protein were decreased as the time prolonged. In addition, SAEW further accelerated the reduction of starch and protein contents after germination for 48 h (Fig. S2B and C). The contents of starch and protein were respectively reduced by 47.75% and 26.40%. Beyond that, the fat content was gradually reduced as time went by and not influenced by SAEW treatment (Fig.S2D). All the results showed that 60 ppm SAEW had the capacity for expediting the germination of watermelon seeds.

Seed germination could also cause the change of enzymatic activities. In order to further explored the effect of SAEW on enzymes, the activities of SOD, POD and CAT were determined in the seed of watermelon. As the time of germination increased, the activity of SOD increased and reached the peak on 48 h in both the Con and the SAEW



**Fig. 1.** The effects of different concentrations of slightly acidic electrolyzed water (SAEW) on watermelon seed germination. (A) performed the phenotype of seed germination with different treatments. The germination rate (B), germination energy (C), germination index (D) and vitality index (E) influenced by different concentrations of SAEW were shown. Watermelon seeds were respectively soaked in deionized water (Control which was presented as Con) or different concentrations of SAEW (20, 40, 60, 80 ppm) at 25°C for 5 h. The germination indices were measured at indicated time (24, 48, 72, 96 and 120 h). Data are shown as the means  $\pm$  SE ( $n=30$ ). Different letters indicate significant differences at  $P < 0.05$  according to Duncan's multiple range test.

treatment. Meanwhile, the activity of SOD was significantly improved by SAEW except 24 h. The activity of SOD was respectively increased over 50% at different germination times (Fig.S3A). In addition, the activities of POD and CAT had the similar change trend. With extended time of germination, the activities of POD and CAT were gradually increased, which remained constantly after 72 and 48 h respectively. Thereinto, the activity of POD was nearly tripled by SAEW after germinated 48 h and remained constantly after 96 h (Fig.S3B and C). In conclusion, the up-regulation of SOD, POD and CAT activities was induced by SAEW, which was benefit for seed germination.


### 3.3. Effect of slightly acidic electrolyzed water (SAEW) on the concentrations of endogenous phytohormone in watermelon seed

ABA could induce and maintain seed dormancy and GA is responsible for the induction of seed germination (Shu et al. 2016; Tuan et al. 2018). In order to figure out whether these plant hormones contributed to the SAEW-induced germination, the concentrations of ABA, GA, IAA and Zeatin were detected severally. As shown in Fig.2A, ABA concentration was markedly decreased by SAEW. Particularly at 24 h, the ABA concentration was reduced by 45.18% with SAEW treatment. Meanwhile, the total GA concentration was gradually improved with the germinating time (Fig.2B). The total GA was the sum of GA1, GA3 and GA4, which was separately shown in Fig.S4A-C. At 48h, the total GA concentration was improved by 64.85% with SAEW treatment. Then, the ratio of GA to ABA was remarkably risen by SAEW. Besides, the concentration of Zeatin was consistent with the changing trend of GA, which was gradually increased as germination time went by and further improved by SAEW (Fig.S4D). In addition, the trend of IAA concentration first increasing and then decreasing was showed with the germination time, and SAEW-induced the increase of IAA concentration was shown at 48 and 72 h (Fig.S4E). In conclusion, the SAEW-induced germination maybe caused by the changing of endogenous ABA and GA.

### 3.4. Effect of slightly acidic electrolyzed water (SAEW) on the expressions of key genes involved in abscisic acid (ABA)/gibberellins (GA) biosynthesis and catabolism


The endogenous ABA and GA level in seed is determined by a balance between its biosynthesis and catabolism. In order to further probe into the interrelation between ABA/GA and SAEW-induced germination, the transcriptional level of relative genes involved in ABA/GA biosynthesis and catabolism were analyzed respectively (Fig.3A). The expression of *ZEP*, encoded a single copy zeaxanthin epoxidase gene that functions in first step of the biosynthesis of ABA, was reduced by SAEW. As the extension of germination time, the expression of *ZEP* showed a descending trend gradually in both the Con and the SAEW-treatment group. *NCEDs* encodes 9-cis-epoxycarotenoid dioxygenase which is a key enzyme in the biosynthesis of ABA (Tan et al., 2003; Lefebvre et al., 2006). Then, the expressions of *NCED2*, *NCED3* and *NCED6* showed the same changing tendency as the *ZEP*, which were declined by degrees no matter in the Con or in SAEW treatment. Thereinto, the expressions of *NCED3* were further lowered by SAEW, but the expressions of *NCED2* and *NCED6* were lower in the control than it in SAEW-treatment. In addition, the expression of *AAO3*, coded the aldehyde oxidase delta isoform catalyzing the final step in ABA biosynthesis, were gradually reduced as the germination of seed and slightly lowered at 48 and 72 h in the control. What's more, the expressions of *CYP707A1* and *CYP707A2*, encoded a protein with ABA 8'-hydroxylase activity involved in ABA catabolism, showed an upward tendency as the extension of germination time and were further significantly elevated by the SAEW treatment, which may contribute to the reduction of ABA content in seeds.

When it comes to the GA biosynthesis and catabolism, the expressions of *GA3ox1*, *GA3ox2*, *GA13ox* and *GA20ox2*, encoded gibberellin oxidases involved in GA biosynthesis, were gradually increased as the process of germination and further induced by SAEW (Fig.3B).



**Fig. 2.** The analyses of endogenous abscisic acid (ABA) and gibberellin (GA) concentrations in watermelon seed after soaked in slightly acidic electrolyzed water (SAEW). The concentrations of ABA (A), total GA (B) and the ratio of GA/ABA (C) influenced by 60 ppm SAEW were shown. Watermelon seeds were respectively soaked in deionized water (Control) or 60 ppm SAEW at 25°C for 5 h. The concentrations of plant hormones were measured at indicated time (0, 24, 48 and 72h). Data are shown as the means  $\pm$  SE ( $n=3$ ). The asterisk indicated significant differences at  $P < 0.05$  according to Duncan's multiple range test and the significant differences were compared between Control and 60 ppm SAEW treatment at indicated time.

Meanwhile, the expressions of *GA2ox3*, *GA2ox6* and *GA3ox1* involved in GA catabolism showed a decline tendency induced by SAEW. However, the effect of SAEW on the expressions of GA-related genes was not significant. In conclusion, the SAEW-induced germination of watermelon seeds may arise from the accelerating catabolism of ABA by increasing the transcriptional levels of *CYP707A1* and *CYP707A2*, which ultimately led to a lower ABA accumulation to break the seed dormancy.




**Fig. 3.** The effects of slightly acidic electrolyzed water (SAEW) treatment on the expressions of genes involved in biosynthesis and catabolism of abscisic acid (ABA) and gibberellin (GA). The key genes of ABA (A) and GA (B) influenced by 60 ppm SAEW were shown. Watermelon seeds were respectively soaked in deionized water (Con) or 60 ppm SAEW at 25°C for 5 h. The expressions of genes were measured at indicated time (0, 24, 48 and 72 h). Data are shown as the means  $\pm$  SE ( $n=9$ ). The asterisk indicated significant differences at  $P < 0.05$  according to Duncan's multiple range test and the significant differences were compared between Con and 60 ppm SAEW treatment at indicated time.

### 3.5. The endogenous reactive oxygen species (ROS) of watermelon seeds was changed by slightly acidic electrolyzed water (SAEW)

Several researches revealed that ROS identified as non-enzymatic factors play a key but largely elusive role in the repression of seed germination in some plant species like *Arabidopsis thaliana* and lettuce (Müller et al., 2009; Lin et al., 2013; Ortiz-Espín et al., 2017; Choudhury et al., 2017; Luo et al., 2021; Zhang et al., 2014; Yang et al., 2020). In order to ensure the roles of ROS in the SAEW-induced germination, the ROS accumulation in different treatments was examined by the content

analyses and staining of  $H_2O_2$  and  $O_2^-$  respectively. As it shown in Fig. 4A-C, the  $H_2O_2$  and  $O_2^-$  contents were significantly reduced by SAEW. The lighter brown and bluish violet were found in SAEW treatment (Fig. 4A). Meanwhile, the results of content determination were in accord with the staining evidence. The contents of  $H_2O_2$  and  $O_2^-$  were markedly decreased by SAEW at 48 h and 96 h. Thereinto, compared with the control, the content of  $H_2O_2$  and  $O_2^-$  was reduced by 54.29% and 55.56% at 48 h respectively with SAEW treatment.

Most of *Rbohs* are shown involved in the process of the response to abiotic stress and stomatal closure (Kámán-Tóth et al., 2019; Ma et al.,



**Fig. 4.** The effects of slightly acidic electrolyzed water (SAEW) on the endogenous reactive oxygen species (ROS). Histochemical staining of  $O_2^-$  (superoxide radicals) by NBT (nitro blue tetrazolium) and  $H_2O_2$  (hydrogen peroxide) by DAB (3,3'-diaminobezidine) were shown in (A). The contents of  $H_2O_2$  (B) and  $O_2^-$  (C) were influenced by SAEW. (D) and (E) showed the relative expressions of *RbohC* and *RbohD* with different treatment. Watermelon seeds were respectively soaked in deionized water (Control) or 60 ppm SAEW at 25°C for 5 h. The concentrations of ROS and the expressions of genes were measured at indicated time (0, 24, 48, 72, 96 and 120 h). Data are shown as the means  $\pm$  SE ( $n=9$ ). The asterisk indicated significant differences at  $P < 0.05$  according to Duncan's multiple range test and the significant differences were compared between Control and 60 ppm SAEW treatment at indicated time. And the histochemical staining was performed at 72 h in the radicles of watermelon seeds.


2012; Scuffi et al., 2018; Wang et al., 2016). However, only *RbohB* and *RbohC* have been identified so far in the process of seed development and germination (Müller et al., 2009; Yang et al., 2020). In the study, *RbohC*, *RbohD*, *RbohF* and *RbohH* were expressed in watermelon seeds (Fig. 4D, E and Fig.S5). Meanwhile, the relative expressions of *Rbohs* were detected at indicated time. Thereinto, the expressions of *RbohC* and *RbohD* were up-regulated first but then them went down with the extension of germination time (Fig. 4D and E). The expressions of *RbohC* were significantly reduced by SAEW after 48 h and the expressions of *RbohD* were markedly inhibited from 0 h to 120 h. That's to say, *RbohD* had lower transcriptional level after seeds imbibed with SAEW than it with deionized water (Con). Beyond that, the expressions of *RbohF* and *RbohH* were measured and shown in Fig.S5. However, it revealed that the expressions of *RbohF* and *RbohH* were not significantly affected by SAEW. To sum up, the reduced ROS accumulation may give rise to the

promoted germination caused by SAEW in watermelon.

SAEW could significantly influence the ROS accumulation (Fig. 4). Therefore, the role of ROS in SAEW-induced germination in watermelon seeds were further explored. DMTU (dimethyl thiourea), an inhibitor of ROS, was used to soak the watermelon seeds. As shown in Fig. 5, DMTU and SAEW could significantly promote the germination of seeds. Furthermore, the co-treatment of SAEW+DMTU accelerated the promotion of germination. The germination energy, germination index and vitality index were markedly increased by SAEW+DMTU. It suggested that the reduced accumulation of ROS may contribute to the SAEW-induced germination.

#### 4. Discussion

To improve the seed germination of watermelon, some treatments



**Fig. 5.** The effects of reactive oxygen species inhibitor (*N,N'*-Dimethylthiourea, DMTU) and slightly acidic electrolyzed water (SAEW) on the germination of watermelon seeds. The phenotype (A), germination energy (B), germination index (C) and vitality index (D) were shown. Watermelon seeds were respectively soaked in deionized water (Control), 60 ppm SAEW, 60 mM DMTU and SAEW+DMTU mixture at 25°C for 5 h. The germination-related indicators were measured. Data are shown as the means  $\pm$  SE ( $n = 30$ ). Different letters indicated significant differences at  $P < 0.05$  according to Duncan's multiple range test.


including seed coat removal, scarification, and seed nicking have been proved to have the capacity of enhancing seed germination and seedling vigor. However, some varieties of watermelon like triploid seeds still had lower seedling vigor than that of diploids (Grange et al., 2000; Acharya et al., 2020). Therefore, the novel methods to improve and uniform seed germination and seedling vigor are needed. Up to now, some exogenous substances such as salicylic acid, oxygenated brackish water and plasma activated water were shown to markedly improve the seed germination in sunflower (*Helianthus annuus* L.), wheat (*Triticum aestivum* L.), alfalfa (*Medicago sativa*) and mung bean (*Vigna radiata*) (Huang et al., 2020; Zhu et al., 2020; Machado-Moreira et al., 2020). However, the research of using exogenous substances to improve the seed germination of watermelon is scarce until now. This study assessed the roles of SAEW in the seed germination of watermelon by the analyses of germination index, nutrient contents, endogenous plant hormone contents and the expression level of related genes. These data manifested that SAEW had the capacity for the promotion of seed germination may be caused by accelerating the catabolism of ABA by the induced-expressions of *CYP707A1* and *CYP707A2*, which further led to a lower ABA accumulation than it in the control. Meanwhile, the decreased-expressions of *RbohC* and *RbohD* caused a less accumulation of ROS and further broke the seed dormancy (Fig. 6).

#### 4.1. Abscisic acid catabolism was involved in the slightly acidic electrolyzed water-induced the promotion of seed germination in watermelon

The germination capacity of watermelon seeds can be reflected by germination parameters, including germination rate, germination energy, germination index and vitality index (Wen et al., 2018). In this study, the germination parameters of SAEW-treatment were significantly higher than those of control, indicating that SAEW could promote

seed germination of watermelon (Fig.1). As the SAEW concentration increased, the acceleration of seed germination was increased by degrees. Compared with the control, 60 ppm SAEW could induce seed germinated one day in advance (Fig.1). In addition, when the time of seed soaked with 60 ppm SAEW from 1 to 10 h, the germination parameters were improved gradually and then decreased (Fig.S1). Meanwhile, SAEW caused a mass of changes to nutrient substances and enzymatic activities in watermelon seeds (Fig.S2-S3), which performed the ability of SAEW to accelerate the seed germination. To sum up, soaked in 60 ppm SAEW with 5 h could significantly promote water seed germination.

ABA and GA are the two major phytohormones regulating seed dormancy and germination of plants. ABA could induce and maintain seed dormancy and GA is responsible for the induction of seed germination (Shu et al. 2016; Tuan et al. 2018). Moreover, the degree of seed dormancy correlates with endogenous ABA levels in imbibed seeds rather than in dry seeds in various species, such as in *Arabidopsis thaliana*, lettuce, barley and tobacco (Ali-Rachedi et al., 2004; Gonai et al., 2004; Jacobsen et al., 2002; Grappin et al., 2000). In addition, the ABA accumulation in dry seed decreased rapidly after seed imbibition (Okamoto et al., 2006). In this study, it revealed that SAEW could dramatically decrease the ABA accumulation while increase the GA content (Fig.2), which gave rise to the accelerated germination. The endogenous ABA and GA level in seed is determined by a balance between its biosynthesis and catabolism. The ABA is mainly biosynthesized by *NCEDs*, *ZEP*, *AAO* and catabolized by *CYP707As* (Nambara and Marion-Poll, 2005). Meanwhile, the GA is biosynthesized by *GA3ox*, *GA13ox*, *GA20ox* and catabolized by *GA2ox* respectively (Tuan et al., 2020). In addition, some previous studies have revealed the roles of genes encoding these enzymes in regulating seed GA levels and germination in several plant species including *Arabidopsis thaliana* and wheat (Ogawa et al., 2003; Mitchum et al., 2006; Kashiwakura et al., 2016;



**Fig. 6.** The mechanistic model of slightly acidic electrolyzed water (SAEW)-induced germination in watermelon seeds. The germination of seeds was significantly promoted by 60 ppm SAEW. SAEW could increase the expressions of *CYP707A1* and *CYP707A2*, which accelerated the ABA catabolism and then reduced the accumulation of ABA. In addition, the expressions of *RbohC* and *RbohD* were further decreased, which gave rise to the reduced accumulation of ROS. Ultimately, the germination of watermelon seeds was improved by the imbibition with SAEW. However, the interrelation between ABA and ROS in seed germination, pointed by the yellow double-headed arrow, need to be further explored. ER stood for the endoplasmic reticulum and PM represented the plasma membrane.

Izydorczyk et al., 2018). According to the expression analyses of primary genes, it suggested that SAEW mostly decreased the transcriptional levels of genes involved in ABA biosynthesis such as *ZEP*, *NCED3* and *AAO3* (Fig. 3A). What's more, the transcriptions of *CYP707A1* and *CYP707A2* were significantly increased by SAEW, which may contribute to the reduction of ABA content in seeds. In addition, as the time went by, the expressions of *CYP707A1* and *CYP707A2* were gradually increased in both Con and SAEW treatment (Fig. 3A). It was consistent with the result of previous study which revealed that the transcription of *CYP707A2* accumulated predominantly in the dry seed and immediately up-regulated after seed imbibition (Kushiro et al., 2004). Beyond that, transcriptions of *GA3ox*, *GA13ox*, *GA20ox*, which were responsible for GA biosynthesis, were mostly induced by SAEW and gradually elevated with the extension of time (Fig. 3B). All the results suggested that SAEW-induced germination may arise from the depressed ABA accumulation while raised GA accumulation. Moreover, SAEW reduced ABA content in seeds mainly by the promotion of ABA catabolism.

#### 4.2. Reactive oxygen species may play a role in the accelerated seed germination caused by slightly acidic electrolyzed water

ROS are increasingly identified as non-enzymatic factors in seed germination in some plant species like *Arabidopsis thaliana* and lettuce (Müller et al., 2009; Zhang et al., 2014; Choudhury et al., 2017; Yang et al., 2020). In this study, ROS accumulation was reduced by SAEW treatment, which was performed by the analyses of  $H_2O_2$  and  $O_2^-$

concentrations (Fig. 4). Moreover, we found that DMTU could significantly improve the seed germination, which was in accordance with the promoted germination caused by SAEW treatment (Fig. 5). That's to say, the reduced ROS accumulation may contribute to the improvement of germination induced by SAEW. The plasma membrane NADPH oxidases, encoded by the *Rbohs* family, are key enzymes for the production of ROS. Most of *Rbohs* are shown involved in the process of the response to abiotic stress and stomatal closure (Ma et al., 2012; Wang et al., 2016; Scuffi et al., 2018; Kámán-Tóth et al., 2019). However, only *RbohB* and *RbohC* have been identified so far in the process of seed development and germination (Müller et al., 2009; Yang et al., 2020). In the study, *RbohC*, *RbohD*, *RbohF* and *RbohH* were expressed in watermelon seeds (Fig. 4D, E and Fig. S6). Thereinto, the expressions of *RbohC* and *RbohD* were up-regulated first but then went down with the extension of germination time (Fig. 4D and E). Furthermore, SAEW had the capacity for inhibition of *RbohC* and *RbohD* transcript, which may lead to the reduced accumulation of  $H_2O_2$  and  $O_2^-$  (Fig. 4B and C). However, *RbohF* and *RbohH* were not significantly influenced by SAEW (Fig. S6). To sum up, the reduced ROS accumulation may give rise to the promoted germination caused by SAEW in watermelon. Even more to the point, the previous research demonstrated that the increased expression of *RbohD* is essential for ABA-mediated induction of ROS accumulation (Kwak et al., 2003). However, the relationship among endogenous ABA, ROS and seed germination in watermelon need to be further explored.

#### 5. Authors' contributions

XW and KC contributed to the conception of the study. XW designed and performed the experiments. XW analyzed the data and wrote the manuscript. CW and ZB helped with graphical edit. ZY and LM modified the grammatical and format errors. LX and EB critically reviewed the manuscript.

#### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this research.

#### Funding

The authors would like to thank Jiangsu Postdoctoral Research Founding Program (2021K469C), Jiangsu Agricultural Science and Technology Innovation Fund [CX(21)2022] and National Natural Science Foundation of China (Grant No.31801903).

#### Acknowledgement

This research was supported financially by the Key Laboratory of Protected Horticultural Engineering in Northwest, Ministry of Agriculture, PR China.

#### Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.scienta.2021.110581.

#### References

- Acharya, P., Jayaprakash, G., Crosby, K., Jifon, J., Patil, B., 2020. Nanoparticle-mediated seed priming improves germination, growth, yield, and quality of watermelons (*Citrullus lanatus*) at multi-locations in Texas. Sci. Rep. 10 (1), 5037. <https://doi.org/10.1038/s41598-020-61696-7>.
- Ali-Rachedi, S., Ouinot, D., Wagner, M., Bonnet, M., Sotta, B., Grappin, P., Jullien, M., 2004. Changes in endogenous abscisic acid levels during dormancy release and maintenance of mature seeds: studies with the Cape Verde Islands ecotype, the dormant model of *Arabidopsis thaliana*. Planta 219 (3), 479–488. <https://doi.org/10.1007/s00425-004-1251-4>.

- Chen, K., Arora, R., 2013. Priming memory invokes seed stress-tolerance. *Environ. Exp. Bot.* 94, 33–45. <https://doi.org/10.1016/j.enexpbot.2012.03.005>.
- Choudhury, F., Rivero, R., Blumwald, E., Mittler, R., 2017. Reactive oxygen species, abiotic stress and stress combination. *Plant J.* 90 (5), 856–867. <https://doi.org/10.1111/tpj.13299>.
- Cutler, S., Rodriguez, P., Finkelstein, R., Abrams, S., 2010. Abscisic acid: Emergence of a core signaling network. *Annu Rev Plant Biol* 61 (1), 651–679. <https://doi.org/10.1146/annurev-arplant-042809-112122>.
- FAOSTAT, 2017. Retrieved from: Available online: <http://www.fao.org/faostat/en/#data/QC>.
- Dabirian, S., Miles, C.A., 2017. Increasing survival of splice-grafted watermelon seedlings using a sucrose application. *HortScience* 52 (4), 579–583. <https://doi.org/10.21273/HORTSCI11667-16>.
- Dahlquist-Willard, R., Marshall, M., Betts, S., Tuell-Todd, C., Vandergheynst, J., Stapleton, J., 2016. Development and validation of a Weibull-Arrhenius model to predict thermal inactivation of black mustard (*Brassica nigra*) seeds under fluctuating temperature regimens. *Biosyst. Eng.* 151 (11), 350–360. <https://doi.org/10.1016/j.biosystemseng.2016.09.015>.
- Gonai, T., Kawahara, S., Tougou, M., Satoh, S., Hashiba, T., Hirai, N., Kawaide, H., Kamiya, Y., Yoshioka, T., 2004. Abscisic acid in the thermoinhibition of lettuce seed germination and enhancement of its catabolism by gibberellin. *J Exp Bot* 55 (8), 111–118. <https://doi.org/10.1093/jxb/erh023>.
- Grange, S., Leskovar, D., Pike, L., Cobb, B., 2000. Excess moisture and seedcoat nicking influence germination of triploid watermelon. *HortScience* 35 (7), 1355–1356. <https://doi.org/10.1086/386232>.
- Grappin, P., Bouinot, D., Sotta, B., Miginiac, E., Jullien, M., 2000. Control of seed dormancy in *Nicotiana plumbaginifolia*: post-imbibition abscisic acid synthesis imposes dormancy maintenance. *Planta* 210 (2), 279–285. <https://doi.org/10.1007/pl00008135>.
- Guan, X., Ramaswamy, H., Zhang, B., Lin, B., Wang, S., 2020. Influence of moisture content, temperature and heating rate on germination rate of watermelon seeds. *Sci. Hortic.* 272, 109528. <https://doi.org/10.1016/j.scienta.2020.109528>.
- Han, Y., Zhang, J., Chen, X., Gao, Z., Xuan, W., Xu, S., Ding, X., Shen, W., 2007. Carbon monoxide alleviates cadmium-induced oxidative damage by modulating glutathione metabolism in the roots of *Medicago sativa*. *New Phytol* 177, 155–166. <https://doi.org/10.1111/j.1469-8137.2007.02251.x>.
- Huang, B., Xu, S., Wei, X., Li, M., Cao, Z., Liu, K., Ling, T., Shen, W., 2006. Carbon monoxide alleviates salt-induced oxidative damage in wheat seedling leaves. *J. Integr. Plant Biol.* 48 (3), 249–254. <https://doi.org/10.1111/j.1744-7909.2006.00220.x>.
- Huang, X., Zhu, S., Zhou, X., He, J., Yu, Y., Ye, Z., 2021. Preservative effects of the combined treatment of slightly acidic electrolyzed water and ice on pomfret. *Int. Jagr. Biol. Eng.* 14 (1), 230–236. <https://doi.org/10.25165/j.ijabe.20211401.5967>.
- Huang, Y., Cai, S., Ruan, X., Chen, S., Mei, G., Ruan, G., Cao, D., 2020. Salicylic acid enhances sunflower seed germination under  $Zn^{2+}$  stress via involvement in  $Zn^{2+}$  metabolic balance and phytohormone interactions. *Sci. Hortic.* 275, 109702. <https://doi.org/10.1016/j.scienta.2020.109702>.
- Hubbard, K., Nishimura, N., Hitomi, K., Getzoff, E., Schroeder, J., 2010. Early abscisic acid signal transduction mechanisms: Newly discovered components and newly emerging questions. *Genes Dev* 24, 1695–1708. <https://doi.org/10.1101/gad.1953910>.
- Hubbard, M., Germida, J., Vujanovic, V., 2012. Fungal endophytes improve wheat seed germination under heat and drought stress. *Botany* 90 (2), 137–149. <https://doi.org/10.1139/b11-091>.
- Huehne, P., Bhinjija, K., 2012. Application of cryoprotectants to improve low temperature storage survival of orchid seeds. *Sci. Hortic.* 135, 186–193. <https://doi.org/10.1016/j.scienta.2011.11.026>.
- Izydorczyk, C., Nguyen, T., Jo, S., Son, S., Tuan, P., Ayele, B., 2018. Spatiotemporal modulation of abscisic acid and gibberellin metabolism and signalling mediates the effects of suboptimal and supraoptimal temperatures on seed germination in wheat (*Triticum aestivum* L.). *Plant Cell and Environ* 41 (5), 1022–1037. <https://doi.org/10.1111/pce.12949>.
- Jacobsen, J., Pearce, D., Poole, A., Pharis, R., Mander, L., 2002. Abscisic acid, phaseic acid and gibberellin contents associated with dormancy and germination in barley. *Physiol. Plant.* 115 (3), 428–441. <https://doi.org/10.1034/j.1399-3054.2002.1150313.x>.
- Jawad, U., Gao, L., Gebremeskel, H., Saifdar, L., Liu, W., 2020. Expression pattern of sugars and organic acids regulatory genes during watermelon fruit development. *Sci. Hortic.* 265, 109102. <https://doi.org/10.1016/j.scienta.2019.109102>.
- Jiao, S., Zhong, Y., Deng, Y., 2016. Hot air-assisted radio frequency heating effects on wheat and corn seeds: quality change and fungi inhibition. *J. Stored Prod. Res.* 69, 265–271. <https://doi.org/10.1016/j.jspr.2016.09.005>.
- Jung, S., Ko, B., Jang, H., Park, H., Oh, S., 2017. Effects of slightly acidic electrolyzed water ice and grapefruit seed extract ice on shelf life of brown sole (*Pleuronectes herzensteini*). *Food Sci. Biotechnol.* 27 (1), 261–267. <https://doi.org/10.1007/s10068-017-0198-8>.
- Kámán-Tóth, E., Dankó, T., Gullner, G., Bozsó, Z., Palkovics, L., Pogány, M., 2019. Contribution of cell wall peroxidases and NADPH oxidase-derived reactive oxygen species to *Alternaria brassicicola*-induced oxidative burst in Arabidopsis. *Mol. Plant Pathol.* 20, 485–499. <https://doi.org/10.1111/mpp.12769>.
- Kashiwakura, Y., Kobayashi, Daisuke, Jikumaru, Yusuke, Takebayashi, Yumiko, Nambara, Eiji, Mitsunori, Kamiya, Yuji, Kushiro, Tetsuo, Kawakami, Naoto, 2016. Highly sprouting-tolerant wheat grain exhibits extreme dormancy and cold imbibition-resistant accumulation of abscisic acid. *Plant Cell Physiol* 57 (4), 715–732. <https://doi.org/10.1093/pcp/pcw051>.
- Kwak, J., Mori, I., Zhen, M., Leonhardt, N., Torres, M., Dangl, J., Bloom, R., Bodde, S., Jones, J., Schroeder, J., 2003. NADPH oxidase *ArbboD* and *ArbboH* genes function in ROS-dependent ABA signaling in Arabidopsis. *EMBO J* 22 (11), 2623–2633. <https://doi.org/10.1093/emboj/cdg277>.
- Kucera, B., Cohn, M., Leubner-Metzger, G., 2005. Plant hormone interactions during seed dormancy release and germination. *Seed Sci Res* 15 (4), 281–307. <https://doi.org/10.1079/SSR2005218>.
- Kushiro, T., Okamoto, M., Nakabayashi, K., Kitamura, S., Hirai, N., Kamiya, Y., Nambara, E., 2004. The Arabidopsis cytochrome P450 CYP707A encodes ABA 8'-hydroxylases: key enzymes in ABA catabolism. *EMBO J* 23, 1647–1656.
- Lefebvre, V., North, H., Frey, A., Sotta, B., Seo, M., Okamoto, M., Nambara, E., Marion-Poll, A., 2006. Functional analysis of Arabidopsis *NCED6* and *NCED9* genes indicates that ABA synthesized in the endosperm is involved in the induction of seed dormancy. *Plant J* 45 (3), 309–319. <https://doi.org/10.1038/sj.emboj.7600121>.
- Li, H., Chang, J., Chen, H., Wang, Z., Gu, X., Wei, C., Zhang, Y., Ma, J., Yang, J., Xian, Z., 2017. Exogenous melatonin confers salt stress tolerance to watermelon by improving photosynthesis and redox homeostasis. *Front. Plant Sci.* 8, 00295. <https://doi.org/10.3389/fpls.2017.00295>.
- Lin, Y., Yang, L., Paul, M., Zu, Y., Tang, Z., 2013. Ethylene promotes germination of Arabidopsis seed under salinity by decreasing reactive oxygen species: Evidence for the involvement of nitric oxide simulated by sodium nitroprusside. *Plant Physiol. Biochem.* 73, 211–218. <https://doi.org/10.1016/j.plaphy.2013.10.003>.
- Luo, X., Dai, Y., Zheng, C., Yang, Y., Chen, W., Wang, Q., Umashankar, C., Du, J., Liu, W., Shu, K., 2021. The ABI4-RboHD/VT2C regulatory module promotes reactive oxygen species (ROS) accumulation to decrease seed germination under salinity stress. *New Phytol* 229 (2), 950–962. <https://doi.org/10.1111/nph.16921>.
- Lv, J., Lin, B., Han, X., Xu, D., Ding, S., Li, C., Ge, Y., Li, J., 2021. Effects of 1-MCP treatment on sprouting and preservation of ginger rhizomes during storage at room temperature. *Food Chem.* 349, 129004. <https://doi.org/10.1016/j.foodchem.2021.129004>.
- Ma, L., Zhang, H., Sun, L., Jiao, Y., Zhang, G., Miao, C., Hao, F., 2012. NADPH oxidase *ArbboD* and *ArbboH* function in ROS-dependent regulation of  $Na^+/K^+$  homeostasis in Arabidopsis under salt stress. *J. Exp. Bot.* 63 (1), 305–317. <https://doi.org/10.1093/jxb/err280>.
- Machado-Moreira, B., Tiwari, B., Richards, K., Abram, F., Burgess, C., 2020. Application of plasma activated water for decontamination of alfalfa and mung bean seeds. *Food Microbiol* 96 (4), 103708. <https://doi.org/10.1016/j.fm.2020.103708>.
- Malcolm, P., Holford, P., McGlasson, W., Newman, S., 2003. Temperature and seed weight affect the germination of peach rootstock seeds and the growth of rootstock seedlings. *Sci. Hortic.* 98 (3), 247–256. [https://doi.org/10.1016/S0304-4238\(03\)00017-7](https://doi.org/10.1016/S0304-4238(03)00017-7).
- Miransari, M., Smith, D., 2014. Plant hormones and seed germination. *Environ. Exp. Bot.* 99 (3), 110–121. <https://doi.org/10.1016/j.enexpbot.2013.11.005>.
- Mitchum, M., Yamaguchi, S., Hanada, A., Kuwahara, A., Yoshioka, Y., Kato, T., Tabata, S., Kamiya, Y., Sun, T., 2006. Distinct and overlapping roles of two gibberellin 3-oxidases in Arabidopsis development. *Plant J* 45 (5), 804–818. <https://doi.org/10.1111/j.1365-313X.2005.02642.x>.
- Mohammed, A., Abdelrahim, H., Soad, K., Alkhalfah, D., Hozzein, W., Selim, S., Abdelgawad, H., Kamis, G., 2020. Influence of elevated  $CO_2$  on nutritive value and health-promoting prospective of three genotypes of Alfalfa sprouts (*Medicago Sativa*). *Food Chem* 340, 128147. <https://doi.org/10.1016/j.foodchem.2020.128147>.
- Morris, D., 1948. Quantitative determination of carbohydrates with dreywood's anthrone reagent. *Science* 107, 254–255. <https://doi.org/10.1126/science.107.2775.254>.
- Müller, K., Carstens, A., Linkies, A., Torres, M., Leubner-Metzger, G., 2009. The NADPH-oxidase *AtrboH* plays a role in Arabidopsis seed after-ripening. *New Phytol* 184, 885–897. <https://doi.org/10.1111/j.1469-8137.2009.03005.x>.
- Ogawa, M., Hanada, A., Yamauchi, Y., Kuwahara, A., Kamiya, Y., Yamaguchi, S., 2003. Gibberellin biosynthesis and response during Arabidopsis seed germination. *Plant Cell* 15 (7), 1591–1604. <https://doi.org/10.1105/tpc.011650>.
- Okamoto, M., Kuwahara, A., Seo, M., Kushiro, T., Asami, T., Hirai, N., Kamiya, Y., Koshiba, T., Nambara, E., 2006. CYP707A1 and CYP707A2, which encode abscisic acid 8'-hydroxylases, are indispensable for proper control of seed dormancy and germination in Arabidopsis. *Plant Physiol* 141 (1), 97–107. <https://doi.org/10.1104/pp.106.079475>.
- Pérez-Jiménez, M., Cantero-Navarro, E., Pérez-Alfocea, F., Cos-Terrer, J., 2014. Endogenous hormones response to cytokinins with regard to organogenesis in explants of peach (*Prunus persica* L. Batsch) cultivars and rootstocks (*P. persica* × *Prunus dulcis*). *Plant Physiol. Bioch.* 84, 197–202. <https://doi.org/10.1016/j.plaphy.2014.09.014>.
- Scuffi, D., Nietzel, T., Fino, L., Meyer, A., Lamattina, L., Schwarzländer, M., Laxalt, A., García-Mata, C., 2018. Hydrogen sulfide increases production of NADPH oxidase-dependent hydrogen peroxide and phospholipase D-derived phosphatidic acid in guard cell signaling. *Plant Physiol* 176 (3), 2532–2542. <https://doi.org/10.1104/pp.17.01636>.
- Su, N., Wu, Q., Cui, J., 2016. Increased sucrose in the hypocotyls of radish sprouts contributes to nitrogen deficiency-induced anthocyanin accumulation. *Front Plant Sci* 7 (927), 1976. <https://doi.org/10.3389/fpls.2016.01976>.
- Tan, B., Josephy, L., Dengy, W., Liu, L., Li, Q., 2003. Molecular characterization of the Arabidopsis 9-cis epoxycarotenoid dioxygenase gene family. *Plant J* 35, 44–56. <https://doi.org/10.1046/j.1365-313X.2003.01786.x>.
- Tuan, P., Nguyen, T., Jordan, M., Ayele, B., 2020. A shift in abscisic acid/gibberellin balance underlies retention of dormancy induced by seed development temperature. *Plant Cell Environ.* 1–15. <https://doi.org/10.1111/pce.13963>.
- Wang, S., Bao, W., Zhang, F., Qi, F., Nan, S., He, J., Zhu, S., Ye, Z., 2017. Disinfection kinetics of slightly acidic electrolyzed water to freshwater under the condition of

- dynamic hybrid. J. Clean Prod. 174, 1136–1146. <https://doi.org/10.1016/j.jclepro.2017.11.033>.
- Wang, X., Zhang, M., Wang, Y., Gao, Y., Li, R., Wang, G., Li, W., Liu, W., Chen, K., 2016. The plasma membrane NADPH oxidase *OsRbohA* plays a crucial role in developmental regulation and drought-stress response in rice. Physiol. Plant. 156 (4), 421–443. <https://doi.org/10.1111/pp.12389>.
- Wang, Y., Li, L., Ye, T., Zhao, S., Liu, Z., Feng, Y., Wu, Y., 2011. Cytokinin antagonizes ABA suppression to seed germination of Arabidopsis by down-regulating *ABF5* expression. Plant J. 68 (2), 249–261. <https://doi.org/10.1111/j.1365-313X.2011.04683.x>.
- Wei, H., Wang, M., Jeong, B., 2020. Effect of supplementary lighting duration on growth and activity of antioxidant enzymes in grafted watermelon seedlings. Agronomy 10 (3), 337. <https://doi.org/10.3390/agronomy10030337>.
- Wen, D., Xu, H., Xie, L., He, M., Hou, H., Wu, C., Li, Y., Zhang, C., 2018. Effects of nitrogen level during seed production on wheat seed vigor and seedling establishment at the transcriptome level. Int. J. Mol. Sci. 19 (11), 3417. <https://doi.org/10.3390/ijms19113417>.
- Wu, X., Zhu, Z., Chen, J., Huang, Y., Liu, Z., Zou, J., Chen, Y., Su, N., Cui, J., 2019. Transcriptome analysis revealed pivotal transporters involved in the reduction of cadmium accumulation in Pak choi (*Brassica chinensis* L.) by exogenous hydrogen-rich water. Chemosphere 216, 684–697. <https://doi.org/10.1016/j.chemosphere.2018.10.152>.
- Yang, X., Zhang, F., Yang, M., He, Y., Li, Z., Yang, J., Wang, X., 2020. The NADPH oxidase *LsRbohC1* plays a role in lettuce (*Lactuca sativa*) seed germination. Plant Physiol. Bioch. 154, 751–757. <https://doi.org/10.1016/j.plaphy.2020.05.042>.
- Ye, Z., Wang, S., Chen, T., Gao, W., Zhu, S., He, J., Han, Z., 2017. Inactivation mechanism of *Escherichia coli* induced by slightly acidic electrolyzed water. Sci. Rep. 7 (1), 6279. <https://doi.org/10.1038/s41598-017-06716-9>.
- Yu, D., Shrestha, B., Baik, O., 2016. Radio frequency (RF) control of red flour beetle (*Tribolium castaneum*) in stored rapeseeds (*Brassica napus* L.). Biosyst. Eng 151, 248–260. <https://doi.org/10.1016/j.biosystemseng.2016.09.006>.
- Yu, Y., Wang, J., Li, S., Kakan, X., Zhou, Y., Miao, Y., Wang, F., Qin, H., Huang, R., 2019. Ascorbic acid integrates the antagonistic modulation of ethylene and abscisic acid in the accumulation of reactive oxygen species. Plant Physiol 179, 1861–1875. <https://doi.org/10.1104/pp.18.01250>.
- Zhang, M., Liu, Q., Yang, X., Xu, J., Liu, G., Yao, X., Ren, R., Xu, J., Lou, L., 2020. CRISPR/Cas9-mediated mutagenesis of *Cipsk1* in watermelon to confer resistance to *Fusarium oxysporum* f.sp. *niveum*. Plant Cell Rep 39 (6), 589–595. <https://doi.org/10.1007/s00299-020-02516-0>.
- Zhang, Y., Chen, B., Xu, Z., Shi, Z., Chen, S., Huang, X., Chen, J., Wang, X., 2014. Involvement of reactive oxygen species in endosperm cap weakening and embryo elongation growth during lettuce seed germination. J. Exp. Bot. 65 (12), 3189–3200. <https://doi.org/10.1093/jxb/eru167>.
- Zhu, M., Wang, Q., Sun, Y., Zhang, J., 2020. Effects of oxygenated brackish water on germination and growth characteristics of wheat. Agr. Water Manage. 245 (9), 106520 <https://doi.org/10.1016/j.agwat.2020.106520>.