

Hardwood recruitment into conifer plantations in Japan: Effects of thinning and distance from neighboring hardwood forests

Etsuko Utsugi, Hiroshi Kanno, Naoto Ueno, Mizuki Tomita, Tomoyuki Saitoh,
Megumi Kimura, Kenich Kanou, Kenji Seiwa *

Laboratory of Forest Ecology, Graduate School of Agricultural Science, Tohoku University, Naruko-onsen, Miyagi 989-6711, Japan

Received 29 November 2005; received in revised form 21 June 2006; accepted 11 September 2006

Abstract

Conservation of plant species diversity in managed and natural forests is a concern worldwide. To clarify the effects of thinning and distance from neighboring hardwood forests on hardwood establishment in conifer plantations, we measured the abundance and diversity of hardwoods at different life stages (seed-fall, soil seed bank, seedling, sapling) and environmental conditions (light, soil temperature, litter accumulation, and composition) from –2 to 40 m from the hardwood-coniferous forest boundary into the coniferous forest interior in five unthinned and four thinned stands in *Cryptomeria japonica* plantations in Japan. In unthinned stands, hardwood abundance and diversity decreased with increasing distance into the coniferous forest interior for all life stages, perhaps because of a distance-dependent decrease in seed-fall and the subsequent soil seed bank for wind- and bird-dispersed seeds. Higher light and soil temperatures and shallower litter accumulation were observed at the hardwood-coniferous forest boundary compared to the coniferous forest interior, suggesting favorable environmental conditions for germination, emergence, and seedling growth for hardwood species, and resulting in higher species diversity near the boundary. In thinned stands, hardwood abundance and diversity also decreased with increasing distance into the coniferous forest interior for seed-fall and the soil seed bank, but did not decrease for seedlings and saplings, probably because of distance-independent improvement of environmental conditions due to thinning. Higher light availability and soil temperature and lower litter accumulation in thinned stands enhance seedling establishment for not only shade-tolerant but also shade-intolerant and light-demanding species. Our results suggest that distance-dependent diversity of juvenile hardwoods in dense coniferous plantations is closely related to distance-dependent gradients in both seed-fall and environmental conditions, but thinning enhances diversity, even in the forest interior.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Edge effects; Light conditions; Litter accumulation; Seed dispersal; Seedling establishment; Soil seed bank; Species diversity

1. Introduction

The maintenance of biodiversity has become an increasingly important management goal, to provide a broad array of ecosystem services that directly or indirectly benefit human endeavors, such as maintaining the hydrological cycle, sustaining and/or increasing primary production, and storing and cycling nutrients (Tilman et al., 1996; Callaway et al., 2003; Hooper et al., 2005). Maintaining and/or increasing plant species diversity in conifer plantations is also a concern (Hansen et al., 1991; Roberts and Gilliam, 1995; Moore and Allen, 1999; Simberloff, 1999; Nagaike, 2002; Gotmark et al., 2005). In Japan, conifer plantations (i.e., *Cryptomeria*

japonica, *Chamaecyparis obtusa*, *Pinus densiflora*, *Larix leptolepis*) were established for timber production after intensive clear-cutting of natural forests, particularly from the 1950s through the 1970s, and now occupy one-quarter of the area of Japan (10 million ha; Forestry Agency, 2000). These monocultures generally have low value for wildlife (Yui, 1999; Osawa, 2004) and timber production, particularly at high elevations (Krauchi et al., 2000; Masaki et al., 2004). Even-aged coniferous forests may also be susceptible to pest and wind damage and effects of global warming (e.g., Lugo, 1997; Bradshaw et al., 2000), and may acidify the soil (Nordborg and Olsson, 1999). Thus, forest managers require techniques to create mixed hardwood-coniferous forests from these monocultures.

Although old plantations have functions comparable to natural forests, thinning is an effective option to create biodiversity sooner and with more consistency. Thinning,

* Corresponding author. Tel.: +81 229 847399; fax: +81 229 846490.

E-mail address: seiwa@bios.tohoku.ac.jp (K. Seiwa).

followed closely by underplanting an understory consisting of several native tree species, is a useful tool for increasing stand structural diversity and plant and wildlife diversity (e.g., Maas-Hebner et al., 2005). To preserve genetic diversity and reduce management costs, the conversion of plantations by spontaneous colonization of hardwood species (i.e., natural regeneration) is more suitable and sustainable than underplanting. However, natural regeneration may not produce the desired species composition, distribution, and density in a timely manner because of spatial and temporal variability in seed sources.

Many studies have explored the patterns and mechanisms of hardwood species recruitment in conifer plantations and have found that hardwood abundance and diversity are closely associated with thinning, stand age, previous land-use history, fertilization, and topography (e.g., Guariguata et al., 1995; Halpern and Spies, 1995; Kohli et al., 1996; Allen, 1997; Keenan et al., 1997; Thomas et al., 1999; Ito et al., 2003, 2004; Nagaike et al., 2003; Ramovs and Roberts, 2003; Jobidon et al., 2004). Forest thinning usually allows a greater number of understory species to persist (e.g., Thomas et al., 1999) because canopy disturbance usually increases available resources (e.g., light, nutrients) and fluctuations in temperature, which enhance seed germination and seedling establishment (Washitani and Takenaka, 1987; Tappeiner and Zasada, 1993; Seiwa, 1997; Sakai et al., 2005). Alternatively, thinning may result in increased dominance by one or a few understory species, thereby reducing understory diversity (Alaback and Herman, 1991; Jobidon et al., 2004). However, the mechanisms and processes underlying hardwood recruitment are not fully understood.

In long-lived tree species, the most drastic changes in population demography occur during the early life stages, particularly at the seed and seedling stages (Harper, 1977; Harcombe, 1987; Seiwa and Kikuzawa, 1996). Spatially heterogeneous demographic events at the juvenile stages strongly affect the abundance and composition of forest communities (e.g., Nakashizuka, 2000; Nagamatsu et al., 2002). Thus, to improve hardwood diversity in conifer plantations through natural regeneration, knowledge of the regeneration processes, including seed dispersal, development of the seed bank, germination, and seedling establishment, is important. However, few studies have monitored the early stages of regeneration (but see Halpern et al., 1999).

Recruitment of hardwood trees into conifer plantations largely depends on seed dispersal via dispersal agents such as wind, birds, and small mammals. Although the number of dispersed seeds usually decreases with increasing distance from the seed source, irrespective of dispersal agents (Hoppes, 1988; Nathan and Muller-Landau, 2000), dispersal agents, particularly birds and small mammals, show different dispersal behavior (i.e., dispersal distance, dispersal stand preference) in response to stand conditions, such as vegetation composition and abundance (e.g., Kikuzawa, 1988; Restrepo et al., 1999; Seiwa et al., 2002). Few researchers have investigated the patterns of seed colonization from hardwood forests to coniferous plantations (but see White et al., 2004), although

distance-dependent hardwood seed abundance and diversity is expected.

In fragmented forests, moderate to dramatic differences in light availability, temperature, soil moisture, litter accumulation, and abundance and activity of seed dispersal agents occur in the edge and interior environments. Most of these environmental conditions strongly affect seed germination and seedling recruitment (e.g., Murcia, 1995; Laurance et al., 1998; Wolf and Batzli, 2004), resulting in distance-dependent diversity and growth of understory vegetation (e.g., Williams-Linera, 1990; Chen et al., 1992; Matlack, 1994; Laurance et al., 1998; McDonald and Urban, 2004). Environmental conditions in conifer plantations change with the distance from the boundary with the neighboring hardwood forest in a manner analogous to fragmented forests, which should affect the regeneration of hardwood species.

The aim of this study was to evaluate the effects of thinning and the distance from the hardwood-coniferous forest boundary on hardwood recruitment success and species diversity in *C. japonica* plantations. We investigated environmental conditions (i.e., light, soil temperature, litter accumulation) and hardwood abundance and diversity at different stages of the regeneration process (i.e., seed-fall, soil seed bank, seedling, and sapling) at various distances from neighboring hardwood forests in both thinned and unthinned *C. japonica* stands. We addressed the following questions: To what extent is hardwood abundance and diversity affected by thinning and the distance from the hardwood-coniferous forest boundary into the interior of the *C. japonica* forest? How do environmental conditions vary with distance and thinning, and how do they affect hardwood regeneration?

2. Materials and methods

2.1. Study area

This study was conducted in *C. japonica* plantations adjacent to deciduous broad-leaved stands in the Experimental Forest of the Integrated Field Research Center of Tohoku University (38°45'N, 140°45'E, 330–540 m asl), northeastern Japan. Most of the forest area had previously been utilized as fuelwood coppices or hay meadows, but beginning in the 1950s, some of these were replanted with *C. japonica*. Other remnant coppices and meadows were abandoned in the 1960s and have regenerated naturally as secondary forests consisting of deciduous broad-leaved trees. The current landscape pattern of the study site is composed of plantation components (each 0.5–20 ha) as a mosaic in the secondary hardwood forests. In 2000, the mean annual temperature and rainfall were 10.7 °C and 1733 mm, respectively. Snow cover lasted from mid-December 2000 to mid-April 2001, with a maximum depth of 1.0 m. The soil is a well-drained Melanic Andosol (FAO-UNESCO, 1998).

We chose four stands in which thinning had been performed 2–3 years before the investigation, and five control (unthinned) stands with dense, closed canopies (Table 1). In both the thinned and control stands, hardwood trees were absent from

Table 1

Characteristics and management history of the study sites of five unthinned and four thinned stands in *Cryptomeria japonica* plantations, and characteristics of neighboring hardwood forests

	Control sites					Thinned sites			
	C1	C2	C3	C4	C5	T1	T2	T3	T4
<i>Cryptomeria japonica</i> stands									
Aspect	SW	S	E	E	S	E	N	–	SSW
Slope (%)	25	40	25	35	35	25	35	0	10
Transect aspect (boundary-interior)	SE-NW	E-W	W-E	W-E	N-S	W-E	W-E	NW-SE	N-S
Years after thinning	–	–	–	–	–	2	2	3	3
Years after planting	35	33	25	23	23	23	30	23	38
Density (number ha ⁻¹)	1725	2050	1675	1800	1475	1175	1900	1500	975
Mean tree height (m)	17.1	20.8	14.5	15.6	13.2	12.5	7.7	10.1	17.4
Ry index ^a	0.88	1.00	0.86	0.91	0.82	0.72	0.63	0.73	0.68
Neighboring hardwood forests									
Basal area of trees with wind-dispersed seeds (m ² ha ⁻¹)	0.17	4.08	17.73	1.97	5.32	11.65	11.25	17.24	0.66
Basal area of trees with bird-dispersed seeds (m ² ha ⁻¹)	0.20	7.07	0.22	9.47	5.97	1.73	6.93	3.39	6.12
Basal area of trees with small mammal dispersed seeds (m ² ha ⁻¹)	9.48	3.88	27.52	30.17	33.73	27.09	36.68	0.28	1.01
Total basal area	9.85	15.03	45.48	41.60	45.01	40.47	54.86	20.91	7.79
Number of tree species, including shrubs ^b (400 m ⁻²)	10	9	11	12	14	14	19	16	12

^a The Ry index is a measure of tree crowding relative to full density at a given stand volume; Ry of 1.00 indicate full density.

^b Trees ≥ 5 cm in diameter at breast height and shrubs > 1.5 m in height.

the canopy layer, and the slope was gentle ($< 5^\circ$). Precommercial thinning had been conducted at least once in all stands. To estimate crowding, we calculated the Ry index (Ando, 1968), which has been used as an index to assess the density of conifer plantations:

$$Ry = \frac{V_{Rc}}{V_{Rfc}},$$

where V_{Rc} and V_{Rfc} are the observed and expected total stem volume (m³ ha⁻¹) at full density, respectively. The Ry index was similar among the five control and four thinned stands (Table 1), indicating comparable crowding within the control and thinned stands. In each of the neighboring hardwood forests, a 20-m \times 20-m quadrat was established along the boundary edge, and the diameter at breast height (DBH) was measured for all trees > 5 cm in DBH and all shrubs > 1.5 m in height. The hardwood forest canopy was typically dominated by *Quercus serrata*, *Q. mongolica* var. *grosseserrata*, *Castanea crenata*, and *Carpinus laxiflora*; the subcanopy layer included *Ilex macropoda*, *Cornus controversa*, *Acer amoenum* var. *matsumurae*, and *A. mono*. *Enkianthus perulatus*, *Viburnum dilatatum*, and *V. furcatum* dominated the shrub layer. The total basal area of hardwood forests neighboring to the control stands did not differ statistically from that neighboring to thinned stands (Table 1; one-way ANOVA, $F = 0.0009$, $P = 0.98$).

2.2. Measurement of seed-fall, seedlings, and saplings

In each thinned and unthinned stand, a straight edge was delimited at the hardwood-coniferous forest boundary. Three parallel transects (45 m \times 2 m), separated by more than 5 m, were established perpendicular to the boundary, with the exception of one thinned stand (T1) in which only two transects were established. Each transect began 5 m inside the edge of the hardwood forest and extended 40 m into the interior of the

managed *C. japonica* forest. In each transect, nine 2-m \times 2-m quadrats were established at -2, 0, 2, 5, 10, 15, 20, 30, and 40 m from the edge of the hardwood-coniferous forest boundary. All woody stems in each quadrat were identified as seedlings (< 0.3 m in height) or saplings (> 0.3 m in height). The measurements were conducted from July to August in 2000 and 2001.

Three replicate seed traps were placed beside each of the nine quadrats in two control (C1, C2) and two thinned (T1, T2) stands in early August 2000 for a total of 108 traps. Each trap consisted of a 0.5-m² circular wire frame supporting a 0.2-mm mesh bag. Each frame was suspended 1.5 m above the ground on three polyvinyl chloride posts. We collected seeds and counted the number of sound seeds monthly from August 2000 to December 2001. During the winter (October 2000–April 2001), all traps were placed on the ground to reduce snow pressure.

2.3. Measurement of the soil seed bank

To estimate the buried viable seed population, we used the seedling emergence method (Halpern et al., 1999; Sakai et al., 2005) and counted the number of seedlings that emerged from sampled soil in a greenhouse. Twelve soil samples (0.5 m \times 0.5 m, 25 mm depth) were collected from quadrats 0, 5, 10, and 30 m from the hardwood-coniferous forest boundary in three control (C1–C3) and three thinned (T1–T3) stands in late May 2001. Stones and roots were removed, and each sample was spread in a plastic tray (28 cm \times 58 cm, 30-mm depth) at a 3–5-mm thickness within 24 h of collection. To prevent drying, silica fiber mattresses were placed in each tray. We identified and counted all hardwood seedlings until no new seedlings emerged (24 August 2001). The trays were occasionally watered to capacity. The minimum and maximum temperatures during the experiment were 9.5 and 45.5 $^\circ\text{C}$, respectively. The range of temperature in the greenhouse was

almost identical to that observed in a thinned stand (T1) of *C. japonica* plantations (10.0–48.1 °C).

2.4. Microenvironment measurements

To estimate light conditions in the forest understory, hemispherical canopy photographs were taken at a height of 1.0 m from the ground at the center of each quadrat at 0, 5, 10, and 30 m from the hardwood-coniferous forest boundary in all study stands. Photographs were taken using a digital camera with a fish-eye lens (Nikon FC-E8 0.21x) before (5 April) and after (July–August) canopy closure occurred in the neighboring hardwood forests. Canopy openness was calculated using Winphoto (ter Steege, 1996). Photosynthetic photon flux density (PPFD) was measured at a height of 0.3 m from the ground beside quadrats at 0, 5, 10, and 30 m from the forest boundary in one control (C2) and one thinned (T2) stand using a photo-sensor (MDS-MkV-L; Alec Electronics Co., Ltd., Kobe, Japan). Measurements were taken at 20-min intervals for 1 month (12 July–11 August 2001). Each sensor recorded the average and cumulative daily PPFD during the census period.

The soil temperature was measured in quadrats 0, 5, 10, and 30 m from the hardwood-coniferous forest boundary in one control (C2) and one thinned (T2) stand. Measurements were taken at a depth of 5 mm at 20-min intervals for 1 month (11 June–12 July 2001) using a thermo-recorder TR-52 (T&D Co., Nagano, Japan). Each sensor recorded the average and cumulative daily soil temperature during the census period.

Litter depth was measured at five points (four corners and the center) in all quadrats using a steel shovel with an engraved scale, and averaged for each quadrat. In June and July, the percent cover of *C. japonica* litter, including needles and woody material (twigs, cones, branches), was estimated at 0.1% increments.

2.5. Grouping of species by habitat and life-history traits

For comparative analyses, plant species were divided into predetermined categories describing their life-history traits (i.e., dispersal agents, seed size, clonality, successional status) according to several previous studies (Appendix A; Kikuzawa, 1983, 1984; Ishizuka and Sugawara, 1989; Seiwa and Kikuzawa, 1991, 1996; Abe et al., 1995; Seiwa et al.,

2006). Dispersal agents were classified into three categories: wind, birds, and small mammals. Seed size was classified as small (<0.5 mg), medium (<300 mg), and large (>300 mg). Our broad definition of clonality included species with rhizomes or runners and those that characteristically layer. Successional status was based on shade tolerance or environmental light availability in the common habitat of the species, and was classified into two categories: early and late successional.

2.6. Species diversity and richness

The species diversity and richness were quantified using Shannon's species diversity index (H' ; Margurran, 1988) and the number of species in each stand.

2.7. Data analysis

To evaluate whether hardwood abundance and diversity in *C. japonica* forests were affected by those of the neighboring hardwood forests, Spearman's rank correlations were calculated between the woody plant basal area of the neighboring hardwood forest (400 m⁻²) and the number of hardwoods in the study stand (100 m⁻²), and between the number of species in the neighboring hardwood forest (400 m⁻²) and the number of species of juveniles (seedlings and saplings) in the study stand (100 m⁻²). The correlations were calculated separately for control ($n = 5$) and thinned ($n = 4$) stands.

To evaluate the effects of thinning and distance from the hardwood-coniferous forest boundary on the measured environmental conditions and hardwood diversity at each life stage, we used analysis of covariance (ANCOVA) with distance as the covariate. To evaluate the effects of season (i.e., before and after canopy closure), thinning, and distance on canopy openness, we used a two-factor ANCOVA with distance as the covariate. The effects of thinning and distance from the hardwood-coniferous forest boundary on hardwood abundance were also analyzed separately for each life stage, using ANCOVA with distance as the covariate. To determine whether distance-dependent patterns of hardwood abundance differed among species grouped by life-history traits, separate ANCOVAs were conducted for control and thinned stands, with distance as the covariate.

Table 2
ANCOVA results for the effects of thinning, season, and distance from the hardwood-coniferous forest boundary into the coniferous forest interior on environmental variables in *Cryptomeria japonica* plantations

Environmental variable	Distance <i>F</i>	Thinning <i>F</i>	Season <i>F</i>	Distance × thinning <i>F</i>	Distance × season <i>F</i>	Thinning × season <i>F</i>
Canopy openness	12.34**	0.31	18.08****	1.17	4.65*	0.74
Photosynthetic photon flux density	1.02	63.95*		0.02		
Soil temperature	0.00	1.58		0.96		
Litter depth	82.26****	0.66		14.85***		
Percent cover of coniferous litter	212.60**	0.30		12.42***		

* $P < 0.05$.

** $P < 0.01$.

*** $P < 0.001$.

**** $P < 0.0001$.




Fig. 1. Changes in understory environmental conditions with distance from the hardwood-coniferous forest boundary into the coniferous forest interior in *Cryptomeria japonica* plantations. (a) Photosynthetic photon flux density (PPFD) after canopy closure; (b) canopy openness before (C, H) and after (E, J) canopy closure in the neighboring hardwood forests; (c) cumulative soil temperature; (d) depth of coniferous litter; and (e) percent cover of coniferous litter (%). Open and closed symbols indicate thinned and unthinned stands, respectively.

When necessary to meet the assumptions of the ANOVA, the data were square-root-transformed for litter cover; ln-transformed for soil temperature, litter depth, number of individuals, and number of species; and arcsine square-root-transformed for canopy openness and species diversity index (H').

To evaluate the effects of litter on seedling establishment, Spearman's rank correlations were calculated between the number of seedlings and litter depth or coniferous litter percent cover for all quadrats.

All analyses were conducted using data pooled across the three replicated quadrats at each distance for each study stand. The number of seeds from seed traps was also pooled across 2 years (2000 and 2001) because of poor seed-fall in 2001. All statistical analyses were performed using JMP (SAS Institute Inc., Cary, NC, USA).

3. Results

3.1. Microenvironment

In the *C. japonica* forests, summer PPFD was higher in thinned than control stands, but did not change with distance into the forest interior (Table 2, Fig. 1a). In both stand types, canopy openness was greatest at the edge and decreased toward the interior in winter, before canopy closure occurred

in the neighboring hardwood forests. Little difference was observed with distance in summer, after canopy closure, resulting in a significant interaction between season and distance (Table 2, Fig. 1b). Soil temperature was not affected by distance, but was higher in thinned than in control stands (Fig. 1c), although not significantly so (Table 2). Both litter depth and coniferous litter percent cover increased with increasing distance into the coniferous forest interior in both control and thinned stands (Table 2, Fig. 1d and e). The increase was much greater in control than in thinned stands, resulting in a significant interaction between thinning and distance (Table 2).

3.2. Hardwood abundance and diversity

We recorded 28,101 individuals of 95 tree species, including 17,653 trapped seeds of 35 species, 3868 buried viable seeds of 18 species, 3436 seedlings of 80 species, and 3255 saplings of 81 species (Table 3).

3.2.1. Seed-fall and soil seed bank

For trapped and buried viable seeds in both control and thinned stands, the number of individuals and number of species were greatest in quadrats at the hardwood-coniferous forest boundary and decreased with increasing distance into the

Table 3

Total number of seeds, viable seeds in the soil, seedlings and saplings per 100 m² in each of control and thinned sites

Life stage	Number of hardwoods									
	Control stands					Thinned stands				Total
	N1	N2	N3	N4	N5	T1	T2	T3	T4	
Seed	2549	52246				9224	6032			17653 (35)
Viable seeds in the soil	6460	4928	75158			13520	9091	19647		3868 (18)
Seedling	661	269	790	347	359	326	124	225	331	3436 (80)
Sapling	285	41	486	152	205	51	197	57	1786	3255 (81)

Numerals in parentheses are number of species observed.

coniferous forest interior, whereas H' did not show distance-dependent patterns (Table 4, Fig. 2a and b).

In both control and thinned stands, decreasing patterns of distance dependency were also observed for most of the dominant species (Fig. 3). The exceptions were three species with small, wind-dispersed seeds: *Alnus hirsuta* var. *sibirica*, *Betula grossa*, and *B. maximowicziana* showed little decline with distance in the number of seeds trapped in both stand types (Fig. 3a). Most bird-dispersed seeds were disseminated near the hardwood-coniferous forest boundary, whereas wind-dispersed seeds showed moderate declining patterns with distance into the coniferous forest interior (Fig. 3 a and b), although the extent of the decline was not statistically different between the two dispersal modes in either control or thinned stands (ANCOVA, control: $F < 0.52$, $P > 0.47$; thinned: $F < 0.07$, $P > 0.79$). In *Q. serrata*, the largest-seeded species, the number of acorns trapped decreased drastically with distance, and all acorns were trapped within 10 m of the forest boundary in the thinned stands (Fig. 3c). As a result, we found a significant difference in the distance-dependent declining patterns among seed sizes in thinned stands, but not control stands, because no acorns were trapped in the control stands (ANCOVA, control: $F < 1.92$, $P > 0.16$; thinned: $F < 5.44$, $P > 0.007$).

3.2.2. Seedlings and saplings

For both seedlings and saplings, the number of individuals, number of species, and H' were greatest in quadrats at the hardwood-coniferous forest boundary and decreased with increasing distance into the coniferous forest interior in control

stands, except in thinned stands (Table 4, Fig. 2c and d). The differences between thinned and control stands were few for seedlings but large for saplings, particularly in the forest interior, compared to the hardwood-coniferous forest boundary (Fig. 2c and d), resulting in a significant interaction between distance and thinning (Table 4).

For most of the species studied, distance-dependent and distance-independent patterns were also observed in control and thinned stands, respectively (Fig. 4). The exceptions were *Lindera umbellata* var. *membranacea* (distant-independent pattern) in control and *V. dilatatum* (distant-dependent pattern) in thinned stands.

In both control and thinned stands, few differences (ANCOVA, $F < 0.55$, $P > 0.58$) in distance-dependent changes among species grouped by life-history traits (dispersal mode, seed size, successional status), were observed in any of the groups for any variable (number of individuals or species, H').

The number of seedlings was negatively correlated with litter depth ($r = -0.423$, $P < 0.0001$) and coniferous litter percent cover ($r = -0.441$, $P < 0.0001$) in control stands, but not in thinned stands (litter depth: $r = -0.053$, $P = 0.624$; coniferous litter percent cover: $r = -0.026$, $P = 0.810$; Fig. 5a and b).

3.3. Effects of neighboring hardwood forests

In both control and thinned stands, neither the number of individuals per 100 m², nor the number of species of seedlings and saplings was correlated with the basal area or

Table 4

ANCOVA results for the effects of thinning and distance from the hardwood-coniferous forest boundary into the coniferous forest interior on the number of individuals, number of species, species diversity index (Shannon's H') at different life stages (seed, seed bank, seedling, and sapling) in *Cryptomeria japonica* plantations

	Number of individuals			Number of species			Shannon's H'		
	Distance F	Thinning F	Distance \times thinning F	Distance F	Thinning F	Distance \times thinning F	Distance F	Thinning F	Distance \times thinning F
Seed	3.51 [†]	0.42	0.25	3.83 [†]	2.78	0.09	0.05	66.44***	0.17
Seed bank	7.86*	0.03	0.27	10.30**	1.17	0.15	0.07	0.24	0.88
Seedling	4.27*	2.46	5.78*	6.21*	2.14	4.08*	0.97	0.17	0.38
Sapling	13.70***	2.72 [†]	10.50**	11.91***	3.88*	7.73**	12.14***	2.22	3.94*

* $P < 0.05$.** $P < 0.01$.*** $P < 0.001$.† $P < 0.1$.




Fig. 2. Changes in the number of individuals, number of species, and species diversity index (H') of hardwood species with distance from the hardwood-coniferous forest boundary into the coniferous forest interior in *Cryptomeria japonica* plantations. (a) Seeds; (b) viable seeds in the soil; (c) seedlings; and (d) saplings. The number of individuals is given as the relative value in each of the thinned and control stands because of large differences among stands. Vertical bars indicate \pm S.E. Open and closed symbols denote thinned and unthinned stands, respectively.

number of species in the neighboring hardwood forests ($-0.95 \leq r_{sp} \leq -0.40$, $0.052 \leq P \leq 0.60$).

4. Discussion

4.1. Distance-dependent species diversity in unthinned *C. japonica* forests

In unthinned *C. japonica* forests, hardwood abundance and diversity were greatest at the hardwood-coniferous forest boundary, but decreased dramatically with increasing distance into the *C. japonica* forest interior. Such distance-dependent patterns in juvenile species diversity were strongly affected by distance-dependent patterns in seed-fall and soil seed banks. Decreasing seed-fall with distance from the parent tree is common in wind-dispersed species (e.g., Augspurger and Franson, 1987; Nathan and Muller-Landau, 2000). However, we observed exceptions for three species that have small, wind-dispersed seeds (*A. hirsuta* var. *sibirica*, *B. grossa*, and *B. maximowicziana*), probably because of their large wing-load (Greene and Johnson, 1995; Houle, 1998), which would enable

them to disperse beyond the traps 40 m from the hardwood-coniferous forest boundary. This would also result in little correlation in species diversity between understory hardwoods in *C. japonica* forests and in neighboring hardwood forests. In contrast, most bird-dispersed seeds were disseminated beneath the neighboring hardwood canopy, probably because of greater avian abundance in hardwood forests than in *C. japonica* plantations (Yui, 1999). Seedlings of *Q. serrata* were observed more than 20 m from the forest boundary, although the acorns were disseminated a very short distance from the forest boundary. Acorns are usually dispersed by wood mice, such as *Apodemus speciosus* and *A. argentius*, but wood mice are rarely active in the absence of vegetative cover, thus inhibiting seed dispersal into open habitats, such as gaps and forest edges, in temperate forests (Kikuzawa, 1988; Miyaki and Kikuzawa, 1988; Wada, 1993; Seiwa et al., 2002). Our results indicate that wood mice cached acorns beneath the dense canopy of unthinned *C. japonica* forests, even though there was little vegetation in the understory. Jays (*Cyanocitta* spp.) are also a probable disperser of acorns. In a heterogeneous landscape, Gomez (2003) found that jays moved acorns nonrandomly,




Fig. 3. Changes in the number of seeds trapped and the number of viable seeds in the soil of hardwood species with distance from the hardwood-coniferous forest boundary into the coniferous forest interior in *Cryptomeria japonica* plantations. (a) Wind-dispersed seeds; (b) bird-dispersed seeds; (c) mammal-dispersed seeds. The number of individuals is given as the relative value in each of the thinned and control stands because of large differences among stands. Only the most abundant species are shown (>500 and >1000 individuals per 100 m² for trapped seeds and viable seeds in the soil, respectively). Vertical bars indicate \pm S.E.

avoiding shrubland and grasslands, and moving most acorns to pine stands.

Distance-dependent recruitment patterns of hardwood species in the coniferous plantations were also largely affected by distance-dependent changes in environmental conditions. Greater accumulation of litter usually inhibits seed germination, particularly for light-demanding species (Seiwa, 1997), because light is reduced exponentially by increasing amounts of litter (Facelli and Pickett, 1991). Litter also inhibits seedling emergence for small-seeded species that cannot penetrate thick layers of litter (Peterson and Facelli, 1992; Seiwa and Kikuzawa, 1996; Kanno and Seiwa, 2004). Therefore, seedling establishment would be strongly inhibited by the greater amounts of litter in the *C. japonica* forest interior than at the forest boundary. Furthermore, favorable light conditions at the hardwood-coniferous forest boundary would promote hardwood

recruitment into the edge of the coniferous forest. Understory light conditions were more favorable for understory hardwoods in the hardwood-coniferous forest boundary than in the coniferous forest interior, particularly in spring before canopy closure, because leaves of the dominant canopy trees (i.e., *Q. serrata*, *Q. mongolica* var. *grosseserrata*, and *C. crenata*) began to unfold approximately 2–6 weeks after those of most hardwood juveniles (Utsugi and Seiwa, personal observation). There is evidence that greater light availability in spring, prior to canopy closure in temperate deciduous forests, strongly enhances the survival and growth of juvenile hardwoods by promoting carbon acquisition and escape from pathogen and/or herbivore attacks (e.g., Seiwa, 1998; Tomita and Seiwa, 2004).

Gradients in the microenvironmental conditions from the forest edge to the interior, or “edge effects,” usually cause distance-dependent changes in vegetation (e.g., Matlack, 1994;




Fig. 4. Changes in the number of (a) seedlings and (b) saplings of hardwood species with distance from the hardwood-coniferous forest boundary into the coniferous forest interior in *Cryptomeria japonica* plantations. The number of individuals is given as the relative value in each of the thinned and control stands because of large differences among stands. Only the most abundant species are shown (>50 individuals per 100 m^2). Vertical bars indicate $\pm\text{S.E.}$

Oosterhoorn and Kappelle, 2000; Euskirchen et al., 2001). Most of the distance-dependent patterns we observed were in accordance with edge effects previously reported for hardwood forest edges adjacent to clearings. However, there were some exceptions. In hardwood forests, early successional species are more abundant near edges adjacent to clearings (Matlack, 1994; Euskirchen et al., 2001), but this did not occur at the hardwood-coniferous forest boundary, probably because favorable conditions for understory plants at forest boundaries are more ephemeral and subtle than at forest edges.

4.2. Effects of thinning

In managed forests, thinning often dramatically increases the abundance and diversity of understory vegetation (e.g., Allen, 1997; Halpern et al., 1999; Thomas et al., 1999; Ito et al.,

2003). In this study, we also observed that the number of species was greater in thinned than unthinned *C. japonica* stands for saplings, but not earlier life stages (seed-fall, seed bank, or seedlings). This suggests that the higher species richness of the saplings was caused by thinning, not by favorable conditions intrinsic to the thinned stands, which were characterized by a higher abundance of wind- and bird-dispersed species in adjacent hardwood forests, compared to the unthinned stands. If favorable conditions influenced species richness, a greater number of species would be present in the seed-fall and soil seed bank stages, as well as the sapling stage.

In thinned stands, seed germination and seedling growth of shade-intolerant species are enhanced by the removal of canopy trees, which increases light availability and soil temperature in the understory, (e.g., *Rhus javanica*, *Ulmus japonica*; Washitani and Takenaka, 1987; Seiwa, 1997). Thinning may also increase




Fig. 5. Relationships between the number of seedlings (per 4 m²) and (a) litter depth and (b) percent cover of coniferous litter.

the availability of nutrients and water for understory plants (Brix and Mitchell, 1986), enhancing the growth of seedlings, saplings, or both. Although we did not monitor the growth of the cohorts in this study, we observed rapid growth of hardwood seedlings immediately after thinning operations in other experimental stands of *C. japonica* plantations near our study stands (Eto and Seiwa, unpublished data). For example, in thinned stands, *Acer rufinerve*, *Alaria elata*, and *Moms bombycis* seedlings grew 25–48 cm in mean height 1–2 years after thinning, whereas the seedlings grew only 0–5 cm in unthinned stands.

Furthermore, in thinned stands, the negative effects of litter accumulation on seedling abundance were far less than in unthinned stands, most likely owing to less accumulation of coniferous litter, which has stronger negative effects than hardwood litter. Thinning usually disturbs the forest floor, providing favorable substrates such as bare soil and shallow coniferous litter accumulation for seedling establishment, particularly for light-demanding and small-seeded species (Peterson and Facelli, 1992; Seiwa and Kikuzawa, 1996; Seiwa, 1997; Kanno and Seiwa, 2004). Thus, reducing the litter accumulation appears to be an important contribution of thinning to species diversity.

Our results strongly suggest that thinning promotes seedling establishment for a variety of hardwood species by providing favorable environmental conditions for seed germination, seedling emergence, and seedling growth, resulting in species richness in *C. japonica* plantations. We observed improved diversity especially in the forest interior of thinned stands, where thinning improved environmental conditions (i.e., light, litter accumulation) more than at the hardwood-coniferous forest boundary. This increased hardwood recruitment in the forest interior and resulted in distance-independent patterns of species richness.

4.3. Management implications

In Japan, most commercial plantations of even-aged conifers are forest fragments that range in area from 0.1 to 100 ha and are adjacent to natural hardwood forests on at least one side. In recent decades, forest managers have been reluctant to thin

plantations because of decreasing timber prices and increasing harvesting costs. Thus, many coniferous plantations are overcrowded and have low species diversity, particularly in large fragments, because of the abrupt decline in species diversity with distance into the forest interior. In contrast, high diversity would be expected in small fragments because most of the area would experience edge effects. However, we evaluated juvenile, rather than adult, species richness, whereas most of the ecosystem function effects of biodiversity are performed by adult plants (Hooper et al., 2005).

Thinning appears to be a reliable method for restoring adult-level hardwood diversity in *C. japonica* plantations because it promptly and dramatically diminishes edge effects and strongly enhances the growth of understory juveniles, regardless of their successional status. However, in this study, we examined the distribution of seedlings and saplings but did not undertake long-term monitoring of our cohorts. To evaluate the duration of thinning effects, further long-term studies are needed.

To improve the species diversity of adult trees in monoculture forests, periodic thinning should be performed because intensive thinning often results in one or a few dominant species (Jobidon et al., 2004). However, periodic thinning is unrealistic economically, due to low timber prices and high operation costs. To meet this challenge, changes in ecosystem functions as newly recruited hardwoods increase in size should be evaluated. Improving the economic value of hardwoods is also important to determine thinning intensity. Further research is warranted to better understand the full range of management activities (e.g., thinning intensity, rotation age, stand size, configuration, timber price, and operation costs) for conversion of artificial coniferous forests to mixed hardwood-coniferous forests.

Acknowledgements

We wish to thank Motonori Sato, Munetaka Tozawa, Kyoko Obayashi, Yoshiko Miwa, Hiroshi Kanno, and Tomoyuki Saitoh for their help with the field observations. Kiyoshi Umeki provided constructive comments on this manuscript. This work was supported by a grant from the Ministry of Education, Science, and Culture (No. 12460064, Seiwa).

	Unthinned stands			Thinned stands			Functional group			
	S [†]	VS [‡]	SL&SA [§]	S [†]	SB [‡]	SL&SA [§]	SDA ^a	SS ^b	CL ^c	SC ^d
	Mean number ± S.D. (n = 2)	Mean number ± S.D. (n = 3)	Mean number ± S.D. (n = 5)	Mean number ± S.D. (n = 2)	Mean number ± S.D. (n = 3)	Mean number ± S.D. (n = 4)				
<i>Acanthopanax sciadophylloides</i>	26.0 ± 5.7		43.4 ± 46.9			4.7 ± 5.5	B	M		L
<i>Acanthopanax spinosus</i>			1.0				B	S		L
<i>Acer amoenum</i> var. <i>matsumurae</i>	1848.5 ± 2457.2	1931.0	9.0 ± 9.8	148.5 ± 115.3		10.5 ± 9.9	W	M		L
<i>Acer cissifolium</i>	16898.5 ± 23876.9	266.0	6.0	44.0			W	M		L
<i>Acer distylum</i>			7.5 ± 7.8			5.0	W	M	+	L
<i>Acer japonicum</i>	96.0		9.0			2.0	W	M		L
<i>Acer micranthum</i>	27.0		6.1 ± 8.3				W	M		L
<i>Acer mono</i>	2382.0 ± 3348.9	216.5 ± 259.5	5.5 ± 4.2	452.5 ± 586.2	33.0	2.8 ± 1.7	W	M		L
<i>Acer rufinerve</i>	296.0 ± 398.8	33.0	19.0 ± 26.4	152.0 ± 120.2	66.5 ± 47.4	2.3 ± 1.3	W	M	+	L
<i>Acer sieboldianum</i>			3.0 ± 2.8			0.5 ± 0.7	W	M		L
<i>Actinidia arguta</i>	166.5 ± 204.4		6.0	252.0		4.0	B	S	+	L
<i>Actinidia polygama</i>			4.0 ± 1.7			0.8 ± 0.3	B	S	+	L
<i>Akebia trifoliata</i>			44.5 ± 48.8			14.8 ± 18	B	M	+	L
<i>Alnus hirsuta</i> var. <i>sibirica</i>	578.0 ± 168.3			70.5 ± 16.3			W	S		E
<i>Aralia elata</i>	278.0 ± 350.7	399.7 ± 547.9	1.3 ± 0.6	92.5 ± 120.9	1399.0	27.8 ± 47.6	B	S		E
<i>Aralia elata</i>						42.0	B	M	+	L
<i>Aucuba japonica</i> var. <i>borealis</i>			32.0			13.5 ± 17.7	B	M	+	L
<i>Betula grossa</i>	1476.0 ± 1821.5	410.3 ± 170.9	1.5 ± 0.7	2089.5 ± 2368.1	1543.0 ± 2326.1	2.0	W	S		E
<i>Betula maximowicziana</i>	340.5 ± 461.7	24309.0 ± 41614.3			189.0 ± 101.8		W	S		E
<i>Broussonetia kazinoki</i>			0.2				B	S		E
<i>Callicarpa japonica</i>			1.0			6.2 ± 8.3	B	S		E
<i>Carpinus cordata</i>	7.0			44.0			W	M		L
<i>Carpinus laxiflora</i>	719.5 ± 743.2	277.3 ± 214.2	5.3 ± 4.2	1159.0 ± 886.7	5883.0 ± 4945.2	1.8 ± 1	W	S		E
<i>Castanea crenata</i>	7.0		8.3 ± 11.9			5.0 ± 5.7	M	L		E
<i>Celastrus orbiculatus</i>	63.0 ± 35.4		4.8 ± 3.9			7.0 ± 5.4	B	M		E
<i>Cledendrum trichotomum</i>			9.5 ± 11.2			1.0	W	S		E
<i>Clerodendron trichotomum</i>				7.0		7.5 ± 3.5	B	M		E
<i>Cornus controversa</i>	181.5 ± 120.9		4.0 ± 4.8	148.0 ± 147.1		2.7 ± 2.1	B	M		E
<i>Corylus heterophylla</i> var. <i>thunbergii</i>			1.0				M	L		E
<i>Corylus sieboldiana</i>			0.5			1.0	M	L		E
<i>Enkianthus perulatus</i>			3.0				W	S		L
<i>Euonymus macropterus</i>			0.2			0.4	B	M		L
<i>Euonymus sieboldianus</i>						3.0	B	M		L
<i>Fagus crenata</i>				11.0 ± 5.7			M	L		L
<i>Fraxinus lanuginosa</i> f. <i>serrata</i>			10.3 ± 13.5			30.0 ± 36.8	W	M		L
<i>Fraxinus sieboldiana</i>			4.0			3.5 ± 3.5	W	M		L
<i>Hamamelis japonica</i>						10.0	A	M		L
<i>Hydrangea paniculata</i>			7.5 ± 7.3			6.9 ± 7.3	W	S		E
<i>Hydrangea petiolaris</i>			2.2 ± 2.5			1.0	W	S	+	L
<i>Hydrangea serrata</i> var. <i>megacarpa</i>			1.5 ± 0.7			1.0	W	S	+	E
<i>Ilex crenata</i> var. <i>paludosa</i>			116.0			2.0	B	M	+	L
<i>Ilex macropoda</i>	611.5 ± 738.9		6.3 ± 4.8	59.0 ± 73.5		15.0	B	M		L
<i>Juglans ailanthifolia</i>						1.0	M	L		E
<i>Kalopanax pictus</i>	296.0		0.5 ± 0.7			1.3 ± 0.6	B	S		L
<i>Lespedeza bicolor</i>		100.0 ± 94.8			144.3 ± 138.9	0.2	A	M		E
<i>Leucothoe grayana</i>			43.0 ± 52.3			0.5	W	S	+	E
<i>Lindera praxox</i>						3.0	M	M		L
<i>Lindera umbellata</i> var. <i>membranacea</i>			9.8 ± 15.6			3.2 ± 4.2	B	M		L

Appendix A (Continued)

	Unthinned stands			Thinned stands			Functional group			
	S [†] Mean number ± S.D. (n = 2)	VS [‡] Mean number ± S.D. (n = 3)	SL&SA [§] Mean number ± S.D. (n = 5)	S [†] Mean number ± S.D. (n = 2)	SB [‡] Mean number ± S.D. (n = 3)	SL&SA [§] Mean number ± S.D. (n = 4)	SDA ^a	SS ^b	CL ^c	SC ^d
<i>Magnolia hypoleuca</i>			0.8 ± 1	7.0		1.5 ± 0.7	B	M		L
<i>Magnolia praecocissima</i>			1.0				B	M		L
<i>Melisoma myriantha</i>						1.0	B	M		L
<i>Menzeisia purpurea</i>			14.0				W	S		L
<i>Morus bombycis</i>		233.0	1.0	52.0	344.0 ± 538.7	22.0	B	S		E
<i>Ostrya japonica</i>				52.0 ± 52.3			W	M		E
<i>Prunus grayana</i>			6.0 ± 1.4			5.0 ± 5.7	B	M		L
<i>Prunus sargentii</i>			1.5 ± 0.7	38.0		3.0	B	M		L
<i>Prunus verecunda</i>	104.0 ± 73.5		13.0 ± 19.1	33.0 ± 15.6		4.0	B	M		E
<i>Pterocarya rhoifolia</i>				15.0			W	M		E
<i>Pueraria lobata</i>		33.0					A	M	+	E
<i>Quercus mongolica</i> var. <i>grosserrata</i>			1.0			2.5 ± 1.9	M	L		L
<i>Quercus serrata</i>	14.5 ± 10.6	33.0	7.4 ± 10.5	2734.5 ± 3793.6	33.0	7.5 ± 5.8	M	L		E
<i>Rhododendron</i> <i>kaempferi</i>			10.9 ± 12.7			1.0	W	S		E
<i>Rhus ambigua</i>			0.8 ± 0.4			1.4 ± 1.3	B	M	+	L
<i>Rhus javanica</i>						0.3	B	M		E
<i>Rhus trichocarpa</i>	22.0		2.8 ± 2.9	44.5 ± 53		6.0	B	M		E
<i>Rubus parvifolius</i> L.						1.0	B	S	+	E
<i>Rubus crataegiofolius</i>						0.0	B	S	+	E
<i>Rubus palmatus</i> var. <i>coptophyllus</i>			63.0 ± 41.4			110.0 ± 210.7	B	S	+	E
<i>Salix vulpine</i>			6.0			1.7 ± 1.8	W	S		E
<i>Sambucus sieboldiana</i>			7.0		33.0	10.7 ± 13.3	B	S		E
<i>Schisandra nigra</i>			0.5				B	S	+	L
<i>Schizophragma</i> <i>hydrangeoides</i>			15.8 ± 10.2			23.5 ± 24.7	W	S	+	L
<i>Smilax china</i>			3.4 ± 4.1			3.1 ± 4.3	B	M	+	E
<i>Sorbus alnifolia</i>			0.5 ± 0.5			3.0	B	M		L
<i>Sorbus commixta</i>			2.0			2.0	B	S		L
<i>Sorbus japonica</i>			5.0			0.4	B	M		L
<i>Stachyurus praecox</i>						0.7 ± 0.5	B	S		E
<i>Staphylea bumalda</i>			0.4				A	M		L
<i>Styrax japonica</i>				30.0	133.0	2.7 ± 3.3	M	M		E
<i>Styrax obassia</i>			0.6 ± 0.4			4.5 ± 4.9	M	L		L
<i>Symplocos chinensis</i>			4.0			0.4	B	M		L
<i>Tilia japonica</i>			2.0	30.0			W	M		L
<i>Tripterygium regelii</i>			1.0				B	S	+	L
<i>Ulmus japonica</i>				7.0		0.7 ± 0.4	W	S		E
<i>Ulmus laciniata</i>						2.5 ± 0.7	W	S		L
<i>Vaccinium japonicum</i>			2.0				B	S		L
<i>Vaccinium oldhamii</i>			3.5 ± 3.5				B	S		E
<i>Viburnum plicatum</i> var. <i>toentosum</i>				89.0		21.0	B	M	+	L
<i>Viburnum wrightii</i>			8.7 ± 7.5				B	M	+	L
<i>Viburnum dilatatum</i>			20.0 ± 20.3			19.8 ± 17.9	B	M	+	L
<i>Viburnum furcatum</i>			247.0			3.0	B	M	+	L
<i>Vitis coignetiae</i>	25.5 ± 26.2	199.5 ± 94	5.2 ± 7.7	7.0	66.3 ± 57.7	7.0 ± 8.5	B	M		E
<i>Weigela hortensis</i>		3163.5 ± 2260.6	4.0 ± 4.2		3141.3 ± 2077.7	2.4 ± 3.1	W	S		E
<i>Wisteria brachybotrys</i>			1.0			3.1 ± 2.5	A	L	+	E
<i>Zelkova serrata</i>	1156.0 ± 1540.1		3.3 ± 3.9	70.5 ± 4.9	233.0	3.0 ± 2.8	W	M		E

[†] S: seeds.[‡] VS: viable seeds in the soil.[§] SL & SA: seedlings and saplings.^a Seed dispersal agents (SDA): A: dispersed automatically; B: bird-dispersed; M: mammal-dispersed; W: wind-dispersed.^b Seed size (SS): S: small <0.5 mg; M: medium <300 mg; L: large ≥300 mg.^c Clonality (CL): + indicate clonal plants which grow clonally by stolon and rhizome.^d Successional status (SC): E: early successional; L: late successional.

References

- Abe, S., Masaki, T., Nakashizuka, T., 1995. Factors influencing sapling composition in canopy gaps of a temperate deciduous forest. *Vegetatio* 120, 21–32.
- Alaback, P.B., Herman, F.R., 1991. Long-term response of understory vegetation to stand density in *Picea-Tsuga* forest. *Can. J. For. Res.* 21, 534–539.
- Allen, J., 1997. Reforestation of bottomland hardwoods and the issue of woody species diversity. *Restor. Ecol.* 5, 125–134.
- Ando, T., 1968. Ecological studies on stand density control in even-aged pure stands. *Bull. For. For. Prod. Res. Inst.* 210, 75–85 (in Japanese).
- Augsburger, C.K., Franson, S.E., 1987. Wind dispersal of artificial fruits varying in mass, area, and morphology. *Ecology* 68, 27–42.
- Bradshaw, R.H.W., Holmquist, B.H., Cowling, S.A., Sykes, M.T., 2000. The effects of climate change on the distribution and management of *Picea abies* in southern Sweden. *Can. J. For. Res.* 30, 1992–1998.
- Brix, H., Mitchell, A.K., 1986. Thinning and nitrogen fertilization effects on soil and tree water stress in a Douglas-fir stand. *Can. J. For. Res.* 16, 1134–1138.
- Callaway, J.C., Sullivan, G., Zedler, J.B., 2003. Species-rich planting increases biomass and nitrogen accumulation in a wetland restoration experiment. *Ecol. Appl.* 13, 1626–1639.
- Chen, J., Franklin, J.F., Spies, T.A., 1992. Vegetation responses to edge environments in old-growth Douglas-fir forests. *Ecol. Appl.* 2, 378–396.
- Euskirchen, E., Chen, J., Bi, R., 2001. Effects of edges on plant communities in a managed landscape in northern Wisconsin. *For. Ecol. Manage.* 148, 93–108.
- Facelli, J.M., Pickett, S.T.A., 1991. Plant litter: its dynamics and effects on plant community structure. *Bot. Rev.* 57, 1–32.
- FAO-UNESCO, 1998. World Reference Base for Soil Resources. World Soil Resources Report. 84, UNESCO, Rome, 88 pp.
- Forestry Agency, 2000. Annual Report on Trends in Forestry. Ministry of Agriculture, Forestry, and Fisheries, Japan, 7 pp.
- Gomez, J.M., 2003. Spatial patterns in long-distance dispersal of *Quercus ilex* acorns by jays in a heterogeneous landscape. *Ecography* 26, 573–584.
- Gotmark, F., Fridman, J., Kenpe, G., Norden, B., 2005. Broad-leaved tree species in conifer-dominated forestry: regeneration and limitation of saplings in southern Sweden. *For. Ecol. Manage.* 214, 142–157.
- Greene, D.F., Johnson, E.A., 1995. Long-distance wind dispersal of tree seeds. *Can. J. Bot.* 73, 1036–1045.
- Guariguata, M., Rheingans, R., Montagnini, F., 1995. Early woody invasion under tree plantations in Costa Rica: implications for forest restoration. *Restor. Ecol.* 3, 252–260.
- Halpern, C.B., Spies, T.A., 1995. Plant species diversity in natural and managed forests of the Pacific Northwest. *Ecol. Appl.* 5, 913–934.
- Halpern, C.B., Evans, S., Nielson, S., 1999. Soil seed banks in young, closed-canopy forests of the Olympic Peninsula, Washington: potential contributions to understory reinitiation. *Can. J. Bot.* 77, 922–935.
- Hansen, A.J., Spies, T.A., Swanson, F.J., Ohmann, J.L., 1991. Conserving biodiversity in managed forests. Lessons from natural forests. *Bioscience* 41, 382–392.
- Harcombe, P.A., 1987. Tree life tables: simple birth, growth, and death data encapsulate life histories and ecological roles. *Bioscience* 37, 558–568.
- Harper, J.L., 1977. Population Biology of Plants. Academic Press, London, UK.
- Hooper, D.U., Chapin III, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.H., Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A.J., Vandermeer, J., Wardle, D.A., 2005. Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.* 75, 3–35.
- Hoppes, W.G., 1988. Seedfall pattern of several species of bird-dispersed plants in an Illinois woodland. *Ecology* 69, 320–329.
- Houle, G., 1998. Seed dispersal and seedling recruitment of *Betula alleghaniensis*: spatial inconsistency in time. *Ecology* 79, 807–818.
- Ishizuka, M., Sugawara, S., 1989. Composition and structure of natural mixed forests in central Hokkaido (I) Composition differences and species characteristics by elevation and from disturbances. *J. Jpn. For. Soc.* 68, 79–86.
- Ito, S., Nakagawa, M., Buckley, G.P., Nogami, K., 2003. Species richness in sugi (*Cryptomeria japonica* D. Don) plantations in southeastern Kyushu, Japan: the effects of stand type and age on understory trees and shrubs. *J. For. Res.* 8, 49–57.
- Ito, S., Nakayama, R., Buckley, G.P., 2004. Effects of previous land-use on plant species diversity in semi-natural and plantation forests in a warm-temperate region in southeastern Kyushu, Japan. *For. Ecol. Manage.* 196, 213–225.
- Jobidon, R., Guillaume, C., Thiffault, N., 2004. Plant species diversity and composition along an experimental gradient of northern hardwood abundance in *Picea mariana* plantations. *For. Ecol. Manage.* 198, 209–221.
- Kanno, H., Seiwa, K., 2004. Sexual vs. vegetative reproduction in relation to forest dynamics in the understory shrub, *Hydrangea paniculata* (Saxifragaceae). *Plant Ecol.* 170, 43–53.
- Keenan, R., Lamb, D., Woldring, O., Irvine, T., Jensen, R., 1997. Restoration of plant biodiversity beneath tropical tree plantations in northern Australia. *For. Ecol. Manage.* 99, 117–131.
- Kikuzawa, K., 1983. Leaf survival of woody plants in deciduous broad-leaved forests. 1. Tall trees. *Can. J. Bot.* 61, 2133–2139.
- Kikuzawa, K., 1984. Leaf survival of woody plants in deciduous broad-leaved forests. 2. Small trees and Shrubs. *Can. J. Bot.* 6.
- Kikuzawa, K., 1988. Dispersal of *Quercus mongolica* acorns in a broadleaved deciduous forest. 1. Disappearance. *For. Ecol. Manage.* 25, 1–8.
- Kohli, R., Singh, H., Rani, D., 1996. Status of floor vegetation under some monoculture and mixculture plantations in North India. *J. For. Res.* 1, 205–209.
- Krauchi, N., Brang, P., Schonenberger, W., 2000. Forest mountainous regions: gaps in knowledge and research needs. *For. Ecol. Manage.* 132, 73–82.
- Laurance, W.F., Ferreira, L.V., Rankin-de Merona, J.M., Laurance, S.G., 1998. Rainforest fragmentation and the dynamics of Amazonian tree communities. *Ecology* 79, 2032–2040.
- Lugo, A., 1997. The apparent paradox of reestablishing species richness on degraded lands with tree monocultures. *For. Ecol. Manage.* 99, 9–19.
- Maas-Hebner, K.G., Emmingham, W.E., Larson, D.J., Chan, S.S., 2005. Establishment and growth of native hardwood and conifer seedlings underplanted in thinned Douglas-fir stands. *For. Ecol. Manage.* 208, 331–345.
- Margurran, A.E., 1988. Ecological Diversity and its Measurements. Princeton University Press, Princeton, NJ.
- Masaki, T., Ota, T., Sugita, H., Oohara, H., Otani, T., Nagaike, T., Nakamura, S., 2004. Structure and dynamics of tree populations within unsuccessful conifer plantations near the Shirakami Mountains, a snowy region of Japan. *For. Ecol. Manage.* 194, 389–401.
- Matlack, G.R., 1994. Vegetation dynamics of the forest edge-trends in space and successional time. *J. Ecol.* 82, 113–123.
- McDonald, R.I., Urban, D.L., 2004. Forest edges and tree growth rates in the North Carolina Piedmont. *Ecology* 85, 2258–2266.
- Miyaki, M., Kikuzawa, K., 1988. Dispersal of *Quercus mongolica* acorns in a broadleaved deciduous forest. 2. Scatter-hoarding by mice. *For. Ecol. Manage.* 25, 9–16.
- Moore, S.E., Allen, E.L., 1999. Plantation forestry. In: Hunter, M.L.J. (Ed.), Maintaining Biodiversity in Forest Ecosystems. Cambridge University Press, Cambridge, UK, pp. 400–433.
- Murcia, C., 1995. Edge effects in fragmented forests: implications for conservation. *TREE* 10, 58–62.
- Nagaike, T., 2002. Differences in plant species diversity between conifer (*Larix kaempferi*) plantations and broad-leaved (*Quercus crispula*) secondary forests in central Japan. *For. Ecol. Manage.* 168, 111–123.
- Nagaike, T., Hayashi, A., Abe, M., Arai, N., 2003. Differences in plant species diversity in *Larix kaempferi* plantations of different ages in central Japan. *For. Ecol. Manage.* 183, 177–193.
- Nagamatsu, D., Seiwa, K., Sakai, A., 2002. Seedling establishment of deciduous trees in various topographic positions. *J. Veg. Sci.* 13, 35–44.
- Nakashizuka, T., 2000. Species coexistence in temperate, mixed deciduous forest. *TREE* 16, 205–210.
- Nathan, R., Muller-Landau, H., 2000. Spatial patterns of seed dispersal, their determinants and consequences for recruitment. *TREE* 15, 278–285.
- Nordborg, F., Olsson, S., 1999. Changes in soil mineralogy and exchangeable cation in stands of Norway spruce planted on former pasture land. *Plant Soil* 207, 219–229.
- Oosterhoorn, M., Kappelle, M., 2000. Vegetation structure and composition along an interior-edge-exterior gradient in a Costa Rican montane cloud forest. *For. Ecol. Manage.* 126, 291–307.

- Osawa, M., 2004. Species richness of Cerambycidae in larch plantations and natural broad-leaved forests of the central mountainous region of Japan. *For. Ecol. Manage.* 189, 375–385.
- Peterson, C., Facelli, J., 1992. Contrasting germination and seedling growth of *Betula alleghaniensis* and *Rhus typhina* subjected to various amounts and types of plant litter. *Am. J. Bot.* 79, 1209–1216.
- Ramovs, B.V., Roberts, M.R., 2003. Understory vegetation and environment responses to tillage, forest harvesting, and conifer plantation development. *Ecol. Appl.* 13, 1682–1700.
- Restrepo, C., Gomez, N., Heredia, S., 1999. Anthropogenic edges, treefall gaps, and fruit-frugivore interactions in a neotropical montane forest. *Ecology* 80, 668–685.
- Roberts, M.R., Gilliam, F.S., 1995. Patterns and mechanisms of plant diversity in forested ecosystems: implications for forest management. *Ecol. Appl.* 5, 969–997.
- Sakai, A., Sato, S., Sakai, T., Kuramoto, S., Tabuchi, R., 2005. A soil seed bank in a mature conifer plantation and establishment of seedlings after clear-cutting in southern Japan. *J. For. Res.* 10, 295–304.
- Seiwa, K., 1997. Variable regeneration behaviour of *Ulmus davidiana* var. *japonica* in response to disturbance regime for risk spreading. *Seed Sci. Res.* 7, 195–207.
- Seiwa, K., 1998. Advantages of early germination for growth and survival of seedlings of *Acer mono* under different overstorey phenologies in deciduous broad-leaved forests. *J. Ecol.* 86, 219–228.
- Seiwa, K., Kikuzawa, K., 1991. Phenology of tree seedlings in relation to seed size. *Can. J. Bot.* 69, 532–538.
- Seiwa, K., Kikuzawa, K., 1996. Importance of seed size for the establishment of seedlings of five deciduous broad-leaved tree species. *Vegetatio* 123, 51–64.
- Seiwa, K., Watanabe, A., Irie, K., Kanno, H., Saitoh, T., Akasaka, S., 2002. Impact of site-induced mouse caching and transport behaviors on regeneration in Japanese chestnuts, *Castanea crenata*. *J. Veg. Sci.* 13, 517–526.
- Seiwa, K., Kikuzawa, K., Kadowaki, T., Akasaka, S., Ueno, N., 2006. Shoot lifespan in relation to successional status in deciduous broad-leaved tree species in a temperate forest. *New Phytol.* 169, 537–548.
- Simberloff, D., 1999. The role of science in the preservation of forest diversity. *For. Ecol. Manage.* 115, 101–111.
- Tappeiner, J.C., Zasada, J.C., 1993. Establishment of salmonberry, salal, vine maple, and bigleaf maple seedlings in the coastal forests of Oregon. *Can. J. For. Res.* 23, 1775–1780.
- ter Steege, H., 1996. Winphoto 5.0. A programme to analyze vegetation indices, light and light quality from hemispherical photographs. Tropenbos, Wageningen, The Netherlands.
- Thomas, S., Halpern, C., Falk, D., Liguori, D., Austin, K., 1999. Plant diversity in managed forests: understory responses to thinning and fertilization. *Ecol. Appl.* 9, 864–879.
- Tilman, D., Wedin, D., Knops, J., 1996. Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature* 379, 718–720.
- Tomita, M., Seiwa, K., 2004. Influence of canopy tree phenology on understory populations of *Fagus crenata*. *J. Veg. Sci.* 15, 379–388.
- Wada, N., 1993. Dwarf bamboos affect the regeneration of zoochorous trees by providing habitats to acorn-feeding rodents. *Oecologia* 94, 403–407.
- Washitani, I., Takenaka, A., 1987. Gap-detecting mechanism in the seed germination of *Mallotus japonicus* (Thunb.) Muell. Arg., a common pioneer tree of secondary succession in temperate Japan. *Ecol. Res.* 2, 191–201.
- White, E., Tucker, N., Meyers, N., Wilson, J., 2004. Seed dispersal to revegetated isolated rainforest patches in North Queensland. *For. Ecol. Manage.* 192, 409–426.
- Williams-Linera, G., 1990. Vegetation structure and environmental conditions of forest edges in Panama. *J. Ecol.* 78, 356–373.
- Wolf, M., Batzli, B., 2004. Forest edge—high or low quality habitat for white-footed mice (*Peromyscus leucopus*)? *Ecology* 85, 756–769.
- Yui, M., 1999. Bird species conservation. In: Fujimori, T., Yui, M., Ishii, N. (Eds.), *Wildlife Conservation in Forests – Conservation of Biodiversity – Nihon Ringyo Chosakai*, pp. 125–127 (in Japanese).