

WEEDS, INSECTS, AND DISEASES

D.T. PATTERSON

NCDA&CS, Raleigh, North Carolina 27611, USA

J.K. WESTBROOK

USDA/ARS, College Station, Texas 77840, USA

R.J.V. JOYCE⁺

P.D. LINGREN

USDA/ARS, Lane, Oklahoma 74555, USA

J. ROGASIK

Agricultural Institute for Climatic Research, Munchenberg, Germany

Abstract. The geographic distribution, vigor, virulence, and agricultural impact of weeds, insects, and plant pathogens will be affected by climatic changes accompanying the global "greenhouse effect." Weed/crop competitive interactions, particularly among species differing in photosynthetic pathway (C_3 v C_4), may be altered, with the C_3 species favored by increasing CO_2 . Physiological and biochemical changes induced in host crop plants by rising CO_2 may affect feeding patterns of pest insects. Compilation of climatic thresholds for phenological development of pest insects reveals the potential for shifts in pest behavior induced by global warming and other climatic change. Generation times may be reduced, enabling more rapid population increases to occur. Poleward migration may be accelerated during the crop season. The epidemiology of plant diseases also will be altered. Prediction of disease outbreaks will be more difficult in periods of rapidly changing climate and unstable weather. Environmental instability and increased incidence of extreme weather may reduce the effectiveness of pesticides on targeted pests or result in more injury to non-target organisms. Biological control may be affected either negatively or positively. Overall, the challenge to agriculture from pests probably will increase.

1. Introduction

Weeds, insects, and pathogen-mediated plant diseases will be affected by global climate change. Alterations in the geographic distribution of these crop pests and their vigor in current ranges likely will occur. Existing research has investigated climatic determinants of the range of many pests, and a large body of literature addresses the effects of warmer drier climates on crops and some of their pests. However, the potential for climatically driven changes in crop

⁺deceased



Climatic Change 43: 711–727, 1999.

©1999 Kluwer Academic Publishers. Printed in the Netherlands.

losses due to pests have not been included in most agricultural impact studies (Prestidge and Pottinger, 1990; Stinner et al., 1987; Waggoner, 1983). This review provides a background for evaluating possible impacts of climate change on crop pests and the resulting effects on crop production.

2. Weeds

Worldwide, crop production losses from weed interference are estimated to be about 12%. Losses are highest in primitive agricultural production systems, averaging 25% of crop production (Parker and Fryer, 1975). Extensive efforts are made to control weeds. Most cultivation and tillage practices are for weed control, and over half of the pesticides applied annually to cropland are herbicides for controlling weeds (Furtick, 1978).

Weeds share the same trophic level (primary producer) with crops and compete with crops for the resources essential for plant growth. Unless controlled, weeds are always present at yield-reducing levels in the agroecosystem, because of the reservoir of weed seeds in agricultural soils. Changes in CO₂ concentration, temperature, and water or nutrient availability affect plant growth. Factors that differentially affect crops and weeds are most likely to affect weed/crop competition and other weed/crop interactions.

2.1 THE PHYSIOLOGICAL EFFECTS OF RISING CO₂ ON WEEDS

Weeds will benefit from the "CO₂ fertilization effect" and from improvements in water use efficiency associated with the partial closure of stomates by increasing CO₂ concentrations. Evapotranspiration on a land area basis may not be altered as much as expected from the effects of rising CO₂ on stomatal aperture, because of interactions with other limiting factors (Patterson, 1995). The different patterns of response of C₃ and C₄ plants to CO₂ are of particular significance in weed/crop competition, because most of the world's crops are C₃ plants, while many major weeds are C₄ plants (Patterson and Flint, 1980). For example, 80 of the 86 "plant species commodities" that contribute 90% of per capita food supplies worldwide are C₃ plants (Prescott-Allen and Prescott-Allen, 1990), while 14 of the 18 "world's worst weeds" are C₄ plants (Holm et al., 1977). In the agroecosystem, the growth and competitive interactions of C₃ and C₄ crops and weeds are synchronized by the timing of planting of the crop and by other cultural manipulations and are not temporally separated as they are in most natural ecosystems. Additionally, the agricultural weed flora has been homogenized through dispersal and distribution by man, so that species combinations which never occur in natural ecosystems occur

together and compete in agroecosystems.

Under controlled conditions, CO₂ enrichment stimulates the growth of C₃ weeds and crops more than that of C₄ species, although a considerable range of response has been reported (Table 1). Temperatures during growth also affect responses to elevated CO₂, and responses vary with species (Patterson, 1995). In competition experiments, C₃ species generally benefit from CO₂ enrichment at the expense of C₄ species (Carter and Peterson, 1983; Wray and Strain, 1987; Bazzaz and Garbutt, 1988; Patterson and Flint, 1990). For example, the competitiveness between C₃ soybean and C₄ johnson grass (*Sorghum halepense* (L.) Pers.) was increased when CO₂ was increased from 350 to 675 ppm (Patterson et al.; 1984). Unfortunately, no studies of CO₂ effects on weed/crop competition under field conditions have been reported to date.

Table 1

Biomass responses of C₃ and C₄ plants to CO₂ doubling.

Category	Range of Response
C ₃ Crops	1.10 to 2.43 X
C ₄ Crops	0.98 to 1.24 X
C ₃ Weeds	0.95 to 2.72 X
C ₄ Weeds	0.56 to 1.61 X

Summarized from a literature survey by Patterson (1993)

2.2 INDIRECT (CLIMATIC) EFFECTS ON WEED GROWTH AND WEED/CROP COMPETITION

In addition to the direct, physiological effects of rising CO₂, climatic change could have significant effects on weed growth, weed/crop competition, and the geographical distribution of weeds. Temperature and precipitation are the primary variables controlling the distribution of vegetation (Woodward and Williams, 1987). High latitude limits of tropical and warm temperate annual species are primarily set by the accumulated heat sum during the growing season. Low latitude limits of species from cool or cold regions are set by competition, cold tolerance apparently being achieved at the expense of competitive ability (Woodward, 1988). Temperature and precipitation patterns influence the distribution of weeds concurrently with the selection of crops. Choice of growing season, manipulative cultural practices, fertilization, and irrigation ameliorate unfavorable conditions for crops, but weeds often benefit

as well (Harlan, 1975).

Most weeds of warm season crops originated in tropical or warm temperate areas. Growth of such species often is very responsive to small increases in temperature. For example, the growth of three leguminous weeds increased significantly as day/night temperature increased from 23/17 to 26/20 or 29/23°C (Flint et al., 1984). Biomass of C₄ smooth pigweed (*Amaranthus hybridus* L.) grown at 29/20°C was 240% of that grown at 26/17°C (Flint and Patterson, 1983). C₄ grass weeds also respond to small temperature increases in the range of 26/20 to 29/23°C (Patterson, 1993). The stimulation of weed growth by 3 to 5°C increments in this range has significant implications for weed/crop competition, because temperature increases of this magnitude may occur within the next 100 yr.

The range expansion of many weeds into higher latitudes may accelerate with global warming (Patterson, 1993; Rahman and Wardle, 1990). Detailed study of the growth of one such species, itchgrass (*Rottboellia cochinchinensis* Lour.), in 36 day/night temperature regimes ranging from 17/11 to 32/26°C confirmed its responsiveness to simulated global warming (Patterson et al., 1979). A warming of 3 °C from 26/20 to 29/23°C increased itchgrass dry weight by 88% and leaf area by 68%. Based on its responses to temperature, itchgrass in the United States' current climate could achieve 75 to 100% of its maximum growth in the Gulf Coast states, the South Atlantic states, the lower Midwest, and the Southwest. Only 50 to 75% of maximum growth would occur in the Middle Atlantic states, the central Midwest and the Sacramento Valley in California. A 3°C warming of these latter areas would effectively shift the zone of 75 to 100% growth northward to include the zone previously supporting only 50 to 75% of maximum growth. Changes in climatic variables other than temperature could influence this hypothetical northward range expansion by itchgrass.

Global warming also could allow the northward range expansion of other exotic noxious weeds in North America such as cogongrass (*Imperata cylindrica* (L.) Beauv), and witchweed (*Striga asiatica* (L.) Ktze) (Patterson, 1995). However, not all exotic weeds will be favored by climatic warming. When the exotic wild proso millet (*Panicum miliaceum* L.) was grown in competition with its native congener, (*Panicum texanum* Buckl.), the latter was superior at 30/24°C while the former was more competitive at 24/18°C (Patterson et al., 1986). Under current climatic conditions, the southward spread of wild proso millet may be at least partially restricted by competition from grass weeds adapted to warm conditions. Climatic warming might further restrict the southward range expansion of wild proso millet and other weeds currently limited either by warm temperatures or by competitive pressure from warm-adapted weeds. The relative advantage conferred to either weed or crop will depend on the competing species and on other environmental factors in addition to temperature.

2.3 IMPLICATIONS FOR WEED MANAGEMENT

A major obstacle to projecting the future impact of climate change on agriculture is a lack of knowledge of the effects on agricultural pests (Prestidge and Pottinger, 1990; Stinner et al., 1987; Waggoner, 1983). If the increasing CO₂ acts as a fertilizer, the growth of weeds as well as crops will be stimulated. Which will benefit the most depends on the species, their photosynthetic pathways, and the interacting effects of other environmental factors. Field experiments with conventional fertilizers have demonstrated that increased weed competition can occur in response to supplemental nutrients and that weed competition can eliminate the yield-enhancing effects of fertilization (Carlson and Hill, 1985).

In addition to influencing the competitiveness and geographical distribution of weeds, increasing CO₂ and climate change probably will require modification of current weed control technology (Patterson, 1993, 1995). Environmental factors such as temperature, precipitation, wind, soil moisture, and atmospheric humidity influence the application of herbicides and their effectiveness (Hatzios and Penner, 1982; Muzik, 1976). Variations in these environmental factors accompanying global warming may make appropriate application more difficult, and may affect herbicide injury of crops and other non-target organisms.

Temperature and moisture availability affect both the uptake and metabolism of herbicides by crops and target weeds. The antitranspirant effects of higher CO₂ levels could reduce plant uptake of soil-applied herbicides. Changes in leaf anatomy and leaf surface characteristics, induced directly by high CO₂ or caused by climatic factors, could also affect plant uptake of herbicides. High leaf starch concentrations which commonly occur in C₃ plants grown under CO₂ enrichment (Wong, 1990) might interfere with herbicide activity. On the other hand, increased temperatures and increased metabolic activity tend to increase uptake, translocation, and effectiveness of many herbicides. Increased rhizome and tuber growth in perennial C₃ weeds is a likely consequence of the increased photosynthate supply in high CO₂ environments (Oechel and Strain, 1985). This could increase the difficulty of controlling perennial weeds either mechanically or chemically.

The effectiveness of natural and implemented biological control of weeds and other pests depends on climate. Synchrony between the growth, development, and reproduction of biocontrol agents and their targets, essential for effective long term control, might not be maintained in periods of climatic change, particularly if climatic extremes occur more frequently. On the other hand, global warming could facilitate overwintering of insect populations and favor earlier poleward migrations in the spring. This could increase the effectiveness of biological control of weeds in some cases.

3. Insect Activity and Distribution

Increasing global atmospheric CO₂ may affect insect feeding activity through effects on host plant physiology and chemical composition (Patterson, 1993). More importantly, the accompanying climatic changes will affect the distribution and degree of infestation of insect pests through both direct effects on the life cycles of insects and indirectly through climatic effects on hosts, predators, competitors, and insect pathogens. There is some evidence that the risk of crop loss will increase due to poleward expansion of insect ranges. Agricultural systems create modified and unstable habitats favoring migratory insects that have higher reproduction and mortality rates (Southwood and Comins, 1976). The population size of insects associated with unstable environments tends to be dominated by density-dependent characteristics, so human alteration of conditions that affect host plant survival, irrigation for example, will affect phytophagous (leaf-eating) insect populations. When confronted by climate change, agricultural insects must diapause (enter a state of dormancy), migrate, otherwise adapt genetically, or die.

Insect life cycle processes affected by climate and weather include lifespan duration, fecundity, diapause, dispersal, mortality and genetic adaptation. Porter et al. (1991) listed the following effects of temperature on insects: limiting geographical ranges; over-wintering; population growth rates; number of generations per annum; length of growing season; crop-pest synchronization; interspecific interactions; dispersal and migration; and availability of host plants and refugia. The effects of climate and weather on insect life cycles have been documented for a wide variety of insect pests of agriculture, rangelands, and forests (Dobzhansky, 1965; Fye and McAda, 1972; Tauber et al., 1986; Mattson and Haack, 1987; Kingsolver, 1989; Cammell and Knight, 1991; Harrington and Stork, 1995). A summary of the climatic thresholds for the phenological development of agricultural pest insects can be found in Table 2. Many effects involve changes in the severity of outbreaks following extreme weather events (on the order of hours to weeks). Freezing temperatures are a major factor in mortality but *Drosophila* sp. insects that survived relatively colder temperatures were found to be more fecund than cohorts that were not exposed to the low temperatures (Dobzhansky, 1965). Temperatures that exceed critical thresholds frequently have adverse effects on fecundity as in the cases of bollworm, *Helicoverpa zea* (Boddie), tobacco budworm, *Heliothis virescens* (F.), beet armyworm, *Spodoptera exigua* (Hübner), cabbage looper, *Trichoplusia ni* (Hübner), saltmarsh caterpillar, *Estigmene acrea* (Drury), and pink bollworm, *Pectinophora gossypiella* (Saunders) (Fye and McAda, 1972).

Table 2

Climatic thresholds and responses for the phenological development of agricultural pest insects.

Species	Common	Thresholds	Response Functions
<i>Aculops lycoperici</i>	Tomato russet mite	$T_a < 0.0^\circ\text{C}$	Lethal to all stages for prolonged exposure; 7 days per generation
<i>Aedes aegypti</i>	Salt Marsh mosquito	$T_a = 27.0^\circ\text{C}$	10-14 days per generation
<i>Anabrus simplex</i>	Mormon cricket		60-90 days per generation
<i>Antonina simplex</i>	Rhodes grass mealybug	$T_x > 42.0^\circ\text{C}$ and $T_n < -2.0^\circ\text{C}$; $0.0^\circ\text{C} < T < 38.0^\circ\text{C}$; $29.0^\circ\text{C} < T < 32.0^\circ\text{C}$	Lethal; Physiological development occurs; Optimal physiological development rate occurs; Confined to $30^\circ\text{S} - 30^\circ\text{N}$ latitude;
<i>Blissus leucopterus</i>	Chinch bugs		Frequent rain during hatch reduces infestations; Extremely low temperatures and sudden changes of temperature kill adults in winter and spring; Thrive in hot, dry weather
<i>Carpophilus lugubris</i>	Dusky sap beetle	$T_a = 24.0^\circ\text{C}$	Eggs: 2.5 days; Larvae: 14 days; Adults: 14 days
<i>Chlorochroa say</i>	Say stinkbug	$T > 21.0^\circ\text{C}$	Oviposition occurs
<i>Cochilomyia hominivorax</i>	Screwworm	$T_a < 12.0^\circ\text{C}$	Two-month exposure is lethal to pupae in the soil
<i>Coleophora laricella</i>	Larch casebearer	No lower thermal limit	
<i>Dasyneura laricis</i>	Larch gall midge	$T_a = 0.0^\circ\text{C}$	1600 DD per generation
<i>Dendroctonus ponderosae</i>	Mountain pine beetle	$T_a = 16.0-30.0^\circ\text{C}$; $21.0^\circ\text{C} < T < 38.0^\circ\text{C}$;	Emergence; Flight initiation;

		$T_n < -18.0^\circ\text{C};$ $T_n < -40.0^\circ\text{C}$	Lethal to eggs; Lethal to larvae; Single generation may require 2 years; Multiple generations may develop in the southern part of the range
<i>Dendroctonus micans</i>	Greater European spruce bark beetle	$T = 20.0 - 23.0^\circ\text{C}$	Flight; Eggs: 10-15 days at 20.0°C ; Larvae: 50-60 days at $19.0-23.0^\circ\text{C}$; Pupae: 4-6 days at $19.0-23.0^\circ\text{C}$; Emergence: 44 days at 20.0°C
<i>Diabrotica undecimpunctata undecimpunctata</i>	Western spotted cucumber beetle	Jan. $T_a < 10.0^\circ\text{C}$	No overwintering
<i>Diprion pini</i>	Pine sawfly	$T_n < 10.0^\circ\text{C}$ or $T_x > 35.0^\circ\text{C};$ $T_n < 0.0^\circ\text{C}$ or $T_x > 35.0^\circ\text{C};$ $12.0^\circ\text{C} < T < 25.0^\circ\text{C};$ $T < -20.0^\circ\text{C}$	Lethal to eggs; Lethal to false caterpillars; False caterpillars develop: Eonymphs in cocoons can survive for at least 1 mo.; Incubation: 40 days at 10.0°C ; 18 days at 18.0°C ; 13-14 days at 20.0°C ; 12 days at 25.0°C ; 7 days at 32.0°C
<i>Epirrita autumnata</i>	Autumnal moth	$T_n < -37.0^\circ\text{C}$	Lethal to eggs
<i>Helicoverpa zea</i>	Corn earworm, Cotton bollworm	$T_a = 12.5^\circ\text{C};$ $T_a = 12.6^\circ\text{C}$	185 DD for larval development (on sweet corn); 557 DD per generation (on cotton); Overwintering confined to $< 39^\circ\text{N}$ latitude
<i>Heliothis virescens</i>	Tobacco budworm	$T_a = 12.6^\circ\text{C}$	448 DD per generation (on cotton)
<i>Hyblaea pueria</i>	Teak defoliator		Larvae 10-12 days at optimal conditions; 21 days average;

			Life cycle 19-36 days
<i>Ips typographis</i>	Spruce bark beetle	$T_x > 47.0^{\circ}\text{C}$	Lethal
<i>Lymantria monacha</i>	Nun moth	$0.0^{\circ}\text{C} < T < 45.0^{\circ}\text{C}$	Vital thermal limits; More frequent outbreaks in low-precipitation sites
<i>Lymantria dispar</i>	Gypsy moth	Precipitation 25-100 cm; $15.0^{\circ}\text{C} < \text{Jul. } T_a < 27.0^{\circ}\text{C};$ $-18.0^{\circ}\text{C} < \text{Jan. } T_a < -12.0^{\circ}\text{C}$ $T_a = 20.0^{\circ}\text{C}$	Photoperiod does not control diapause; Dispersal delayed by precipitation or $T < 7.0^{\circ}\text{C};$ Larvae: 44-50 days at 20.0°C
<i>Monochamus alternatus</i>	Japanese pine sawyer	$T_a = 12.5^{\circ}\text{C};$ $T < 18.0^{\circ}\text{C}$	625 DD per generation; No feeding or dispersal; Flight Index (FI) is a measure of infestation potential: $\text{FI} = \# \text{ days with } T_{\text{avg}} > 18.0^{\circ}\text{C} \text{ and } P_{\text{avg}} < 10 \text{ mm, No infestation for } \text{FI} < 60 \text{ days, Severe infestation for } \text{FI} > 80 \text{ days}$
<i>Musca domestica</i>	House fly	$T_a = 27.0^{\circ}\text{C}$	10-12 days per generation
<i>Ostrinia nubilalis</i>	European corn borer	$T_a = 10.0^{\circ}\text{C}$	726 DD per generation; Eggs 7 days; Larvae 35 days; Diapause of mature larvae induced at photophase $\text{LD} < 14.75:9.25;$ Cool, rainy weather in June inhibits oviposition and washes emerging Larvae from plants thus reducing infestations; Dry summers and extremely cold winters increase mortality
<i>Panolis flammea</i>	Pine beauty moth		Obilgatory pupal diapause 120-130 days; Oviposition: $T < 6.0^{\circ}\text{C}$ none; $T = 10.0^{\circ}\text{C}$ 50%; $T = 15.0-20.0^{\circ}\text{C}$ maximal; Eggs: 6.3 days at $20.0^{\circ}\text{C};$ 19.7 days at $10.0^{\circ}\text{C};$

			Larvae: 28 days at 20.0°C; 71 days at 10.0°C; Pupae: susceptible to prolonged water logging, desiccation and min. 1 day exposure to T = -20.0°C
<i>Pectinophora gossypiella</i>	Pink bollworm		25-30 days per generation (midsummer)
<i>Perkinsiella saccharicida</i>	Sugarcane delpacid		Life cycle: 48 days at 25.0°C; 56 days at 22.2°C
<i>Phthorimaea operculella</i>	Potato tuberworm	$T_n < 10.0^\circ\text{C}$	No damage occurs to plants
<i>Pissodes strobi</i>	Spruce weevil	$T_a = 7.2^\circ\text{C}$	785 DD per generation
<i>Pseudaletia unipuncta</i>	Armyworm		Outbreaks follow cold, wet spring weather
<i>Schizaphis graminum</i>	Greenbug	$T_n > 4.5^\circ\text{C}$	
<i>Scolytus ventralis</i>	Fir engraver beetle	$T > 24.0^\circ\text{C}$	Flight initiation
<i>Sirex noctilio</i>	Woodwasp	$T > 20.0^\circ\text{C}$	Enhanced flight and mating; Outbreaks during drought
<i>Spodoptera exigua</i>	Beet armyworm	$T_a = 12.2^\circ\text{C}$	517 DD per generation
<i>Spodoptera frugiperda</i>	Fall armyworm	$T_a = 13.8^\circ\text{C}$	340 DD per generation
<i>Trypodendron lineatum</i>	Striped ambrosia beetle	$T > 15.5^\circ\text{C};$ $19.0^\circ\text{C} < T < 26.0^\circ\text{C};$ $T = 30.0^\circ\text{C}$	Dispersal flight initiated; Optimal for flight; Maximum for flight
<i>Zeiraphera diniana</i>	Larch budmoth	$T_n < -51.3^\circ\text{C};$ $T > 7.0^\circ\text{C}$	Lethal to eggs; Flight initiation

Abnormally cool, wet conditions were associated with high subsequent infestations of cotton by cotton tipworm, *Crociosema plebejana* (Zeller) (Hamilton and Gage, 1986). November precipitation and April temperature were the best indicators of mean grasshopper densities in southern Idaho

(Fielding and Brusven, 1990). Drought can affect various physiological processes of plants which may increase the plants' attractiveness and susceptibility to phytophagous insects (Mattson and Haack, 1987). Intense precipitation has been noted as a deterrent to the occurrence and success of oviposition by insects such as the European corn borer, *Ostrinia nubilalis* (Hübner) (Davidson and Lyon, 1987). Abundant precipitation can affect mortality through, for example, drowning of soil-dwelling insects (Watt and Leather, 1986), but is more likely to indirectly affect insects through climatic effects on insect pathogens, predators and parasites as has been shown for *H. zea* pupae especially under persistently saturated soil conditions (Raulston et al., 1992).

While the specific ways in which climate change could affect persistent wind patterns such as nocturnal wind jets in the U.S. and convergence systems such as the Inter-Tropical Convergence Zone (ITCZ) are poorly predicted by GCMs, changes in the strength, timing, and geographical extent of these systems have been hypothesized. Some of the most important insect pests of agriculture currently use these systems to disperse widely from decaying habitats to viable habitats (Pedgley, 1982). Corn earworm, *H. zea*, moths in the U.S. (Westbrook et al., 1985; Scott and Achtemeier, 1987; Lingren et al., 1993), grasshoppers or locusts (*Ailopus simulatrix* Walker), old world bollworm (*Heliothis armigera* (Hübner)) and whitefly (*Bemisia tabaci* (Gennadius)) disperse via atmospheric transport. In the case of the ITCZ and locusts, the winds also contribute ephemeral precipitation for host plants as reviewed by Joyce (1983) and Rainey (1989). Changes in these systems would affect the speed and range of dispersal of these pests.

Many of the documented effects of climate on insects are based on unusual weather events affecting the severity of insect outbreaks within their normal range; the (unintended) introduction of exotic species into new environments; or the extension of (host) crops into new environments (e.g., through irrigation). Human-induced local climate change as has occurred within urban areas provides evidence of how insects can adapt to changes in their environment. Examples of genetic adaptation include: Queensland fruitfly (*Dacus tryoni* Froggatt), lucerne flea (*Sminthurus viridis* L.) and sheep blowfly (*Lucilia cuprina*) (Wied.) in Australia after introduction from Europe and with the expansion of crop areas; the European corn borer, *O. nubilalis* (Hübner) and the European spruce sawfly, *Gilpinia hercyniae* (Hartig) in North America after introduction from Europe; and of *Drosophila serrata* Malloch across latitudes (Birch, 1965). *Drosophila serrata* Malloch also has responded by natural selection to the changed conditions in large cities (Dobzhansky, 1965). Selection to tighten host-insect interaction (such as codling moth, *Cydia pomonella* (L.), in fruit trees) appears to favor the evolution of isolated host races (Pashley and Bush, 1979). Climatic, agronomic, political, and economic factors jointly led to populations of boll weevil, *Anthonomus grandis grandis* Boheman, that adapted differently across cotton areas in the U.S. (Terranova et

al., 1990). Although insect populations have been redistributed during periods of major climatic change, a continuous intermixing of beetle species' gene pools in the Rocky Mountains of the U.S. was determined to have prevented speciation (Elias, 1991).

Understanding of physiological development and behavior has allowed development of numerical models that estimate insect growth, movement and mortality in response to potential changes in climate (Goodenough and McKinion, 1992). Minimum and maximum temperatures, required cumulative degree days, and, where important, the effects of drought or wet conditions, which are the basis for such models, have been established for many common agricultural pests (e.g., Fye and McAda, 1972; Davidson and Lyon, 1987). Simulation models predict *potential* redistribution of insects under simulated climates.

Specific studies of the likely impacts of climate change were reviewed by Cammel and Knight (1991), Porter *et al.* (1991), Sutherst (1990, 1991) and Sutherst *et al.* (1995). They demonstrate that impacts could be severe in many different environments and involve numerous different species of insect pests. Principal concern is with species that can increase their population size by undergoing an extra generation each year in warmer climates or expand their geographical distributions. For example, Porter *et al.* (1991) found that in Europe *O. nubilalis* would shift 1220 km northward if temperature increased by 3° to 6°C by 2025-2070. For a 3°C temperature increase in Japan, Mochida (1991) predicted expanded ranges for tobacco cut worm (*Spodoptera litura*), southern green stink bug (*Nezera viridula*), rice stink bug (*Lagynotomus eleongatus*), lima-bean pod borer (*Etiella zinckenella*), common green stink bug (*Nezera antennata*), soybean stem gall (*Asphondylia* sp.), rice weevil (*Sitophilus oryzae*), and soybean pod borer (*Legumunuvora glycinivorella*), but a decreased range for rice leaf beetle (*Oulema oryzae*) and rice leaf miner (*Agromyza oryzae*). Vegetation subzones were linked with microclimates to extrapolate spruce weevil hazard zones in British Columbia (Spittlehouse and Sieben, 1994). Models that match discrete ranges of temperature and precipitation, with the presence of particular species (such as CLIMEX), may be especially appropriate for projecting the effects of climate change on insect redistribution (Worner, 1988; Sutherst *et al.*, 1995).

Actual insect distributions under climate change will also depend on host distributions (Rainey, 1989), competition with existing species (DeBach, 1965), adaptability to new conditions, and the presence of natural enemies in the area. Since climate effects on insect life cycles frequently depend on extreme events (e.g., freezes, intense precipitation) and climatic features such as the persistent winds of the Inter-Tropical Convergence system, the reliability of predicted redistribution of insects depends, in part, on the reliability of predictions of these climatic features.

4. Plant Diseases

The occurrence of plant fungal and bacterial pests depends on temperature, rainfall, humidity, radiation, and dew. Climatic conditions affect the survival, growth and spread of pathogens as well as the resistance of their hosts to infection. Friedrich (1994) summarized the observed relationship between climatic conditions and important plant diseases. Among these, mild winters have been associated with more rapid and stronger outbreaks of powdery mildew (*Erysiphe graminis*), brown leaf rust of barley (*Puccinia hordei*), and strip rust of cereals (*Puccinia striiformis*) (Meier, 1985). Mild winters combined with very warm weather conditions provide optimal growth conditions for cercospora leaf spot disease (*Cercospora beticola*), powdery mildew (*Erysiphe betae*), and rizomania disease (*Rizomania*) (Treharne, 1989).

Warm, humid conditions lead to earlier and stronger outbreaks of late potato blight (*Phytophthora infestans*) (Löpmeyer, 1990; Parry et al., 1990). Dry and hot summers generally reduce infestations of most fungal diseases because plant resistance is increased. Summer dryness, particularly in early summer, also decreases rhynchosporium leaf blotch (*Rhynchosporium secalis*) and septoria leaf spot diseases (*Septoria tritici* and *S. nodorum*) but more frequent summer precipitation, particularly heavy storms, would increase incidences of these diseases because rain and rain-born splash water are the means by which disease spores are spread (Royle et al., 1986). Warmer temperatures would likely also shift the occurrence of these diseases into presently cooler regions (Treharne, 1989).

5. Conclusions

Although weeds will be affected directly by increasing atmospheric CO₂ levels, the most significant effects on pests will result from changes in temperature, precipitation distribution, and wind patterns. Climatic change will alter the geographic range and impact of many crop pests. New pest/crop associations may occur. Some old problems may worsen, but others may become less severe. Efforts to control or manage pests will be affected. Environmental variation and increased incidence of extreme weather may reduce the effectiveness of pesticides on targeted pests or result in more injury to non-target organisms. Natural and managed biological control may be affected either negatively or positively. Overall, the challenge to agriculture from pests probably will increase.

References

- Bazzaz, F.A., and Garbutt, K.: 1988, 'The Response of Annuals in Competitive Neighborhoods: Effects of Elevated CO₂'. *Ecology* 69, 937-946.
- Birch, L.C.: 1965, 'Evolutionary Opportunity for Insects and Mammals in Australia'. in Baker, H.G., and Stebbins, G.L. (eds.), *The Genetics of Colonizing Species*, Academic Press, New York, N.Y., pp. 197-211.
- Cammell, M.E., and Knight, J.D.: 1991, 'Effects of Climate Change on the Population Dynamics of Crop Pests'. *Advances in Ecological Research* 22, 117-162.
- Carlson, H. L., and Hill, J.E.: 1985, 'Wild Oat (*Avena fatua*) Competition with Spring Wheat: Effects of Nitrogen Fertilization'. *Weed Sci.* 34, 29-33.
- Carter, D.R., and Peterson, K.M.: 1983, 'Effects of a CO₂ Enriched Atmosphere on the Growth and Competitive Interaction of a C₃ and a C₄ Grass'. *Oecologia* 58, 188-193.
- Davidson, R.H., and Lyon, W.F.: 1987, *Insect Pests of Farm, Garden and Orchard*, John Wiley & Sons, New York, 640 pp.
- DeBach, P.: 1965, 'Some Biological and Ecological Phenomena Associated with Colonizing Entomophagous Insects', in Baker, H.G., and Stebbins, G.L. (eds.), *The Genetics of Colonizing Species*, Academic Press, New York, N.Y., pp. 287-303.
- Dobzhansky, T.: 1965, 'Wild' and 'Domestic' Species of *Drosophila*', in Baker, H.G., and Stebbins, G.L. (eds.), *The Genetics of Colonizing Species*, Academic Press, New York, N.Y., pp. 533-546.
- Elias, S.A.: 1991, 'Insects and Climate Change: Fossil Evidence from the Rocky Mountains'. *BioSci Am Inst Biol Sci.* 41(8), 552-559.
- Fielding, D.J., and Brusven, M.A.: 1990, 'Historical Analysis of Grasshopper (Orthoptera: Acrididae) Population Responses to Climate in Southern Idaho, 1950-1980'. *Environmental Entomology* 19(6), 1786-1791.
- Flint, E.P., and Patterson, D.T.: 1983, 'Interference and Temperature Effects on Growth in Soybean (*Glycine max*) and Associated C₃ and C₄ Weeds'. *Weed Science* 31, 193-199.
- Flint, E.P., Patterson, D.T., Mortensen, D. A., Riechers, G. H., and Beyers, J. L.: 1984, 'Temperature Effects on Growth and Leaf Production in Three Weed Species'. *Weed Science* 32, 655-663.
- Friedrich, S.: 1994, 'Wirkung Veränderter Klimatischer Faktoren auf Pflanzenschädlinge', in Brunnert, H., and Dämmgen, U. (eds.), *Klimaveränderungen und Landwirtschaft*, Part II, Lanbauforsch, Völknerode, Spec. 148, pp. 17-26.
- Fye, R.E., and McAda, W.C.: 1972, *Laboratory Studies on the Development, Longevity, and Fecundity of Six Lepidopterous Pests of Cotton in Arizona*, Agricultural Research Service, Technical Bulletin No. 1454, U.S. Department of Agriculture, Washington, DC.
- Furtick, W. R.: 1978, 'Weeds and World Food Production', in Pimentel, D. (ed.), *World Food, Pest Losses, and the Environment*, AAAS Selected Symp. 13., Westview Press, Boulder, CO., pp. 51-62.
- Goodenough, J.L., and Mckinion, J.M. (eds.): 1992, *Basics of Insect Modeling*, ASAE Monograph No. 10., Amer. Soc. Agric. Engrs., St. Joseph, MI., 221 pp.
- Hamilton, J.G., and Gage, S.H.: 1986, 'Outbreaks of the Cotton Tipworm, *Crociosema plebejana* (Lepidoptera: Tortricidae), Related to Weather in Southeast Queensland, Australia'. *Environmental Entomology* 15, 1078-1082.
- Harlan, J. R.: 1975, *Crops and Man*, American Society of Agronomy, and Crop Science Society of America, Madison, WI.
- Harrington, R., and Stork, N.E. (eds.): 1995, *Insects in a Changing Environment*, Academic Press, London, 535 pp.
- Hatzios, K.K., and Penner, D.: 1982, *Metabolism of Herbicides in Higher Plants*, CEPSCO iv.,

- Burgess Publ., Edina, MN.
- Holm, L.G., Plucknett, D.L., Pancho, J.V., and Herberger, J.P.: 1977, *The World's Worst Weeds: Distribution and Biology*, Univ. of Hawaii Press, Honolulu.
- Joyce, R.J.V.: 1983, 'Aerial Transport of Pests and Pest Outbreaks'. *EPPO Bulletin* 13(2), 111-119.
- Kingsolver, J.G.: 1989, 'Weather and the Population Dynamics of Insects: Integrating Physiological and Population Ecology'. *Physiol. Ecol.* 62(2), 314-334.
- Lingren, P.D., Bryant, V.M., Jr., Raulston, J.R., Pendleton, M., Westbrook, J., and Jones, G.D.: 1993, 'Adult Feeding Host Range and Migratory Activities of Corn Earworm, Cabbage Looper, and Celery Looper (Lepidoptera: Noctuidae) Moths as Evidenced by Attached Pollen'. *Journal of Economic Entomology* 86, 1429-1439.
- Löpmeier, F.J.: 1990, 'Klimaimpakforschung aus Agrarmeteorologischer'. *Sicht. Bayer. Landw. Jarhb.* 67(1), 185-190.
- Mattson, W.J., and Haack, R.A.: 1987, 'The Role of Drought in Outbreaks of Plant-eating Insects'. *BioScience* 37(2), 110-118.
- Meier, W.: 1985, *Pflanzenschutz im Feldbau*. Tierische Schädlinge und Pflanzenkrankheiten, Eidgenössische Forschungsanstalt für Landwirtschaftlichen Pflanzenbau, 8 Auflage, Zürich-Reckenholz.
- Mochida, O.: 1991, 'Impact of CO₂-climate Change on Pests Distribution'. *Agr. Horticult.* 66, 128-136.
- Muzik, T.J.: 1976, 'Influence of Environmental Factors on Toxicity to Plants', in Audus, L.J. (ed), *Herbicides: Physiology, Biochemistry, Ecology*, 2nd ed., 2, Academic Press, New York, pp. 204-247.
- Oechel, W.C., and Strain, B.R.: 1985, 'Native Species Responses to Increased Atmospheric Carbon Dioxide Concentration', in Strain, B.R., and Cure, J.D. (eds), *Direct Effects of Increasing Carbon Dioxide on Vegetation*, DOE/ER-0238, Department of Energy, Washington, DC.
- Parker, C., and Fryer, J.D.: 1975, 'Weed Control Problems Causing Major Reductions in World Food Supplies'. *FAO Plant Protection Bulletin* 23, 83-95.
- Parry, M.L., Porter, J.H., and Carter, T.R.: 1990, 'Agriculture: Climatic Change and its Implications'. *Trends in Ecology and Evolution* 5, 318-322.
- Pashley, D.P., and Bush, G.L.: 1979, 'The Use of Allozymes in Studying Insect Movement with Special Reference to the Codling Moth, *Laspeyresia pomonella* (L.) (Olethreutidae)', in Rabb, R.L., and Kennedy, G.G. (eds), *Movement of Highly Mobile Insects: Concepts and Methodology in Research*, North Carolina State University Press, Raleigh, North Carolina, pp. 333-341.
- Patterson, D.T.: 1993, 'Implications of Global Climate Change for Impact of Weeds, Insects, and Plant Diseases', in *International Crop Science*, vol. 1, Crop Science Society of America, Madison, Wisconsin, pp. 273-280.
- Patterson, D.T.: 1995, 'Weeds in a Changing Climate.' *Weed Science* 43, 685-701.
- Patterson, D.T., and Flint, E. P.: 1980, 'Potential Effects of Global Atmospheric CO₂ Enrichment on the Growth and Competitiveness of C₃ and C₄ Weed and Crop Plants'. *Weed Science* 28, 71-75.
- Patterson, D.T. and Flint, E.P.: 1990, 'Implications of Increasing Carbon Dioxide and Climate Change for Plant Communities and Competition in Natural and Managed Ecosystems', in Kimball, B.A., Rosenberg, N.J., and Allen, L.H., Jr. (eds), *Impact of CO₂, Trace Gases, and Climate Change on Global Agriculture*, Special Publication 53, American Society of Agronomy, Madison, WI, pp. 83-110.
- Patterson, D.T., Meyer, C.R., Flint, E.P., and Quimby, P.C., Jr.: 1979, 'Temperature Responses and Potential Distribution of Itchgrass (*Rottboellia exaltata*) in the United States'. *Weed Science* 27, 77-82.
- Patterson, D.T., Flint, E.P., and Beyers, J.L.: 1984, 'Effects of CO₂ Enrichment on Competition Between a C₄ Weed and a C₃ Crop'. *Weed Science* 32, 101-105.
- Patterson, D. T., Russell, A.E., Mortensen, D. A., Coffin, R. D., and Flint, E.P.: 1986, 'Effects of Temperature and Photoperiod on Texas Panicum (*Panicum texanum*) and Wild Proso Millet

- (*Panicum miliaceum*)', *Weed Science* 32, 101-105.
- Pedgley, D.E.: 1982, *Windborne Pests and Diseases: Meteorology of Airborne Organisms*, Ellis Horwood Ltd., West Sussex, England, 250 pp.
- Porter, J.H., Parry, M.L., and Carter, T.R.: 1991, 'The Potential Effects of Climatic Change on Agricultural Insect Pests'. *Agricultural Forest Meteorology* 57, 221-240.
- Prescott-Allen, R., and Prescott-Allen, C.: 1990, 'How Many Plants Feed the World?' *Conservation Biology* 4, 365-374.
- Prestidge, R. A., and Pottinger, R. P.: 1990, 'The Impact of Climate Change on Pests, Diseases, Weeds and Beneficial Organisms Present in New Zealand Agricultural and Horticultural Systems', New Zealand Ministry for the Environment, Wellington.
- Rahman, A., and Wardle, D. A.: 1990, 'Effects of Climate Change on Cropping Weeds in New Zealand', in Prestidge, R. A., and Pottinger, R. P. (eds.), *The Impact of Climate Change on Pests, Diseases, Weeds and Beneficial Organisms Present in New Zealand Agricultural and Horticultural Systems*, New Zealand Ministry for the Environment, Wellington, pp. 107-112.
- Rainey, R.C.: 1989, *Migration and Meteorology: Flight Behavior and the Atmospheric Environment of Migrant Pests*, Oxford University Press, New York, 314 pp.
- Raulston, J.R., Pair, S.D., Loera, J., and Cabanillas, H.E.: 1992, 'Prepupal and Pupal Parasitism of *Helicoverpa zea* and *Spodoptera frugiperda* (Lepidoptera: Noctuidae) by *Steinernema* sp. in Corn Fields in the Lower Rio Grande Valley'. *Journal of Economic Entomology* 85, 1666-1670.
- Royle, D.J., Shaw, M.W., and Cook, R.J.: 1986, 'Patterns of Development of *Septoria nodorum* and *S. tritici* in Some Winter Wheat Crops in Western Europe, 1981-1983'. *Plant Pathology* 35, 466-476.
- Scott, R.W., and Achtemeier, G.L.: 1987, 'Estimating Pathways of Migrating Insects Carried in Atmospheric Winds'. *Environmental Entomology* 16(6), 1244-1254.
- Southwood, T.R.E., and Comins, H.N.: 1976, 'A Synoptic Population Model'. *Journal of Animal Ecology* 45(3), 949-965.
- Spittlehouse, D.L., and Sieben, B.: 1994, 'Mapping the Effect of Climate on Spruce Weevil Infestation Hazard', in *Preprints of the 11th Conference on Biometeorology and Aerobiology*, American Meteorology Society, Boston, MA, pp. 448-450.
- Stinner, B.R., Taylor, R.A.J., Hammond, R.B., Purrington, F.F., McCartney, D.A., Rodenhouse, N., and Barrett, G.W.: 1987, 'Potential Effects of Climate Change on Plant-pest Interactions', in Smith, J., and Tirpak, D. (eds.), *The Potential Effects of Global Climate Change on the United States*, U.S. Environmental Protection Agency, Washington, DC, pp. 8.1-8.35.
- Sutherst, R.W.: 1990, 'Impact of Climate Change on Pests and Diseases in Australasia'. *Search* 21, 230-232.
- Sutherst, R.W.: 1991, 'Pest Risk Analysis and the Greenhouse Effect'. *Review of Agricultural Entomology* 79, 1177-1187.
- Sutherst, R.W., Maywald, G.F., and Skarrate, D.B.: 1995, 'Predicting Insect Distributions in a Changed Climate' in Harrington, R., and Stork, N.E. (eds.), *Insects in a Changing Environment*, Academic Press, London, pp. 59-91.
- Tauber, M.J., Tauber, C.A., and Masaki, S.: 1986, *Seasonal Adaptations of Insects*, Oxford University Press, New York, NY, 411 pp.
- Terranova, A.C., Jones, R.G., and Bartlett, A.C.: 1990, 'The Southeastern Boll Weevil: An Allozyme Characterization of its Population Structure'. *Southwestern Entomology* 15(4), 481-496.
- Treharne, K.: 1989, 'The Implications of the 'Greenhouse Effect' for Fertilizers and Agrochemicals', in Bennet, R.M. (ed.), *The 'Greenhouse Effect and UK Agriculture*, No. 19, Center for Agricultural Strategy, University of Reading, Reading, UK, pp. 67-78.
- Waggoner, P.E.: 1983, 'Agriculture and a Climate Changed by More Carbon Dioxide'. in *Changing Climate: Report of the Carbon Dioxide Assessment Committee*, National Academy Press, Washington, DC, pp. 383-418.
- Waggoner, P.E.: 1993, 'Assessing Nonlinearities and Surprises in the Links of Farming to Climate and Weather', in Darmstadter, J., and Toman, M.A. (eds.), *Assessing Surprises and*

- Nonlinearities in Greenhouse Warming*, Resources for the Future, Washington, D.C. pp. 45-65.
- Westbrook, J.K., Wolf, W.W., Pair, S.D., Sparks, A.N., and Raulston, J.R.: 1985, 'An Important Atmospheric Vehicle for the Long Range Migration of Serious Agricultural Pests of the South Central United States', in *Preprints of the 7th Conf. on Biometeorol. and Aerobiol.*, American Meteorological Society, Boston, MA, pp. 281-282.
- Watt, A.D., and Leather, S.R.: 1986, 'The Pine Beauty in Scottish Lodgepole Pine Plantations', in Berryman, A.A. (ed.), *Dynamics of Forest Insect Populations: Patterns, Causes, Implications*, Plenum Press, New York, pp. 243-266.
- Woodward, F.I.: 1988, 'Temperature and the Distribution of Plant Species', in Long, S.P., and Woodward, F.I. (eds.), *Plants and Temperature*, The Company of Biologists Ltd., Univ. of Cambridge, Cambridge, UK, pp. 59-75.
- Woodward, F.I., and Williams, B.G.: 1987, 'Climate and Plant Distribution at Global and Local Scales'. *Vegetatio* **69**, 189-197.
- Wong, S.C.: 1990, 'Elevated Atmospheric Partial Pressure of CO₂ and Plant Growth. II. Non-structural Carbohydrate Content in Cotton Plants and its Effect on Growth Parameters'. *Photosynthesis Research* **23**, 171-180.
- Worner, S.P.: 1988, 'Ecoclimatic Assessment of Potential Establishment of Exotic Pests'. *Journal Economic Entomology* **81**(4), 973-983.
- Wray, S. M., and Strain, B. R.: 1987, 'Competition in Old-field Perennials Under CO₂ Enrichment'. *Functional Ecology* **1**, 145-149.

Received 18 December 1997; in revised form 11 June 1999