

## Acute toxicity of carbon dioxide to juvenile marine shrimp *Litopenaeus vannamei* (Boone 1931)

Plínio S. Furtado, Carlos A. P. Gaona, Fabiane P. Serra, Luis H. Poersch & Wilson Wasielesky Jr.

To cite this article: Plínio S. Furtado, Carlos A. P. Gaona, Fabiane P. Serra, Luis H. Poersch & Wilson Wasielesky Jr. (2017): Acute toxicity of carbon dioxide to juvenile marine shrimp *Litopenaeus vannamei* (Boone 1931), *Marine and Freshwater Behaviour and Physiology*, DOI: 10.1080/10236244.2017.1371568

To link to this article: <http://dx.doi.org/10.1080/10236244.2017.1371568>



Published online: 04 Sep 2017.



Submit your article to this journal [↗](#)



Article views: 4



View related articles [↗](#)



View Crossmark data [↗](#)



SHORT COMMUNICATION



## Acute toxicity of carbon dioxide to juvenile marine shrimp *Litopenaeus vannamei* (Boone 1931)

Plínio S. Furtado, Carlos A. P. Gaona, Fabiane P. Serra, Luis H. Poersch and Wilson Wasielesky Jr.

Marine Station of Aquaculture, Institute of Oceanography, Federal University of Rio Grande, Rio Grande, Brazil

### ABSTRACT

Elevated concentrations of dissolved carbon dioxide (CO<sub>2</sub>) and reduced pH levels are observed during the culture and transportation of aquatic organisms. Studies on the toxicity effects of CO<sub>2</sub> in penaeid shrimp are scarce when compared to the amount of research in fish. The objective of the present study was to determine the lethal concentration and safety levels of CO<sub>2</sub> for juvenile white shrimp *Litopenaeus vannamei*. Juveniles (1.76 ± 0.36 g) were exposed for 96 h to one of six concentrations of dissolved CO<sub>2</sub> (14.5, 23.8, 59.0, 88.0, 115.0, and 175.0 mg/L) or a control condition (without the addition of CO<sub>2</sub>), and their survival was monitored for 96 h. The LC<sub>50</sub> values with 95% confidence limits at 24, 48, 72, and 96 h were 130.05 (104.2–162.1), 77.2 (73.8–80.02), 69.65 (65.47–74.32), and 59.12 (53.08–66.07) mg/L of CO<sub>2</sub>, respectively. The calculated safety level was 5.9 mg/L of CO<sub>2</sub>, and the highest concentration that did not induce significantly higher mortality than that observed in controls (NOEC) was 23.8 mg/L of CO<sub>2</sub>. We recommend that CO<sub>2</sub> levels should be kept below the safety level obtained in this study.

### ARTICLE HISTORY

Received 24 November 2016  
Accepted 22 August 2017

### KEYWORDS

Toxicity; carbon dioxide; safety level; pH; *Litopenaeus vannamei*

## Introduction

Researchers have recently observed an increase in the concentration of dissolved carbon dioxide (CO<sub>2</sub>) in the oceans and a reduction of 0.1 pH unit in the water surface since the pre-industrial era (Caldeira and Wickett 2003). The pH in the oceans is predicted to be reduced by 0.2–0.3 units by the year 2100 (Feely et al. 2009).

Several studies were conducted to analyze the effects of ocean acidification on larval, juvenile, and adult invertebrates (Kurihara 2008; Bierbower and Cooper 2010; Carter et al. 2013). The results obtained by Kurihara (2008) showed that, with chronic exposure of the shrimp *Palaemon pacificus* to the predicted concentrations of CO<sub>2</sub> over the next centuries, the survival, growth and possibly the reproduction of these organisms may be reduced.

The consumption of oxygen and the release of CO<sub>2</sub> occur during the transportation of aquatic organisms in closed environments such as plastic bags and generate a stressful condition of hypoxia and hypercapnia (Jensen et al. 2014). In fish and shrimp cultures, CO<sub>2</sub>

concentrations are increased mainly due to their biomass and decaying organic matter in the culture environment (Timmons and Ebeling 2010). The Pacific white shrimp, *Litopenaeus vannamei*, is the most widely farmed species of shrimp in superintensive bioflocs technology (BFT) systems. Stocking densities in these systems are between 150 and 600 shrimp/m<sup>2</sup>, and the biomass is between 2 and 8 kg/m<sup>2</sup> (Silva et al. 2013).

In a BFT system, if the number of heterotrophic organisms exceeds the number of photoautotrophic microorganisms, then CO<sub>2</sub> production also exceeds the production of dissolved oxygen. This may accumulate CO<sub>2</sub> and reduce the pH levels of the culture water (Vinatea et al. 2010; Furtado et al. 2011). Under fault conditions in the aeration system or power failure, dissolved oxygen concentrations decrease rapidly, CO<sub>2</sub> concentrations increase and the water is acidified (Furtado et al. 2014). According to Van Wyk and Scarpa (1999), CO<sub>2</sub> concentrations below 5 mg/L in the water are ideal and 20 mg/L is considered acceptable for penaeid shrimp. Concentrations between 20 and 60 mg/L are not lethal but can interfere in the exchange of CO<sub>2</sub> in the gills, and concentrations above 60 mg/L can be lethal. Dissanayake and Ishimatsu (2011) found a synergistic effect of elevated CO<sub>2</sub> concentrations on temperature, which can impair the breathing and swimming capacities of penaeids.

The CO<sub>2</sub> is an essential carbon source for photosynthesis but is also the final product of the respiration of aquatic organisms (Boyd 2008). Initially, the conditions for culture in a BFT system are predominantly photoautotrophic. CO<sub>2</sub> concentrations can fluctuate between 5 and 10 mg/L in the morning but may exceed 20 mg/L in ponds with high biomass (Boyd 2008). Over a 24-h period, dissolved oxygen (DO), pH and carbon dioxide levels can oscillate. During the light phase of the day, CO<sub>2</sub> is removed and pH and DO increase as a result of photosynthetic processes; however, at night, photosynthesis stops and respiration predominates, and the release of CO<sub>2</sub> by aerobic organisms results in reduced pH and DO in the culture water (Vinatea 2004). Over time, CO<sub>2</sub> is permanently produced, and there is minimal production of DO by phytoplankton within the dominant heterotrophic communities in the BFT system.

Excretion of CO<sub>2</sub> in the gill epithelium is hampered when CO<sub>2</sub> concentrations exceed 20 mg/L, resulting in a decline of the hemolymph pH. This process adversely affects the transport oxygen from the hemolymph, reduces tissue oxygenation and increases the ventilation rate (Taylor and Whiteley 1989; Van Wyk and Scarpa 1999). An intra- or extra-cellular maladjustment in the acid-base balance can damage the metabolism of shrimp (Wang et al. 2012). This is possible even if decapod crustaceans possess the ability to tolerate basic and slightly acidic conditions, due to the exchange of Na<sup>+</sup>/H<sup>+</sup> and Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> occurring in their gills (Henry et al. 1981).

Studies on the effects of CO<sub>2</sub> in penaeid shrimp are scarce when compared to the amount of research in fish (Grottum and Sigholt 1996; Santos et al. 2013). The aim of this study was to determine the lethal concentrations and safety level of carbon dioxide for *L. vannamei* juveniles.

## Materials and methods

### Location and facilities

This study was conducted at the Marine Aquaculture Station Prof. Marcos Alberto Marchiori (EMA) of the Oceanographic Institute – Federal University of Rio Grande (FURG), located at Cassino Beach, Rio Grande (32°11' S, 52°10' W), Rio Grande do Sul, Brazil.

## Water, shrimp, and experimental design

The water used in the experiment was pumped from the beach and filtered through a sand filter and a cartridge with a pore size of 5  $\mu\text{m}$ . No water was exchanged throughout the study. Dechlorinated fresh water using 1 ppm ascorbic acid and intense aeration for 12 h and was added to restore the volume lost through evaporation.

The shrimp used in this study was obtained from the laboratory Aquatec Ltda (Canguaretama, Rio Grande do Norte, Brazil). After the *L. vannamei* nauplii arrived at EMA, they were maintained in the larviculture sector until they reached the post-larvae 15 (pL15) stage. Thereafter, they were cultivated in a nursery (1500 pL/m<sup>2</sup>) in a BFT system, where they reached a mean weight of  $1.76 \pm 0.36$  g. A total of 210 *L. vannamei* juveniles in the intermolt stage (2 days after molting) were acclimated to the experimental conditions for 48 h to reduce the stress of capture and handling. During this period, they were fed twice per day with commercial feed (Guabi® 38% crude protein and 8% ethereal extract) in the proportion of 8% of biomass/day. The experimental photoperiod was 12 h light: 12 h dark at 27 °C.


The acute test to determine the median lethal concentration (96 h-LC<sub>50</sub>) was conducted in a semi-static system. Animals were not fed during the test. Each rectangular tank had a usable volume of 30 L and was stocked with 10 shrimp, and each treatment had three replicates ( $n = 30$  shrimp). Six concentrations of carbon dioxide were tested (14.5, 23.8, 59.0, 88.0, 115.0, 175.0 mg/L) in addition to a control treatment (without the addition of CO<sub>2</sub>).

CO<sub>2</sub> was injected into the experimental units through a pressurized gas cylinder weighing 30 kg and equipped with a manometer, control valve and carbon dioxide flow meters (scale of 0–15 L/min). Each treatment consisted of a recirculation system with three rectangular tanks of 30 L and a 200 L circular reservoir, which had a submerged pump (capacity of 1000 L/h) with a Venturi tube that injected CO<sub>2</sub> from the cylinder (Figure 1). The pump transferred water from the reservoir to the rectangular tanks, solubilizing CO<sub>2</sub> into the water during the process. The water exchange rate in each experimental unit was 150 L/h and water returned to the reservoir by gravity. Six air diffusers provided aeration in each 200 L reservoir, and a thermostat was used to maintain the temperature at 27 °C.

Temperature, pH and oxygen were measured every 2 h using a pH meter model 100 YSI® (Yellow Springs Instruments, USA) and an oximeter 55 YSI® (Yellow Springs Instruments, USA). Salinity was verified at the beginning and end of the experiment with an optical refractometer (Atago®, Japan). Total ammonia (TAN) concentrations (NH<sub>3</sub> + NH<sub>4</sub><sup>+</sup>) and alkalinity were measured every 24 h according to the methodologies recommended by UNESCO (1983) and APHA (1998), respectively. The dissolved carbon dioxide concentrations were measured according to APHA (1998) and calculated with the software CO<sub>2</sub> Analysis Salt® (Timmons and Ebeling 2010). CO<sub>2</sub> concentrations and shrimp survival were monitored every 2 h (12 times per day) over a 96 h period. Shrimp were considered dead when they were still and did not respond to mechanical stimulation with a glass rod (Lin and Chen 2003). The dead shrimp were removed from the tank as soon as they were detected.

## Statistical analyses

The cumulative data of mortality for the periods of 24, 48, 72, and 96 h were used to estimate the median lethal concentration (LC<sub>50</sub>) and its confidence limits (95%) using the Trimmed Spearman Karber Method software (Hamilton et al. 1977). The safety level was calculated



**Figure 1.** Diagram illustrating the experimental design where: (1) a carbon dioxide cylinder ( $\text{CO}_2$ ); (2) a pressure gage and regulating valve; (3) flow meters; (4) a circular tank with 200 L of usable volume; (5) a submerged pump with a Venturi tube for  $\text{CO}_2$  entry; (6) an air diffuser for water oxygenation; (7) a heater with a thermostat; (8) a water register for flow adjustment; (9) 30 L experimental tanks for the exposure of the shrimp to  $\text{CO}_2$ ; and (10) a water pipe to return water from the experimental units to the circular tank.

using a 0.1 factor, according to Sprague (1971). The NOEC (i.e. the highest concentration that did not induce significantly higher mortality than that observed in controls) (EPA 2002) was calculated by comparing the number of survivors in each treatment at the end of the test.

We used the software STATISTICA 7.0<sup>®</sup> (StatSoft Inc. 2004, Tulsa, Oklahoma, USA) for the statistical analysis. To check for significant differences between the data, an analysis of variance (ANOVA, one-way) was performed after the assumptions of homocedasticity of variances (Levene) and data distribution normality (Kolmogorov-Smirnov) were confirmed. Tukey's test was used to compare the means when significant differences were detected between treatments ( $p < 0.05$ ).


## Results

The mean values  $\pm$  the standard deviation of the water quality parameters monitored throughout the test are shown in Table 1. There were no significant differences in the salinity, temperature, and alkalinity ( $p > 0.05$ ) between treatments. However, significant differences were found ( $p < 0.05$ ) for dissolved oxygen, pH, and TAN. The mean values of dissolved oxygen were above 6 mg/L at  $\text{CO}_2$  concentrations below 59.0 mg/L and differed significantly from other treatments with higher  $\text{CO}_2$  concentrations. The treatment with 175.0 mg/L showed the lowest dissolved oxygen level (5.6 mg/L). There were significant differences in pH among all of the treatments (Figure 2), and pH showed an inverse relationship between the values of  $\text{CO}_2$ . Thus, pH values decreased as the concentrations of

**Table 1.** Water quality parameters (mean  $\pm$  SD) measured during a 96 h test with different concentrations of carbon dioxide (CO<sub>2</sub>) with juvenile *L. vannamei*.

CO <sub>2</sub> treatment (mg/L)	Salinity (‰)	Temperature (°C)	DO (mg/L)	pH	TAN (mg/L)	Alkalinity (mgCaCO <sub>3</sub> /L)
0.6 $\pm$ 0.3 <sup>a</sup>	30.2 $\pm$ 0.3	27.2 $\pm$ 0.6	6.24 $\pm$ 0.18 <sup>a</sup>	8.06 $\pm$ 0.05 <sup>a</sup>	0.20 $\pm$ 0.13 <sup>a</sup>	105.2 $\pm$ 1.4
14.5 $\pm$ 0.9 <sup>b</sup>	30.3 $\pm$ 0.3	27.0 $\pm$ 0.54	6.21 $\pm$ 0.15 <sup>a</sup>	6.79 $\pm$ 0.08 <sup>b</sup>	0.14 $\pm$ 0.09 <sup>a</sup>	105.1 $\pm$ 1.7
23.8 $\pm$ 1.5 <sup>c</sup>	30.5 $\pm$ 0.2	26.9 $\pm$ 0.87	6.22 $\pm$ 0.14 <sup>a</sup>	6.57 $\pm$ 0.13 <sup>c</sup>	0.18 $\pm$ 0.11 <sup>a</sup>	104.7 $\pm$ 2.6
59.0 $\pm$ 3.6 <sup>d</sup>	30.1 $\pm$ 0.6	26.96 $\pm$ 0.7	6.0 $\pm$ 0.12 <sup>ab</sup>	6.17 $\pm$ 0.1 <sup>d</sup>	0.12 $\pm$ 0.06 <sup>a</sup>	104.7 $\pm$ 2.5
88.0 $\pm$ 6.5 <sup>e</sup>	30.4 $\pm$ 0.3	27.1 $\pm$ 0.70	5.75 $\pm$ 0.25 <sup>b</sup>	6.03 $\pm$ 0.04 <sup>e</sup>	0.08 $\pm$ 0.04 <sup>ab</sup>	104.6 $\pm$ 2.5
115.0 $\pm$ 10.5 <sup>f</sup>	30.6 $\pm$ 0.3	26.95 $\pm$ 0.9	5.65 $\pm$ 0.27 <sup>b</sup>	5.88 $\pm$ 0.1 <sup>f</sup>	0.06 $\pm$ 0.03 <sup>b</sup>	104.2 $\pm$ 2.0
175.0 $\pm$ 15.6 <sup>g</sup>	30.8 $\pm$ 0.5	26.81 $\pm$ 0.7	5.61 $\pm$ 0.3 <sup>b</sup>	5.7 $\pm$ 0.05 <sup>g</sup>	0.05 $\pm$ 0.01 <sup>b</sup>	103.9 $\pm$ 1.5

Notes: Different superscript letters in each column indicate significant differences ( $p < 0.05$ ) among the treatments. DO: dissolved oxygen. TAN: total ammonia nitrogen.



**Figure 2.** pH in the water during 96 h of study in the different concentrations of CO<sub>2</sub> (mg/L).

CO<sub>2</sub> increased. The measured TAN concentrations were higher in the treatments from 0.6 to 88.0 mg/L, which differed from the treatments of 115 and 175 mg/L.


The accumulated mortality data along the 96-h test presented in the Figure 3. All *L. vannamei* juveniles exposed to the concentrations of 175, 115, and 88 mg/L of CO<sub>2</sub> died within 36, 48, and 84 h of exposure, respectively. There was mortality of shrimp exposed to 59.0 mg/L of CO<sub>2</sub> for 64 h, and it reached 33% after 96 h. No mortalities were observed in the treatments with 0.6, 14.5 and 23.8 mg/L of CO<sub>2</sub>.

The calculated 96 h LC<sub>50</sub> was 59.12 mg/L of CO<sub>2</sub> with a 95% confidence interval from 53.08 to 66.07 mg/L of CO<sub>2</sub>. LC<sub>50</sub> values of 24, 48, 72, and 96 h are shown in Table 2. The safety level was 5.9 mg/L of CO<sub>2</sub>, approximately 25% of the highest concentration tested which did not cause mortality (NOEC) (23.8 mg/L of CO<sub>2</sub>).

## Discussion

The chemical and physical parameters of water quality measured in this study remained within the range considered adequate for the survival of juvenile *L. vannamei*, except for the treatments with CO<sub>2</sub> concentrations greater than 23.8 mg/L (Ponce-Palafox et al. 1997; Van Wyk and Scarpa 1999; Lin and Chen 2001).

TAN concentrations increased over time but did not reach levels that could affect the survival of shrimp (Lin and Chen 2001). There was a difference in TAN levels due to the premature mortality of shrimp in the treatments with concentrations greater than 88.0 mg/L of CO<sub>2</sub>; therefore, the treatments with lower CO<sub>2</sub> concentrations showed higher TAN levels because the shrimp survived and excreted for a longer time.



**Figure 3.** Cumulative mortality (%) of shrimp during 96 h of exposure to different concentrations of CO<sub>2</sub>.

**Table 2.** Median lethal concentration (LC<sub>50</sub>) data for marine shrimp *L. vannamei* juveniles exposed to CO<sub>2</sub>. Confidence limits are 95%.

Time (h)	LC <sub>50</sub> CO <sub>2</sub> (mg/L)	Confidence limits (mg/L)
24	130.05 <sup>a</sup>	(104.2–162.1)
48	77.2 <sup>b</sup>	(73.8–80.02)
72	69.65 <sup>bc</sup>	(65.47–74.32)
96	59.12 <sup>c</sup>	(53.08–66.07)

Note: Different superscript letters in the same column indicate significant differences ( $p < 0.05$ ).

According to Van Wyk and Scarpa (1999), dissolved oxygen concentrations above 5.0 mg/L are considered ideal for the growth of shrimp. In the present study, DO concentrations remained above 5.0 mg/L in all of the treatments, even those with higher CO<sub>2</sub> concentrations. DO concentrations decreased with the increase in CO<sub>2</sub>, which was also verified by Hall and Van Ham (1998). Under hypoxic conditions, the toxicity of CO<sub>2</sub> was increased, as it interfered in the oxygen absorption capacity (Boyd and Tucker 1998). To avoid lethal concentrations of CO<sub>2</sub> during the transportation of shrimp, which typically has hypoxic conditions, the biomass must be reduced and the pH must be adjusted with alkalizing compounds (Jensen et al. 2014).

Suitable pH values for the culture of penaeid shrimp are in the range of 7–9 (Van Wyk and Scarpa 1999). However, penaeids have a resistance to acidic conditions generated by the addition of hydrochloric acid, the values of lethal pH<sub>50</sub> at 24, 48, 72, and 96 h for juvenile *L. vannamei* had acidic pH values of 3.86, 3.92, 3.94, and 4.04 (Furtado et al. 2015). Hall and Van Ham (1998) stated that the incorporation of CO<sub>2</sub> reduces water pH from 8.5 to 5.9, generating behavioral changes and increasing glucose levels in the hemolymph of *Penaeus monodon* when compared to the same pH reduction by the addition of sulfuric acid. The lowest pH values measured in this study were higher than those generating mortality in the studies developed by Furtado et al. (2015). However, mortality rates in this study were 100% at a pH 6.03.

This difference is due to a higher diffusion capacity of CO<sub>2</sub> than that of hydrogen ions (H<sup>+</sup>), as molecular CO<sub>2</sub> directly diffuses faster than protons through cell membranes (Gutknecht et al. 1977). A change in the hemolymph pH creates favorable conditions for the production of reactive oxygen species, resulting in oxidative stress and damage to cell structures such as lipid membranes, proteins, and DNA (Lushchak 2011; Wang et al. 2012).



Kikkawa et al. (2008) evaluated the tolerance of juvenile Kuruma prawn, *Marsupenaeus japonicus*, to acute exposure to CO<sub>2</sub> and calculated the 72 h LC<sub>50</sub> as 14.3% of CO<sub>2</sub>. A LC<sub>50</sub> value of approximately 50% for the same exposure period was found in this study, which was lower than those obtained in the Kikkawa et al. study. Therefore, juvenile *L. vannamei* are more sensitive to CO<sub>2</sub>. The results of this study contradict the literature recommendations (Van Wyk and Scarpa 1999; Boyd 2008) because shrimp exposed to concentrations below 60 mg/L of CO<sub>2</sub> had a mortality rate of 30%. During the present study, we observed a loss of equilibrium in the shrimp before death. Such behavior was also observed by Kikkawa et al. (2008).

An acute toxicity test provides information about the relative lethality of a toxin but cannot predict the concentration that has sublethal and chronic effects on organisms (Buikema et al. 1982). In the present study, we calculated the safety level to be 5.9 mg/L of CO<sub>2</sub> using the factor 0.1 as suggested by Sprague (1971); therefore, *L. vannamei* juveniles can be cultivated in the presence of that level of CO<sub>2</sub> without mortalities. However, we do not recommend levels as NOEC of 23.8 mg/L of CO<sub>2</sub> to be maintained in farming systems for longer periods than those tested in this study.

Wickins (1984) studied the chronic effects of hypercapnia and found that juvenile *P. occidentalis* exposed for 56 days to pH 7.3 showed reduced growth when compared to shrimp farmed in sea water without the addition of CO<sub>2</sub> (pH 7.6). That study also showed a decrease in the growth of *P. monodon* juveniles as the pH decreased with the injection of CO<sub>2</sub> and longer intervals between molts (8.2 and 6.6 days for pH 6.4 and 7.4, respectively).

Circadian monitoring of pH and DO during culture is important for decision-making to help prevent high concentrations of CO<sub>2</sub> and hypoxic conditions, which can cause shrimp mortality and hinder the success of the culture. Several alkalizing compounds may be applied to correct pH and CO<sub>2</sub> levels such as calcium hydroxide, sodium carbonate and bicarbonate (Furtado et al. 2011). Based on chemical reactions, approximately 0.84 mg/L of calcium hydroxide or 0.64 mg/L of calcium oxide are needed to remove 1 mg/L of CO<sub>2</sub>. As calcium hydroxide does not readily dissolve to react completely with CO<sub>2</sub>, the concentrations mentioned above must be doubled (Boyd 2008). Another strategy to remove CO<sub>2</sub> is the use of degassing columns (Moran 2010).

## Conclusion

In this study of the acute toxicity of CO<sub>2</sub>, the 96 h LC<sub>50</sub> was found to be 59.12 mg/L of CO<sub>2</sub>, and the safety level was 5.9 mg/L of CO<sub>2</sub>. The highest non-lethal concentration was 23.8 mg/L of CO<sub>2</sub>. We recommend that CO<sub>2</sub> levels are maintained below the safety limit calculated for the species to avoid hypercapnia conditions, which can cause immunosuppression in the shrimp, making them more susceptible to diseases.

## Acknowledgements

The authors are grateful for the financial support provided by the National Council for Scientific and Technological Development (CNPq), Ministry of Fisheries and Aquaculture (MPA) and Higher Level Personnel Training Coordination (CAPES). Wilson Wasielesky Jr. and Luis Henrique Poersch are fellow researchers of CNPq.



## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by the National Council for Scientific and Technological Development (CNPq), Ministry of Fisheries and Aquaculture (MPA) and Higher Level Personnel Training Coordination (CAPES).

## References

- [APHA] American Public Health Association. 1998. Standard methods for the examination of water and wastewater. 20th ed. Washington (DC): American Public Health Association; p. 1193.
- Bierbower SM, Cooper RL. 2010. The effects of acute carbon dioxide on behavior and physiology in *Procambarus clarkii*. J Exp Zool Part A 313(8): 484–497.
- Boyd CE. 2008 Jul/Aug. Carbon dioxide: waste, nutrient. Glob Aquac Advoc. 62–65.
- Boyd CE, Tucker CS. 1998. Pond aquaculture water quality management. Boston (MA): Kluwer Academic; p. 700.
- Buikema AL, Niederlehner BR, Cairns JJ. 1982. Biological monitoring part IV – toxicity testing. Water Res. 16:239–262.
- Caldeira K, Wickett ME. 2003. Oceanography: anthropogenic carbon and ocean pH. Nature. 425:365.
- Carter HA, Ceballos-Osuna L, Miller NA, Stillman JH. 2013. Impact of ocean acidification on metabolism and energetics during early life stages of the intertidal porcelain crab *Petrolisthes cinctipes*. J Exp Biol. 216:1412–1422.
- Dissanayake A, Ishimatsu A. 2011. Synergistic effects of elevated CO<sub>2</sub> and temperature on the metabolic scope and activity in a shallow-water coastal decapod (*Metapenaeus joyneri*; Crustacea: Penaeidae). J Mar Sci. 68(6):1147–1154.
- [EPA] Environmental Protection Agency. 2002. Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms. Washington (DC): US EPA.
- Feely RA, Doney SC, Cooley SR. 2009. Ocean acidification: present conditions and future changes in a high-CO<sub>2</sub> world. Oceanography. 22(4):36–47.
- Furtado PS, Poersch LH, Wasielesky W. 2011. Effect of calcium hydroxide, carbonate and sodium bicarbonate on water quality and zootechnical performance of shrimp *Litopenaeus vannamei* reared in bio-flocs technology (BFT) systems. Aquaculture. 321:130–135.
- Furtado PS, Serra FP, Poersch LH, Wasielesky Jr. W. 2014. Short communication: acute toxicity of hydrogen peroxide in juvenile white shrimp *Litopenaeus vannamei* reared in biofloc technology systems. Aquacult Int. 22(2):653–659.
- Furtado PS, Fugimura MMS, Monserrat JM, Souza DM, Garcia LO, Wasielesky Jr. W. 2015. Acute effects of extreme pH and its influences on the survival and biochemical biomarkers of juvenile White Shrimp, *Litopenaeus vannamei*. Mar Freshwater Behav Physiol. 48(6):417–429. DOI: <https://doi.org/10.1080/10236244.2015.1086539>.
- Grottum JA, Sigholt T. 1996. Acute toxicity of carbon dioxide on European seabass (*Dicentrarchus labrax*): mortality and effects on plasma ions. Comp Biochem Physiol A. 115(4):323–327.
- Gutknecht J, Bisson MA, Tosteson FC. 1977. Diffusion of carbon dioxide through lipid bilayer membranes: effects of carbonic anhydrase, bicarbonate, and unstirred layers. J Gen Physiol. 69:779–794.
- Hall MR, Van Ham EH. 1998. The effects of different types of stress on blood glucose in the Giant Tiger Prawn *Penaeus monodon*. J World Aquacul Soc. 29(3):290–299.
- Hamilton MA, Russo RC, Thurston RV. 1977. Trimmed Spearman-Kärber method for estimating median lethal concentrations in toxicity bioassays. Environ Sci Technol. 11(7):714–719.
- Henry R, Kormanik G, Smatresk N, Cameron J. 1981. The role of CaCO<sub>3</sub> dissolution as a source of HCO<sub>3</sub><sup>-</sup> for the buffering of hypercapnic acidosis in aquatic and terrestrial decapod crustaceans. J Exp Biol. 94:269–274.

- Jensen L, Furtado PS, Fugimura MMS, Garcia LO, Poersch LH, Verani JR, Wasielesky Jr. W. 2014. The effect of stocking density on the transport of pink shrimp *Farfantepenaeus brasiliensis* (Crustacea: Decapoda), as live bait for sport fishing in Brazil. *Lat Am J Aquat Res.* 42(1):204–212.
- Kikkawa T, Watanabe Y, Katayama Y, Kita J, Ishimatsu A. 2008. Acute CO<sub>2</sub> tolerance limits of juveniles of three marine invertebrates, *Sepia lycidas*, *Sepioteuthis lessoniana*, and *Marsupenaeus japonicus*. *Plankton and Benthos Res.* 3(3):184–187.
- Kurihara H. 2008. Effects of CO<sub>2</sub>-driven ocean acidification on the early developmental stages of invertebrates. *Mar Ecol Prog Ser.* 373:275–284.
- Lin YC, Chen JC. 2001. Acute toxicity of ammonia on *Litopenaeus vannamei* Boone juveniles at different salinity levels. *J Exp Mar Biol Ecol.* 259:109–119.
- Lin YC, Chen JC. 2003. Acute toxicity of nitrite on *Litopenaeus vannamei* (Boone) juveniles at different salinity levels. *Aquaculture.* 224:193–201.
- Lushchak VI. 2011. Environmentally induced oxidative stress in aquatic animals. *Aquat Toxicol.* 101:13–30.
- Moran D. 2010. Carbon dioxide degassing in fresh and saline water. I. Degassing performance of a cascade column. *Aquacult Eng.* 43:29–36.
- Ponce-Palafox J, Martinez-Palacios CA, Ross LG. 1997. The effects of salinity and temperature on the growth and survival rates of juvenile white shrimp, *Penaeus vannamei*, Boone, 1931. *Aquaculture.* 157:107–115.
- Santos GA, Schrama JW, Capelle J, Rombout JHWM, Verreth JAJ. 2013. Effects of dissolved carbon dioxide on energy metabolism and stress responses in European seabass (*Dicentrarchus labrax*). *Aquacult Res.* 44:1370–1382.
- Silva AF, Lara GR, Ballester ELC, Krummenauer D, Abreu PC, Wasielesky Jr. W. 2013. Effect of high stocking densities on growth and survival of *Litopenaeus vannamei* in final growout phase, reared in biofloc technology (BFT) system. *Ciênc anim bras.* 14(3):279–287.
- Sprague JB. 1971. Measurement of pollutant toxicity to fish – III. Sublethal effects and “safe” concentrations. *Water Res.* 5:245–266.
- Taylor EW, Whiteley NM. 1989. Oxygen transport and acid-base balance in the haemolymph of the lobster, *Homarus gammarus*, during aerial exposure and resubmersion. *J Exp Biol.* 144:417–436.
- Timmons MB, Ebeling JM. 2010. *Recirculating Aquaculture*. 2nd ed. Ithaca (NY): Cayuga Aqua Ventures; 998 pp. <http://www.bee.cornell.edu/aquain18/11/2010>.
- UNESCO. 1983. Chemical methods for use in marine environmental monitoring. Manual and Guides 12, Intergovernmental Oceanographic Commission. Paris, France.
- Van Wyk P, Scarpa J. 1999. Water quality and management. In: Van Wyk P, Davis-Hodgkins M, Laramore R, Main KL, Mountain J, Scarpa J. editors. *Farming marine shrimp in recirculating freshwater systems*. Tallahassee: Florida Department of Agriculture and Consumer Services; p. 141–161.
- Vinatea LA. 2004. Princípios químicos de qualidade da água em aquicultura: uma revisão para peixes e camarões [Chemical Principles of Aquaculture Water Quality: A Review for Fish and Shrimp]. 2ª edição revisada e ampliada. Florianópolis: Editora UFSC; 231 pp.
- Vinatea L, Galvez AO, Browdy CL, Stokes A, Venero J, Haveman J, Lewis BL, Lawson A, Shuler A, Leffler JW. 2010. Photosynthesis, water respiration and growth performance of *Litopenaeus vannamei* in a super-intensive raceway culture with zero water exchange: Interaction of water quality variables. *Aquacult Eng.* 42:17–24.
- Wang W-N, Li B-S, Liu J-J, Shi L, Alam MJ, Su S-J, Wu J, Wang L, Wang A-L. 2012. The respiratory burst activity and expression of catalase in white shrimp, *Litopenaeus vannamei*, during long-term exposure to pH stress. *Ecotoxicology.* 21:1609–1616.
- Wickins JF. 1984. The effect of hypercapnic sea water on growth and mineralization in penaeid prawns. *Aquaculture.* 41:37–48.