

Accepted Manuscript

Erythrocytes as a biological model for screening of xenobiotics toxicity

Mayada Ragab Farag, Mahmoud Alagawany

PII: S0009-2797(17)30848-7

DOI: [10.1016/j.cbi.2017.11.007](https://doi.org/10.1016/j.cbi.2017.11.007)

Reference: CBI 8143

To appear in: *Chemico-Biological Interactions*


Received Date: 2 August 2017

Revised Date: 24 October 2017

Accepted Date: 7 November 2017

Please cite this article as: M.R. Farag, M. Alagawany, Erythrocytes as a biological model for screening of xenobiotics toxicity, *Chemico-Biological Interactions* (2017), doi: 10.1016/j.cbi.2017.11.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Erythrocytes as a biological model for screening of xenobiotics toxicity**Mayada Ragab Farag^{1*}, Mahmoud Alagawany^{2*}**

¹Forensic Medicine and Toxicology Department, Veterinary Medicine Faculty, Zagazig University, Zagazig 44511, Egypt; ²Poultry Department, Agriculture Faculty, Zagazig University, Zagazig 44511, Egypt

***Corresponding authors:** mragabfa@zu.edu.eg, mmalagwany@zu.edu.eg

Abstract

Erythrocytes are the main cells in circulation. They are devoid of internal membrane structures and easy to be isolated and handled providing a good model for different assays. Red blood cells (RBCs) plasma membrane is a multi-component structure that keeps the cell morphology, elasticity, flexibility and deformability. Alteration of membrane structure upon exposure to xenobiotics could induce various cellular abnormalities and releasing of intracellular components. Therefore the morphological changes and extracellular release of haemoglobin [hemolysis] and increased content of extracellular adenosine triphosphate (ATP) [as signs of membrane stability] could be used to evaluate the cytotoxic effects of various molecules. The nucleated RBCs from birds, fish and amphibians can be used to evaluate genotoxicity of different xenobiotics using comet, DNA fragmentation and micronucleus assays. The RBCs could undergo programmed cell death (eryptosis) in response to injury providing a useful model to analyze some mechanisms of toxicity that could be implicated in apoptosis of nucleated cells. Erythrocytes are vulnerable to peroxidation making it a good biological membrane

model for analyzing the oxidative stress and lipid peroxidation of various xenobiotics. 23
 The RBCs contain a large number of enzymatic and non-enzymatic antioxidants. The 24
 changes of the RBCs antioxidant capacity could reflect the capability of xenobiotics to 25
 generate reactive oxygen species (ROS) resulting in oxidative damage of tissue. These 26
 criteria make RBCs a valuable *in vitro* model to evaluate the cytotoxicity of different 27
 natural or synthetic and organic or inorganic molecules by cellular damage measures. 28

Keywords: Erythrocytes, xenobiotic, toxicity, ROS, eryptosis, blood. 29

1. Introduction 30

Red blood cells (RBCs) represent the main cells in the body circulatory system, and they 31
 act as oxygen transporters. The RBCs of mammals are devoid of nucleus and other 32
 cellular organelles while non-mammalian RBCs (birds, fish and amphibians) are 33
 nucleated cells however, both types of RBCs have a plasma membrane with specific 34
 composition and structure which is highly correlated to their biological functions [1,2]. 35

The main component in the RBCs membrane is protein which represents about 39.5% 36
 followed by lipids [35.1%] then water [19.5%] and finally carbohydrates [5.8%] [3]. The 37
 structure of RBCs membrane resembles that of other eukaryotic cells. It contains a 38
 phospholipids bilayer, integral proteins stabilized in the lipid leaflet by covalent bonds 39
 and hydrophobic bonds and a membrane skeleton which is a multi-protein complex 40
 composed of some structural proteins including spectrin [α and β], actin, ankyrin and 41
 protein 4.1. [4]. The interaction between the membrane components make it elastic and 42
 soft and enable its fluctuations and thus provide RBC with elasticity, flexibility and 43
 deformability which are very important for RBC to keep its structural integrity and 44
 original shape and to protect it from the force of the circulating fluid applied on it during 45

their passage into the microcirculation [5]. This interaction is metabolically active and need [occur in] the presence of Adenosine Triphosphate [ATP] which is important to protect the cells from fragmentation and the membrane from vesiculation [6, 7]. The cytoskeleton is also very important to keep the cellular components particularly, haemoglobin [Hb] which is the major protein in RBC and it is responsible for transporting oxygen [O₂] and carbon dioxide [CO₂] to and from tissues. Therefore, the level of intra cellular haemoglobin could be used for determining the cytoplasmic viscosity. Meanwhile, the fluidity of membrane lipids depends generally on the type of phospholipids, the acyl chain length, the degree of fatty acid saturation, the presence of free or esterified cholesterol in addition to some amphipathic compounds such as lisophosphatides [8, 9]. Therefore, the changes in the RBCs membrane could be used as indicators for the cell physiological conditions or their possible alterations. The RBCs could be isolated and handled easily so that they could provide a good model for many assays [10, 11]. The application of *in vitro* cytotoxicity assays in RBCs could be used as alternative tools to screen and evaluate the toxicity of different xenobiotic [12]. Additionally, the high concentration of polyunsaturated fatty acids in RBCs membrane, the high oxygen tension, and redox active hemoglobin molecules [the source of reactive oxygen species in RBC] make them a good biological lipid membrane model especially for screening the oxidative stress conditions induced by xenobiotic [13].

Exposure of erythrocytes to oxidative stress lead to lipid peroxidation that could alter the membranes of RBCs inducing membrane protein conformation and protein cross-linking by decreasing membrane protein content and consequently lead to abnormal cell morphology and hemolysis that could disturb the microcirculation [14, 15]. Xenobiotics

including oxidants could trigger the RBCs programmed death which is known as eryptosis and exhaust their antioxidant defense mechanism [16]. Some cysteine proteases-caspases present in RBCs could be also activated after exposure to oxidative stress such as caspase 8 which is a mediator bounded to RBC membrane that could initiate the cellular cascade for apoptosis and caspase 3 which is an effector mediator causing proteolysis of cellular proteins [17]. Energy depletion in the cell beside osmotic shock could also trigger eryptosis [18]. This review is an overview on the different changes that could occur to RBCs in response to different xenobiotics and the role of these alterations in understanding the different mechanisms of xenobiotics toxic actions with an enumeration of some methods for determination of such effects used in previous studies depended on RBCs as a biological model for cytotoxicity assays.

2. Erythrocyte cellular abnormalities

The hematological changes serve as an early indicator for screening the toxic impacts of xenobiotics on tissues [19]. Toxic substances could induce either direct or indirect damage to the RBC cytoskeleton and disturb cell metabolism and ion permeability of RBCs, thus lead to abnormalities in the cellular morphology [20].

The nucleated RBCs from birds, fish and amphibians considered as important models for studying cellular abnormalities. The abnormality in RBCs morphology of such species could be classified in to two main categories: the first one is nuclear abnormalities [ENA] which include: binucleates [BN], lobed nuclei [LB], notched [NT], nuclear bud [NBu] and vacuolated nuclei [VN] according to the identification of Fenech et al. [21]. The second category is cytoplasmic abnormalities [ECAs] such as acanthocytes [AC], echinocytes [EC], vacuolated cytoplasm [VC], notched cytoplasm [NC] and microcytes

[MC] [22]. The shape of RBCs from different animal species and structure of RBCs 92
 membrane are represented in Figures (1) and (2). 93

Most of erythrocytes abnormalities were detected after exposure to toxic substances such 94
 as pesticides, metals, chemical agents, irradiation and some types of drugs [8]. Immature 95
 pycnotic and mitotic RBCs and with severe anemia were observed in chicken after lead 96
 poisoning and poikilocytosis and anisocytosis in swans [23]. Anisocytosis, Poikilocytosis 97
 and altered hematocrit and mean corpuscular volume with changes in hemoglobin were 98
 also observed in rats exposed to lead [24]. Exposure to lead shot changes nuclear 99
 morphology in the blood of domestic fowl (young chickens) [increased RBCs with 100
 pycnotic nuclei, enucleated RBCs and reticulocytes] [25]. Exposure of Swiss albino mice 101
 to lead during gestation and lactation led to various hematological disorders in RBCs of 102
 neonates and abnormal types and sizes [Macrocytosis] and reduced their life span 103
 resulting in anemia [26]. 104

Fragmented RBCs [schistocytes] and RBCs with larger sizes were reported after 105
 exposure to phenylhydrazine in calf and aluminum in rats [27]. Comelekoglu et al. [28] 106
 stated that some pesticides may cause alterations in surface shapes and size of human 107
 RBCs. Zeni et al. [29] reported the occurrence of echinocytes in *Ictalurus melas* after 108
 exposure to the anionic detergent [sodium dodecyl benzene sulphonate] as a result of 109
 adaptation of the cellular physiological parameters required for shape maintenance. In 110
 addition, Koc et al. [30] reported changes in surface shapes and structural defects of 111
 RBCs of rats exposed to malathion and endosulfan. Suwalsky et al. [31] have reported 112
 that human RBCs when incubated with aqueous extract of *Aristotelia chilensis* showed 113
 morphological alterations represented by echinocytic form. Chlorpyrifos provoked 114

alterations in the cytoskeleton [protein and lipids] of RBCs from Wistar rats thus
affecting the cell surface area [32].

3. Characterization of erythrocyte morphology and size

Isolation of RBCs occurs by centrifugation of heparinized blood at 4 °C for 10 minutes at
3000 rpm, and then buffy coat and plasma were discarded. The erythrocytes were
pelleted by centrifugation after being washed once with 0.9% NaCl solution and twice
with ice-cold phosphate buffered saline [PBS] [33].

The abnormalities of RBCs morphology could be detected by smearing of pelleted RBCs
on glass slides and left to dry then the slides were fixed in absolute methanol for 15
minutes and stained with freshly prepared Giemsa stain and examined under microscope
as described by Sharma et al. [26]. The morphological abnormalities of RBCs could be
also detected by examining the packed erythrocytes under scanning electron microscope
[SEM] after fixation with glutaraldehyde according to method of Agrawal and Sultana
[34]. While, alterations in the relative size of RBCs could be determined using flow
cytometry using forward scatter [FSC] correlating with cell volume and size [35].

4. Haemolysis, osmotic fragility and protein content in the hemolysate

The *in vitro* haemolytic assay using spectrophotometer represents an effective and easy
test for the quantitative measuring of haemolysis [36]. The *in vitro* haemolytic activity
test has been reported to be an alternative method in screening for the cytotoxicity of
various compounds. It is rapid, reproducible and costless test so it could decrease the use
of experimental animals for *in vivo* testing [37]. Many researchers have used the
haemolysis assay for evaluating the cytotoxicity of some herbal plants, mushrooms [36,

38] and extracts from different algae [39] to exclude the natural or synthetic products with possible cytotoxic impact which has been prepared for pharmaceutical uses [40].

The changes in osmotic pressure of RBCs could also change their osmotic fragility and cell integrity so it is could be also used as a diagnostic tool in hemolytic conditions [41]. The *in vitro* osmotic fragility of erythrocytes could be determined by the method of Chikezie [42].

The RBCs membrane resistance has been used by many authors as tool in assessment of toxicity for example chlorpyrifos [43], fluoride [44] and 2,4-dichlorophenoxyacetic [45] toxicities in rat RBCs. The haemolytic assay is based on measuring the release of haemoglobin from RBCs suspended in solution with gradual reducing the concentration and detecting the cells that showed osmotic lysis [concentration-response]. The concentration of released hemoglobin [protein] is correlated to the percentage of lysed cells. Hemoglobin and protein contents in the hemolysate could be determined photometrically at 540 nm. The absorption of the hemolysate of RBCs lysed in distilled H₂O was defined as being 100% haemolysis. The percentage of haemolysis could be calculated by the following equation:

$$Haemolysis(\%) := \frac{Abs_{sample}}{Abs_{erythrocytes \text{ in water}}} \times 100$$

5. Changes in cellular energy [ATP]

The ATP is used by RBCs to maintain osmotic stability and keep submembrane skeletal-network proteins thereby maintain membrane shape and control deformation [46]. Meyers and Hendricks [47] reported that cellular injuries decreased the oxidative phosphorylation process inside the cell resulting in reduced ATP content which consequently forces cells to

show vacuolization and lead also to unequal distributed hemoglobin which resulted in RBCs cytoplasmic vacuoles [48]. Exposure of rat RBCs to lead [Pb] shortened their lifespan due to inhibition of the Na-K-ATPase and loss of membrane integrity [49]. A more recent study on cyadox suggested its role in induction of energy depletion in isolated rabbit RBCs by decreasing their ATP contents [50]. Sikora et al. [51] stated that the only source of extracellular ATP is cell lysis. So determination of ATP content of RBCs can give idea about the status of their membrane integrity and energy charge as well.

Measurement of the ATP and determination of erythrocytes energy charge: The measurement of the ATP content could be performed according to the method developed by Adams [52] with ATP expressed as $\mu\text{mol/g Hb}$ or through measuring the intracellular ATP content using luciferin–luciferase assay kit. To determine the adenylates contents, ADP and AMP were measured as the difference after their enzymatic conversion to ATP [53]. The adenylate energy charge [EC] was calculated by the equation; $EC = \frac{[ATP] + 1/2[ADP]}{[ATP] + [ADP] + [AMP]}$ [54].

6. Erythrocytes as a tool in genotoxicity assays

Screening for genotoxic impact is of great importance during the evaluation of xenobiotic cytotoxicity as the genotoxic potential is usually implicated in carcinogenic and reproductive toxicities as a primary risk factor therefore; genotoxicity testing are helpful for describing the ability of different xenobiotics to damage the cellular genetic information and to induce mutation. For analyzing the effect of a genotoxic molecule, DNA damage in cells exposed should be evaluated. The DNA damage can be in the form of single-strand, double-strand breaks, cross-linking, loss of excision repair, point

mutations, alkali-labile sites and chromosomal aberrations [structural and numerical] 181
[55]. 182

Genotoxic effect of xenobiotics on isolated RBCs could be evaluated using different 183
assays including micronucleus assay, comet assay and DNA fragmentation assay as 184
follow: 185

The micronucleus assay [MN]: The MN is an easy test used as a marker of genotoxic 186
effect of different pollutants in bio-monitoring studies for assessing of their 187
cytogenotoxic potential in field and in laboratory conditions. The appearance of 188
micronucleus in a cell indicated the occurrence of chromosomal aberrations during 189
mitosis [numerical or structural] [21]. The MN assay was performed earlier on the 190
mammalian RBCs [especially rodents], then it has been widely used along with other 191
cellular abnormalities [ENA and ECA] as endpoints in several species of fish and other 192
aquatic organisms [56] and also in different species of birds [57] exposed to various 193
toxicants. Tan et al. [58] reported that there is a positive correlation between the 194
frequencies of MN and other nuclear abnormalities. The MN test was helpful in 195
evaluating genotoxicity of CdCl₂ in gill cells and haemocytes of *Dreissena polymorpha* 196
[59]. The MN was observed in RBCs of chickens exposed to endosulfan [57]. To detect 197
micronuclei in erythrocytes, few drops of whole blood were directly smeared on clean 198
slides. The slides were left for 24 h to dry, fixed for 10 min in methanol, stained by 10% 199
Giemsa , then analyzed using a 1000· oil-immersion lens [60]. 200

The comet assay: The comet assay is rapid, potent and economic test for studying DNA 201
damage induced by chemical and physical agents [61]. This assay is sensitive in 202
detecting DNA damage even if the level of damage is low, and could be performed on 203

small number of cells and could generate data at individual cell level. It has been widely employed in environmental cytogenotoxicity monitoring in a variety of fish species. Comet assay in RBCs was helpful in detecting cadmium chloride induced genotoxicity and cytotoxicity in freshwater fish *Labeo rohita* [35].

DNA fragmentation assay: The DNA fragmentation of the RBCs could be isolated according to the method developed by Weil et al. [62]. Then the fragmented DNA was analyzed by agarose gel electrophoresis at 40 V for 5 h using agarose gel [1%] then examined and photographed as described by Wang et al. [63].

7. Erythrocyte programmed death [Eryptosis]

The death of RBCs caused by injury is a more complex process than haemolysis and it resembles the apoptosis [suicidal death of nucleated cells] which is important for the disposal of deformed cells without rupturing the cellular membranes [18]. Apoptosis of nucleated cells has been used for evaluating the toxic effects of various xenobiotics that could be implicated in disrupting the cell function and tissue destruction [64]. Apoptosis occurs through a chain of events, including nuclear condensation and shrinkage, DNA fragmentation, mitochondrial depolarization, cell membrane blebbing and exposure of phosphatidyl serine on the plasma membrane [breakdown of phosphatidylserine asymmetry of the plasma membrane] [65].

Several researches showed that mammalian RBCs however lack nuclei they have the ability to undergo a programmed death [eryptosis] upon exposure to injurious materials and endogenous challenges which affect the cell integrity and arrest their life cycle [39, 66]. Therefore, eryptosis could be used as a model for analyzing different mechanisms that are of the same importance for the apoptosis of nucleated cells. Simpson and Kling

[67] reported that erythroblasts of dogs exposed to phenylhydrazine showed denucleation without condensation of chromatin. The RBCs of embryonic and newly hatched chickens exposed to cycloheximide or staurosporine showed DNA fragmentation and pycnosis [62]. The RBCs of human respond to lead by surface exposure of phosphatidylserine and shrinkage of cells [68].

During apoptosis, caspases function either as initiators such as caspase-8 and -9 (the membrane-bound mediator initiates the cellular cascade for apoptosis) in response to proapoptotic signals or as effectors such as caspase-3 (the effector mediator leading to cellular proteins proteolysis) . Mature erythrocytes contain considerable amounts of caspase-3 and caspase-8 whereas other essential components of the mitochondrial apoptotic cascade such as caspase-9, Apaf-1 and cytochrome c are absent [69]. Eryptosis could be triggered by osmotic shock and energy depletion in RBCs [17]. Generation of ROS, depletion of antioxidants and formation of adducts are important factors promote death process and lead to the cell disintegration [70]. Chronic arsenic exposure led to ROS generation in the RBCs of rats and consequently led to the activation of caspase 3 [71]. The significant increase in caspase 3 and 8 were also observed in rabbit erythrocytes exposed to cyadox [50].

Some toxic substances can lead to formation of ceramid in RBCs of some animals like rat and mice [44, 72]. This could be returned to the activation of Gardos channel. The Gardos channel activation induces the shrinkage of RBCs and consequently activates the sphingomyelinase [aSMase] and enhance more ceramid generation which triggers suicidal erythrocyte death [73]. Ceramid formation has been also reported to help the super aggregation [capping] of Fas receptors [translocated to lipid raft of RBCs

membrane] and are highly essential for formation of death inducing signaling complex [DISC] and other downstream events associated with Fas induced apoptosis [74]. So, determination of caspase activities, ceramid formation, Gardos channel activation, aSMase activity and Fas aggregation could be used to evaluate the apoptotic effects of xenobiotics on RBCs.

Determining the lifespan of RBCs: Survival of RBC could be measured from the half-life of erythrocyte over time according to the standard method of Sen et al. [75].

Determination of ceramide formation: Ceramide contents in RBCs could be determined by antibody based fluorimetric method [76].

Determination of Gardos channel activity: Gardos channel activity could be determined by the method of Wolff et al. [77].

Determination of aSMase activity in RBCs: aSMase activity could be quantified according to the method of Petrache et al. [78].

Fas aggregation on erythrocyte membrane: For determination of Fas aggregation, the lipid raft should be firstly isolated and this could be done by the method of discontinuous density gradient ultracentrifugation according to Muppidi and Siegel [79], then Fas aggregation [translocation of Fas to lipid raft] could be determined by immunohistochemistry with Fas antibody [74].

Determination of caspase activities: The proteolytic activity of caspase8 and caspase 3 could be measured in RBCs lysate as described in Mukherjee et al. [72].

8. Erythrocytes as an initial screen for oxidative stress

Erythrocytes have been extensively used as a biological membrane model to analyse the oxidative damage as they are highly vulnerable to peroxidation owing to the high content of polyunsaturated fatty acid [PUFA] in their membrane, their role as O_2 and CO_2 transporters and the presence of redox active hemoglobin molecule, which is a potent source of reactive oxygen species [ROS]. They also contain heme-iron which is ferrous hemoglobin essential for the hydrogen peroxide-stimulated oxidation of lipids in the RBCs membrane [13, 80]. Redox regulation in RBCs is shown in Figure (3).

Oxidative damage to RBCs after exposure to xenobiotics [chemicals, drugs, metals, pesticides, and irradiations] induced alterations in morphology of cells, membrane protein conformation, protein cross-linking, lipid peroxidation and consequently hemolysis of RBCs [14, 15]. Oxyhemoglobin and autoxidation are the main sources of intracellular ROS in the RBCs [81]. Oxidative stress impairs oxygen delivery and induces RBCs aging [82]. Different effects of xenobiotics on erythrocytes are shown in Figure (4).

Erythrocytes are protected against oxidative damage by various biological mechanisms including antioxidant enzymes as superoxide dismutase [SOD] and catalase [CAT] [83]. SOD and CAT protect cells by scavenging the free radicals and ROS elimination. SOD converts the highly reactive superoxide anion to a less reactive species, H_2O_2 and to O_2 . SOD contains zinc to keep its stability and copper to maintain its activity [84]. Concerning CAT, more than 98% of blood CAT is present in the RBCs and it is able to remove extra and intra-cellular H_2O_2 giving protecting for tissues deficient in CAT activity [85]. Therefore, blood CAT activity could be used as a good bioindicator for the overall protection against the diffusible H_2O_2 .

Reduced glutathione [GSH] is a non-enzymatic antioxidant that protects lipids and proteins in RBCs membrane and keeps its stability, enhances the RBCs survival against oxidative injury and provides the primary antioxidant defense for the stored RBCs [86]. GSH protects important proteins in RBCs against oxidation and acts as vital sulfhydryl buffer, which keeps SH groups in enzymes and Hb in the reduced state [84]. Therefore, GSH depletion could increase oxidative stress and modify the RBCs components.

4-Hydroxynonenal [4-HNE], is one of the major alpha beta unsaturated aldehydes formed in erythrocytes as a result of peroxidation of membrane lipids after oxidative injury. 4-HNE induced reduction of intracellular GSH by forming GSH-HNE adducts in cytosol and interacted with cytosolic and membrane proteins forming HNE-protein adducts. These adducts resulted in depletion and modulation of antioxidant activities, increasing production of ROS and disruption of RBCs redox status [87]. HNE-protein adducts formations were observed in erythrocytes of rats after exposure to chronic arsenic toxicity [71]. While GSH-HNE adducts were used as a tool to detect lead toxicity on mice RBCs by Mukherjee et al. [72].

Lipid peroxidation [LPO] is one of the consequences of oxidative damage, and it is one of the chief mechanism for cell injury and death [i.e., hemolysis] [84]. Malondialdehyde [MDA], the well-characterized product of the LPO of RBCs, is a highly reactive bifunctional molecule, that could impair various membrane functions by cross-linking the RBCs proteins and phospholipids leading to diminished survival and death [induce hemolysis] [88, 89]. Moreover, LPO of RBCs may be implicated in cell aging, and variable pathological conditions. The determination of MDA level provides a good measure of LPO [90].

Additionally, erythrocytes SOD contains copper and zinc [CuZn-SOD]. The interaction between CuZn-SOD and MDA modified the residues of histidine amino acid and produced protein–protein cross-linked derivatives therefore each type of ROS exhibits a different pattern of protein oxidation. So the protein carbonyl content could be used as an indicator of protein oxidation in erythrocytes as reported by Reddy et al. [91].

Erythrocytes were proved to be a good tool for analyzing the oxidative stress and lipid peroxidation as mechanism of toxic action in various studies. RBCs help in assessing the toxicity of arsenic [71] and 2, 4-dichlorophenoxyacetic [45] in rats; lead toxicity in mice [72] and Lambda-cyhalothrin [92] and cyadox [50] in rabbits.

Some toxic substances could induce pronounced changes in the fatty acid profiles of RBCs membrane due to free radical generation which transferred the both types of membrane fatty acids [saturated and unsaturated] from superficial neutral lipids into phosphatidylethanolamine. For example 2,4-dichlorophenoxyacetic induced cellular deterioration through peroxidation of PUFA ,degradation of membrane phospholipids and increasing the index of fatty acid unsaturation [SFA to UFA ratio] [45]. Alterations of fatty acid composition of RBCs membrane could result in changing the membrane viscosity [93]. Exposure of erythrocytes to ROS generated during arsenic toxicity could result in alterations of membrane proteins [71].

A- Oxidative stress parameters

Preparation of RBCs membrane [erythrocyte ghosts]: erythrocyte ghost could be prepared by the method of Dodge et al. [94].

Fatty acid composition of erythrocytes: The fatty acid composition of erythrocyte was analyzed by gas chromatography after trans-esterification following the method of Giacometti et al. [95]

Electrophoresis of membrane proteins: polyacrylamide gel electrophoresis with dodecyl sulphate [SDS-PAGE] of RBCs membrane proteins could be determined according to Laemmli [96].

Assessment of lipid hydroperoxide content in RBCs membrane: could be estimated using the FOX2 method [97].

Determination of RBCs redox potential: Redox potential in RBC could be determined by the protocol of Biswas et al. [98] from the ratios of pyridine nucleotides. Redox potential was assayed from $\text{NADH}/[\text{NAD}^{++} + \text{NADH}]$ and $\text{NADPH}/[\text{NADP}^{++} + \text{NADPH}]$ ratios.

Estimation of intracellular reactive oxygen species [ROS]: The ROS content of RBCs could be measured by incubating with fluorescent probes dihydroethidium [DHE] and 2',7'-dichlorofluorescein diacetate [DCFDA] according to Lopez-Revuelta et al. [99]; and Zhao et al. [100], then fluorescent-positive cells were detected by flow cytometer according to [101].

The H_2O_2 and $\text{O}_2^{\bullet-}$ concentrations of RBCs: could be determined by the method of Qian et al. [102]. While sulfhydryl group [SH] groups could be measured in RBCs after reaction with 5, 50-dithiobis-[2-nitrobenzoic acid] using the method of Ellman [103].

Measurement of lipid peroxidation of erythrocytes: There are many different methods that could be used for measuring the lipid peroxidation in RBCs. Malondialdehyde

[MDA], a product of lipid peroxidation was measured at 532 nm by using 2-thiobarbituric acid [2, 6-dihydroxypyrimidine-2-thiol; TBA] [104] where MDA concentration was expressed either as nmol/mg protein for the membranes or nmol/gHb for whole erythrocytes. Lipid peroxidation could be assessed indirectly through measurement of the thiobarbituric acid [TBA] reaction [105] where MDA values were expressed as pmol/g Hb. MDA levels could be also measured by the method of Bartosz [106] using the thiobarbituric acid [TBA] method, after the reaction with TBA, the reaction product was measured spectrophotometrically at 535nm where the MDA level was expressed in M/gHb. The method of Tedesco et al. [107] could be also used to measure the level of Lipid peroxidation in RBCs in terms of MDA equivalents using the thiobarbituric acid reaction.

Protein oxidation: Protein carbonyl content is used as a marker of protein oxidation in RBCs and it could be measured using the method described by Uchida and Stadtman[108] or according to Jiang et al[109] .]

Measurement of GSH–HNE and HNE–protein adduct formation: GSH–HNE adduct could be measured using Mass spectrometry as described by Biswas et al. [71]. While HNE–protein adduct formation could be analyzed according to Arguelles et al. [110].

B- Determination of erythrocyte antioxidants

Reduced glutathione [GSH] and oxidized glutathione [GSSG] estimation: GSH content in RBCs could be detected by different methods. It could be measured at 25 °C according to the standard methods using DTNB as described by Tietze [111] where GSH content was expressed as $\mu\text{mol/g Hb}$ or by the method of Dise and Goodman [112]

where GSH was expressed as nmol GSH per mg of hemoglobin. While, intra-cellular 381
GSSG content could be measured at 340 nm as described by Beutler [113]. 382

**Glutathione peroxidase [GSH-Px], Glutathione-s-transferase [GST] and glutathione 383
reductase [GR] activities:** The GSH-Px could be evaluated using Ransel reagents based 384
on the method of Paglia and Valentine [114]. Results were expressed in U/g Hgb. While 385
GST activity was measured spectrophotometrically by the method of Habig et al. [115] 386
using S-2,4-dinitro phenyl glutathione [CDNB] as a substrate. The activity of GST was 387
expressed in terms of nmol/mg protein. The glutathione reductase GR activity was 388
assayed using reagent from Randox Laboratories [GR 2368]; the assay was adapted from 389
the method of Beutler [113]. Results were expressed in U/g Hgb. 390

SOD and CAT activity: The SOD activity could be determined according to the method 391
of Marklund and Marklund [116] by measuring the inhibition of pyrogallol autoxidation. 392
One unit of SOD was defined as the amount of enzyme that inhibits the rate of pyrogallol 393
autoxidation in 50%. The SOD activity in RBCs could be also determined by using the 394
classical NBT method [117] where SOD activity expressed as U/mg Hb. while CAT 395
activity in erythrocyte lysate could be determined according to the method of Aebi [118]. 396
The method is based on the decomposition of H_2O_2 by catalase. Enzyme activity was 397
expressed as units per mg of Hb [U/mg Hb]. 398

Types of damage resulted from interaction of some xenobiotics and RBCs from different 399
animal species are illustrated in Table 1. 400

Conclusion 401

This review shows that erythrocytes could be considered as a valuable model in studying
 the cytotoxic effect of xenobiotics owing to its particular structure and the important
 components of their plasma membrane especially lipids and proteins. The disturbance of
 plasma membrane integrity leads haemoglobin to be released extracellular so the
 hemolytic activity of RBCs could provide useful information about the interaction of
 different molecules and the biological activities on the cell level and could be used to
 exclude the pharmaceutical products with possible cytotoxic effects. The RBCs
 membrane could be altered and deformed in response to injurious xenobiotics resulting in
 cellular abnormalities in morphology and size which could be used as initial testing for
 cytotoxicity. ATP content of RBCs can give idea about the status of their membrane
 integrity and energy charge as the only source of extracellular ATP is lysis of cells.
 Exposure of the body to different physical and chemical agents can result in genotoxic
 and DNA damaging effects and lead to carcinogenic and reproductive toxicities. RBCs
 can be used for evaluating the genotoxic impacts of different xenobiotics using different
 assays including micronucleus assay, comet assay and DNA fragmentation assay.
 The RBCs however lack nuclei they have the ability to undergo a programmed death
 [eryptosis] upon exposure to injurious materials and endogenous challenges. Therefore,
 eryptosis could be used as a model for analyzing different mechanisms that are of the
 same importance for the apoptosis of nucleated cells. Erythrocytes were proved to be a
 good biological membrane model for analyzing the oxidative stress and lipid
 peroxidation as possible mechanisms of toxic action of various xenobiotics because of
 their vulnerability to peroxidation as RBCs contain polyunsaturated fatty acid [PUFA] in
 their membrane and hemoglobin which is a source of ROS and due to its capacity to

transport O₂ and CO₂. Oxidative damage could alter the membrane lipid and protein structure and cellular morphology and induce lipid peroxidation and hemolysis therefore; lipid peroxidation could be also of value in understanding the mechanism of action of various xenobiotics. The RBCs are well equipped by various biological mechanisms of antioxidants including enzymes and non-enzymatic antioxidants which also could be altered by oxidative damage. The changes of component of RBCs, the structure of plasma membrane and the antioxidant capacity of RBCs could be used as good indicators of the ROS generating activity of xenobiotics and their oxidative damaging effects on tissues. All these findings make the RBCs a valuable in vitro initial screening model for the evaluation of cytotoxic mechanisms of various substances and prove their ability to be used as alternative testing methods to decrease animal experimentation, helping reach the goal of reducing, refining and replacing studies conducted with animals which requires major ethical and financial regulations.

Acknowledgement

All the authors of the manuscript thank and acknowledge their respective universities and institutions

Competing interests:

Authors declare that there are no competing interests.

Author contributions:

MRF and MA wrote, revised and reviewed the manuscript.

References

1. Mohandas N, Chasis JA, Shohet SB. The influence of membrane skeleton on red cell 447
deformability, membrane material properties, and shape. *Sem Hematol* 1983; 20: 448
225–242. 449
2. De Oliveira S, Saldanha C. An overview about erythrocyte membrane. *Clin Hemorheol* 450
Micro 2010; 44: 6. 451
3. Bruno-Franco M, Mazzei C. The red blood cell membrane: structure and functions. 452
Blood Transf 2004; 2, 160–180. 453
4. Mohandas N, Gallagher PG. 2008. Red cell membrane: past, present, and future. 454
Blood; 112: 3939–3948 455
5. Park Y, Best CA, Badizadegan K, 2010. Measurement of red blood cell mechanics 456
during morphological changes. *Proc Natl Acad Sci*; 107: 6731–6736. 457
6. Mohandas N, Evans E. 1994. Mechanical properties of the red cell membrane in 458
relation to molecular structure and genetic defects. *Annu Rev Biophys Biomol Struct* 459
23: 787–818. 460
7. Betz T, Lenz M, Joanny J-F, Sykes C. ATP-dependent mechanics of red blood cells. 461
Proc Natl Acad Sci 2009; 106: 15320–15325. 462
8. Mohandas N, Chasis J. 1993. Red blood cell deformability, membrane material 463
properties and shape: regulation by transmembrane, skeletal and cytosolic proteins 464
and lipids. *Sem Hematol*; 30: 171–192 465
9. Plasenzotti R, Windberger U, Ulberth F, Osterode W, Losert U. 2007 Influence of 466
fatty acid composition in mammalian erythrocytes on cellular aggregation. *Clin* 467
Hemorheol Micro; 37: 237–243. 468

10. Alagawany M., M.R. Farag, M.S. El-Kholy, S.A.A. El-Sayed and K. Dhama 2016. 469
 Effect of resveratrol, cinnamaldehyde and their combinations on the antioxidant 470
 defense system and ATP release of rabbit erythrocytes: in vitro study. Asian J Anim 471
 Sci Vet Adv 12,1-9. 472
11. Alagawany M., M.R. Farag, Mohamed Ezzat Abd El-Hack, Elisabetta Casalino, 473
 Vincenzo Tufarelli, Maryam Sayab and Kuldeep Dhama 2016. Assessment of cyadox 474
 effects on the antioxidant defense system and hemolysis of isolated rabbit 475
 erythrocytes. Int J Pharmacol 13, 183-190. 476
12. Pagano M. and Faggio C. The use of erythrocyte fragility to assess xenobiotic 477
 cytotoxicity. cell biochemistry and function. Cell Biochem Funct. 2015; DOI: 478
 10.1002/cbf.3135 479
13. Shiva, S., Subramanyam, M.V., Vani, R., Asha, D., 2007. In vitro models of 480
 oxidative stress in rat erythrocytes: effect of antioxidant supplements. Toxicol. In 481
 Vitro 21, 1355–1364. 482
14. Okamoto, K., Maruyama, T., kaji, Y., Harada, M., Mawatari, S., Fujino,T., Uyesaka, 483
 N., 2004. Verapamil prevents impairment in filterability of human erythrocytes 484
 exposed to oxidative stress. Japanese Journal of Physiology 54, 39–46. 485
15. Asha, D.S., Subramanyam, M.V.V., Vani, R., Jeevaratnam, K., Adaptations of the 486
 antioxidant system in erythrocytes of trained adult rats: Impact of intermittent 487
 hypobaric-hypoxia at two altitudes. Comparative Biochemistry and Physiology. Part 488
 C 2005; 140: 59–67. 489

16. Lang, F., Gulbins, E., Lang, P.A., Ceramide in suicidal death of erythrocytes. *Cell* 490
Physiol Biochem 2010; 26: 21–28. 491
17. Mandal, S., Mukherjee, S., Chowdhury, K. D., Sarkar, A., Basu, K., Paul, S., Karmakar, 492
D., Chatterjee, M., Biswas, T., Chandr, G., Khan, S., Sen, G. 2012 S-allyl cysteine in 493
combination with clotrimazole downregulates Fas induced apoptotic events in 494
erythrocytes of mice exposed to lead. *Biochim Biophys Acta* 1820: 9–23. 495
18. Lang E, Qadri SM, Lang F. 2012. Killing me softly – suicidal erythrocyte death. *Int J* 496
Biochem Cell Biol; 44: 1236–1243. 497
19. Paprika MV and Sharma BB. Effect of oral administration of herbicide diclofop on 498
some hematological parameters in mouse. *J. Cell Tissue Res.* 2003; 3: 12-17. 499
20. Nikinmaa M. How does environmental pollution affect red cell in fish? *Aquat* 500
Toxicol 1992; 22: 227-238. 501
21. Fenech, M., Kirsch-Volders, M., Natarajan, A.T., Surrallés, J., Crott, J.W., Parry, J., 502
Norppa, H., Eastmond, D.A., Tucker, J.D., Thomas, P., 2011. Molecular mechanisms 503
of micronucleus, nucleoplasmic bridge and nuclear bud formation in mammalian and 504
human cells. *Mutagenesis* 26, 125–132. 505
22. Harabawy, A.S.A., Mosleh, Y.Y.I., 2014. The role of vitamins A, C, E and selenium 506
as antioxidants against genotoxicity and cytotoxicity of cadmium, copper, lead and 507
zinc on erythrocytes of Nile tilapia, *Oreochromis niloticus*. *Ecotoxicol. Environ. Saf.* 508
104, 28–35. 509

23. Ochiai, K., Jin, K., Itakura, C., Goryo, M., Yamashita, K., Mizuno, N., Fujinaga, T., 510
Tsuzuki, T., 1992. Pathological study of lead poisoning in whooper swans *Cygnus* 511
cygnus in Japan. *Avian Diseases* 36, 313–323. 512
24. Gurer H, Ozgunes H, Neal R, Spitz D R, Erçal N. 1998. Antioxidant effects of N- 513
acetylcysteine and succimer in red blood cells from lead-exposed rats, *Toxicology*; 514
128,3:181–189 515
25. Hirag T., Kei Ohyama, Ayano Hashigay, Tomoya Ishikawa, Wakana Muramoto, 516
Hiroshi Kitagaw, Naoharu Mizun, Hiroki Teraoka 2008 Lead exposure induces 517
pycnosis and enucleation of peripheral erythrocytes in the domestic fowl. *The* 518
Veterinary Journal 178 2008 109–114. 519
26. Sharma R., Khushbu Panwar and Sheetal Mogra. 2013 Alterations in developing 520
RBCs after prenatal and postnatal exposure to lead acetate and vitamins. *International* 521
Journal of Pharmaceutical Sciences and Research 4 8 : 3214-3224. 522
27. Vittori D, Nesse A, Perez G, Garbossa G. 1999. Morphologic and functional 523
alterations of erythroid cells induced by long-term ingestion of aluminium. *J* 524
Inorganic Biochem 76: 113-120.23. 525
28. Comelekoglu U, Mazmanci B, Arpacı A. 2000 Investigation of the liver functions in 526
agricultural workers chronically exposed to pesticides. *Turk J Biol* 24: 461-466. 527
29. Zeni C, Bovolenta MR, Stagni A. 2002. Occurrence of echinocytosis in circulating 528
RBC of black bullhead, *Ictalurus melas* Rafinesque , following exposure to an 529
anionic detergent at sublethal concentrations. *Aquat Toxicol*; 57: 217-224. 530

30. Koc ND, Muslu MN, Sesal C, Kayhan FE. 2008. Histopathological effects of malathion and endosulfan on blood cells of Wistar albino rats *Rattus norvegicus* J Appl Biol Sci; 2: 105-108.
31. Suwalsky M, Vargas P, Avello M, Villena F, Sotomayor CF. 2008. Human erythrocytes are affected in vitro by flavonoids of *Aristotelia chilensis* Maqui leaves. International J Pharmaceutics; 363: 85-9033.
32. Tripathi S, Srivastav A. 2010. Alterations in the profile of blood cells of wistar rats induced by long-term ingestion of chlorpyrifos. International Journal of Pharma and Bio Sciences 1, B-322 ref.39
33. Yang, H.L., Chen, S.C., Chang, N.W., Chang, J.M., Lee, M., Tsai, P.C., Fu, H.H., Kao, W.W., Chiang, H.C., Wang, H.H., Hseu, Y.C. 2006 Protection from oxidative damage using *Bidenspilosa* extracts in normal human erythrocytes. Food Chem Toxicol 44: 1513–1521.
34. Agrawal D., P. Sultana, [1993] Biochemical and structural alterations in rat erythrocytes due to hexachlorocyclohexane exposure. Food Chem Toxicol., 31 443–448.
35. Jindal R and S Verma 2015 In vivo genotoxicity and cytotoxicity assessment of cadmium chloride in peripheral erythrocytes of *Labeo rohita* Hamilton Ecotoxicology and Environmental Safety 118 2015 1–10
36. Kumar G, Karthik L, Rao KVB. 2011. Hemolytic activity of Indian medicinal plants towards human erythrocytes: an in vitro study. Elixir Appl Botany; 40: 5534–5537.

37. Orsine JVC, Costa R, Silva R, Santos M, Novaes M. The acute cytotoxicity and lethal concentration LC50 of *Agaricus sylvaticus* through hemolytic activity on human erythrocyte. *Int J Nutr Met* 2012; 4:19–23.
38. Budan A., Tessier N., Saunier M., 2013 Effect of several saponin containing plant extracts on rumen fermentation in vitro, *Tetrahymena pyriformis* and sheep erythrocytes. *J Food Agric Environ* 11, 576–582.
39. Faggio C, Morabito M, Armeli Minicante S, Lo Piano G, Pagano M, Genovese G. 2015; Potential use of polysaccharides from the brown alga *Undaria pinnatifida* as anticoagulant. *Braz Arch Biol Technol* 58, 798–804.
40. Faggio C, Pagano M, Morabito M, Minicante Armeli S, Arfuso F, Genovese G. 2014. In vitro assessment of the effect of *Undaria pinnatifida* extracts on erythrocytes membrane and blood coagulation parameters of *Equus caballus*. *J Coast Life Med*; 2, 614–616.
41. Kolanjiappana K, Manoharana S, Kayalvizhib M. 2002. Measurement of erythrocyte lipids, lipid peroxidation, antioxidants and osmotic fragility in cervical cancer patients. *Clin Chim Acta*. 326:143–149.
42. Chikezie PC Osmotic fragility index of HbAA red blood cells in the presence of aqueous extracts of three medicinal plants *Aframomum melegueta*, *Garina kola*, and *Cymbopogon Citracus* *Global J. Pure Applied Sci.* 2007; 13, 496-499
43. Mansour S.A., Abdel-Tawab H. Mossa. [2009] Lipid peroxidation and oxidative stress in rat erythrocytes induced by chlorpyrifos and the protective effect of zinc . *Pesticide Biochemistry and Physiology* 93 34–39

44. Agalakova N., Gennadii P. Gusev. 2011 Fluoride-induced death of rat erythrocytes 574
in vitro. *Toxicology in Vitro*, 25 1609–1618. 575
45. Wafa T, Nakbi Amela, Chargui Issama, Cheraief Imed a, Miled Abdelhedib , 576
Hammami Mohamed 2011 Subacute effects of 2,4-dichlorophenoxyacetic herbicide 577
on antioxidant defensesystem and lipid peroxidation in rat erythrocytesq. *Pesticide* 578
Biochemistry and Physiology 99 2011 256–264 579
46. Rendell, M., Luu, T., Quinlan, E., Knox, S., Fox, M., Kelly, S.,Kahler, K. 1992 Red 580
cell filterability determined using the cell transit timeanalyzer CTTA : effects of ATP 581
depletion and change in calciumconcentration. *Biochim Biophys Acta* 1133: 293– 582
300. 583
47. Meyers, T.R., Hendricks, J.D., 1985. Histopathology In: Rand, G.M., Petrocelli, S.R. 584
Eds. , *Fundamentals of Aquatic Toxicity: Methods and Applications*. Hemi sphere 585
Publishing Corporation, Washington, USA, pp. 283–331. 586
48. Mekkawy, I.A., Mahmoud, U.M., Sayed, A.E.-D.H., 2011. Effects of 4-nonylphenol 587
on blood cells of the African catfish *Clarias gariepinus* Burchell, 1822 *Tissue Cell* 588
43, 223–229. 589
49. Teijon C, Delsocorro JM, Martin JA, Lozano M, Bernardo V, Blaco D. 2000 Lead 590
accumulation in rats at non acute doses and short periods of time. Hepatic, renal and 591
hematological effects. *Ecotoxico Environm Restor* 3, 36 –41. 592
50. Farag MR, Mahmoud Alagawany & Vincenzo Tufarelli 2017 In vitro antioxidant 593
activities of resveratrol, cinnamaldehyde and their synergistic effect against cyadox- 594

- induced cytotoxicity in rabbit erythrocytes, Drug and Chemical Toxicology, 40 2 595
:196-205. 596
51. Sikora, J.; Sergei N. O., Furuya, K. and Grygorczyk, R. 2014. Hemolysis is a 597
primary ATP-release mechanism in human erythrocytes. Blood, 124: 2150-2157. 598
52. Adams, H. 1963 In: Bergmeyer, H.U. Ed. Methods Enzymatic Analysis. Academic 599
Press, New York, pp. 539–543.43. 600
53. Yoshino M, Chikashi Y, Keiko M, 1992 Stabilization of the adenylate energy charge 601
in erythrocytes of rats and humans at high altitude hypoxia. Comp Biochem Physiol 602
A Comp Physiol 101:65–68. 603
54. Atkinson DE, Walton GM. 1967 Adenosine triphosphate conservation in metabolic 604
regulation. Rat liver citrate cleavage enzyme. J Biol Chem 242, 3239–3241. 605
55. Kim S, Ryu DY. 2013. Silver nanoparticle-induced oxidative stress, genotoxicity and 606
apoptosis in cultured cells and animal tissues. J Appl Toxicol 33:78–89. 607
56. Bolognesi, C., Hayashi, M., 2011. Micronucleus assay in aquatic animals. 608
Mutagenesis 26, 205–213. 609
57. Farag MR., Mahmoud M. Alagawany and Kuldeep Dhama. 2014. Antidotal effect of 610
turmeric Curcuma longa against endosulfan-induced cytogenotoxicity and 611
immunotoxicity in broiler chicks. International journal of Pharmacology, , 10 8 : 612
429-439. 613
58. Tan, D., Li, L., Wang, S., Wei, B., Zhang, X., Sun, B., Ji, S., 2013. The cytogenetic 614
effectsof acrylamide on Carassius auratus periperial blood cells. Food Chem. Toxicol. 615
62, 318–322. 616

59. Vincent-Hubert, F., Arini, A., Gourlay-France, C., 2011. Early genotoxic effects in
gill cells and haemocytes of *Dreissena polymorpha* exposed to cadmium, B a P and a
combination of B a P and Cd. *Mutat. Res.* 723, 26–35.
60. Holden, E.H., Jenness, B.M., Debbie, S.A., 1997. Direct comparison of mouse and rat
bone marrow and blood as target tissues in the micronucleus assay. *Mutagen
Research/Genetic Toxicology and the Environmental Mutagen* 391, 87–90.
61. Singh, N.P., McCoy, M.T., Tice, R.R., Schneider, E.L., 1988. A simple technique for
quantitation of low levels of DNA damage in individual cells. *Exp. Cell Res.*
175,184–191.
62. Weil, M., Jacobson, M.D., Raff, M.C., 1998. Are caspases involved in the death
of cells with a transcriptionally inactive nucleus ?Sperm and chicken erythrocytes
Journal of Cell Science 111, 2707–2715.
63. Wang Y.-C, Chaung R.-H, Tung L. 2004 . Comparison of the cytotoxicity induced
by different exposure to sodium arsenite in two fish cell lines. *Aquatic
Toxicology*, 69, 67-79.
64. Orrenius S, Nicotera P, Zhivotovsky B. 2011. Cell death mechanisms and their
implications in toxicology. *Toxicol Sci.*, 119:3–19.
65. Webb, S.J., Harrison, D.J., Wyllie, A.H., 1997. Apoptosis: an overview of the process
and its relevance in disease. In: Kaufmann, S.H. Ed, .Apoptosis: Pharmacological
Implications and Therapeutic Opportunities, *Advances in Pharmacology*, vol. 41.
Academic Press, San Diego,pp. 1–34.

66. Jacobi J, Lang E, Bissinger R, 2014. Stimulation of erythrocyte cell membrane 638
scrambling by mitotane. *Cell Physiol Biochem*; 33: 1516–1516. 639
67. Simpson, C.F., Kling, J.M., 1967. The mechanism of denucleation in circulating 640
erythroblasts. *Journal Cell Biology* 35, 237–245. 641
68. Kempe, D.S., Lang, P.A., Eisele, K., Klarl, B.A., Wieder, T., Huber, S.M Duranton, 642
C., Lang, F., 2005. Stimulation of erythrocyte phosphatidylserine exposure by lead 643
ions. *American Journal of Physiology Cell Physiology* 288, C396–C402. 644
69. Berg, C.P., Engels, I.H., Rothbart, A., Lauber, K., Renz, A., Schlosser, S.F., Schulze- 645
Osthoff, K., Wesselborg, S. 2001 Human mature red blood cells express caspase-3 646
and caspase-8, but are devoid of mitochondrial regulators of apoptosis. *Cell Death* 647
Differ 8, 1197-1206. 648
70. Bejarano, I., Lozano, G.M., Ortiz, A., Garcia, J.F., Paredes, S.D., Rodriguez, A.B., 649
Pariente, J.A., 2008. Caspase 3 activation in human spermatozoa in response to 650
hydrogen peroxide and progesterone. *Fertil. Steril.* 90, 1340–1347. 651
71. Biswas D, Gargi Sen, Tuli Biswas 2010 Reduced cellular redox status induces 4- 652
hydroxynonenal-mediated caspase 3 activation leading to erythrocyte death during 653
chronic arsenic exposure in rats. *Toxicology and Applied Pharmacology* 244, 315– 654
327 655
72. Mukherjee S., Kaustav Dutta Chowdhury, Avik Sarkara, Kankana Basu, Soumosish 656
Paulb, Debasish Karmakarb, Mahasweta Chatterjee , Puli Biswas Gobinda, Chandra 657
Sadhukhan, Gargi Sen 2012 S-allyl cysteine in combination with clotrimazole 658

- downregulates Fas induced apoptotic events in erythrocytes of mice exposed to lead. 659
 Samir Mandal *Biochimica et Biophysica Acta* 1820 9–23. 660
73. Senapati S.K., S. Dey, S.K. Dwivedi, D. Swarup, 2001 Effect of garlic *Allium* 661
sativum L. extract on tissue lead level in rats, *J. Ethnopharmacol.* 76 229–232. 662
74. Cremesti A., F.O. Paris, H. Grassme, N. Holler, J. Tschopp, Z. Fuksi, E. Gulbins, R. 663
 Kolesnick, 2001 Ceramide enables Fas to cap and kill. *J. Biol. Chem.* 276, 23954– 664
 23961. 665
75. Sen G., D. Biswas, M. Ray, T. Biswas, 2007 Albumin–quercetin combination offers 666
 a therapeutic advantage in the prevention of reduced survival of erythrocytes in 667
 visceral leishmaniasis, *Blood Cells Mol. Dis.* 39 245–254. 668
76. Chowdhury K.D., G. Sen, A. Sarkar, T. Biswas, 2011 Role of endothelial 669
 dysfunction in modulating the plasma redox homeostasis in visceral leishmaniasis, 670
Biochim. Bio-phys. Acta 1810, 652–665. 671
77. Wolff D., X. Cecchi, A. Spalvins, M. Canessa, 1988 Charybdotoxin blocks with 672
 high affinity the Ca-activated K⁺channel of Hb A and Hb S red cells: individual 673
 differences in the number of channels, *J. Membr. Biol.* 106, 243–252. 674
78. Petrache I., V. Natarajan, L. Zhen, T.R. Medler, A.T. Richter, C. Cho, W.C. Hubbard, 675
 E.V. Berdyshev, R.M. Tudor, 2005 Ceramide up regulation causes pulmonary cell 676
 apoptosis and emphysema-like disease in mice, *Nat. Med.* 11 491–498. 677
79. Muppidi J.R., R.M. Siegel, 2004 Ligand-independent redistribution of Fas CD95 678
 into lipid rafts mediates clonotypic T cell death, *Nat. Immunol.* 5 182–189. 679

80. Ahmad, S., Beg, Z.H. 2013 Alleviation of plasma, erythrocyte and liver lipidemic- 680
oxidative stress by by thymoquinone and limonene in atherogenic suspension fed rats. 681
J Functional Foods 5, 251-259. 682
81. Nagababu, E., Rifkind, J.M., 2000. Reaction of hydrogen peroxide with 683
ferrylhemoglobin: superoxide production and heme degradation. Biochemistry 39, 684
12503–12511. 685
82. Mohanty J.G., Enika Nagababu, and Joseph M. Rifkind 2014. Red blood cell 686
oxidative stress impairs oxygen delivery and induces red blood cell aging. Frontiers 687
in Physiology .; 5: 84. 688
83. Rossen van, M.E.E., Sluiter, W., Bonthuis, F., Jeekel, H., Marquet, R.L., Van Eijck, 689
C.H.J .2000. Scavenging of reactive oxygen species leads to diminished peritoneal 690
tumor recurrence. Cancer Res. 60, 5625–5629. 691
84. Çimen, M., 2008. Free radical metabolism in human erythrocytes. Clinica 692
Chimica Acta 390, 1–11. 693
85. Goth, L., Vitai, M., 2003. The effects of hydrogen peroxide promoted by homocys- 694
teine and inherited catalase deficiency on human hypocatalasemic patients. Free 695
Radic. Biol. Med. 35, 882–888. 696
86. Yamamoto, Y., Niki, E., Eguchi, J., Kamiya, Y., Shimasaki, H., 1985. Oxidation of 697
biological membranes and its inhibition. Free radical chain oxidation of erythrocyte 698
ghost membranes by oxygen. Biochimica et Biophysica Acta 819, 29–36. 699

87. Carini, M., Aldini, G., Facino, R.M., 2004. Mass spectrometry for detection of 4- 700
hydroxy-trans-2-nonenal HNE adducts with peptides and proteins. Mass Spectrom. 701
Rev. 23, 281–305. 702
88. Hebbel R.P., A. Leung, N. Mohandas, 1990 Oxidation-induced changes in 703
microrheologic properties of the red blood cell membrane, Blood 76 1015–1020. 704
89. Sugihara, T., Rawicz, W.E.A., Hebbel, R.P. 1991 Lipid hydroperoxides permit 705
deformation-dependent leak of monovalent cation from erythrocytes. Blood 77: 706
2757–2763. 707
90. Sadowska-Wodaa I., Barbara Sychtaa, Marta Rachelb, Edyta Bieszczad-Bedrejcuka. 708
2010 Protective effect of desloratadine against oxidative stress in human erythrocytes 709
in vitro. Environmental Toxicology and Pharmacology 30 141–146 710
91. Reddy C.S. Shiva Shankar a ,M.V.V. Subramanyam b ,R. Vania ,S. Asha Devia 2007 711
In vitro models of oxidative stress in rat erythrocytes: Effect of antioxidant 712
supplements. Toxicology in Vitro 21 1355–1364 713
92. El-Demerdash F.M. 2007 Lambda-cyhalothrin-induced changes in oxidative stress 714
biomarkers in rabbit erythrocytes and alleviation effect of some antioxidants. 715
Toxicology in Vitro 21, 392–397. 716
93. Lukivskaya O.Y., A.A. Maskevish, V.U. Buko, 2001 Effect of ursodeoxycholic acid 717
on prostaglandin metabolism and microsomal membranes in alcoholic fatty liver. 718
Alcohol 25 99–105. 719

94. Dodge, J.T., Mitchell, C., Hanahan, D.J., 1963. The preparation and chemical 720
characteristics of hemoglobin free ghosts of human erythrocytes. Arch. Biochem. 721
Biophys. 100, 119–130. 722
95. Giacometti J., A. Milošević, C[˘]. Milin, 2002 Gas chromatography determination of 723
fatty acids contained in different classes after their separation by solid-phase 724
extraction, J. Chromatogr. A 976 47–54. 725
96. Laemmli, U.K., 1970. Cleavage of structural proteins during the assembly of the head 726
of bacteriophage T4. Nature 227 5259 , 680–685. 727
97. Wolff, S.P., 1994. Ferrous ion oxidation in presence of ferric ion indicator xylenol 728
orange for measurement of hydroperoxides. Methods Enzymol 233 Part C , 182–189. 729
98. Biswas, D., Banerjee, M., Sen, G., Das, J.K., Banerjee, A., Sau, T.J., Pandit, S., Giri, 730
A.K., Biswas, T., 2008. Mechanism of erythrocyte death in human population exposed 731
to arsenic through drinking water. Toxicol. Appl. Pharmacol. 230, 57–66. 732
99. Lopez-Revuelta, A., Sanchez-Gallego, J.I., Hernandez-Hernandez, A., Sanchez- 733
Yague, J., Llanillo, M., 2005. Increase in vulnerability to oxidative damage in 734
cholesterol-modified erythrocytes exposed to t-BuOOH. Biochim. Biophys. Acta 735
1734, 74–85. 736
100. Zhao, H., Joseph, J., Fales, H.M., Sokoloski, E.A., Levine, R.L., Vasquez-Vivar, J., 737
Kalyanaraman, B., 2005 Detection and characterization of the product of 738
hydroethidium and intracellular superoxide by HPLC and limitations of fluorescence. 739
Proc. Natl. Acad. Sci. 102, 5727–5732. 740

101. Setsukinai, K., Urano, Y., Kakinuma, K., Majima, H.J., Nagano, T., 2003. 741
Development of novel fluorescence probes that can reliably detect reactive oxygen 742
species and distinguish specific species. *J. Biol. Chem.* 278 5 , 3170–3175. 743
102. Qian, H., Chen, W., Li, J., Wang, J., Zhou, Z., Liu, W., Fu, Z., 2009. The 744
effect of exogenous nitric oxide on alleviating herbicide damage in *Chlorella* 745
vulgaris. *Aquatic Toxicology* 92, 250–257.103. 746
103. Ellman, G.L. 1959 Tissue sulfhydryl groups. *Arch Biochem Biophys* 82: 70–77. 747
104. Ohkawa, H., Ohishi, N., Yagi, K., 1979. Assay for lipid peroxides in animal tissues 748
by thiobarbituric acid reaction. *Analytical Biochemistry* 95, 351–358. 749
105. Yagi, K., 1984. Assay for blood plasma or serum. *Methods in Enzymology* 105, 750
328–331. 751
106. Bartosz, G. 2004 *Druga twarz tlenu. Wolnorodnikiw przyrodzie*, Wyd.Nauk. PWN, 752
Warszawa. 753
107. Tedesco, S., Doyle, H., Blasco, J., Redmond, G., Sheehan, D., 2010. Oxidative stress 754
and toxicity of gold nanoparticles in *Mytilus edulis*. *Aquatic Toxicology* 100, 178– 755
186. 756
108. Uchida, K., Stadtman, E.R. 1993. Covalent attachment of 4-hydroxy- nonenal to 757
glyceraldehyde-3-phosphate dehydrogenase. *J Biol Chem* 268, 6388–6393. 758
109. Jiang, W.D., Wu, P., Kuang, S.Y., Liu, Y., Jiang, J., Hu, K., Li, S.H., Tang, 759
L., Feng, L., Zhou, X.Q., 2011. Myo-inositol prevents copper-induced oxidative 760
damage and changes in antioxidant capacity in various organs and the 761

- enterocytes of juvenile Jian carp *Cyprinus carpio* var. Jian *Aquatic Toxicology* 105, 543–551.
110. Arguelles, S., Machado, A., Ayala, A., 2009 Adduct formation of 4-hydroxynonenal and malondialdehyde with elongation factor-2 in vitro and in vivo. *Free Radic. Biol. Med.* 47, 324–330.
111. Tietze, F., 1969. Enzymatic method for quantitative determination of nanogram amounts of total and oxidized glutathione: applications to mammalian blood and other tissues. *Anal. Biochem.* 27 3, 502–522.
112. Dise CA, Goodman DBP 1986. t-Butyl hydroperoxide alters fatty acid incorporation into erythrocyte membrane phospholipid. *Biochim Biophys Acta* 859:69–78
113. Beutler, E., 1971. In: Beutler, E. Ed., *Red Cell Metabolism. A Manual of Biochemical Methods*. Grune and Stratton Inc., New York.
114. Paglia D.E., W.N.J. Valentine, 1967 Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase, *Lab. Clin. Med.* 70 158–169.
115. Habig, W.H., Pabst, M.J., Jakoby, W.B., 1974. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *Journal of Biological Chemistry* 249, 7130–7139.
116. Marklund, S.L., Marklund, G., Involvement of the superoxide anion radical in the autoxidation of pirogallol and a convenient assay for superoxide dismutase. *Eur. J. Biochem.* 1974; 47: 469–474.

117. Spitz, D.R., Oberley, L.W., An assay for superoxide dismutase activity in 783
mammalian tissue homogenates. *Analytical Biochemistry* 1989; 179: 8–18. 784
118. Aebi, H.E. Catalase in vitro. *Method Enzymol* 1984; 105:121–126. 785
119. Simpson, C.F., Damron, B.L., Harms, R.H., 1970. Abnormalities of erythrocytes 786
and renal tubules of chickens poisoned with lead. *American Journal of Veterinary* 787
Research 31, 515–523. 788
120. Monteiroa V., D.G.S.M. Cavalcantea, M.B.F.A. Vilélaa, S.H. Sofiab, C.B.R. 789
Martineza 2011. In vivo and in vitro exposures for the evaluation of the 790
genotoxic effects of lead on the Neotropical freshwater fish *Prochilodus* 791
lineatus. *Aquatic Toxicology* 104 291–298. 792
121. A. Rendo ´n-Ramírez, J. Cerbo ´n-Solo ´rzano, M. Maldonado-Vega, M.A. 793
Quintanar-Escorza, J.V. Caldero ´n-Salinas, Vitamin-E reduces the oxidative damage 794
on δ -aminolevulinic dehydratase induced by lead intoxication in rat erythrocytes. 795
Toxicology in Vitro 21 (2007) 1121–1126 796
122. Mousa, H.M., Al-Qarawi, A.A., Ali, B.H., Abdel-Rahman, H.A., ElMougy, S.A., 797
2002. Effect of lead exposure on the erythrocytic antioxidant levels in goats. *Journal* 798
of Veterinary Medicine, Series A Physiology, Pathology, Clinical Medicine 49, 531– 799
534. 800
123. L.E. Eriksson, H. Beving, Calcium- and lead-activated morphological changes in 801
human erythrocytes: a spin label study of the cytoplasm, *Arch. Biochem. Biophys.* 802
303 (1993) 296–301. 803
124. M. Shields, R. Grygorczyk, G.F. Fuhrmann, W. Schwarz, H. Passow, Lead-induced 804

- activation and inhibition of potassium-selective channels in the human red blood cell, 805
 Biochim. Biophys. Acta. 815 (1985) 223–232. 806
125. Ferraro M.V.M., Fenocchio A.S., Mantovani M.S., Ribeiro C.O. and Cestari M.M. 807
 (2004) Mutagenic effects of tributyltin and in-organic lead (Pb II) on the fish *H.* 808
Malabaricus as evaluated using the comet assay and piscine micronucleus and chro- 809
 mosome aberration tests. *Genet Mol Biol* 27:103-107. 810
126. S. Kousar, M. Javed, 2015. Studies on induction of nuclear abnormalities in 811
 peripheral blood erythrocytes of fish exposed to copper. *Turkish Journal of Fisheries* 812
 and Aquatic Sciences 15: 879-886 (2015). 813
127. Çavas T., Garanko N.N., Arkhipchuk V.V. (2005) Induction of micronuclei and 814
 binuclei in blood, gill and liver cells of fishes subchronically exposed to cadmium 815
 chlorid and copper sulphate. *Food Chem Toxicol* 43:569-574. 816
128. Harabawy A.S.A., Mosleh Y.Y.I., 2014. The role of vitamins A, C, E and selenium 817
 as antioxidants against genotoxicity and cytotoxicity of cadmium, copper, lead and 818
 zinc on erythrocytes of Nile tilapia, *Oreochromis niloticus*. *Ecotoxicol. En-viron. Saf.* 819
 104, 28–35. 820
129. Nepomuceno, J.C., Ferrari, I., Spano, M.A., Centeno, A.J., 1997. Detection of 821
 micronuclei in peripheral erythrocytes of *Cyprinus carpio* exposed to metallic 822
 mercury. *Environ. Mol. Mutagen.* 30, 293–297. 823
130. Guilherme, S., Valega, M., Pereira, M.E., Santos, M.A., Pacheco, M., 2008. 824
 Erythrocytic nuclear abnormalities in wild and caged fish (*Liza aurata*) along an 825

- environmental mercury contamination gradient. *Ecotoxicol. Environ. Saf.* 70, 411- 826
421. 827
131. L. Stana, A. Trif, L.G. Stana, S. Petrovici, C. Gravila, comparative study of the 828
potassium dichromate effect on the osmotic resistance of rat erythrocyte membrane. 829
Animal Science and Biotechnologies 43 (1), 425-428. 830
132. D. Biswas, G. Sen, A. Sarkar, T. Biswas, Atorvastatin acts synergistically with N- 831
acetyl cysteine to provide therapeutic advantage against Fas-activated erythrocyte 832
apoptosis during chronic arsenic exposure in rats, *Toxicol. Appl. Pharmacol.* 250 833
(2011) 39–53. 834
133. P. F. Zatta, D. Cervellin, P. Zambenedetti. Effects of the Aluminium Speciation on 835
theMorphology of Rabbit Erythrocytes: aToxicological Model Toxicology in Vitro, 836
12 (1998) 287±293 837
134. Abou-Seif, M.A., 1998. Oxidative stress of vanadium-mediated oxygen free radical 838
generation stimulated by aluminium on human erythrocytes. *Ann. Clin. Biochem.* 35, 839
254–260. 840
135. Rice-evans, C., Baysal, E., 1987. Iron-mediated oxidative stress in 841
erythrocytes. *Bio-chemical Journal* 244, 191–196. 842
136. M. Sinha, P. Manna, P.C. Sil, A 43 k D protein from the herb, *Cajanus indicus* L., 843
protects against fluoride induced oxidative stress in mice erythrocytes, 844
Pathophysiology 14 (2007) 47–54. 845

137. A. Scibior, Halina Zaporowska, Effects of combined vanadate and magnesium
treatment on erythrocyte antioxidant defence system in rats. *Environmental
Toxicology and Pharmacology* 30 (2010) 153–161.
138. Heller, K.B., Jahn, B., Deuticke, B., 1987. Peroxidative membrane damage in
human erythrocytes induced by a concerted action of iodoacetate, vanadate and
ferricyanide. *Biochim. Biophys. Acta* 901, 67–77.
139. Cavas, T. and Ergene-Gozukara, S. (a) (2003) Evaluation of the genotoxic potential
of lambda-cyhalothrin using nuclear and nucleolar biomarkers on fish cells. *Mutat.
Res.*, 534, 93–99.
140. H. Fetoui, E.M. Garoui, F. Makni-Ayadi, N. Zeghal, Oxidative stress induced by
lambda-cyhalothrin (LTC) in rat erythrocytes and brain: attenuation by vitamin C,
Environ. Toxicol. Pharmacol. 26 (2008) 225–231.
141. Fulya Dilek Gökalp Muranli, Utku Güner, Induction of micronuclei and nuclear
abnormalities in erythrocytes of mosquito fish (*Gambusia affinis*) following exposure
to the pyrethroid insecticide lambda-cyhalothrin *Mutation Research* 726 (2011) 104–
108.
142. Campana, M.A., Panzeri, A.M., Moreno, V.J. and Dulout, F.N., 1999. Genotoxic
evaluation of the pyrethroid lambda-cyhalothrin using the micronucleus test in
erythrocytes of the fish *Cheirodon interruptus*. *Mutat. Res.*, 438: 155-61.
143. Gul-e-Zehra Naqvi, Nafisa Shoaib, Aisha Majid Ali, genotoxic potential of
pesticides in the peripheral blood erythrocytes of fish (*Oreochromis mossambicus*).
Pakistan J. Zool., vol. 48(6), pp. 1643-1648, 2016.

144. M. Kale, N. Rathore, S. John, D. Bhatnagar, Lipid peroxidative damage on 868
pyrethroid exposure and alterations in antioxidant status in rat RBCs: a possible 869
involvement of reactive oxygen species, *Toxicol. Lett.* 105 (1999) 197–205. 870
145. B. Bukowska, B. Rychlik, A. Krokosz, J. Michalowicz, Phenoxyherbicides induce 871
production of free radicals in human erythrocytes: oxidation of dichlorofluorescein 872
and dihydrorhodamine 123 by 2,4-D-Na and MCPA-Na, *Food Chem. Toxicol.* 46 873
(2008) 359–367. 874
146. B. Bukowska, Jaromir Michalowicz, Aneta Wojtaszek, Agnieszka Marczak², 875
Comparison of the effect of phenoxyherbicides on human erythrocyte membrane (in 876
vitro). *Biologia*, 66(2): 379—385, 2011. 877
147. Ateeq B, Farah MA, Ali MN, Ahmad W (2002) Induction of micronuclei and 878
erythrocyte alterations in the catfish *Clarias batrachus* by 2,4-dichlorophenoxyacetic 879
acid and butachlor. *Mutat Res* 518:135-144. 880
148. Ateeq, B., Abul Farah, M. and Ahmad, W. (2005) Detection of DNA damage by 881
alkaline single cell gel electrophoresis in 2,4-dichlorophenoxy-acetic-acid- and 882
butachlor-exposed erythrocytes of *Clarias batrachus*. *Ecotoxicol. Environ. Saf.*, 62, 883
348–354 884
149. Ali D, Nagpure NS, Kumar S, Kumar R, Kushwaha B., Genotoxicity assessment of 885
acute exposure of chlorpyrifos to freshwater fish *Channa punctatus* (Bloch) using 886
micronucleus assay and alkaline single-cell gel electrophoresis. *Chemosphere*. 2008 887
71(10):1823-31. 888

150. A.C.D. Bainy, A.C.M. Arisi, L.A. Azzalis, K. Simizu, S.B.M. Barios, L.A. Videla, 889
V.B.C. Jungueira, Differential effects of short-term lindane administration on 890
parameters related to oxidative stress in rat liver and erythrocytes, J. Biochem. 891
Toxicol. 8 (1993) 187–194. 892
151. Prasanthi, K., Muralidhara, Rajini, P.S., 2005. Morphological and biochemical 893
perturbations in rat erythrocytes following *in vitro* exposure to Fenvalerate and its 894
metabolite. Toxicology in Vitro 19, 449–456. 895
152. Tolga C avas, Serpil Ko  nen., Detection of cytogenetic and DNA damage in 896
peripheral erythrocytes of goldfish (*Carassius auratus*) exposed to a glyphosate 897
formulation using the micronucleus test and the comet assay Mutagenesis, 22: 4 263– 898
268, 2007. 899
153. Clements, C., Ralph, S., Pertas, M. (1997) Genotoxicity of select herbicides in Rana 900
catesbeiana tadpoles using the alkaline single-cell gelDNA electrophoresis (comet) 901
assay. Environ. Mol. Mutagen., 29, 277–288. 902
154. Pallavi Srivastava and Ajay Singh 2015.Evidence of micronuclei in fish blood as a 903
biomarker of genotoxicity due to surface run off agriculture fungicide 904
(Propiconazole) Journal of toxicology and environmental health sciences, 7 (1): 4-8 , 905
2015. 906
155. Cavaş T1, Ergene-G z kara S., (b) Micronuclei, nuclear lesions and interphase 907
silver-stained nucleolar organizer regions (AgNORs) as cyto-genotoxicity indicators 908
in Oreochromis niloticus exposed to textile mill effluent. Mutat Res. 2003 8, 538(1- 909
2):81-91. 910

156. Çavas T and Ergene-Gözükara S (2005) Induction of micronuclei and nuclear
abnormalities in *Oreochromis niloticus* following exposure to petroleum refinery and
chromium processing plant effluents. *Aquat Toxicol* 74:264-271.
157. Da Silva Souza, T., Fontanetti, C. S. (2006) Micronucleus test and observation of
nuclear alterations in erythrocytes of Nile tilapia exposed to waters affected by
refinery effluent. *Mutat. Res.*, 605, 87–93.
- 158- Belinda C. Gómez-Meda, Ana L. Zamora-Perez, Jaime Luna-Aguirre, Andrés
González-Rodríguez, M. Luisa Ramos-Ibarra, Olivia Torres-Bugarín, Cecilia M.
Batista-González, Guillermo M. Zúñiga-González, Nuclear abnormalities in
erythrocytes of parrots (*Aratinga canicularis*) related to genotoxic damage. *Avian
Pathology* (2006) 35(3), 206-210.
159. Winter M.J., Ellis, L.C.J., Hutchinson, T.H., 2007. Formation of micronuclei in
erythrocytes of the fathead minnow (*Pimephales promelas*) after acute treatment
with mitomycin C or cyclophosphamide. *Mutat. Res.* 629, 89–99.
160. Toni P. Galindo and Lilia M. Moreira. Evaluation of genotoxicity using the
micronucleus assay and nuclear abnormalities in the tropical sea fish *Bathygobius
soporator* (Valenciennes, 1837) (Teleostei, Gobiidae). *Genetics and Molecular
Biology*, 32, 2, 394-398 (2009).
161. Zúñiga-González, G.M., Torres-Bugarín, O., Zamora-Perez, A., Gómez-Meda,
B.C., Ramos-Ibarra, M.L., Gallegos-Arreola, M.P., Flores-García, A. & López-
Uribe, A. (2003b). Induction of micro-nucleated erythrocytes in mouse peripheral

- blood after cutaneous application of 5-fluorouracil. Archives of Medical Research, 34 932
141-144. 933
162. Masuda S1, Deguchi Y, Masuda Y, Watanabe T, Nukaya H, Terao Y, Takamura 934
T, Wakabayashi K, Kinae N. Genotoxicity of 2-[2-(acetylamino)-4-[bis(2- 935
hydroxyethyl)amino]-5-methoxyphenyl]-5-amino-7-bromo-4-chloro-2H- benzotriazole 936
(PBTA-6) and 4-amino-3,3'-dichloro-5,4'-dinitro-biphenyl (ADDB) in goldfish 937
(*Carassius auratus*) using the micronucleus test and the comet assay. Mutat Res. 2004 938
9, 560(1):33-40. 939
163. Bolognesi C., Perrone E, Roggieri P, Pampanin DM and Sciutto A (2006) 940
Assessment of micronuclei induction in peripheral erythrocytes of fish exposed to 941
xenobiotics under controlled conditions. Aquat Toxicol 78S:S93-S98. 942
164. Teles M, Pacheco M and Santos MA (2003) *Anguilla anguilla* L. Liver 943
ethoxyresorufin O-deethylation, glutathione S-transferase, erythrocytic nuclear 944
abnormalities, and endocrine responses to naphthalene and β -naphthoflavone. 945
Ecotoxicol. Environ. Saf. 55:98-107 946
165. Hooftman, R. N., de Raat, W. K. (1982) Induction of nuclear anomalies 947
(micronuclei) in the peripheral blood erythrocytes of the eastern mudminnow *Umbra* 948
pygmaea by ethyl methanesulphonate. Mutat. Res., 104, 147–152. 949
166. I. Dobrzynska, B. Szachowicz-Petelska, J. Ostrowska, E. Skrzydlewska, Z. 950
Figaszewski, Protective effect of green tea on erythrocyte membrane of different age 951
rats intoxicated with ethanol, Chem. Biol. Interact. 156 (2005) 41–53. 952

- 167- Ajila C., Prasadaraao, U., 2008. Protection against hydrogen peroxide induced 953
 oxida-tive damage in rat erythrocytes by *Mangifera indica* L. peel extract. 954
Food and Chemical Toxicology 46, 303–309. 955
- 168- Kim, Yu-Kyung, Kwon, Eun-Hee, Kim, Dong-Hyun, Won, Dong-I, Shin, Sehyun, 956
 Suh, Jang-Soo, Susceptibility of oxidative stress on red blood cells exposed to gamma 957
 rays: Hemorheological evaluation . *Clinical Hemorheology and Microcirculation* 40, 958
 4, 315-324, 2008. 959
169. Gustavino, B., Scornajenghi, K.A., Minissi, S., Ciccotti, E., 2001. Micronuclei 960
 induced in erythrocytes of *Cyprinus carpio* (teleostei, pisces) by X-rays and 961
 colchicines. *Mutat. Res.* 494, 151–159. 962
 963




Figure 1. A, erythrocytes of mammals (rounded, non-nucleated). B, erythrocytes of camel (oval, non-nucleated). C, erythrocytes of birds, fish and amphibians (oval, nucleated). D, the structure of erythrocyte membrane (Figures are collected from the internet)




Figure2. Erythrocytes as a model for evaluating the cytogenotoxic effects of xenobiotics

975




Figure 3. Redox regulation in RBCs

976

977

978

979

980

981

982




Figure 4. Different effects of xenobiotics on erythrocytes

983

984

985

986

Table 1. Type of damage resulted from interaction of some xenobiotics and RBCs from different animal species.

987
988

The xenobiotic	The type of damage to RBC	The species of the RBC	References
Lead	Oxidative stress , apoptotic events (Fas aggregation in lipid rafts led to Fas-dependant death of RBCs) GSH and GSH-HNE adduct Gardos channel Ceramide formation inhibition of acid sphingomyelinase (aSMase)	Mice	[17]
	poikilocytosis and anisocytosis	Whooper swans (<i>Cygnus Cygnus</i>)	[23]
	Anisocytosis, Poikilocytosis ,changes in hemoglobin and oxidative damage	Rats	[24]
	Pycnosis and enucleation of peripheral erythrocytes	Chicken	[25]
	shortened lifespan , inhibited Na-K-ATPase and loss of membrane integrity	Rat	[49]
	Stimulation of phosphatidylserine exposure at the erythrocyte surface	Human	[68]
	Fas induced apoptotic events in erythrocytes	Mice	[72]
	Abnormalities of RBCs	Chickens	[119]
	genotoxicity (micronuclei and other nuclear abnormalities	Freshwater fish <i>Prochilodus lineatus</i>	[120]
	oxidative damage	Rat	[121]
	Decreased antioxidant levels	Goats	[122]
	Morphological changes	Human	[123]
	Altered activity of potassium-selective channels	Human	[124]
	Induction of micronucleus	The fish H. Malabaricus	[125]
Copper	Genotoxicity (increased micronuclei and other nuclear abnormalities)	Freshwater fish species viz. <i>Labeo rohita</i> , <i>Cirrhina mrigala</i> , <i>Catla catla</i> and <i>Ctenopharyngodon idella</i>	[126]
Cadmium chloride and copper sulphate	micronuclei and binuclei	Fish Common carp (<i>Cyprinus carpio</i>), Prussian carp (<i>Carassius gibelio</i>)	[127]

		and Peppered cory(<i>Corydoras paleatus</i>)	
Cadmium chloride	genotoxicity and cytotoxicity cellular abnormalities and DNA damage	Fresh water fish Labeo rohita (Hamilton)	[35]
	Genotoxic effects	<i>Dreissena polymorpha</i>	[59]
Cadmium, copper, lead and zinc	Genotoxicity and cytotoxicity	Nile tilapia, (<i>Oreochromis niloticus</i>)	[128]
Mercury	Induction of micronuclei	Cyprinus carpio	[129]
	Nuclear abnormalities	Wild and caged fish (<i>Liza aurata</i>)	[130]
Potassium dichromat	hemolytic effect	Rats	131
Arsenic	Fas-activated erythrocyte apoptosis	Rats	[132]
	Reduced cellular redox status, induced 4- hydroxynonenal-mediated caspase 3 activation, erythrocyte death	Rats	[71]
	erythrocyte death	Human	[98]
Aluminium	Morphologic and functional alterations		[27]
	Alter Morphology	Rabbit	[133]
Vanadium aluminium	Oxidative stress	Human	[134]
Iron	oxidative stress	Human	[135]
Fluoride	Apoptosis, altered membrane integrity, altered cell morphology and size, induced moderate ceramide formation,	Rat	[44]
	Oxidative stress	Mice	[136]
Vanadate and magnesium	Attenuated defence system	Wistar rats	[137]
Iodoacetate, vanadate and ferri cyanide	Peroxidative membrane damage	human	[138]
Lambda-cyhalothrin	Genotoxicity (micronucleus)	Fish (Garra rufa (Pisces: Cyprinidae)	[139]
	Oxidative stress	Rat	[140]
	Oxidative stress decrease the activity of	Rabbit	[92]

	acetylcholinesterase (AChE)		
	Micronuclei and nuclear abnormalities	Mosquito fish (<i>Gambusia affinis</i>)	[141]
	Micronucleus	Erythrocytes of the fish (<i>Cheirodon interruptus</i>)	[142]
Organophosphate pesticides (chlorpyrifos and malathion), synthetic pyrethroid pesticide (cypermethrin, lambda-cyhalothrin)	Genotoxicity (micronuclei)	Fish (<i>Oreochromis mossambicus</i>)	[143]
Pyrethroid	Lipid peroxidative damage and alterations in antioxidant status	Rat	[144]
Phenoxyherbicides	Production of free radicals	Human	[145]
Phenoxyherbicides (e.g., 2,4-dichlorophenoxyacetic acid (2,4-D-Na), 2,4,5-trichlorophenoxyacetic acid (2,4,5-T-Na) and 4-chloro-2-methylphenoxyacetic acid (MCPA-Na))	Altered erythrocyte membrane fluidity and changed membrane proteins content	Erythrocytes	[146]
2,4-dichlorophenoxyacetic herbicide	Attenuated antioxidant defense system and induced lipid peroxidation	Rat	[45]
2,4-dichlorophenoxyacetic acid and butachlor.	Induced micronuclei and erythrocyte alterations	Catfish (<i>Clarias batrachus</i>)	[147]
-2,4-dichlorophenoxy acetic-acid- and butachlor	DNA damage by alkaline single cell gel electrophoresis	Catfish (<i>Clarias batrachus</i>)	[148]
Chlorpyrifos	Alterations in the profile of blood cells, alterations in the cytoskeleton [protein and lipids] of RBCs from wistar rats thus affecting the cell surface area	Wistar rats	[32]
	Lipid peroxidation and oxidative stress	Rat	[43]
	Genotoxicity, micronucleus	Freshwater fish (<i>Channa punctatus</i>)	[149]
Lindane	Oxidative stress	Rat	[150]
hexachlorocyclohexane	Biochemical and structural	Rat	[34]

	alterations		
Fenvalerate and its metabolite.	Morphological and biochemical perturbations	Rat	[151]
Endosulfan	The MN was observed in RBCs of chickens exposed to endosulfan	Chicks	[57]
Malathion and endosulfan	Changes in surface shapes and structural defects of RBCs of exposed to malathion and endosulfan	Rats	[30]
The herbicide (Roundup)	Micronuclei, nuclear abnormalities and DNA damage	Freshwater goldfish (<i>Carassius auratus</i>)	[152]
The herbicides AAtrex Nine-O-, Dual-960E-, Roundup-, and Sencor-500F	DNA damage by comet (DNA fragmentation)	Erythrocytes from <i>Rana catesbeiana</i> (bullfrog) tadpoles	[153]
The fungicide (Propiconazole)	Genotoxicity (micronuclei)	Fresh water fish (<i>Clarias batrachus</i>)	[154]
Tributyltin	genotoxicity (micronucleus)	The fish <i>H. Malabaricus</i>	[125]
Anionic detergent	Echinocytosis	Bullhead, <i>Ictalurus melas Rafinesque</i>	[9]
Textile mill effluent	Micronuclei and other nuclear abnormalities	<i>Oreochromis niloticus</i>	[155]
Petroleum refinery and chromium processing plant effluents	Micronuclei and nuclear abnormalities	<i>Oreochromis niloticus</i>	[156]
Waters affected by refinery effluent	Micronucleus test and observation of nuclear alterations	Nile tilapia	[157]
Mitomycin-C (cytotoxic antineoplastic agent)	Nuclear abnormalities (micronuclei and nuclear buds)	Parrots (<i>Aratinga canicularis</i>)	[158]
Mitomycin C and cyclophosphamide	Micronuclei	Fathead minnow (<i>Pimephales promelas</i>)	[159]
Cyclophosphamide	Genotoxicity using the micronucleus assay and nuclear abnormalities	The tropical sea fish (<i>Bathygobius soporator</i>)	[160]
5-Fuorouracil	Micronuclei	Mouse	[161]
Cyadox	Altered antioxidant defense system and induced hemolysis	Rabbit	[11]

	Induced energy depletion by decreasing their ATP contents	Rabbit	[50]
2-[2-(acetylamino)-4-[bis(2-hydroxyethyl)amino]-5-methoxyphenyl]-5-amino-7-bromo-4-chloro-2H-benzotriazole (PBTA-6) and 4-amino-3,3'-dichloro-5,4'-dinitro-biphenyl (ADDB)	Genotoxicity DNA damage	Goldfish (<i>Carassius auratus</i>)	[162]
Dialkyl phthalate, bisphenol A, tetrabromodiphenyl ether	Micronuclei	Fish (<i>Scophthalmus maximus</i>)	[163]
Tert-Butyl hydroperoxide	Alters fatty acid incorporation into erythrocyte membrane phospholipid.	Human	[112]
4-nonylphenol	ATP content which consequently forces cells to show vacuolization and lead also to unequal distributed hemoglobin which resulted in RBCs cytoplasmic vacuoles	African catfish (<i>Clarias gariepinus</i> Burchell)	[48]
acrylamide	Cytogenetic effects	<i>Carassius auratus</i>	[58]
Naphthalene and β -naphthoflavone	Erythrocytic nuclear abnormalities	<i>Anguilla anguilla</i> L.	[164]
Ethyl methane sulphonate	Nuclear anomalies (micronuclei)	Fish (eastern mudminnow <i>Umbra pygmaea</i>)	[165]
Ethanol	erythrocyte membrane	Rats	[166]
Hydrogen peroxide	Oxidative damage	Rat	[167]
Irradiation (gamma-ray)	Induced biochemical changes, generate reactive oxygen species (ROS), deformability, lipid peroxidation	Human	[168]
X-rays and colchicines	Micronuclei induced	<i>Cyprinus carpio</i> (teleostei, pisces)	[169]
Flavonoids of <i>Aristotelia chilensis</i> Maqui leaves	Morphological alterations represented by echinocytic form	Human	[31]

Agaricus sylvaticus	Hemolytic activity	Human	[37]
Thymoquinone and limonene	Alleviation of plasma, erythrocyte and liver lipidemic-oxidative stress	Rats	[80]

989

990

991

992

993

994

995

996

997

998

Highlights

- Exposure of the body to xenobiotics results in cytotoxicity and genotoxicity
- Erythrocytes contain specific structure and antioxidant system
- Erythrocytes with its structure are valuable in screening of xenobiotics toxicity