

## Transactions of the Royal Society of South Africa

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/ttrs20>

### AN ACCOUNT OF THE PLANT ECOLOGY OF THE DUNE FOREST AT LAKE SIBAYI

C. M. Breen <sup>a</sup>

<sup>a</sup> Department of Botany and Microbiology , Rhodes University)

Published online: 22 Mar 2010.

To cite this article: C. M. Breen (1971) AN ACCOUNT OF THE PLANT ECOLOGY OF THE DUNE FOREST AT LAKE SIBAYI, Transactions of the Royal Society of South Africa, 39:3, 223-234, DOI: [10.1080/00359197109519117](https://doi.org/10.1080/00359197109519117)

To link to this article: <http://dx.doi.org/10.1080/00359197109519117>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

# AN ACCOUNT OF THE PLANT ECOLOGY OF THE DUNE FOREST AT LAKE SIBAYI

By C. M. BREEN

(Department of Botany and Microbiology, Rhodes University)

(With 3 text-figures and 2 plates)

(Read April 15, 1970)

## OPSOMMING

Die legging, fisiografie en klimaat van die duinwoud in die omgewing van Sibayimeer, word beskryf. Die algemene ekologie van die woud, met besondere verwysing na die samestelling en digtheid van die bome en struik, word uiteengesit. Die waarnemings word bespreek met betrekking tot die oorlewing van die belangrikste bome wat die daklaag vorm. Dit lei tot die gevolgtrekking dat die woud 'n stabiele tipe plantegroei, d.w.s. klimaksduinwoud, verteenwoordig. 'n Voorlopige lys van plante, wat in die duinwoud versamel is, word gegee.

## SUMMARY

The location, physiography and climate of the dune forest in the vicinity of Lake Sibayi is given. The general ecology of the forest, with particular reference to composition and density of trees and shrubs, is outlined. The results are discussed in relation to survival of the most important canopy trees. It is concluded that the forest represents a stable vegetation type, i.e. Climax Dune Forest. A preliminary list of plants collected in the dune forest is appended.

## INTRODUCTION

The coastal dune forest (Coastal Tropical Forest, Acocks, 1953) in the vicinity of Lake Sibayi ( $32^{\circ}43'E$  between approximately  $27^{\circ}19'S$  and  $27^{\circ}25'S$ ) is part of the Bantu Trust area of Tongaland. The area is administered by the Department of Bantu Administration and, consequently, provision has been made for proper control of the indigenous forests. However, cutting has not been strictly controlled because, until recently, there was no alternative source of timber for the Bantu in the area. The forestry project, which was started in 1959, now serves as a source of employment and as a supply of timber for construction and fuel. The demands on the indigenous forests have, therefore, decreased gradually and, at present, little cutting occurs in the forest.

In some parts of Zululand, the Department of Bantu Administration has successfully made large plantings of *Casuarina* to stabilize the dune system. Between 1948 and 1957, the Department planted *Casuarina* on some of the larger exposed areas at Sibayi (fig. 1) and these areas have been successfully stabilized.

The soil in the Sibayi area is sandy and will not support crops; consequently the Bantu cultivated areas in the forest to utilize the accumulated organic debris which resulted in improved crops. As with cutting in the forest, control had to be applied gradually as alternative sources of livelihood were developed and improved farming methods introduced, e.g. use of fertilizer. At the present time, agriculture in the dune system has been stopped.

In view of the history of the area, it seemed desirable to start ecological investi-

gations on the dune forest. This paper is the result of the preliminary investigations designed to describe the general structure of the forest as a whole and to determine the extent to which the canopy-forming trees have been affected by the practices outlined above.

### TOPOGRAPHY

The area consists of a ridge of stabilized sand, rising steeply (slope of about  $30^\circ$ ) from the lake shore in the west (70 feet above sea-level) to a maximum elevation of 449 feet (Beacon, fig. 1). Eastwards, the surface undulates so that the slope down to the Indian Ocean is less steep than in the west (pl. 3). Towards the NNE., the altitude of the dune system decreases gradually so that the northernmost part of the dune forest investigated is at an altitude of approximately 200 feet.

The dune forest can be divided into three major ecological areas: A steep western aspect, an undulating area eastwards and a marginal coastal fringe.




FIG. 1. The distribution of dune forest in the vicinity of Lake Sibayi.

### CLIMATE

Climatic data are available from the Weather Bureau reports for three stations in the vicinity of Lake Sibayi: Mbazwane ( $27^\circ 30'S$ ,  $32^\circ 36'E$ ), Mseleni ( $27^\circ 22'S$ ,  $32^\circ 32'E$ ) and Ngutshana ( $27^\circ 13'S$ ,  $32^\circ 34'E$ ). The last station has few records and consequently the data available are not representative. The records for Mbazwane (12 years) and Mseleni (20 years) provide some idea of the climate and are summarized in figure 2. The rainfall of the coastal plains of northern Natal is highest near the sea and decreases inland. Thus, the coastal dune forest in the vicinity of Lake




FIG. 2. Summary of climatic data available from stations in the vicinity of Lake Sibayi.

Sibayi is likely to receive more rain than either Mseleni or Mbazwane; probably of the order of 1,000 mm annually.

Generally speaking, the climate is humid with warm, wet summers, and winters which are mild and dry by comparison.

The prevailing winds are north-east (sea breezes) and south-west (rain-bearing) and the greater proportion of the rain results from thunderstorms. Although predominantly a summer-rainfall area, on the average no month is likely to have less than about 30 mm of rain, distributed over three days.

#### METHOD

The Random Pairs Method (Cottom & Curtis, 1956) was chosen in preference to other plotless methods because the measurements are easy to take in the field and the method allows workers to cover considerable distances, enhancing the overall

impression gained as a result of field observation. There is evidence that, provided sufficient points are taken, the results obtained do not differ significantly from those obtained with the Quarter Method (Cottam & Curtis, 1956).

The method involves taking sampling points along a predetermined line. In order to obtain a random distribution of these lines within the dune forest, ten sample areas were selected as follows: the area was transferred to a grid and, using random numbers, ten areas were selected. Each area was sampled by at least 40 points. The total number of points available for analysis was 430.

Since the main object of the investigation was the structure and survival of the canopy and subcanopy, only trees greater than 6 feet in height were recorded. The height was estimated and recorded in different height classes and the girth at breast height was measured and recorded in girth classes (Table 1).

## RESULTS

### 1. General observations

The forest is composed of three strata: canopy, subcanopy and the shrub and herbaceous layer. The height of the canopy varies considerably in relation to aspect, the highest canopy (40–50 feet) occurring on the level ground. On the steep western slope (pl. 1), the canopy is low (10–20 feet) with occasional emergent trees, e.g. *Balanites maughamii*, *Ziziphus mucronata* and *Mimusops caffra*, consequently there is little evidence of a subcanopy. Few of the large trees on this slope remain vertical as the substrate is loose and easily dislodged.

Where the canopy is continuous, the herbaceous stratum is not well developed. However, in areas where the canopy has been broken (e.g. margins of exposed areas) dense undergrowth develops. Thickets of *Isoglossa* sp. up to 10 feet tall are common in these areas. The shrubs *Tricalysia lanceolata*, *Clerodendrum glabrum*, *Grewia occidentalis* and tree saplings, principally *Croton gratissimus*, *Deinbollia oblongifolia* and *Strychnos* sp., are common.

Climbers occur infrequently in the forest, although in areas of succession *Grewia caffra*, *Grewia occidentalis* and *Acacia schweinfurthii* are common. Other creepers collected were *Dalbergia armata*, *Popowia caffra*, *Rhoicissus* spp., *Secamone frutescens* and *Smilax kraussiana*.

Where the forest approaches the lake shore a fringe of *Acacia karroo* occurs. During the course of this work, it became apparent that *A. karroo* is common in the dune forest (Table 1). However, it appeared that *A. karroo* was distributed mainly along the drainage areas of the dune system, in areas where the original forest cover had been removed. The Random Pairs Method of ecological investigation is not suited to illustrate the tendency for plants to occur in such groves. However, even with this method, of the 63 occurrences of *A. karroo*, 36 were recorded as random pairs, i.e. at 18 points only *A. karroo* was recorded. *A. karroo* occurred 27 times in conjunction with other species. In view of the large number of points taken (430), this result shows the tendency for *A. karroo* to occur in stands. Beneath the stands of *A. karroo*, there is little evidence of regeneration of either *A. karroo* or of other forest species.

## 2. Results of Plotless Sampling

To determine the survival of the canopy-forming trees, it is necessary to establish firstly that sufficient members of potential canopy-formers are available to replace any dying in the canopy, and secondly that the species composition of the canopy is unlikely to change, i.e. that the potential canopy-formers in the lower strata are the same species as those presently forming the canopy.

Since height and girth are to some extent representative of age, the use of height and girth classes can supply an index of survival. The prevalence of plants in the middle height and, to a lesser extent, in the middle girth classes (fig. 3) suggests that there is available an abundance of potential canopy-forming plants. However, plants which seldom, if ever, contribute to the canopy, could contribute to this general pattern. A consideration of the numbers of canopy-forming species only (i.e. those recorded in the greater than 30 feet height category), however, reveals a strikingly similar pattern (fig. 3). Thus, there are a large number of young canopy plants.




FIG. 3. The distribution of stems in height and girth classes.

TABLE 1

Data compiled from plotless sampling

SPECIES	Number of individuals	Relative density	GIRTH (inches)										HEIGHT (feet)				
			<2	2	3	4	5	6	7-12	13-18	19-24	>24	6	7-10	10-20	21-30	>30
<i>Drypetes natalensis</i>	86	10.0	1	—	3	10	6	4	32	15	11	4	—	8	41	29	8
<i>Acacia karroo</i>	63	7.3	—	2	2	2	4	1	22	12	6	12	—	4	11	21	27
<i>Teclea natalensis</i>	58	6.7	1	5	11	7	5	5	16	3	2	3	—	19	27	10	2
<i>Mimusops caffra</i>	50	5.8	—	—	2	5	6	—	9	4	4	20	—	3	15	9	23
<i>Cassipourea gerrardii</i>	41	4.8	—	1	9	5	5	3	12	3	—	3	—	10	20	6	5
<i>Plectroniella armata</i>	39	4.5	1	2	5	6	3	7	13	1	1	—	—	11	20	8	—
<i>Croton gratusimus</i>	25	2.9	—	—	2	1	2	2	7	3	2	6	—	2	9	7	7
<i>Diospyros inhaecensis</i>	25	2.9	—	—	4	3	4	3	3	5	1	2	—	6	13	4	2
<i>Acalypha glabrata</i>	24	2.8	—	3	6	2	1	5	3	2	1	1	—	8	11	4	1
<i>Placoxylon obliquum</i>	22	2.5	1	1	5	4	1	1	5	2	1	1	—	6	9	7	—
<i>Euclea natalensis</i>	21	2.4	—	1	1	2	1	—	7	4	4	1	—	4	7	5	5
<i>Clausena anisata</i>	20	2.3	2	4	2	2	2	2	4	—	2	—	—	2	13	5	—
<i>Diospyros scabrida</i>	17	2.0	2	1	3	3	2	1	4	—	—	1	—	2	15	—	—
<i>Cassine aethiopica</i>	16	1.9	—	3	4	3	6	—	7	2	1	—	—	3	7	6	—
<i>Deinbollia oblongifolia</i>	16	1.9	—	—	2	1	2	—	—	—	—	—	—	1	7	8	—
<i>Diospylos longispina</i>	16	1.9	1	2	2	4	3	1	1	1	1	—	—	2	9	3	1
<i>Pavetta gerrardii</i>	15	1.7	1	—	1	1	2	1	2	3	1	2	—	3	9	1	—
<i>Celtis africana</i>	13	1.5	—	—	—	1	2	1	2	—	—	—	—	2	4	3	4
<i>Tricalysia lanceolata</i>	13	1.5	1	2	3	4	2	—	1	—	—	—	—	1	5	1	—
<i>Clerodendrum glabrum</i>	12	1.4	—	—	—	1	5	—	6	—	—	—	—	—	11	1	—
<i>Drypetes gerrardii</i>	11	1.3	—	1	1	6	1	3	3	1	—	—	—	—	7	2	1
<i>Tapura fischeri</i>	11	1.3	1	—	1	1	1	2	2	1	1	—	—	5	6	—	—
<i>Morus mesozygia</i>	10	1.2	—	1	—	1	1	1	3	1	—	—	—	1	6	1	1
<i>Erythroxylum emarginatum</i>	9	1.0	—	1	4	—	—	—	1	2	1	2	—	5	2	1	1
<i>Mimusops obovata</i>	9	1.0	—	—	—	—	1	1	1	2	1	—	—	—	4	3	2
<i>Rausonia lucida</i>	9	1.0	—	1	1	—	—	2	—	1	1	2	—	—	4	1	2
<i>Tricalysia capensis</i>	9	1.0	—	1	3	3	—	—	—	1	—	—	—	1	5	1	—
<i>Xeromphus obovata</i>	9	1.0	—	—	3	1	1	1	1	1	—	5	—	2	4	2	1
<i>Ziziphus mucronata</i>	9	1.0	—	—	—	2	1	—	5	—	—	—	—	1	3	1	4
<i>Allophylus natalensis</i>	8	0.9	—	1	—	1	1	—	4	3	—	—	—	2	4	1	1
<i>Linociera peglerae</i>	8	0.9	—	—	—	—	—	1	—	—	—	—	—	—	4	4	1
<i>Maylenus nemorosa</i>	8	0.9	1	—	1	1	—	2	3	2	—	1	—	—	4	2	—
<i>Drypetes arguta</i>	7	0.8	—	1	2	1	1	—	3	—	—	—	—	2	3	1	1
<i>Ochna arborea</i>	7	0.8	1	2	1	1	1	—	2	—	—	—	—	2	3	1	—
<i>Strychnos innocua</i>	7	0.8	—	—	1	1	1	—	—	—	2	2	—	2	1	2	2
<i>Vepria lanceolata</i>	7	0.8	2	—	—	1	—	—	1	3	—	—	—	2	1	4	—
<i>Balanites manghamii</i>	6	0.7	—	—	—	1	1	1	4	2	—	—	—	1	3	—	2
<i>Diospyros natalensis</i>	6	0.7	—	—	—	—	—	—	—	—	—	—	—	—	3	3	—
<i>Coffea racemosa</i>	5	0.6	—	2	—	—	2	1	—	—	—	—	—	1	3	1	—
<i>Cola natalensis</i>	5	0.6	—	—	—	—	—	2	—	—	—	—	—	1	4	—	—
<i>Ephippiocarpa orientalis</i>	5	0.6	—	—	2	1	—	—	1	—	—	—	—	1	3	—	—
<i>Ficus natalensis</i>	5	0.6	—	—	—	1	—	—	2	1	—	—	—	—	2	1	—
<i>Lecomtedoxa henriquesii</i>	5	0.6	—	—	1	—	1	—	—	—	2	1	—	—	1	1	2

Mean distance apart 5' 4"  
Density per acre 2361



The data in Table 1 indicate that *Drypetes natalensis*, *Acacia karroo*, *Teclea natalensis*, *Mimusops caffra*, *Cassipourea gerrardii*, *Croton gratissimus*, *Diospyros inhacaensis*, *Ptaeroxylon obliquum* and *Euclea natalensis* are the most common plants which contribute to the canopy. However, since their relative density does not correspond directly with their occurrence in the greater than 20 feet classes (Table 2), they may be separated into the components of the canopy (greater than 20 feet) and the subcanopy (less than 20 feet) (Table 3).

Of the ten more common plants of the subcanopy listed in Table 3, five are typical of the canopy. Thus the canopy-forming species are present in the lower strata. Not only are the canopy-forming species present in the lower strata, they are common enough to ensure the continuation of the canopy in its present form. It would seem unlikely, therefore, that there will be any significant change in the canopy in the near future.

TABLE 2

*The major components of the greater than 20 feet height categories*

SPECIES	Relative density	Number of occurrences
<i>Acacia karroo</i>	63	48
<i>Drypetes natalensis</i>	86	37
<i>Mimusops caffra</i>	50	32
<i>Croton gratissimus</i>	25	14
<i>Teclea natalensis</i>	58	12
<i>Cassipourea gerrardii</i>	41	11
<i>Euclea natalensis</i>	21	10
<i>Ptaeroxylon obliquum</i>	22	7
<i>Celtis africana</i>	13	7

TABLE 3

*For explanation, see text*

Canopy	Subcanopy
<i>Acacia karroo</i>	<i>Drypetes natalensis</i>
<i>Mimusops caffra</i>	<i>Teclea natalensis</i>
<i>Drypetes natalensis</i>	<i>Cassipourea gerrardii</i>
<i>Croton gratissimus</i>	<i>Plectroniella armata</i>
<i>Cassipourea gerrardii</i>	<i>Mimusops caffra</i>
<i>Euclea natalensis</i>	<i>Croton gratissimus</i>
<i>Celtis africana</i>	<i>Clausena anisata</i>
<i>Ziziphus mucronata</i>	<i>Acacia karroo</i>
	<i>Acalypha glabrata</i>
	<i>Clerodendrum glabrum</i>

The shrub layer is formed mainly by young canopy formers and by other predominantly shrub types such as *Plectroniella armata*, *Acalypha glabrata*, *Clausena anisata*, *Clerodendrum glabrum* and *Cassine aethiopica*.

#### DISCUSSION

The areas of exposed sand evident in aerial photographs taken in 1942 (pl. 2) are generally confined to the hollows between the sand dunes. The drainage and accumulation of organic material in these hollows provide the most suitable substrate for agriculture. Thus, there can be little doubt that the exposed areas are the result

of agricultural practices which were common in the dune system prior to 1948. These practices have been controlled since about 1948 and had certainly ceased by 1959 when the forestry project was started in the area by Mr. B. H. Groenewald. However, Bantu were still resident along the forest margin in 1959. These families had moved by the author's first visit in 1966 although there are still families near the northern margin. These latter families are probably responsible for the slight encroachment in the area (pl. 3 and 4).

In traverses through the forest, there is little evidence of tree felling and discussions with the Bantu and with a former Chief Forester (Mr. B. H. Groenewald) suggest that the Bantu are not 'species' selective in their cutting. It is likely, therefore, that they restrict most of their felling to the forest margin, particularly in the north.

Control has been effected for about twenty years. During this period, the exposed areas have been recolonized and there has been a general increase in the density of the vegetation. Whether exposed areas still exist is not possible to discover since the latest available aerial photographs were taken in 1964. The exposed areas evident in these photographs have, in all probability, been further reduced.

The presence of groves of *A. karroo* in areas which were previously exposed suggests they may represent a seral stage in the development of dune forest, as has been suggested by Edwards (1967). The lack of regeneration of climax species in these groves may be due to the relatively short period of recolonization.

The marginal fringe of *A. karroo* along the lake shore probably represents a subclimax and not a seral stage in dune forest formation. This subclimax is developed in conjunction with the greater availability of water, a factor which may be of importance in the recolonization of the exposed areas in the drainage system.

In spite of evidence of tree-felling and encroachment in the north, the canopy-forming species are abundant in the middle girth and height categories. This suggests that the future of the canopy is assured. However, at this stage it is not possible to discover whether this will result in an increase in the density of the canopy.

#### CONCLUSIONS

1. The control of agriculture and cutting in the dune forest has resulted in considerable improvement in the forest.
2. Regeneration is such that the character of the forest is unlikely to change in the near future and, therefore, the forest represents a Climax Forest. The canopy and subcanopy trees have a high survival rate as evidenced by their distribution in the height and girth classes.
3. Apart from the limited encroachment as a result of agricultural practices since 1942, cutting by the local inhabitants, on the present scale, does not appear to be having any detrimental effect on the survival of the canopy trees.

#### ACKNOWLEDGEMENTS

I should like to express my thanks to Professor B. R. Allanson, the Staff of the Botany Department and the students who have accompanied me; Mrs. Estelle Brink of the Albany Museum, Grahamstown and the staff of the National Herbarium,

Pretoria; Mr. McKay, Chief Agricultural Extension Officer (Zululand) for the Department of Bantu Affairs, and Mr. B. H. Groenewald, Chief Forester, Port Durnford.

This work has been financed by grants from C.S.I.R. and Rhodes University Council.

## REFERENCES

- ACOCKS, J. P. H. 1953. Veld Types of South Africa. *Mem. Bot. Surv. S. Afr. No. 28*.  
 COTTAM, G. & CURTIS, J. T. 1956. The use of distance measures in Phytosociological sampling. *Ecology*, 37, 741-757.  
 EDWARDS, D. 1967. A Plant Ecological Survey of the Tugela River Basin. *Mem. Bot. Surv. S. Afr. No. 36*.  
 Weather Bureau. 1954. *Climate of South Africa*. Part 1. W.B. 19 Govt. Printer.  
 Weather Bureau (various). *Annual Reports of the Weather Bureau*. Govt. Printer.

## APPENDIX

### LIST OF ANGIOSPERMS COLLECTED IN THE DUNE FOREST

#### GRAMINEAE

- Panicum deustum* Thunb.  
*Pennisetum typhoides* (Burm.) Stapf & Hubb.

#### CYPERACEAE

- Cyperus maritimus* Poir.

#### DIOSCOREACEAE

- Dioscorea sylvatica* Eckl.

#### ORCHIDACEAE

- Eulophidium tainioides* (Schltr.) Summerh.

#### ULMACEAE

- Celtis africana* Burm. f.  
*Trema guineensis* (Schum.) Ficalho  
*Chaetachme aristata* Planch.

#### MORACEAE

- Morus mesozygia* Stapf.  
*Ficus burtt-davayi* Hutch.  
*Ficus natalensis* Hochst.

#### LORANTHACEAE

- Loranthus kraussianus* Meissn.

#### SANTALACEAE

- Colpoen compressum* Berg.

#### OLACACEAE

- Ximenia caffra* Sond.

#### PHYTOLACCACEAE

- Limeum viscosum* (Gay) Fenzl

#### ANNONACEAE

- Popowia caffra* (Sond.) Benth.

#### LAURACEAE

- Cryptocarya* sp.

#### CAPPARIDACEAE

- Cladostemon kirkii* (Oliv) Pax & Gilg  
*Capparis citrifolia* Lam.

#### PITTOSPORACEAE

- Pittosporum undulatum* Vent.

#### LEGUMINOSAE

- Albizia adianthifolia* (Schum.)  
 W. F. Wright  
*Acacia karroo* Hayne  
*Acacia schweinfurthii* Bren. & Exell  
*Bauhinia tomentosa* L.  
*Sophora inhambanensis* Klotzsch  
*Crotalaria capensis* Jacq.  
*Indigofera* ? *endecaphylla* Jacq.  
*Dalbergia armata* E. Mey.  
*Dalbergia nitidula* Welw. ex Benth.  
*Craibia zimmermannii* (Harms)  
 Harms ex Dunn

#### ERYTHROXYLACEAE

- Erythroxylum emarginatum* Thonn.  
*Erythroxylum pictum* E. Mey

#### ZYGOPHYLLACEAE

- Balanites maughanii* Sprague

#### RUTACEAE

- Fagara capensis* Thunb.  
*Calodendron capensis* (L.f.) Thunb.  
*Vepris lanceolata* G. Don  
*Teclea natalensis* (Sond.) Engl.  
*Clausena anisata* (Willd.) Hook f. ex Benth.

#### MELIACEAE

- Ptaeroxylon obliquum* (Thunb.) Radlk.  
*Turraea floribunda* Hochst.  
*Ekebergia capensis* Sparrm.  
*Trichilia emetica* Vahl.

#### DICHAPETALACEAE

- Tapura fischeri* Engl.

**EUPHORBIACEAE**

- Phyllanthus* sp.  
*Drypetes arguta* Hutch.  
*Drypetes gerrardii* Hutch.  
*Drypetes natalensis* Hutch.  
*Antidesma venosum* E. Mey. ex Tul.  
*Cleistanthus schlechteri* (Pax)  
 Hutch. var. *schlechteri*  
*Bridelia cathartica* Bertol. f. subsp. *cathartica*  
 (?*Bridelia schlechteri* in Tinley)  
*Croton gratissimus* Burch.  
*Erythrococca berberidea* Prain.  
*Acalypha glabrata* Thunb.  
*Tragia meyeriana* Meull.-Arg.  
*Suregada africana* (Sond.) Kuntze  
*Spirostachys africana* Sond.

**CELASTRACEAE**

- Maytenus acuminata* (L.f.) Loes.  
*Maytenus nemorosa* (Eckl. Zeyh.) Marais  
*Allocassine laurifolia* (Harv.) N. Robson  
*Cassine aethiopica* Thunb.  
*Cassine peragua* L.

**ICACINACEAE**

- Apodytes dimidiata* E. Mey. ex Arn.

**SAPINDACEAE**

- Allophylus melanocarpus* (Sond.) Radlk.  
*Allophylus natalensis* (Sond.) De Winter  
*Deinbollia oblongifolia* (Sond.) Radlk.  
*Pancovia gologensis* (Hiern) Exell & Mendonca  
*Blighia unijugata* Bak.  
*Dodonaea viscosa* Jacq.

**RHAMNACEAE**

- Ziziphus mucronata* Willd.

**VITACEAE**

- Rhoicissus digitata* (L.f.) Gilg & Brand  
*Rhoicissus rhomboidea* (E. Mey. ex Harv.)  
 Planch.

**TILIACEAE**

- Grewia caffra* Meissn.  
*Grewia occidentalis* L.

**MALVACEAE**

- Thespesia acutifolia* (Bak.f.) Exell &  
 Mendonca

**STERCULIACEAE**

- Cola natalensis* Oliv.

**OCHNACEAE**

- Ochna arborea* Burch. ex DC.  
*Ochna natalita* (Meissn.) Walp.

**VIOLACEAE**

- Rinorea ilicifolia* (Welw. ex Oliv.)  
 Kuntze var. *ilicifolia*

**FLACOURTIACEAE**

- Rawsonia lucida* Harv. & Sond.  
*Xylotheca kraussiana* Hochst. var. *glabrifolia*  
 Wild

- Scolopia zeyheri* (Nees) Harv.  
*Dovyalis longispina* (Harv.) Harv.  
*Dovyalis* sp.

**OLINIACEAE**

- Olinia radiata* J. Hofmeyr & Phill.

**THYMELEACEAE**

- Peddiea africana* Harv.

**RHIZOPHORACEAE**

- Cassipourea gerrardii* Alston

**MYRTACEAE**

- Eugenia natalita* Sond.  
*Syzygium cordatum* Hochst.

**ARALIACEAE**

- Cussonia spicata* Thunb.

**SAPOTACEAE**

- Sideroxylon inerme* L.  
*Mimusops caffra* E. Mey. ex A. DC.  
*Mimusops henriquesiana* Sim  
*Mimusops obovata* Sond.  
*Austromimusops* sp. cf. *marginata* (N.E. Br.)  
 A. Meeuse  
*Lecomtedoxa henriquesii* (Engl. & Warb.)  
 A. Meeuse  
*Manilkara concolor* (Harv. ex C.H. Wr.)  
 Gerstner

**EBENACEAE**

- Diospyros inhacaensis* F. White  
*Diospyros natalensis* (Harv.) Brenan  
*Diospyros scabrida* (Harv. ex Hiern) De  
 Winter  
*Diospyros whyteana* (Hiern) F. White  
*Euclea divinorum* Hiern  
*Euclea natalensis* A. DC.

**OLEACEAE**

- Linociera peglerae* (C.H. Wr.) Gilg &  
 Schellenb.  
*Olea capensis* L.

**LOGANIACEAE**

- Strychnos innocua* Del. subsp.  
*gerrardii* (N.E. Br.) Verdoorn  
*Strychnos henningsii* Gilg  
*Strychnos spinosa* Lam.

**APOCYNACEAE**

- Acokanthera oppositifolia* (Lam.) L.E. Codd  
*Acokanthera schimperi* (A. DC.)  
 Schweinf. var. *rotundata* Codd  
*Carissa bispinosa* (L.) Desf. ex Brenan var.  
*acuminata* (E. Mey.) Codd  
*Conopharyngia elegans* (Stapf) Stapf  
*Ephippiocarpa orientalis* (S. Moore) Markgf.

**ASCLEPIADACEAE**

- Secamone frutescens* Decne.

**BORAGINACEAE**

- Cordia caffra* Sond.

**VERBENACEAE**

*Clerodendrum glabrum* E. Mey.

**ACANTHACEAE**

*Sclerochiton harveyanus* Nees

*Isoglossa* sp.

**RUBIACEAE**

*Tarenna barbertonensis* (Brem.) Brem.

*Tarenna junodii* (Schinz.) Brem.

*Enterospermum littorale* Hiern

*Xeromphis obovata* (Hochst.) Keay

*Xeromphis rudis* (E. Mey. ex Harv.) L. E. Codd

*Rothmannia* sp.

*Tricalysia capensis* (Meissn.) Sim

*Tricalysia lanceolata* (Sond.) Burtt-Davy

*Vangueria chartacea* Robyns

*Vangueria infausta* Burch.

*Lagynias* sp.

*Canthium obovatum* Klotzsch

*Canthium setiflorum* Hiern

*Canthium spinosum* (Klotzsch) Kuntze

*Canthium ventosum* (L.) S. Moore

*Plectroniella armata* (K. Schum.) Robyns

*Pavetta* ? *gerstneri* Brem.

*Coffea racemosa* Lour.

*Psychotria capensis* (Eckl.) Vatke

**CUCURBITACEAE**

*Momordica involucrata* E. Mey. ex Sond.

**CAMPANULACEAE**

*Wahlenbergia undulata* A. DC.

**COMPOSITAE**

*Brachylaena discolor* A. DC.

*Senecio* ? *littoreus* Thunb.



PLATE I, FIG. 1. Dune forest viewed from Lake Sibayi. Note marginal fringe of *Acacia karroo* and Steep slope (Approximately  $30^\circ$ )




PLATE I, FIG. 2. Aerial photograph taken in 1942. Note considerable areas of unstable sand, distributed mainly in the hollows.

Aerial photographs reproduced under Government Printer's copyright authority No. 4084 of 21.11.68.




PLATE 2. Aerial photograph taken in 1964. Note colonization of exposed areas.