

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/49660985>

In vitro antiparasmodial activity and cytotoxicity of nine plants traditionally used in Gabon

Article in *Journal of Ethnopharmacology* · February 2011

DOI: 10.1016/j.jep.2010.11.056 · Source: PubMed

CITATIONS

48

READS

284

10 authors, including:



Jean Bernard Lekana-Douki

International Centre of Medical Research of Franceville

114 PUBLICATIONS 1,400 CITATIONS

SEE PROFILE



jean-bernard Bongui

Universite des Sciences et Techniques de Masuku

13 PUBLICATIONS 373 CITATIONS

SEE PROFILE



Sandrine Lydie Oyegue Liabagui

Ecole Doctorale régionale; Universite des Sciences et Techniques de Masuku

27 PUBLICATIONS 270 CITATIONS

SEE PROFILE



Ulrick Bisvigou

University of Health Sciences in Libreville

63 PUBLICATIONS 529 CITATIONS

SEE PROFILE



In vitro antiplasmodial activity and cytotoxicity of nine plants traditionally used in Gabon

Jean Bernard Lekana-Douki^{a,c,*,1,3}, Jean Bernard Bongui^{b,2}, Sandrine Lydie Oyegue Liabagui^{a,c,1,3}, Sonya Estelle Zang Edou^{a,1}, Rafika Zatra^{a,1}, Ulrich Bisvigou^{a,d,1,3}, Pierre Druilhe^{e,4}, Jacques Lebibi^{b,2}, Fousseyni Samba Toure Ndouo^{a,1}, Maryvonne Kombila^{c,3}

^a Unité de Parasitologie Médicale (UPARAM), Centre International de Recherches Médicales de Franceville (CIRMF), B.P. 769, Franceville, Gabon

^b Unité de Recherches en chimie, Faculté des Sciences, Université des Sciences et Techniques de Masuku, B.P. 943, Franceville, Gabon

^c Département de Parasitologie-Mycologie Médecine Tropicale, Faculté de Médecine, Université des Sciences de la Santé, B.P. 4009, Libreville, Gabon

^d Département de Santé Publique et de Médecine légale et du travail, Faculté de Médecine, Université des Sciences de la Santé, B.P. 4009, Libreville, Gabon

^e Biochemical Parasitology Unit, Institut Pasteur, 25 rue du docteur Roux, 75724 Paris, France

ARTICLE INFO

Article history:

Received 24 September 2010

Received in revised form

15 November 2010

Accepted 23 November 2010

Available online 4 December 2010

Keywords:

Plasmodium falciparum

Plant

Antiplasmodial activity

Cytotoxicity

Gabon

ABSTRACT

Aim of the study: As part of a project to identify new compounds active on malarial parasites, we tested the *in vitro* antiplasmodial activity of nine plants traditionally used to treat malaria symptoms in Haut-Ogooué Province, South-East Gabon.

Materials and methods: Dichloromethane and methanolic extracts of each plant were tested for their antiplasmodial activity on two chloroquine-resistant strains of *Plasmodium falciparum* (FCB and W2), based on lactate dehydrogenase activity. Cytotoxicity was assessed with the MTT test on MRC-5 human diploid embryonic lung cells.

Results: The methanolic extract of *Staudtia gabonensis* and the dichloromethane extract of *Adhatoda latibracteata* showed high antiplasmodial activity ($IC_{50} < 1 \mu\text{g/ml}$) and low cytotoxicity, with selectivity indexes of about 58.25 and 16.43, respectively. The methanolic extract of *Monodora myristica* and the dichloromethane extract of *Afromomum giganteum* also showed promising activity ($1 < IC_{50} < 10 \mu\text{g/ml}$) and low cytotoxicity, with selectivity indexes about 15.70 and 12.48, respectively. Dichloromethane extracts of *Monodora myristica* and *Leonotis Africana* showed moderate activity ($10 < IC_{50} < 40 \mu\text{g/ml}$), with selectivity indexes about 6.07 and 28.89, respectively. Both extracts of *Culcasia lancifolia* had IC_{50} values of 10–40 $\mu\text{g/ml}$ but high cytotoxicity (selectivity indexes < 2.77). The methanolic extract of *Dorstenia klaineana* had moderate antiplasmodial activity (IC_{50} around 17 $\mu\text{g/ml}$) but strong cytotoxicity (0.43 $\mu\text{g/ml}$), giving a selectivity index of about 0.03.

Conclusions: Most extracts of nine selected plants traditionally used to treat malaria in Gabon had interesting antiplasmodial activity *in vitro*. This supports continued investigations of traditional medicines in the search for new antimalarial agents. The compounds responsible for the observed antiplasmodial effects are under investigation.

© 2010 Elsevier Ireland Ltd. All rights reserved.

* Corresponding author at: Département de Parasitologie-Mycologie Médecine Tropicale, Faculté de Médecine Université des Sciences de la Santé, Centre International de Recherches Médicales de Franceville (CIRMF), B.P. 769, Franceville, Gabon. Tel.: +241 06 25 95 90; fax: +241 70 48 96.

E-mail address: Lekana.jb@yahoo.fr (J.B. Lekana-Douki).

¹ Tel.: +241 06 25 95 90.

² Tel.: +241 07 17 33 60.

³ Tel.: +241 70 48 96; fax: +241 70 48 96.

⁴ Tel.: +33 1 45 68 85 78.

1. Introduction

Malaria remains a major public health problem, especially in tropical and subtropical regions, despite new funds for research and control programs to roll back malaria. Two of the main reasons of malaria persistence are increasing insecticide resistance of mosquito vectors and *Plasmodium falciparum* drug resistance (Wellemans and Plowe, 2001). Morbidity and mortality due to malaria have fallen in recent years with the advent of artemisinin-based combination therapy (ACT) and widespread use of impregnated bed nets (World Health Organisation, 2008). However, ACT treatment failures have been reported in some countries (Alker et al., 2007; Holmgren et al., 2007) and artemisinin-resistant malaria parasites

Table 1

Studied plants from Haut-Ogooué, families and traditional medicinal uses.

Botanic name	Family	Voucher number Gabon*	Site of plants collection	Traditional medicinal uses
<i>Adhatoda latibracteata</i>	Acanthaceae	Bongui, No. 001	Epila	Fever, Headache, Vomiting
<i>Aframomum giganteum</i>	Zingiberaceae	Bourboubou, No. 225	Omoi	Fever, Headache, Deworming, Tooth pains
<i>Culcasia lancifolia</i>	Araceae	Bongui, No. 002	Kele	Fever, Headache, Vomiting
<i>Dorstenia klaineana</i>	Moraceae	Bongui, No. 003	Kele	Fever, Headache, Vomiting
<i>Leonotis africana</i>	Lamiaceae	Bongui, No. 004	Mvouna	Headache, Vomiting, Tummyache, Healing, Diarrhea
<i>Monodora myristica</i>	Annonaceae	Sosef M, No. 2022	Epila	Fever, Headache, Vomiting, Deworming, Constipation
<i>Rinorea subintegrifolia</i>	Violaceae	Sosef M, No. 1357	Onkoua	Headache, Rheumatism, Tummyache
<i>Staudtia gabonensis</i>	Myristicaceae	ReitsmaB, No. 1531	Mvouna	Fever, Headache, Healing, Gonorrhea, Rheumatism, Eye's pains
<i>Nauclea latifolia</i>	Rubiaceae	Rei, No. 1648	Maboukou	Fever, Headache, Tummyache

Voucher number Gabon*: name of author who collected the specimen and number of collection order.

were recently detected in Cambodia (Maude et al., 2009), justifying the search for new antimalarial drugs. In addition, the use of these drugs is very limited in some rural areas, where the populations prefer less expensive traditional plant remedies (Newman et al., 2002). Several plants have been shown to possess strong antiparasmodial activity (for review see Kaur et al., 2009). Indeed, the best two current antimalarials (quinine and artemisinin) are both derived from plants.

Several families of phytochemical compounds have antimalarial activity, including alkaloids, terpenes, quassinoids, flavonoids, limonoids, chalcones, xanthonoids and quinoleines (Chhabra et al., 1987; Schwikard and van Heerden, 2002; Batista et al., 2009).

Malaria transmission is hyperendemic and perennial in Gabon, a country with a population of about 1.5 million people. Up to 80% of Gabon is covered by forest, representing a potential source of natural compounds of medical value. Some medicinal plants used in northern and southern Gabon were listed in the 1990s (Akendengue, 1992; Akendengue and Louis, 1994), of which *Uvaria klaineana* has been shown to have antiparasmodial activity (Akendengue et al., 2002).

In this paper, using ethnobotanical information obtained from traditional healers living in Haut-Ogooué Province, south-eastern Gabon, we selected nine Gabonese medicinal plants used to treat malaria symptoms and tested their extracts for *in vitro* antiparasmodial activity and cytotoxicity.

2. Materials and methods

2.1. Plants and extraction procedures

The plants were collected in April 2008 in Haut-Ogooué Province, south-eastern Gabon, where malaria is endemic and the vegetation is a savanna-forest mosaic. The plants were collected in the Plateaux Batéké region composed mainly of shrubland. The families and traditional uses of the nine selected plants are indicated in Table 1.

The protocol of traditional preparation and of use of each tested plant were obtained from notices of traditional healers (unpublished information).

Each species was identified by comparison with voucher specimens deposited at the Gabonese herbarium in Libreville. Plant stems (100 g) were dried for 5 days then crushed and macerated in solvents at room temperature for about 24 h. The material was then extracted first with dichloromethane and then with methanol. The amount of solvent was at least 10 times the volume of plant material. Filtrates were prepared and evaporated to dryness under reduced pressure with a rotary evaporator (Rotavapor®) at 30 °C. Extracts of *Nauclea latifolia* were used as positive controls (Zirihi et al., 2005).

2.2. Parasite culture

Plasmodium falciparum strains FCB and W2 (both chloroquine-resistant) from MR4® (Malaria Research and Reference Reagent Resource Center) were grown under standard conditions as previously described (Trager and Jensen, 1976). The parasites were synchronized by repeated 5% sorbitol treatment. The plant extracts were dissolved in 100 µL of DMSO at an initial concentration of 200 mg/ml and then serially diluted with culture medium before being added to synchronous parasite cultures. The concentration range was 500–0.05 µg/ml. Two hundred microliters of synchronized trophozoite suspension (1.5% final hematocrit in RPMI 1640 + 0.5% Albumax®) was incubated in triplicate with the different concentrations of plant extracts in 96-well flat-bottom plates (NUNC, VWR International, Strasbourg, France). The plates were then placed at 37 °C as previously described (Douki et al., 2003) for 42 h, before being frozen at –20 °C for 3 h to stop parasite growth.

Dihydroartemisinin (Sigma–Aldrich) and chloroquine were used as positive and negative controls of parasite growth inhibition in all experiments.

2.3. Antiparasmodial activity

Antiparasmodial activity was analyzed by measuring *Plasmodium* lactate dehydrogenase (pLDH) activity with a commercial ELISA method as recommended by the manufacturer (ELISA-Malaria antigen test; DiaMed AG) (Kaddouri et al., 2006). The ELISA plates are coated with a monoclonal antibody (MAb) against pan-*Plasmodium* LDH. The absorbance of each well of plate was read with a microplate spectrophotometer (LP400; Bio-Rad) at 450 nm, with a reference wavelength of 620 nm. All experiments were performed at least in triplicate.

The results were expressed as the mean IC₅₀ (the drug concentration that reduced parasitaemia to 50%).

2.4. Cytotoxicity assay

The cytotoxicity of the extracts was assessed with MRC-5 human diploid embryonic lung cells, using a tetrazolium salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Sigma®) colorimetric method based on reagent cleavage by mitochondrial dehydrogenase in viable cells (Mosmann, 1983). Briefly, 5000 cells per well were seeded in 96-well microplates in culture medium (DMEM + 10% inactivated SVF + 2 mM L-glutamine + penicillin/streptomycin/neomycin (0.5/0.5/1 µg/ml)). After 24 h, the cells were washed and incubated with eight concentrations (from 500 to 0.05 µg/ml) of each extract for 7 days at 37 °C in 5% CO₂-air. Cytotoxicity was scored as the percentage reduction in absorbance at 540 nm versus untreated

Table 2*In vitro* antiparasmodial activity, cytotoxicity and selectivity index of the selected plant extracts.

Plants species	Extract	Antiplasmodial activity (IC ₅₀ , µg/ml)*		Cytotoxicity MRC-5 (LD ₅₀ , µg/ml)*	Selectivity index ^a	
		FCB	W2		FCB	W2
<i>Adhatoda latibracteata</i>	CH ₂ Cl ₂	0.7 ± 0.2	1.6 ± 0.1	11.5 ± 2.4	16.43	7.19
	CH ₃ OH	NT	NT	NT	ND	ND
<i>Aframomum giganteum</i>	CH ₂ Cl ₂	8.3 ± 2.7	13.5 ± 1.9	168.5 ± 31.8	20.30	12.48
	CH ₃ OH	410 ± 25	460 ± 75	47.2 ± 9.3	0.12	0.10
<i>Culcasia lancifolia</i>	CH ₂ Cl ₂	8.9 ± 1.6	10.0 ± 2.0	10.25 ± 3.1	1.15	1.3
	CH ₃ OH	25.0 ± 0.9	16.0 ± 2.5	44.3 ± 10.3	1.77	2.77
<i>Dorstenia klaineana</i>	CH ₂ Cl ₂	16.7 ± 5.3	17.0 ± 3.8	0.43 ± 0.2	0.03	0.03
	CH ₃ OH	30.9 ± 2.8	33.7 ± 4.1	29.3 ± 3.7	0.95	0.87
<i>Leonotis africana</i>	CH ₂ Cl ₂	27.1 ± 7.2	15.2 ± 4.0	439.1 ± 7.5	16.20	28.89
	CH ₃ OH	NT	NT	NT	ND	ND
<i>Monodora myristica</i>	CH ₂ Cl ₂	25.0 ± 4.3	28.1 ± 1.5	170.5 ± 30.6	6.82	6.07
	CH ₃ OH	5.5 ± 2.0	6.1 ± 1.5	95.7 ± 12.3	17.4	15.7
<i>Rinorea subintegrifolia</i>	CH ₂ Cl ₂	NT	NT	NT	ND	ND
	CH ₃ OH	360.7 ± 46.9	408.0 ± 64.0	520 ± 21.7	1.44	1.27
<i>Staudtia gabonensis</i>	CH ₂ Cl ₂	8.1 ± 1.5	15.5 ± 2.6	123.2 ± 10.6	15.20	7.95
	CH ₃ OH	0.8 ± 0.4	0.8 ± 0.4	46.6 ± 9.5	58.25	58.25
<i>Nauclea latifolia</i>	CH ₂ Cl ₂	6.6 ± 2.4	8.0 ± 2.7	75.5 ± 12.9	11.44	9.44
	CH ₃ OH	17.1 ± 2.8	18.4 ± 1.9	56.5 ± 23.5	3.30	3.07
<i>Chloroquine</i>		0.6 ± 0.4	0.4 ± 0.2	ND	ND	ND
<i>Artemisinin</i>		0.04 ± 0.01	0.002	ND	ND	ND

NT: not tested; ND: not determined. (IC₅₀, µg/ml)* = Inhibition concentration 50% ± standard deviation. (LD₅₀, µg/ml)** = Lethal dose 50% ± standard deviation.^a Selectivity index = LD₅₀/IC₅₀.

control cultures. All experiments were performed at least in triplicate.

The results were expressed as the mean lethal dose 50 (LD₅₀ = the drug concentration that reduced the number of viable cells by 50%).

A selectivity index (SI), corresponding to the ratio between the cytotoxic and antiparasitic activities of each plant extract, was calculated as follows:

$$SI \text{ Plasmodium} = \frac{LD_{50} \text{ MRC-5}}{IC_{50} \text{ Plasmodium}}$$

3. Results and discussion

We investigated the *in vitro* activity of extracts of nine plants on two *Plasmodium falciparum* strains (FCB and W2). The plants were selected because their antiparasmodial activities had not previously been described. We have not tested leaf extracts because of excessive chlorophyll levels, which hamper the isolation of other compounds. Based on WHO guidelines and previous data (Jonville et al., 2008) antiparasmodial activity was classified as follows: high (IC₅₀ < 5 µg/ml), promising (5 < IC₅₀ < 15 µg/ml), moderate (15 < IC₅₀ < 50 µg/ml) and inactive (IC₅₀ > 50 µg/ml). Only two extracts showed high antiparasmodial activity, while seven showed promising activity. Although methanolic extracts of some plants were more active than the corresponding dichloromethane extracts, the latter tended to be more active overall, and also more selective (Table 2).

Although the traditional healers used some plants to treat malaria symptoms these had no antiparasmodial activity *in vitro*. It should be noted that we only tested antiparasmodial activity on the asexual erythrocytic stage of *Plasmodium falciparum*, and the extracts that we found inactive might possibly inhibit other parasite stages. Alternatively, some plants without *in vitro* antiparasmodial activity may stimulate the immune response; this is the case of *Markhamia lutea*, which inhibits *Plasmodium berghei* growth in mice but is inactive *in vitro* (Hakizamungu and Weri, 1988).

3.1. *Adhatoda latibracteata*

People in Haut-Ogooué use this plant to treat fever, headache and vomiting; the stems and leaves are boiled and then drunk, while crushed leaves are applied to the head and stomach.

The dichloromethane extract of *Adhatoda latibracteata* had very high antimalarial activity, with IC₅₀ values of 0.7 ± 0.2 and 1.6 ± 0.1 µg/ml on strains FCB and W2, respectively. It was weakly cytotoxic, yielding selectivity indexes of 16.43 and 7.19. *Adhatoda vasica*, a specie of the same genus containing certain alkaloids, has been shown to be active against fever, cestodal disease, respiratory tract infections, and airway inflammation (Claeson et al., 2000; Sharafkhaneh et al., 2007; Yadav and Tangpu, 2008). It was recently reported that *Adhatoda justicia*, other member of the *Adhatoda* genus, is used to treat bronchitis in India (Rajakumar and Shivanna, 2009).

Other members of the Acanthaceae family, such as *Andrographis paniculata*, *Hypoestes rosea* and *Clinacanthus siamensis*, contain terpenes and xanthones with antiparasmodial activity (Tuntiwachwuttikul et al., 2003; Ojo-Amaize et al., 2007; Mishra et al., 2009). Thus, compounds with antiparasmodial activity found in Acanthaceae might also be present in *Adhatoda latibracteata*.

3.2. *Aframomum giganteum*

This plant is used as a boiled leaf decoction by traditional healers in Haut-Ogooué to treat headache and fever. It is used as a deworming and purgative agent. It is also used to treat tooth pains. The dichloromethane extract of *Aframomum giganteum* exhibited promising and selective antiparasmodial activity, with IC₅₀ values of 8.3 ± 2.7 and 13.5 ± 1.9 µg/ml on strains FCB and W2, respectively, and selectivity indexes of about 20.30 and 12.48. In contrast, methanolic extracts had no antimalarial activity (IC₅₀ = 410 ± 25 and 460 ± 75 µg/ml) and were poorly selective (SI 0.12 and 0.10), suggesting that most of the compounds responsible for the antiparasmodial activity of *Aframomum giganteum* are extracted in dichloromethane.

Aframomum giganteum is an aromatic plant containing several terpenes (De Bernardi et al., 1981) that are very active on asexual blood stages of *Plasmodium falciparum*. This plant is a potential

source of a new antimalarial drug. The genus *Aframomum* includes several species with strong antiplasmodial activity (Duker-Eshun et al., 2002; Kenmogne et al., 2006) mediated by flavonoids, diaryl heptanoids and sesquiterpenes (De Bernardi et al., 1981; Kamnaing et al., 2003). Our results are consistent with the use of this plant for traditional medicine in Haut-Ogooué.

3.3. *Culcasia lancifolia*

This is a highly perfumed plant used by women as a cosmetics. It is also used by traditional healers to treat malaria symptoms (headache, fever and vomiting) consistently with our *in vitro* data. The dichloromethane extract of *Culcasia lancifolia* had promising antimalarial activity, with IC_{50} values of 8.9 ± 1.6 and 10.0 ± 2.0 $\mu\text{g/ml}$ on strains FCB and W2, respectively, although its selectivity was limited (1.15 and 1.03, respectively). The methanolic extract of *Culcasia lancifolia* had moderate antimalarial activity. Several sesquiterpenes have been isolated from *Culcasia scandens*, another member of the *Culcasia* genus which has analgesic and anti-inflammatory properties (Okoli and Akah, 2000, 2004; Okoli et al., 2006), but has not previously been studied for its antiplasmodial activity.

3.4. *Dorstenia klaineana*

Dorstenia klaineana (synonymous *Dorstenia klainei*) is used to treat malaria symptoms in Gabon. Some traditional healers make an infusion containing *Dorstenia klaineana* and *Leonotis africana* to treat malarial symptoms. This plant is also used in aromatic perfume in spiritual ceremonies. We found that its extracts had no antiplasmodial activity and were strongly toxic for cultured human cells: the dichloromethane extract had IC_{50} values of 16.7 ± 5.3 and 17.0 ± 3.8 $\mu\text{g/ml}$ on strains FCB and W2, respectively, and selectivity indexes of about 0.03, while the methanolic extract had IC_{50} values of 30.9 ± 2.8 and 33.7 ± 4.1 $\mu\text{g/ml}$ FCB and W2, respectively, and poor selectivity indexes (0.95 and 0.87). This is consistent with a recent report that methanolic extracts of *Dorstenia klaineana* have no antiplasmodial activity, although such activity was found after fractionation with hexane–ethyl acetate (Boyom et al., 2009). Other members of the Moraceae family (*Artocarpus rigidus* BLUME subsp. *Rigidus* and *Ficus fistulosa* Reinw.) exhibit antiplasmodial activity driven by the flavonoid artonin F, the flavonoid 7-demethylartanol E, and the flavonoid cycloartobiloxanthone, Verrucaridin L acetate (Zhang et al., 2002; Namdaung et al., 2006).

3.5. *Leonotis africana*

Rural populations in Haut-Ogooué Province use this plant as a healing agent after circumcision. Another use of this plant is in the contraction of loving pact between lovers. The leaves of this plant are used against malaria symptoms, in the form of infusions and baths. The same plant is used to treat diarrhea too. We found that the dichloromethane extract of *Leonotis africana* had moderate antimalarial activity (IC_{50} 27.1 ± 7.2 and 15.2 ± 4.0 $\mu\text{g/ml}$ on strains FCB and W2, respectively), and weak cytotoxicity (LD_{50} 439.1 ± 7.5 $\mu\text{g/ml}$), giving selectivity indexes of 16.20 and 28.89. This plant is a member of the Lamiaceae family. It was recently reported that other Lamiaceae are used in traditional medicine in the Yanasha region of Peru. We found that *Leonotis Africana* exhibited moderate but selective antiplasmodial activity, in keeping with the reported activity of Lamiaceae from Yanasha (Valadeau et al., 2009). The Lamiaceae family may thus be a future source of antiplasmodial compounds, as several antiplasmodial terpenes and flavonoids have been isolated from these species (van Baren et al., 2006; Agnihotri et al., 2008; HENCHIRI et al., 2009). One, *Plectran-*

thus amboinicus, was found to have antiplasmodial activity in albino mice (Periyanayagam et al., 2008).

3.6. *Monodora myristica*

In Haut-Ogooué region, this member of the Annonaceae is used in an infusion containing *Nauclea latifolia* too, to treat malaria symptoms. Some people used this against constipation. To treat headache leaves of *Monodora myristica* are applied on the face. It is also used in Western Africa as a deworming agent. We found that the dichloromethane extract exhibited moderate antimalarial activity (IC_{50} of 25.0 ± 4.3 and 28.1 ± 1.5 $\mu\text{g/ml}$ on strains FCB and W2, respectively) in keeping with previously published data (Okpekon et al., 2004), and also weak cytotoxicity (LD_{50} of 170.5 ± 30.6 $\mu\text{g/ml}$), giving selectivity indexes of 6.82 and 6.07. In contrast, the methanolic extract of *Monodora myristica* had high antimalarial activity (IC_{50} of 5.5 ± 2.0 and 6.1 ± 1.5 $\mu\text{g/ml}$ on strains FCB and W2, respectively) and weak cytotoxicity (LD_{50} 95.7 ± 12.3 $\mu\text{g/ml}$), giving good selectivity indexes (17.40 and 15.69). Terpenes, alkaloids and other families of compounds with antiplasmodial activity have been isolated from several members of the Annonaceae (Mahiou et al., 2000; Boyom et al., 2003; Kanokmedhakul et al., 2007). The presence of such compounds in *Monodora myristica* could explain the antiplasmodial activity observed here.

3.7. *Staudtia gabonensis*

The methanolic extract of *Staudtia gabonensis* showed high antimalarial activity (IC_{50} of 0.8 ± 0.4 and 0.8 ± 0.6 $\mu\text{g/ml}$ on strains FCB and W2, respectively) and the best selectivity indexes (58.25 for both strains). The dichloromethane extract showed promising antimalarial activity (IC_{50} of 8.1 ± 1.5 and 15.5 ± 2.6 $\mu\text{g/ml}$ on strains FCB and W2, respectively) and high selectivity indexes of 15.20 and 7.95. *Staudtia gabonensis* is also used in other regions of Gabon, and notably in Libreville region, where it is combined with *Enantia chlorantha* Oliv or *Combretodendron africanum* Exell. in an infusion used to treat malaria (Lamidi et al., 2002). This plant is also used as healing agent because of its red sap. Eye pain, gonorrhea and articular rheumatism are also treated by the decoction of leaves of this plant. This plant is a member of the Myristicaceae family, other members of which also exhibit antiplasmodial activity. Several compounds with antiplasmodial properties have been isolated from Myristicaceae (Ancolio et al., 2002; Abrantes et al., 2008; Rangkaew et al., 2009). Whereas we found that the methanolic extract was most active, dichloromethane extracts and lignans from *Pycnanthus angolensis* have been reported to have strong antiplasmodial activity (Abrantes et al., 2008).

3.8. *Rinorea subintegrifolia*

This plant is used by traditional healer in decoction with lemons to treat malaria symptoms. But, it is also used as an aroma agent during ancestral cults because of its smell. We only tested a methanolic extract of *Rinorea subintegrifolia* because insufficient material was collected. This extract was not active against *Plasmodium falciparum* *in vitro*. The activity of this plant has not previously been tested, although *Viola websteri* Hemsl, another member of the Violaceae family, has antiplasmodial activity and active compounds have been isolated from it (Chung et al., 2009; Lee et al., 2009).

3.9. *Nauclea latifolia*

Some previous data have showed the antiplasmodial activity of this plant. The dichloromethane extract of *Nauclea latifolia* showed promising antimalarial activity (IC_{50} 6.6 ± 2.4 and 8.0 ± 2.7 $\mu\text{g/ml}$

on strains FCB and W2, respectively) with selectivity indexes of 11.44 and 9.44. The methanolic extract showed moderate anti-malarial activity (IC_{50} 17.1 ± 2.8 and 18.4 ± 1.9 μ g/ml on strains FCB and W2, respectively) but was poorly selective (selectivity indexes both around 3.3). These results are consistent with other published data from central West Africa (Traore-Keita et al., 2000; Zirihi et al., 2005). Although the active components have yet to be identified, this plant shows hypoglycemic, anticonvulsant, anxiolytic, antinociceptive, anti-inflammatory and anti-pyretic properties that justify its use to treat malaria symptoms (Gidado et al., 2008; Ngo Bum et al., 2009; Abbah et al., 2010). Compounds with antifungal properties have also been isolated from *Nauclea latifolia* (Ata et al., 2009).

4. Conclusion

Overall, the *in vitro* activities of these Gabonese plant extracts are compatible with their use as traditional remedies for malaria. We are currently attempting to isolate compounds with antiparasitological activity from the most promising of these plants, prior to testing in animal models of malaria.

Acknowledgments

We thank the traditional practitioners who collaborated with us by showing us the traditional plants; J. Mbandza who identified the plants; and F. Lekoulou who provided technical assistance. MR4 and ATCC kindly provided the parasite reference strains and MRC-5 cells. CIRMF supported the biological analyses. CIRMF is supported by the Gabonese government, Total, and the French Foreign Ministry.

References

- Abbah, J., Amos, S., Chindo, B., Ngazal, I., Vongtau, H.O., Adzu, B., Farida, T., Odutola, A.A., Wambele, C., Gamaniel, K.S., 2010. Pharmacological evidence favouring the use of *Nauclea latifolia* in malaria ethnopharmacy: effects against nociception, inflammation, and pyrexia in rats and mice. *Journal of Ethnopharmacology* 127, 85–90.
- Abrantes, M., Mil-Homens, T., Duarte, N., Lopes, D., Cravo, P., Madureira, M., Ferreira, M.J., 2008. Antiplasmodial activity of lignans and extracts from *Pycnanthus angolensis*. *Planta Medica* 74, 1408–1412.
- Agnihotri, V.K., Elshohly, H.N., Smillie, T.J., Khan, I.A., Walker, L.A., 2008. New labdane diterpenes from *Leonurus cardiaca*. *Planta Medica* 74, 1288–1290.
- Akendengue, B., 1992. Medicinal plants used by the Fang traditional healers in Equatorial Guinea. *Journal of Ethnopharmacology* 37, 165–173.
- Akendengue, B., Louis, A.M., 1994. Medicinal plants used by the Masango people in Gabon. *Journal of Ethnopharmacology* 41, 193–200.
- Akendengue, B., Ngou-Milama, E., Roblot, F., Laurens, A., Hocquemiller, R., Grelhier, P., Frappier, F., 2002. Antiplasmodial activity of *Uvaria klaineana*. *Planta Medica* 68, 167–169.
- Alker, A.P., Lim, P., Sem, R., Shah, N.K., Yi, P., Bouth, D.M., Tsuyuoka, R., Maguire, J.D., Fandeur, T., Arie, F., Wongsrichanalai, C., Meshnick, S.R., 2007. Pfmdr1 and *in vivo* resistance to artesunate-mefloquine in falciparum malaria on the Cambodian–Thai border. *American Journal of Tropical Medicine and Hygiene* 76, 641–647.
- Ancolio, C., Azas, N., Mahiou, V., Ollivier, E., Di Giorgio, C., Keita, A., Timon-David, P., Balansard, G., 2002. Antimalarial activity of extracts and alkaloids isolated from six plants used in traditional medicine in Mali and Sao Tome. *Phytotherapy Research* 16, 646–649.
- Ata, A., Udenigwe, C.C., Matochko, W., Holloway, P., Eze, M.O., Uzoegwu, P.N., 2009. Chemical constituents of *Nauclea latifolia* and their anti-GST and anti-fungal activities. *Natural products communications* 4, 1185–1188.
- Batista, R., Silva Ade Jr., J., de Oliveira, A.B., 2009. Plant-derived antimalarial agents: new leads and efficient phytomedicines. Part II. Non-alkaloidal natural products. *Molecules* 14, 3037–3072.
- Boyom, F.F., Kemgne, E.M., Tpongning, R., Ngouana, V., Mbacham, W.F., Tsamo, E., Zollo, P.H., Gut, J., Rosenthal, P.J., 2009. Antiplasmodial activity of extracts from seven medicinal plants used in malaria treatment in Cameroon. *Journal of Ethnopharmacology* 123, 483–488.
- Boyom, F.F., Ngouana, V., Zollo, P.H., Menut, C., Bessiere, J.M., Gut, J., Rosenthal, P.J., 2003. Composition and anti-plasmodial activities of essential oils from some Cameroonian medicinal plants. *Phytochemistry* 64, 1269–1275.
- Chhabra, S.C., Mahunnah, R.L., Mshiu, E.N., 1987. Plants used in traditional medicine in eastern Tanzania. I. Pteridophytes and angiosperms (Acanthaceae to Canelaceae). *Journal of Ethnopharmacology* 21, 253–277.
- Chung, I.M., Seo, S.H., Kang, E.Y., Park, W.H., Moon, H.I., 2009. Anti-malarial activity of 6-(8-Z-pentadecenyl)-salicylic acid from *Viola websteri* in mice. *Malaria Journal* 8, 151.
- Claeson, U.P., Malmfors, T., Wikman, G., Bruhn, J.G., 2000. Adhatoda vascica: a critical review of ethnopharmacological and toxicological data. *Journal of Ethnopharmacology* 72, 1–20.
- De Bernardi, M., Mellerio, G., Paternoster-Colombo, M., Vidari, G., Vita-Finzi, P., 1981. Constituents of essential oil of *Aframomum giganteum*. *Planta Medica* 41, 359–365.
- Douki, J.B., Sterkers, Y., Lepolard, C., Traore, B., Costa, F.T., Scherf, A., Gysin, J., 2003. Adhesion of normal and *Plasmodium falciparum* ring-infected erythrocytes to endothelial cells and the placenta involves the rhoptry-derived ring surface protein-2. *Blood* 101, 5025–5032.
- Duker-Eshun, G., Jaroszewski, J.W., Asomaning, W.A., Oppong-Boachie, F., Olsen, C.E., Christensen, S.B., 2002. Antiplasmodial activity of labdanes from *Aframomum latifolium* and *Aframomum sceptrum*. *Planta Medica* 68, 642–644.
- Gidado, A., Ameh, D.A., Atawodi, S.E., Ibrahim, S., 2008. Hypoglycaemic activity of *Nauclea latifolia* Sm. (Rubiaceae) in experimental animals. *African Journal of Traditional, Complementary and Alternative Medicine* 5, 201–208.
- Hakizamungu, E., Weri, M., 1988. L'usage de plantes médicinales dans le traitement du paludisme en médecine traditionnelle Rwandaise. *Bulletin de Médecine et Pharmacie* 2, 11–17.
- Henchiri, H., Bodo, B., Deville, A., Dubost, L., Zourgui, L., Raies, A., Grelhier, P., Mambu, L., 2009. Sesquiterpenoids from *Teucrium ramosissimum*. *Phytochemistry* 70, 1435–1441.
- Holmgren, G., Hamrin, J., Svard, J., Martensson, A., Gil, J.P., Bjorkman, A., 2007. Selection of pfmdr1 mutations after amodiaquine monotherapy and amodiaquine plus artemisinin combination therapy in East Africa. *Infection, Genetics and Evolution* 7, 562–569.
- Jonville, M.C., Kodja, H., Humeau, L., Fournel, J., De Mol, P., Cao, M., Angenot, L., Frederich, M., 2008. Screening of medicinal plants from Reunion Island for anti-malarial and cytotoxic activity. *Journal of Ethnopharmacology* 120, 382–386.
- Kaddouri, H., Nakache, S., Houze, S., Mentre, F., Le Bras, J., 2006. Assessment of the drug susceptibility of *Plasmodium falciparum* clinical isolates from Africa by using a *Plasmodium lactate* dehydrogenase immunodetection assay and an inhibitory maximum effect model for precise measurement of the 50-percent inhibitory concentration. *Antimicrobial Agents and Chemotherapy* 50, 3343–3349.
- Kammaing, P., Tsopmo, A., Tanifum, E.A., Tchuendem, M.H., Tane, P., Ayafor, J.F., Sterner, O., Rattendi, D., Iwu, M.M., Schuster, B., Bacchi, C., 2003. Trypanocidal diarylheptanoids from *Aframomum lesteuianum*. *Journal of Natural Products* 66, 364–367.
- Kanokmedhakul, S., Kanokmedhakul, K., Lekphrom, R., 2007. Bioactive constituents of the roots of *Polyalthia cerasoides*. *Journal of Natural Products* 70, 1536–1538.
- Kaur, K., Jain, M., Kaur, T., Jain, R., 2009. Antimalarials from nature. *Bioorganic & Medicinal Chemistry* 17, 3229–3256.
- Kenmogne, M., Prost, E., Harakat, D., Jacquier, M.J., Frederich, M., Sondengam, L.B., Zeche, M., Waffo-Teguo, P., 2006. Five labdane diterpenoids from the seeds of *Aframomum zambesiaceum*. *Phytochemistry* 67, 433–438.
- Lamidi, M., Eyle, M., Mba, C., Nze-Ekekang, L.B.G., 2002. Inquiry next to the traditional healers in three regions of Gabon (Des sources du savoir aux médicaments du futur), pp. 289–291.
- Lee, S.J., Park, W.H., Moon, H.I., 2009. Bioassay-guided isolation of antiparasitological anacardic acids derivatives from the whole plants of *Viola websteri* Hemsl. *Parasitology Research* 104, 463–466.
- Mahiou, V., Roblot, F., Fournet, A., Hocquemiller, R., 2000. Bisbenzylisoquinoline alkaloids from *Guatteria boliviana* (Annonaceae). *Phytochemistry* 54, 709–716.
- Maude, R.J., Pontavornpinyo, W., Saralamba, S., Aguas, R., Yeung, S., Dondorp, A.M., Day, N.P., White, N.J., White, L.J., 2009. The last man standing is the most resistant: eliminating artemisinin-resistant malaria in Cambodia. *Malaria Journal* 8, 31.
- Mishra, K., Dash, A.P., Swain, B.K., Dey, N., 2009. Anti-malarial activities of *Andropogon paniculata* and *Hedyotis corymbosa* extracts and their combination with curcumin. *Malaria Journal* 8, 26.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* 65, 55–63.
- Namdaung, U., Aroonrerk, N., Suksamrarn, S., Danwisetkanjana, K., Saenboonrueng, J., Arjchomphu, W., Suksamrarn, A., 2006. Bioactive constituents of the root bark of *Artocarpus rigidus* subsp. *rigidus*. *Chemical & Pharmaceutical Bulletin (Tokyo)* 54, 1433–1436.
- Newman, D.J., Cragg, G.M., Holbeck, S., Sausville, E.A., 2002. Natural products and derivatives as leads to cell cycle pathway targets in cancer chemotherapy. *Current Cancer Drug Targets* 2, 279–308.
- Ngo Bum, E., Taiwe, G.S., Moto, F.C., Ngoupaye, G.T., Nkanchoua, G.C., Pelanken, M.M., Rakotonirina, S.V., Rakotonirina, A., 2009. Anticonvulsant, anxiolytic, and sedative properties of the roots of *Nauclea latifolia* Smith in mice. *Epilepsy Behaviour* 15, 434–440.
- Ojo-Amaize, E.A., Nchekwube, E.J., Cottam, H.B., Oyemade, O.A., Adesomoku, A.A., Okogun, J.I., 2007. *Plasmodium berghei*: antiparasitic effects of orally administered hypoxostoxide in mice. *Experimental Parasitology* 117, 218–221.

- Okoli, C.O., Akah, P.A., 2000. A pilot evaluation of the anti-inflammatory activity of *Culcasia scandens*, a traditional antirheumatic agent. *Journal of Alternative and Complementary Medicine* 6, 423–427.
- Okoli, C.O., Akah, P.A., 2004. Mechanisms of the anti-inflammatory activity of the leaf extracts of *Culcasia scandens* P. Beauv (Araceae). *Pharmacology Biochemistry and Behavior* 79, 473–481.
- Okoli, C.O., Akah, P.A., Egbuniwe, O.N., 2006. Analgesic activity of leaf extracts of *Culcasia scandens* P. Beauv. *Indian Journal of Experimental Biology* 44, 422–424.
- Okpekon, T., Yolou, S., Gleye, C., Roblot, F., Loiseau, P., Bories, C., Grellier, P., Frappier, F., Laurens, A., Hocquemiller, R., 2004. Antiparasitic activities of medicinal plants used in Ivory Coast. *Journal of Ethnopharmacology* 90, 91–97.
- Periyannayagam, K., Nirmala Devi, K., Suseela, L., Uma, A., Ismail, M., 2008. In vivo antimalarial activity of leaves of *Plectranthus amboinicus* (Lour) Spreng on *Plasmodium berghei yoelii*. *Journal of Communicable Diseases* 40, 121–125.
- Rajakumar, N., Shivanna, M.B., 2009. Ethno-medicinal application of plants in the eastern region of Shimoga District, Karnataka, India. *Journal of Ethnopharmacology* 126, 64–73.
- Rangkaew, N., Suttisri, R., Moriyasu, M., Kawanishi, K., 2009. A new acyclic diterpene acid and bioactive compounds from *Knema glauca*. *Archives of Pharmacological Research* 32, 685–692.
- Schwikkard, S., van Heerden, F.R., 2002. Antimalarial activity of plant metabolites. *Natural Product Reports* 19, 675–692.
- Sharafkhaneh, A., Velamuri, S., Badmaev, V., Lan, C., Hanania, N., 2007. The potential role of natural agents in treatment of airway inflammation. *Therapeutic Advances in Respiratory Disease* 1, 105–120.
- Trager, W., Jensen, J.B., 1976. Human malaria parasites in continuous culture. *Science* 193, 673–675.
- Traore-Keita, F., Gasquet, M., Di Giorgio, C., Ollivier, E., Delmas, F., Keita, A., Doumbo, O., Balansard, G., Timon-David, P., 2000. Antimalarial activity of four plants used in traditional medicine in Mali. *Phytotherapy Research* 14, 45–47.
- Tuntiwachwuttikul, P., Pootaeng-on, Y., Pansa, P., Srisanpang, T., Taylor, W.C., 2003. Sulfur-containing compounds from *Clinacanthus siamensis*. *Chemical & Pharmaceutical Bulletin (Tokyo)* 51, 1423–1425.
- Valadeau, C., Pabon, A., Deharo, E., Alban-Castillo, J., Estevez, Y., Lores, F.A., Rojas, R., Gamboa, D., Sauvain, M., Castillo, D., Bourdy, G., 2009. Medicinal plants from the Yanesha (Peru): evaluation of the leishmanicidal and antimalarial activity of selected extracts. *Journal of Ethnopharmacology* 123, 413–422.
- van Baren, C., Anao, I., Leo Di Lira, P., Debenedetti, S., Houghton, P., Croft, S., Martino, V., 2006. Triterpenic acids and flavonoids from *Satureja parvifolia*. Evaluation of their antiprotozoal activity. *Zeitschrift Fur Naturforschung C* 61, 189–192.
- Wellems, T.E., Plowe, C.V., 2001. Chloroquine-resistant malaria. *Journal of Infectious Diseases* 184, 770–776.
- World Health Organisation, W., 2008. Malaria Report 2008. WHO Report.
- Yadav, A.K., Tangpu, V., 2008. Anticestodal activity of *Adhatoda vasica* extract against *Hymenolepis diminuta* infections in rats. *Journal of Ethnopharmacology* 119, 322–324.
- Zhang, H.J., Tamez, P.A., Aydogmus, Z., Tan, G.T., Saikawa, Y., Hashimoto, K., Nakata, M., Hung, N.V., Xuan, T., Cuong, N.M., Soejarto, D.D., Pezzuto, J.M., Fong, H.H., 2002. Antimalarial agents from plants. III. Trichothecenes from *Ficus fistulosa* and *Rhaphidophora decursiva*. *Planta Medica* 68, 1088–1091.
- Zirih, G.N., Mambu, L., Guede-Guina, F., Bodo, B., Grellier, P., 2005. In vitro antiplasmodial activity and cytotoxicity of 33 West African plants used for treatment of malaria. *Journal of Ethnopharmacology* 98, 281–285.