Exercise Sheet 5

Discrete Mathematics by Hongfei Fu, 2023.9.28

Name:_________ Student ID:_________ Class: ____________

Note:The following exercises involve the logical operators $NAND$ and NOR. The proposition $p\ NAND\ q$ is true when either p or q, or both, are false; and it is false when both p and q are true. The proposition $p\ NOR\ q$ is true when both p and q are false, and it is false otherwise. The propositions $p\ NAND\ q$ and $p\ NOR\ q$ are denoted by $p|q$ and $p \downarrow q$, respectively. (The operators — and $\downarrow$ are called the Sheffer stroke and the Peirce arrow after H. M. Sheffer and C. S. Peirce, respectively.)

1. In this exercise we will show that $\{\downarrow\}$ is a functionally complete collection of logical operators.

   a) Show that $p \downarrow p$ is logically equivalent to $\neg p$.

   b) Show that $(p \downarrow q) \downarrow (p \downarrow q)$ is logically equivalent to $p \vee q$.

   c) Conclude from parts (a) and (b), and Exercise 49 , that $\{\downarrow\}$ is a functionally complete collection of logical operators.

   Note::Show that $p \downarrow q$ is logically equivalent to $\neg(p \vee q)$

   Answer Area:

2. Show that $p|(q|r)$ and $(p|q)|r$ are not equivalent, so that the logical operator $|$ is not associative.
   Answer Area:

3. Put these statements in prenex normal form, where the domain of all variables are the same and nonempty. [Hint: Use logical equivalence from Tables 6 and 7 in Section 1.3, Table 2 in Section 1.4, Example 19 in Section 1.4, Exercises 45 and 46 in Section 1.4, and Exercises 48 and 49.]
b) $\neg(\forall x P(x) \vee \forall x Q(x))$
c) $(\exists x P(x)) \rightarrow (\exists x Q(x))$

Answer Area:

4. Consider the compound proposition $\phi = p \rightarrow (q \oplus r)$ where p, q, r are propositional variables.

   (a) Give a disjunctive normal form of ϕ.

   (b) Give a conjunctive normal form of ϕ.

Answer Area: