# Internal Explanatory Paper:

## The Evolution of the GLLM Framework

From Raw Geometry to Emergent Gauge Structure and Curvature

The Geometrica Team

February 13, 2026

## 1 Where It Began — Pure Geometry, No Physics Assumed

The Geometrica Large Language Model (GLLM) did not begin as a particle theory, nor as a modification of the Standard Model. It began with abstract relational geometry. The earliest stage of the program examined seven families of high-dimensional relational bundles—purely combinatorial connectivity structures defined by adjacency alone.

There was no metric, no coordinates, no background spacetime, and no assumed gauge group. These were simply transport networks: relational node structures capable of supporting paths. The initial question was: *What must survive if we impose only transport admissibility?*

This filtering process (Phases 1–4) eliminated unstable relational structures and revealed persistent geometric features. At this stage, the project was discovering which relational configurations remain coherent under structural transport constraints.

## 2 From Geometry to Operator Structure (Part I)

As transport constraints were enforced, ordering forced algebraic consistency, and spinorial structure appeared. The complex phase $i$ emerged from internal transport closure.

### 2.1 The Core Construction (Phases 33–36)

The framework derived a global triad–spinor basis (Phase 33), a physical subspace projector $B$ (Phase 35), and a transport operator `A_mode` (Phase 36). From these, the central object was defined:

$$A_{\text{phys}} = B^{\dagger} A_{\text{mode}} B \tag{1}$$

Everything that followed was derived from the operator's internal spectral properties. No Lagrangian or particle ontology was inserted.

### 2.2 Chirality and Generations — Spectral Structure

**Chirality (Phase 39):** A true chirality operator emerged internally as an intrinsic spectral feature.

**Generation Structure (Phase 42):** Spectral analysis revealed three stable eigenmodes aligned with representation stability. A "No-Go" condition for $N = 4$ emerged from spectral collapse.

## 3 Part II — Structural Necessity, Probability, and Law

Part II reframed the program fundamentally. Rather than asking what physics is, the framework addressed: *What must exist if a bounded observer interacts with relational structure?*

- **Probability** as consistency of partial access.
- **Law** as stability under admissible update.
- **Dynamics** as stabilized operator structure.
- **Physics** as stability under structural admissibility constraints.

## 4 Part III — Emergent Gauge Structure

Spectral coarse-graining was performed on $A_{\text{phys}}$ to form $\tilde{A}(\delta)$ by clustering eigenvalues $|\lambda_i - \lambda_j| \leq \delta$.

### 4.1 Emergent $SU(3)$ (Phases 40–42)

At $\delta = 0.10$, a stable 3-dimensional degenerate block emerged. Within that block, $\mathfrak{su}(3)$ closure held exactly on $\tilde{A}$ with a maximum residual of $1.39 \times 10^{-16}$.

### 4.2 Emergent $SU(2)_L$ (Phase 43) and $U(1)$ (Phase 44)

A stable 2D degenerate block closed exactly as $\mathfrak{su}(2)$ on $\tilde{A}$ acting exclusively on L-like states. Phase 44 identified an Abelian commuting direction $H$, allowing for the definition of the mixing family $Q(\theta) = \cos(\theta)H + \sin(\theta)H_3$.

### 4.3 Phase 45 — Mass-Proxy Structure

Mass-like structure appeared as spectral deformation when comparing restricted $A_{\text{phys}}$ against a block-diagonal reference $A_{\text{bd}}$:

$$\text{mass_proxy} = \text{mean}_k |\text{Re}(\lambda_{\text{full},k}) - \text{Re}(\lambda_{\text{bd},k})| \tag{2}$$

## 5 Part IV — Structural Interpretation

Part IV consolidated the interpretation of spacetime as the representational organization of transport. Curvature was defined via the transport commutator structure:

$$R_{\mu\nu} = [T_\mu, T_\nu] = T_\mu T_\nu - T_\nu T_\mu \tag{3}$$

Gravity is interpreted as emergent non-commuting transport generators.

## 6 Independent Python Validation

To eliminate environment dependence, `A_mode` and $B$ were exported to a NumPy/SciPy environment.

1. **SU(3) Emergence Paper:** Confirmed exact $\mathfrak{su}(3)$ closure ($R < 10^{-16}$) on the coarse-grained operator.
2. **Spacetime & Gravity Paper:** Reconstructed transport along triad axes and computed a non-trivial Frobenius norm $\|[T_{e1}, T_{e2}]\|_F \approx 3.117$.

The Python replication confirms these structures are operator-internal and numerically reproducible.

## 7 One Clean Statement

The GLLM evolved from abstract relational bundles to a canonical operator whose internal spectral structure forces chirality, three-generation stability, $SU(3) \times SU(2) \times U(1)$ gauge structure, and curvature via commutator structure—all without assuming fields, particles, or ontological spacetime.