

Differential and Integral Calculus

Aakash Jog

2014-15

Contents

1	Lecturer Information	3
2	Required Reading	3
3	Additional Reading	3
I	Sequences and Series	4
1	Sequences	4
1.1	Limit of a Sequence	6
1.2	Sub-sequences	9
1.3	Cauchy Characterisation of Convergence	10
2	Series	12
2.1	Convergence Criteria	14
2.1.1	Leibniz's Criteria	14
2.1.2	Comparison Test	15
2.1.3	d'Alembert Criteria (Ratio Test)	16
2.1.4	Cauchy Criteria (Cauchy Root Test)	16



This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>.

2.1.5	Integral Test	17
3	Power Series	18
3.1	Differentiation and Integration of Power Series	19
3.2	Taylor Series	20
4	Series of Real-valued Functions	22

1 Lecturer Information

Dr. Yakov Yakubov

Office: Schreiber 233

Telephone: +972 3-640-5357

E-mail: yakubov@post.tau.ac.il

2 Required Reading

Protter and Morrey: *A first Course in Real Analysis*, UTM Series, Springer-Verlag, 1991

3 Additional Reading

Thomas and Finney, *Calculus and Analytic Geometry*, 9th edition, Addison-Wesley, 1996

Part I

Sequences and Series

1 Sequences

Definition 1 (Sequence). A sequence of real numbers is a set of numbers which are written in some order. There are infinitely many terms in a sequence. It is denoted by $\{a_n\}_{n=1}^{\infty}$ or $\{a_n\}$.

Example 1. $1, \frac{1}{2}, \frac{1}{3}, \dots$ is called the harmonic sequence.

$$a_n = \frac{1}{n}$$

Example 2. $1, -\frac{1}{2}, \frac{1}{3}, \dots$ is called the alternating harmonic sequence.

$$a_n = (-1)^{n+1} \frac{1}{n}$$

Example 3. $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots$

$$a_n = \frac{n}{n+1}$$

Example 4. $\frac{2}{3}, \frac{3}{9}, \frac{4}{27}, \dots$

$$a_n = \frac{n+1}{3^n}$$

Example 5. The Fibonacci sequence is given by

$$f_n = \begin{cases} 1 & ; \quad n = 1, 2 \\ f_{n-1} + f_{n-2} & ; \quad n \geq 3 \end{cases}$$

Example 6. A geometric sequence is given by

$$a_n = a_1 q^{n-1}$$

where q is called the common ratio.

Example 7. A geometric sequence is given by

$$a_n = a_1 + d(n - 1)$$

where d is called the common difference.

Definition 2 (Equal sequences). Two sequences $\{a_n\}$ and $\{b_n\}$ are said to be equal if $a_n = b_n, \forall n \in \mathbb{N}$.

Definition 3 (Sequences bounded from above). $\{a_n\}$ is said to be bounded from above if $\exists M \in \mathbb{R}$, s.t. $a_n \leq M, \forall n \in \mathbb{N}$. Each such M is called an upper bound of $\{a_n\}$.

Definition 4 (Sequences bounded from below). $\{a_n\}$ is said to be bounded from below if $\exists m \in \mathbb{R}$, s.t. $a_n \geq m, \forall n \in \mathbb{N}$. Each such m is called a lower bound of $\{a_n\}$.

Definition 5. $\{a_n\}$ is said to be bounded if it is bounded from below and bounded from above.

Example 8. The sequence $a_n = n^2 + 2$ is not bounded from above but is bounded from below, by all $m \leq 3$.

Example 9. $\left\{ \frac{2n-1}{3n} \right\}$ is bounded.

$$m = 0 \leq \frac{2n-1}{3n} \leq \frac{2n}{3n} = \frac{2}{3} = M$$

Definition 6 (Monotonic increasing sequence). A sequence $\{a_n\}$ is called monotonic increasing if $\exists n_0 \in \mathbb{N}$, s.t. $a_n \leq a_{n+1}, \forall n \geq n_0$.

Definition 7 (Monotonic decreasing sequence). A sequence $\{a_n\}$ is called monotonic decreasing if $\exists n_0 \in \mathbb{N}$, s.t. $a_n \geq a_{n+1}, \forall n \geq n_0$.

Definition 8 (Strongly increasing sequence). A sequence $\{a_n\}$ is called monotonic increasing if $\exists n_0 \in \mathbb{N}$, s.t. $a_n < a_{n+1}, \forall n \geq n_0$.

Definition 9 (Strongly decreasing sequence). A sequence $\{a_n\}$ is called monotonic decreasing if $\exists n_0 \in \mathbb{N}$, s.t. $a_n > a_{n+1}, \forall n \geq n_0$.

Example 10. The sequence $\left\{\frac{n^2}{2^n}\right\}$ is strongly decreasing. However, this is not evident by observing the first few terms. $\frac{1}{2}, 1, \frac{9}{8}, 1, \frac{25}{32}, \dots$

$$\begin{aligned}
 & a_n > a_{n+1} \\
 \iff & \frac{n^2}{2^n} > \frac{(n+1)^2}{2^{n+1}} \\
 \iff & 2n^2 > (n+1)^2 \\
 \iff & \sqrt{2}n > n+1 \\
 \iff & n(\sqrt{2}-1) > 1 \\
 \iff & n > \frac{1}{\sqrt{2}-1} \\
 \iff & n > 3
 \end{aligned}$$

Exercise 1.

Is $a_n = (-1)^n$ monotonic?

Solution 1.

The sequence $-1, 1, -1, 1, \dots$ is not monotonic.

1.1 Limit of a Sequence

Definition 10. Let $\{a_n\}$ be a given sequence. A number L is said to be the limit of the sequence if $\forall \varepsilon > 0, \exists n_0 \in \mathbb{N}$, s.t. $|a_n - L| < \varepsilon, \forall n \geq n_0$. That is, there are infinitely many terms inside the interval and a finite number of terms outside it.

Example 11. The sequence $\left\{\frac{1}{n}\right\}$ tends to 0, i.e. for any open interval $(-\varepsilon, \varepsilon)$, there are finite number of terms of the sequence outside the interval, and therefore there are infinitely many terms inside the interval.

Exercise 2.

Prove

$$\lim_{n \rightarrow \infty} \frac{n+2}{2n-1} = \frac{1}{2}$$

Solution 2.

$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N}$

Exercise 3.

Prove that 2 is not a limit of $\left\{ \frac{3n+1}{n} \right\}$.

Solution 3.

If possible, let

$$\lim_{n \rightarrow \infty} \frac{3n+1}{n} = 2$$

Then, $\forall \varepsilon > 0, \exists n_0 \in \mathbb{N}$, s.t. $\left| \frac{3n+1}{n} - 2 \right| < \varepsilon, \forall n \geq n_0$. However,

$$\left| \frac{3n+1}{n} - 2 \right| = 1 + \frac{1}{n} > 1$$

This is a contradiction for $\varepsilon = \frac{1}{2}$. Therefore, 2 is not a limit.

Theorem 1. *If a sequence $\{a_n\}$ has a limit L then the limit is unique.*

Proof. If possible let there exist two limits L_1 and L_2 . Therefore, $\forall \varepsilon > 0$, there exist a finite number of terms in the interval $(L_1 - \varepsilon, L_1 + \varepsilon)$. Therefore, there exist a finite number of terms in the interval $(L_2 - \varepsilon, L_2 + \varepsilon)$. This contradicts the definition of a limit. Therefore, the limit is unique. \square

Theorem 2. *If a sequence $\{a_n\}$ has limit L , then the sequence is bounded.*

Theorem 3. *Let*

$$\begin{aligned} \lim_{n \rightarrow \infty} a_n &= a \\ \lim_{n \rightarrow \infty} b_n &= b \end{aligned}$$

and let c be a constant. Then,

$$\begin{aligned} \lim c &= c \\ \lim(ca_n) &= c \lim a_n \\ \lim(a_n \pm b_n) &= \lim a_n \pm \lim b_n \\ \lim(a_n b_n) &= \lim a_n \lim b_n \\ \lim\left(\frac{a_n}{b_n}\right) &= \frac{\lim a_n}{\lim b_n} \quad (\text{if } \lim b_n \neq 0) \end{aligned}$$

Theorem 4. Let $\{b_n\}$ be bounded and let $\lim a_n = 0$. Then,

$$\lim(a_nb_n) = 0$$

Theorem 5 (Sandwich Theorem). Let $\{a_n\}$, $\{b_n\}$, $\{c_n\}$ be three sequences. If

$$\lim a_n = \lim b_n = L$$

and $\exists n_0 \in \mathbb{N}$, s.t. $\forall n \geq n_0$, $a_n \leq b_n \leq c_n$. Then,

$$\lim b_n = L$$

Exercise 4.

Calculate $\lim_{n \rightarrow \infty} \sqrt[n]{2^n + 3^n}$

Solution 4.

$$\begin{aligned} \sqrt[n]{3^n} &\leq \sqrt[n]{2^n + 3^n} \leq \sqrt[n]{3^n + 3^n} = \sqrt[n]{2 \cdot 3^n} \\ \therefore 3 &\leq \sqrt[n]{2^n + 3^n} \leq 3\sqrt[n]{2} \end{aligned}$$

Therefore, by the Sandwich Theorem, $\lim_{n \rightarrow \infty} \sqrt[n]{2^n + 3^n} = 3$.

Theorem 6. Any monotonically increasing sequence which is bounded from above converges. Similarly, any monotonically decreasing sequence which is bounded from below converges.

Exercise 5.

Prove that there exists a limit for $a_n = \underbrace{\sqrt{2 + \sqrt{2 + \sqrt{2 + \dots}}}}_{n \text{ times}}$ and find it.

Solution 5.

$$a_1 = \sqrt{2} < \sqrt{2 + \sqrt{2}} = a_2$$

If possible, let

$$\begin{aligned} a_{n-1} &< a_n \\ \therefore \sqrt{2 + a_{n-1}} &< \sqrt{2 + a_n} \\ \therefore a_n &< a_{n+1} \end{aligned}$$

Hence, by induction, $\{a_n\}$ is monotonically increasing.

$$a_1 = \sqrt{2} \leq 2$$

If possible, let

$$\begin{aligned} a_n &\leq 2 \therefore \sqrt{2 + a_n} && \leq \sqrt{2 + 2} \\ \therefore a_{n+1} &\leq 2 \end{aligned}$$

Hence, by induction, $\{a_n\}$ is bounded from above by 2. Therefore, by , $\{a_n\}$ converges.

Definition 11 (Limit in a wide sense). The sequence $\{a_n\}$ is said to converge to $+\infty$ if $\forall M \in \mathbb{R}, \exists n_0 \in \mathbb{N}$, s.t. $\forall n \geq n_0, a_n > M$.

The sequence $\{a_n\}$ is said to converge to $-\infty$ if $\forall M \in \mathbb{R}, \exists n_0 \in \mathbb{N}$, s.t. $\forall n \geq n_0, a_n < M$.

1.2 Sub-sequences

Definition 12 (Sub-sequence). Let $\{a_n\}_{n=1}^{\infty}$ be a sequence. Let $\{n_k\}_{k=1}^{\infty}$ be a strongly increasing sequence of natural numbers. Let $\{b_k\}_{k=1}^{\infty}$ be a sequence such that $b_k = a_{n_k}$. Then $\{b_k\}_{k=1}^{\infty}$ is called a sub-sequence of $\{a_n\}_{n=1}^{\infty}$.

Example 12.

$$a_n = \frac{1}{n}$$

If we choose $n_k = k^2$,

$$b_k = a_{n_k} = a_{k^2} = \frac{1}{k^2}$$

Therefore,

$$\{b_k\} = 1, \frac{1}{4}, \frac{1}{9}, \dots$$

Theorem 7. *If the sequence $\{a_n\}$ converges to L in a wide sense, i.e. L can be infinite, then any sub-sequence of $\{a_n\}$ converges to the same limit L .*

Definition 13 (Partial limit). A real number a , which may be infinite, is called a partial limit of the sequence $\{a_n\}$ if there exists a sub-sequence of $\{a_n\}$ which converges to a .

Example 13. Let

$$a_n = (-1)^n$$

Therefore, $\nexists \lim_{n \rightarrow \infty} a_n$. Let

$$b_k = a_{n_k} = a_{2k-1}$$

Therefore,

$$\begin{aligned} \{b_k\} &= -1, -1, -1, \dots \\ \therefore \lim_{k \rightarrow \infty} b_k &= -1 \end{aligned}$$

Therefore, -1 is a partial limit of $\{a_n\}$.

Theorem 8 (Bolzano-Weierstrass Theorem). *For any bounded sequence there exists a subsequence which is convergent, s.t. there exists at least one partial limit.*

Definition 14 (Upper partial limit). The largest partial limit of a sequence is called the upper partial limit. It is denoted by $\overline{\lim} a_n$ or $\limsup a_n$.

Definition 15 (Lower partial limit). The smallest partial limit of a sequence is called the lower partial limit. It is denoted by $\underline{\lim} a_n$ or $\liminf a_n$.

Theorem 9. *If the sequence $\{a_n\}$ is bounded and*

$$\overline{\lim} a_n = \underline{\lim} a_n = a$$

then

$$\exists \lim a_n = a$$

1.3 Cauchy Characterisation of Convergence

Definition 16. A sequence $\{a_n\}$ is called a Cauchy sequence if $\forall \varepsilon > 0$, $\exists n_0 \in \mathbb{N}$, s.t. $\forall m, n \geq n_0$, $|a_n - a_m| < \varepsilon$.

Theorem 10 (Cauchy Characterisation of Convergence). *A sequence $\{a_n\}$ converges if and only if it is a Cauchy sequence.*

Proof. Let

$$\lim_{n \rightarrow \infty} a_n = L$$

Then $\forall \varepsilon > 0$, $\exists n_0 \in \mathbb{N}$, such that $\forall n \geq n_0$, $|a_n - L| < \frac{\varepsilon}{2}$. Therefore if $n \geq n_0$ and $m \geq n_0$, then

$$\begin{aligned} |a_n - a_m| &= |a_n - L + L - a_m| \\ &\leq |a_n - L| + |L - a_m| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ \therefore |a_n - a_m| &= \varepsilon \end{aligned}$$

Similarly, the converse can be proved by Theorem 9. \square

Theorem 11 (Another Formulation of the Cauchy Characterisation Theorem). *The sequence $\{a_n\}$ converges if and only if $\forall \varepsilon > 0$, $\exists n_0 \in \mathbb{N}$, such that $\forall n \geq n_0$ and $\forall p \in \mathbb{N}$, $|a_{n+p} - a_n| < \varepsilon$.*

Exercise 6.

Prove that the sequence

$$a_n = \frac{1}{1^2} + \frac{1}{2^2} + \cdots + \frac{1}{n^2}$$

is convergent.

Solution 6.

$$\begin{aligned} |a_{n+p} - a_n| &= \left| \frac{1}{1^2} + \frac{1}{2^2} + \cdots + \frac{1}{(n+p)^2} - \left(\frac{1}{1^2} + \frac{1}{2^2} + \cdots + \frac{1}{n^2} \right) \right| \\ &= \frac{1}{(n+1)^2} + \frac{1}{(n+2)^2} + \cdots + \frac{1}{(n+p)^2} \\ \therefore |a_{n+p} - a_n| &< \frac{1}{n(n+1)} + \frac{1}{(n+1)(n+2)} + \cdots + \frac{1}{(n+p-1)(n+p)} \\ \therefore |a_{n+p} - a_n| &< \frac{1}{n} - \frac{1}{\cancel{n+1}} + \frac{1}{\cancel{n+1}} + \cdots + \frac{1}{\cancel{n+p-1}} - \frac{1}{n+p} \\ \therefore |a_{n+p} - a_n| &< \frac{1}{n} - \frac{1}{n+p} \\ \therefore |a_{n+p} - a_n| &< \frac{1}{n} \end{aligned}$$

Therefore, $\forall \varepsilon > 0$, $\exists n_0 \in \mathbb{N}$, s.t. $\forall n \geq n_0$ and $\forall p \in \mathbb{N}$, $|a_{n+p} - a_n| < \varepsilon$, where $n_0 > \frac{1}{\varepsilon}$. \square

Exercise 7.

Prove that the sequence

$$a_n = \frac{1}{1} + \frac{1}{n} + \cdots + \frac{1}{n}$$

diverges.

Solution 7.

If possible, let the sequence converge. Then, by the Cauchy Characterisation of Convergence, $\forall \varepsilon > 0$, $\exists n_0 \in \mathbb{N}$, s.t. $\forall n \geq n_0$ and $\forall p \in \mathbb{N}$, $|a_{n+p} - a_n| < \varepsilon$. Therefore,

$$\begin{aligned} |a_{n+p} - a_n| &= \left| \frac{1}{1} + \frac{1}{2} + \cdots + \frac{1}{n} + \frac{1}{n+p} - \left(\frac{1}{n} + \cdots + \frac{1}{n} \right) \right| \\ &= \frac{1}{n+1} + \cdots + \frac{1}{n+p} \\ &\geq p \cdot \frac{1}{n+p} \\ \therefore |a_{n+p} - a_n| &> \frac{p}{n+p} \end{aligned}$$

If $n = p$,

$$\frac{p}{n+p} = \frac{1}{2}$$

This contradicts the result obtained from the Cauchy Characterisation of Convergence, for $\varepsilon = \frac{1}{4}$.

Therefore, the sequence diverges.

2 Series

Definition 17 (Series). Given a sequence $\{a_n\}$, the sum $a_1 + \cdots + a_n + \cdots$ is called an infinite series or series. It is denoted as $\sum_{n=1}^{\infty} a_n$ or $\sum a_n$.

Definition 18 (Partial sum). The partial sum of the series $\sum a_n$ is defined as

$$S_i = a_1 + \cdots + a_i$$

Definition 19 (Convergent and divergent series). If the sequence $\{S_n\}_{n=1}^{\infty}$ converges, then the series is called convergent. Otherwise, the series is called divergent.

Definition 20 (Sum of a series). If the sequence $\{S_n\}_{n=1}^{\infty}$ converges to $S \neq \pm\infty$, the number S is called the sum of the series.

$$\sum_{n=1}^{\infty} a_n = S$$

Example 14.

$$\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \frac{1}{2^n}$$

Therefore,

$$S_1 = \frac{1}{2} \tag{1}$$

$$S_2 = \frac{1}{2} + \frac{1}{2^2} \tag{2}$$

$$\vdots S_n = \frac{1}{2} + \cdots + \frac{1}{2^n} \tag{3}$$

$$= \frac{a_1(1 - q^n)}{1 - q} \tag{4}$$

$$= \frac{1/2(1 - 1/2^n)}{1 - 1/2} \tag{5}$$

$$= 1 - \frac{1}{2^n} \tag{6}$$

$$\lim_{n \rightarrow \infty} S_n = 1 \tag{7}$$

Therefore, the series converges.

$$S = \sum_{n=1}^{\infty} = 1$$

Theorem 12. A geometric series $\sum_{n=1}^{\infty} a_1 q^{n-1}$, $a_1 \neq 0$ converges if $|q| < 1$ and then,

$$S = \sum_{n=1}^{\infty} a_1 q^{n-1} = \frac{a_1}{1 - q}$$

Definition 21 (p -series). The series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ is called the p -series.

Theorem 13. *The p -series converges for $p > 1$ and diverges for $p \leq 1$.*

Theorem 14. *If $\sum a_n$ converges, then*

$$\lim_{n \rightarrow \infty} a_n = 0$$

Proof.

$$\begin{aligned} a_n &= S_n - S_{n-1} \\ \therefore \lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} S_n - \lim_{n \rightarrow \infty} S_{n-1} \\ &= S - S \\ &= 0 \end{aligned}$$

□

Theorem 15. *If $\sum a_n$ and $\sum b_n$ converge, then $\sum(a_n \pm b_n)$ and $\sum ca_n$, where c is a constant, also converge. Also,*

$$\begin{aligned} \sum(a_n \pm b_n) &= \sum a_n \pm \sum b_n \\ \sum(ca_n) &= c \sum a_n \end{aligned}$$

2.1 Convergence Criteria

2.1.1 Leibniz's Criteria

Definition 22 (Alternating series). The series $\sum_{n=1}^{\infty} (-1)^{n-1} a_n$, where all $a_n > 0$ or all $a_n < 0$ is called an alternating series.

Theorem 16 (Leibniz's Criteria for Convergence). *If an alternating series $\sum (-1)^{n-1} a_n$ with $a_n > 0$ satisfies*

1. $a_{n+1} \leq a_n$, i.e. $\{a_n\}$ is monotonically decreasing.

2. $\lim_{n \rightarrow \infty} a_n = 0$

then the series $\sum (-1)^{n-1} a_n$ converges.

Proof. Consider the even partial sums of the series $\sum_{n=1}^{\infty} (-1)^{n-1} a_n$.

$$S_{2m} = (a_1 - a_2) + (a_3 - a_4) + \cdots + (a_{2m-1} - a_{2m})$$

As $\{a_n\}$ is monotonically increasing, all brackets are non-negative. Therefore,

$$S_{2m+2} \geq S_{2m}$$

Therefore, $\{S_{2m}\}$ is increasing.

Also,

$$S_{2m} = a_1 - (a_2 - a_3) - (a_4 - a_5) - \cdots - (a_{2m-2} - a_{2m-1}) - a_{2m}$$

All brackets and a_{2m} are non-negative. Therefore,

$$S_{2m} \leq a_1$$

Therefore, $\{S_{2m}\}$ is bounded from above by a_1 . Hence,

$$\exists \lim_{m \rightarrow \infty} S_{2m} = S$$

For S_{2m+1} ,

$$\begin{aligned} S_{2m+1} &= S_{2m} + a_{2m+1} \\ \therefore \lim_{m \rightarrow \infty} S_{2m+1} &= \lim_{m \rightarrow \infty} S_{2m} + \lim_{m \rightarrow \infty} a_{2m+1} \\ &= S + 0 \\ &= S \end{aligned}$$

Therefore,

$$\lim_{n \rightarrow \infty} S_n = S$$

□

Example 15. The alternating harmonic series $\sum \frac{(-1)^{n-1}}{n}$ converges as $a_n = \frac{1}{n} > 0$, a_n decreases and $\lim a_n = 0$.

2.1.2 Comparison Test

Theorem 17 (Comparison Test for Convergence). Assume $\exists n_0 \in \mathbb{N}$, such that $a_n \geq 0$, $b_n \geq 0$, $\forall n \geq n_0$.

1. If $a_n \leq b_n$, $\forall n \geq n_0$ and $\sum_{n=1}^{\infty} b_n$ converges, then $\sum_{n=1}^{\infty} a_n$ converges.
2. If $a_n \geq b_n$, $\forall n \geq n_0$ and $\sum_{n=1}^{\infty} b_n$ diverges, then $\sum_{n=1}^{\infty} a_n$ diverges.

Theorem 18 (Another Formulation of the Comparison Test for Convergence). Assume $\exists n_0 \in \mathbb{N}$, such that $a_n \geq 0$, $b_n \geq 0$, $\forall n \geq n_0$ and

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = a > 0$$

where a is a finite number. Then $\sum_{n=1}^{\infty} a_n$ converges if and only if $\sum_{n=1}^{\infty} b_n$ converges.

2.1.3 d'Alembert Criteria (Ratio Test)

Definition 23 (Absolute and conditional convergence). The series $\sum a_n$ is said to converge absolutely if $\sum |a_n|$ converges. The series $\sum a_n$ is said to converge conditionally if it converges but $\sum |a_n|$ diverges.

Example 16. The series $\sum \frac{(-1)^{n-1}}{n^2}$ converges absolutely, as $\sum \left| \frac{(-1)^{n-1}}{n^2} \right| = \sum \frac{1}{n^2}$ converges.

Example 17. The series $\sum \frac{(-1)^{n-1}}{n}$ converges conditionally, as it converges, but $\sum \left| \frac{(-1)^{n-1}}{n} \right| = \sum \frac{1}{n}$ diverges.

Theorem 19. *If the series $\sum a_n$ converges absolutely then it converges.*

Theorem 20 (d'Alembert Criteria (Ratio Test)). 1. *If*

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n-1}}{a_n} \right| = L < 1$$

then $\sum a_n$ converges absolutely.

2. *If*

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n-1}}{a_n} \right| = L > 1$$

(including $L = \infty$), then $\sum a_n$ converges diverges.

3. *If $L = 1$, the test does not apply.*

2.1.4 Cauchy Criteria (Cauchy Root Test)

Theorem 21 (Cauchy Criteria (Cauchy Root Test)). 1. *If*

$$\overline{\lim} \sqrt[n]{|a_n|} = L < 1$$

then $\sum a_n$ converges absolutely.

2. *If*

$$\overline{\lim} \sqrt[n]{|a_n|} = L > 1$$

(including $L = \infty$), then $\sum a_n$ diverges.

3. *If $L = 1$, the test does not apply.*

2.1.5 Integral Test

Theorem 22 (Integral Test for Series Convergence). *Let $f(x)$ be a continuous, non-negative, monotonic decreasing function on $[1, \infty)$ and let $a_n = f(n)$. Then the series $\sum_{n=1}^{\infty} a_n$ converges if and only if the improper integral $\int_1^{\infty} f(x) dx$ converges.*

Exercise 8.

Does $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converge or diverge?

Solution 8.

Let

$$f(x) = \frac{1}{x^p}$$

with $p > 0$.

Therefore, $f(x)$ is continuous, non-negative and monotonic decreasing on $[1, \infty)$. Therefore, the Integral Test for Series Convergence is applicable.

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx$$

If $p \neq 1$,

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^p} &= \lim_{t \rightarrow \infty} \left. \frac{x^{-p+1}}{-p+1} \right|_1^t \\ &= \lim_{t \rightarrow \infty} \left(\frac{t^{-p+1}}{-p+1} - \frac{1}{-p+1} \right) \\ &= \frac{1}{p-1} \end{aligned}$$

If $p = 1$,

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^p} &= \lim_{t \rightarrow \infty} \ln x \Big|_1^t \\ &= \infty \end{aligned}$$

Therefore, the series converges for $p > 1$ and diverges for $p \leq 1$.

Theorem 23. *If the series $\sum a_n$ absolutely converges and the series $\sum b_n$ is obtained from $\sum a_n$ by changing the order of the terms in $\sum a_n$ then $\sum b_n$ also absolutely converges and $\sum b_n = \sum a_n$.*

Theorem 24. *If a series converges then the series with brackets without changing the order of terms also converges. That is, if $\sum a_n$ converges, then any series of the form $(a_1 + a_2) + (a_3 + a_4 + a_5) + a_6 + \dots$ also converges.*

Theorem 25. *If a series with brackets converges and the terms in the brackets have the same sign, then the series without brackets also converges.*

3 Power Series

Definition 24 (Power series). The series $\sum_{n=0}^{\infty} a_n(x - c)^n$ is called a power series.

Theorem 26 (Cauchy-Hadamard Theorem). *For any power series $\sum_{n=0}^{\infty} a_n(x - c)^n$ there exists the limit, which may be infinity,*

$$R = \frac{1}{\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|}}$$

and the series converges for $|x - c| < R$ and diverges for $|x - c| > R$. The end points of the interval, i.e. $x = c - R$ and $x = c + R$ must be separately checked for series convergence.

Definition 25 (Radius of convergence and convergence interval). The number R is called the radius of convergence and the interval $|x - c| < R$ is called the convergence interval of the series. The point c is called the centre of the convergence interval.

Theorem 27. *If $\exists \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$, which may be infinite, then,*

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

Theorem 28 (Stirling's Approximation). *For $n \rightarrow \infty$,*

$$n! \approx \left(\frac{n}{e} \right)^n \sqrt{2\pi n}$$

3.1 Differentiation and Integration of Power Series

Theorem 29. If R is a radius of convergence of the power series $\sum_{n=0}^{\infty} a_n(x - c)^n$ then the function $f(x) = \sum_{n=0}^{\infty} a_n(x - c)^n$ is differentiable on $(c - R, c + R)$ and the derivative is

$$f'(x) = \sum_{n=0}^{\infty} n a_n (x - c)^{n-1}$$

Theorem 30. If R is a radius of convergence of the series $\sum_{n=0}^{\infty} a_n(x - c)^n$ then the function $f(x) = \sum_{n=0}^{\infty} a_n(x - c)^n$ is integrable in $(c - R, c + R)$ and

$$\int f(x) dx = \sum_{n=0}^{\infty} a_n \frac{(x - c)^{n+1}}{n + 1} + A$$

where $c - R < x < c + R$.

Exercise 9.

Find $\int_0^x e^{-t^2} dt$.

Solution 9.

$\forall s \in \mathbb{R}$,

$$\begin{aligned} e^s &= 1 + \frac{s}{1!} + \frac{s^2}{2!} + \cdots + \frac{s^n}{n!} + \cdots \\ \therefore e^{-t^2} &= 1 - \frac{t^2}{1!} + \frac{t^4}{2!} + \cdots + (-1)^n \frac{t^{2n}}{n!} + \cdots \\ \therefore \int_0^x e^{-t^2} dt &= x - \frac{x^3}{1!3} + \frac{x^5}{2!5} + \cdots + (-1)^n \frac{x^{2n+1}}{n!(2n+1)} + \cdots \end{aligned}$$

Theorem 31. If the series $A(x) = \sum_{n=0}^{\infty} a_n x^n$ and $B(x) = \sum_{n=0}^{\infty} B_n x^n$ absolutely converge for $|x| < R$ and $c_n = \sum_{k=0}^n a_k b_{n-k}$, then the series $C(x) = \sum_{n=0}^{\infty} c_n x^n$ also absolutely converges for $|x| < R$ and $C(x) = A(x)B(x)$.

3.2 Taylor Series

Definition 26 (Taylor series). Let $f(x)$ be infinitely differentiable on an open interval about a and let x be an arbitrary point in the interval. Then the power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$ is called the Taylor series of $f(x)$ at a . If $a = 0$ then it is called the Maclaurin series of $f(x)$ at 0.

Theorem 32. *If there exists a power series which converges to $f(x)$, i.e. if, for $|x-a| < R$,*

$$f(x) = \sum_{n=0}^{\infty} a_n (x-a)^n$$

then, for $|x-a| < R$,

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

that is, $\forall n$,

$$a_n = \frac{f^{(n)}(a)}{n!}$$

Exercise 10.

Show that

$$f(x) = \begin{cases} 0 & ; \quad x = 0 \\ e^{-\frac{1}{x^2}} & ; \quad x \neq 0 \end{cases}$$

is not equal to its Taylor series at $a = 0$.

Solution 10.

If $n = 1$,

$$\begin{aligned} f^{(1)}(0) &= \lim_{\Delta x \rightarrow 0} \frac{f(0 + \Delta x) - f(0)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{e^{-\frac{1}{(\Delta x)^2}}}{\Delta x} \end{aligned}$$

Let $t = \frac{1}{\Delta x}$

$$\begin{aligned}\therefore f'(0) &= \lim_{t \rightarrow \infty} \frac{e^{-t^2}}{\frac{1}{t}} \\ &= \lim_{t \rightarrow \infty} \frac{t}{e^{t^2}} \\ &= \lim_{t \rightarrow \infty} \frac{1}{e^{t^2} 2t} \\ &= 0\end{aligned}$$

Therefore,

$$f'(x) = \begin{cases} 0 & ; \quad x = 0 \\ e^{-\frac{1}{x^2} \cdot 2 \cdot x^{-3}} & ; \quad x \neq 0 \end{cases}$$

Similarly, $\forall n \geq 1, f^{(n)}(0) = 0$

Therefore, the Taylor series is not equal to $f(x)$.

Exercise 11.

Find the Maclaurin series of $f(x) = e^x$ and prove that the series converges to $f(x)$ for any $x \in \mathbb{R}$.

Solution 11.

$\forall n \geq 1, f^{(n)}(x) = e^x$.

Therefore,

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \cdots + \frac{x^n}{n!} + \frac{e^c x^{n+1}}{(n+1)!}$$

where c is between 0 and x .

Therefore, as

$$0 \leq |R_n(x)| \leq \frac{|x|^{n+1}}{(n+1)!}$$

by the Sandwich Theorem

$$\lim_{n \rightarrow \infty} |R_n(x)| = 0$$

Therefore,

$$e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \cdots + \frac{x^n}{n!} + \cdots$$

4 Series of Real-valued Functions

Definition 27 (Sequence of functions). A sequence $\{f_n\} = f_1(x), f_2(x), \dots$ defined on $D \subseteq \mathbb{R}$ is called a sequence of functions.

Definition 28 (Pointwise convergence and domain of convergence). $\{f_n\}$ converges pointwise in some domain $E \subseteq D$ if for every $x \in E$, the sequence of $\{f_n(x)\}$ converges. In such a case, E is said to be a domain of convergence of $\{f_n\}$.

Exercise 12.

Find the domain of convergence of $f_n(x) = x^n$, defined on some $D \subseteq \mathbb{R}$.

Solution 12.

$$\lim_{n \rightarrow \infty} f_n(x) = \begin{cases} 0 & ; \quad -1 < x < 1 \\ 1 & ; \quad x = 1 \\ \text{diverges} & ; \quad x \notin (-1, 1] \end{cases}$$

Therefore, the domain of convergence of $\{f_n\}$ is $(-1, 1]$.

Exercise 13.

Let $f(x) : (0, \infty) \rightarrow \mathbb{R}$ be some function such that $\lim_{x \rightarrow \infty} f(x) = 0$. Let $f_n(x) = f(nx)$. What is the domain of convergence of f_n ? What is the limit function?

Solution 13.

Let x have some fixed value in $(0, \infty)$. Therefore, as $\lim_{x \rightarrow \infty} f(x) = 0$,

$$\begin{aligned} \lim_{n \rightarrow \infty} f_n(x) &= \lim_{n \rightarrow \infty} f(nx) \\ &= 0 \end{aligned}$$

Therefore, the domain of convergence is $(0, \infty)$ and the limit function is a constant function with value 0.