

Viscereality: Bioresponsive Virtual Reality for Box Breathing and Altered-State Experience

George Fejer^{1,2,3*}, Till Holzapfel⁴, Taru Hirvonen,
Anestis Lalidis Mateo⁶, Johannes Blum⁵, Michael Gaebler²,
Bigna Lenggenhager^{7,8}

^{1*}University of Konstanz, Konstanz, Germany.

²Max Planck Institute for Human Cognitive and Brain Sciences,
Germany.

³ALIUS Research Network, Germany.

⁴University of Duisburg-Essen, Research Center Trustworthy Data
Science and Security, Duisburg, Germany.

⁵Pädagogische Hochschule Schaffhausen, Schaffhausen, Switzerland.

⁶Institute for Building Intelligence, Hochschule Bielefeld - University of
Applied Sciences and Arts, Bielefeld, Germany.

⁷AIR—Association for Independent Research, Zurich, Switzerland.

⁸Department of Psychology, University of Zurich, Zurich, Switzerland.

*Corresponding author(s). E-mail(s): george.fejer@aliusresearch.org;
Contributing authors: tillholzapfel@gmail.com; 2taruhi@gmail.com;
alalidis@hsbi.de; johannes.blum@phsh.ch; gaebler@mpib-berlin.mpg.de;
bigna.lenggenhager@gmail.com;

Abstract

Breath-based interactions in virtual reality typically map breathing onto discrete visual objects or physiological proxies. Viscereality instead translates respiratory state into a surrounding particle field, mapping lung volume onto external space so that inhalation expands the environment and exhalation contracts it. Because particle fields have no fixed boundary, they can convey information through the motion and interactions of many simple elements. We tested whether this virtual-reality paradigm could reliably elicit altered-state experiences during box breathing, whether transient symmetric particle coupling would further influence subjective experience, and whether participants could identify the manipulation.

In a preregistered within-subjects study, 39 participants completed three counterbalanced 10-minute audio-guided box-breathing sessions: a high-symmetry virtual-reality condition, an asymmetric virtual-reality condition, and a dark-screen control. We assessed breathing adherence, altered states of consciousness, pictographic self-related measures, and sensitivity to the symmetry manipulation. Relative to the dark-screen control, the pooled virtual-reality conditions improved breathing synchronization, increased ratings of the perceived spatial extension of the self, and produced altered-state profiles with strong Perceptual Effects, moderate Positive Effects, and weaker Distressing Effects. Participants detected some symmetry-relevant features above chance, but neither reliably identified which session was symmetric nor showed outcome differences between the two virtual-reality mappings. These findings show that the overall virtual-reality paradigm produced reliable experiential differences from the control, but they do not establish that bioresponsiveness rather than visual engagement caused those differences. The specific affective contribution of particle symmetry, and whether it requires explicit awareness, therefore remain unresolved.

Keywords: biofeedback, breath-based interaction, breathwork, altered states of consciousness, particle systems

1 Introduction

1.1 Breath as an Affective Sensory Modality

Breathing is a unique and multifaceted sensory modality, as it carries both interoceptive and exteroceptive streams of information. One stream concerns internal sensory information about respiratory mechanics and blood-gas status, including signals related to lung inflation, airway and respiratory-muscle activity, and blood O_2/CO_2 balance (Chan et al. 2024; Engelen et al. 2023; Weng et al. 2021). Because inhaled air stems from the external environment, breathing also samples exteroceptive information about ambient conditions, including odours, airflow, and air temperature, through its coupling with orthonasal olfaction and thermo- and mechanosensory receptors in the upper airways (Mori and Sakano 2022a,b; Zhou et al. 2011; Sabnis et al. 2008; Kytikova et al. 2021). Breathing is also closely tied to affective experience. People sigh with relief after stress (Balban et al. 2023), hold their breath during freeze responses (Boiten et al. 1994), and hyperventilate during fear or panic (Jerath and Beveridge 2020; Kreibig 2010). Arousing negative emotions such as fear and anger are linked to shallower, more rapid breathing, whereas relaxation and contentment correspond to slower, deeper respiratory patterns (Boiten et al. 1994; Homma and Masaoka 2008; Jerath and Beveridge 2020). Unlike most visceral signals such as heartbeat, breathing is also amenable to voluntary control (Sarkar 2017), making this relationship bidirectional. Voluntarily changing respiratory patterns can modulate affective state (Jerath and Beveridge 2020; Ashhad et al. 2022), a principle that underlies breathwork as a therapeutic modality (Fincham et al. 2023; Balban et al. 2023; Siebieszuk

et al. 2025). Breathwork techniques range from slow-paced protocols that target relaxation, stress regulation, and improvements in heart rate variability through sustained cardiovascular-respiratory coupling (Balban et al. 2023; Ahmed et al. 2021; Siebieszuk et al. 2025), to high-ventilation protocols such as cyclic hyperventilation that can facilitate more pronounced alterations in affect and consciousness (Bahi et al. 2024a; Lewis-Healey et al. 2024a; Havenith et al. 2025a), effects increasingly posited as therapeutic, for instance in facilitating emotional breakthroughs (Fincham et al. 2026, 2024). Breathwork is already established as a therapeutic modality for stress, anxiety, and depression (Fincham et al. 2023; Bentley et al. 2023; Banushi et al. 2023), and a growing number of virtual-reality applications also incorporate breathing for therapeutic and health-related purposes (Cortez-Vázquez et al. 2024; Pancini et al. 2025). Yet this literature has so far explored only a relatively limited portion of the broader design space that breathing affords as an interaction modality in VR.

1.2 Breath-Based Interactions in VR

Existing breath-based VR systems span meditation (Prpa et al. 2020; Shamekhi and Bickmore 2018), anxiety regulation (Bossenbroek et al. 2020), stress management through heart-rate-variability biofeedback (Blum et al. 2019; Rockstroh et al. 2019), and rehabilitation (Shi et al. 2023; Alhammad 2024). In the context of our framework, we distinguish between two motifs in the design space regarding the mapping of the user’s respiratory signal onto discrete visual objects, in contrast to environmental mappings. Object-oriented approaches typically place respiratory feedback in a discrete visual target. Third-person variants let the user watch an external entity respond to breathing, such as a cloud that approaches during inhalation and recedes during exhalation (Tinga et al. 2019). In first-person variants, the mapping is applied to a virtual body seen from the user’s own vantage point. The “embreathment” illusion demonstrated that synchronising a virtual avatar’s chest movements with the user’s breathing enhances body ownership and agency (Monti et al. 2020a), and mapping respiratory signals onto the expansion and contraction of a virtual avatar bubble viewed from a third-person perspective has been found to enhance agency and body awareness (Wald et al. 2023). What these object-oriented designs share is that respiratory feedback is concentrated in a single visual target, whether an external creature or a self-representation, which the user attends to as an indicator of their breathing. These designs are typically evaluated in terms of embodiment, measuring whether the temporal coupling between breathing and visual response enhances body ownership, felt agency, and awareness of the respiratory signal (Monti et al. 2020a; Wald et al. 2023).

We contrast these with designs that embed respiratory feedback in the surrounding virtual environment rather than in a discrete object. DEEP (Bossenbroek et al. 2020; van Rooij et al. 2016) and Flowborne (Rockstroh et al. 2021) use diaphragmatic breathing to drive locomotion through immersive landscapes, rewarding paced breathing with movement and environmental change (Bossenbroek et al. 2020; Rockstroh et al. 2021). In a seated variant, Blum et al. (2020) mapped exhalation onto colour changes in flowers, trees, and rocks, keeping the feedback environmental but locally anchored to scene objects. A further group of environmental designs couples not the breathing signal directly but a slower downstream variable, typically heart

139 rate variability, to ambient scene properties. In [Blum et al. \(2019\)](#) and *Heart Garden*
 140 ([Rook et al. 2026](#)), increasing coherence or heart rate variability gradually transforms
 141 the scene. These designs create calming feedback loops, but because the environment
 142 changes through a slower physiological proxy rather than respiration itself, moment-
 143 to-moment respiratory agency is attenuated. The locomotion-based designs provide
 144 immediate agency, but mainly as a progression mechanic rather than as a way of mak-
 145 ing surrounding space itself feel breath-coupled. In our design, we sought to synthesize
 146 these two approaches by mapping the user’s breathing state onto an environment that
 147 expands and contracts with respiration. This provides immediate feedback and a sense
 148 of agency while presenting breathing as an ambient property of the surrounding space
 149 rather than as a distinct object requiring continuous attention. Specifically, the envi-
 150 ronment is composed of particles that respond to the user’s breath such that inhalation
 151 expands the space outward and exhalation contracts it inward. In this way, changes
 152 in internal lung volume are mapped onto changes in external space. This mapping is
 153 closely related to what [Goral et al. \(2024b\)](#) describe as exteroceptive–interoceptive
 154 sensory substitution, in which internal bodily signals are translated into external sen-
 155 sory events that remain experientially linked to the body rather than appearing as
 156 separate readouts. By mirroring the felt expansion and contraction of the thorax,
 157 our system was intended to preserve the natural structure of breathing experience
 158 and thereby function as an intuitive interface that can be understood with minimal
 159 explicit instruction ([Antle et al. 2009](#)). This choice was also informed by evidence
 160 from embodied cognition showing that people judge quantities as larger after inhala-
 161 tion than after exhalation ([Belli et al. 2021](#)), which may indicate a possible, though
 162 still speculative, link between respiratory state and the perception of space. More
 163 broadly, this embodied metaphor appears across a range of immersive biofeedback
 164 systems that map respiration onto expanding and contracting visual stimuli ([Wald
 165 et al. 2023](#); [Goral et al. 2024b](#); [Barton et al. 2024](#); [Stepanova et al. 2020](#)). Two exist-
 166 ing systems are conceptually closest to the approach presented here. [Barton et al.
 167 \(2024\)](#) immersed participants in an abstract fractal-like environment and used a cen-
 168 tral breathing guide whose size participants controlled by spreading and closing their
 169 arms in time with the guide; the interaction was therefore structured as an explicit
 170 movement-synchronisation task rather than as an environment that responded directly
 171 to respiration. [Goral et al. \(2024b\)](#) introduced exteroceptive–interoceptive sensory
 172 substitution in a system that translated respiratory signals into dynamic visual and
 173 auditory stimuli intended to feel continuous with the breathing body; however, their
 174 implementation externalized the signal through projection rather than embedding it in
 175 an immersive three-dimensional environment. Our system was designed to combine the
 176 strengths of these two approaches by pairing real-time environmental responsiveness
 177 that affords immediate respiratory agency with feedback distributed across surround-
 178 ing space rather than concentrated in a single object or task. Rather than relying on
 179 semantic cues such as naturalistic scenery, skyboxes, or rendered objects, we pursued
 180 a bottom-up design in which the sense of surrounding space emerges from relations
 181 among many simple particles, drawing on Gestalt principles such as proximity, com-
 182 mon fate, and continuity ([Wagemans et al. 2012](#)) so that their dynamic arrangement,
 183
 184

and specifically their responses to expansion and contraction, conveys depth and closeness dynamically within an abstract particle-based environment. This design rationale also motivates the broader question of whether coupling an internal bodily signal such as respiration to the surrounding sense of space can also reshape the sense of self, which we consider next through a review of relevant VR experiments.

1.3 VR interventions testing the malleability of the self

Virtual reality has been used extensively to investigate the malleability of bodily self-experience through multisensory manipulation. In full-body illusion paradigms, synchronous visuotactile stimulation between a participant’s physical body and a seen virtual body can shift perceived self-location and induce a sense of ownership over the virtual body (Lenggenhager et al. 2007; Petkova and Ehrsson 2008). Later work showed that interoceptive signals can serve a similar synchronising function. Presenting a virtual body or silhouette that flashes in synchrony with the participant’s heartbeat enhances body ownership and alters self-location even in the absence of tactile stimulation (Aspell et al. 2013; Suzuki et al. 2013). Respiratory signals have been used in a similar way. For example, synchronising a virtual avatar’s chest movements, viewed from a first-person perspective, with the user’s breathing enhances body ownership (Monti et al. 2020a). Similarly, mapping respiratory signals onto the expansion and contraction of a virtual bubble, viewed from a third-person perspective, enhances agency and body awareness (Wald et al. 2023). Collectively, these studies show that a range of sensory modalities, including touch, breathing, and cardiac activity, can be used to reinforce visual feedback and thereby strengthen the sense of embodiment toward a virtual object or avatar from both first-person and third-person perspectives (Lenggenhager et al. 2007; Petkova and Ehrsson 2008; Aspell et al. 2013; Suzuki et al. 2013; Monti et al. 2020a; Wald et al. 2023).

A related but distinct line of research concerns peripersonal space, the region immediately surrounding the body where the brain integrates multisensory signals in body-centered coordinates to maintain a dynamic boundary between self and environment (Serino 2019; Bufacchi and Iannetti 2018). This boundary is not fixed; it extends with tool use (Maravita and Iriki 2004), shifts with emotional valence and social context (Bufacchi and Iannetti 2018; Serino 2019), and interacts with interoceptive aspects of bodily self-representation (Tsakiris et al. 2011; Azzalini et al. 2019). Recent virtual-reality work further suggests that embodiment and multisensory manipulations can reshape peripersonal space in VR (Petrizzo et al. 2024; Karakoc et al. 2025). While the embodiment paradigms reviewed above manipulate the felt relationship between self and a visual body representation, peripersonal space research highlights that self-experience also depends on how the brain represents the spatial relationship between the body and the surrounding environment (Serino 2019; Bufacchi and Iannetti 2018).

Notably absent from both lines of research are paradigms that intervene through the surrounding environment rather than through a representation of the body itself. In most VR studies of embodiment and self-location, self-related experience is altered by manipulating how the body is represented, with any effects on perceived space arising indirectly through changes in how the body is seen in space (Guy et al. 2023). Although VR research has also developed techniques that manipulate space itself, such

as scaling, translation, and redirected movement, these are typically used to support navigation and interaction rather than to investigate self-experience (Abtahi et al. 2022). By contrast, the present approach couples respiration not to a virtual body or discrete object, but to the spatial extent of the surrounding environment itself. This may engage the boundary between body-centred and environment-centred space in a way that body-focused paradigms do not.

There is a broad consensus that the sense of self is multidimensional (Millière 2020; Taves 2020), and the present study assessed three complementary pictographic measures. These were body-boundary salience (Dambrun 2016), spatial frame of reference (Hanley and Garland 2019), and perceived self-size (Vidal et al. 2025). At the same time, the intervention was expected to affect not only self-experience, but potentially the broader structure of subjective experience. The present study therefore also considered its effects in relation to psychometric frameworks from altered-states research, which informed both measurement and the design rationale of the VR system described next.

1.4 Breathwork as an experiential design space for virtual reality

Embedding a breathing protocol in a responsive virtual environment introduces specific design constraints. Different breathing practices produce different respiratory patterns, including variations in the pace, duration, and force of inhalation and exhalation. Visual feedback coupled to these dynamics must therefore avoid becoming dizzying, claustrophobic, disorienting, or overwhelming. In VR, large-amplitude radial scene motion, particularly in the visual periphery, is a documented source of cybersickness (Chang et al. 2020; Pöhlmann et al. 2021), and contracting environments can elicit involuntary defensive postural responses even in non-phobic users (Kodithuwakku Arachchige et al. 2022). These risks are compounded when the user is in a heightened or unfamiliar experiential state, given that negative affect and arousal are associated with increased VR sickness severity (Kaufeld et al. 2022). The choice of breathing protocol therefore shapes not only what physiological and experiential effects to expect, but also what the visual environment can safely and comfortably do in response.

High-ventilation breathwork practices such as holotropic breathing, conscious connected breathing, and cyclic hyperventilation involve sustained increases in breathing rate and depth that, over extended sessions, induce pronounced physiological changes, including respiratory alkalosis and reductions in cerebral blood flow, and can give rise to marked changes in subjective experience (Lewis-Healey et al. 2024a; Kartar et al. 2025; Fincham et al. 2024). Participants report experiences of oceanic boundlessness, in which the felt boundary between self and world dissolves into a sense of unity or merging with the surroundings; visionary restructuralisation, involving vivid geometric imagery, complex visual scenes, and synesthetic blending of sensory modalities; and dread of ego dissolution, characterised by a felt loss of control over one’s thoughts and sense of identity (Bahi et al. 2024a; Havenith et al. 2025a; Lewis-Healey et al. 2024a). These experiential dimensions overlap with those reported under medium- to high-dose psilocybin as measured by the 11D-ASC (Studerus et al. 2010; Havenith et al. 2025a),

and are often accompanied by emotional catharsis and strong bodily sensations such as tingling, dizziness, and autonomic arousal (Havenith et al. 2025a; Lewis-Healey et al. 2024a; Fincham et al. 2024, 2026). However, the rapid, large-amplitude chest excursions characteristic of high-ventilation protocols would produce correspondingly rapid expansion and contraction of a breath-coupled particle environment, and the accompanying physiological arousal and altered experiential state would coincide with precisely the visual conditions most likely to provoke cybersickness and spatial discomfort. For these reasons, we chose to facilitate slow-paced breathing rather than high-ventilation breathwork in the present system.

Slow-paced breathing occupies a different region of this subjective and affective space. Its established effect profile is characterised by relaxation, reduced anxiety, increased comfort, and calm alertness, supported by parasympathetic dominance and enhanced heart rate variability (Zaccaro et al. 2018; Balban et al. 2023; Fincham et al. 2023). Slow breathing is also one of the most common attentional anchors in guided meditation practice (Gerritsen and Band 2018; Zaccaro et al. 2018), and evidence from experienced pranayama practitioners indicates that very slow nasal breathing can induce altered states of consciousness characterised by altered body perception, distortions in time sense, unusual salience, increased joy, and reduced anxiety (Zaccaro et al. 2022). The mild but documented non-ordinary experiential profile of slow-paced breathing provided a low baseline against which any augmentation by the visual environment would be easier to detect, while keeping the environment’s spatial dynamics within a comfortable perceptual range.

Augmenting subjective experience is clinically relevant because experiential intensity during altered-state interventions predicts therapeutic outcomes (Roseman et al. 2018; Romeo et al. 2025; Havenith et al. 2025a). We chose a slow-paced protocol precisely because its mild baseline provides sensitivity for detecting whether we can enhance specific attributes of the subjective experience through VR. To characterise this experiential profile and compare it with existing breathwork and psychedelic literatures, we adopted the 11D-ASC (Studerus et al. 2010; Dittrich 1998), the standard instrument across these fields. Its subscales capture the phenomenological features (e.g., boundary dissolution, changed imagery, synesthetic blending) that a breath-coupled environment might selectively amplify, and can be summarised into three higher-order dimensions reflecting positive, distressing, and perceptual effects (Stocker et al. 2025). Because we also examine how these dimensions evolve over the course of a breathing session, we include psychometric time-series ratings, which we refer to as temporal experience tracers (Jachs 2021). We match the tracers one-to-one with the 11D-ASC dimensions to compare global ratings with time-resolved intensity profiles of the same experiential constructs. The motivation to enhance subjective experience is also reflected in the aesthetic design of the visual environment, which we describe next.

1.5 Particle dynamics as a bioresponsive design medium

Viscereality is a bioresponsive VR system developed through an iterative design process, parts of which have been reported previously with respect to its design rationale, technical implementation, and interaction mechanics in human-computer interaction

contexts (Fejer et al. 2025; Fejer 2026). The present study provides its first controlled empirical evaluation, testing whether the system’s particle dynamics measurably shape subjective experience during breathwork. As described above, the system maps lung volume onto a surrounding particle field through interoceptive-exteroceptive sensory substitution (Goral et al. 2024b). Inhalation expands the environment outward, whereas exhalation contracts it inward. Existing implementations of this type of coupling typically use modelled surfaces, either presented as localized targets linked to the user (Wald et al. 2023) or projected onto the walls of a physical room (Goral et al. 2024b). In both cases, the user watches a representation change size, but a uniform surface that grows or shrinks does not, on its own, strongly convey that the surrounding space itself is approaching or receding; that impression depends on motion-based depth cues that arise when many elements move relative to one another (Rogers and Graham 1979). We designed Viscereality as a particle system specifically to afford this. The environment is composed of approximately 2,500 particles distributed on a geodesic icosphere, and the sense of spatial expansion and contraction arises from Gestalt grouping among these elements, specifically proximity, common fate, and continuity (Wagemans et al. 2012), rather than from a texture mapped onto a surface. Because each particle independently carries multiple visual properties (position, size, colour, transparency, motion trajectory), the same elements that produce the breath-coupled spatial dynamics can simultaneously encode additional biofeedback channels, such as heart rate variability coherence (Fejer 2026), without requiring separate visual layers. And because inter-particle interaction rules, specifically oscillator coupling across neighbours, generate emergent aesthetic properties such as dynamic symmetry and geometric coherence, these can be parameterised independently of the breathing mechanics, making the aesthetic texture of the environment an experimentally controllable variable. Throughout the breathing cycle, the particle field remains continuously active. Surface-wave oscillations, chromatic cycles, and shape fluctuations continue during inhalations, exhalations, and both breath-hold phases. The within-VR manipulation therefore does not switch visual motion on or off; rather, during breath holds the high-symmetry condition transiently aligns these ongoing particle cycles into coherent geometric organisations, whereas the asymmetric condition leaves the same dynamics desynchronised. To our knowledge, no existing interoceptive-exteroceptive sensory substitution system takes this particle-based approach. Aesthetics in VR are often treated as incidental to intervention design, yet recent work identifies them as among the strongest predictors of emotional response and experience quality in immersive environments (Marcolin et al. 2021; Cheiran et al. 2025). In Viscereality, aesthetics are not a cosmetic layer but emerge from the same particle dynamics that carry the respiratory coupling, drawing on a longer history of expressive and aesthetic design in particle-based systems applied in visual effects animation.

1.5.1 Particle systems as a generative medium

Particle systems were introduced by Reeves (1983) to represent visual phenomena that have no fixed boundary and no stable shape, including fire, smoke, clouds, and water. In his formulation, each particle is born with stochastically assigned properties (position, velocity, colour, size, transparency, lifetime), moves and changes over time,

and eventually vanishes. The overall form is never fully specified; it emerges from the aggregate of many simple elements, each governed by a small set of controllable parameters. This is the property that makes particle systems useful as a bioresponsive medium because each particle independently carries multiple visual channels, so a single population of elements can simultaneously respond to physiological signals through spatial position, colour shifts, size, transparency, or motion trajectory without requiring separate visual layers for each mapping. Building on this work, Reynolds (1987) introduced interaction between elements, enabling coordinated collective behaviour to emerge from local neighbour rules, whereas in a standard particle system each element responds only to global parameters and physics without awareness of its neighbours. Reynolds specified three local rules. They were to avoid collision with nearby elements, match their velocity, and maintain proximity to them. This became foundational for the ability of 3D animation to produce the fluid, coordinated group motion recognisable in flocking birds, schooling fish, or stampeding herds, all arising bottom-up from neighbour interactions without a centrally scripted choreography.

1.5.2 Particle systems as an affective medium

Research on abstract emotion representation shows that affective meaning can be conveyed through low-level perceptual parameters of abstract expressions (de Rooij et al. 2013). In particular, affective valence can be mapped through semantically minimal features such as motion speed and trajectory smoothness (Bartram and Nakatani 2009; Pollick et al. 2001; Burger and Toiviainen 2020), angular or rounded shape contours (Blazhenkova and Kumar 2018; Aronoff 2006), and line orientation (Damiano et al. 2023). In conceptual terms, Khatin-Zadeh et al. (2019) argue that motion provides an effective vehicle for representing emotion because both unfold dynamically over time, allowing changes in affective state to be expressed through movement, direction, and position in space. This claim has also been borne out empirically in swarm robotics, where the collective motion of distributed abstract disc-shaped agents has been shown to alter the observer’s state of flow and sense of time relative to observing a single moving agent (Kaduk et al. 2024). Dietz et al. (2017) found that higher speed in swarm motion increased perceived emotional intensity, while smooth trajectories, relative to jittery motion, were associated with more positive affect. Extending this approach, Santos and Egerstedt (2021) translated trajectory and motion into swarm choreographies for Ekman’s six basic emotions (Ekman 1993). Rounded and smooth configurations marked positive or low-arousal states, while angular, clustered, or dispersive formations with fast or slow trajectories marked negative and higher-arousal states. Participants identified the intended emotions above chance (Santos and Egerstedt 2021). What makes these findings from swarm robotics relevant here is that their expressive effects arise from relational motion among many simple elements rather than from the form of any individual agent. The same principle holds in *Viscerality*, where local coupling among particles governs trajectory speed, smoothness, synchrony, clustering, and dispersal, producing higher-order collective motion patterns from which affective qualities may emerge. The particle field can therefore be treated as a medium for conveying affective states through abstract visual dynamics rather than semantic or figurative cues.

1.5.3 Particle systems as an aesthetic medium

Beyond serving as a medium for affective mapping, we designed the particle field as an aesthetic means of rendering affective states through abstract spatiotemporal organization. Research on abstract visual art suggests that non-figurative forms evoke aesthetic and affective responses through the isolation of simple features, the grouping of many elements into emergent wholes, and the amplification of perceptual qualities through Gestalt grouping principles (de Rooij et al. 2013; Ramachandran and Hirstein 1999). In our particle system, these same grouping principles also help convey depth when approximately 2,500 ring-shaped particles expand and contract around the user. Because each particle is visually minimal, higher-order forms are not pre-given but emerge through local relations among neighbouring elements, allowing circles, lattices, polygons, and other geometric organisations to arise through changes in three-dimensional location. The appearance of such forms can itself become aesthetically salient, as the resolution of an initially indeterminate field into a coherent Gestalt is known to produce pleasurable moments of insight or ‘aesthetic Aha’ moments (Muth and Carbon 2013; Spehar and van Tonder 2017; Spee et al. 2024). In Viscerality, this made it possible to translate oscillator-level control into changes in overall visual order, symmetry, and coherence without moving to figurative imagery.

Another design influence was the geometric and kaleidoscopic qualities of hallucinatory and psychedelic imagery, which are often marked by geometric patterns, bilateral symmetry, and perceptual ambiguity (Klüver 1966; Billock and Tsou 2012; Lawrence et al. 2022; Rubin et al. 2010). This motivated our use of ring-shaped particles, as their overlaps can give rise to distinctive emergent patterns during the breathing cycle. Because these qualities emerge from shifting relations among many simple particles rather than from fixed symbolic forms, they remain open in meaning and foreground a defining affordance of the medium, namely its capacity to continually reorganise and transform its aesthetic structure over time.

In the context of VR design, Glowacki (2024) terms such non-figurative scenes a weak representational aesthetic, in contrast to photorealistic depictions that render scenes and bodies as recognizable real-world entities. In practice, this aesthetic has been applied in VR scenarios designed to induce altered states of consciousness through the representation of participants’ bodies as luminous, semi-corporeal avatar bodies composed of particle clouds with mergeable body boundaries, contributing to the malleability of self-experience and altered subjective states (Glowacki et al. 2020, 2022; Glowacki 2024; Vidal et al. 2025). Our approach adopts a related design aesthetic while focusing more specifically on diminishing the felt boundary between self and environment.

To evaluate the particle system’s capacity to manipulate aesthetic properties, we operationalized symmetry as an additional independent variable that varied across conditions. Previous studies have linked visual symmetry to more positive affect and perceptual fluency (Bertamini et al. 2013; Pecchinenda et al. 2014; Makin et al. 2012), while synchrony in group movement has been linked to greater enjoyment and more positive affective appraisal (Vicary et al. 2017; Mogan et al. 2017). We therefore asked whether higher degrees of symmetry would enhance positively valenced experiences

in the context of Viscerality (Johnson 2023). We were also interested in whether participants would be able to detect and discriminate between the two conditions.

1.6 Present Study

The present study provides the first controlled empirical evaluation of Viscerality with naive participants. We used a preregistered within-subjects design with $N = 39$. Each participant completed three 10-minute audio-guided box-breathing sessions in counterbalanced order. These were a high-symmetry condition, in which positive oscillator coupling transiently aligned ongoing particle oscillations into dynamic geometric coherence during breath-hold phases; an anti-symmetry condition, in which negative coupling produced a visually irregular particle field throughout; and a dark-screen control (see Section 2.3.4). The two particle conditions used identical particles, breath-coupled spatial dynamics, colour palette, and audio guidance, differing only in oscillator coupling parameters. The central within-VR contrast therefore concerned the aesthetic organisation of an otherwise identical breath-responsive scene.

We assessed four substantive outcome domains together with a methodological blinding assessment. Breathing adherence was quantified as the phase-locking value between the observed respiratory signal and an ideal reference trajectory. Self-related experience was assessed with three pictographic measures, namely body-boundary salience (Dambrun 2016), spatial frame of reference (Hanley and Garland 2019), and perceived self-size (Vidal et al. 2025). Altered states of consciousness were assessed retrospectively with the 11D-ASC (Studerus et al. 2010; Dittrich 1998). Blinding efficacy between the two active VR conditions was assessed retrospectively with a two-stage procedure. Before unmasking, participants completed 12 probe questions covering condition-specific, shared, and sham features. After being told that the two active scenes differed, they indicated in which order the high-symmetry and anti-symmetry scenes had been presented, together with confidence ratings. These responses were used to estimate condition-identification blinding with a standard single-group Bang blinding index and an exploratory confidence-weighted directional variant (Bang et al. 2004).

We hypothesised that, relative to the dark-screen control, the bioresponsive VR conditions would increase breathing adherence and altered-state intensity and would also influence the three preregistered self-related measures. Within VR, we further expected the high-symmetry condition to enhance positively valenced experiences and produce stronger alterations in self-related experience than the anti-symmetry condition. The preregistration did not specify a separate directional prediction for each pictographic outcome. We therefore treated the three outcomes as one family and controlled the false-discovery rate across them.

2 Methods

2.1 Participants

Participants were recruited from the University of Konstanz psychology student participant pool and received course credit for participation. The study used a within-subject

507 design in which each participant completed all three conditions in one of six coun-
508 terbalanced block-order permutations, with a preregistered target of 6 completers per
509 permutation and 36 total. We did not systematically assess prior experience with
510 virtual reality, meditation, breathwork, or psychedelics. We use recruitment-naïve to
511 mean that participants came from the general psychology participant pool through an
512 advertisement for a VR breathing-biofeedback study rather than an altered-state or
513 experience-seeking recruitment channel; it does not mean that prior experience was
514 screened or absent. We tested 47 participants; 8 were excluded based on breathing
515 signal quality (phase-locking value (PLV) outlier criterion described below), yielding
516 a final sample of $N = 39$ (6–7 per permutation), preserving approximately equal rep-
517 resentation across all six condition sequences. Among the included participants, mean
518 age was 20.8 years; 37 were female (mean age 20.7 years) and 2 were male (mean age
519 23.0 years).

520

521 2.2 Design, Conditions, and Procedure

522

523 Participants were seated in a bean bag, wore noise-blocking headphones, held a con-
524 troller against their abdomen, and completed all breathing blocks, questionnaires,
525 and outcome measures fully immersed in virtual reality. The three-session procedure
526 was automated end-to-end in the headset: once the headset was put on, participants
527 followed in-VR instructions for the breathing blocks and all post-block outcome mea-
528 sures while remaining seated, with minimal experimenter intervention beyond initial
529 setup and debriefing. Before the experimental sequence, participants completed a
530 1-minute guided breathing practice block for acclimation, familiarizing them with con-
531 troller placement and the paced-breathing rhythm. Each participant then completed
532 three conditions, with session order assigned automatically by random selection from
533 the six counterbalanced block-order permutations. Because the dynamic symmetry
534 manipulation was embedded in the software-controlled scene logic and the condi-
535 tion sequence was assigned automatically at runtime, neither the participant nor the
536 experimenter knew which condition would appear in which session. In all conditions,
537 participants followed a 10-minute audio-guided box-breathing exercise (sourced from
538 a publicly available recording) consisting of repeated 4-4-4-4 cycles with four seconds
539 each of inhalation, breath hold, exhalation, and breath hold. In the high-symmetry
540 condition, the bioresponsive particle environment formed dynamic symmetric patterns
541 during breath-hold phases. In the asymmetric condition, visual features of the particles
542 remained asymmetric to one another throughout. In the dark-screen control condition,
543 the particle environment was not displayed; participants viewed a dark field contain-
544 ing only a central fixation marker that also provided breathing-performance feedback
545 (Section 2.3.4). We refer to this condition as the “dark-screen control” throughout.
546 After each block, participants completed the full assessment battery (questionnaires
547 and pictographic scales) while remaining in virtual reality. Figure 1 summarizes the
548 full procedure, including the repeated condition loop and blinding sequence, using the
549 same condition-color mapping as the Results figures.

550

551

552




Fig. 1 Study procedure. After onboarding, each participant completed three 10-minute box-breathing sessions (dark-screen control, symmetric VR, asymmetric VR) in one of six counterbalanced orders, with the full assessment battery administered in VR after each block. A blinding questionnaire followed the third block before the study concluded.

2.3 Virtual-Reality Intervention

Viscereality is a bioresponsive virtual-reality system that maps respiratory signals to dynamic visual feedback within an immersive 3D particle environment (Fejer et al. 2025; Fejer 2026). A video summary of the system is available online.¹ The system runs on a Meta Quest 3 headset (standalone, untethered) and was built in Unity, with CPU-optimized parallel processing (Unity Job System with Burst compilation) maintaining a stable 72 Hz frame rate. The two active virtual-reality conditions used the same visual environment, audio guidance, and logging pipeline; they differed only in a single engine parameter controlling oscillator coupling dynamics. The dark-screen control used identical audio guidance and data logging but omitted the particle visualization. We focus here on the experiment-relevant functional logic; fuller technical

¹<https://www.youtube.com/watch?v=Qm-lmC-xYCI>; archived at <https://osf.io/4enfz>

599 and implementation details of the Viscereality engine are described in the prior system
600 papers (Fejer et al. 2025; Fejer 2026).

601

602 **2.3.1 Breath Detection**

603

604 Breath detection follows the approach introduced by Flowborne (Rockstroh et al.
605 2021), using the VR controller’s positional tracking to infer respiratory displacement
606 without additional sensors. During an initial calibration, the user places the controller
607 against the lower abdomen to establish a tracking axis. Each frame, the algorithm reads
608 the controller’s position and orientation, projects motion onto this breathing-relevant
609 axis, and updates a short-term versus long-term smoothed motion estimate. The dif-
610 ference between these smoothed signals functions as a breathing trend signal. Positive
611 values indicate inhalation, negative values indicate exhalation, and near-zero values
612 indicate a pause. Samples are rejected as tracking artifacts if controller rotation is too
613 abrupt or if the inferred trend is implausibly large. The resulting classification (inhale,
614 exhale, pause, or bad tracking) drives the visual feedback pipeline described below
615 (Figure 2). Detailed implementation choices and parameterization are documented in
616 the Viscereality technical descriptions (Fejer et al. 2025; Fejer 2026).

617

618 **2.3.2 Sphere Radius and Visual Feedback**

619

620 The user is positioned at the center of a large particle-covered sphere. The primary
621 feedback loop couples sphere radius to breathing. Inhalation causes the sphere to
622 expand outward, exhalation causes it to contract inward, and breath holds main-
623 tain the current radius. Functionally, the controller-derived breathing-state label is
624 converted into a signed expansion/contraction command whose magnitude is scaled
625 to the configured radius range and the intended inhale/exhale timing. That com-
626 mand updates a target sphere radius, and the rendered radius is then smoothed with
627 a damped interpolation step (0.25 s smooth time; overshoot prevention enabled). In
628 practice, this means participants do not see a raw controller-motion trace; they see a
629 bounded, temporally stable visual signal that preserves breathing direction while sup-
630 pressing jitter. Full implementation details of the radius controller are described in
631 the prior Viscereality system reports (Fejer et al. 2025; Fejer 2026).

632

633

634

635

636

637

638

639

640

641

642

643

644




Fig. 2 Signal processing pipeline from abdomen-mounted controller motion to rendered sphere radius. Raw 3D movement is projected onto a calibrated breathing axis, classified into inhale, exhale, or hold phases, and mapped onto expansion and contraction dynamics that update the visible sphere at a fixed velocity.

The sphere surface is composed of approximately 2,500 view-aligned ring particles distributed on an icosphere mesh for uniform geodesic spacing. Each particle is rendered with a radial gradient that produces depth cues through overlap, and during expansion and contraction, particles generate cone-like tracer effects that reinforce the perception of three-dimensional movement. Surface wave oscillations, structured noise, and unsynchronized chromatic cycles maintain continuous visual dynamics across all breathing phases, preventing the environment from appearing static during breath holds.

691 2.3.3 Oscillator Network and Condition Manipulation

692 The second layer of the system implements a Kuramoto oscillator network across the
693 sphere’s vertices (Acebrón et al. 2005; Rodrigues et al. 2016). Each particle has an
694 internal oscillation phase (analogous to a clock angle) and interacts with its neigh-
695 bors through coupling weights that determine whether nearby particles tend toward
696 synchronization or away from it. If the coupling weight is positive, neighbors are
697 pulled toward shared timing (phase attraction). If negative, the interaction becomes
698 *anti-coupling*, and neighbors are pushed away from alignment (phase repulsion).
699

700 The network uses a two-tier neighbor structure reflecting the icosphere topology.
701 Most vertices have 6 immediate (tier-1) neighbors and 12 next-nearest (tier-2) neigh-
702 bors, while 12 vertices at the original icosahedron poles have 5 tier-1 neighbors. At
703 each update step, each oscillator advances according to its own intrinsic rhythm plus
704 a neighborhood interaction term. In plain terms, the algorithm compares an oscilla-
705 tor’s current phase with those of its neighbors and then adjusts it toward or away
706 from them depending on the sign and magnitude of the tier-specific coupling weights.
707 This is the standard Kuramoto logic (phase-offset-dependent attraction/repulsion),
708 implemented here with two neighbor tiers and a global coupling gain, and updated in
709 parallel each frame together with particle position, size, and color. The formal imple-
710 mentation and parameterization are described in the prior Viscerality publications
711 (Fejer et al. 2025; Fejer 2026).

712 The two active conditions differed in how coupling weights were assigned. In the
713 high-symmetry condition, tier-1 and tier-2 coupling weights were interpolated dynam-
714 ically as a function of a coherence scalar (0–1), together with a coherence-dependent
715 frequency scaling term. Verbally, low coherence pushes the system toward anti-coupled,
716 irregular dynamics, whereas higher coherence progressively shifts interactions toward
717 phase attraction and more synchronized motion. As a result, particle motion, color,
718 and size periodically coordinate into coherent geometric patterns, such as flower-of-
719 life-like lattices, during breath-hold phases before dissolving back into disorder when
720 breathing resumes.

721 In the asymmetric condition, the system did not use this dynamic interpolation.
722 Instead, coupling weights were held at fixed negative values (anti-coupling), so local
723 interactions were permanently biased toward phase repulsion and the particle field
724 remained visually irregular throughout. The manipulation was thus dynamical rather
725 than geometric. The same scene, particles, and audio were used in both conditions,
726 and only the oscillator interaction rules changed.

727 2.3.4 Breathing Performance Indicator

728 All three conditions displayed a minimal breathing performance indicator at the cen-
729 ter of the participant’s field of view. This consisted of a white pentagon ring whose
730 individual line segments scaled with sphere radius: at half-volume (midpoint between
731 full inhale and full exhale), the pentagon was fully visible; as the participant inhaled or
732 exhaled toward the extremes, the segments progressively shortened and disappeared
733 entirely at maximum and minimum volume. The indicator thus functioned as an unob-
734 trusive loading-bar-style monitor of breathing excursion. Participants were instructed
735
736

that the goal was to make the pentagon disappear completely on each inhalation and exhalation, confirming sufficient breathing depth.




Fig. 3 Central breathing-performance indicator shown in all three conditions. The pentagon ring was fully visible at the neutral sphere radius and progressively disappeared toward full contraction or full expansion, providing condition-matched feedback about breathing excursion.

2.4 Outcome Measures

We retained the five preregistered measurement blocks used in the assessment battery, but organize them here into four recurring outcome families that also structure the Results section. These were adherence to box breathing, self-related measures, altered states of consciousness, and the blinding questionnaire. Within the altered-states family, the main text presents the derived 3D-ASCr composites as a dimensionality-reduced summary; Online Resource 1, Section S3 reports the complete preregistered 11D-ASC analyses.

2.4.1 Adherence to Box Breathing

Adherence quantifies how closely each participant’s breathing followed the instructed 4-4-4-4 box-breathing rhythm across each 10-minute block. The preregistered plan was to compute this from discrete phase labels produced by the breath-detection algorithm (Section 2.3.1), which classifies each frame as inhale, exhale, hold, or bad tracking. The intended metric was phase-duration error, defined as the absolute deviation from the 4-second target per breathing phase, averaged across valid cycles per condition, with participant exclusion if fewer than 70% of cycles were valid or if mean error exceeded 2 standard deviations above the sample mean.

The recorded phase sequences did not yield the expected clean succession of 4-second phases. Rather than discrete blocks of inhale, hold, exhale, and hold, the recorded data consisted of highly fragmented segments interspersed with brief transitional or ambiguous intervals. Computing phase-duration error from this data required first reconstructing plausible box-breathing cycles from the raw phase stream, a step that was not anticipated at preregistration. We explored several reconstruction strategies, including rule-based merging of adjacent micro-segments and temporal smoothing algorithms, but each approach introduced its own arbitrary parameter choices, such as minimum segment duration thresholds, smoothing window widths, and rules for combining adjacent same-phase fragments, that could materially influence the resulting error estimates. Because these post-hoc binning decisions carried more researcher degrees of freedom than the original preregistered metric implied, we adopted an alternative approach.

We quantified adherence as the phase-locking value (PLV) between two continuous time series. One was the sphere-radius signal that participants actually saw as visual feedback during the breathing session. The other was an ideal reference radius trajectory derived from the audio-instruction timestamps for perfect 4-4-4-4 breathing. PLV measures how consistently the phase relationship between two signals remains stable over time. A value near 1 indicates that the participant’s breathing closely tracked the instructed rhythm throughout the block. A value near 0 indicates little consistent alignment. This approach circumvents the phase-reconstruction problem entirely, because it evaluates alignment at the level of the continuous visual feedback signal rather than attempting to recover discrete phase boundaries from a fragmented controller trace. PLV thus captures the same underlying construct as the preregistered metric, namely how closely breathing tracked the instructed rhythm, while avoiding the measurement artifacts that any post-hoc phase-segmentation strategy would have introduced. Figure 4 illustrates the relationship between observed and reference trajectories for a representative participant; the fragmented phase-label structure visible in the lower bars of each panel motivates the PLV-based approach described above.

Because PLV does not decompose into per-cycle validity counts, the preregistered 70%-valid-cycle plus 2-SD mean-error exclusion rule could not be transferred directly. Outlier exclusion instead used a lower-tail global IQR rule on condition-level PLV values, elevated to participant-level exclusion if any condition was flagged. Online Resource 1, Section S2 describes PLV computation and quality screening in detail.




Fig. 4 Breathing dynamics for a representative participant (P01) across all three conditions. Observed sphere-radius traces (orange) are overlaid on the ideal 4-4-4-4 reference (blue), with per-block PLV values shown at right. Fragmented phase-label bars below each time series illustrate the segmentation noise that motivated the switch from phase-duration error to PLV as the adherence metric. Equivalent figures for all participants are provided in the OSF repository.

2.4.2 Self-Related Measures

Three single-item pictographic measures assessed self-related experience after each condition. The adapted Perceived Body Boundaries item (Dambrun 2016) presents seven silhouettes progressing from diffuse (A) to sharply delineated (G). Responses were mapped from A–G to 1–7, so higher analyzed scores indicate stronger boundary salience. For construct-facing exploratory correlations, the item was reverse-scored so that higher values indicate greater boundary dissolution. The Spatial Frame of Reference Continuum (SFoRC; Hanley et al. 2020) depicts a figure in concentric spatial zones from body-centered to environment-encompassing, indexing the degree to which the spatial sense of self extended beyond the body. The Small Self pictographic item was adapted from Vidal et al. (2025), who presented it as an ad hoc item. We included it because Viscerality is organized around a spatial metaphor and the item represents self-related experience through the spatial size of the self relative to its surroundings. PBBS and Small Self use A–G responses mapped to 1–7, whereas SFoRC uses A–H responses mapped to 1–8. Each measure yields a single ordinal score per condition. We recreated all three pictographs as SVG graphics for VR presentation, using the previously published versions as visual references. The vector format allowed direct and flexible control over graded visual properties, including transparency, line width, dash spacing, relative figure size, and spatial extent, while enabling sharp rendering at headset resolution without the resolution loss associated with scaling raster images. The SVG source was refined with large language model assistance. The OSF reproducibility package (<https://osf.io/64mwk/>) provides the recreated stimulus set, SVG source files, administered item keys, and study materials; Online Resource 1, Section S6 provides the corresponding reproducibility map.

2.4.3 Altered States of Consciousness

3D-ASC_r composites. The revised 3D-ASC_r higher-order factor model became available after preregistration. It provides a parsimonious contemporary summary of the same preregistered 11D-ASC responses by organizing them into Positive, Distressing, and Perceptual Effects. This level of aggregation matches our broad theoretical interest in positively versus negatively valenced experience, while retaining a perceptual dimension, and avoids overinterpreting lower-order dimensions for which no specific predictions were made. We therefore use the 3D-ASC_r composites as the main-text summary. Online Resource 1, Section S3 reports complete analyses of all 11 preregistered 11D-ASC dimensions.

11D-ASC Questionnaire. We assessed altered states with the 11-Dimensional Altered States of Consciousness Rating Scale (11D-ASC; 42 items, 0–100 visual analogue scales), a validated lower-order factorization of the OAV/APZ framework (Studerus et al. 2010; Dittrich 1998; Hovmand et al. 2024; Prugger et al. 2022). The 11 subscales were Experience of Unity (EU), Spiritual Experience (SE), Blissful State (BS), Insightfulness (IS), Disembodiment (DIS), Impaired Control and Cognition (ICC), Anxiety (ANX), Complex Imagery (CI), Elementary Imagery (EI), Audio-Visual Synesthesia (AVS), and Changed Meaning of Percepts (CMP). Participants completed one condition-specific rating after each block.

875 **Temporal Experience Tracers (Exploratory).** Temporal experience tracers
876 provide psychometric time-series ratings of the same 11 experiential dimensions.

877 For each dimension, one prompt con-
878 solidated the constituent questionnaire
879 items into a construct-level description.
880 After each condition, participants retro-
881 spectively drew the 11 intensity curves
882 on a world-space time-by-intensity sur-
883 face using the VR controller. Controller
884 position was projected onto the draw-
885 ing plane, input was gated through a
886 trigger hold, and points were emitted
887 at constant spatial spacing regardless of
888 hand speed. Each trace was reduced to
889 peak intensity, representing the strongest
890 reported moment, and area under the
891 curve (AUC), representing intensity accu-
892 mulated across reconstructed session time
893 (Figure 5). We focused on these two sum-
894 maries because no hypothesis required
895 finer temporal features such as onset, tim-
896 ing, slope, or variability. We treat the
897 tracer analyses as exploratory; Online
898 Resource 1, Section S5 provides the com-
899 plete implementation, data reduction,
900 analyses, and results.

901

902 2.4.4 Blinding Questionnaire

903

904 Blinding was assessed in two phases. In the pre-unmasking phase, participants were
905 told that the study investigated how subtle differences in visual aesthetics might influ-
906 ence perception in virtual reality. We constructed a 12-item ad hoc masked-comparison
907 questionnaire for this study. For each item, participants chose whether the feature was
908 stronger in Scenario 1, stronger in Scenario 2, equal in both scenarios, or absent in both
909 scenarios. The chance baseline was 0.25. Scenario labels referred to the two virtual-
910 reality conditions in encounter order. The item set covered three hidden domains.
911 These were manipulation-relevant symmetry/coordination cues that genuinely differed
912 between the two VR conditions, matched comparison features that were present simi-
913 larly in both conditions and therefore should be judged as equal, and foil (sham)
914 features that were not present in either condition and therefore should be judged as
915 absent. There were four items per domain.

916 In shorthand, the four symmetry-related items asked whether one scene more
917 strongly showed transient coordination and ordering of the particle field, including
918 particles moving as a coordinated group, becoming more organized or structured at
919 certain moments, showing wave-like coordinated motion-and-color changes, or shift-
920 ing from chaotic to ordered movement during breath holds. The four neutral matched

**Temporal Trace Profiles:
Peak versus Area Under the Curve**




Fig. 5 Temporal experience tracers: peak and cumulative intensity. These hypothetical profiles illustrate information retained by psychometric time-series intensity ratings but not by a single retrospective visual-analogue-scale response. The red profile has a higher momentary peak, whereas the broader green profile has a larger area under the curve (AUC), reflecting greater cumulative intensity. Analyses in this paper focus on these two summaries because no hypothesis required finer-grained temporal features.

items asked about features that were present in both VR conditions to a similar degree, namely particles becoming brighter during breath holds, leaving faint motion trails during exhalation, remaining in gentle ongoing motion across the surface, or appearing to rise and fall like ocean waves. The four sham items were constructed as absent cross-modal or pulsing foils, asking about the music becoming quieter, the music seeming to slow down, particle motion and color changing in response to the music, or particles showing a size-based pulsing effect. To keep administration uniform while avoiding an obviously grouped structure, items from all three domains were intermixed in a single randomized order that was generated once in advance and then presented in that same order to all participants.

In the post-unmasking phase, participants were told that one condition featured brief visual synchronization during breath holds while the other remained irregular throughout. They indicated which session was more symmetric via a three-option forced choice (*first*, *second*, or *I don't know*), followed by a confidence rating.

2.5 Preregistration and Reproducibility

The study was preregistered on AsPredicted (#262,545). The unchanged registry record, all deidentified participant-level data, statistical analysis scripts and outputs, figure-generation scripts, and study materials were uploaded to OSF as a reproducibility package (<https://doi.org/10.17605/OSF.IO/64MWK>). We retained all preregistered design elements and outcome families. The preregistered analysis plan specified a repeated-measures ANOVA with condition as the within-subjects factor and presentation order as a covariate for each outcome, two planned contrasts (pooled VR versus dark-screen control, and symmetric versus asymmetric), and parallel Bayesian analyses. Adherence was to be quantified as phase-duration error with a cycle-validity exclusion rule, as described and motivated in Section 2.4.1. We deviated from this plan in several respects. Online Resource 1, Section S1 summarizes the deviations from preregistration and their rationale.

Adherence metric. Phase-duration error was replaced by PLV because the recorded phase-label data were too fragmented to yield reliable per-phase error estimates without introducing arbitrary binning decisions (see Section 2.4.1 for a full account). The associated exclusion rule was changed from 70%-valid-cycle plus 2-SD mean-error to a lower-tail global IQR rule on condition-level PLV values, because PLV does not decompose into per-cycle validity counts.

Omnibus condition test. For every non-PLV outcome, the preregistered order-adjusted repeated-measures analysis tests the condition effect after accounting for participant and numeric block position. It is implemented as a partial- F comparison between a participant-and-order model and the same model plus condition. Participant fixed effects provide within-participant blocking, and block position adjusts for presentation order. The finalized PLV omnibus remains unchanged. A condition-only repeated-measures ANOVA without the order covariate is retained in the OSF outputs as a sensitivity check.

Planned contrasts. The two preregistered planned contrasts were implemented as standalone paired t -tests, which yield the same paired contrast statistics as ANOVA

967 follow-ups in this within-subject setting. These serve as the confirmatory anchor for all
968 multi-test outcome families in the Results section, and for significant planned contrasts
969 in the altered-state families we additionally report paired common-language effect sizes
970 (CLES), defined as the proportion of participants whose score was higher in the VR
971 condition than in the dark-screen control (McGraw and Wong 1992).

972 **Robustness and multiplicity.** Because Shapiro–Wilk checks indicated non-
973 normality for several condition-level and difference-score distributions, we added
974 Wilcoxon signed-rank and Friedman tests as secondary distribution-free checks. The
975 preregistration did not specify a multiplicity procedure. Nominal p values there-
976 fore govern the primary interpretation of the preregistered planned contrasts, with
977 Benjamini–Hochberg q values reported as sensitivity information within the 11D-
978 ASC, 3D-ASCr, and self-related outcome families. FDR-adjusted q values govern
979 exploratory and post hoc screens. The OSF package contains the complete adjusted
980 and distribution-free output; the paper reports material divergences from the primary
981 planned-contrast interpretation.
982

983 **3D-ASCr composites.** The revised 3D-ASCr scoring scheme (Stocker et al. 2025)
984 became available after preregistration. Because these composites are deterministic
985 averages of the preregistered 11D-ASC subscale scores, they add no new participant
986 data but do constitute a post-registration summary choice. We therefore use them to
987 organize the main-text narrative while retaining the complete preregistered 11D-ASC
988 analyses in Online Resource 1, Section S3.

989 **Bayesian analyses.** Bayesian analyses were preregistered and run in parallel on the
990 same included sample ($N = 39$). Omnibus Bayes factors compared a point-null model
991 without condition (H_0 : outcome $\sim$ order + participant random intercept) against
992 a model that additionally included condition (H_1 : outcome $\sim$ condition + order +
993 participant random intercept), using the BayesFactor R package with explicit prior
994 settings (fixed-effect $r = 0.50$, continuous-covariate $r = \sqrt{2}/4 \approx 0.354$, random-effect
995 nuisance $r = 1.00$; Morey and Rouder 2024; van Doorn et al. 2021; Kruschke 2021).
996 For the two planned paired contrasts, we compared H_0 ($\delta = 0$) against two-sided H_1
997 ($\delta \neq 0$) using a Bayesian paired t -test with the default Cauchy prior on standardized
998 effect size ($r = \sqrt{2}/2 \approx 0.707$; Morey and Rouder 2024; van Doorn et al. 2021). For
999 EU, BS, and AVS, the preregistered joint ordering symmetric > asymmetric > dark-
1000 screen control was tested after block-position adjustment with an encompassing-prior
1001 Bayes factor against the equal-means null. Under the exchangeable condition-factor
1002 prior, each of the six strict condition orderings had prior probability 1/6. We report
1003 BF_{10} values (and $BF_{01} = 1/BF_{10}$ when the data favor the null) as relative evidence
1004 rather than as posterior probabilities or effect sizes (van Doorn et al. 2021; Kruschke
1005 2021). Posterior median standardized effect sizes with 95% credible intervals and prior-
1006 robustness checks are reported in the OSF repository.
1007

1008 **Blinding analyses.** Pre-unmasking accuracy was scored using the documented
1009 answer key for the four symmetry, four neutral, and four sham items and across
1010 all 12 items. Mean accuracy was compared with the 0.25 chance baseline using
1011 Wilcoxon signed-rank tests, with Benjamini–Hochberg FDR correction across the
1012 four comparisons. Post-unmasking guesses were classified as correct (R), incorrect

(W), or uncertain (U) from the participant’s selected active-VR session and condition order. The standard single-group Bang blinding index was calculated as $BBI = (R - W)/(R + W + U)$ (Bang et al. 2004; Kolahi et al. 2009), with a participant-bootstrap 95% confidence interval. Effective blinding was considered established only if the entire interval fell within $[-0.2, 0.2]$ (Qin et al. 2024; Kolahi et al. 2009). We also report an exploratory confidence-weighted directional index, calculated as the confidence-weighted proportion of correct directional guesses minus 0.5. The standard Bang interpretation bands were not applied to this modified index. To test whether pre-unmasking symmetry accuracy predicted subsequent condition identification, we calculated a point-biserial correlation between the 0–4 symmetry score and final-guess correctness, classified as correct versus incorrect or uncertain, with an exact two-sided permutation p value.

Exploratory analyses. We fit condition $\times$ PLV interaction models to test whether breathing synchronization moderated subjective effects. We also conducted an exploratory analysis of whether VR-related changes in the pictographic measures covaried with changes in ASC outcomes. For focused, construct-based interpretation, we examined SFoRC in relation to Experience of Unity and Disembodiment, including items concerning unity with the environment, floating, bodylessness, and altered self-location outside the body. We examined perceived body-boundary dissolution primarily in relation to Disembodiment and the same bodily-self items, and Small Self in relation to Experience of Unity, Spiritual Experience, and Blissful State. These thematic selections were post hoc and guided interpretation rather than defining separate inferential families. We additionally conducted broad exploratory screens across all 3D-ASCr composites, 11D-ASC subscales, and ASC items. Because only SFoRC showed a reliable pooled-VR-versus-dark-screen-control condition effect, the correlations were treated as exploratory tests of cross-measure correspondence rather than as evidence that all three pictographic measures changed under VR. Online Resource 1, Section S4 reports the complete analysis, with false-discovery-rate correction across all 9 composite, 33 subscale, and 126 item-level associations, together with symmetric-minus-asymmetric change correlations and order-adjusted participant-aware condition-interaction models.

3 Results

Across the four recurring outcome families, condition effects were consistently driven by the contrast between virtual-reality conditions and the dark-screen control. No outcome showed a reliable symmetric-versus-asymmetric difference. We report the results in the same family order used in Methods: adherence, self-related measures, altered states of consciousness, and the blinding questionnaire. For the non-PLV multi-test families, confirmatory interpretations are anchored in the preregistered planned contrasts evaluated as paired t -tests, with the registered order-adjusted participant-blocked partial- F omnibus noted for each family as context; PLV retains its finalized mixed-effects omnibus. Wilcoxon signed-rank robustness checks, Friedman omnibus tests, and FDR-adjusted q values are documented in the OSF repository and are mentioned here only when they materially qualify the paired t pattern.

1059 3.1 Sample and Exclusions

1060 The dataset contained records from all 47 tested participants. Condition-level PLV val-
1061 ues from all participants were used to apply the quality-screening procedure described
1062 in Section 2.4.1 and illustrated in Figure 6. This procedure excluded 8 participants
1063 and yielded a final analytic sample of $N = 39$. The 39 included participants each con-
1064 tributed a complete within-subject triad and constituted the final sample used for all
1065 inferential analyses.
1066

1067 3.2 Outcome Families

1069 3.2.1 Adherence (PLV)

1070 To test whether visual biofeedback improved adherence to the guided box-breathing
1071 rhythm, we compared PLV across conditions. Mean PLV was $M = 0.787$ ($SD =$
1072 0.141) for the dark-screen control, $M = 0.836$ ($SD = 0.127$) for symmetric VR, and
1073 $M = 0.860$ ($SD = 0.094$) for asymmetric VR. The planned pooled-VR-versus-dark-
1074 screen-control contrast confirmed this difference, $t(38) = 3.47$, $p = .001$, $d_z = 0.56$,
1075 95% CI for the mean PLV difference $[0.026, 0.097]$. The symmetric-versus-asymmetric
1076 contrast was not significant, $t(38) = -1.13$, $p = .267$, $d_z = -0.18$. The order-adjusted
1077 mixed-effects omnibus supported a condition effect, $\chi^2(2) = 12.66$, $p = .002$. The
1078 corresponding Bayesian planned contrasts showed that the pooled-VR-versus-dark-
1079 screen-control difference was about 24.1 times more likely than no difference ($BF_{10} =$
1080 24.11), whereas the symmetric-versus-asymmetric data were about 3.2 times more
1081 likely under a no-difference model ($BF_{01} = 3.22$).
1082

1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104




Fig. 6 Controller-derived feedback synchrony (PLV) by condition for the selected sample ($N = 39$), with condition-wise distributions and paired individual trajectories in gray. The dashed red line marks the pooled lower-tail IQR exclusion cutoff (0.366); hollow circles and red crosses indicate flagged participants and their below-cutoff values, respectively. If any single condition fell below the cutoff, the participant was excluded entirely from all analyses across all conditions.

3.2.2 Self-Related Measures

To assess whether the breath-responsive virtual environment shifted subjective self-representation, we compared three pictographic measures across conditions. Only the spatial frame-of-reference continuum showed a condition effect. Participants reported a more spatially extended sense of self during virtual-reality conditions, with $M = 3.15$ ($SD = 1.41$) for the dark-screen control, $M = 4.18$ ($SD = 1.82$) for symmetric VR, and $M = 4.26$ ($SD = 1.73$) for asymmetric VR. The planned pooled-VR-versus-dark-screen-control contrast confirmed the effect, $t(38) = 4.28$, $p < .001$, $d_z = 0.69$, 95% CI [0.56, 1.57]. The symmetric-versus-asymmetric contrast was not

significant, $t(38) = -0.32$, $p = .752$. The registered order-adjusted omnibus result is $F(2, 75) = 10.99$, $p < .001$, $q_{FDR} < .001$, $\eta_p^2 = .227$. Neither perceived body boundary salience ($t(38) = -0.17$, $p = .864$) nor small self ($t(38) = 1.39$, $p = .173$) showed a significant VR-versus-dark-screen-control contrast, and no self-related measure showed a significant symmetric-versus-asymmetric contrast. Figure 7 shows the response formats together with the condition-wise distributions and paired participant trajectories. Online Resource 1, Section S4 provides the complete registered frequentist and Bayesian condition-analysis inventory for all three pictographic measures.




Fig. 7 Condition-wise distributions for the three pictographic self-related measures ($N = 39$). The upper row shows the administered response formats, and the lower row shows condition-wise distributions, paired participant trajectories, and means (diamonds), from left to right: Perceived Body Boundaries, the Spatial Frame of Reference Continuum (SForC), and Small Self. Perceived Body Boundaries is displayed in its administered A–G direction, from diffuse to sharply delineated, so higher scores indicate stronger boundary salience. Only SForC showed a reliable pooled-VR-versus-dark-screen-control difference. Perceived Body Boundaries and Small Self showed no reliable pooled-VR-versus-dark-screen-control differences, and no pictographic measure differed reliably between the symmetric and asymmetric VR conditions. Colors indicate the dark-screen control, symmetric VR, and asymmetric VR.

Bayesian comparisons agreed with this pattern at the planned-contrast level. For spatial frame of reference, the observed data were approximately 204 times more likely under a pooled-VR-versus-dark-screen-control difference model than under the no-difference model ($BF_{10} = 204.44$). Perceived body boundaries favored the no-difference model ($BF_{01} = 5.71$), whereas the Small Self result provided only weak and inconclusive evidence in the null direction ($BF_{01} = 2.39$).

Exploratory pictograph–ASC change-score analyses are reported in Online Resource 1, Section S4. Their clearest pattern was coordinated participant-level variation between spatial self-extension and positive or unity-related experience, but these post hoc associations do not alter the primary condition-effect conclusions.

3.2.3 Altered States of Consciousness

We report the altered-state family in the main text using the 3D-ASCr composites as a dimensionality-reduced summary. Online Resource 1, Section S3 reports the complete preregistered 11D-ASC analyses. The clearest pattern is stronger positive and perceptual VR-related effects, with a weaker distress-related component and no reliable symmetric-versus-asymmetric difference.

3D-ASCr composites.

Summarizing the 11D-ASC data through the 3D-ASCr composites (see Section 2.4.3 for derivation), the three higher-order dimensions showed the following pattern. Perceptual Effects had $M = 11.41$ ($SD = 12.61$) in the dark-screen control, $M = 27.25$ ($SD = 19.92$) in symmetric VR, and $M = 25.59$ ($SD = 17.24$) in asymmetric VR. Positive Effects had $M = 18.88$ ($SD = 15.39$) in the dark-screen control, $M = 25.64$ ($SD = 19.48$) in symmetric VR, and $M = 26.46$ ($SD = 17.84$) in asymmetric VR. Distressing Effects had $M = 12.33$ ($SD = 10.26$) in the dark-screen control, $M = 17.33$ ($SD = 12.66$) in symmetric VR, and $M = 17.02$ ($SD = 13.18$) in asymmetric VR. All three composites showed significant pooled-VR-versus-dark-screen-control planned contrasts: Perceptual Effects $t(38) = 4.90$, $p < .001$, $d_z = 0.79$; Positive Effects $t(38) = 3.22$, $p = .003$, $d_z = 0.51$; Distressing Effects $t(38) = 2.56$, $p = .014$, $d_z = 0.41$. Expressed on the native 0–100 scale, pooled VR increased Perceptual Effects by 15.0 points, Positive Effects by 7.2 points, and Distressing Effects by 4.8 points. These corresponded to a large effect for Perceptual Effects, a moderate effect for Positive Effects, and a smaller effect for Distressing Effects, with 76.9%, 69.2%, and 76.9% of participants, respectively, scoring higher in VR than in the dark-screen control.

No composite showed a symmetric-versus-asymmetric difference (all $p \geq .357$). The registered order-adjusted omnibus results are $F(2, 75) = 19.63$, $p < .001$, $q_{FDR} < .001$, $\eta_p^2 = .344$ for Perceptual Effects; $F(2, 75) = 7.56$, $p = .001$, $q_{FDR} = .002$, $\eta_p^2 = .168$ for Positive Effects; and $F(2, 75) = 4.64$, $p = .013$, $q_{FDR} = .013$, $\eta_p^2 = .110$ for Distressing Effects. Bayesian omnibus comparisons favored condition models for Perceptual Effects ($BF_{10} = 121,810$), Positive Effects ($BF_{10} = 30.49$), and, more modestly, Distressing Effects ($BF_{10} = 3.45$). The planned Bayesian contrasts showed the same hierarchy. For pooled VR versus control, the observed data were approximately 1,178 times more likely under a difference model for Perceptual Effects ($BF_{10} = 1,178.41$), 13 times more likely for Positive Effects ($BF_{10} = 12.96$), and only 3 times more likely for Distressing Effects ($BF_{10} = 3.02$). For symmetric versus asymmetric VR, the observed data favored the no-difference models by factors of approximately 3.9, 5.1, and 5.7, respectively ($BF_{01} = 3.87$, 5.07, and 5.66; Figure 8). Thus, the Perceptual and Positive pooled-VR effects had clear relative support, whereas the smaller Distressing effect had only modest Bayesian support.




Fig. 8 Condition-wise distributions for the three 3D-ASCr composites (N = 39). All three dimensions increased in VR relative to the dark-screen control, with the largest effect for Perceptual Effects; no symmetric-versus-asymmetric differences emerged.

11D-ASC questionnaire.

Complete condition descriptives and preregistered frequentist and Bayesian results for all 11 lower-order ASC dimensions, including the registered directional hypotheses, are reported in Online Resource 1, Section S3.

3.2.4 Blinding Questionnaire

Pre-unmasking responses showed selective but incomplete sensitivity to the symmetry manipulation. Mean symmetry-item accuracy was 44.2%, equivalent to approximately 1.77 of four items, and exceeded the 25% chance baseline (FDR-adjusted $q < .001$). Neutral-item accuracy was 27.6% ($q = .610$), and sham-item accuracy was 32.7% ($q = .100$); neither was clearly above chance. Across all 12 items, mean accuracy was 34.8% ($q < .001$). Participants therefore detected some genuine symmetry features, but performance remained well below perfect identification.

Post-unmasking guesses were nearly balanced: 17 participants guessed correctly (43.6%), 19 guessed incorrectly (48.7%), and 3 were uncertain (7.7%). The standard Bang blinding index was close to zero ($BBI = -.051$, 95% bootstrap CI $[-.359, .256]$). Confidence weighting changed little: the exploratory confidence-weighted index was $-.016$ (95% bootstrap CI $[-.203, .168]$, $n = 35$ directional responses). Both intervals crossed zero, indicating no reliable tendency toward either correct or opposite identification. However, the standard BBI interval extended beyond the predefined $[-0.2, 0.2]$ blinding region. The results were therefore consistent with effective blinding but too uncertain to rule out meaningful identification in either direction.

To test whether pre-unmasking sensitivity predicted subsequent condition identification, we related the 0–4 symmetry score to final-guess correctness, classified as correct versus incorrect or uncertain. Pre-unmasking symmetry-detection accuracy was descriptively higher among participants who subsequently identified the correct condition (2.00 versus 1.59 of four items), but the association was small and inconclusive (point-biserial $r = .18$, exact two-sided permutation $p = .33$; Figure 9d). Thus, better masked detection did not reliably predict the final condition guess in this sample.




Fig. 9 Blinding and manipulation-awareness summary ($N = 39$). (a) Mean pre-unmasking accuracy for the four symmetry, four neutral, and four sham items and across all 12 items. The dashed line marks the 25% chance baseline; asterisks denote FDR-adjusted $q < .001$. (b) Post-unmasking guesses: 17 correct, 19 incorrect, and 3 uncertain. (c) Standard Bang blinding index (filled blue circle) and exploratory confidence-weighted directional index (open green diamond), with participant-bootstrap 95% confidence intervals. The shaded regions indicate incorrect, blinded, and correct identification and apply only to the standard Bang index. (d) Participant-level pre-unmasking accuracy across the three four-item categories, grouped by final-guess correctness. Thin lines and small markers represent participants; heavier lines and larger markers represent group means. Incorrect and uncertain guesses are combined.

3.3 Exploratory Checks

Exploratory mixed-model checks for order and PLV moderation are reported in the OSF repository. No FDR-corrected order main effects, condition $\times$ order, condition $\times$ PLV, or condition $\times$ order $\times$ PLV interactions emerged for any ASC, 3D-ASCr, or self-related outcome family. For PLV itself, there was a main effect of order, $\chi^2(1) = 8.28$, $p = .004$, but no condition $\times$ order interaction, $\chi^2(2) = 0.37$, $p = .833$, indicating that block order did not change the between-condition adherence pattern. A planned linear contrast across session position was significant, $t(38) = -3.16$, $p = .003$, 95% CI $[-0.100, -0.022]$, indicating that PLV decreased rather than increased across sessions and therefore aligning more with waning engagement or fatigue than with a practice-improves account.

4 Discussion

This study provided a first controlled empirical evaluation of Viscereality with naive participants. Relative to audio-guided box breathing in the dark-screen control, the two VR conditions improved breathing adherence, shifted spatial self-experience toward a more expanded frame of reference, and produced an altered-state profile in which perceptual and positive effects outweighed mild distressing effects. None of the 47 tested participants discontinued the experiment because of discomfort or inability to tolerate the VR exposure. However, the preregistered aesthetic manipulation that varied the symmetry of the particle dynamics did not reliably differentiate the two active VR conditions. Taken together, these findings indicate that the paradigm produced a structured experiential profile during a brief breathing session with naive participants, but stronger control comparisons and objective measures are needed to determine the specificity and depth of these effects.

4.1 Breathing adherence and self-related experience

Participants in both active VR conditions followed the instructed breathing pattern more closely than in the dark-screen control. The display responded to participants' ongoing breathing rather than to the audio guide, so it made respiration continuously visible without telling participants how well they were matching the instructed pacing. One plausible interpretation is that the breath-responsive environment helped adherence by keeping participants engaged and by giving them the sense that their breathing was producing the visual experience. Previous findings are consistent with this. Breath-coupled VR feedback has been linked to improved focus on breathing and greater diaphragmatic movement (Blum et al. 2020; Rockstroh et al. 2021), greater focused attention and less distraction than non-VR biofeedback (Rockstroh et al. 2019), and increased interoceptive sensibility (Goral et al. 2024a). A simpler explanation also remains possible. Guided breathing in the dark-screen control is presumably less engaging than the same task in an immersive environment, and the adherence metric itself was derived from how closely breathing matched the trajectory expressed by the visual display. The study would have benefited from a more stringent control, such as a visually matched scene that does not respond to breathing or one that replays

1427 a prerecorded animation, because the engaging visual environment on its own may
1428 have contributed to the effect. This caveat matters because controlled comparisons in
1429 the VR breathing literature have not consistently shown VR-based interventions to
1430 outperform matched non-VR approaches on clinical outcomes (Cortez-Vázquez et al.
1431 2024; Pancini et al. 2025). The present adherence finding should therefore be treated
1432 as preliminary until tested against a more stringent control.

1433 Of the three self-related measures, only the spatial frame-of-reference continuum
1434 showed a condition effect. We had reason to expect body-boundary salience to shift as
1435 well because prior work reports reduced perceived boundary salience during medita-
1436 tion (Dambrun 2016); in one study that administered both scales, mindfulness training
1437 reduced perceived body boundaries and expanded allocentric spatial framing in tan-
1438 dem, with the boundary change mediating the spatial shift (Hanley et al. 2020). That
1439 pattern did not emerge here, perhaps because the two constructs respond to partly
1440 different triggers. Body-boundary dissolution appears to shift most reliably during
1441 practices that direct sustained attention toward the body itself, such as body-scan
1442 meditation (Dambrun 2016), mindfulness training (Hanley et al. 2020), and floatation-
1443 REST (Hruby et al. 2024). Spatial frame of reference, by contrast, has been linked
1444 not only to body-directed contemplative practice (Hanley et al. 2020) but also to
1445 perceived expansion of the physical frame of reference beyond the body (Dambrun
1446 2023). Because Viscerality acts on the surrounding visual-spatial environment rather
1447 than directing attention toward the body, it may have engaged the spatial-frame
1448 construct through this environmental route without producing the body-focused con-
1449 ditions under which boundary dissolution typically occurs. Of the three scales, spatial
1450 frame of reference is also the one most closely aligned with the broader question
1451 motivating this research programme, namely whether breath-coupled changes in sur-
1452 rounding space can alter the felt extension of self into the environment. Previous work
1453 has shown that mapping respiration onto an externalized virtual avatar can modulate
1454 body ownership (Monti et al. 2020b) and self-location (Adler et al. 2014; Betka et al.
1455 2020) through afferent respiratory pathways (Betka et al. 2020).

1456 However, this result requires cautious interpretation. The concentric depiction of
1457 the SFoRC scale may have resembled the expanding and contracting particle sphere
1458 shown in VR, which raises the possibility that participants selected more spatially
1459 expansive depictions because the scale imagery matched what they had just seen
1460 rather than because their felt self-boundary had shifted. This demand-characteristics
1461 concern cannot be ruled out with pictographic self-report alone and represents a clear
1462 limitation of the present measurement approach. The finding should accordingly be
1463 treated as preliminary pending corroboration from measures that do not share visual
1464 features with the stimulus, such as peripersonal space tasks. If confirmed by such
1465 measures, the spatial-frame shift would support continued investigation of whether
1466 mapping respiration onto surrounding space can modulate the felt extension of self
1467 into the environment.

1468 Within the context of a 10-minute box-breathing exercise with ambient background
1469 music, *Viscerality* increased all three higher-order dimensions of the Altered States
1470 of Consciousness Rating Scale relative to the dark-screen control, which provided
1471 only a central fixation marker and minimal visual breathing-performance feedback.
1472

Perceptual Effects showed the largest increase, Positive Effects a moderate increase, and Distressing Effects a smaller increase. None differed reliably between the two VR mappings that manipulated visual symmetry, and participants showed only limited feature-level sensitivity to this manipulation without reliable condition-level identification.

The fine-grained subscale decomposition and item-content analysis reveal a more specific experiential profile (Online Resource 1, Section S3.4). The perceptual shift primarily concerned simple patterns, colors, lights, and flashes, together with impressions that sound changed visual shapes or colors. These elementary visual phenomena closely matched features supplied directly by the particle environment, so their increase partly reflects overlap between the intervention and the questionnaire criteria. Participants also attributed visual changes to sound even though the same background music was presented in every condition and did not control the display. One plausible interpretation is that the richer and more dynamic visual stream provided more perceptual material that could be bound to, or retrospectively attributed to, concurrent sound, rather than producing audio-visual synaesthesia in the stricter sense. A similar distinction has been raised in psychedelic research, where LSD increased spontaneous synaesthesia-like reports that lacked the consistency and inducer specificity associated with genuine synaesthesia (Terhune et al. 2016).

The Positive Effects increase was concentrated in items reflecting a loosening of separation between self, surroundings, and time, heightened significance and emotional salience of ordinary objects, sensations of floating or diminished bodily presence, and moments of awe.

The smaller Distressing Effects increase had the character of diffuse destabilization, encompassing reduced agency, difficulty organizing thought, isolation, unexplained apprehension, and an altered or threatening quality of the surroundings. Neither constituent distress-related subscale remained reliable after correction across the fine-grained analyses. We therefore interpret the composite increase as an accumulation of mild and compatible unpleasant experiences rather than as a pronounced or clearly localized adverse response.

Descriptively, the pooled-VR ASC profile was compared with five breathwork datasets: conscious connected breathing (Bahi et al. 2024b), the 90-minute breathwork arm of Airways to Alteration (Fincham et al. 2026), facilitated circular breathwork (Havenith et al. 2025b), high-ventilation breathwork with music in an MRI/laboratory setting (Kartar et al. 2025), and Holotropic Breathwork (Uthaug et al. 2021). The Viscerality profile fell within the range of several of these longer breathwork datasets and showed comparatively lower distress than facilitated circular breathwork. Differences in breathing technique, duration, sample, music, facilitation, setting, and control structure make this comparison contextual rather than evidence of equivalence. Figure 10 shows the full profile overlay, and Online Resource 1, Section S3 retains the fine-grained subscale commentary.

Cross-Study ASC Profile Comparison




Fig. 10 Descriptive cross-study comparison of the present dataset with five named breathwork datasets. The left panel shows 11D-ASC subscale means and the right panel shows 3D-ASCr composite means. The Viscerality profile pools the symmetric and asymmetric VR conditions. Lines show raw 0–100 profile means, not standardized effect sizes; differences in technique, duration, sample, music, facilitation, and study design preclude inferential comparison. Legend order is purely graphical and does not imply a numbered study sequence.

At subscale level, the profiles differed most in Experience of Unity, Spiritual Experience, Blissful State, Disembodiment, Insightfulness, and Changed Meaning of Percepts. Changed Meaning of Percepts increased in pooled VR relative to the dark-screen control in the present study, whereas Insightfulness did not change reliably. Because the reference studies differ on multiple procedural and contextual dimensions, these profile differences do not identify the mechanism responsible for any between-study contrast.

The blinding results distinguish masked feature-level sensitivity from post-unmasking condition identification. Before unmasking, symmetry-item accuracy exceeded chance, while neutral and sham accuracy did not. Participants therefore detected some genuine symmetry features, but only imperfectly. After unmasking, correct and incorrect guesses were nearly balanced, and both blinding indices were close to zero. Correct guessers had descriptively higher symmetry accuracy than incorrect or uncertain guessers, but the association was small and inconclusive ($r = .18$, exact

$p = .33$). Overall, the results were consistent with effective blinding, but the wide standard BBI interval prevented a conclusive determination.

Participants did not reliably identify the symmetric condition. Although this is consistent with blinding, it may also limit the test of the symmetry hypothesis, because explicit attention to and awareness of the manipulated feature may be necessary for it to influence emotional experience. Symmetric coherence appeared only during the breath-hold phases, approximately half of each breathing cycle, and remained secondary to the more salient breath-coupled expansion and contraction. The null difference therefore does not distinguish between an absence of an affective effect and insufficient focused exposure or awareness. We therefore plan a follow-up study that makes symmetry or particle coherence the focal stimulus dimension, manipulates it in isolation under controlled exposure, verifies that participants perceive the difference, and assesses its effects on emotional valence using physiological markers alongside direct ratings rather than relying only on retrospective altered-state ratings.

4.2 Limitations and future directions

Except for breathing-adherence PLV, the study’s outcomes relied on retrospective reports completed after each condition. The observed differences therefore establish how participants subsequently rated their experience, not whether the intervention produced corresponding changes in ongoing neural, autonomic, or multisensory processes.

Retrospective psychometric scales remain valuable because subjective content is not directly observable. The 11D-ASC belongs to a widely used questionnaire family conceived for standardized comparison of altered-state phenomenology across induction methods (Dittrich 1998; Hovmand et al. 2023). The Altered States Database demonstrates this comparative role by bringing standardized questionnaire data from pharmacological and non-pharmacological inductions into a common framework that supports direct comparisons across methods (Schmidt and Berkemeyer 2018; Prugger et al. 2022).

Standardization does not remove the limitations of retrospective psychometrics. Scores remain sensitive to recall, item wording, construct coverage, and assumptions built into the instrument. Reviews of altered-state measures identify substantial variation across scales and incomplete evidence for their psychometric quality and coverage, while critiques of mystical-experience instruments note their religious framing (Sanders and Zijlmans 2021; Hovmand et al. 2023; de Deus Pontual et al. 2023). The present 11D-ASC results should therefore be interpreted as standardized questionnaire-defined reports that support cross-study comparison, not as a complete or modality-neutral measure of altered consciousness.

Future studies should triangulate retrospective ratings with concurrent measures rather than treat any single physiological signal as a direct readout of subjective experience. Candidate approaches include EEG complexity measures associated with breathwork phenomenology (Lewis-Healey et al. 2024b), the perturbational complexity index developed to distinguish levels of consciousness across wakefulness, sleep, anaesthesia, and disorders of consciousness (Casali et al. 2013), heart-rate entropy

1611 examined as a low-cost correlate of psychedelic states (Rosas et al. 2023), and mul-
1612 tisensory peripersonal-space tasks that quantify changes in the represented boundary
1613 around the body (Serino 2019; Bufacchi and Iannetti 2018). These measures remain
1614 indirect and construct-specific, but convergence across subjective, neural, autonomic,
1615 and behavioural domains would provide stronger evidence than questionnaires alone.
1616 No-report paradigms offer a separate way to reduce report-related confounds
1617 in neural measurements. They infer conscious content from previously validated
1618 behavioural or physiological signatures, but they do not eliminate the need for
1619 first-person validation (Tsuchiya et al. 2015). Laukkonen and Slagter identify neu-
1620 rophenomenology and no-report paradigms as complementary future directions for
1621 studying meditation states in which ordinary report may be difficult (Laukkonen and
1622 Slagter 2021).

1623 Viscerality was designed to test whether breath-coupled changes in surround-
1624 ing space can alter the felt boundary between self and environment. The spatial
1625 frame-of-reference finding provides preliminary support for this possibility, but a
1626 peripersonal-space task would offer a more direct behavioural test. Peripersonal space
1627 represents the multisensory boundary surrounding the body and can be estimated from
1628 response changes as stimuli approach or recede (Serino 2019; Bufacchi and Iannetti
1629 2018; Karakoc et al. 2025).

1630 A visuotactile peripersonal-space task could be administered during the 4-second
1631 breath holds without reusing the intervention’s visual features. Comparing responses
1632 after inhalation and exhalation would test whether respiratory expansion and contrac-
1633 tion are accompanied by corresponding changes in the represented boundary around
1634 the body. A reversed-mapping condition could then test whether any shift depends on
1635 congruence between respiratory phase and spatial direction.

1636 The particle system in the present experiment was used to convey the expansion
1637 and contraction of surrounding space in response to breathing, but particles can also
1638 encode visceral signals directly onto the body or its surroundings, or construct abstract
1639 visual configurations that convey affective states through movement, colour, shape, or
1640 synchrony. Our manipulation of dynamic particle symmetry through oscillator cou-
1641 pling explored only one such parameter and did not yield observable effects. Our
1642 lesson from this is that mapping aesthetic features of particle displays onto affective
1643 dimensions requires a more fine-grained approach than a single global variable assessed
1644 through altered-state questionnaires. Short aesthetic judgment tasks offer a tractable
1645 starting point, following swarm robotics research where trajectory smoothness, speed,
1646 synchronization, and spatial coherence map onto separable affective dimensions (Dietz
1647 et al. 2017; Santos and Egerstedt 2021). The immersive 3D particle environment affords
1648 considerably more degrees of freedom than planar swarm displays, with particles mov-
1649 ing through three spatial dimensions under both local neighbour-coupling and global
1650 synchronization parameters, and testing specific feature combinations in controlled
1651 affective assessments would provide a more solid basis for these mappings than retro-
1652 spective questionnaires after prolonged sessions. Because the medium is abstract rather
1653 than representational, the same particle field could in principle simultaneously encode
1654 multiple biosignals and serve as the basis for biofeedback displays that are intuitive
1655 to the user and conducive to the affective states the feedback is meant to support.

1656

The present study found that the particle system was associated with higher Positive and Perceptual Effects during box breathing, alongside a smaller Distressing Effects increase. The cross-study profile provides descriptive context only and does not establish equivalence, superior tolerability, or clinical benefit. Because the system is self-administered and the session lasted 10 minutes, future studies could test whether it supports breathing practice in settings where access to a facilitator is limited or time is scarce. High-ventilation techniques are contraindicated for individuals vulnerable to panic, syncope, or hypocapnia (Fincham et al. 2023), and a slow-paced breath-coupled VR system offers a lower-intensity entry point into breathing as a therapeutic modality.

One possible population for such testing is stroke survivors, given that respiratory dysfunction and sleep-disordered breathing are common after stroke and associated with poorer recovery (Patrizz et al. 2023; Seiler et al. 2019), while structured respiratory rehabilitation remains underemphasized in clinical practice (Menezes et al. 2016). Slow-paced breathing exercises are already used to retrain respiratory function after stroke (Abdullahi et al. 2024; Yoon et al. 2022), and coupling breathing to an intuitive visual mapping of respiratory depth could engage an additional learning pathway that supports recovery of motor control over respiration. These remain illustrative possibilities. The present sample consisted of a homogeneous group of university psychology students, and testing usability, tolerability, and efficacy in clinical populations is a necessary next step.

5 Conclusion

During a brief self-administered box-breathing session, Viscerality was associated with closer synchronization to the instructed rhythm, a more spatially extended self-rating, and higher Positive, Perceptual, and smaller Distressing Effects than the dark-screen control. No measured outcome differed reliably between the symmetric and asymmetric VR mappings. These results come from a recruitment-naïve university sample and do not establish equivalence to longer facilitator-supported breathwork or clinical efficacy.

Supplementary Information

Online Resource 1: Supplementary methods and analyses, including deviations from preregistration, phase-locking value computation and quality screening, complete 11D-ASC results, exploratory pictograph-ASC correspondence, temporal experience tracer analyses, and reproducibility information.

Statements and Declarations

Funding: The study reported here was funded by the BIAL Foundation through Grant 308/22, “Altered States of Viscerality: Augmenting Breathwork with Bio-Responsive Virtual Reality,” awarded under the 2022 Grants Programme for Scientific Research. Continued development of Viscerality received support from the Baden-Württemberg

1703 Stiftung through the Forschungswerkstatt – Deine Idee für Morgen programme for
1704 the project “Atemräume: Integration atem- und herzratenbasierter Interaktionen in
1705 therapeutische VR-Umgebungen.”

1706 Competing interests: The authors declare no competing interests.

1707 Ethics approval: The Ethics Committee of the University of Konstanz approved
1708 the study, which was conducted in accordance with the Declaration of Helsinki.

1709 Consent to participate: All participants provided informed consent before partici-
1710 pation.

1711 Consent for publication: Participants consented to the publication of anonymized,
1712 non-identifiable data.

1713 Availability of data and materials: The deidentified participant-level data, origi-
1714 nal preregistration record, statistical analysis outputs, figure-generation scripts, study
1715 materials, and reproducibility documentation are available as a reproducibility package
1716 at OSF: <https://doi.org/10.17605/OSF.IO/64MWK>.

1717 Code availability: The statistical analysis and figure-generation scripts, exact envi-
1718 ronment specifications, and replay instructions are available at OSF: <https://doi.org/10.17605/OSF.IO/64MWK>.

1720 Authors’ contributions: George Fejer: Conceptualization, Data curation, Formal
1721 analysis, Funding acquisition, Investigation, Methodology, Project administration,
1722 Software, Validation, Visualization, Writing – original draft, Writing – review &
1723 editing. Till Holzapfel: Conceptualization, Methodology, Resources, Software, Valida-
1724 tion, Visualization. Taru Hirvonen: Methodology, Resources, Software, Visualization.
1725 Anestis Lalidis Mateo: Methodology, Resources, Software. Johannes Blum: Method-
1726 ology, Resources, Software, Validation. Michael Gaebler: Conceptualization, Project
1727 administration, Supervision, Writing – review & editing. Bigna Lenggenhager: Con-
1728 ceptualization, Funding acquisition, Supervision, Writing – review & editing.

1729

1730 References

1731

1732 Abdullahi A, Wong TWL, Ng SSM (2024) Efficacy of diaphragmatic breathing exercise
1733 on respiratory, cognitive, and motor function outcomes in patients with stroke: a sys-
1734 tematic review and meta-analysis. *Frontiers in Neurology* 14:1233408. <https://doi.org/10.3389/fneur.2023.1233408>, URL <https://doi.org/10.3389/fneur.2023.1233408>
1735
1736

1737 Abtahi P, Hough SQ, Landay JA, et al (2022) Beyond being real: A sensorimotor
1738 control perspective on interactions in virtual reality. In: *Proceedings of the 2022*
1739 *CHI Conference on Human Factors in Computing Systems*. ACM, pp 1–17, <https://doi.org/10.1145/3491102.3517706>, URL <https://doi.org/10.1145/3491102.3517706>
1740

1741 Acebrón JA, Bonilla LL, Pérez Vicente CJ, et al (2005) The kuramoto model: A simple
1742 paradigm for synchronization phenomena. *Reviews of modern physics* 77(1):137–
1743 185. <https://doi.org/10.1103/RevModPhys.77.137>
1744

1745 Adler D, Herbelin B, Similowski T, et al (2014) Breathing and sense of self:
1746 Visuo-respiratory conflicts alter body self-consciousness. *Respiratory Physiology*
1747 & *Neurobiology* 203:68–74. <https://doi.org/10.1016/j.resp.2014.08.003>, URL <https://doi.org/10.1016/j.resp.2014.08.003>
1748

//doi.org/10.1016/j.resp.2014.08.003	1749
Ahmed A, Devi RG, Priya AJ (2021) Effect of box breathing technique on lung function test. <i>Journal of Pharmaceutical Research International</i> 33(58A):25–31. https://doi.org/10.9734/JPRI/2021/v33i58A34085 , URL https://doi.org/10.9734/JPRI/2021/v33i58A34085	1750 1751 1752 1753 1754 1755
Alhammad SA (2024) Advocating for action: Exploring the potential of virtual reality in breathing exercise – a review of the clinical applications. <i>Patient Preference and Adherence</i> 18:695–707. https://doi.org/10.2147/ppa.s451609 , URL https://doi.org/10.2147/ppa.s451609	1756 1757 1758 1759 1760
Antle AN, Corness G, Droumeva M (2009) What the body knows: Exploring the benefits of embodied metaphors in hybrid physical digital environments. <i>Interacting with Computers</i> 21(1–2):66–75. https://doi.org/10.1016/j.intcom.2008.10.005 , URL https://doi.org/10.1016/j.intcom.2008.10.005	1761 1762 1763 1764 1765
Aronoff J (2006) How we recognize angry and happy emotion in people, places, and things. <i>Cross-Cultural Research</i> 40(1):83–105. https://doi.org/10.1177/1069397105282597 , URL https://doi.org/10.1177/1069397105282597	1766 1767 1768 1769
Ashhad S, Kam K, Del Negro CA, et al (2022) Breathing rhythm and pattern and their influence on emotion. <i>Annual Review of Neuroscience</i> 45:223–247. https://doi.org/10.1146/annurev-neuro-090121-014424	1770 1771 1772
Aspell JE, Heydrich L, Marillier G, et al (2013) Turning body and self inside out. <i>Psychological Science</i> 24(12):2445–2453. https://doi.org/10.1177/0956797613498395 , URL https://doi.org/10.1177/0956797613498395	1773 1774 1775 1776
Azzalini D, Rebollo I, Tallon-Baudry C (2019) Visceral signals shape brain dynamics and cognition. <i>Trends in Cognitive Sciences</i> 23(6):488–509. https://doi.org/10.1016/j.tics.2019.03.007	1777 1778 1779 1780
Bahi C, Irrmischer M, Franken K, et al (2024a) Effects of conscious connected breathing on cortical brain activity, mood and state of consciousness in healthy adults. <i>Current Psychology</i> 43(12):10578–10589. https://doi.org/10.1007/s12144-023-05119-6	1781 1782 1783 1784
Bahi C, Irrmischer M, Franken K, et al (2024b) Effects of conscious connected breathing on cortical brain activity, mood and state of consciousness in healthy adults. <i>Current Psychology</i> 43(12):10578–10589. https://doi.org/10.1007/s12144-023-05119-6	1785 1786 1787 1788
Balban MY, Neri E, Kogon MM, et al (2023) Brief structured respiration practices enhance mood and reduce physiological arousal. <i>Cell Reports Medicine</i> 4(1):100895. https://doi.org/10.1016/j.xcrm.2022.100895	1789 1790 1791
Bang H, Ni L, Davis CE (2004) Assessment of blinding in clinical trials. <i>Controlled Clinical Trials</i> 25(2):143–156. https://doi.org/10.1016/j.cct.2003.10.016	1792 1793 1794

- 1795 Banushi B, Brendle M, Ragnhildstveit A, et al (2023) Breathwork interventions for
1796 adults with clinically diagnosed anxiety disorders: A scoping review. *Brain Sciences*
1797 13(2):256. <https://doi.org/10.3390/brainsci13020256>
1798
- 1799 Barton AC, Do M, Sheen J, et al (2024) The restorative and state enhancing potential
1800 of abstract fractal-like imagery and interactive mindfulness interventions in virtual
1801 reality. *Virtual Reality* 28(1):53. <https://doi.org/10.1007/s10055-023-00916-7>
1802
- 1803 Bartram L, Nakatani A (2009) Distinctive parameters of expressive motion.
1804 In: *Computational Aesthetics in Graphics, Visualization, and Imag-*
1805 *ing*. The Eurographics Association, pp 129–136, [https://doi.org/10.2312/](https://doi.org/10.2312/COMPAESTH/COMPAESTH09/129-136)
1806 [COMPAESTH/COMPAESTH09/129-136](https://doi.org/10.2312/COMPAESTH/COMPAESTH09/129-136), URL [http://diglib.eg.org/handle/10.](http://diglib.eg.org/handle/10.2312/COMPAESTH.COMPAESTH09.129-136)
1807 [2312/COMPAESTH.COMPAESTH09.129-136](http://diglib.eg.org/handle/10.2312/COMPAESTH.COMPAESTH09.129-136)
1808
- 1809 Belli F, Felisatti A, Fischer MH (2021) BreaThink: breathing affects produc-
1810 tion and perception of quantities. *Experimental Brain Research* 239(8):2489–
1811 2499. <https://doi.org/10.1007/s00221-021-06147-z>, URL [https://doi.org/10.1007/](https://doi.org/10.1007/s00221-021-06147-z)
1812 [s00221-021-06147-z](https://doi.org/10.1007/s00221-021-06147-z)
- 1813 Bentley TGK, D’Andrea-Penna G, Rakic M, et al (2023) Breathing practices for stress
1814 and anxiety reduction: Conceptual framework of implementation guidelines based
1815 on a systematic review of the published literature. *Brain Sciences* 13(12):1612. <https://doi.org/10.3390/brainsci13121612>
1816 <https://doi.org/10.3390/brainsci13121612>
1817
- 1818 Bertamini M, Makin ADJ, Rampone G (2013) Implicit association of symmetry with
1819 positive valence, high arousal and simplicity. *i-Perception* 4(5):317–327. [https://doi.](https://doi.org/10.1068/i0601jw)
1820 [org/10.1068/i0601jw](https://doi.org/10.1068/i0601jw)
1821
- 1822 Betka S, Canzoneri E, Adler D, et al (2020) Mechanisms of the breathing con-
1823 tribution to bodily self-consciousness in healthy humans: Lessons from machine-
1824 assisted breathing? *Psychophysiology* 57(8):e13564. [https://doi.org/10.1111/psyp.](https://doi.org/10.1111/psyp.13564)
1825 [13564](https://doi.org/10.1111/psyp.13564), URL <https://doi.org/10.1111/psyp.13564>
1826
- 1827 Billock VA, Tsou BH (2012) Elementary visual hallucinations and their relation-
1828 ships to neural pattern-forming mechanisms. *Psychological Bulletin* 138(4):744–774.
1829 <https://doi.org/10.1037/a0027580>, URL <https://doi.org/10.1037/a0027580>
1830
- 1831 Blazhenkova O, Kumar MM (2018) Angular versus curved shapes: Correspon-
1832 dences and emotional processing. *Perception* 47(1):67–89. [https://doi.org/10.1177/](https://doi.org/10.1177/0301006617731048)
1833 [0301006617731048](https://doi.org/10.1177/0301006617731048), URL <https://doi.org/10.1177/0301006617731048>
1834
- 1835 Blum J, Rockstroh C, Göritz AS (2019) Heart rate variability biofeedback based on
1836 slow-paced breathing with immersive virtual reality nature scenery. *Frontiers in*
1837 *psychology* 10:2172. <https://doi.org/10.3389/fpsyg.2019.02172>
1838
- 1839 Blum J, Rockstroh C, Göritz AS (2020) Development and pilot test of a virtual real-
1840 ity respiratory biofeedback approach. *Applied Psychophysiology and Biofeedback*

45(3):153–163. https://doi.org/10.1007/s10484-020-09468-x	1841
Boiten FA, Frijda NH, Wientjes CJE (1994) Emotions and respiratory patterns: review and critical analysis. <i>International Journal of Psychophysiology</i> 17(2):103–128. https://doi.org/10.1016/0167-8760(94)90027-2	1842
Bossenbroek R, Wols A, Weerdmeester J, et al (2020) Efficacy of a virtual reality biofeedback game (DEEP) to reduce anxiety and disruptive classroom behavior: single-case study. <i>JMIR Mental Health</i> 7(3):e16066. https://doi.org/10.2196/16066 , URL https://doi.org/10.2196/16066	1843
Bufacchi RJ, Iannetti GD (2018) An action field theory of peripersonal space. <i>Trends in Cognitive Sciences</i> 22(12):1076–1090. https://doi.org/10.1016/j.tics.2018.09.004	1844
Burger B, Toiviainen P (2020) See how it feels to move: Relationships between movement characteristics and perception of emotions in dance. <i>Human Technology</i> 16(3):233–256. https://doi.org/10.17011/ht/urn.202011256764 , URL https://doi.org/10.17011/ht/urn.202011256764	1845
Casali AG, Gosseries O, Rosanova M, et al (2013) A theoretically based index of consciousness independent of sensory processing and behavior. <i>Science Translational Medicine</i> 5(198):198ra105. https://doi.org/10.1126/scitranslmed.3006294	1846
Chan PYS, Lee LY, Davenport PW (2024) Neural mechanisms of respiratory interoception. <i>Autonomic Neuroscience: Basic and Clinical</i> 253:103181. https://doi.org/10.1016/j.autneu.2024.103181	1847
Chang E, Kim HT, Yoo B (2020) Virtual reality sickness: a review of causes and measurements. <i>International Journal of Human–Computer Interaction</i> 36(17):1658–1682. https://doi.org/10.1080/10447318.2020.1778351	1848
Cheiran JFP, Bandeira DR, Pimenta MS (2025) Measuring the key components of the user experience in immersive virtual reality environments. <i>Frontiers in Virtual Reality</i> 6:1585614. https://doi.org/10.3389/frvir.2025.1585614 , URL https://doi.org/10.3389/frvir.2025.1585614	1849
Cortez-Vázquez G, Adriaanse M, Burchell GL, et al (2024) Virtual reality breathing interventions for mental health: A systematic review and meta-analysis of randomized controlled trials. <i>Applied Psychophysiology and Biofeedback</i> 49(1):1–21. https://doi.org/10.1007/s10484-023-09611-4 , URL https://doi.org/10.1007/s10484-023-09611-4	1850
Dambrun M (2016) When the dissolution of perceived body boundaries elicits happiness: The effect of selflessness induced by a body scan meditation. <i>Consciousness and Cognition</i> 46:89–98. https://doi.org/10.1016/j.concog.2016.09.013	1851

1887 Dambrun M (2023) Beyond the body/self-consciousness matrix: Scale of the physical
 1888 frame of reference and self-transcendence experience. *Psychology of Consciousness:*
 1889 *Theory, Research, and Practice* 10(3):292–312. <https://doi.org/10.1037/cns0000314>,
 1890 URL <https://doi.org/10.1037/cns0000314>
 1891
 1892 Damiano C, Gayen P, Rezanejad M, et al (2023) Anger is red, sadness is blue: Emotion
 1893 depictions in abstract visual art by artists and non-artists. *Journal of Vision* 23(4):1.
 1894 <https://doi.org/10.1167/jov.23.4.1>, URL <https://doi.org/10.1167/jov.23.4.1>
 1895
 1896 de Deus Pontual AA, Senhorini HG, Corradi-Webster CM, et al (2023) Systematic
 1897 review of psychometric instruments used in research with psychedelics. *Journal of*
 1898 *Psychoactive Drugs* 55(3):359–368. <https://doi.org/10.1080/02791072.2022.2079108>
 1899
 1900 Dietz G, Jane L. E., Washington P, et al (2017) Human perception of swarm robot
 1901 motion. In: *Proceedings of the 2017 CHI Conference Extended Abstracts on Human*
 1902 *Factors in Computing Systems*. ACM, pp 2520–2527, <https://doi.org/10.1145/3027063.3053220>, URL <https://doi.org/10.1145/3027063.3053220>
 1903
 1904 Dittrich A (1998) The standardized psychometric assessment of altered states of con-
 1905 sciousness (ascs) in humans. *Pharmacopsychiatry* 31(S2):80–84. [https://doi.org/10.](https://doi.org/10.1055/s-2007-979351)
 1906 [1055/s-2007-979351](https://doi.org/10.1055/s-2007-979351)
 1907
 1908 van Doorn JB, van den Bergh D, Bohm U, et al (2021) The JASP guidelines for
 1909 conducting and reporting a bayesian analysis. *Psychonomic Bulletin & Review*
 1910 28(3):813–826. <https://doi.org/10.3758/s13423-020-01798-5>
 1911
 1912 Ekman P (1993) Facial expression and emotion. *American psychologist* 48(4):384–392.
 1913 <https://doi.org/10.1037/0003-066X.48.4.384>
 1914
 1915 Engelen T, Solcà M, Tallon-Baudry C (2023) Interoceptive rhythms in
 1916 the brain. *Nature Neuroscience* 26(10):1670–1684. [https://doi.org/10.1038/](https://doi.org/10.1038/s41593-023-01425-1)
 1917 [s41593-023-01425-1](https://doi.org/10.1038/s41593-023-01425-1)
 1918
 1919 Fejer G (2026) Breathing space: Spatial mapping of breath and cardiac biofeedback
 1920 for affective state representation and coherence training in viscerality. In: *Proceed-*
 1921 *ings of the 1st International Conference on Human-Computer Interaction in the*
 1922 *Alps (AlpCHI 2026)*, <https://doi.org/10.1145/3780045.3780061>, added to support
 1923 manuscript introduction citation; verify publisher metadata before submission
 1924
 1925 Fejer G, Holzapfel T, Gómez-Emilsson A, et al (2025) Viscerality: A bio-responsive
 1926 vr system for breath-based interactions and coupled oscillator dynamics to augment
 1927 altered states of consciousness. In: *Mensch und Computer 2025 Workshopband,*
 1928 *Gesellschaft für Informatik e.V.*, <https://doi.org/10.18420/muc2025-mci-ws11-174>,
 1929 URL <https://dl.gi.de/handle/20.500.12116/46829>
 1930
 1931 Fincham GW, Strauss C, Montero-Marin J, et al (2023) Effect of breathwork on
 1932 stress and mental health: A meta-analysis of randomised-controlled trials. *Scientific*

Reports 13(1):432. https://doi.org/10.1038/s41598-022-27247-y	1933
	1934
Fincham GW, Epel E, Colasanti A, et al (2024) Effects of brief remote high ventilation breathwork with retention on mental health and wellbeing: a randomised placebo-controlled trial. <i>Scientific Reports</i> 14(1):16893. https://doi.org/10.1038/s41598-024-64254-7	1935
	1936
	1937
	1938
	1939
Fincham GW, Caddy E, Kartar AA, et al (2026) An exploratory study of breathwork-induced altered states of consciousness in experienced practitioners: the airways to alteration (A2A) trial. <i>Frontiers in Psychology</i> 17:1851882. https://doi.org/10.3389/fpsyg.2026.1851882	1940
	1941
	1942
	1943
	1944
Gerritsen RJS, Band GPH (2018) Breath of life: The respiratory vagal stimulation model of contemplative activity. <i>Frontiers in Human Neuroscience</i> 12:397. https://doi.org/10.3389/fnhum.2018.00397	1945
	1946
	1947
	1948
Glowacki DR (2024) VR models of death and psychedelics: An aesthetic paradigm for design beyond day-to-day phenomenology. <i>Frontiers in Virtual Reality</i> 4:1286950. https://doi.org/10.3389/frvir.2023.1286950 , URL https://doi.org/10.3389/frvir.2023.1286950	1949
	1950
	1951
	1952
Glowacki DR, Wonnacott MD, Freire R, et al (2020) Isness: Using multi-person VR to design peak mystical type experiences comparable to psychedelics. In: <i>Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems</i> . ACM, pp 1–14, https://doi.org/10.1145/3313831.3376649 , URL https://doi.org/10.1145/3313831.3376649	1953
	1954
	1955
	1956
	1957
	1958
Glowacki DR, Williams RR, Wonnacott MD, et al (2022) Group VR experiences can produce ego attenuation and connectedness comparable to psychedelics. <i>Scientific Reports</i> 12(1):8995. https://doi.org/10.1038/s41598-022-12637-z , URL https://doi.org/10.1038/s41598-022-12637-z	1959
	1960
	1961
	1962
	1963
Goral O, Wald IY, Maimon A, et al (2024a) Enhancing interoceptive sensibility through exteroceptive–interoceptive sensory substitution. <i>Scientific Reports</i> 14(1):14855. https://doi.org/10.1038/s41598-024-63231-4 , URL https://doi.org/10.1038/s41598-024-63231-4	1964
	1965
	1966
	1967
	1968
Goral O, Wald IY, Maimon A, et al (2024b) Enhancing interoceptive sensibility through exteroceptive–interoceptive sensory substitution. <i>Scientific Reports</i> 14(1):14855. https://doi.org/10.1038/s41598-024-63231-4 , URL https://doi.org/10.1038/s41598-024-63231-4	1969
	1970
	1971
	1972
	1973
Guy M, Normand JM, Jeunet-Kelway C, et al (2023) The sense of embodiment in virtual reality and its assessment methods. <i>Frontiers in Virtual Reality</i> 4:1141683. https://doi.org/10.3389/frvir.2023.1141683 , URL https://doi.org/10.3389/frvir.2023.1141683	1974
	1975
	1976
	1977
	1978

- 1979 Hanley AW, Garland EL (2019) Spatial frame of reference as a phenomenological
1980 feature of self-transcendence: Measurement and manipulation through mindfulness
1981 meditation. *Psychology of Consciousness: Theory, Research, and Practice* 6(4):329–
1982 345. <https://doi.org/10.1037/cns0000204>
1983
- 1984 Hanley AW, Dambrun M, Garland EL (2020) Effects of mindfulness meditation on
1985 self-transcendent states: Perceived body boundaries and spatial frames of reference.
1986 *Mindfulness* 11(5):1194–1203. <https://doi.org/10.1007/s12671-020-01330-9>
1987
- 1988 Havenith MN, Leidenberger M, Brasanac J, et al (2025a) Decreased co2 saturation
1989 during circular breathwork supports emergence of altered states of consciousness.
1990 *Communications Psychology* 3(1):59. <https://doi.org/10.1038/s44271-025-00247-0>
1991
- 1992 Havenith MN, Leidenberger M, Brasanac J, et al (2025b) Decreased co2 saturation
1993 during circular breathwork supports emergence of altered states of consciousness.
1994 *Communications Psychology* 3(1). <https://doi.org/10.1038/s44271-025-00247-0>,
1995 URL <https://doi.org/10.1038/s44271-025-00247-0>
- 1996 Homma I, Masaoka Y (2008) Breathing rhythms and emotions. *Experimental Physi-*
1997 *ology* 93(9):1011–1021. <https://doi.org/10.1113/expphysiol.2008.042424>
1998
- 1999 Hovmand OR, Poulsen ED, Arnfred S (2023) Assessment of the acute subjective
2000 psychedelic experience: A review of patient-reported outcome measures in clinical
2001 research on classical psychedelics. *Journal of Psychopharmacology* 38(1):19–32.
2002 <https://doi.org/10.1177/02698811231200019>
2003
- 2004 Hovmand OR, Madsen M, Fisher PM, et al (2024) Altered states of consciousness
2005 in danish healthy volunteers and recreational users of psilocybin and the possible
2006 impact of setting and intention: Danish validation of the five-dimensional altered
2007 states of consciousness questionnaire. *Journal of Psychopharmacology* 38(9):924–
2008 932. <https://doi.org/10.1177/02698811241269669>
2009
- 2010 Hruby H, Schmidt S, Feinstein JS, et al (2024) Induction of altered states of conscious-
2011 ness during Floatation-REST is associated with the dissolution of body boundaries
2012 and the distortion of subjective time. *Scientific Reports* 14(1):9316. <https://doi.org/10.1038/s41598-024-59642-y>, URL <https://doi.org/10.1038/s41598-024-59642-y>
2013
- 2014 Jachs B (2021) The neurophenomenology of meditative states: Introducing temporal
2015 experience tracing to capture subjective experience states and their neural corre-
2016 lates. PhD thesis, University of Cambridge, <https://doi.org/10.17863/CAM.80153>,
2017 URL <https://www.repository.cam.ac.uk/handle/1810/332709>
2018
- 2019 Jerath R, Beveridge C (2020) Respiratory rhythm, autonomic modulation, and the
2020 spectrum of emotions: The future of emotion recognition and modulation. *Frontiers*
2021 *in Psychology* 11:1980. <https://doi.org/10.3389/fpsyg.2020.01980>
2022
2023
2024

- Johnson ME (2023) Qualia formalism and a symmetry theory of valence. OpenTheory, URL https://opentheory.net/Qualia_Formalism_and_a_Symmetry_Theory_of_Valence.pdf
- Kaduk J, Cavdan M, Drewing K, et al (2024) From one to many: How active robot swarm sizes influence human cognitive processes. In: 2024 33rd IEEE International Conference on Robot and Human Interactive Communication (RO-MAN). IEEE, pp 1207–1212, <https://doi.org/10.1109/RO-MAN60168.2024.10731232>, URL <https://doi.org/10.1109/RO-MAN60168.2024.10731232>
- Karakoc C, Lucifora C, Massimino S, et al (2025) Extending peri-personal space in immersive virtual reality: A systematic review. Virtual Worlds 4(1):5. <https://doi.org/10.3390/virtualworlds4010005>, URL <https://doi.org/10.3390/virtualworlds4010005>
- Kartar AA, Horinouchi T, Örsik B, et al (2025) Neurobiological substrates of altered states of consciousness induced by high ventilation breathwork accompanied by music. PLOS One 20(8):e0329411. <https://doi.org/10.1371/journal.pone.0329411>, URL <https://doi.org/10.1371/journal.pone.0329411>
- Kaufeld M, Bourdeinik J, Prinz LM, et al (2022) Emotions are associated with the genesis of visually induced motion sickness in virtual reality. Experimental Brain Research 240(10):2757–2771. <https://doi.org/10.1007/s00221-022-06454-z>
- Khatin-Zadeh O, Eskandari Z, Vahdat S, et al (2019) Why are motions effective in describing emotions? Polish Psychological Bulletin 50(2):119–124. <https://doi.org/10.24425/ppb.2019.126027>
- Klüver H (1966) Mescal, and Mechanisms of Hallucinations. University of Chicago Press, Chicago, reprint edition; originally published 1928
- Kodithuwakku Arachchige SNK, Chander H, Freeman H, et al (2022) The walls are closing in: postural responses to a virtual reality claustrophobic simulation. Clinical and Translational Neuroscience 6(2):15. <https://doi.org/10.3390/ctn6020015>
- Kolahi J, Bang H, Park J (2009) Towards a proposal for assessment of blinding success in clinical trials: Up-to-date review. Community Dentistry and Oral Epidemiology 37(6):477–484. <https://doi.org/10.1111/j.1600-0528.2009.00494.x>
- Kreibig SD (2010) Autonomic nervous system activity in emotion: A review. Biological Psychology 84(3):394–421. <https://doi.org/10.1016/j.biopsycho.2010.03.010>
- Kruschke JK (2021) Bayesian analysis reporting guidelines. Nature Human Behaviour 5(10):1282–1291. <https://doi.org/10.1038/s41562-021-01177-7>
- Kytikova O, Novgorodtseva T, Denisenko Y, et al (2021) Thermosensory transient receptor potential ion channels and asthma. Biomedicines 9(7):816. <https://doi.org/>

2071 [10.3390/biomedicines9070816](https://doi.org/10.3390/biomedicines9070816)
 2072
 2073 Laukkonen RE, Slagter HA (2021) From many to (n)one: Meditation and the plasticity
 2074 of the predictive mind. *Neuroscience & Biobehavioral Reviews* 128:199–217. <https://doi.org/10.1016/j.neubiorev.2021.06.021>
 2075
 2076 Lawrence DW, Carhart-Harris R, Griffiths R, et al (2022) Phenomenology and content
 2077 of the inhaled N, N-dimethyltryptamine (N, N-DMT) experience. *Scientific Reports*
 2078 12(1):8562. <https://doi.org/10.1038/s41598-022-11999-8>, URL [https://doi.org/10.](https://doi.org/10.1038/s41598-022-11999-8)
 2079 [1038/s41598-022-11999-8](https://doi.org/10.1038/s41598-022-11999-8)
 2080
 2081 Lenggenhager B, Tadi T, Metzinger T, et al (2007) Video ergo sum: Manipulating
 2082 bodily self-consciousness. *Science* 317(5841):1096–1099. [https://doi.org/10.1126/](https://doi.org/10.1126/science.1143439)
 2083 [science.1143439](https://doi.org/10.1126/science.1143439), URL <https://doi.org/10.1126/science.1143439>
 2084
 2085 Lewis-Healey E, Tagliazucchi E, Canales-Johnson A, et al (2024a) Breathwork-induced
 2086 psychedelic experiences modulate neural dynamics. *Cerebral Cortex* 34(8):bhae347.
 2087 <https://doi.org/10.1093/cercor/bhae347>
 2088
 2089 Lewis-Healey E, Tagliazucchi E, Canales-Johnson A, et al (2024b) Breathwork-induced
 2090 psychedelic experiences modulate neural dynamics. *Cerebral Cortex* 34(8):bhae347.
 2091 <https://doi.org/10.1093/cercor/bhae347>
 2092
 2093 Makin ADJ, Wilton MM, Pecchinenda A, et al (2012) Symmetry perception and
 2094 affective responses: A combined eeg/emg study. *Neuropsychologia* 50(14):3250–3261.
 2095 <https://doi.org/10.1016/j.neuropsychologia.2012.10.003>
 2096
 2097 Maravita A, Iriki A (2004) Tools for the body (schema). *Trends in Cognitive Sciences*
 2098 8(2):79–86. <https://doi.org/10.1016/j.tics.2003.12.008>
 2099
 2100 Marcolin F, Wally Scurati G, Ulrich L, et al (2021) Affective virtual reality: How to
 2101 design artificial experiences impacting human emotions. *IEEE Computer Graphics*
 2102 *and Applications* 41(6):171–178. <https://doi.org/10.1109/MCG.2021.3115015>, URL
 2103 <https://doi.org/10.1109/MCG.2021.3115015>
 2104
 2105 McGraw KO, Wong SP (1992) A common language effect size statistic. *Psychological*
 2106 *Bulletin* 111(2):361–365. <https://doi.org/10.1037/0033-2909.111.2.361>, URL <https://doi.org/10.1037/0033-2909.111.2.361>
 2107
 2108 Menezes KKP, Nascimento LR, Ada L, et al (2016) Respiratory muscle training
 2109 increases respiratory muscle strength and reduces respiratory complications after
 2110 stroke: a systematic review. *Journal of Physiotherapy* 62(3):138–144. [https://doi.](https://doi.org/10.1016/j.jphys.2016.05.014)
 2111 [org/10.1016/j.jphys.2016.05.014](https://doi.org/10.1016/j.jphys.2016.05.014), URL <https://doi.org/10.1016/j.jphys.2016.05.014>
 2112
 2113 Millière R (2020) The varieties of selflessness. *Philosophy and the Mind Sci-*
 2114 *ences* 1(I):1–41. <https://doi.org/10.33735/phimisci.2020.i.48>, URL [https://doi.org/](https://doi.org/10.33735/phimisci.2020.i.48)
 2115 [10.33735/phimisci.2020.i.48](https://doi.org/10.33735/phimisci.2020.i.48)
 2116

Mogan R, Fischer R, Bulbulia JA (2017) To be in synchrony or not? a meta-analysis of synchrony's effects on behavior, perception, cognition and affect. *Journal of Experimental Social Psychology* 72:13–20. <https://doi.org/10.1016/j.jesp.2017.03.009>

Monti A, Porciello G, Tieri G, et al (2020a) The “embreathment” illusion highlights the role of breathing in corporeal awareness. *Journal of Neurophysiology* 123(1):420–427. <https://doi.org/10.1152/jn.00617.2019>, URL <https://doi.org/10.1152/jn.00617.2019>

Monti A, Porciello G, Tieri G, et al (2020b) The “embreathment” illusion highlights the role of breathing in corporeal awareness. *Journal of Neurophysiology* 123(1):420–427. <https://doi.org/10.1152/jn.00617.2019>, URL <https://doi.org/10.1152/jn.00617.2019>

Morey RD, Rouder JN (2024) BayesFactor: Computation of Bayes Factors for Common Designs. <https://doi.org/10.32614/CRAN.package.BayesFactor>, URL <https://CRAN.R-project.org/package=BayesFactor>, r package version 0.9.12-4.7

Mori K, Sakano H (2022a) Neural circuitry for stress information of environmental and internal odor worlds. *Frontiers in Behavioral Neuroscience* 16:943647. <https://doi.org/10.3389/fnbeh.2022.943647>

Mori K, Sakano H (2022b) Processing of odor information during the respiratory cycle in mice. *Frontiers in Neural Circuits* 16:861800. <https://doi.org/10.3389/fncir.2022.861800>

Muth C, Carbon CC (2013) The aesthetic aha: On the pleasure of having insights into gestalt. *Acta psychologica* 144(1):25–30. <https://doi.org/10.1016/j.actpsy.2013.05.001>

Pancini E, Di Natale AF, Villani D (2025) Breathing in virtual reality for promoting mental health: a scoping review. *Virtual Reality* 29(1):29. <https://doi.org/10.1007/s10055-024-01096-8>, URL <https://doi.org/10.1007/s10055-024-01096-8>

Patrizz A, El Hamamy A, Maniskas M, et al (2023) Stroke-induced respiratory dysfunction is associated with cognitive decline. *Stroke* 54(7):1863–1874. <https://doi.org/10.1161/STROKEAHA.122.041239>, URL <https://doi.org/10.1161/STROKEAHA.122.041239>

Pecchinenda A, Bertamini M, Makin AD, et al (2014) The pleasantness of visual symmetry: always, never or sometimes. *PLoS ONE* 9(3):e92685. <https://doi.org/10.1371/journal.pone.0092685>

Petkova VI, Ehrsson HH (2008) If I were you: Perceptual illusion of body swapping. *PLoS ONE* 3(12):e3832. <https://doi.org/10.1371/journal.pone.0003832>, URL <https://doi.org/10.1371/journal.pone.0003832>

2163 Petrizzo I, Mikellidou K, Avraam S, et al (2024) Reshaping the periper-
 2164 sonal space in virtual reality. *Scientific Reports* 14(1). <https://doi.org/10.1038/s41598-024-52383-y>, URL <https://doi.org/10.1038/s41598-024-52383-y>
 2165
 2166
 2167 Pöhlmann KMT, Föcker J, Dickinson P, et al (2021) The effect of motion direction
 2168 and eccentricity on vection, VR sickness and head movements in virtual reality.
 2169 *Multisensory Research* 34(6):623–662. <https://doi.org/10.1163/22134808-bja10049>
 2170
 2171 Pollick FE, Paterson HM, Bruderlin A, et al (2001) Perceiving affect from arm move-
 2172 ment. *Cognition* 82(2):B51–B61. [https://doi.org/10.1016/S0010-0277\(01\)00147-0](https://doi.org/10.1016/S0010-0277(01)00147-0),
 2173 URL [https://doi.org/10.1016/S0010-0277\(01\)00147-0](https://doi.org/10.1016/S0010-0277(01)00147-0)
 2174
 2175 Prpa M, Fdili-Alaoui S, Schiphorst T, et al (2020) Articulating experience. In: Pro-
 2176 ceedings of the 2020 CHI Conference on Human Factors in Computing Systems.
 2177 ACM, pp 1–14, <https://doi.org/10.1145/3313831.3376664>, URL <https://doi.org/10.1145/3313831.3376664>
 2178
 2179 Prugger J, Derdiyok E, Dinkelacker J, et al (2022) The altered states database:
 2180 Psychometric data from a systematic literature review. *Scientific Data* 9(1):720.
 2181 <https://doi.org/10.1038/s41597-022-01822-4>
 2182
 2183 Qin Z, Yu Y, Gu H, et al (2024) Sas macro programme for Bang’s Blinding Index to
 2184 assess and visualise the success of blinding in randomised controlled trials. *General*
 2185 *Psychiatry* 37:e101578. <https://doi.org/10.1136/gpsych-2024-101578>
 2186
 2187 Ramachandran VS, Hirstein W (1999) The science of art: A neurological theory of
 2188 aesthetic experience. *Journal of Consciousness Studies* 6(6–7):15–31
 2189
 2190 Reeves WT (1983) Particle systems—a technique for modeling a class of fuzzy
 2191 objects. *ACM Transactions on Graphics* 2(2):91–108. <https://doi.org/10.1145/357318.357320>, URL <https://doi.org/10.1145/357318.357320>
 2192
 2193 Reynolds CW (1987) Flocks, herds, and schools: A distributed behavioral model. In:
 2194 Proceedings of the 14th Annual Conference on Computer Graphics and Interactive
 2195 Techniques. ACM, pp 25–34, <https://doi.org/10.1145/37401.37406>, URL <https://doi.org/10.1145/37401.37406>
 2196
 2197
 2198 Rockstroh C, Blum J, Göritz AS (2019) Virtual reality in the application of heart
 2199 rate variability biofeedback. *International Journal of Human-Computer Studies*
 2200 130:209–220. <https://doi.org/10.1016/j.ijhcs.2019.06.011>, URL <https://doi.org/10.1016/j.ijhcs.2019.06.011>
 2201
 2202
 2203 Rockstroh C, Blum J, Göritz AS (2021) A mobile VR-based respiratory biofeed-
 2204 back game to foster diaphragmatic breathing. *Virtual Reality* 25(2):539–
 2205 552. <https://doi.org/10.1007/s10055-020-00471-5>, URL <https://doi.org/10.1007/s10055-020-00471-5>
 2206
 2207
 2208

Rodrigues FA, Peron TKDM, Ji P, et al (2016) The kuramoto model in complex networks. <i>Physics Reports</i> 610:1–98. https://doi.org/10.1016/j.physrep.2015.10.008	2209 2210 2211
Rogers B, Graham M (1979) Motion parallax as an independent cue for depth perception. <i>Perception</i> 8(2):125–134. https://doi.org/10.1068/p080125 , URL https://doi.org/10.1068/p080125	2212 2213 2214 2215
Romeo B, Kervadec E, Fauvel B, et al (2025) The intensity of the psychedelic experience is reliably associated with clinical improvements: A systematic review and meta-analysis. <i>Neuroscience & Biobehavioral Reviews</i> 172:106086. https://doi.org/10.1016/j.neubiorev.2025.106086 , URL https://doi.org/10.1016/j.neubiorev.2025.106086	2216 2217 2218 2219 2220
de Rooij A, Broekens J, Lamers MH (2013) Abstract expressions of affect. <i>International Journal of Synthetic Emotions</i> 4(1):1–31. https://doi.org/10.4018/jse.2013010101 , URL https://doi.org/10.4018/jse.2013010101	2221 2222 2223 2224
van Rooij M, Lobel A, Harris O, et al (2016) DEEP: A biofeedback virtual reality game for children at-risk for anxiety. In: <i>Proceedings of the 2016 CHI Conference Extended Abstracts on Human Factors in Computing Systems</i> . ACM, pp 1989–1997, https://doi.org/10.1145/2851581.2892452 , URL https://doi.org/10.1145/2851581.2892452	2225 2226 2227 2228 2229
Rook C, Panda S, Sice P, et al (2026) Heart garden: Visualizing heart rate data using the metaphor of a garden. <i>International Journal of Human–Computer Interaction</i> 42(1):114–130. https://doi.org/10.1080/10447318.2025.2504201 , URL https://doi.org/10.1080/10447318.2025.2504201	2230 2231 2232 2233 2234
Rosas FE, Mediano PAM, Timmermann C, et al (2023) The entropic heart: Tracking the psychedelic state via heart rate dynamics. <i>bioRxiv</i> p 2023.11.07.566008. https://doi.org/10.1101/2023.11.07.566008 , URL https://doi.org/10.1101/2023.11.07.566008	2235 2236 2237 2238 2239
Roseman L, Nutt DJ, Carhart-Harris RL (2018) Quality of acute psychedelic experience predicts therapeutic efficacy of psilocybin for treatment-resistant depression. <i>Frontiers in Pharmacology</i> 8:974. https://doi.org/10.3389/fphar.2017.00974	2240 2241 2242 2243
Rubin DS, Morgan RC, Pinchbeck D (2010) <i>Psychedelic: Optical and Visionary Art Since the 1960s</i> . MIT Press, https://doi.org/10.5860/choice.48-0074	2244 2245 2246
Sabnis AS, Shadid M, Yost GS, et al (2008) Human lung epithelial cells express a functional cold-sensing TRPM8 variant. <i>American Journal of Respiratory Cell and Molecular Biology</i> 39(4):466–474. https://doi.org/10.1165/rcmb.2007-0440OC	2247 2248 2249
Sanders JW, Zijlmans J (2021) Moving past mysticism in psychedelic science. <i>ACS Pharmacology & Translational Science</i> 4(3):1253–1255. https://doi.org/10.1021/acscptsci.1c00097 , URL https://doi.org/10.1021/acscptsci.1c00097	2250 2251 2252 2253 2254

2255 Santos M, Egerstedt M (2021) From motions to emotions: Can the fundamental emo-
 2256 tions be expressed in a robot swarm? *International Journal of Social Robotics*
 2257 13(4):751–764. <https://doi.org/10.1007/s12369-020-00665-6>, URL [https://doi.org/](https://doi.org/10.1007/s12369-020-00665-6)
 2258 [10.1007/s12369-020-00665-6](https://doi.org/10.1007/s12369-020-00665-6)
 2259
 2260 Sarkar AA (2017) Functional correlation between breathing and emotional states.
 2261 *MOJ Anatomy & Physiology* 3(5):157–158. [https://doi.org/10.15406/mojap.2017.](https://doi.org/10.15406/mojap.2017.03.00108)
 2262 [03.00108](https://doi.org/10.15406/mojap.2017.03.00108)
 2263
 2264 Schmidt TT, Berkemeyer H (2018) The altered states database: Psychometric data of
 2265 altered states of consciousness. *Frontiers in Psychology* 9:1028. [https://doi.org/10.](https://doi.org/10.3389/fpsyg.2018.01028)
 2266 [3389/fpsyg.2018.01028](https://doi.org/10.3389/fpsyg.2018.01028)
 2267
 2268 Seiler A, Camilo M, Korostovtseva L, et al (2019) Prevalence of sleep-
 2269 disordered breathing after stroke and TIA. *Neurology* 92(7):e648–e654. [https:](https://doi.org/10.1212/WNL.0000000000006904)
 2270 [/doi.org/10.1212/WNL.0000000000006904](https://doi.org/10.1212/WNL.0000000000006904), URL [https://doi.org/10.1212/WNL.](https://doi.org/10.1212/WNL.0000000000006904)
 2271 [0000000000006904](https://doi.org/10.1212/WNL.0000000000006904)
 2272
 2273 Serino A (2019) Peripersonal space (PPS) as a multisensory interface between
 2274 the individual and the environment, defining the space of the self. *Neuroscience*
 2275 & *Biobehavioral Reviews* 99:138–159. [https://doi.org/10.1016/j.neubiorev.2019.01.](https://doi.org/10.1016/j.neubiorev.2019.01.016)
 2276 [016](https://doi.org/10.1016/j.neubiorev.2019.01.016)
 2277
 2278 Shamekhi A, Bickmore T (2018) Breathe deep. In: *Proceedings of the 12th EAI Inter-*
 2279 *national Conference on Pervasive Computing Technologies for Healthcare*. ACM, pp
 2280 108–117, <https://doi.org/10.1145/3240925.3240940>, URL [https://doi.org/10.1145/](https://doi.org/10.1145/3240925.3240940)
 2281 [3240925.3240940](https://doi.org/10.1145/3240925.3240940)
 2282
 2283 Shi L, Liu F, Liu Y, et al (2023) Biofeedback respiratory rehabilitation training system
 2284 based on virtual reality technology. *Sensors* 23(22):9025. [https://doi.org/10.3390/](https://doi.org/10.3390/s23229025)
 2285 [s23229025](https://doi.org/10.3390/s23229025), URL <https://doi.org/10.3390/s23229025>
 2286
 2287 Siebieszuk A, Płoński AF, Baranowski M (2025) Breathwork for chronic stress
 2288 and mental health: Does choosing a specific technique matter? *Medical Sciences*
 2289 13(3):127. <https://doi.org/10.3390/medsci13030127>
 2290
 2291 Spee BT, Arato J, Mikuni J, et al (2024) The dynamics of experiencing gestalt and
 2292 aha in cubist art: pupil responses and art evaluations show a complex interplay of
 2293 task, stimuli content, and time course. *Frontiers in Psychology* 15:1192565. [https:](https://doi.org/10.3389/fpsyg.2024.1192565)
 2294 [/doi.org/10.3389/fpsyg.2024.1192565](https://doi.org/10.3389/fpsyg.2024.1192565)
 2295
 2296 Spehar B, van Tonder GJ (2017) Koffka’s aesthetic gestalt. *Leonardo* 50(1):53–57.
 2297 https://doi.org/10.1162/leon.a_01020
 2298
 2299 Stepanova ER, Desnoyers-Stewart J, Pasquier P, et al (2020) Jel: Breathing together to
 2300 connect with others and nature. In: *Proceedings of the 2020 ACM Designing Inter-*
 active Systems Conference. ACM, pp 641–654, <https://doi.org/10.1145/3357236>.

3395532, URL https://doi.org/10.1145/3357236.3395532	2301
Stocker K, Hartmann M, Schmid Y, et al (2025) The 3d-ascr scale: A revalidation of the core dimensions of the altered states of consciousness rating scale 5d(11)-ascr for psychedelic research. <i>Journal of Psychopharmacology</i> https://doi.org/10.1177/02698811251397328	2302
Studerus E, Gamma A, Vollenweider FX (2010) Psychometric evaluation of the altered states of consciousness rating scale (oav). <i>PLOS ONE</i> 5(8):e12412. https://doi.org/10.1371/journal.pone.0012412	2303
Suzuki K, Garfinkel SN, Critchley HD, et al (2013) Multisensory integration across exteroceptive and interoceptive domains modulates self-experience in the rubber-hand illusion. <i>Neuropsychologia</i> 51(13):2909–2917. https://doi.org/10.1016/j.neuropsychologia.2013.08.014	2304
Taves A (2020) Mystical and other alterations in sense of self: An expanded framework for studying nonordinary experiences. <i>Perspectives on Psychological Science</i> 15(3):669–690. https://doi.org/10.1177/1745691619895047 , URL https://doi.org/10.1177/1745691619895047	2305
Terhune DB, Luke DP, Kaelen M, et al (2016) A placebo-controlled investigation of synaesthesia-like experiences under LSD. <i>Neuropsychologia</i> 88:28–34. https://doi.org/10.1016/j.neuropsychologia.2016.04.005	2306
Tinga AM, Nyklíček I, Jansen MP, et al (2019) Respiratory biofeedback does not facilitate lowering arousal in meditation through virtual reality. <i>Applied Psychophysiology and Biofeedback</i> 44(1):51–59. https://doi.org/10.1007/s10484-018-9421-5 , URL https://doi.org/10.1007/s10484-018-9421-5	2307
Tsakiris M, Tajadura-Jiménez A, Costantini M (2011) Just a heartbeat away from one’s body: Interoceptive sensitivity predicts malleability of body-representations. <i>Proceedings of the Royal Society B: Biological Sciences</i> 278(1717):2470–2476. https://doi.org/10.1098/rspb.2010.2547	2308
Tsuchiya N, Wilke M, Frässle S, et al (2015) No-report paradigms: Extracting the true neural correlates of consciousness. <i>Trends in Cognitive Sciences</i> 19(12):757–770. https://doi.org/10.1016/j.tics.2015.10.002	2309
Uthaug MV, Mason NL, Havenith MN, et al (2021) An experience with Holotropic Breathwork is associated with improvement in non-judgement and satisfaction with life while reducing symptoms of stress in a Czech-speaking population. <i>Journal of Psychedelic Studies</i> 5(3):176–189. https://doi.org/10.1556/2054.2021.00193 , URL https://doi.org/10.1556/2054.2021.00193	2310
Vicary S, Sperling M, von Zimmermann J, et al (2017) Joint action aesthetics. <i>PLOS ONE</i> 12(7):e0180101. https://doi.org/10.1371/journal.pone.0180101	2311
	2312
	2313
	2314
	2315
	2316
	2317
	2318
	2319
	2320
	2321
	2322
	2323
	2324
	2325
	2326
	2327
	2328
	2329
	2330
	2331
	2332
	2333
	2334
	2335
	2336
	2337
	2338
	2339
	2340
	2341
	2342
	2343
	2344
	2345
	2346

2347 Vidal J, Andreu CI, Wrzesien M, et al (2025) Inducing selflessness through
 2348 a numadelic virtual reality experience: a preliminary study. *Virtual Reality*
 2349 30(1). <https://doi.org/10.1007/s10055-025-01293-z>, URL <https://doi.org/10.1007/s10055-025-01293-z>
 2350
 2351
 2352 Wagemans J, Elder JH, Kubovy M, et al (2012) A century of Gestalt psychol-
 2353 ogy in visual perception: I. perceptual grouping and figure-ground organization.
 2354 *Psychological Bulletin* 138(6):1172–1217. <https://doi.org/10.1037/a0029333>, URL
 2355 <https://doi.org/10.1037/a0029333>
 2356
 2357 Wald IY, Maimon A, Keniger de Andrade Gensas L, et al (2023) Breathing based
 2358 immersive interactions for enhanced agency and body awareness: A claustrophobia
 2359 motivated study. In: *Extended Abstracts of the 2023 CHI Conference on Human*
 2360 *Factors in Computing Systems*, pp 1–7, <https://doi.org/10.1145/3544549.3585897>,
 2361 URL <https://doi.org/10.1145/3544549.3585897>
 2362
 2363 Weng HY, Feldman JL, Leggio L, et al (2021) Interventions and manipulations of
 2364 interoception. *Trends in Neurosciences* 44(1):52–62. <https://doi.org/10.1016/j.tins.2020.09.010>
 2365
 2366 Yoon JM, Im SC, Kim K (2022) Effects of diaphragmatic breathing and pursed lip
 2367 breathing exercises on the pulmonary function and walking endurance in patients
 2368 with chronic stroke: a randomised controlled trial. *International Journal of Therapy*
 2369 *and Rehabilitation* 29(8):1–11. <https://doi.org/10.12968/ijtr.2021.0027>, URL <https://doi.org/10.12968/ijtr.2021.0027>
 2370
 2371
 2372 Zaccaro A, Piarulli A, Laurino M, et al (2018) How breath-control can change your life:
 2373 a systematic review on psycho-physiological correlates of slow breathing. *Frontiers*
 2374 *in Human Neuroscience* 12:353. <https://doi.org/10.3389/fnhum.2018.00353>
 2375
 2376 Zaccaro A, Piarulli A, Melosini L, et al (2022) Neural correlates of non-ordinary
 2377 states of consciousness in pranayama practitioners: the role of slow nasal breathing.
 2378 *Frontiers in Systems Neuroscience* 16:803904. <https://doi.org/10.3389/fnsys.2022.803904>
 2379
 2380
 2381 Zhou Y, Sun B, Li Q, et al (2011) Sensitivity of bronchopulmonary receptors to cold
 2382 and heat mediated by transient receptor potential cation channel subtypes in an ex
 2383 vivo rat lung preparation. *Respiratory Physiology & Neurobiology* 177(3):327–332.
 2384 <https://doi.org/10.1016/j.resp.2011.05.011>
 2385
 2386
 2387
 2388
 2389
 2390
 2391
 2392