

Supplementary Information

Viscereality: Bioresponsive Virtual Reality for Box Breathing and Altered-State Experience

George Fejer, Till Holzapfel, Taru Hirvonen, Anestis Lalidis Mateo, Johannes Blum, Michael Gaebler, and Bigna Lenggenhager

Virtual Reality

Corresponding author: George Fejer

University of Konstanz, Konstanz, Germany

Max Planck Institute for Human Cognitive and Brain Sciences, Germany

ALIUS Research Network, Germany

george.fejer@aliusresearch.org

S1 Deviations from preregistration

The preregistered phase-duration adherence measure was replaced by phase-locking value (PLV) because the recorded phase labels were too fragmented for stable phase-duration estimation without additional segmentation rules. The associated requirement of at least 70% valid cycles and the two-standard-deviation phase-error exclusion rule were replaced by a pooled lower-tail interquartile range (IQR) rule applied to condition-level PLV values.

For every non-PLV outcome, the preregistered order-adjusted repeated-measures analysis tests the condition effect after accounting for participant and numeric block position. It is implemented as a partial- F comparison between a participant-and-order model and the same model plus condition. Participant fixed effects provide within-participant blocking, and block position adjusts for presentation order. The finalized PLV omnibus remains unchanged. The two preregistered contrasts, pooled VR versus dark-screen control and symmetric versus asymmetric VR, were implemented as paired t -tests.

Wilcoxon signed-rank and Friedman tests were added as distribution-free checks. Benjamini–Hochberg false-discovery-rate correction was applied within each multi-test outcome family because the preregistration did not specify a multiplicity procedure. The subsequently published revised three-dimensional scoring model for the Altered States of Consciousness Rating Scale (3D-ASC_r; [Stocker et al. 2025](#)) was applied to the registered 11D-ASC scores as a post-registration higher-order summary; complete analyses of all 11 subscales remain in Section S3. Pictograph–ASC correspondence, condition-by-PLV models, and ASC–tracer correspondence analyses were treated as exploratory. All inferential analyses used the final included sample of $N = 39$.

The preregistration also specified correlations between each tracer summary and the corresponding retrospective ASC dimension. These were not implemented as inferential correlations because the three condition-level observations were nested within participants and no dependence-aware correspondence model had been specified. Section S5 therefore reports descriptive profile convergence only; formal tracer validity remains to be established in a dedicated dependence-aware analysis.

S2 PLV computation and quality screening

S2.1 Phase-locking value

PLV was calculated between the normalized sphere-radius trajectory displayed during each block and the ideal sphere-radius trajectory generated from the audio-instruction timing. Samples for which either series was non-finite were removed. Each series was mean-centred, and instantaneous phase was obtained from its analytic signal. Blocks with fewer than 20 paired finite samples were treated as missing.

For participant j in condition c ,

$$PLV_{jc} = \left| \frac{1}{T} \sum_{t=1}^T \exp(i[\phi_{jct}^{\text{observed}} - \phi_t^{\text{reference}}]) \right|, \quad (\text{S1})$$

where T is the number of paired finite samples. PLV ranges from 0 to 1, with larger values indicating a more stable phase relationship. PLV does not measure respiratory depth or tidal volume.

S2.2 Screening rule and exclusions

The screening distribution contained the three condition-level PLV values from each of the 47 tested participants. The lower fence was calculated as

$$Q_1 - 1.5(Q_3 - Q_1), \quad (\text{S2})$$

with $Q_1 = 0.695346$ and $Q_3 = 0.914776$, producing a cutoff of 0.366201. A participant was excluded from all conditions if any condition-level PLV was below the cutoff. Table S1 lists the excluded participants and their condition-level values. The remaining 39 participants formed the analytic sample used for all inferential analyses.

Table S1 Condition-level PLV values for participants excluded by the pooled lower-tail IQR quality-screening rule. Values below the cutoff of 0.366201 are marked with an asterisk. If any condition was flagged, the participant was excluded from all three conditions.

Participant	Dark-screen control	Symmetric VR	Asymmetric VR	Below-cutoff condition(s)
P08	0.487994	0.641091	0.363136*	Asymmetric
P12	0.045540*	0.858897	0.651905	Dark-screen control
P16	0.322388*	0.424615	0.923065	Dark-screen control
P18	0.046679*	0.424962	0.105354*	Dark-screen control; asymmetric
P23	0.950082	0.033076*	0.048964*	Symmetric; asymmetric
P27	0.859589	0.488851	0.336871*	Asymmetric
P32	0.723235	0.493400	0.101153*	Asymmetric
P40	0.441212	0.267064*	0.245893*	Symmetric; asymmetric

S3 11D-ASC: Altered States of Consciousness Rating Scale

This section reports analyses of the 11 dimensions of the Altered States of Consciousness Rating Scale (11D-ASC; Studerus et al. 2010), grouped under the Positive, Distressing, and Perceptual Effects composites reported in the main article. Within each composite, we compare the constituent dimensions’ pooled-VR-minus-dark-screen-control mean differences to identify the principal contributors to the higher-order result. These comparisons are descriptive and do not estimate independent or causal effects. Benjamini–Hochberg q_{FDR} values control the false-discovery rate across all 11 dimensions within each test family, rather than separately within each higher-order composite.

S3.1 Positive Effects

Table S2 ranks the six Positive Effects dimensions by paired effect size for the pooled-VR-versus-dark-screen-control contrast.

Table S2 Positive Effects dimensions ranked by paired Cohen’s d_z for the pooled-VR-versus-dark-screen-control contrast.

Dimension	Mean difference [95% CI]	Paired Cohen’s d_z	p	q_{FDR}
Experience of Unity	13.10 [5.53, 20.68]	0.56	.001	.004
Changed Meaning of Percepts	9.56 [3.69, 15.42]	0.53	.002	.006
Disembodiment	8.45 [1.84, 15.07]	0.41	.014	.025
Spiritual Experience	6.60 [1.43, 11.77]	0.41	.014	.025
Insightfulness	3.74 [−0.91, 8.39]	0.26	.112	.136
Blissful State	1.56 [−6.00, 9.13]	0.07	.678	.746

Note. Mean differences are pooled-VR-minus-dark-screen-control differences on the 0–100 scale; brackets show 95% confidence intervals. d_z is Cohen’s standardized effect size for paired observations. Benjamini–Hochberg q values are calculated across all 11 dimensions, not separately within each higher-order composite.

Experience of Unity.

Ratings were higher in symmetric VR ($M = 34.36$, $SD = 27.78$) and asymmetric VR ($M = 35.68$, $SD = 25.81$) than in the dark-screen control ($M = 21.92$, $SD = 22.33$). The registered order-adjusted three-condition result is [REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]. The corresponding Bayesian evidence was $\text{BF}_{10} = 64.02$; the data were approximately 64 times more likely under a model containing condition than under the order-only model. Pooled VR exceeded the control, $t(38) = 3.50$, $d_z = 0.56$, $q_{\text{FDR}} = .004$, $\text{BF}_{10} = 26.15$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.48$, $d_z = -0.08$, $q_{\text{FDR}} = .766$, $\text{BF}_{01} = 5.20$. The repeated-measures ANOVA and distribution-free analyses supported the same conclusion. The directional tests favored asymmetric VR over control, $\text{BF}_{10} = 87.69$, but not symmetric over asymmetric VR, $\text{BF}_{01} = 8.07$.

Exploratory checks found neither an order effect ($q_{\text{FDR}} = .934$) nor a condition-by-order interaction ($q_{\text{FDR}} = .872$). Both VR conditions increased Experience of Unity relative to control, with no meaningful difference between the two VR mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Changed Meaning of Percepts.

Ratings were higher in symmetric VR ($M = 19.93$, $SD = 19.65$) and asymmetric VR ($M = 20.41$, $SD = 20.62$) than in the dark-screen control ($M = 10.62$, $SD = 10.77$). The registered order-adjusted three-condition result is [REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]. The corresponding Bayesian evidence was $\text{BF}_{10} = 12.56$; the data were approximately 13 times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 3.30$, $d_z = 0.53$, $q_{\text{FDR}} = .006$, $\text{BF}_{10} = 15.81$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.17$, $d_z = -0.03$, $q_{\text{FDR}} = .866$, $\text{BF}_{01} = 5.72$. The repeated-measures ANOVA and distribution-free analyses supported the same conclusion. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .345$) nor a condition-by-order interaction ($q_{\text{FDR}} = .819$). Both VR conditions increased Changed Meaning of Percepts relative to control, without evidence of a difference between the two mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Disembodiment.

Ratings were higher in symmetric VR ($M = 29.55$, $SD = 27.87$) and asymmetric VR ($M = 30.40$, $SD = 25.49$) than in the dark-screen control ($M = 21.52$, $SD = 22.40$). The registered order-adjusted three-condition result is [REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]. The corresponding Bayesian evidence was $\text{BF}_{10} = 4.94$; the data were approximately five times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 2.59$, $d_z = 0.41$, $q_{\text{FDR}} = .025$, $\text{BF}_{10} = 3.17$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.39$, $d_z = -0.06$, $q_{\text{FDR}} = .766$, $\text{BF}_{01} = 5.39$. The repeated-measures ANOVA supported the condition effect, while the FDR-adjusted Friedman result was weaker. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .817$) nor a condition-by-order interaction ($q_{\text{FDR}} = .872$). VR increased Disembodiment relative to control, without a difference between the VR mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Spiritual Experience.

Ratings were higher in symmetric VR ($M = 17.97$, $SD = 22.77$) and asymmetric VR ($M = 18.91$, $SD = 22.91$) than in the dark-screen control ($M = 11.85$, $SD = 17.12$). The registered order-adjusted three-condition result is [REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]. The corresponding Bayesian evidence was $\text{BF}_{10} = 5.60$; the data were approximately six times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 2.58$, $d_z = 0.41$, $q_{\text{FDR}} = .025$, $\text{BF}_{10} = 3.13$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.60$, $d_z = -0.10$, $q_{\text{FDR}} = .766$,

$BF_{01} = 4.90$. The repeated-measures ANOVA and distribution-free analyses supported the same conclusion. Exploratory checks found neither an order effect ($q_{FDR} = .817$) nor a condition-by-order interaction ($q_{FDR} = .872$). VR increased Spiritual Experience relative to control, without evidence of a difference between the two mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Insightfulness.

Mean ratings were 17.54 ($SD = 18.11$) in the dark-screen control, 22.42 ($SD = 21.33$) in symmetric VR, and 20.14 ($SD = 18.56$) in asymmetric VR. The registered order-adjusted three-condition result is **[REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]**. The Bayesian analysis favored the order-only model, $BF_{01} = 3.31$; the observed data were approximately three times more likely without a condition effect. Neither pooled VR versus control, $t(38) = 1.63$, $d_z = 0.26$, $q_{FDR} = .136$, $BF_{01} = 1.73$, nor symmetric versus asymmetric VR, $t(38) = 0.84$, $d_z = 0.13$, $q_{FDR} = .766$, $BF_{01} = 4.16$, provided reliable evidence of a difference. The sensitivity and distribution-free analyses agreed. Exploratory checks found neither an order effect ($q_{FDR} = .980$) nor a condition-by-order interaction ($q_{FDR} = .819$). Insightfulness was descriptively higher during VR, but the tests did not establish a reliable condition effect. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

Blissful State.

Mean ratings were 29.86 ($SD = 21.60$) in the dark-screen control, 29.63 ($SD = 23.30$) in symmetric VR, and 33.22 ($SD = 25.51$) in asymmetric VR. The registered order-adjusted three-condition result is **[REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]**. The Bayesian analysis favored the order-only model, $BF_{01} = 7.62$; the data were approximately eight times more likely without a condition effect. Neither pooled VR versus control, $t(38) = 0.42$, $d_z = 0.07$, $q_{FDR} = .746$, $BF_{01} = 5.34$, nor symmetric versus asymmetric VR, $t(38) = -1.31$, $d_z = -0.21$, $q_{FDR} = .766$, $BF_{01} = 2.61$, provided reliable evidence of a difference. The sensitivity and distribution-free analyses agreed, and the directional tests did not support either registered step. Exploratory checks found neither an order effect ($q_{FDR} = .934$) nor a condition-by-order interaction ($q_{FDR} = .872$). Blissful State did not differ reliably among the three conditions. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

S3.2 Distressing Effects

Table S3 ranks the two Distressing Effects dimensions by paired effect size for the pooled-VR-versus-dark-screen-control contrast.

Table S3 Distressing Effects dimensions ranked by paired Cohen’s d_z for the pooled-VR-versus-dark-screen-control contrast.

Dimension	Mean difference [95% CI]	Paired Cohen’s d_z	p	q_{FDR}
Impaired Control and Cognition	5.47 [0.32, 10.62]	0.34	.038	.060
Anxiety	4.22 [−0.23, 8.67]	0.31	.063	.086

Note. Mean differences are pooled-VR-minus-dark-screen-control differences on the 0–100 scale; brackets show 95% confidence intervals. d_z is Cohen’s standardized effect size for paired observations. Benjamini–Hochberg q values are calculated across all 11 dimensions, not separately within each higher-order composite.

Impaired Control and Cognition.

Mean ratings were 19.17 ($SD = 16.50$) in the dark-screen control, 24.17 ($SD = 20.02$) in symmetric VR, and 25.12 ($SD = 18.97$) in asymmetric VR. The registered order-adjusted three-condition result is **[REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]**. The preregistered pooled-VR contrast likewise provided weak nominal evidence, $t(38) = 2.15$, $p = .038$, $d_z = 0.34$, but not correction-robust evidence ($q_{FDR} = .060$). The corresponding Bayes factors did not clearly distinguish the effect and null models ($BF_{10} = 1.10$ and 1.35, respectively). Symmetric and asymmetric VR did not differ, $t(38) = -0.51$, $p = .615$, $d_z = -0.08$, $q_{FDR} = .766$, $BF_{01} = 5.14$. The Friedman check detected a distributional difference, but this secondary rank-based test did not assess the same mean-difference estimand or adjust for presentation order. Exploratory checks found neither an FDR-retained order effect ($q_{FDR} = .682$) nor a condition-by-order interaction ($q_{FDR} = .872$). Impaired Control and Cognition was higher during pooled VR by a modest amount, but the frequentist evidence was weak and sensitive to family correction, and the Bayesian evidence was inconclusive. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

Anxiety.

Mean ratings were 5.49 ($SD = 7.54$) in the dark-screen control, 10.49 ($SD = 13.83$) in symmetric VR, and 8.92 ($SD = 13.66$) in asymmetric VR. The registered order-adjusted three-condition result is **[REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]**. The Bayesian analysis slightly favored the order-only model, $BF_{01} = 1.34$. The pooled-VR contrast was also inconclusive, $t(38) = 1.92$, $d_z = 0.31$, $q_{FDR} = .086$, $BF_{01} = 1.11$, while symmetric and asymmetric VR did not differ, $t(38) = 0.86$, $d_z = 0.14$, $q_{FDR} = .766$, $BF_{01} = 4.10$. The distribution-free results were similarly not significant after correction. The exploratory order effect was nominally significant before correction but not after it ($p = .015$, $q_{FDR} = .085$); the condition-by-order interaction was not supported ($q_{FDR} = .872$). Anxiety was descriptively higher during VR, but the condition difference did not provide correction-robust evidence. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

S3.3 Perceptual Effects

Table S4 ranks the three Perceptual Effects dimensions by paired effect size for the pooled-VR-versus-dark-screen-control contrast.

Table S4 Perceptual Effects dimensions ranked by paired Cohen's d_z for the pooled-VR-versus-dark-screen-control contrast.

Dimension	Mean difference [95% CI]	Paired Cohen's d_z	p	q_{FDR}
Audio-Visual Synesthesia	25.56 [15.52, 35.61]	0.82	< .001	< .001
Elementary Imagery	19.92 [11.35, 28.49]	0.75	< .001	< .001
Complex Imagery	-0.45 [-7.73, 6.83]	-0.02	.901	.901

Note. Mean differences are pooled-VR-minus-dark-screen-control differences on the 0–100 scale; brackets show 95% confidence intervals. d_z is Cohen's standardized effect size for paired observations. Benjamini–Hochberg q values are calculated across all 11 dimensions, not separately within each higher-order composite.

Audio-Visual Synesthesia.

Ratings were markedly higher in symmetric VR ($M = 38.85$, $SD = 33.23$) and asymmetric VR ($M = 34.44$, $SD = 30.60$) than in the dark-screen control ($M = 11.09$, $SD = 17.82$). The registered order-adjusted three-condition result is [REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]. The corresponding Bayesian evidence was $BF_{10} = 416,663.96$; the data were over 400,000 times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 5.15$, $d_z = 0.82$, $q_{FDR} < .001$, $BF_{10} = 2,408.46$, whereas symmetric and asymmetric VR did not differ reliably, $t(38) = 1.41$, $d_z = 0.23$, $q_{FDR} = .766$, $BF_{01} = 2.32$. The repeated-measures and distribution-free analyses supported the condition effect. The directional analysis strongly favored asymmetric VR over control, $BF_{10} = 745.10$, but did not clearly distinguish symmetric from asymmetric VR. The exploratory order effect was nominally significant before correction but not afterward ($p = .010$, $q_{FDR} = .085$); the condition-by-order interaction was not supported ($q_{FDR} = .819$). Both VR conditions produced a large increase in Audio-Visual Synesthesia, with no reliable difference between mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Elementary Imagery.

Ratings were higher in symmetric VR ($M = 26.51$, $SD = 28.07$) and asymmetric VR ($M = 28.78$, $SD = 26.81$) than in the dark-screen control ($M = 7.73$, $SD = 13.43$). The registered order-adjusted three-condition result is [REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]. The corresponding Bayesian evidence was $BF_{10} = 7,796.37$; the data were almost 7,800 times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 4.71$, $d_z = 0.75$, $q_{FDR} < .001$, $BF_{10} = 669.92$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.69$, $d_z = -0.11$, $q_{FDR} = .766$, $BF_{01} = 4.65$. The repeated-measures and distribution-free analyses

supported the same conclusion. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .421$) nor a condition-by-order interaction ($q_{\text{FDR}} = .872$). Both VR conditions strongly increased Elementary Imagery relative to control, without evidence of a difference between the mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Complex Imagery.

Mean ratings were 15.41 ($SD = 17.99$) in the dark-screen control, 16.38 ($SD = 17.87$) in symmetric VR, and 13.55 ($SD = 16.31$) in asymmetric VR. The registered order-adjusted three-condition result is [REGISTERED OMNIBUS STATISTICS PENDING OSF SOURCE LOCK]. The Bayesian analysis favored the order-only model, $BF_{01} = 8.83$; the data were approximately nine times more likely without a condition effect. Neither pooled VR versus control, $t(38) = -0.12$, $d_z = -0.02$, $q_{\text{FDR}} = .901$, $BF_{01} = 5.75$, nor symmetric versus asymmetric VR, $t(38) = 1.22$, $d_z = 0.20$, $q_{\text{FDR}} = .766$, $BF_{01} = 2.91$, provided reliable evidence of a difference. The sensitivity and distribution-free analyses agreed. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .980$) nor a condition-by-order interaction ($q_{\text{FDR}} = .872$). Complex Imagery was comparable across the three conditions. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

S3.4 Summary of fine-grained subscale contributions

The Positive Effects increase was principally associated with Experience of Unity and Changed Meaning of Percepts, followed by Disembodiment and Spiritual Experience. Together, these four fine-grained subscales account arithmetically for nearly nine-tenths of the composite increase. Their items assess experiences such as reduced separation between oneself and the environment, time or apparent contradictions merging into a unified whole, ordinary objects acquiring unusual significance or emotional salience, diminished bodily presence or floating sensations, and feelings of awe or spiritual connection. Blissful State and Insightfulness contributed comparatively little and did not change reliably, indicating that the Positive Effects result did not primarily reflect greater pleasure, inner peace, love, cognitive clarity, or original thought. The resulting profile is therefore better characterized as increased unity, altered meaning, bodily-boundary changes, and modest transcendence than as a general increase in positive mood.

The smaller Distressing Effects increase was distributed across Impaired Control and Cognition and Anxiety, with Impaired Control and Cognition making the somewhat larger contribution. Their items assess experiences such as diminished personal agency, difficulty making decisions or sustaining a coherent thought, feelings of paralysis or isolation, unexplained fear, perceived threat, strangely altered surroundings, and anticipation that something harmful might occur. Both subscales were descriptively higher during VR, but neither remained significant after correction across the 11 dimensions, and their Bayesian results did not clearly distinguish effect from null models. The significant composite result is therefore best interpreted as the accumulation of two modest and compatible distress-related tendencies rather than as a pronounced or clearly localized adverse effect.

The Perceptual Effects increase was almost entirely associated with Audio-Visual Synesthesia and Elementary Imagery. These subscales assess phenomena such as sounds appearing to influence perceived shapes or colors and the perception of simple patterns, colors, lights, or flashes. Complex Imagery, which instead assesses vivid internally generated scenes, memories, fantasy images, and imaginative content, remained unchanged. The perceptual result therefore reflects elementary visual phenomena and perceived cross-modal influence rather than a general intensification of internally generated imagery. Because the strongest item themes closely resemble properties directly supplied by the visually rich VR environment, the magnitude of this composite also reflects overlap between the intervention content and the questionnaire’s perceptual criteria. Across all fine-grained subscales, there was no reliable symmetric-versus-asymmetric difference and no FDR-corrected evidence that the condition pattern was influenced by the order in which participants underwent the three conditions.

S4 Pictographic self-related measures and ASC correspondence

S4.1 Scoring and analytic approach

These post hoc analyses examined whether participant-level changes in the three pictographic self-related measures corresponded with changes in ASC outcomes. They were not specified in the preregistration and are treated as exploratory. Only the Spatial Frame of Reference Continuum (SFoRC) showed a reliable pooled-VR-versus-dark-screen-control condition effect; the analyses below therefore do not assume that VR changed all three pictographic outcomes. Instead, they test whether individual differences in their change scores corresponded with changes in subjective experience. For construct-oriented interpretation, we focused on ASC subscales and items concerning unity with the environment, spatial or bodily disembodiment, blissful experience, and spiritual experience. This thematic focus determined which findings were highlighted in the main article, but all significance decisions were based on correction across the complete composite, subscale, or item-level screen. The released Perceived Body Boundaries pictograph progresses from a diffuse silhouette at A to a sharply delineated silhouette at G. Because the stored A–G coding maps these responses to 1–7, higher stored values indicate stronger boundary salience. We reversed this score for the analyses in this section so that higher values indicate greater boundary dissolution. Higher SFoRC and Small Self ratings respectively indicate greater perceived extension of the self into surrounding space and greater self-diminishment relative to the surroundings.

The primary estimand was the Spearman correlation between pooled-VR-minus-dark-screen-control changes in each pictograph and the corresponding change in each ASC outcome. This tests whether participants showing larger VR-related changes in pictographic self-representation also showed larger VR-related changes in subjective effects. Two-sided permutation p values used 10,000 random permutations, and percentile confidence intervals used 2,000 participant-bootstrap resamples. To examine mapping specificity, we separately correlated symmetric-minus-asymmetric changes

and tested condition-by-pictograph interactions in order-adjusted participant fixed-effects models with participant-clustered covariance. Pictograph ratings were centered within participant for the interaction models. Benjamini–Hochberg correction was applied separately across the 9 composite, 33 subscale, and 126 item-level associations and separately within each interaction family.

S4.2 Composite-level correspondence

Table S5 Pooled-VR-minus-dark-screen-control change correlations between pictographic measures and 3D-ASCr composites.

Composite	Boundary dissolution	SFoRC	Small Self
Positive Effects	.25 (.276)	.70 (.001)	.46 (.015)
Distressing Effects	−.04 (.885)	.07 (.842)	.02 (.885)
Perceptual Effects	.20 (.334)	.21 (.334)	.28 (.234)

Note. Cells report Spearman ρ with the Benjamini–Hochberg q value in parentheses. Correction was applied across all nine pictograph–composite tests. Boundary dissolution is reverse-scored so that higher values indicate greater dissolution.

Table S5 reports the complete pictograph–3D-ASCr composite screen. Greater SFoRC extension covaried strongly with greater Positive Effects, $\rho = .70$, $q = .001$. Greater Small Self ratings also covaried with Positive Effects, $\rho = .46$, $q = .015$. Boundary dissolution did not show a reliable association with Positive Effects, and none of the pictographs showed a correction-robust association with Distressing or Perceptual Effects. Rank-partial follow-ups controlling the other two pictographs retained the Positive Effects associations for SFoRC, partial $r = .64$, $q < .001$, and Small Self, partial $r = .38$, $q = .028$, but not boundary dissolution, partial $r = .04$, $q = .802$; these three follow-ups were corrected together.

S4.3 Fine-grained ASC correspondence

Table S6 reports all 33 pictograph–subscale associations. The SFoRC association with Positive Effects was distributed across Experience of Unity, Blissful State, Insightfulness, and Spiritual Experience. The association with Disembodiment was more moderate and did not survive correction across the full subscale family, $\rho = .36$, $q = .081$. Greater Small Self ratings covaried with Blissful State, Spiritual Experience, and Audio-Visual Synesthesia. Boundary dissolution did not show a correction-robust association with any ASC subscale, including Disembodiment, $\rho = .11$, $q = .591$.

Table S6 Pooled-VR-minus-dark-screen-control change correlations between pictographic measures and all 11 ASC subscales.

ASC subscale	Boundary dissolution	SFoRC	Small Self
Experience of Unity	.14 (.492)	.68 (.002)	.36 (.081)
Spiritual Experience	.19 (.357)	.41 (.049)	.46 (.021)
Blissful State	.24 (.277)	.65 (.002)	.52 (.012)
Insightfulness	.29 (.174)	.51 (.012)	.36 (.081)
Disembodiment	.11 (.591)	.36 (.081)	.22 (.292)
Impaired Control and Cognition	.19 (.357)	.28 (.188)	.15 (.469)
Anxiety	-.25 (.259)	-.32 (.114)	-.09 (.644)
Complex Imagery	.21 (.331)	.04 (.867)	.01 (.969)
Elementary Imagery	-.09 (.660)	.22 (.295)	.01 (.969)
Audio-Visual Synesthesia	.23 (.277)	.33 (.114)	.47 (.021)
Changed Meaning of Percepts	.13 (.538)	.36 (.081)	.20 (.331)

Note. Cells report Spearman ρ with the Benjamini–Hochberg q value in parentheses. Correction was applied across all 33 pictograph–subscale tests.

Item-level follow-ups clarified the content of these associations. Within Disembodiment, SFoRC change covaried with floating, $\rho = .55$, $q = .015$, and altered self-location outside the body, $\rho = .45$, $q = .044$, but not with bodylessness. The broader item-level SFoRC pattern also included all three Blissful State items and three Experience of Unity items. Small Self associations included items concerning inner peace, pleasure, spiritual connection, and sound-related colour changes. Table S7 reports every association surviving correction across all 126 item-level tests; no boundary-dissolution association survived this correction.

Table S7 Item-level pictograph–ASC associations surviving correction across 126 tests.

Pictograph	Item	Subscale	Brief content	ρ (q)
SFoRC	Q39	Blissful State	Inner peace	.69 (.008)
SFoRC	Q15	Experience of Unity	Unity with environment	.64 (.008)
SFoRC	Q18	Experience of Unity	Timelessness	.58 (.008)
SFoRC	Q19	Experience of Unity	Resolution of conflict	.57 (.013)
SFoRC	Q29	Disembodiment	Floating	.55 (.015)
SFoRC	Q03	Blissful State	Boundless pleasure	.51 (.016)
Small Self	Q39	Blissful State	Inner peace	.51 (.016)
Small Self	Q34	Audio-Visual Synesthesia	Sound-related colour change	.49 (.017)
SFoRC	Q12	Changed Meaning of Percepts	Altered meaning	.49 (.028)
SFoRC	Q24	Insightfulness	Profundity	.47 (.028)
SFoRC	Q36	Impaired Control and Cognition	Reduced personal will	.48 (.033)
SFoRC	Q16	Impaired Control and Cognition	Difficulty judging importance	.45 (.039)
Small Self	Q02	Spiritual Experience	Connection to a superior power	.46 (.044)
Small Self	Q03	Blissful State	Boundless pleasure	.45 (.044)
SFoRC	Q41	Blissful State	All-embracing love	.44 (.044)
SFoRC	Q28	Disembodiment	Location outside the body	.45 (.044)

Note. Item descriptions are concise content labels rather than replacements for the administered questionnaire wording. q values control the false-discovery rate across all 126 pictograph–item tests.

507 S4.4 Intervention and condition specificity

508 The primary change-score results explicitly concern intervention-related change: they
509 correlate each participant’s pooled-VR-minus-control pictograph change with that
510 participant’s pooled-VR-minus-control ASC change. To test whether correspondence
511 instead depended on the visual mapping, we repeated the analysis using symmetric-
512 minus-asymmetric change scores. No association survived correction among the nine
513 3D-ASCr composite tests (minimum $q = .142$) or the 33 subscale tests (minimum
514 $q = .082$). Thus, the data did not establish that participants showing a larger pic-
515 tographic response to one VR mapping also showed a larger ASC response to that
516 mapping.

517 The participant fixed-effects interaction models gave the same overall conclusion
518 about mapping specificity. No symmetric-versus-asymmetric pictograph-slope interac-
519 tion survived correction for any composite (all $q \geq .959$) or subscale (all $q \geq .999$).
520 The omnibus condition-by-SFoRC interaction for Positive Effects survived correction
521 across the nine composite models, $q = .038$, but neither the planned pooled-VR-versus-
522 control slope contrast, $q = .111$, nor the symmetric-versus-asymmetric slope contrast,
523 $q = .959$, survived correction. We therefore treat that omnibus result as inconclusive
524 rather than as evidence that condition reliably changed the SFoRC–Positive Effects
525 relationship.

526 Taken together, the exploratory analyses show that participants with larger pooled
527 VR-related increases in spatial self-extension also tended to show larger increases in
528 Unity and Positive Effects. SFoRC was also the only pictographic measure showing
529 a reliable pooled-VR-versus-dark-screen-control condition effect, making these asso-
530 ciations the clearest evidence of coordinated intervention-related variation across the
531 two measurement domains. Small Self change covaried with blissful and spiritual
532 experience, but no overall VR-related Small Self effect was established. Boundary dis-
533 solution showed no correction-robust association with its thematically closest ASC
534 subscale, Disembodiment. None of these relationships was specific to the symmetric or
535 asymmetric mapping. These exploratory associations may highlight overlap between
536 the measurement domain of the sense of self assessed through pictographs and the
537 subjective phenomenology captured by ASC subdomains.

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

541 S5 Temporal experience tracer analyses

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

543 S5.1 Task and data reduction

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

Temporal experience tracers are psychometric time-series ratings in which participants
retrospectively draw intensity curves for specified experiential dimensions over a ses-
sion’s time course (Jachs 2021; Damci et al. 2023; Lewis-Healey et al. 2024). Unlike
global retrospective ratings, tracers preserve temporal dynamics. For each of the 11
ASC dimensions, one prompt consolidates the constituent questionnaire items into a
construct-level description, yielding one tracer per dimension rather than 42 item-level
ratings.

Figure S1 illustrates the distinction between peak intensity and cumulative AUC.

After each condition, participants draw the 11 intensity curves on a world-space time-by-intensity surface using the VR controller. We project controller position onto the drawing plane, gate input through a trigger hold, and emit points at constant spatial spacing regardless of hand speed. We map reconstructed time and intensity to normalized 0–1 coordinates. Peak is the maximum sampled intensity. AUC is the trapezoidal integral of intensity after ordering points by reconstructed time, normalized to the 0–1 plotting interval. We analyse peak and AUC as separate exploratory families.

S5.2 Condition contrasts

Tracer peak and AUC profiles broadly follow the condition profiles observed for the 11D-ASC questionnaire, with the strongest pooled-VR differences clustering on Experience of Unity, Elementary Imagery, and Audio-Visual Synesthesia (Figure S2). After correction across the 11 dimensions within each tracer family, pooled VR exceeded the dark-screen control for all three dimensions for both peak and AUC. The paired effects were moderate for Experience of Unity and Audio-Visual Synesthesia and large for Elementary Imagery. Several smaller nominal differences did not survive FDR correction. No symmetric-versus-asymmetric comparison survived correction. Table S8 reports all peak and AUC contrasts.

Temporal Trace Profiles:
Peak versus Area Under the Curve




Fig. S1 Temporal experience tracers: peak and cumulative intensity. These hypothetical profiles illustrate information retained by psychometric time-series intensity ratings but not by a single retrospective visual-analogue-scale response. The red profile has a higher momentary peak, whereas the broader green profile has a larger area under the curve (AUC), reflecting greater cumulative intensity. Analyses in this paper focus on these two summaries because no hypothesis required finer-grained temporal features.

Table S8 Exploratory paired contrasts for tracer peak and AUC summaries ($N = 39$). VR denotes the mean of the symmetric and asymmetric VR conditions. All tests have 38 degrees of freedom. Benjamini–Hochberg adjustment was performed separately across the 11 dimensions for each tracer summary and contrast.

Summary	Dimension	Pooled VR vs control				Symmetric vs asymmetric			
		t	d_z	p	q	t	d_z	p	q
Peak	EU	2.93	0.47	.006	.021	0.83	0.13	.413	.780
	SE	0.50	0.08	.622	.684	−0.94	−0.15	.352	.780
	BS	−0.67	−0.11	.508	.621	0.28	0.04	.784	.784
	IS	2.18	0.35	.035	.090	0.78	0.12	.440	.780
	DIS	2.12	0.34	.041	.090	−1.48	−0.24	.146	.780
	ICC	1.80	0.29	.079	.124	0.56	0.09	.579	.780
	ANX	1.84	0.29	.074	.124	0.78	0.12	.441	.780
	CI	0.10	0.02	.920	.920	0.57	0.09	.574	.780
	EI	6.98	1.12	< .001	< .001	0.37	0.06	.713	.784
	AVS	3.34	0.54	.002	.010	0.89	0.14	.377	.780
	CMP	1.63	0.26	.112	.155	0.47	0.08	.638	.780
AUC	EU	3.26	0.52	.002	.009	1.31	0.21	.199	.850
	SE	0.76	0.12	.453	.553	−0.70	−0.11	.490	.850
	BS	−0.38	−0.06	.704	.774	0.39	0.06	.695	.850
	IS	2.24	0.36	.031	.057	0.80	0.13	.430	.850
	DIS	2.37	0.38	.023	.057	−1.19	−0.19	.240	.850
	ICC	2.29	0.37	.028	.057	0.26	0.04	.794	.873
	ANX	1.55	0.25	.129	.178	0.88	0.14	.382	.850
	CI	0.14	0.02	.886	.886	0.67	0.11	.507	.850
	EI	7.25	1.16	< .001	< .001	0.49	0.08	.627	.850
	AVS	3.58	0.57	< .001	.005	0.02	0.00	.985	.985
	CMP	1.62	0.26	.113	.178	0.42	0.07	.677	.850

S5.3 ASC–tracer correspondence

The questionnaire, tracer-peak, and tracer-AUC profiles show broadly similar relative patterns across the 11 dimensions (Figure S2). This comparison is descriptive: observations from the three conditions are nested within 39 participants, and no dependence-aware correspondence model was specified. The present results therefore support profile-level convergence, rather than demonstrating that the measurement formats are interchangeable.

Collecting one explained tracer for each dimension could, in principle, reduce item-level responses to 11 construct-level ratings while retaining distinctions between momentary peaks and cumulative experience. This may make assessment faster and more engaging, but the present study did not directly compare completion time, burden, enjoyment, reliability, or validity. Whether explanatory tracer prompts can replace or complement the 11D-ASC therefore requires dedicated validation, including dependence-aware correspondence analyses and tests of whether peak, AUC, or finer temporal features explain global retrospective ratings.




Fig. S2 Condition-wise altered-state profiles across all 11 dimensions for retrospective 11D-ASC questionnaire ratings (top; 0–100 visual-analogue scale), tracer peak intensity (middle; normalized 0–1 intensity), and tracer AUC (bottom; normalized 0–1 area). Lines show condition means for the included sample ($N = 39$). Separate radial scales are used; comparisons should concern profile shape and condition separation rather than absolute height across measurement formats.

710 S6 Reproducibility and file access

711 The study materials and reproducibility resources are available through three comple-
712 mentary records:

- 714 • **OSF project.** The preregistration record, deidentified participant-level data, inclu-
715 sion indicator, statistical outputs, analysis scripts, figure-generation scripts, study
716 materials, environment specifications, and replay instructions are available at <https://doi.org/10.17605/OSF.IO/64MWK>.
- 718 • **Zenodo archival release.** An immutable archival version of the reproducibility
719 package corresponding to the submitted manuscript is available at [https://doi.org/](https://doi.org/ZENODO_VERSION_DOI)
720 [ZENODO_VERSION_DOI](https://doi.org/ZENODO_VERSION_DOI).
- 721 • **GitHub repository.** The version-controlled reproducibility repository is
722 available at https://github.com/GeorgeFejer91/0.OSF_reproducibility_package.
723 The exact commit corresponding to the submitted manuscript is avail-
724 able at [https://github.com/GeorgeFejer91/0.OSF_reproducibility_package/tree/](https://github.com/GeorgeFejer91/0.OSF_reproducibility_package/tree/FINAL.SUBMISSION_COMMIT)
725 [FINAL.SUBMISSION_COMMIT](https://github.com/GeorgeFejer91/0.OSF_reproducibility_package/tree/FINAL.SUBMISSION_COMMIT).

726 The GitHub repository’s Start Here documentation identifies the analysis entry
727 points, output directories, environment files, and replay sequence.

729

730 References

731

732 Damci A, Hoeijmakers J, Hollander M, et al (2023) Acceptability, usability and feasi-
733 bility of experience sampling method in chronic secondary pain syndromes. *Frontiers*
734 in *Neurology* 14. <https://doi.org/10.3389/fneur.2023.1219236>

735

736 Jachs B (2021) The neurophenomenology of meditative states: Introducing temporal
737 experience tracing to capture subjective experience states and their neural corre-
738 lates. PhD thesis, University of Cambridge, <https://doi.org/10.17863/CAM.80153>,
739 URL <https://www.repository.cam.ac.uk/handle/1810/332709>

740

741 Lewis-Healey E, Tagliazucchi E, Canales-Johnson A, et al (2024) Breathwork-induced
742 psychedelic experiences modulate neural dynamics. *Cerebral Cortex* 34(8):bhae347.
743 <https://doi.org/10.1093/cercor/bhae347>

744

745 Stocker K, Hartmann M, Schmid Y, et al (2025) The 3d-ascr scale: A revalidation of
746 the core dimensions of the altered states of consciousness rating scale 5d(11)-asc
747 for psychedelic research. *Journal of Psychopharmacology* [https://doi.org/10.1177/](https://doi.org/10.1177/02698811251397328)
748 [02698811251397328](https://doi.org/10.1177/02698811251397328)

749

750 Studerus E, Gamma A, Vollenweider FX (2010) Psychometric evaluation of the altered
751 states of consciousness rating scale (oav). *PLOS ONE* 5(8):e12412. [https://doi.org/](https://doi.org/10.1371/journal.pone.0012412)
752 [10.1371/journal.pone.0012412](https://doi.org/10.1371/journal.pone.0012412)

753

754

755