

Supplementary Information

Viscerality: Bioresponsive Virtual Reality for Box Breathing and Altered-State Experience

George Fejer, Till Holzapfel, Taru Hirvonen, Anestis Laidis Mateo, Johannes Blum, Michael Gaebler, and Bigna Lenggenhager

Virtual Reality

Corresponding author: George Fejer

University of Konstanz, Konstanz, Germany

Max Planck Institute for Human Cognitive and Brain Sciences, Germany

ALIUS Research Network, Germany

george.fejer@aliusresearch.org

S1 Deviations from preregistrations

The preregistered phase-duration adherence measure was replaced by phase-locking value (PLV) because the recorded phase labels were too fragmented for stable phase-duration estimation without additional segmentation rules. The associated requirement of at least 70% valid cycles and the two-standard-deviation phase-error exclusion rule were replaced by a pooled lower-tail IQR rule applied to condition-level PLV values.

The preregistered repeated-measures ANOVA with presentation order as a covariate was replaced by an order-adjusted mixed-effects likelihood-ratio test comparing a condition-plus-order model with an order-only model and including participant as a random intercept. The two preregistered contrasts, pooled VR versus dark-screen control and symmetric versus asymmetric VR, were implemented as paired *t*-tests.

Wilcoxon signed-rank and Friedman tests were added as distribution-free checks. Benjamini–Hochberg false-discovery-rate correction was applied within each multi-test outcome family because the preregistration did not specify a multiplicity procedure. The subsequently published 3D-ASCr scoring system was applied to the registered 11D-ASC scores as a post-registration higher-order summary; complete analyses of all 11 subscales remain in Section S3. Condition-by-PLV models and ASC–tracer correspondence analyses were treated as exploratory. All inferential analyses used the final included sample of $N = 39$.

S2 PLV computation and quality screening

S2.1 Phase-locking value

PLV was calculated between the normalized sphere-radius trajectory displayed during each block and the ideal sphere-radius trajectory generated from the audio-instruction timing. Samples for which either series was non-finite were removed. Each series was mean-centred, and instantaneous phase was obtained from its analytic signal. Blocks with fewer than 20 paired finite samples were treated as missing.

For participant j in condition c ,

$$PLV_{jc} = \left| \frac{1}{T} \sum_{t=1}^T \exp(i[\phi_{jct}^{\text{observed}} - \phi_t^{\text{reference}}]) \right|, \quad (\text{S1})$$

where T is the number of paired finite samples. PLV ranges from 0 to 1, with larger values indicating a more stable phase relationship. PLV does not measure respiratory depth or tidal volume.

S2.2 Screening rule and exclusions

The screening distribution contained the three condition-level PLV values from each of the 47 tested participants. The lower fence was calculated as

$$Q_1 - 1.5(Q_3 - Q_1), \quad (\text{S2})$$

with $Q_1 = 0.695346$ and $Q_3 = 0.914776$, producing a cutoff of 0.366201. A participant was excluded from all conditions if any condition-level PLV was below the cutoff. Table S1 lists the excluded participants and their condition-level values. The remaining 39 participants formed the analytic sample used for all inferential analyses.

Table S1 Condition-level PLV values for participants excluded by the pooled lower-tail IQR quality-screening rule. Values below the cutoff of 0.366201 are marked with an asterisk. If any condition was flagged, the participant was excluded from all three conditions.

Participant	Black screen	Symmetric VR	Asymmetric VR	Below-cutoff condition(s)
P08	0.487994	0.641091	0.363136*	Asymmetric
P12	0.045540*	0.858897	0.651905	Black screen
P16	0.322388*	0.424615	0.923065	Black screen
P18	0.046679*	0.424962	0.105354*	Black screen; asymmetric
P23	0.950082	0.033076*	0.048964*	Symmetric; asymmetric
P27	0.859589	0.488851	0.336871*	Asymmetric
P32	0.723235	0.493400	0.101153*	Asymmetric
P40	0.441212	0.267064*	0.245893*	Symmetric; asymmetric

S3 11D-ASC: Altered States of Consciousness Rating Scale

This section reports analyses of the 11 dimensions of the Altered States of Consciousness Rating Scale (11D-ASC; Studerus et al. 2010), grouped under the Positive, Distressing, and Perceptual Effects composites reported in the main article. Within each composite, we compare the constituent dimensions' pooled-VR-minus-dark-screen-control mean differences to identify the principal contributors to the higher-order result. These comparisons are descriptive and do not estimate independent or causal

effects. Benjamini–Hochberg q_{FDR} values control the false-discovery rate across all 11 dimensions within each test family, rather than separately within each higher-order composite.

S3.1 Positive Effects

Table S2 Positive Effects dimensions ranked by paired Cohen’s d_z for the pooled-VR-versus-dark-screen-control contrast.

Dimension	Mean difference [95% CI]	Paired Cohen’s d_z	p	q_{FDR}
Experience of Unity	13.10 [5.53, 20.68]	0.56	.001	.004
Changed Meaning of Percepts	9.56 [3.69, 15.42]	0.53	.002	.006
Disembodiment	8.45 [1.84, 15.07]	0.41	.014	.025
Spiritual Experience	6.60 [1.43, 11.77]	0.41	.014	.025
Insightfulness	3.74 [−0.91, 8.39]	0.26	.112	.136
Blissful State	1.56 [−6.00, 9.13]	0.07	.678	.746

Note. Mean differences are pooled-VR-minus-dark-screen-control differences on the 0–100 scale; brackets show 95% confidence intervals. d_z is Cohen’s standardized effect size for paired observations. Benjamini–Hochberg q values are calculated across all 11 dimensions, not separately within each higher-order composite.

Experience of Unity.

Ratings were higher in symmetric VR ($M = 34.36$, $SD = 27.78$) and asymmetric VR ($M = 35.68$, $SD = 25.81$) than in the dark-screen control ($M = 21.92$, $SD = 22.33$). The order-adjusted three-condition test showed a condition effect, $\chi^2(2) = 16.07$, $q_{\text{FDR}} = .001$, with corresponding Bayesian evidence, $\text{BF}_{10} = 64.02$; the data were approximately 64 times more likely under a model containing condition than under the order-only model. Pooled VR exceeded the control, $t(38) = 3.50$, $d_z = 0.56$, $q_{\text{FDR}} = .004$, $\text{BF}_{10} = 26.15$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.48$, $d_z = -0.08$, $q_{\text{FDR}} = .766$, $\text{BF}_{01} = 5.20$. The repeated-measures ANOVA and distribution-free analyses supported the same conclusion. The directional tests favored asymmetric VR over control, $\text{BF}_{10} = 87.69$, but not symmetric over asymmetric VR, $\text{BF}_{01} = 8.07$. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .934$) nor a condition-by-order interaction ($q_{\text{FDR}} = .872$). Both VR conditions increased Experience of Unity relative to control, with no meaningful difference between the two VR mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Changed Meaning of Percepts.

Ratings were higher in symmetric VR ($M = 19.93$, $SD = 19.65$) and asymmetric VR ($M = 20.41$, $SD = 20.62$) than in the dark-screen control ($M = 10.62$, $SD = 10.77$). The order-adjusted three-condition test showed a condition effect, $\chi^2(2) = 11.99$, $q_{\text{FDR}} = .007$, with corresponding Bayesian evidence, $\text{BF}_{10} = 12.56$; the data were approximately 13 times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 3.30$, $d_z = 0.53$, $q_{\text{FDR}} =$

.006, $BF_{10} = 15.81$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.17$, $d_z = -0.03$, $q_{FDR} = .866$, $BF_{01} = 5.72$. The repeated-measures ANOVA and distribution-free analyses supported the same conclusion. Exploratory checks found neither an order effect ($q_{FDR} = .345$) nor a condition-by-order interaction ($q_{FDR} = .819$). Both VR conditions increased Changed Meaning of Percepts relative to control, without evidence of a difference between the two mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Disembodiment.

Ratings were higher in symmetric VR ($M = 29.55$, $SD = 27.87$) and asymmetric VR ($M = 30.40$, $SD = 25.49$) than in the dark-screen control ($M = 21.52$, $SD = 22.40$). The order-adjusted three-condition test showed a condition effect, $\chi^2(2) = 9.96$, $q_{FDR} = .013$, with corresponding Bayesian evidence, $BF_{10} = 4.94$; the data were approximately five times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 2.59$, $d_z = 0.41$, $q_{FDR} = .025$, $BF_{10} = 3.17$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.39$, $d_z = -0.06$, $q_{FDR} = .766$, $BF_{01} = 5.39$. The repeated-measures ANOVA supported the condition effect, while the FDR-adjusted Friedman result was weaker. Exploratory checks found neither an order effect ($q_{FDR} = .817$) nor a condition-by-order interaction ($q_{FDR} = .872$). VR increased Disembodiment relative to control, without a difference between the VR mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Spiritual Experience.

Ratings were higher in symmetric VR ($M = 17.97$, $SD = 22.77$) and asymmetric VR ($M = 18.91$, $SD = 22.91$) than in the dark-screen control ($M = 11.85$, $SD = 17.12$). The order-adjusted three-condition test showed a condition effect, $\chi^2(2) = 10.38$, $q_{FDR} = .012$, with corresponding Bayesian evidence, $BF_{10} = 5.60$; the data were approximately six times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 2.58$, $d_z = 0.41$, $q_{FDR} = .025$, $BF_{10} = 3.13$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.60$, $d_z = -0.10$, $q_{FDR} = .766$, $BF_{01} = 4.90$. The repeated-measures ANOVA and distribution-free analyses supported the same conclusion. Exploratory checks found neither an order effect ($q_{FDR} = .817$) nor a condition-by-order interaction ($q_{FDR} = .872$). VR increased Spiritual Experience relative to control, without evidence of a difference between the two mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Insightfulness.

Mean ratings were 17.54 ($SD = 18.11$) in the dark-screen control, 22.42 ($SD = 21.33$) in symmetric VR, and 20.14 ($SD = 18.56$) in asymmetric VR. The order-adjusted three-condition test did not show a condition effect, $\chi^2(2) = 3.33$, $q_{FDR} = .231$, and the Bayesian analysis favored the order-only model, $BF_{01} = 3.31$; the observed data were approximately three times more likely without a condition effect. Neither pooled VR versus control, $t(38) = 1.63$, $d_z = 0.26$, $q_{FDR} = .136$, $BF_{01} = 1.73$,

nor symmetric versus asymmetric VR, $t(38) = 0.84$, $d_z = 0.13$, $q_{\text{FDR}} = .766$, $\text{BF}_{01} = 4.16$, provided reliable evidence of a difference. The sensitivity and distribution-free analyses agreed. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .980$) nor a condition-by-order interaction ($q_{\text{FDR}} = .819$). Insightfulness was descriptively higher during VR, but the tests did not establish a reliable condition effect. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

Blissful State.

Mean ratings were 29.86 ($SD = 21.60$) in the dark-screen control, 29.63 ($SD = 23.30$) in symmetric VR, and 33.22 ($SD = 25.51$) in asymmetric VR. The order-adjusted three-condition test did not show a condition effect, $\chi^2(2) = 1.23$, $q_{\text{FDR}} = .594$, and the Bayesian analysis favored the order-only model, $\text{BF}_{01} = 7.62$; the data were approximately eight times more likely without a condition effect. Neither pooled VR versus control, $t(38) = 0.42$, $d_z = 0.07$, $q_{\text{FDR}} = .746$, $\text{BF}_{01} = 5.34$, nor symmetric versus asymmetric VR, $t(38) = -1.31$, $d_z = -0.21$, $q_{\text{FDR}} = .766$, $\text{BF}_{01} = 2.61$, provided reliable evidence of a difference. The sensitivity and distribution-free analyses agreed, and the directional tests did not support either registered step. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .934$) nor a condition-by-order interaction ($q_{\text{FDR}} = .872$). Blissful State did not differ reliably among the three conditions. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

S3.2 Distressing Effects

Table S3 Distressing Effects dimensions ranked by paired Cohen's d_z for the pooled-VR-versus-dark-screen-control contrast.

Dimension	Mean difference [95% CI]	Paired Cohen's d_z	p	q_{FDR}
Impaired Control and Cognition	5.47 [0.32, 10.62]	0.34	.038	.060
Anxiety	4.22 [-0.23, 8.67]	0.31	.063	.086

Note. Mean differences are pooled-VR-minus-dark-screen-control differences on the 0–100 scale; brackets show 95% confidence intervals. d_z is Cohen's standardized effect size for paired observations. Benjamini–Hochberg q values are calculated across all 11 dimensions, not separately within each higher-order composite.

Impaired Control and Cognition.

Mean ratings were 19.17 ($SD = 16.50$) in the dark-screen control, 24.17 ($SD = 20.02$) in symmetric VR, and 25.12 ($SD = 18.97$) in asymmetric VR. The order-adjusted three-condition result, $\chi^2(2) = 6.45$, $q_{\text{FDR}} = .062$, and pooled-VR contrast, $t(38) = 2.15$, $d_z = 0.34$, $q_{\text{FDR}} = .060$, did not remain significant after FDR correction. Bayesian evidence for these effects was correspondingly inconclusive ($\text{BF}_{10} = 1.10$ and 1.35, respectively). Symmetric and asymmetric VR did not differ, $t(38) = -0.51$, $d_z = -0.08$, $q_{\text{FDR}} = .766$, $\text{BF}_{01} = 5.14$. The Friedman check detected a distributional difference, but this secondary rank-based test did not assess the same mean-difference

estimand or adjust for presentation order and therefore did not alter the primary FDR-qualified conclusion. Exploratory checks found neither an order effect ($q_{\text{FDR}} = .682$) nor a condition-by-order interaction ($q_{\text{FDR}} = .872$). Impaired Control and Cognition tended to be higher during VR, but the evidence did not survive correction across the 11 dimensions and did not differ reliably between the mappings. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

Anxiety.

Mean ratings were 5.49 ($SD = 7.54$) in the dark-screen control, 10.49 ($SD = 13.83$) in symmetric VR, and 8.92 ($SD = 13.66$) in asymmetric VR. The order-adjusted three-condition test did not show a corrected condition effect, $\chi^2(2) = 5.35$, $q_{\text{FDR}} = .095$, and the Bayesian analysis slightly favored the order-only model, $\text{BF}_{01} = 1.34$. The pooled-VR contrast was also inconclusive, $t(38) = 1.92$, $d_z = 0.31$, $q_{\text{FDR}} = .086$, $\text{BF}_{01} = 1.11$, while symmetric and asymmetric VR did not differ, $t(38) = 0.86$, $d_z = 0.14$, $q_{\text{FDR}} = .766$, $\text{BF}_{01} = 4.10$. The distribution-free results were similarly not significant after correction. The exploratory order effect was nominally significant before correction but not after it ($p = .015$, $q_{\text{FDR}} = .085$); the condition-by-order interaction was not supported ($q_{\text{FDR}} = .872$). Anxiety was descriptively higher during VR, but the condition difference did not provide correction-robust evidence. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

S3.3 Perceptual Effects

Table S4 Perceptual Effects dimensions ranked by paired Cohen's d_z for the pooled-VR-versus-dark-screen-control contrast.

Dimension	Mean difference [95% CI]	Paired Cohen's d_z	p	q_{FDR}
Audio-Visual Synesthesia	25.56 [15.52, 35.61]	0.82	< .001	< .001
Elementary Imagery	19.92 [11.35, 28.49]	0.75	< .001	< .001
Complex Imagery	-0.45 [-7.73, 6.83]	-0.02	.901	.901

Note. Mean differences are pooled-VR-minus-dark-screen-control differences on the 0–100 scale; brackets show 95% confidence intervals. d_z is Cohen's standardized effect size for paired observations. Benjamini-Hochberg q values are calculated across all 11 dimensions, not separately within each higher-order composite.

Audio-Visual Synesthesia.

Ratings were markedly higher in symmetric VR ($M = 38.85$, $SD = 33.23$) and asymmetric VR ($M = 34.44$, $SD = 30.60$) than in the dark-screen control ($M = 11.09$, $SD = 17.82$). The order-adjusted three-condition test showed a condition effect, $\chi^2(2) = 35.71$, $q_{\text{FDR}} < .001$, with overwhelming Bayesian evidence, $\text{BF}_{10} = 416,663.96$; the data were over 400,000 times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 5.15$, $d_z = 0.82$, $q_{\text{FDR}} < .001$, $\text{BF}_{10} = 2,408.46$, whereas symmetric and asymmetric VR did

not differ reliably, $t(38) = 1.41$, $d_z = 0.23$, $q_{FDR} = .766$, $BF_{01} = 2.32$. The repeated-measures and distribution-free analyses supported the condition effect. The directional analysis strongly favored asymmetric VR over control, $BF_{10} = 745.10$, but did not clearly distinguish symmetric from asymmetric VR. The exploratory order effect was nominally significant before correction but not afterward ($p = .010$, $q_{FDR} = .085$); the condition-by-order interaction was not supported ($q_{FDR} = .819$). Both VR conditions produced a large increase in Audio-Visual Synesthesia, with no reliable difference between mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Elementary Imagery.

Ratings were higher in symmetric VR ($M = 26.51$, $SD = 28.07$) and asymmetric VR ($M = 28.78$, $SD = 26.81$) than in the dark-screen control ($M = 7.73$, $SD = 13.43$). The order-adjusted three-condition test showed a condition effect, $\chi^2(2) = 26.50$, $q_{FDR} < .001$, with overwhelming Bayesian evidence, $BF_{10} = 7,796.37$; the data were almost 7,800 times more likely under the condition model than under the order-only model. Pooled VR exceeded the control, $t(38) = 4.71$, $d_z = 0.75$, $q_{FDR} < .001$, $BF_{10} = 669.92$, whereas symmetric and asymmetric VR did not differ, $t(38) = -0.69$, $d_z = -0.11$, $q_{FDR} = .766$, $BF_{01} = 4.65$. The repeated-measures and distribution-free analyses supported the same conclusion. Exploratory checks found neither an order effect ($q_{FDR} = .421$) nor a condition-by-order interaction ($q_{FDR} = .872$). Both VR conditions strongly increased Elementary Imagery relative to control, without evidence of a difference between the mappings. There was no reliable evidence that the result was influenced by the order in which participants underwent the three conditions.

Complex Imagery.

Mean ratings were 15.41 ($SD = 17.99$) in the dark-screen control, 16.38 ($SD = 17.87$) in symmetric VR, and 13.55 ($SD = 16.31$) in asymmetric VR. The order-adjusted three-condition test did not show a condition effect, $\chi^2(2) = 0.74$, $q_{FDR} = .690$, and the Bayesian analysis favored the order-only model, $BF_{01} = 8.83$; the data were approximately nine times more likely without a condition effect. Neither pooled VR versus control, $t(38) = -0.12$, $d_z = -0.02$, $q_{FDR} = .901$, $BF_{01} = 5.75$, nor symmetric versus asymmetric VR, $t(38) = 1.22$, $d_z = 0.20$, $q_{FDR} = .766$, $BF_{01} = 2.91$, provided reliable evidence of a difference. The sensitivity and distribution-free analyses agreed. Exploratory checks found neither an order effect ($q_{FDR} = .980$) nor a condition-by-order interaction ($q_{FDR} = .872$). Complex Imagery was comparable across the three conditions. There was no reliable evidence that it was influenced by the order in which participants underwent the three conditions.

S3.4 Summary of fine-grained subscale contributions

The Positive Effects increase was principally associated with Experience of Unity and Changed Meaning of Percepts, followed by Disembodiment and Spiritual Experience. Together, these four fine-grained subscales account arithmetically for nearly nine-tenths of the composite increase. Their items assess experiences such as reduced separation between oneself and the environment, time or apparent contradictions

merging into a unified whole, ordinary objects acquiring unusual significance or emotional salience, diminished bodily presence or floating sensations, and feelings of awe or spiritual connection. Blissful State and Insightfulness contributed comparatively little and did not change reliably, indicating that the Positive Effects result did not primarily reflect greater pleasure, inner peace, love, cognitive clarity, or original thought. The resulting profile is therefore better characterized as increased unity, altered meaning, bodily-boundary changes, and modest transcendence than as a general increase in positive mood.

The smaller Distressing Effects increase was distributed across Impaired Control and Cognition and Anxiety, with Impaired Control and Cognition making the somewhat larger contribution. Their items assess experiences such as diminished personal agency, difficulty making decisions or sustaining a coherent thought, feelings of paralysis or isolation, unexplained fear, perceived threat, strangely altered surroundings, and anticipation that something harmful might occur. Both subscales were descriptively higher during VR, but neither remained significant after correction across the 11 dimensions, and their Bayesian results did not clearly distinguish effect from null models. The significant composite result is therefore best interpreted as the accumulation of two modest and compatible distress-related tendencies rather than as a pronounced or clearly localized adverse effect.

The Perceptual Effects increase was almost entirely associated with Audio-Visual Synesthesia and Elementary Imagery. These subscales assess phenomena such as sounds appearing to influence perceived shapes or colors and the perception of simple patterns, colors, lights, or flashes. Complex Imagery, which instead assesses vivid internally generated scenes, memories, fantasy images, and imaginative content, remained unchanged. The perceptual result therefore reflects elementary visual phenomena and perceived cross-modal influence rather than a general intensification of internally generated imagery. Because the strongest item themes closely resemble properties directly supplied by the visually rich VR environment, the magnitude of this composite also reflects overlap between the intervention content and the questionnaire’s perceptual criteria. Across all fine-grained subscales, there was no reliable symmetric-versus-asymmetric difference and no FDR-corrected evidence that the condition pattern was influenced by the order in which participants underwent the three conditions.

355

356

357

358

359

S4 Temporal experience tracer analyses

360

361

S4.1 Task and data reduction

Temporal experience tracers are psychometric time-series ratings in which participants retrospectively draw intensity curves for specified experiential dimensions over a session’s time course (Jachs 2021; Damci et al. 2023; Lewis-Healey et al. 2024). Unlike global retrospective ratings, tracers preserve temporal dynamics. For each of the 11 ASC dimensions, one prompt consolidates the constituent questionnaire items into a construct-level description, yielding one tracer per dimension rather than 42 item-level ratings.

After each condition, participants draw the 11 intensity curves on a world-space time-by-intensity surface using the VR controller. We project controller position onto the drawing plane, gate input through a trigger hold, and emit points at constant spatial spacing regardless of hand speed. We map reconstructed time and intensity to normalized 0–1 coordinates. Peak is the maximum sampled intensity. AUC is the trapezoidal integral of intensity after ordering points by reconstructed time, normalized to the 0–1 plotting interval. We analyse peak and AUC as separate exploratory families.

S4.2 Condition contrasts

Tracer peak and AUC profiles broadly follow the condition profiles observed for the 11D-ASC questionnaire, with the strongest pooled-VR differences clustering on Experience of Unity, Elementary Imagery, and Audio-Visual Synesthesia (Figure S2). After correction across the 11 dimensions within each tracer family, pooled VR exceeded the dark-screen control for all three dimensions for both peak and AUC. The paired effects were moderate for Experience of Unity and Audio-Visual Synesthesia and large for Elementary Imagery. Several smaller nominal differences did not survive FDR correction. No symmetric-versus-asymmetric comparison survived correction.

**Temporal Trace Profiles:
Peak versus Area Under the Curve**




Fig. S1 Temporal experience tracers: peak and cumulative intensity. These hypothetical profiles illustrate information retained by psychometric time-series intensity ratings but not by a single retrospective visual-analogue-scale response. The red profile has a higher momentary peak, whereas the broader green profile has a larger area under the curve (AUC), reflecting greater cumulative intensity. Analyses in this paper focus on these two summaries because no hypothesis required finer-grained temporal features.

Table S5 Exploratory paired contrasts for tracer peak and AUC summaries ($N = 39$). VR denotes the mean of the symmetric and asymmetric VR conditions. All tests have 38 degrees of freedom. Benjamini–Hochberg adjustment was performed separately across the 11 dimensions for each tracer summary and contrast.

Summary	Dimension	Pooled VR vs control				Symmetric vs asymmetric			
		t	d_z	p	q	t	d_z	p	q
Peak	EU	2.93	0.47	.006	.021	0.83	0.13	.413	.780
	SE	0.50	0.08	.622	.684	−0.94	−0.15	.352	.780
	BS	−0.67	−0.11	.508	.621	0.28	0.04	.784	.784
	IS	2.18	0.35	.035	.090	0.78	0.12	.440	.780
	DIS	2.12	0.34	.041	.090	−1.48	−0.24	.146	.780
	ICC	1.80	0.29	.079	.124	0.56	0.09	.579	.780
	ANX	1.84	0.29	.074	.124	0.78	0.12	.441	.780
	CI	0.10	0.02	.920	.920	0.57	0.09	.574	.780
	EI	6.98	1.12	< .001	< .001	0.37	0.06	.713	.784
	AVS	3.34	0.54	.002	.010	0.89	0.14	.377	.780
	CMP	1.63	0.26	.112	.155	0.47	0.08	.638	.780
AUC	EU	3.26	0.52	.002	.009	1.31	0.21	.199	.850
	SE	0.76	0.12	.453	.553	−0.70	−0.11	.490	.850
	BS	−0.38	−0.06	.704	.774	0.39	0.06	.695	.850
	IS	2.24	0.36	.031	.057	0.80	0.13	.430	.850
	DIS	2.37	0.38	.023	.057	−1.19	−0.19	.240	.850
	ICC	2.29	0.37	.028	.057	0.26	0.04	.794	.873
	ANX	1.55	0.25	.129	.178	0.88	0.14	.382	.850
	CI	0.14	0.02	.886	.886	0.67	0.11	.507	.850
	EI	7.25	1.16	< .001	< .001	0.49	0.08	.627	.850
	AVS	3.58	0.57	< .001	.005	0.02	0.00	.985	.985
	CMP	1.62	0.26	.113	.178	0.42	0.07	.677	.850

S4.3 ASC–tracer correspondence

The questionnaire, tracer-peak, and tracer-AUC profiles show broadly similar relative patterns across the 11 dimensions (Figure S2). This comparison is descriptive: observations from the three conditions are nested within 39 participants, and no dependence-aware correspondence model was specified. The present results therefore support profile-level convergence, rather than demonstrating that the measurement formats are interchangeable.

Collecting one explained tracer for each dimension could, in principle, reduce item-level responses to 11 construct-level ratings while retaining distinctions between momentary peaks and cumulative experience. This may make assessment faster and more engaging, but the present study did not directly compare completion time, burden, enjoyment, reliability, or validity. Whether explanatory tracer prompts can replace or complement the 11D-ASC therefore requires dedicated validation, including dependence-aware correspondence analyses and tests of whether peak, AUC, or finer temporal features explain global retrospective ratings.




Fig. S2 Condition-wise altered-state profiles across all 11 dimensions for retrospective 11D-ASC questionnaire ratings (top; 0–100 visual-analogue scale), tracer peak intensity (middle; normalized 0–1 intensity), and tracer AUC (bottom; normalized 0–1 area). Lines show condition means for the included sample ($N = 39$). Separate radial scales are used; comparisons should concern profile shape and condition separation rather than absolute height across measurement formats.

526 S5 Reproducibility and file access

527
528 The preregistration record, deidentified participant-level data, inclusion indicator,
529 statistical outputs, analysis scripts, figure-generation scripts, study materials, environ-
530 ment specifications, and replay instructions are available at [https://doi.org/10.17605/](https://doi.org/10.17605/OSF.IO/64MWK)
531 [OSF.IO/64MWK](https://doi.org/10.17605/OSF.IO/64MWK).

532 The exact GitHub repository version corresponding to the submitted manuscript
533 is available at:

534
535 [https://github.com/GeorgeFejer91/0-OSF_reproducability_package/tree/](https://github.com/GeorgeFejer91/0-OSF_reproducability_package/tree/FINAL_SUBMISSION_COMMIT)
536 [FINAL_SUBMISSION_COMMIT](https://github.com/GeorgeFejer91/0-OSF_reproducability_package/tree/FINAL_SUBMISSION_COMMIT)

537
538 The repository’s Start Here documentation identifies the analysis entry points,
539 output directories, environment files, and replay sequence.

541 References

542
543 Studerus, E., Gamma, A., Vollenweider, F.X.: Psychometric evaluation of the altered
544 states of consciousness rating scale (oav). PLOS ONE **5**(8), 12412 (2010) [https:](https://doi.org/10.1371/journal.pone.0012412)
545 [//doi.org/10.1371/journal.pone.0012412](https://doi.org/10.1371/journal.pone.0012412)

546
547 Jachs, B.: The neurophenomenology of meditative states: Introducing temporal expe-
548 rience tracing to capture subjective experience states and their neural correlates.
549 PhD thesis, University of Cambridge (2021). <https://doi.org/10.17863/CAM.80153>
550 . <https://www.repository.cam.ac.uk/handle/1810/332709>

551
552 Damci, A., Hoeijmakers, J., Hollander, M., Köke, A., De Mooij, M., Faber, C., Verbunt,
553 J.: Acceptability, usability and feasibility of experience sampling method in chronic
554 secondary pain syndromes. Frontiers in Neurology **14** (2023) [https://doi.org/10.](https://doi.org/10.3389/fneur.2023.1219236)
555 [3389/fneur.2023.1219236](https://doi.org/10.3389/fneur.2023.1219236)

556
557 Lewis-Healey, E., Tagliazucchi, E., Canales-Johnson, A., Bekinschtein, T.A.:
558 Breathwork-induced psychedelic experiences modulate neural dynamics. Cerebral
559 Cortex **34**(8), 347 (2024) <https://doi.org/10.1093/cercor/bhae347>

560
561
562
563
564
565
566
567
568
569
570
571