

- (c) specific latent heat of fusion and specific latent heat of vaporisation; $E = mL$
- (d) (i) an electrical experiment to determine the specific latent heat of fusion and vaporisation
- (ii) techniques and procedures used for an electrical method to determine the specific latent heat of a solid and a liquid.

- (6) M - Define the terms: specific latent heat of fusion and specific latent heat of vaporisation
- (7) S - Select and apply the equation for specific latent heat
- (8) C - Describe an electrical experiment to determine the specific latent heat of fusion and vaporisation

Specific latent heat



STARTER:

Peer mark the HWK.

Use the **practical guidance booklet** for more information.

WWW/EBI SR in response

	Expected Answers	Mark	Additional Guidance
A1.1	Sets up the experiment safely and correctly without help. ✓ Determines the temperature of the water and 100g mass without help. ✓	2	Any help given should be recorded on the front cover of the task sheet. Any help given should be recorded on the front cover of the task sheet.
A1.2	Calculates a value for E correctly. ✓ Repeats the experiment for one further value of starting temperature. ✓	2	Penalise power of 10 error / inconsistent unit. Any help given should be recorded on the front cover of the task sheet. θ_M must be in the range 60 °C to 70 °C
A1.3	Identifies one hazard and describes appropriate safety precaution. ✓	1	e.g. avoid splashing as the mass is inserted and so put in beaker slowly, or breaking the beaker and so put in beaker slowly/ carefully. Do not allow 'use goggles' / 'use lab coats' / 'tie hair back' / 'string' / '(oven) gloves'.
B1.1	Records all data required in a suitable table. ✓ Correctly determines both values for s.h.c. ✓	2	The table must include the following quantities: θ_M , θ_2 , $(\theta_M - \theta_2)$, $(\theta_2 - \theta_1)$, E . Ignore units, sig. fig. and d.p. One value should be centre value $\pm 50\%$
B1.2	Discusses with appropriate justification the value of the s.h.c. of the metal block. Supports the observations by sound reasoning in terms of the energy losses. ✓✓✓	3	Credit the following marking points and ignore reference to one of the s.h.c. values being greater/smaller than the other. The s.h.c. values are different because the 100g mass: 1. loses more energy when transferred from the higher temperature hot water; 2. also warms up the beaker/container (and the cold water); 3. may not have reached temperature θ_M /thermal equilibrium. 4. Allow one other detailed correct statement that supports the observations. Do not allow loses energy (to surroundings) when transferred from hot water to cold water.
	Total	10	

	Expected Answers	Mark	Additional Guidance
A1.1	Sets up the experiment safely and correctly without help. ✓ Determines the temperature of the water and 100g mass without help. ✓	2	Any help given should be recorded on the front cover of the task sheet. Any help given should be recorded on the front cover of the task sheet.
A1.2	Calculates a value for E correctly. ✓ Repeats the experiment for one further value of starting temperature. ✓	2	Penalise power of 10 error / inconsistent unit. Any help given should be recorded on the front cover of the task sheet. θ_M must be in the range 60 °C to 70 °C
A1.3	Identifies one hazard and describes appropriate safety precaution. ✓	1	e.g. avoid splashing as the mass is inserted and so put in beaker slowly, or breaking the beaker and so put in beaker slowly/ carefully. Do not allow 'use goggles' / 'use lab coats' / 'tie hair back' / 'string' / '(oven) gloves'.
B1.1	Records all data required in a suitable table. ✓ Correctly determines both values for s.h.c. ✓	2	The table must include the following quantities: θ_M , θ_2 , $(\theta_M - \theta_2)$, $(\theta_2 - \theta_1)$, E . Ignore units, sig. fig. and d.p. One value should be centre value $\pm 50\%$
B1.2	Discusses with appropriate justification the value of the s.h.c. of the metal block. Supports the observations by sound reasoning in terms of the energy losses. ✓✓✓	3	Credit the following marking points and ignore reference to one of the s.h.c. values being greater/smaller than the other. The s.h.c. values are different because the 100g mass: 1. loses more energy when transferred from the higher temperature hot water; 2. also warms up the beaker/container (and the cold water); 3. may not have reached temperature θ_M /thermal equilibrium. 4. Allow one other detailed correct statement that supports the observations. Do not allow loses energy (to surroundings) when transferred from hot water to cold water.
	Total	10	

- (6) M - Define the terms: specific latent heat of fusion and specific latent heat of vaporisation
 (7) S - Select and apply the equation for specific latent heat
 (8) C - Describe an electrical experiment to determine the specific latent heat of fusion and vaporisation

Specific latent heat

Specific latent heat, L is

defined as the energy requires to change the phase per unit mass - while at constant temperature.




Key point



$$L = \frac{E}{m} \rightarrow \frac{J}{kg}$$




Key point



When a substance changes from a solid to a liquid it is called: **Specific latent heat of fusion**

When a substance changes from a liquid to a gas it is called: **Specific latent heat of vaporisation**

- (6) M - Define the terms: specific latent heat of fusion and specific latent heat of vaporisation
(7) S - Select and apply the equation for specific latent heat
(8) C - Describe an electrical experiment to determine the specific latent heat of fusion and vaporisation



Activity: Complete summary Q1-6

Q4 / Q6

Kilo 10^3

Mega 10^6 Start at 3

Giga 10^9 Start at 5




Key
point

Oxford A Level Sciences
OCR Physics A

14.5 Specific latent heat Stretch and challenge

An old principle saves energy

- (6) M - Define the terms: specific latent heat of fusion and specific latent heat of vaporisation
 (7) S - Select and apply the equation for specific latent heat
 (8) C - Describe an electrical experiment to determine the specific latent heat of fusion and vaporisation



Mini: plenary:

1. What is the definition for **specific latent heat of fusion**....?
2. The SLH of F for water is $3.3 \times 10^5 \text{ Jkg}^{-1}$. A 25g ice cube is left on a desk. The flow of heat energy to the ice is equivalent to 5W. How long until the ice cube melts?

Kilo 10^3

Mega 10^6

Giga 10^9

$$E = mL = 0.025 \times 3.3 \times 10^5 = 8250 \text{ J}$$

Why is this time likely to be inaccurate?

$$E = Pt =$$



$$t = \frac{E}{P} = \frac{8250}{5} = 1650 \text{ s}$$

- (6) M - Define the terms: specific latent heat of fusion and specific latent heat of vaporisation
 (7) S - Select and apply the equation for specific latent heat
 (8) C - Describe an electrical experiment to determine the specific latent heat of fusion and vaporisation

Measuring specific latent heat - demonstration

1. For both demonstrations, complete the method write up sheet.
2. Find and describe a more accurate method to measure the mass of evaporated water for practical 2.
3. Find similarities between the methods and highlighting certain parts of your text.

Kilo 10^3

Mega 10^6

Giga 10^9

Why is this time likely to be inaccurate?



Key
point

Specific latent heat of fusion	Specific latent heat of vaporisation
Labelled diagram of setup	Labelled diagram of setup
Method – basic procedure (the measurements)	Method – basic procedure (the measurements)
Equations and calculations	Equations and calculations
Uncertainties and solutions	Uncertainties and solutions

1.2×10^6

6.02×10^5


Substance	Specific latent heat of fusion (L_f) (J/kg)	Melting point ($^{\circ}\text{C}$)	Specific latent heat of vaporization (L_v) (J/kg)	Boiling point ($^{\circ}\text{C}$)
aluminum	6.6×10^5	2519	4.0×10^5	10 900
ethyl alcohol	1.1×10^5	-114	8.6×10^5	78.3
carbon dioxide	1.8×10^5	-78	5.7×10^5	-57
gold	1.1×10^5	2856	6.4×10^4	1 645
lead	2.5×10^4	327.5	8.7×10^5	1 750
water	3.4×10^5	0	2.3×10^6	100

% difference

- (6) M - Define the terms: specific latent heat of fusion and specific latent heat of vaporisation
 (7) S - Select and apply the equation for specific latent heat
 (8) C - Describe an electrical experiment to determine the specific latent heat of fusion and vaporisation

Plenary

The diagram below shows the variation of the temperature of 200 g of lead as it is heated at a steady rate.



- Use the graph to state the melting point of lead. [1]
- Explain why the graph is a straight line at the start. [1]
- Explain what happens to the energy supplied to the lead as it melts at a constant temperature. [1]
- The initial temperature of the lead is 0 °C. Use the graph to determine the total energy supplied to the lead before it starts to melts. (The specific heat capacity of lead is 130 J kg⁻¹ K⁻¹.) [3]
- Use your answer to **d** to determine the rate of heating of the lead. [2]

Kilo 10³

Mega 10⁶

Giga 10⁹

Calculate the specific latent heat of fusion of lead.

- (6) M - Define the terms: specific latent heat of fusion and specific latent heat of vaporisation
 (7) S - Select and apply the equation for specific latent heat
 (8) C - Describe an electrical experiment to determine the specific latent heat of fusion and vaporisation

Plenary - MS



- a Melting point = 600 °C [1]
 (There is no change in temperature during change of state.)
- b The lead is being heated at a steady rate and therefore the temperature also increases at a steady rate. [1]
- c The energy supplied to the lead is used to break the atomic bonds and increase the separation between the atoms of lead (and hence their potential energy increases). [1]
- d $E = mc\Delta\theta$ [1]
 $E = 200 \times 10^{-3} \times 130 \times (600 - 0)$ [1]
 $E = 1.56 \times 10^4 \text{ J} \approx 1.6 \times 10^4 \text{ J}$ [1]
- e In a time of 300 s, $1.56 \times 10^4 \text{ J}$ of energy is supplied to the lead.
- Rate of heating = power
- $\text{power} = \frac{1.56 \times 10^4}{300}$ [1]
 power = 52 W [1]

next lesson - melting ice cube to find SLH.

combining SHC and SLH