

Shear Mapping in Python (SMPy): Modular, Extensible, and Accessible Dark Matter Mapping

Georgios N. Vassilakis¹, Jacqueline E. McCleary¹, Maya Amit¹, and Sayan Saha¹

¹ Department of Physics, Northeastern University, USA ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Understanding the universe's large-scale distribution of dark matter is a central objective in the era of precision cosmology. A key tool for the study of dark matter is weak gravitational lensing; a phenomenon where light from distant galaxies is sheared as it passes through the gravitational field of a massive object, like a galaxy cluster. This shear, which manifests as a slight (weak) distortion of shapes over thousands of galaxies, allows astrophysicists to infer the distribution of total matter, including both luminous and dark matter.

Obtaining a mass distribution from a catalog of galaxy shears requires an intermediate step. A common tool for this step is the mapping of convergence (κ), which quantifies how much a gravitational lens converges the light from distant galaxies, resulting in a magnification of their shapes. This value is directly proportional to the projected mass density, enabling easy visualization of the overall distribution. For a comprehensive review of weak gravitational lensing, please refer to ([Umetsu, 2020](#)).

The **Shear Mapping in Python (SMPy)** package provides a standardized, well-documented, and open-source solution for creating convergence maps from weak lensing galaxy shear measurements. SMPy was initially developed to support the Superpressure Balloon-borne Imaging Telescope (SuperBIT), a stratospheric, near-UV to near-IR observing platform which completed its 45-night observing run in spring 2023 with over 30 galaxy cluster observations ([Gill et al., 2024](#)). SMPy has since evolved into a general-purpose tool suitable for analyzing the weak lensing data from any source of galaxies.

Statement of Need

Mass maps are a critical deliverable of many cosmological analyses [ACTDR62024], ([Jeffrey et al., 2021](#)), ([Oguri et al., 2017](#)). While some tools exist for mass mapping, such as `lenspack` and `jax-lensing` ([Remy et al., 2022](#)), they are either not well documented and lack algorithmic rigor, or they are highly specialized for specific use cases, such as Neural Score Matching to create Deep Learning-based maps. This leaves a significant gap in the field for a user-friendly, well-documented tool that can handle both astrometric and pixel-space convergence mapping while maintaining mathematical rigor.

SMPy fills this void by providing a robust, well-documented, and open-source tool to construct publication-quality mass maps from galaxy shear data. It is, to our knowledge, the first convergence mapping software that combines accessibility with the flexibility to compute convergence in either astrometric or pixel space.

SMPy was built with the following design principles in mind:

1. **Accessibility:** SMPy is written entirely in Python and deliberately relies only on widely-used

scientific Python packages (numpy, scipy, pandas, astropy, matplotlib, and pyyaml). This choice of standard dependencies ensures that users can easily install the packages without complex dependency chains, and that the codebase is maintainable and familiar to the scientific Python community.

2. **Extensibility:** SMPy is built with a modular architecture that allows for easy implementation of new mass mapping techniques beyond the currently implemented Kaiser-Squires inversion algorithm (Kaiser & Squires, 1993). An example convergence map is shown in Figure 1, created from simulated galaxy cluster observations from SuperBIT (McCleary et al., 2023). Aperture mass mapping (Leonard et al., 2012) and KS+ (Pires, 2020) algorithms are currently planned to be added to the codebase.
3. **Usability:** Creating convergence maps with SMPy requires minimal input - users need only provide a catalog of galaxies with their associated shears (g_1 & g_2) and coordinates. This straightforward input requirement makes the tool accessible to researchers at all levels. A flexible configuration system is integrated via a single YAML file that defines file paths, convergence map algorithm settings, plotting parameters, and more.
4. **Robustness:** Designed to be mathematically and algorithmically accurate, allowing the user to create convergence maps with any galaxy shear data. The coordinate system abstraction handles both RA/Dec celestial coordinates (with proper spherical geometry approximations) or pixel-based coordinates through a unified interface. Signal-to-noise maps can be generated using either spatial shuffling (randomizing galaxy positions while preserving shear values) or orientation shuffling (randomizing shear orientations while preserving positions) to distinguish real signals from noise.




Figure 1: Example convergence map created with SMPy showing the mass distribution of a simulated galaxy cluster. The map was generated using the Kaiser-Squires inversion method on simulated weak lensing data from SuperBIT. The color scale represents the dimensionless surface mass density (convergence), with brighter regions indicating higher mass concentrations.

Software References

SMPy is written in Python 3.8+ and uses the following packages:

- NumPy (Harris et al., 2020)
- SciPy (Virtanen et al., 2020)
- Pandas (team, 2024)

- 67 ▪ Astropy (Astropy Collaboration et al., 2022) (Astropy Collaboration et al., 2018) (Astropy
- 68 Collaboration et al., 2013)
- 69 ▪ Matplotlib (Hunter, 2007)
- 70 ▪ PyYAML (Simonov, 2024)

71 Acknowledgements

72 This material is based upon work supported by a Northeastern University Undergraduate
73 Research and Fellowships PEAK Summit Award.

74 Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., Earl, N., Starkman, N., Bradley, L.,
75 Shupe, D. L., Patil, A. A., Corrales, L., Brasseur, C. E., Nöthe, M., Donath, A., Tollerud,
76 E., Morris, B. M., Ginsburg, A., Vaher, E., Weaver, B. A., Tocknell, J., Jamieson, W., ...
77 Astropy Project Contributors. (2022). The Astropy Project: Sustaining and Growing a
78 Community-oriented Open-source Project and the Latest Major Release (v5.0) of the Core
79 Package. *935*(2), 167. <https://doi.org/10.3847/1538-4357/ac7c74>

80 Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., Lim, P. L.,
81 Crawford, S. M., Conseil, S., Shupe, D. L., Craig, M. W., Dencheva, N., Ginsburg, A.,
82 VanderPlas, J. T., Bradley, L. D., Pérez-Suárez, D., de Val-Borro, M., Aldcroft, T. L.,
83 Cruz, K. L., Robitaille, T. P., Tollerud, E. J., ... Astropy Contributors. (2018). The Astropy
84 Project: Building an Open-science Project and Status of the v2.0 Core Package. *156*(3),
85 123. <https://doi.org/10.3847/1538-3881/aabc4f>

86 Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., Greenfield, P., Droettboom, M., Bray,
87 E., Aldcroft, T., Davis, M., Ginsburg, A., Price-Whelan, A. M., Kerzendorf, W. E., Conley,
88 A., Crighton, N., Barbary, K., Muna, D., Ferguson, H., Grollier, F., Parikh, M. M., Nair, P.
89 H., ... Streicher, O. (2013). Astropy: A community Python package for astronomy. *558*.

90 Gill, A. S., Benton, S. J., Damaren, C. J., Everett, S. W., Fraisse, A. A., Hartley, J. W.,
91 Harvey, D., Holder, B., Huff, E. M., Jauzac, M., Jones, W. C., Lagattuta, D., Leung, J.
92 S.-Y., Li, L., Luu, T. V. T., Massey, R., McCleary, J. E., Nagy, J. M., Netterfield, C. B., ...
93 Vitorelli, A. Z. (2024). SuperBIT superpressure flight instrument overview and performance:
94 Near-diffraction-limited astronomical imaging from the stratosphere. *The Astronomical*
95 *Journal*, *168*(2), 85. <https://doi.org/10.3847/1538-3881/ad5840>

96 Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D.,
97 Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk,
98 M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant,
99 T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>

101 Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science &*
102 *Engineering*, *9*(3), 90–95. <https://doi.org/10.1109/MCSE.2007.55>

103 Jeffrey, N., Gatti, M., Chang, C., Whiteway, L., Demirbozan, U., Kovacs, A., Pollina, G.,
104 Bacon, D., Hamaus, N., Kacprzak, T., Lahav, O., Lanusse, F., Mawdsley, B., Nadathur, S.,
105 Starck, J. L., Vielzeuf, P., Zeurcher, D., Alarcon, A., Amon, A., ... Collaboration, D. (2021).
106 Dark energy survey year 3 results: Curved-sky weak lensing mass map reconstruction.
107 *Monthly Notices of the Royal Astronomical Society*, *505*(3), 4626–4645. <https://doi.org/10.1093/mnras/stab1495>

109 Kaiser, N., & Squires, G. (1993). Mapping the dark matter with weak gravitational lensing.
110 *404*, 441–450. <https://doi.org/10.1086/172297>

111 Leonard, A., Pires, S., & Starck, J.-L. (2012). Fast calculation of the weak lensing aperture
112 mass statistic. *Monthly Notices of the Royal Astronomical Society*, *423*(4), 3405–3412.
113 <https://doi.org/10.1111/j.1365-2966.2012.21133.x>

- 114 McCleary, J. E., Everett, S. W., Shaaban, M. M., Gill, A. S., Vassilakis, G. N., Huff, E.
115 M., Massey, R. J., Benton, S. J., Brown, A. M., Clark, P., Holder, B., Fraisse, A.
116 A., Jauzac, M., Jones, W. C., Lagattuta, D., Leung, J. S.-Y., Li, L., Luu, T. V. T.,
117 Nagy, J. M., ... Tam, S. I. (2023). Lensing in the blue. II. Estimating the sensitivity of
118 stratospheric balloons to weak gravitational lensing. *The Astronomical Journal*, 166(3),
119 134. <https://doi.org/10.3847/1538-3881/ace7ca>
- 120 Oguri, M., Miyazaki, S., Hikage, C., Mandelbaum, R., Utsumi, Y., Miyatake, H., Takada,
121 M., Armstrong, R., Bosch, J., Komiyama, Y., Leauthaud, A., More, S., Nishizawa, A. J.,
122 Okabe, N., & Tanaka, M. (2017). Two- and three-dimensional wide-field weak lensing
123 mass maps from the hyper supprime-cam subaru strategic program S16A data. *Publications*
124 *of the Astronomical Society of Japan*, 70(SP1), S26. <https://doi.org/10.1093/pasj/psx070>
- 125 Pires, S. (2020). Euclid: Reconstruction of weak-lensing mass maps for non-gaussianity studies.
126 *Astronomy & Astrophysics*, 638, A141. <https://doi.org/10.1051/0004-6361/201936865>
- 127 Remy, B., Lanusse, F., Jeffrey, N., Liu, J., Starck, J.-L., Osato, K., & Schrabback, T. (2022).
128 Probabilistic mass-mapping with neural score estimation. *Astronomy & Astrophysics*, 672.
129 <https://doi.org/10.1051/0004-6361/202243054>
- 130 Simonov, K. (2024). PyYAML. <https://pyyaml.org/>
- 131 team, T. pandas development. (2024). *Pandas-dev/pandas: pandas* (Version v2.2.2). Zenodo.
132 <https://doi.org/10.5281/zenodo.10957263>
- 133 Umetsu, K. (2020). Cluster-galaxy weak lensing. *The Astronomy and Astrophysics Review*,
134 28(1), 106. <https://doi.org/10.1007/s00159-020-00129-w>
- 135 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
136 Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson,
137 J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., ... SciPy
138 1.0 Contributors. (2020). SciPy 1.0: Fundamental Algorithms for Scientific Computing in
139 Python. *Nature Methods*, 17, 261–272. <https://doi.org/10.1038/s41592-019-0686-2>