

Shear Mapping in Python (SMPy): Modular, Extensible, and Accessible Dark Matter Mapping

Georgios N. Vassilakis¹, Jacqueline E. McCleary¹, Maya Amit¹, and Sayan Saha¹

¹ Department of Physics, Northeastern University, Boston, MA, USA ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Understanding the universe's large-scale distribution of dark matter is a central objective in the era of precision cosmology. A key technique for the study of dark matter is weak gravitational lensing: a phenomenon where light from distant galaxies is sheared as it passes through the gravitational field of a massive object, like a galaxy cluster. This shear, which manifests as a slight (weak) distortion of shapes over thousands of galaxies, allows astrophysicists to infer the distribution of total matter, including both luminous and dark matter.

Obtaining a mass distribution from a catalog of galaxy shears requires an intermediate step. A common tool for this step is the mapping of convergence (κ), which quantifies how much a gravitational lens converges the light from distant galaxies, resulting in a magnification of their shapes. This value is directly proportional to the projected mass density, enabling easy visualization of the overall mass distribution. For a comprehensive review of weak gravitational lensing refer to ([Umetsu, 2020](#)).

The **Shear Mapping in Python (SMPy)** package provides a standardized, well-documented, and open-source solution for creating convergence maps from weak lensing galaxy shear measurements. SMPy was initially developed to support the Superpressure Balloon-borne Imaging Telescope (SuperBIT), a stratospheric, near-UV to near-IR observing platform which completed its 45-night observing run in spring 2023 with over 30 galaxy cluster observations ([Gill et al., 2024](#)), ([Sirks et al., 2023](#)). SMPy has since evolved into a general-purpose tool suitable for analyzing the weak lensing data from cosmological surveys.

Statement of Need

While mass maps are a key deliverable of many cosmological analyses ([Madhavacheril et al., 2024](#)) ([Jeffrey et al., 2021](#)) ([Oguri et al., 2017](#)), scientists are often left to make these maps from scratch. The weak lensing community is served by publicly available mapping tools like lenspack and jax-lensing ([Remy et al., 2022](#)), each with their own strengths. jax-lensing excels at neural network-based approaches and deep learning methods, while lenspack has a well-documented module with stand-alone mass-mapping functions. While both tools are powerful, the steep learning curve of jax-lensing and the rigid architecture of lenspack motivated the development of SMPy as an accessible and extensible alternative.

SMPy addresses an outstanding need for the lensing community: an accessible, well-documented, and extensible tool to construct publication-quality mass maps from galaxy shear data. Built on standard scientific Python packages, it provides an easy entry point for researchers new to mass mapping, while also being robust for more senior scientific use. There are currently three separate mapping methods implemented into SMPy: the classic Kaiser-Squires inversion ([Kaiser & Squires, 1993](#)), aperture mass mapping ([Leonard et al., 2012](#)) ([J. McCleary et al.,](#)

2020), and notably, to our knowledge, the first publicly available implementation of the KS+ algorithm (Pires, 2020). KS+ improves reconstruction quality by correcting for systematic effects including missing data, field borders, and reduced shear. SMPy also offers specialized and unique features valuable for mass mapping, such as flexible coordinate system support (both celestial and pixel space) and comprehensive signal-to-noise analysis with multiple noise randomization techniques. An example convergence map, created from simulated SuperBIT galaxy cluster observations (J. E. McCleary et al., 2023), is shown in Figure 1. SMPy is, to our knowledge, the first convergence mapping software to prioritize both accessibility and advanced features.

50 Software Features

51 SMPy was built with the following design principles in mind:

- 52 1. **Accessibility:** SMPy is written entirely in Python and deliberately relies only on widely-used
53 scientific Python packages (numpy, scipy, pandas, astropy, matplotlib, and pyyaml).
54 This choice of standard dependencies ensures that users can easily install the packages
55 without complex dependency chains, and that the codebase is maintainable and familiar
56 to the scientific Python community.
- 57 2. **Extensibility:** SMPy is built with a modular architecture that facilitates easy implemen-
58 tation of new mass mapping techniques, encouraging open-source contribution. The
59 framework's standardized abstract base class architecture allows for the integration of
60 different mapping methods with minimal effort.
- 61 3. **Usability:** Creating convergence maps with SMPy requires minimal input—users need to
62 only provide a catalog of galaxies with their associated shear components and coordinates.
63 This straightforward input requirement makes the tool accessible to researchers at all
64 levels. A flexible configuration system is integrated via a single YAML file that defines
65 file paths, convergence map algorithm settings, plotting parameters, and more. With
66 this configuration file, the user can create convergence & SNR maps with one line, either
67 via terminal or within code.
- 68 4. **Robustness:** SMPy is designed to be mathematically and algorithmically accurate, allowing
69 the user to create convergence maps with any galaxy shear data. The coordinate
70 system abstraction handles both celestial coordinates (with proper spherical geometry
71 approximations) or pixel-based coordinates through a unified interface. To quantify the
72 significance of the weak lensing detection, multiple noise realizations can be generated
73 using either spatial shuffling (randomizing galaxy positions while preserving shear values)
74 or orientation shuffling (randomizing shear orientations while preserving positions). These
75 noise realizations are used to create a signal-to-noise map with the observed convergence.




Figure 1: Example convergence map created with SMPy showing the mass distribution of a simulated galaxy cluster. The map was generated using the Kaiser-Squires inversion method on simulated weak lensing data from SuperBIT. The color scale represents the dimensionless surface mass density (convergence), with brighter regions indicating higher mass concentrations.

Software References

SMPy is written in Python 3.8+ and uses the following packages:

- NumPy ([Harris et al., 2020](#))
- SciPy ([Virtanen et al., 2020](#))
- Pandas ([team, 2024](#))
- Astropy ([Astropy Collaboration et al., 2022](#)) ([Astropy Collaboration et al., 2018](#)) ([Astropy Collaboration et al., 2013](#))
- Matplotlib ([Hunter, 2007](#))
- PyYAML ([Simonov, 2024](#))

Acknowledgements

This material is based upon work supported by a Northeastern University Undergraduate Research and Fellowships PEAK Summit Award.

References

- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., Earl, N., Starkman, N., Bradley, L., Shupe, D. L., Patil, A. A., Corrales, L., Brasseur, C. E., Nöthe, M., Donath, A., Tollerud, E., Morris, B. M., Ginsburg, A., Vaher, E., Weaver, B. A., Tocknell, J., Jamieson, W., ... Astropy Project Contributors. (2022). The Astropy Project: Sustaining and Growing a Community-oriented Open-source Project and the Latest Major Release (v5.0) of the Core Package. *935*(2), 167. <https://doi.org/10.3847/1538-4357/ac7c74>
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., Lim, P. L., Crawford, S. M., Conseil, S., Shupe, D. L., Craig, M. W., Dencheva, N., Ginsburg, A., VanderPlas, J. T., Bradley, L. D., Pérez-Suárez, D., de Val-Borro, M., Aldcroft, T. L., Cruz, K. L., Robitaille, T. P., Tollerud, E. J., ... Astropy Contributors. (2018). The Astropy Project: Building an Open-science Project and Status of the v2.0 Core Package. *156*(3), 123. <https://doi.org/10.3847/1538-3881/aabc4f>

- 101 Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., Greenfield, P., Droettboom, M., Bray,
102 E., Aldcroft, T., Davis, M., Ginsburg, A., Price-Whelan, A. M., Kerzendorf, W. E., Conley,
103 A., Crighton, N., Barbary, K., Muna, D., Ferguson, H., Grollier, F., Parikh, M. M., Nair, P.
104 H., ... Streicher, O. (2013). Astropy: A community Python package for astronomy. 558.
- 105 Gill, A. S., Benton, S. J., Damaren, C. J., Everett, S. W., Fraisse, A. A., Hartley, J. W.,
106 Harvey, D., Holder, B., Huff, E. M., Jauzac, M., Jones, W. C., Lagattuta, D., Leung, J.
107 S.-Y., Li, L., Luu, T. V. T., Massey, R., McCleary, J. E., Nagy, J. M., Netterfield, C. B., ...
108 Vitorelli, A. Z. (2024). SuperBIT superpressure flight instrument overview and performance:
109 Near-diffraction-limited astronomical imaging from the stratosphere. *The Astronomical*
110 *Journal*, 168(2), 85. <https://doi.org/10.3847/1538-3881/ad5840>
- 111 Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D.,
112 Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk,
113 M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant,
114 T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- 115
- 116 Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science &*
117 *Engineering*, 9(3), 90–95. <https://doi.org/10.1109/MCSE.2007.55>
- 118 Jeffrey, N., Gatti, M., Chang, C., Whiteway, L., Demirbozan, U., Kovacs, A., Pollina, G.,
119 Bacon, D., Hamaus, N., Kacprzak, T., Lahav, O., Lanusse, F., Mawdsley, B., Nadathur, S.,
120 Starck, J. L., Vielzeuf, P., Zeurcher, D., Alarcon, A., Amon, A., ... Collaboration, D. (2021).
121 Dark energy survey year 3 results: Curved-sky weak lensing mass map reconstruction.
122 *Monthly Notices of the Royal Astronomical Society*, 505(3), 4626–4645. <https://doi.org/10.1093/mnras/stab1495>
- 123
- 124 Kaiser, N., & Squires, G. (1993). Mapping the dark matter with weak gravitational lensing.
125 404, 441–450. <https://doi.org/10.1086/172297>
- 126 Leonard, A., Pires, S., & Starck, J.-L. (2012). Fast calculation of the weak lensing aperture
127 mass statistic. *Monthly Notices of the Royal Astronomical Society*, 423(4), 3405–3412.
128 <https://doi.org/10.1111/j.1365-2966.2012.21133.x>
- 129 Madhavacheril, M. S., Qu, F. J., Sherwin, B. D., MacCrann, N., Li, Y., Abril-Cabezas, I., Ade,
130 P. A. R., Aiola, S., Alford, T., Amiri, M., Amodeo, S., An, R., Atkins, Z., Austermann, J. E.,
131 Battaglia, N., Battistelli, E. S., Beall, J. A., Bean, R., Beringue, B., ... Zheng, K. (2024). The
132 atacama cosmology telescope: DR6 gravitational lensing map and cosmological parameters.
133 *The Astrophysical Journal*, 962(2), 113. <https://doi.org/10.3847/1538-4357/acff5f>
- 134 McCleary, J. E., Everett, S. W., Shaaban, M. M., Gill, A. S., Vassilakis, G. N., Huff, E.
135 M., Massey, R. J., Benton, S. J., Brown, A. M., Clark, P., Holder, B., Fraisse, A.
136 A., Jauzac, M., Jones, W. C., Lagattuta, D., Leung, J. S.-Y., Li, L., Luu, T. V. T.,
137 Nagy, J. M., ... Tam, S. I. (2023). Lensing in the blue. II. Estimating the sensitivity of
138 stratospheric balloons to weak gravitational lensing. *The Astronomical Journal*, 166(3),
139 134. <https://doi.org/10.3847/1538-3881/ace7ca>
- 140 McCleary, J., dell'Antonio, I., & Linden, A. von der. (2020). Dark matter distribution of four
141 low-z clusters of galaxies. *The Astrophysical Journal*, 893(1), 8. <https://doi.org/10.3847/1538-4357/ab7c58>
- 142
- 143 Oguri, M., Miyazaki, S., Hikage, C., Mandelbaum, R., Utsumi, Y., Miyatake, H., Takada,
144 M., Armstrong, R., Bosch, J., Komiyama, Y., Leauthaud, A., More, S., Nishizawa, A. J.,
145 Okabe, N., & Tanaka, M. (2017). Two- and three-dimensional wide-field weak lensing
146 mass maps from the hyper supprime-cam subaru strategic program S16A data. *Publications*
147 *of the Astronomical Society of Japan*, 70(SP1), S26. <https://doi.org/10.1093/pasj/psx070>
- 148 Pires, S. (2020). Euclid: Reconstruction of weak-lensing mass maps for non-gaussianity studies.
149 *Astronomy & Astrophysics*, 638, A141. <https://doi.org/10.1051/0004-6361/201936865>

- 150 Remy, B., Lanusse, F., Jeffrey, N., Liu, J., Starck, J.-L., Osato, K., & Schrabback, T. (2022).
151 Probabilistic mass-mapping with neural score estimation. *Astronomy & Astrophysics*, 672.
152 <https://doi.org/10.1051/0004-6361/202243054>
- 153 Simonov, K. (2024). *PyYAML*. <https://pyyaml.org/>
- 154 Sirks, E. L., Massey, R., Gill, A. S., Anderson, J., Benton, S. J., Brown, A. M., Clark, P.,
155 English, J., Everett, S. W., Fraisse, A. A., Franco, H., Hartley, J. W., Harvey, D., Holder,
156 B., Hunter, A., Huff, E. M., Hynous, A., Jauzac, M., Jones, W. C., ... Vassilakis, G. N.
157 (2023). Data downloaded via parachute from a NASA super-pressure balloon. *Aerospace*,
158 10(11). <https://doi.org/10.3390/aerospace10110960>
- 159 team, T. pandas development. (2024). *Pandas-dev/pandas: pandas* (Version v2.2.2). Zenodo.
160 <https://doi.org/10.5281/zenodo.10957263>
- 161 Umetsu, K. (2020). Cluster-galaxy weak lensing. *The Astronomy and Astrophysics Review*,
162 28(1), 106. <https://doi.org/10.1007/s00159-020-00129-w>
- 163 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
164 Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson,
165 J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., ... SciPy
166 1.0 Contributors. (2020). SciPy 1.0: Fundamental Algorithms for Scientific Computing in
167 Python. *Nature Methods*, 17, 261–272. <https://doi.org/10.1038/s41592-019-0686-2>