Approximation of second-order derivatives

Central difference scheme

$$T_1 + T_2 \Rightarrow \left(\frac{\partial^2 u}{\partial x^2}\right)_i = \frac{u_{i+1} - 2u_i + u_{i-1}}{(\Delta x)^2} + \mathcal{O}(\Delta x)^2$$

Alternative derivation

$$\left(\frac{\partial^2 u}{\partial x^2}\right)_i = \left[\frac{\partial}{\partial x} \left(\frac{\partial u}{\partial x}\right)\right]_i = \lim_{\Delta x \to 0} \frac{\left(\frac{\partial u}{\partial x}\right)_{i+1/2} - \left(\frac{\partial u}{\partial x}\right)_{i-1/2}}{\Delta x}$$

$$\approx \frac{\frac{u_{i+1} - u_i}{\Delta x} - \frac{u_i - u_{i-1}}{\Delta x}}{\Delta x} = \frac{u_{i+1} - 2u_i + u_{i-1}}{(\Delta x)^2}$$

Variable coefficients
$$f(x) = d(x) \frac{\partial u}{\partial x}$$
 diffusive flux

$$\left(\frac{\partial f}{\partial x}\right)_{i} \approx \frac{f_{i+1/2} - f_{i-1/2}}{\Delta x} = \frac{d_{i+1/2} \frac{u_{i+1} - u_{i}}{\Delta x} - d_{i-1/2} \frac{u_{i} - u_{i-1}}{\Delta x}}{\Delta x}$$

$$= \frac{d_{i+1/2} u_{i+1} - (d_{i+1/2} + d_{i-1/2}) u_{i} + d_{i-1/2} u_{i-1}}{(\Delta x)^{2}}$$


Approximation of mixed derivatives

2D:
$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial y} \right) = \frac{\partial}{\partial y} \left(\frac{\partial u}{\partial x} \right)$$

$$\left(\frac{\partial^2 u}{\partial x \partial y}\right)_{i,j} = \frac{\left(\frac{\partial u}{\partial y}\right)_{i+1,j} - \left(\frac{\partial u}{\partial y}\right)_{i-1,j}}{2\Delta x} + \mathcal{O}(\Delta x)^2$$

$$\left(\frac{\partial u}{\partial y}\right)_{i+1,j} = \frac{u_{i+1,j+1} - u_{i+1,j-1}}{2\Delta y} + \mathcal{O}(\Delta y)^2$$

$$\left(\frac{\partial u}{\partial y}\right)_{i-1,j} = \frac{u_{i-1,j+1} - u_{i-1,j-1}}{2\Delta y} + \mathcal{O}(\Delta y)^2$$



Second-order difference approximation

$$\left(\frac{\partial^2 u}{\partial x \partial y}\right)_{i,j} = \frac{u_{i+1,j+1} - u_{i+1,j-1} - u_{i-1,j+1} + u_{i-1,j-1}}{4\Delta x \Delta y} + \mathcal{O}[(\Delta x)^2, (\Delta y)^2]$$