

Saint-Venant equations: fluid dynamics

$$\frac{\partial}{\partial t} h + \frac{\partial}{\partial x} (uh) + \frac{\partial}{\partial y} (vh) = 0,$$

$$\frac{\partial}{\partial t} (uh) + \frac{\partial}{\partial x} \left(u^2 + \frac{1}{2} gh^2 \right) + \frac{\partial}{\partial y} (uvh) = 0,$$


$$\frac{\partial}{\partial t} (vh) + \frac{\partial}{\partial t} (uvh) + \frac{\partial}{\partial y} \left(v^2 h + \frac{1}{2} gh^2 \right) = 0,$$

- depth-integrating the Navier-Stokes equations considering the horizontal length scale is much greater than the vertical length scale.


- conservation of mass implies that the vertical velocity of the fluid is small.

- shallow water equation models have only one vertical level, so they cannot directly encompass any factor that varies with height.

**Scholarship
of Geosciences**








Saint-Venant equations: fluid dynamics




- depth-integrating the Navier-Stokes equations considering the horizontal length scale is much greater than the vertical length scale.
- conservation of mass implies that the vertical velocity of the fluid is small.
- shallow water equation models have only one vertical level, so they cannot directly encompass any factor that varies with height.



$$\begin{aligned}\frac{\partial}{\partial t} h + \frac{\partial}{\partial x} (uh) + \frac{\partial}{\partial y} (vh) &= 0, \\ \frac{\partial}{\partial t} (uh) + \frac{\partial}{\partial x} \left(u^2 + \frac{1}{2} gh^2 \right) + \frac{\partial}{\partial y} (uvh) &= 0, \\ \frac{\partial}{\partial t} (vh) + \frac{\partial}{\partial x} (uvh) + \frac{\partial}{\partial y} \left(v^2 h + \frac{1}{2} gh^2 \right) &= 0,\end{aligned}$$

Saint-Venant equations: fluid dynamics

Tohoku-Oki region, Japan, on 11 March 2011



[youtube](#)