

¹ eo-tides: Tide modelling tools for large-scale satellite Earth observation analysis

³ **Robbi Bishop-Taylor**  ¹¶, **Claire Phillips**  ¹, **Stephen Sagar**  ¹, **Vanessa Newey**¹, and **Tyler Sutterley**  ²

⁵ 1 Geoscience Australia, Australia  ² University of Washington Applied Physics Laboratory, United States of America  ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).¹⁸¹⁹²⁰²¹

Summary

The eo-tides package provides powerful parallelized tools for integrating satellite Earth observation (EO) data with ocean tide modelling. The package provides a flexible Python-based toolkit for modelling and attributing tide heights to a time-series of satellite images based on the spatial extent and acquisition time of each satellite observation (Figure 1).

eo-tides leverages advanced tide modelling functionality from the pyTMD tide prediction software (Sutterley et al., 2017), combining this fundamental tide modelling capability with EO spatial analysis tools from odc-geo (odc-geo contributors, 2024). This allows tides to be modelled in parallel automatically using over 50 supported tide models, and returned in standardised pandas (McKinney, 2010; pandas development team, 2020) and xarray (Hoyer & Joseph, 2017) data formats for further analysis.

Tools from eo-tides are designed to be applied directly to petabytes of freely available satellite data loaded from the cloud using Open Data Cube's odc-stac or datacube packages (e.g. using [Digital Earth Australia](#) or [Microsoft Planetary Computer's SpatioTemporal Asset Catalogues](#)). Additional functionality enables evaluating potential satellite-tide biases, and validating modelled tides using external tide gauge data — both important considerations for assessing the reliability and accuracy of coastal EO workflows. In combination, these open source tools support the efficient, scalable and robust analysis of coastal EO data for any time period or location globally.²²²³²⁴²⁵




Figure 1: An example of a typical eo-tides coastal EO workflow, with tide heights being modelled into every pixel in a spatio-temporal stack of satellite data (for example, from ESA's Sentinel-2 or NASA/USGS Landsat), then combined to derive insights into dynamic coastal environments.

26 Statement of need

27 Satellite remote sensing offers an unparalleled method to view and examine dynamic coastal
28 environments over large temporal and spatial scales (Turner et al., 2021; Vitousek et al.,
29 2023). However, the variable and sometimes extreme influence of ocean tides in these regions
30 can complicate analyses, making it difficult to separate the influence of changing tides from
31 patterns of true coastal change over time (Vos et al., 2019). This is a particularly significant
32 challenge for continental- to global-scale coastal EO analyses, where failing to account for
33 complex tide dynamics can lead to inaccurate or misleading insights into coastal processes
34 observed by satellites.

35 Conversely, information about ocean tides can also provide unique environmental insights that
36 can greatly enhance the utility of coastal EO data. Conventionally, satellite data dimensions
37 consider the geographical “where” and the temporal “when” of data acquisition. The addition
38 of tide height as a new analysis dimension allows data to be filtered, sorted and analysed with
39 respect to tidal processes, delivering a powerful re-imagining of traditional multi-temporal EO
40 data analysis (Sagar et al., 2017). For example, satellite data can be analysed to focus on
41 specific ecologically-significant tidal stages (e.g. high, low tide, spring or neap tides) or on
42 particular tidal processes (e.g. ebb or flow tides; Sent et al. (2025)).

43 This concept has been used to map tidally-corrected annual coastlines from Landsat satellite
44 data at continental scale (Bishop-Taylor et al., 2021), generate maps of the extent and elevation
45 of the intertidal zone (Bishop-Taylor et al., 2019; Murray et al., 2012; Sagar et al., 2017), and
46 create tidally-constrained imagery composites of the coastline at low and high tide (Sagar et
47 al., 2018). However, these approaches have been historically based on bespoke, closed-source
48 or difficult to install tide modelling tools, limiting the reproducibility and portability of these
49 techniques to new coastal EO applications. To support the next generation of coastal EO
50 workflows, there is a pressing need for new open-source approaches for combining satellite data
51 with tide modelling.

52 The `eo-tides` package aims to address these challenges by providing a set of performant
53 open-source Python tools for attributing satellite EO data with modelled ocean tides. This
54 functionality is provided in five main analysis modules (`utils`, `model`, `eo`, `stats`, `validation`)
55 which are described briefly below.

56 Key functionality

57 Setting up tide models

58 The `eo_tides.utils` module simplifies the setup of global ocean tide models, addressing a
59 common barrier in coastal EO workflows. Tools like `list_models` provide feedback on available
60 and supported models (Figure 2), while `clip_models` can be used to improve performance by
61 clipping large model files to smaller regions of interest, significantly reducing processing times
62 for high-resolution models like FES2022.

63 Comprehensive documentation is available to [guide users in setting up commonly used tide](#)
64 [models](#), including downloading, uncompressing, and organizing data files.

	Model	Expected path
✓	EOT20	tide_models/EOT20/ocean_tides
✗	FES2014	tide_models/fes2014/ocean_tide
✓	HAMTIDE11	tide_models/hamtide
✗	TPXO9.1	tide_models/TPXO9.1/DATA
...

Summary:
Available models: 2/50

Figure 2: An example output from `list_tides`, providing a useful summary table which clearly identifies available and supported tide models.

65 Modelling tides

66 The `eo_tides.model` module is powered by advanced tide modelling functionality from the
 67 pyTMD Python package ([Sutterley et al., 2017](#)).

68 pyTMD is an open-source tidal prediction software that aims to simplify the calculation of ocean
 69 and earth tides. Tides are frequently decomposed into harmonic constants (or constituents)
 70 associated with the relative positions of the sun, moon and Earth. For ocean tides, pyTMD.io
 71 contains routines for reading major constituent values from commonly available tide models,
 72 and interpolating those values to spatial locations. Information for each of the supported tide
 73 models is stored within a JSON database that is supplied with pyTMD. pyTMD.astro contains
 74 routines for computing the positions of celestial bodies for a given time. Namely for ocean
 75 tides, pyTMD computes the longitudes of the sun (S), moon (H), lunar perigee (P), ascending
 76 lunar node (N) and solar perigee (PP). pyTMD.arguments contains routines for combining the
 77 astronomical coefficients with the “Doodson number” of each constituent, along with routines
 78 for adjusting the amplitude and phase of each constituent based on their modulations over the
 79 18.6 year nodal period. Finally, pyTMD.predict uses results from those underlying functions to
 80 predict tidal values at a given location and time.

81 The `model_tides` function from `eo_tides.model` wraps pyTMD functionality to return tide
 82 predictions in a standardised pandas.DataFrame format, enabling integration with satellite
 83 EO data and parallelized processing for improved performance. Parallelisation in eo-tides
 84 is automatically optimised based on available workers and requested tide models and tide
 85 modelling locations. This built-in parallelisation can significantly improve tide modelling
 86 performance, especially for large-scale analyses run on a large multi-core machine ([Table 1](#)).
 87 Additional functions like `model_phases` classify tides or determine flow/ebb tides, critical for
 88 interpreting satellite-observed coastal processes like changing turbidity and ocean colour ([Sent](#)
 89 [et al., 2025](#)).

Table 1: A [benchmark comparison](#) of tide modelling performance with parallelisation on vs. off. This comparison was performed on an 8-core and 32-core Linux machine, for a typical large-scale analysis involving a month of hourly tides modelled at 10,000 modelling locations using three tide models (FES2022, TPXO10, GOT5.6).

Cores	Parallelisation	No parallelisation	Speedup
8	2min 46s ± 663 ms	9min 28s ± 536 ms	3.4x

Cores	Parallelisation	No parallelisation	Speedup
32	55.9 s ± 560 ms	9min 24s ± 749 ms	10.1x

90 **Combining tides with satellite data**

91 The `eo_tides.eo` module integrates modelled tides with xarray-format satellite data. For
92 tide attribution, eo-tides offers two approaches that differ in complexity and performance:
93 `tag_tides` assigns a single tide height per timestep for small-scale studies, while `pixel_tides`
94 models tides spatially and temporally for larger-scale analyses, producing a unique tide height
95 for each pixel in a dataset {tab:tide_stats}).

96 These functions can be applied to free and open satellite data for any coastal or ocean location
97 on the planet, for example using data loaded from the cloud using the [Open Data Cube](#) and
98 SpatioTemporal Asset Catalogue ([STAC contributors, 2024](#)).

Table 2: Comparison of the `tag_tides` and `pixel_tides` functions.

tag_tides	pixel_tides
<ul style="list-style-type: none"> - Assigns a single tide height to each timestep/satellite image - Ideal for local or site-scale analysis - Fast, low memory use - Single tide height per image can produce artefacts in complex tidal regions 	<ul style="list-style-type: none"> - Assigns a tide height to every individual pixel through time to capture spatial tide dynamics - Ideal for regional to global-scale coastal product generation - Slower, higher memory use - Produce spatially seamless results across large extents by applying analyses at the pixel level




Figure 3: An example tide height output produced by the `pixel_tides` function, showing spatial variability in tides across Australasia for a single timestep.

99 **Calculating tide statistics and satellite biases**

100 The `eo_tides.stats` module identifies biases caused by complex tide aliasing interactions
101 interactions between tidal dynamics and satellite observations. These interactions can prevent

satellites from observing the entire tide cycle (Eleveld et al., 2014; Sent et al., 2025), and cause coastal EO studies to produce biased or misleading results (Bishop-Taylor et al., 2019). The module produces a range of useful statistics that summarise how well a satellite time series captures real-world tidal conditions, include spread (coverage of tide range) and high/low-tide offsets (missed tidal extremes). Automated reports and plots provide insights further insights into potential biases affecting the analysis.




Figure 4: An example of tidally-biased satellite coverage, where the sensor only observes ~68% of the modelled astronomical tide range and never observes the lowest 24% of tides. Satellite bias plots show satellite observed tides as black dots, overlaid over the full range of modelled tides (blue lines).

108 Validating modelled tides

109 The `eo_tides.validation` module validates modelled tide heights using high-quality sea-level
 110 measurements from the GESLA Global Extreme Sea Level Analysis (Haigh et al., 2023) archive,
 111 providing error metrics like RMSE and MAE (Figure 5). It enables comparison of multiple tide
 112 models against observed data, allowing users to choose optimal tide models for their specific
 113 study area or application (Figure 5).




Figure 5: An example comparison of modelled tides from multiple global ocean tide models (EOT20, GOT5.5, HAMTIDE11) against observed sea level data from the Broome 62650 GESLA tide gauge, Western Australia.

114 Research projects

115 Early versions of eo-tides functions have been used for continental-scale intertidal zone
 116 elevation and exposure mapping ([Bishop-Taylor et al., 2024](#)), multi-decadal shoreline mapping
 117 across Australia ([Bishop-Taylor et al., 2021](#)) and [Africa](#), and to support tide correction for
 118 satellite-derived shorelines as part of the CoastSeg Python package ([Fitzpatrick et al., 2024](#)).

119 Acknowledgements

120 Functions from eo-tides were originally developed in the Digital Earth Australia Notebooks
 121 and Tools repository ([Krause et al., 2021](#)). We thank all DEA Notebooks contributors and
 122 maintainers for their invaluable assistance with code review, feature suggestions and code
 123 edits. This paper is published with the permission of the Chief Executive Officer, Geoscience
 124 Australia. Copyright Geoscience Australia (2025).

125 References

- 126 Bishop-Taylor, R., Nanson, R., Sagar, S., & Lymburner, L. (2021). Mapping Australia's
 127 Dynamic Coastline at Mean Sea Level using Three Decades of Landsat Imagery. *Remote
 128 Sensing of Environment*, 267, 112734. <https://doi.org/10.1016/j.rse.2021.112734>
- 129 Bishop-Taylor, R., Phillips, C., Newey, V., & Sagar, S. (2024). *Digital Earth Australia Intertidal*.
 130 Commonwealth of Australia (Geoscience Australia). <https://doi.org/10.26186/149403>
- 131 Bishop-Taylor, R., Sagar, S., Lymburner, L., & Beaman, R. J. (2019). Between the tides:
 132 Modelling the elevation of australia's exposed intertidal zone at continental scale. *Estuarine,
 133 Coastal and Shelf Science*, 223, 115–128. <https://doi.org/10.1016/j.ecss.2019.03.006>
- 134 Eleveld, M. A., Van der Wal, D., & Van Kessel, T. (2014). Estuarine suspended par-
 135 ticulate matter concentrations from sun-synchronous satellite remote sensing: Tidal
 136 and meteorological effects and biases. *Remote Sensing of Environment*, 143, 204–215.
 137 <https://doi.org/10.1016/j.rse.2013.12.019>
- 138 Fitzpatrick, S., Buscombe, D., Warrick, J. A., Lundine, M. A., & Vos, K. (2024). CoastSeg: An
 139 accessible and extendable hub for satellite-derived-shoreline (SDS) detection and mapping.
 140 *Journal of Open Source Software*, 9(99), 6683. <https://doi.org/10.21105/joss.06683>
- 141 Haigh, I. D., Marcos, M., Talke, S. A., Woodworth, P. L., Hunter, J. R., Hague, B. S.,
 142 Arns, A., Bradshaw, E., & Thompson, P. (2023). GESLA version 3: A major update to

- 143 the global higher-frequency sea-level dataset. *Geoscience Data Journal*, 10(3), 293–314.
144 <https://doi.org/10.1002/gdj3.174>
- 145 Hoyer, S., & Joseph, H. (2017). xarray: N-d labeled arrays and datasets in python. *Journal of*
146 *Open Research Software*, 5(1). <https://doi.org/10.5334/jors.148>
- 147 Krause, C., Dunn, B., Bishop-Taylor, R., Adams, C., Burton, C., Alger, M., Chua, S., Phillips, C.,
148 Newey, V., Kouzoubov, K., Leith, A., Ayers, D., & Hicks, A. (2021). *Digital Earth Australia*
149 *notebooks and tools repository*. <https://github.com/GeoscienceAustralia/dea-notebooks/>;
150 Commonwealth of Australia (Geoscience Australia). <https://doi.org/10.26186/145234>
- 151 McKinney, Wes. (2010). Data Structures for Statistical Computing in Python. In Stéfan van
152 der Walt & Jarrod Millman (Eds.), *Proceedings of the 9th Python in Science Conference*
153 (pp. 56–61). <https://doi.org/10.25080/Majora-92bf1922-00a>
- 154 Murray, N. J., Phinn, S. R., Clemens, R. S., Roelfsema, C. M., & Fuller, R. A. (2012).
155 Continental scale mapping of tidal flats across east asia using the landsat archive. *Remote*
156 *Sensing*, 4(11), 3417–3426. <https://doi.org/10.3390/rs4113417>
- 157 odc-geo contributors. (2024). Opendatacube/odc-geo. In *Github repository*. GitHub.
158 <https://github.com/opendatacube/odc-geo>
- 159 pandas development team. (2020). *Pandas-dev/pandas: pandas* (latest). Zenodo. <https://doi.org/10.5281/zenodo.3509134>
- 160 Sagar, S., Phillips, C., Bala, B., Roberts, D., Lymburner, L., & Beaman, R. J. (2018).
161 Generating continental scale pixel-based surface reflectance composites in coastal regions
162 with the use of a multi-resolution tidal model. *Remote Sensing*, 10(3), 480. <https://doi.org/10.3390/rs10030480>
- 163 Sagar, S., Roberts, D., Bala, B., & Lymburner, L. (2017). Extracting the intertidal extent and
164 topography of the australian coastline from a 28 year time series of landsat observations.
165 *Remote Sensing of Environment*, 195, 153–169. <https://doi.org/10.1016/j.rse.2017.04.009>
- 166 Sent, G., Antunes, C., Spyракος, E., Jackson, T., Atwood, E. C., & Brito, A. C. (2025). What
167 time is the tide? The importance of tides for ocean colour applications to estuaries. *Remote*
168 *Sensing Applications: Society and Environment*, 37, 101425. <https://doi.org/10.2139/ssrn.4858713>
- 169 STAC contributors. (2024). *SpatioTemporal Asset Catalog (STAC) specification*. <https://stacspec.org>
- 170 Sutterley, T. C., Alley, K., Brunt, K., Howard, S., Padman, L., & Siegried, M. (2017). *pyTMD:*
171 *Python-based tidal prediction software*. Zenodo. <https://doi.org/10.5281/zenodo.5555395>
- 172 Turner, I. L., Harley, M. D., Almar, R., & Bergsma, E. W. J. (2021). Satellite optical imagery
173 in Coastal Engineering. *Coastal Engineering*, 167, 103919. <https://doi.org/10.1016/j.coastaleng.2021.103919>
- 174 Vitousek, S., Buscombe, D., Vos, K., Barnard, P. L., Ritchie, A. C., & Warrick, J. A. (2023).
175 The future of coastal monitoring through satellite remote sensing. *Cambridge Prisms: Coastal*
176 *Futures*, 1, e10. <https://doi.org/10.1017/cft.2022.4>
- 177 Vos, K., Splinter, K. D., Harley, M. D., Simmons, J. A., & Turner, I. L. (2019). CoastSat: A
178 Google Earth Engine-enabled Python toolkit to extract shorelines from publicly available
179 satellite imagery. *Environmental Modelling & Software*, 122, 104528. <https://doi.org/10.1016/j.envsoft.2019.104528>