

¹ eo-tides: Tide modelling tools for large-scale satellite Earth observation analysis

³ **Robbi Bishop-Taylor**  ¹¶, **Claire Phillips**  ¹, **Stephen Sagar**  ¹, **Vanessa Newey**¹, and **Tyler Sutterley**  ²

⁵ 1 Geoscience Australia, Australia  ² University of Washington Applied Physics Laboratory, United States of America  ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright¹⁸ and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).¹⁹²⁰

Summary

The eo-tides package provides powerful parallelized tools for integrating satellite Earth observation (EO) data with ocean tide modelling. The package provides a flexible Python-based API that facilitates the modelling and attribution of tide heights to a time-series of satellite images based on the spatial extent and acquisition time of each satellite observation (Figure 1).

eo-tides leverages advanced tide modelling functionality from the pyTMD tide prediction software (Sutterley et al., 2017), combining this fundamental tide modelling capability with EO spatial analysis tools from odc-geo ([odc-geo development team, 2024](#)). This allows tides to be modelled in parallel automatically using over 50 supported tide models, and returned in standardised pandas ([McKinney, 2010](#); [pandas development team, 2020](#)) and xarray ([Hoyer & Joseph, 2017](#)) data formats for further analysis.

Tools from eo-tides are designed to be applied directly to petabytes of freely available satellite data loaded from the cloud using Open Data Cube's odc-stac or datacube packages (e.g. using [Digital Earth Australia](#) or [Microsoft Planetary Computer's SpatioTemporal Asset Catalogue](#)). Additional functionality enables evaluating potential satellite-tide biases, and validating modelled tides using external tide gauge data — both important considerations for assessing the reliability and accuracy of coastal EO workflows. In combination, these open source tools support the efficient, scalable and robust analysis of coastal EO data for any time period or location globally.




Figure 1: An example of a typical eo-tides coastal EO workflow, with tide heights being modelled into every pixel in a spatio-temporal stack of satellite data (for example, from ESA's Sentinel-2 or NASA/USGS Landsat), then combined to derive insights into dynamic coastal environments.

27 Statement of need

28 Satellite remote sensing offers an unparalleled method to view and examine dynamic coastal
29 environments over large temporal and spatial scales (Turner et al., 2021; Vitousek et al.,
30 2023). However, the variable and sometimes extreme influence of ocean tides in these regions
31 can complicate analyses, making it difficult to separate the influence of changing tides from
32 patterns of true coastal change over time (Vos et al., 2019). This is a particularly significant
33 challenge for continental- to global-scale coastal EO analyses, where failing to account for
34 complex tide dynamics can lead to inaccurate or misleading insights into coastal processes
35 observed by satellites.

36 Conversely, information about ocean tides can also provide unique environmental insights that
37 can greatly enhance the utility of coastal EO data. Conventionally, satellite data dimensions
38 consider the geographical “where” and the temporal “when” of data acquisition. The addition
39 of tide height as a new analysis dimension allows data to be filtered, sorted and analysed with
40 respect to tidal processes, delivering a powerful re-imagining of traditional multi-temporal EO
41 data analysis (Sagar et al., 2017). For example, satellite data can be analysed to focus on
42 specific ecologically-significant tidal stages (e.g. high, low tide, spring or neap tides) or on
43 particular tidal processes (e.g. ebb or flow tides).

44 This concept has been used to map tidally-corrected annual coastlines from Landsat satellite
45 data at continental scale (Bishop-Taylor et al., 2021), generate maps of the extent and elevation
46 of the intertidal zone (Bishop-Taylor et al., 2019; Murray et al., 2012; Sagar et al., 2017), and
47 create tidally-constrained imagery composites of the coastline at low and high tide (Sagar et
48 al., 2018). However, these approaches have been historically based on bespoke, closed-source
49 or difficult to install tide modelling tools, limiting the reproducibility and portability of these
50 techniques to new coastal EO applications. To support the next generation of coastal EO
51 workflows, there is a pressing need for new open-source approaches for combining satellite data
52 with tide modelling.

53 The `eo-tides` package aims to address these challenges by providing a set of performant
54 open-source Python tools for attributing satellite EO data with modelled ocean tides. This
55 functionality is provided in five main analysis modules (`utils`, `model`, `eo`, `stats`, `validation`)
56 which are described briefly below.

57 Setting up tide models

58 A key barrier to utilising tide modelling in EO workflows is the complexity and difficulty of
59 initially setting up global ocean tide models for analysis. To address this, the `eo_tides.utils`
60 module contains useful tools for preparing tide model data files for use in `eo-tides`. This
61 includes the `list_models` function that provides visual feedback on the tide models a user has
62 available in their system, while highlighting the naming conventions and directory structures
63 required by the underlying pyTMD tide prediction software (Figure 2).

64 Running tide modelling using the default tide modelling data provided by external providers can
65 be slow due to the large size of these files — especially for recent high-resolution models like
66 FES2022 (Carrere et al., 2022). To improve tide modelling performance, it can be extremely
67 useful to clip tide model files to a smaller region of interest (e.g. the extent of a country
68 or coastal region). The `clip_models` function can be used to automatically clip all suitable
69 NetCDF-format model data files to a user-supplied bounding box, potentially improving tide
70 modelling performance by over an order of magnitude.

71 These tools are accompanied by comprehensive documentation explaining how to set up several
72 of the most commonly used global ocean tide models, including details on how to download or
73 request access to model files, and how to uncompress and arrange the data on disk.

	Model	Expected path
	EOT20	tide_models/EOT20/ocean_tides
	FES2014	tide_models/fes2014/ocean_tide
	HAMTIDE11	tide_models/hamtide
	TPXO9.1	tide_models/TPXO9.1/DATA
...

Summary:

Available models: 2/50

Figure 2: An example output from `list_tides`, providing a useful summary table which clearly identifies available tide models.

74 Modelling tides

75 The `eo_tides.model` module builds upon advanced tide modelling capability provided by the
 76 pyTMD tide prediction software ([Sutterley et al., 2017](#)).

77 [TODO Tyler: Insert brief paragraph here about the core capability of the pyTMD package, with
 78 general background to the science used to predict tides and the range of supported global tide
 79 models]

80 [TODO Robbi: Insert brief paragraph here about how `eo-tides` wraps pyTMD functionality to
 81 model tides in parallel and return data in pandas/xarray format required for EO analysis]

82 Tide modelling functionality in the `model_tides` function is primarily intended to support
 83 more complex EO-related capability in the `eo_tides.eo` module. However it can also be
 84 used independently of EO data, for example for any application that requires a time series of
 85 modelled tide heights. In addition to modelling tide heights, the `model_phases` function allows
 86 users to calculate tidal phases at any location and time. This can be used to classify tides
 87 into high and low tide observations, or determine whether the tide was rising (i.e. flow tide) or
 88 falling (i.e. ebb tide) at any point in time.

89 Combining tides with satellite data

90 The `eo_tides.eo` module contains the package's core functionality, focusing on tools for
 91 attributing satellite data with modelled tide heights. For tide attribution, `eo-tides` offers two
 92 approaches that differ in complexity and performance: `tag_tides` and `pixel_tides` ([Table 1](#)).

93 The `tag_tides` function provides a fast and efficient method for small scale applications where
 94 tides are unlikely to vary across a study area. This approach allocates a single tide height
 95 to each satellite data timestep, based on the geographic-centroid of the dataset and the
 96 acquisition time of each image. Having tide height as a variable allows the selection and
 97 analysis of satellite data based on tides. For example, all available satellite observations for an
 98 area of interest could be sorted by tide height, or used to extract and compare the lowest and
 99 highest tide images in the time series.

100 However, in reality tides vary spatially – potentially by many metres in areas of complex and
 101 extreme tidal dynamics. This means that an individual satellite image can capture a range of
 102 contrasting tide conditions. For larger scale coastal EO analysis, the `pixel_tides` function can
 103 be used to seamlessly model tides through both time and space, producing three-dimensional

¹⁰⁴ “tide height” datacube that can be integrated with satellite data. For efficient processing,
¹⁰⁵ pixel_tides ‘models tides into a customisable low resolution grid surrounding each satellite
¹⁰⁶ image in the time series. These modelled tides are then re-projected back into the original
¹⁰⁷ resolution of the input satellite image, returning a unique tide height for every individual
¹⁰⁸ satellite pixel through time (Figure 3).

Table 1: Comparison of the tag_tides and pixel_tides functions.

tag_tides	pixel_tides
<ul style="list-style-type: none"> - Assigns a single tide height to each timestep/satellite image - Ideal for local or site-scale analysis - Fast, low memory use - Single tide height per image can produce artefacts in complex tidal regions 	<ul style="list-style-type: none"> - Assigns a tide height to every individual pixel through time to capture spatial tide dynamics - Ideal for regional to global-scale coastal product generation - Slower, higher memory use - Produce spatially seamless results across large extents by applying analyses at the pixel level




Figure 3: An example tide height output produced by the pixel_tides function, showing spatial variability in tides across the Australasia for a single timestep.

¹⁰⁹ Calculating tide statistics and satellite biases

¹¹⁰ The eo_tides.stats module contains tools for calculating statistics describing local tide
¹¹¹ dynamics, as well as biases caused by interactions between tidal processes and satellite orbits.
¹¹² Complex tide aliasing interactions between temporal tide dynamics and the regular overpass
¹¹³ timing of sun-synchronous satellite sensors mean that satellites often do not always observe
¹¹⁴ the entire tidal cycle (Eleveld et al., 2014). Biases in satellite coverage of the tidal cycle can
¹¹⁵ mean that tidal extremes (e.g. the lowest or highest tides at a location) or particular tidal
¹¹⁶ processes may either never be captured by satellites, or be over-represented in the satellite
¹¹⁷ record. Local tide dynamics can cause these biases to vary greatly both through time and space
¹¹⁸ (Bishop-Taylor et al., 2019), making it challenging to compare coastal processes consistently -
¹¹⁹ particularly for large-scale coastal EO analyses.

120 To ensure that coastal EO analyses are not inadvertently affected by tide biases, it is important
 121 to understand and compare how well the tides observed by satellites match the full range of
 122 modelled tides at a location. The `tide_stats` function compares the subset of tides observed
 123 by satellite data against the full range of tides modelled at a regular interval through time
 124 across the entire time period covered by the satellite dataset. This comparison is used to
 125 calculate several useful statistics that summarise how well a satellite time series captures the
 126 full range of real-world tidal conditions (Bishop-Taylor et al., 2019). These statistics include:
 127
 128 1. Spread: The proportion of the modelled astronomical tidal range that was observed by
 129 satellites. A high value indicates good coverage of the tide range.
 130 2. High-tide offset: The proportion of the highest tides never observed by satellites, relative
 131 to the modelled astronomical tidal range. A high value indicates that the satellite data
 132 never captures the highest tides.
 133 3. Low-tide offset: The proportion of the lowest tides never observed by satellites, relative
 134 to the modelled astronomical tidal range. A high value indicates that the satellite data
 135 never captures the lowest tides.
 136 A satellite tide bias investigation for a coastal area of interest will return an automated report
 and plot (Figure 4), adding insightful tide-based context to a coastal EO analysis:




Figure 4: In this example satellite time series, the data captured a biased proportion of the tide range: only observing ~68% of the modelled astronomical tide range, and never observing the lowest 24% of tides. The plot visually demonstrates the relationships between satellite observed tide heights (black dots) and modelled astronomical tide height (blue lines) at this location.

137 **Validating modelled tide heights**

138 The `eo_tides.validation` module contains tools for validating modelled tides against observed
139 sea level data. The tide models supported by `eo-tides` can vary significantly in accuracy
140 across the world's coastlines. Evaluating the accuracy of modelled tides is critical for ensuring
141 that resulting marine or coastal EO analyses are reliable and useful.

142 Validation functionality in `eo-tides` provides a convenient tool for loading high-quality sea-level
143 measurements from the GESLA Global Extreme Sea Level Analysis ([Haigh et al., 2023](#)) archive
144 – a global dataset of almost 90,713 years of sea level data from 5,119 records across the world.
145 The `load_gauge_gesla` function allows GESLA data to be loaded for the same location and
146 time period as a satellite time series. Differences between modelled and observed tide heights
147 can then be quantified through the calculation of accuracy statistics that include the Root
148 Mean Square Error (RMSE), Mean Absolute Error (MAE), R-squared and bias.

149 Furthermore, different ocean tide models perform differently in different coastal locations.
150 `eo-tides` allows multiple tide models to be compared against GESLA data simultaneously,
151 empowering users to make informed decisions and choose the optimal tide model that best
152 suits their specific location or application.

153 **Research projects**

154 Early versions of functions provided in `eo-tides` has been for continental-scale modelling of
155 the elevation and exposure of Australia's intertidal zone ([Bishop-Taylor et al., 2024](#)), and to
156 support tide correction for satellite-derived shorelines as part of the CoastSeg Python package
157 ([Fitzpatrick et al., 2024](#)).

158 **Acknowledgements**

159 Functions from `eo-tides` were originally developed in the Digital Earth Australia Notebooks
160 and Tools repository ([Krause et al., 2021](#)). The authors would like to thank all DEA Notebooks
161 contributors and maintainers for their invaluable assistance with code review, feature suggestions
162 and code edits. This paper is published with the permission of the Chief Executive Officer,
163 Geoscience Australia.

164 **References**

- 165 Bishop-Taylor, R., Nanson, R., Sagar, S., & Lymburner, L. (2021). Mapping Australia's
166 Dynamic Coastline at Mean Sea Level using Three Decades of Landsat Imagery. *Remote
167 Sensing of Environment*, 267, 112734. <https://doi.org/10.1016/j.rse.2021.112734>
- 168 Bishop-Taylor, R., Phillips, C., Newey, V., & Sagar, S. (2024). *Digital Earth Australia Intertidal*.
169 Commonwealth of Australia (Geoscience Australia). <https://doi.org/10.26186/149403>
- 170 Bishop-Taylor, R., Sagar, S., Lymburner, L., & Beaman, R. J. (2019). Between the tides:
171 Modelling the elevation of australia's exposed intertidal zone at continental scale. *Estuarine,
172 Coastal and Shelf Science*, 223, 115–128. <https://doi.org/10.1016/j.ecss.2019.03.006>
- 173 Carrere, L., Lyard, F., Cancet, M., Allain, D., Dabat, M.-L., Fouchet, E., Sahuc, E., Faugere,
174 Y., Dibarboire, G., & Picot, N. (2022). A new barotropic tide model for global ocean:
175 FES2022. *2022 Ocean Surface Topography Science Team Meeting*, 43.
- 176 Eleveld, M. A., Van der Wal, D., & Van Kessel, T. (2014). Estuarine suspended partic-
177 ulate matter concentrations from sun-synchronous satellite remote sensing: Tidal and
178 meteorological effects and biases. *Remote Sensing of Environment*, 143, 204–215.

- 179 Fitzpatrick, S., Buscombe, D., Warrick, J. A., Lundine, M. A., & Vos, K. (2024). CoastSeg: An
180 accessible and extendable hub for satellite-derived-shoreline (SDS) detection and mapping.
181 *Journal of Open Source Software*, 9(99), 6683. <https://doi.org/10.21105/joss.06683>
- 182 Haigh, I. D., Marcos, M., Talke, S. A., Woodworth, P. L., Hunter, J. R., Hague, B. S.,
183 Arns, A., Bradshaw, E., & Thompson, P. (2023). GESLA version 3: A major update to
184 the global higher-frequency sea-level dataset. *Geoscience Data Journal*, 10(3), 293–314.
185 <https://doi.org/https://doi.org/10.1002/gdj3.174>
- 186 Hoyer, S., & Joseph, H. (2017). xarray: N-d labeled arrays and datasets in python. *Journal of
187 Open Research Software*, 5(1). <https://doi.org/10.5334/jors.148>
- 188 Krause, C., Dunn, B., Bishop-Taylor, R., Adams, C., Burton, C., Alger, M., Chua, S., Phillips, C.,
189 Newey, V., Kouzoubov, K., Leith, A., Ayers, D., & Hicks, A. (2021). *Digital Earth Australia
190 notebooks and tools repository*. <https://github.com/GeoscienceAustralia/dea-notebooks/>;
191 Commonwealth of Australia (Geoscience Australia). <https://doi.org/10.26186/145234>
- 192 McKinney, Wes. (2010). Data Structures for Statistical Computing in Python. In Stéfan van
193 der Walt & Jarrod Millman (Eds.), *Proceedings of the 9th Python in Science Conference*
194 (pp. 56–61). <https://doi.org/10.25080/Majora-92bf1922-00a>
- 195 Murray, N. J., Phinn, S. R., Clemens, R. S., Roelfsema, C. M., & Fuller, R. A. (2012).
196 Continental scale mapping of tidal flats across east asia using the landsat archive. *Remote
197 Sensing*, 4(11), 3417–3426.
- 198 Sagar, S., Phillips, C., Bala, B., Roberts, D., Lymburner, L., & Beaman, R. J. (2018).
199 Generating continental scale pixel-based surface reflectance composites in coastal regions
200 with the use of a multi-resolution tidal model. *Remote Sensing*, 10(3), 480. <https://doi.org/10.3390/rs10030480>
- 202 Sagar, S., Roberts, D., Bala, B., & Lymburner, L. (2017). Extracting the intertidal extent and
203 topography of the australian coastline from a 28 year time series of landsat observations.
204 *Remote Sensing of Environment*, 195, 153–169.
- 205 Sutterley, T. C., Alley, K., Brunt, K., Howard, S., Padman, L., & Siegried, M. (2017). *pyTMD:
206 Python-based tidal prediction software*. Zenodo. <https://doi.org/10.5281/zenodo.5555395>
- 207 team, odc-geo development. (2024). Opendatacube/odc-geo. In *GitHub repository*. GitHub.
208 <https://github.com/opendatacube/odc-geo>
- 209 team, pandas development. (2020). *Pandas-dev/pandas: pandas* (latest). Zenodo. <https://doi.org/10.5281/zenodo.3509134>
- 210 Turner, I. L., Harley, M. D., Almar, R., & Bergsma, E. W. J. (2021). Satellite optical imagery
211 in Coastal Engineering. *Coastal Engineering*, 167, 103919. <https://doi.org/10.1016/j.coastaleng.2021.103919>
- 212 Vitousek, S., Buscombe, D., Vos, K., Barnard, P. L., Ritchie, A. C., & Warrick, J. A. (2023).
213 The future of coastal monitoring through satellite remote sensing. *Cambridge Prisms:
214 Coastal Futures*, 1, e10. <https://doi.org/10.1017/cft.2022.4>
- 215 Vos, K., Splinter, K. D., Harley, M. D., Simmons, J. A., & Turner, I. L. (2019). CoastSat: A
216 Google Earth Engine-enabled Python toolkit to extract shorelines from publicly available
217 satellite imagery. *Environmental Modelling & Software*, 122, 104528. <https://doi.org/10.1016/j.envsoft.2019.104528>