

2. A. R. Jones *et al.*, *Nat. Biotechnol.* **25**, 1127 (2007).
3. S. A. Sansone *et al.*, *OMICS* **12**, 143 (2008).
4. B. Smith *et al.*, *Nat. Biotechnol.* **25**, 1251 (2007).
5. C. F. Taylor *et al.*, *Nat. Biotechnol.* **26**, 889 (2008).
6. "Protocol for implementing open access data," <http://sciencecommons.org/projects/publishing/open-access-data-protocol>.
7. European Science Foundation (ESF), *Shared Responsibilities in Sharing Research Data: Policies and Partnerships*, Report of an ESF–Deutsch Forschungsgemeinschaft workshop, Padua, Italy, 21 September 2007 (ESF, Strasbourg, France, 2008).
8. European Commission (EC), "On scientific information in the digital age: Access, dissemination and preservation"; http://ec.europa.eu/research/science-society/document_library/pdf_06/communication-022007_en.pdf
9. FDA, "Genomic data submission"; www.fda.gov/Drugs/ScienceResearch/ResearchAreas/Pharmacogenetics/ucm083641.htm.
10. EMEA, *Guideline on Pharmacogenetics Briefing Meetings*; www.emea.europa.eu/pdfs/human/pharmacogenetics/2022704en.pdf.
11. EPA, *Potential Implications of Genomics for Regulatory and Risk Assessment Applications at EPA*; www.epa.gov/osa/genomics.htm.
12. "Pistoia vision," www.pistoiaalliance.org/.
13. Organisation for Economic Co-operation and Development, *OECD Principles and Guidelines for Access to Research Data from Public Funding* (OECD, Paris, 2007); www.oecd.org/dataoecd/9/61/38500813.pdf.
14. B. Tiwari, D. Field, J. Snape, *Nature* **439**, 912 (2006).
15. D. Field, B. Tiwari, J. Snape, *PLoS Biol.* **3**, e297 (2005).
16. D. Field *et al.*, *Nat. Biotechnol.* **24**, 801 (2006).
17. Economic and Social Data Service, www.esds.ac.uk/.
18. D. Field, S. A. Sansone, *OMICS* **10**, 84 (2006).
19. Standardizing data. *Nat. Cell Biol.* **10**, 1123 (2008).
20. Cancer Biomedical Informatics Grid, (caBIG), National Cancer Institute, NIH; <http://cabig.cancer.gov>.
21. Biomedical Informatics Research Network, www.nbirn.net/.
22. ELIXIR, <http://www.elixir-europe.org>.
23. EC, "INSPIRE Directive," <http://inspire.jrc.ec.europa.eu/index.cfm>.
24. DCC, www.dcc.ac.uk/.
25. C. Thomas, *Science* **324**, 1632 (2009).
26. S. Wiley, *Scientist* **23**, 33 (2009).
27. E. Pennisi, *Science* **324**, 1000 (2009).
28. Earth System Science Data, www.earth-system-science-data.net/.
29. D. Howe *et al.*, *Nature* **455**, 47 (2008).
30. International Society for Biocuration, www.biocurator.org.
31. H. Barsnes *et al.*, *Nat. Biotechnol.* **27**, 598 (2009).
32. Investigation/Study/Assay (ISA) Infrastructure for Managing Experimental Metadata, <http://isatab.sf.net>.
33. C. Brooksbank, J. Quackenbush, *OMICS* **10**, 94 (2006).
34. Z. Merali, J. Giles, *Nature* **435**, 1010 (2005).
35. Biosharing, <http://biosharing.org/>.
36. Toronto International Data Release Workshop Authors, *Nature* **461**, 168 (2009).
37. P. N. Schofield *et al.*, *Nature* **461**, 171 (2009).
38. "Summary of principles agreed at the First International Strategy Meeting on Human Genome Sequencing, Bermuda, 25 to 28 February 1996 (Human Genome Organisation, Singapore, 1996); available at www.ornl.gov/sci/techresources/Human_Genome/research/bermuda.shtml.
39. Sharing Data from Large-Scale Biological Research Projects, *A System of Tripartite Responsibility*, Fort Lauderdale, FL, 14 and 15 January 2003 (Wellcome Trust, 2003); available at www.wellcome.ac.uk/stellent/groups/corporatesite/@policy_communications/documents/web_document/wtd003207.pdf.

Supporting Online Material

www.sciencemag.org/cgi/content/full/326/5950/234/DC1

10.1126/science.1180598

GENOMICS

Genome Project Standards in a New Era of Sequencing

P. S. G. Chain,^{1,2,3*}† D. V. Grafham,⁴† S. R. Fulton,⁵† M. G. FitzGerald,⁶† J. Hostettler,⁷† D. Muzny,⁸† J. Ali,⁹ B. Birren,⁶ D. C. Bruce,^{1,10} C. Buhay,⁸ J. R. Cole,³ Y. Ding,⁸ S. Dugan,⁸ D. Field,¹¹ G. M. Garrity,³ R. Gibbs,⁸ T. Graves,⁵ C. S. Han,^{1,10} S. H. Harrison,^{3*} S. Highlander,⁸ P. Hugenholtz,¹ H. M. Khouri,¹² C. D. Kodira,^{6*} E. Kolker,^{13,14} N. C. Kyrpides,¹ D. Lang,¹² A. Lapidus,¹ S. A. Malfatti,¹² V. Markowitz,¹⁵ T. Metha,⁶ K. E. Nelson,⁷ J. Parkhill,⁴ S. Pitluck,¹ X. Qin,⁸ T. D. Read,¹⁶ J. Schmutz,¹⁷ S. Sozhamannan,¹⁸ P. Sterk,¹¹ R. L. Strausberg,⁷ G. Sutton,⁷ N. R. Thomson,⁴ J. M. Tiedje,³ G. Weinstock,⁵ A. Wollam,⁵ Genomic Standards Consortium Human Microbiome Project JUMPstart Consortium,† J. C. Detter¹⁰†‡

For over a decade, genome sequences have adhered to only two standards that are relied on for purposes of sequence analysis by interested third parties (1, 2). However, ongoing developments in revolutionary sequencing technologies have resulted in a redefinition of traditional whole-genome sequencing that requires reevaluation of such standards. With commercially available 454 pyrosequencing (followed by Illumina, SOLiD, and now Helicos), there has been an explosion of genomes sequenced under the moniker "draft"; however, these can be very poor quality genomes (due to inherent errors in the sequencing technologies, and the inability of assembly programs to fully address these errors). Further, one can only infer that such

draft genomes may be of poor quality by navigating through the databases to find the number and type of reads deposited in sequence trace repositories (and not all genomes have this available), or to identify the number of contigs or genome fragments deposited to the database. The difficulty in assessing the quality of such deposited genomes has created some havoc for genome analysis pipelines and has contributed to many wasted hours. Exponential leaps in raw sequencing capability and greatly reduced prices have further skewed the time- and cost-ratios of draft data generation versus the painstaking process of improving and finishing a genome. The result is an ever-widening gap between drafted and finished genomes that only promises to continue (see

More detailed sequence standards that keep up with revolutionary sequencing technologies will aid the research community in evaluating data.

the figure, page 236); hence, there is an urgent need to distinguish good from poor data sets.


The sequencing institutes and consortia whom we represent believe that a new set of standards is required for genome sequences. The following represents community-defined categories of standards that better reflect the quality of the genome sequence, based on our understanding of the technologies, available assemblers, and efforts to improve upon drafted genomes. Due to the increasingly rapid pace of genomics, we avoided rigid numerical thresholds in our definitions to take into account products achieved by any combination of technology, chemistry, assembler, or improvement and/or finishing process.

Standard Draft: minimally or unfiltered data, from any number of different sequencing platforms, that are assembled into contigs. This is the minimum standard for a submission to the public databases. Sequence of this quality will likely harbor many regions of poor quality and can be relatively incomplete. It may not always be possible to remove contaminating sequence data. Despite its shortcomings, Standard Draft is the least expensive to produce and still possesses useful information.

High-Quality Draft: overall coverage representing at least 90% of the genome or target region. Efforts should be made to exclude contaminating sequences. This is still a draft assembly with little or no manual review of the product. Sequence errors and misassem-

¹U.S. Department of Energy Joint Genome Institute. ²Lawrence Livermore National Laboratory. ³Michigan State University. ⁴The Sanger Institute. ⁵Washington University School of Medicine. ⁶The Broad Institute. ⁷J. Craig Venter Institute. ⁸Baylor College of Medicine. ⁹Ontario Institute for Cancer Research. ¹⁰Los Alamos National Laboratory. ¹¹Natural Environment Research Council Centre for Ecology and Hydrology. ¹²National Center for Biotechnology Information. ¹³Seattle Children's Hospital and Research Institute. ¹⁴University of Washington School of Medicine. ¹⁵Lawrence Berkeley National Laboratory. ¹⁶Emory GRA (Georgia Research Alliance) Genomics Center. ¹⁷HudsonAlpha Institute. ¹⁸Naval Medical Research Center.

*Full affiliations are available on Science Online. †Finishing in the Future Working Group members. ‡These authors contributed equally to organizing this work. §Authors for correspondence. E-mail: pchain@lanl.gov (P.S.G.C.); dg1@sanger.ac.uk (D.V.G.)



Trends in generation of drafted and finished genomes. A conservative estimate of future projects is shaded in light blue. Data were derived from (6, 7).

blies are possible, with no implied order and orientation to contigs. This is appropriate for general assessment of gene content.

Improved High-Quality Draft: additional work has been performed beyond the initial shotgun sequencing and High-Quality Draft assembly, by using either manual or automated methods. This should contain no discernable misassemblies and should have undergone some form of gap resolution to reduce the number of contigs and supercontigs (or scaffolds). Undetectable misassemblies are still possible, particularly in repetitive regions. Low-quality regions and potential base errors may also be present. This standard is normally adequate for comparison with other genomes.

Annotation-Directed Improvement: may overlap with the previous standards, but the term emphasizes the verification and correction of anomalies within coding regions, such as frameshifts, and stop codons. It will most often be used in cases involving complex genomes where improvement beyond this category fails to outweigh the associated costs. Gene models (gene calls, including intron-exon determination for eukaryotes) and annotation of the genomic content should fully support the biology of the organism and the scientific questions being investigated. Exceptions to this gene-specific finishing standard should be noted in the submission. Repeat regions at this level are not resolved, so errors in those regions are much more likely. This standard is useful for gene comparisons, alternative splicing analysis, and pathway reconstruction.

Noncontiguous Finished: describes high-quality assemblies that have been subject to automated and manual improvement, and where closure approaches have been successful for almost all gaps, misassemblies, and low-quality regions. Attempts have been

made to resolve all gap and sequence uncertainties, and only those recalcitrant to resolution remain (with notations in the genome submission as to the nature of the uncertainty). This product is thus of “Finished” quality with the only exception being repetitive or intractable gaps, along with heterochromatic sequence for eukaryotic applications. Thus, it is appropriate for most analyses. For nearly all higher organisms, this is the grade previously called “Finished.”

Finished: refers to the current gold standard; genome sequences with less than 1 error per 100,000 base pairs and where each replicon is assembled into a single contiguous sequence with a minimal number of possible exceptions commented in the submission record. All sequences are complete and have been reviewed and edited, all known misassemblies have been resolved, and repetitive sequences have been ordered and correctly assembled. Remaining exceptions to highly accurate sequence within the euchromatin are commented in the submission. The Finished product is appropriate for all types of detailed analyses and acts as a high-quality reference genome for comparative purposes. Some microbial genome sequences where multiple platforms have been used for the same genome have exceeded this standard, and it is believed that no bases are incorrect except for natural, low-level biological variation.

Intermediate standards often overlap, and although we do not advocate any one standard, we recommend that the target standard be based on the needs and goals of each project. There may be cases where select regions will be targeted for improvement and more than one standard may apply (such regionally improved sequences should be identified). This approach is most often used for eukaryotic whole-genome sequencing projects,

where the cost of complete finishing remains prohibitive, and allows improvement to be directed at euchromatic sequence, because heterochromatic sequence remains largely recalcitrant to available approaches. Legacy eukaryotic tiling path standards will remain in use for a time.

Here, we have attempted to capture in a technology-independent fashion the types of whole-genome sequencing projects that are beginning to populate databases, and we have defined a set of standards that accommodate a growing list of alternative genome products that have been obtained via less conventional means, such as environmental (metagenomic) or single-cell sequencing. Ongoing discussions with genome database repositories have been met with enthusiasm, and the implementation of these standards as a requirement for genome submissions is expected. To aid in adoption of this classification of sequence finishing standards, we have added this classification to the Sequence Ontology (3) where it can now be used to comply with the Genomic Standards Consortium’s (GSC) “Minimum Information about a Genome Sequence” standard (4) “sequencing status” descriptor. Furthermore, the efforts described here recently have been adopted under the umbrella of the GSC (5). This common currency in defining the products of genome projects enables better management of expectations and allows users of genomic data to assess the quality of the deposited available sequences and decide whether these meet their needs.

References and Notes

1. International standards for sequence fidelity, established at the Second International Strategy Meeting on Human Genome Sequencing in Bermuda in 1997 (27 February to 2 March). “Finished” quality standards, commonly known as the Bermuda standards, defined finished sequence as a contiguous sequence with less than one error per 10,000 bases. Almost everything else was “draft.”
2. R. W. Blakesley *et al.*, *Genome Res.* **14**, 2235 (2004).
3. K. Eilbeck *et al.*, *Genome Biol.* **6**, R44 (2005).
4. D. Field *et al.*, *Nat. Biotechnol.* **26**, 541 (2008).
5. Genomic Standards Consortium, <http://gensc.org>.
6. GOLD, Genomes OnLine Database, www.genomesonline.org.
7. K. Liolios *et al.*, *Nucleic Acids Res.* **36**, D475 (2008).
8. For Joint Genome Institute members, work was performed under the auspices of the U.S. Department of Energy Office of Science, Biological and Environmental Research Program; by the University of California, Lawrence Berkeley National Laboratory, contract no. DE-AC02-05CH11231; Lawrence Livermore National Laboratory contract no. DE-AC52-07NA27344; and Los Alamos National Laboratory, contract no. DE-AC02-06NA25396. Naval Medical Research Center work was supported by grant TMTI0068_07_NM_T from the Joint Science and Technology Office for Chemical and Biological Defense, Defense Threat Reduction Agency Initiative to T.D.R.

Supporting Online Material

www.sciencemag.org/cgi/content/full/326/5950/236/DC1

10.1126/science.1180614

Genome Project Standards in a New Era of Sequencing

P. S. G. Chain, D. V. Grafham, R. S. Fulton, M. G. FitzGerald, J. Hostetler, D. Muzny, J. Ali, B. Birren, D. C. Bruce, C. Buhay, J. R. Cole, Y. Ding, S. Dugan, D. Field, G. M. Garrity, R. Gibbs, T. Graves, C. S. Han, S. H. Harrison, S. Highlander, P. Hugenholtz, H. M. Khouiri, C. D. Kodira, E. Kolker, N. C. Kyrpides, D. Lang, A. Lapidus, S. A. Malfatti, V. Markowitz, T. Metha, K. E. Nelson, J. Parkhill, S. Pitluck, X. Qin, T. D. Read, J. Schmutz, S. Sozhamannan, P. Sterk, R. L. Strausberg, G. Sutton, N. R. Thomson, J. M. Tiedje, G. Weinstock, A. Wollam, Genomic Standards Consortium Human Microbiome Project Jumpstart Consortium and J. C. Detter

Science **326** (5950), 236-237.
DOI: 10.1126/science.1180614

ARTICLE TOOLS

<http://science.sciencemag.org/content/326/5950/236>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2009/10/08/326.5950.236.DC1>

REFERENCES

This article cites 4 articles, 1 of which you can access for free
<http://science.sciencemag.org/content/326/5950/236#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science* is a registered trademark of AAAS.