

$$p(A \mid B) = \frac{p(B \mid A)p(A)}{p(B)}$$

$$p(A | B, C) = \frac{p(B | A, C)p(A | C)}{p(B | C)}$$

$$p(\theta | D, M) = \frac{p(D | \theta, M)p(\theta | M)}{p(D | M)}$$

$$\text{Posterior} = \frac{\text{Likelihood} \quad \text{Prior}}{\text{Evidence}}$$

What we design:

$$p(D \mid \theta, M), p(\theta \mid M)$$

What we design:

$$p(D \mid \theta, M), p(\theta \mid M)$$

What we want:

$$p(\theta \mid D, M), p(D \mid M)$$

What we design:

$$p(D \mid \theta, M), p(\theta \mid M)$$

What we want:


$$p(\theta \mid D, M), p(D \mid M)$$

It is straightforward to compute $p(D \mid \theta, M)p(\theta \mid M)$ anywhere in $\{\theta\}$, but that doesn't give us much insight

$p(\theta | D, M)$ is a probability density function


$p(\theta | D, M)$ is a probability density function

One way to get insight into a PDF is by computing moments


$$\text{e.g., } \langle \theta \rangle = \int \theta \, p(\theta | D, M) \, d\theta$$


$p(\theta | D, M)$ is a probability density function

One way to get insight into a PDF is by computing moments

$$\text{e.g., } \langle \theta \rangle = \int \theta \, p(\theta | D, M) \, d\theta$$


Even if we wanted to look at just one parameter

$$p(\theta^i | D) = \int p(\theta | D, M) \, d\theta^j d\theta^k d\theta^l \dots$$


But if somebody gave you a set of θ values drawn from $p(\theta^i | D, M)$

$$\{\theta^i\} \rightarrow \{\theta_0^i, \theta_1^i, \theta_2^i, \dots, \theta_N^i\}$$

But if somebody gave you a set of θ values drawn from $p(\theta^i | D, M)$

$$\{\theta^i\} \rightarrow \{\theta_0^i, \theta_1^i, \theta_2^i, \dots, \theta_N^i\}$$

we can compute moments etc. the old fashioned way

$$\langle \theta^i \rangle \simeq \frac{\sum_n \theta_n^i}{N} \quad \text{☺}$$

We want a procedure for generating independent identically distributed (iid) samples from $p(\theta | D, M)$

We want a procedure for generating independent identically distributed (iid) samples from $p(\theta | D, M)$

i.e., a random number generator for $p(\theta | D, M)$

`numpy.random.normal()` returns i.i.d. samples from $N[0,1]$

`numpy.random.uniform()` returns i.i.d. samples from $U[0,1]$

We want a procedure for generating independent identically distributed (iid) samples from $p(\theta | D, M)$

i.e., a random number generator for $p(\theta | D, M)$

`numpy.random.normal()` returns i.i.d. samples from $N[0,1]$

`numpy.random.uniform()` returns i.i.d. samples from $U[0,1]$

Sets from RNGs are a subset of a more general class called Markovian processes

$$p(\theta_N | \theta_0, \theta_1, \dots, \theta_{N-1}) = p(\theta_N | \theta_{N-1})$$

Markov Chain Monte Carlo (MCMC):

Markov Chain Monte Carlo (MCMC):

A class of algorithms for generating (approximately) i.i.d. samples (i.e., a *Markov chain*) from some target distribution

The *Markov chain* must satisfy the **detailed balance** condition

$$p(y | x)p(x) = p(x | y)p(y)$$

The *Markov chain* must satisfy the **detailed balance** condition

$$p(y | x)p(x) = p(x | y)p(y)$$

This is just Bayes' theorem again!

The *Markov chain* must satisfy the **detailed balance** condition

$$p(y|x)p(x) = p(x|y)p(y) \quad \textit{This is just Bayes' theorem again!}$$

Probability of y following x in the Markov Chain. We don't know this.

The *Markov chain* must satisfy the **detailed balance** condition

$$\boxed{p(y|x)}p(x) = p(x|y)p(y) \quad \textit{This is just Bayes' theorem again!}$$

Probability of y following x in the Markov Chain. We don't know this.

Replace $p(y|x)$ with something we can calculate: $q(y|x)$

The *Markov chain* must satisfy the detailed balance condition

$$p(y|x)p(x) = p(x|y)p(y) \quad \text{This is just Bayes' theorem again!}$$

Probability of y following x in the Markov Chain. We don't know this.

Replace $p(y|x)$ with something we can calculate: $q(y|x)$

Proposal Distribution

e.g. $N[\mu, \sigma^2]$, $U[x_{\min}, x_{\max}]$

Our detailed balance equation becomes in inequality

$$q(y | x)p(x) > q(x | y)p(y)$$

Our detailed balance equation becomes an inequality

$$q(y | x)p(x) > q(x | y)p(y)$$

The chain transitions from x to y too often, and y to x too rarely

Our detailed balance equation becomes in inequality

$$q(y | x)p(x) \geq q(x | y)p(y)$$

The chain transitions from x to y too often, and y to x too rarely

To recover detailed balance, reweight by transition probability $\alpha \in [0,1]$

$$q(y | x)p(x)\alpha(x, y) = q(x | y)p(y)\alpha(y, x)$$

Our detailed balance equation becomes in inequality

$$q(y | x)p(x) > q(x | y)p(y)$$

The chain transitions from x to y too often, and y to x too rarely

To recover detailed balance, reweight by transition probability $\alpha \in [0,1]$

$$q(y | x)p(x)\alpha(x, y) = q(x | y)p(y)\alpha(y, x)$$

probability of transitioning from x to y


probability of transitioning from y to x

Our detailed balance equation becomes an inequality

$$q(y | x)p(x) > q(x | y)p(y)$$

The chain transitions from x to y too often, and y to x too rarely

To recover detailed balance, reweight by transition probability $\alpha \in [0,1]$

$$q(y | x)p(x)\alpha(x, y) = q(x | y)p(y)\alpha(y, x)$$


Our detailed balance equation becomes an inequality

$$q(y | x)p(x) > q(x | y)p(y)$$

The chain transitions from x to y too often, and y to x too rarely

To recover detailed balance, reweight by transition probability $\alpha \in [0,1]$

$$\alpha(x, y) = \frac{p(y)q(x | y)}{p(x)q(y | x)}$$

Our detailed balance equation becomes an inequality

$$q(y | x)p(x) > q(x | y)p(y)$$

The chain transitions from x to y too often, and y to x too rarely

To recover detailed balance, reweight by transition probability $\alpha \in [0,1]$

$$\alpha(x, y) = \frac{p(y)q(x | y)}{p(x)q(y | x)} \equiv H(x, y)$$

Our detailed balance equation becomes an inequality

$$q(y | x)p(x) > q(x | y)p(y)$$

The chain transitions from x to y too often, and y to x too rarely

To recover detailed balance, reweight by transition probability $\alpha \in [0,1]$

$$\alpha(x, y) = \frac{p(y)q(x | y)}{p(x)q(y | x)} \equiv H(x, y)$$

Hasting's Ratio

Now reverse the inequality

$$H(x, y) > 1 \quad \text{but} \quad \alpha(x, y) \in [0, 1]$$

Now reverse the inequality

$$H(x, y) > 1 \quad \text{but} \quad \alpha(x, y) \in [0, 1]$$

To generalize:

$$\alpha(x, y) = \min [1, H(x, y)]$$

Markov Chain Monte Carlo (MCMC):

- Start at some state x

Markov Chain Monte Carlo (MCMC):

- Start at some state x
- Propose new state y with $q(y | x)$

Markov Chain Monte Carlo (MCMC):

- Start at some state x
- Propose new state y with $q(y | x)$
- Accept with probability $\alpha(x, y) = \min [1, H(x, y)]$

Markov Chain Monte Carlo (MCMC):

- Start at some state x
- Propose new state y with $q(y | x)$
- Accept with probability $\alpha(x, y) = \min [1, H(x, y)]$
 - if ACCEPT state $\rightarrow y$
 - if REJECT state $\rightarrow x$

Markov Chain Monte Carlo (MCMC):

- Start at some state x
- Propose new state y with $q(y | x)$
- Accept with probability $\alpha(x, y) = \min [1, H(x, y)]$
 - if ACCEPT state $\rightarrow y$
 - if REJECT state $\rightarrow x$

Repeat



Markov Chain Monte Carlo (MCMC):

Now map to our inference problem:

State: Model parameters θ

Target Distribution: Posterior $p(\theta | D, M)$

Markov Chain Monte Carlo (MCMC):

Now map to our inference problem:

State: Model parameters θ

Target Distribution: Posterior $p(\theta | D, M)$

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

To gain insight, always
consider limiting cases:


$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

To gain insight, always consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#1. Symmetric proposal, uniform priors

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M)}{p(D | \theta, M)}$$




To gain insight, always consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#1. Symmetric proposal, uniform priors

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M)}{p(D | \theta, M)}$$

$p(D | \theta') > p(D | \theta)$ ACCEPT → Always go “uphill”



To gain insight, always consider limiting cases:


$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#1. Symmetric proposal, uniform priors

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M)}{p(D | \theta, M)}$$

$p(D | \theta') > p(D | \theta)$ ACCEPT → Always go “uphill”

$p(D | \theta'') < p(D | \theta')$ ROLL THE DICE → Go “downhill” w/
probability $\alpha(\theta', \theta'')$



To gain insight, always consider limiting cases:


$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#1. Symmetric proposal, uniform priors

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M)}{p(D | \theta, M)}$$

$p(D | \theta') > p(D | \theta)$ ACCEPT → Always go “uphill”

$p(D | \theta'') < p(D | \theta')$ ROLL THE DICE → Go “downhill” w/
probability $\alpha(\theta', \theta'')$



This guarantees that MCMCs *eventually* converge

To gain insight, always
consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#2. Proposal distribution matches target distribution

To gain insight, always
consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#2. Proposal distribution matches target distribution

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) q(\theta | \theta')}{p(D | \theta, M) q(\theta' | \theta)}, \quad q(\theta | \theta') = p(\theta | D)$$

To gain insight, always
consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#2. Proposal distribution matches target distribution

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) q(\theta | \theta')}{p(D | \theta, M) q(\theta' | \theta)}, \quad q(\theta | \theta') = p(\theta | D)$$

$$\alpha(\theta, \theta') = 1$$

Everything gets accepted!
Maximally efficient

To gain insight, always
consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#2. Proposal distribution matches target distribution

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) q(\theta | \theta')}{p(D | \theta, M) q(\theta' | \theta)}, \quad q(\theta | \theta') = p(\theta | D)$$

$$\alpha(\theta, \theta') = 1$$

Everything gets accepted!
Maximally efficient

**Designing good proposal distributions is where the work
hides**

To gain insight, always
consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#3. Hammer at the true value θ^{TRUTH}

To gain insight, always
consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#3. Hammer at the true value θ^{TRUTH}

$$q(\theta' | \theta) = \delta(\theta' - \theta^{\text{TRUTH}})$$

To gain insight, always
consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#3. Hammer at the true value θ^{TRUTH}

$$q(\theta' | \theta) = \delta(\theta' - \theta^{\text{TRUTH}})$$

$$\alpha(\theta, \theta^{\text{TRUTH}}) = \frac{p(\theta^{\text{TRUTH}} | D)}{p(\theta | D)} \frac{\delta(\theta - \theta^{\text{TRUTH}})}{\delta(0)} = 0?!?!?$$

To gain insight, always consider limiting cases:

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M) q(\theta | \theta')}{p(D | \theta, M) p(\theta | M) q(\theta' | \theta)}$$

#3. Hammer at the true value θ^{TRUTH}

$$q(\theta' | \theta) = \delta(\theta' - \theta^{\text{TRUTH}})$$

$$\alpha(\theta, \theta^{\text{TRUTH}}) = \frac{p(\theta^{\text{TRUTH}} | D)}{p(\theta | D)} \frac{\delta(\theta - \theta^{\text{TRUTH}})}{\delta(0)} = 0?!?!?$$

The proposal is a teenager: The harder you push the harder it pushes back.

$$\alpha(\theta, \theta') = \frac{p(D | \theta', M) p(\theta' | M)}{p(D | \theta, M) p(\theta | M)} \frac{q(\theta | \theta')}{q(\theta' | \theta)}$$

Always go uphill, but try
going down as well

Counterbalance how
hard you're pushing

INTERMISSION

What about $p(D | M)$?

$$p(D \mid M) = \int d\theta \, p(D \mid \theta, M) p(\theta \mid M)$$

$$p(D | M) = \int d\theta \, p(D | \theta, M) p(\theta | M)$$

Large-dimensional
integral == HARD

Recast as 1D integral

$$Z = \int L \, dX$$

Evidence \swarrow Z \searrow $dX = p(\theta | M) d\theta$
Likelihood \downarrow Prior volume element

Recast as 1D integral

$$Z = \int L \, dX$$

Evidence \swarrow Likelihood \downarrow $dX = p(\theta | M) d\theta$ \searrow Prior volume element

$$X \equiv X(\lambda) = \int_{L(\theta) > \lambda} p(\theta | M) d\theta$$

Recast as 1D integral

$$Z = \int L \, dX$$

Evidence \swarrow Likelihood \downarrow $dX = p(\theta | M) d\theta$ \swarrow Prior volume element

$$X \equiv X(\lambda) = \int_{L(\theta) > \lambda} p(\theta | M) d\theta$$

\swarrow Cumulant prior mass covering all likelihood values $> \lambda$


As λ increases, X goes from $1 \rightarrow 0$

λ is small: likelihood is low \rightarrow prior



so,
$$Z = \int_0^1 L(X) dX$$

λ is large: likelihood is high



$L(X)$ is positive and monotonically decreasing. We can do this.





$L(X)$ is positive and monotonically decreasing. We can do this.





$L(X)$ is positive and monotonically decreasing. We can do this.



$L(X)$ is positive and monotonically decreasing. We can do this.



$L(X)$ is positive and monotonically decreasing. We can do this.



Nested Sampling:

- Start w/ N points sampling prior $\theta_1, \dots, \theta_N$

$$Z = 0, X_0 = 1, i = 1$$

Nested Sampling:

- Start w/ N points sampling prior $\theta_1, \dots, \theta_N$

$$Z = 0, X_0 = 1, i = 1$$

- Record the lowest $L(\theta_n)$ as L_i

Nested Sampling:

- Start w/ N points sampling prior $\theta_1, \dots, \theta_N$

$$Z = 0, X_0 = 1, i = 1$$

- Record the lowest $L(\theta_n)$ as L_i
- $X_i \sim e^{-i/N}, Z = Z + L_i(X_{i-1} - X_i)$

Nested Sampling:

- Start w/ N points sampling prior $\theta_1, \dots, \theta_N$

$$Z = 0, X_0 = 1, i = 1$$

- Record the lowest $L(\theta_n)$ as L_i
- $X_i \sim e^{-i/N}$, $Z = Z + L_i(X_{i-1} - X_i)$
- Draw new θ_N from prior constrained by $L(\theta_n) > L_i$

Nested Sampling:

- Start w/ N points sampling prior $\theta_1, \dots, \theta_N$


$$Z = 0, X_0 = 1, i = 1$$

- Record the lowest $L(\theta_n)$ as L_i

- $X_i \sim e^{-i/N}, Z = Z + L_i(X_{i-1} - X_i)$

- Draw new θ_N from prior constrained by $L(\theta_n) > L_i$

Repeat
 $i = i + 1, i < i_{\max}$




Nested Sampling:

- Start w/ N points sampling prior $\theta_1, \dots, \theta_N$

$$Z = 0, X_0 = 1, i = 1$$

- Record the lowest $L(\theta_n)$ as L_i
- $X_i \sim e^{-i/N}, Z = Z + L_i(X_{i-1} - X_i)$
- Draw new θ_N from prior constrained by $L(\theta_n) > L_i$

Repeat
 $i = i + 1, i < i_{\max}$



- $Z = Z + \sum_n L(\theta_n) X_i / N$

Nested Sampling:

Generate posterior samples by reweighting $\{\theta_n\}$

$$p_i = L_i(X_{i-1} - X_i)/Z$$

ENCORE

Other evidence calculators/estimates

Other sampling algorithms