

Module Todo

Title Todo

Author:

John-Frederik Reeg

Student ID: 8108244

Submission Date: Todo

Supervisor: Prof. Dr. Christian Fischer

JUSTUS LIEBIG UNIVERSITY GIESSEN
INSTITUTE FOR THEORETICAL PHYSICS

Contents

1	Introduction	5
2	Theory	7
2.1	Herleitung Esther	7
2.2	Herleitung Stephan	10
3	Pion BSE	13
4	Conclusion and Outlook	15
A	Mathematical Relations	17
A.1	Dirac algebra	17
A.1.1	Minkowski representation	17
A.1.2	Euclidean representation	18
A.2	Hyperspherical Coordinates	19
A.3	Dirac Spinors in Euclidean Space	19
B	Kinematics	21
B.1	Breit System	21
B.1.1	Pion-Nucleon-Nucleon Coupling	21
B.1.2	Kaon-Nucleon-Lambda Coupling	21
B.2	in the Center of Mass System	22

Chapter 1

Introduction

Theory

$$\text{---}\bullet\text{---}^{-1} = \text{---}^{-1} - \text{---}\overset{\text{---}\bullet\text{---}}{\text{---}} \quad (2.1)$$

2.1 Herleitung Esther

$$G^{(n)}[\phi] = \langle \Omega | T \{ \phi(x_1) \phi(x_2) \dots \phi(x_n) \} | \Omega \rangle \quad (2.2)$$

$$\langle \Omega | T \phi(x) \phi(y) | \Omega \rangle \equiv \begin{cases} \langle \Omega | \phi(x) \phi(y) | \Omega \rangle, & \text{for } x^0 > y^0 \\ \langle \Omega | \phi(y) \phi(x) | \Omega \rangle, & \text{for } y^0 > x^0 \end{cases} \quad (2.3)$$

$$G^{(n)}[\phi] = \frac{\int \mathcal{D}\phi \, \phi(x_1) \dots \phi(x_n) e^{-S[\phi]}}{\int \mathcal{D}\phi \, e^{-S[\phi]}} \quad (2.4)$$

$$Z[J] = \int \mathcal{D}\phi \, e^{-S[\phi] + J\phi} \quad (2.5)$$

$$G[\phi] = \langle \Omega | T G[\phi] | \Omega \rangle = \left[G \left[\frac{\delta}{\delta J} \right] \frac{Z[J]}{Z[0]} \right]_{J=0} \quad (2.6)$$

$$Z[J] = e^{-W[J]} = e^{-(\Gamma[\tilde{\phi}] - \int_x \tilde{\phi} J(x))}, \quad \Gamma[\tilde{\phi}] = W[J] - \tilde{\phi} J \quad (2.7)$$

$$\tilde{\phi}(x) = -\frac{\delta W[J]}{\delta J(x)} = \frac{1}{Z[J]} \frac{\delta}{\delta J(x)} Z[J] = \langle \phi(x) \rangle \quad (2.8)$$

$$S_{\text{QCD}} = \int d^4x \left(\bar{\psi}(\not{D} + m)\psi + \frac{1}{4}F_{\mu\nu}^a F_a^{\mu\nu} \right) \quad (2.9)$$

$$\psi' = U\psi, \quad A'_\mu = UA_\mu U^\dagger - \frac{i}{g}U\partial_\mu U^\dagger, \quad F'_{\mu\nu} = UF_{\mu\nu}U^\dagger \quad (2.10)$$

$$\frac{\delta S[\phi]}{\delta \phi} = 0 \quad (2.11)$$

$$Z[J] = \int \mathcal{D}\phi \, e^{-S[\phi] + J\phi} \quad (2.12)$$

$$\int \mathcal{D}\phi \, e^{-S[\phi] + J\phi} = \int \mathcal{D}\phi \, e^{-S[\phi + \delta\phi] + J(\phi + \delta\phi)} \quad (2.13)$$

$$\left\langle \frac{\delta S[\phi]}{\delta \phi(x)} \right\rangle_J = J(x) \quad (2.14)$$

$$\langle f(\phi) \rangle_J \equiv Z[J]^{-1} \int \mathcal{D}\phi \, f(\phi) e^{-S[\phi] + J\phi} = Z[J]^{-1} f\left(\frac{\delta}{\delta J}\right) Z[J] \quad (2.15)$$

$$\left\langle \frac{\delta S[\phi]}{\delta \phi(x)} \right\rangle = Z[J]^{-1} \frac{\delta S}{\delta \phi} \left[\frac{\delta}{\delta J(x)} \right] Z[J] = J(x) \quad (2.16)$$

$$\frac{\delta \Gamma}{\delta \tilde{\phi}(x)} = \frac{\delta S}{\delta \phi} \left[\tilde{\phi}(x) + \int_y \frac{\delta^2 \Gamma}{\delta \tilde{\phi}(x) \delta \tilde{\phi}(y)} \frac{\delta}{\delta \tilde{\phi}(y)} \right] \quad (2.17)$$

$$S^{-1} = S_0^{-1} + \int g \gamma_\mu D^{\mu\nu} S \Gamma_\nu \quad (2.18)$$

$$S(p) = G^{(2)}(p) = \int d^4x \, e^{-ipx} \langle \Omega | T \bar{\psi}(x) \psi(0) | \Omega \rangle \quad (2.19)$$

$$S_{\alpha\beta abAB}(p) = S_{\alpha\beta}(p) \otimes \delta^{ab} \otimes \delta^{AB} \quad (2.20)$$

$$S_{\alpha\beta}(p) = -i\sigma_v(p)\not{p} + \sigma_s(p)\mathbb{1}_D \quad (2.21)$$

$$S^{-1}(p) = i\not{p}A(p^2) + B(p^2) \quad (2.22)$$

$$S^{-1}(p^2) = S_0^{-1}(p^2) + \int d^4q D_{ab}^{\mu\nu}(p-q) \left(g \frac{\lambda^a}{2} \gamma_\mu \right) S(q^2) (g\Gamma_\nu^b(p, q)) \quad (2.23)$$

$$m_0(\Lambda^2) = Z_m(\mu^2, \Lambda^2) m(\mu^2) \quad (2.24)$$

$$S(p^2, \Lambda^2) = Z_2(\mu^2, \Lambda^2) S_R(p^2, \mu^2) \quad (2.25)$$

$$A^{-1}(p^2, \Lambda^2) = Z_2(\mu^2, \Lambda^2) A_R^{-1}(p^2, \mu^2) \quad (2.26)$$

$$Z_{1F} = \frac{Z_2}{\tilde{Z}_3}, \quad Z_g \tilde{Z}_3 Z_3^{1/2} = 1 \quad (2.27)$$

$$S^{-1}(p^2, \mu^2) = Z_2(i\not{p} + Z_m m) \quad (2.28)$$

$$+ Z_g^2 g^2 \int d^4q Z_3 D_{ab}^{\mu\nu}(p-q, \mu^2) \left(\frac{\lambda^a}{2} \gamma_\mu \right) Z_2^2 S(q^2, \mu^2) (Z_{1F}^{-1} \Gamma_\nu^b(p, q, \mu^2)) \quad (2.29)$$

$$g(\mu^2) = Z_g^{-1}(\mu^2, \Lambda^2) g(\Lambda^2) \quad (2.30)$$

$$Z_g^2 Z_3 Z_2^2 \frac{\tilde{Z}_3}{Z_2} = \frac{Z_2}{\tilde{Z}_3} \cdot [Z_g^2 \tilde{Z}_3^2 Z_3] = \frac{Z_2}{\tilde{Z}_3} = Z_{1F} \quad (2.31)$$

$$S^{-1}(p^2, \mu^2) = Z_2(i\not{p} + Z_m m) \quad (2.32)$$

$$+ \frac{Z_2}{\tilde{Z}_3} g^2 \int d^4q D_{ab}^{\mu\nu}(p-q, \mu^2) \left(\frac{\lambda^a}{2} \gamma_\mu \right) S(q^2, \mu^2) (\Gamma_\nu^b(p, q, \mu^2)) \quad (2.33)$$

$$A(\mu^2) \stackrel{!}{=} 1, \quad B(\mu^2) \stackrel{!}{=} m(\mu^2) \quad (2.34)$$

$$Z_2 = 1 - \Sigma_A(\mu^2, \Lambda^2), \quad Z_4 = 1 - \frac{\Sigma_B(\mu^2, \Lambda^2)}{m(\mu^2)} \quad (2.35)$$

$$\frac{Z_2}{Z_3} g^2 D_{ab}^{\mu\nu}(p-q) \Gamma_\nu^b(p, q) \rightarrow Z_2^2 \mathcal{G}((p-q)^2) \frac{t^{\mu\nu}(p-q)}{(p-q)^2} \frac{\lambda^a}{2} \gamma_\nu \quad (2.36)$$

$$t^{\mu\nu}(k) = \delta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2}$$

$$\frac{\mathcal{G}(k^2)}{4\pi} = \alpha(k^2) = \frac{D}{\omega^6} \pi k^4 e^{-\frac{k^2}{\omega^2}} + \frac{2\pi\gamma_m (1 - e^{-k^2})}{\ln \left[e^2 - 1 + \left(1 + \frac{k^2}{\Lambda_{\text{QCD}}^2} \right)^2 \right]} \quad (2.37)$$

$$\frac{\alpha(k^2)}{k^2} \rightarrow \frac{\alpha(k^2)}{k^2} \left(\frac{\Lambda_{\text{PV}}^2}{k^2 + \Lambda_{\text{PV}}^2} \right) \quad (2.38)$$

$$A(p^2) = Z_2 + \frac{16\pi}{3} Z_2^2 \int \mathrm{d}^4 q \frac{A(q^2)}{q^2 A^2(q^2) + B^2(q^2)} \left[(pq) + 2 \frac{(pk)(qk)}{k^2} \right] \frac{\alpha(k^2)}{k^2} \frac{\Lambda_{\text{PV}}^2}{\Lambda_{\text{PV}}^2 + k^2} \quad (2.39)$$

$$B(p^2) = Z_4 m + 16\pi Z_2^2 \int \mathrm{d}^4 q \frac{B(q^2)}{q^2 A^2(q^2) + B^2(q^2)} \frac{\alpha(k^2)}{k^2} \frac{\Lambda_{\text{PV}}^2}{\Lambda_{\text{PV}}^2 + k^2} \quad (2.40)$$

$$\alpha(k^2) = \alpha_{\text{UV}}(k^2) + \alpha_{\text{IR}}(k^2) \quad (2.41)$$

$$\alpha_{\text{IR}}(k^2) = \frac{D}{\omega^6} \pi k^4 e^{-\frac{k^2}{\omega^2}} \quad (2.42)$$

$$\alpha_{\text{UV}}(k^2) = \frac{2\pi\gamma_m (1 - e^{-k^2})}{\ln \left[e^2 - 1 + \left(1 + \frac{k^2}{\Lambda_{\text{QCD}}^2} \right)^2 \right]} \quad (2.43)$$

2.2 Herleitung Stephan

$$S_{\text{QCD}} = \int \mathrm{d}^4 x \left[\sum_{i=1}^{N_f} \bar{\psi}_i (\not{D} + m_i) \psi_i + \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu} \right] \quad (2.44)$$

$$S^{-1}(p) = Z_2 S_0^{-1}(p) - \Sigma(p) \quad (2.45)$$

$$\Sigma(p) = -Z_{1F}(\mu^2, \Lambda^2) g^2 C_f \int \mathrm{d}^4 q \gamma_\mu S(q) \Gamma_\nu(p, q) D^{\mu\nu}(p-q) \quad (2.46)$$

$$D^{\mu\nu}(k) = T^{\mu\nu}(k) \frac{Z(k^2)}{k^2}, \quad T^{\mu\nu}(k) = \delta^{\mu\nu} - k^\mu k^\nu / k^2 \quad (2.47)$$

$$\Gamma_{\text{RL}}^\mu(p, q) = Z_{1F} \gamma^\mu \lambda_1(k^2) \quad (2.48)$$

$$\Sigma(p) = -C_f Z_{1F}^2 g^2 \int \mathrm{d}^4 q \, \gamma_\mu S(q) \gamma_\nu T^{\mu\nu} \frac{\lambda_1(k^2) Z(k^2)}{k^2} \quad (2.49)$$

$$\alpha(k^2) := \left(\frac{\tilde{Z}_1}{\tilde{Z}_3} \right)^2 \frac{g^2}{4\pi} Z(k^2) \lambda_1(k^2) \quad (2.50)$$

$$Z_{1F}^2 = Z_2^2 \left(\frac{\tilde{Z}_1}{\tilde{Z}_3} \right)^2 \quad (2.51)$$

$$\Sigma(p) = -4\pi C_f Z_2^2 \int \mathrm{d}^4 q \, \gamma_\mu S(q) \gamma_\nu T^{\mu\nu} \frac{\alpha(k^2)}{k^2} \quad (2.52)$$

$$S^{-1}(p) = Z_2 (\mathrm{i}\not{p} + Z_m m_c) + 4\pi C_f Z_2^2 \int \mathrm{d}^4 q \, \gamma_\mu S(q) \gamma_\nu T^{\mu\nu} \frac{\alpha(k^2)}{k^2} \quad (2.53)$$

$$C_f = \frac{4}{3}$$

$$S^{-1}(p) = \mathrm{i}\not{p} A(p^2) + B(p^2) \quad (2.54)$$

$$A(p^2) = Z_2 + \Sigma_A(p^2) \quad (2.55)$$

$$B(p^2) = Z_2 Z_m m_c + \Sigma_B(p^2) \quad (2.56)$$

$$\Sigma_A(p^2) = \frac{16\pi}{3} Z_2^2 \int \mathrm{d}^4 q \, \frac{A(q^2)}{q^2 A^2(q^2) + B^2(q^2)} \left[(pq) + 2 \frac{(pk)(qk)}{k^2} \right] \frac{\alpha(k^2)}{k^2} \frac{\Lambda_{\text{PV}}^2}{\Lambda_{\text{PV}}^2 + k^2} \quad (2.57)$$

$$\Sigma_B(p^2) = 16\pi Z_2^2 \int \mathrm{d}^4 q \, \frac{B(q^2)}{q^2 A^2(q^2) + B^2(q^2)} \frac{\alpha(k^2)}{k^2} \frac{\Lambda_{\text{PV}}^2}{\Lambda_{\text{PV}}^2 + k^2} \quad (2.58)$$

$$\alpha(k^2) = \alpha_{\text{UV}}(k^2) + \alpha_{\text{IR}}(k^2) \quad (2.59)$$

$$\alpha_{\text{IR}}(k^2) = \frac{D}{\omega^6} \pi k^4 e^{-\frac{k^2}{\omega^2}} \quad (2.60)$$

$$\alpha_{\text{UV}}(k^2) = \frac{2\pi\gamma_m (1 - e^{-k^2})}{\ln \left[e^2 - 1 + \left(1 + k^2/\Lambda_{\text{QCD}}^2 \right)^2 \right]} \quad (2.61)$$




Figure 2.1

Chapter 3

Pion BSE

$$\Gamma_{\text{PS}}(p, P) = \gamma^5 \left(E(p, P) - i \not{P} F(p, P) - i(p \cdot P) \not{p} G(p, P) + [\not{p}, \not{P}]_- H(p, P) \right) \quad (3.1)$$

$$\Gamma(p, P) = -C_f Z_2^2 4\pi \int \mathrm{d}^4 q \gamma^\mu S(q_+) \Gamma(q, P) S(q_-) \gamma^\nu T_{\mu\nu}(k) \frac{\alpha(k^2)}{k^2} \quad (3.2)$$

Inserting our ansatz, applying projector p_E , and taking the Dirac trace, we get:

$$\begin{aligned} E(p, P) = \text{tr} \left[-C_f Z_2^2 4\pi \frac{1}{4} \gamma^5 \int \mathrm{d}^4 q \gamma^\mu S(q_+) \gamma^5 (E(q, P) - i \not{P} F(q, P) \right. \\ \left. - i(q \cdot P) \not{q} G(q, P) + [\not{q}, \not{P}]_- H(q, P)) S(q_-) \gamma^\nu T_{\mu\nu}(k) \frac{\alpha(k^2)}{k^2} \right] \end{aligned} \quad (3.3)$$

$$\begin{aligned} = -C_f Z_2^2 \pi \int \mathrm{d}^4 q \text{tr} \left[\gamma^5 \gamma^\mu S(q_+) \gamma^5 (E(q, P) - i \not{P} F(q, P) \right. \\ \left. - i(q \cdot P) \not{q} G(q, P) + [\not{q}, \not{P}]_- H(q, P)) S(q_-) \gamma^\nu T_{\mu\nu}(k) \right] \frac{\alpha(k^2)}{k^2} \end{aligned} \quad (3.4)$$

Observing the alone-standing trace:

$$\text{tr} \left[\gamma^5 \gamma^\mu \frac{i \not{q}_+ A(q_+^2) + B(q_+^2)}{q_+^2 A^2(q_+^2) + B^2(q_+^2)} \gamma^5 (E(q, P) - i \not{P} F(q, P) - i(q \cdot P) \not{q} G(q, P) + [\not{q}, \not{P}]_- H(q, P)) \frac{i \not{q}_- A(q_-^2) + B(q_-^2)}{q_-^2 A^2(q_-^2) + B^2(q_-^2)} \gamma^\nu T_{\mu\nu}(k) \right] \quad (3.5)$$

$$\begin{pmatrix} E(p) \\ F(p) \\ G(p) \\ H(p) \end{pmatrix} = \begin{pmatrix} \kappa_{EE}(p, q) & \kappa_{EF}(p, q) & \kappa_{EG}(p, q) & \kappa_{EH}(p, q) \\ \kappa_{FE}(p, q) & \kappa_{FF}(p, q) & \kappa_{FG}(p, q) & \kappa_{FH}(p, q) \\ \kappa_{GE}(p, q) & \kappa_{GF}(p, q) & \kappa_{GG}(p, q) & \kappa_{GH}(p, q) \\ \kappa_{HE}(p, q) & \kappa_{HF}(p, q) & \kappa_{HG}(p, q) & \kappa_{HH}(p, q) \end{pmatrix} \begin{pmatrix} E(q) \\ F(q) \\ G(q) \\ H(q) \end{pmatrix} \quad (3.6)$$

$$\lambda(P^2) \begin{pmatrix} E \\ F \\ G \\ H \end{pmatrix} (p, P^2) = \mathcal{K}(p, q, P^2) \begin{pmatrix} E \\ F \\ G \\ H \end{pmatrix} (p, P^2) \quad (3.7)$$

$$\lambda(-m_\pi^2) \stackrel{!}{=} 1 \quad (3.8)$$

Projectors:

$$p_E = \frac{1}{4} \gamma^5 \quad (3.9)$$

$$p_F = \frac{i}{4((p \cdot P)^2 - p^2 P^2)} \gamma^5 (p^2 \not{P} - (p \cdot P) \not{p}) \quad (3.10)$$

$$p_G = \frac{i}{4((p \cdot P)^2 - p^2 P^2)} \gamma^5 \left(\frac{P^2}{p \cdot P} \not{p} - \not{P} \right) \quad (3.11)$$

$$p_H = \frac{1}{16((p \cdot P)^2 - p^2 P^2)} \gamma^5 [\not{p}, \not{P}]_- \quad (3.12)$$

Chapter 4

Conclusion and Outlook

Appendix A

Mathematical Relations

A.1 Dirac algebra

The Dirac algebra is the algebra of the gamma matrices γ^μ used in the Dirac equation. It is defined by the anticommutation relation

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} \mathbf{1}, \quad (\text{A.1})$$

where $\eta^{\mu\nu}$ is the defining metric of the space for which the gamma matrices are formulated, or the signature of the dot product of two four vectors. This ensures the Lorentz covariance of the equation. The gamma matrices are represented by four 4×4 matrices and generate the Clifford Algebra $\text{Cl}_{1,3}(\mathbb{C})$. It is also customary to define a fifth matrix, γ^5 , which satisfies the following relations:

$$\{\gamma^5, \gamma^\mu\} = 0, \quad \gamma^5 = (\gamma^5)^\dagger, \quad (\gamma^5)^2 = \mathbf{1} \quad (\text{A.2})$$

Independent of the representation, we define the Feynman-Slash operator to be:

$$\not{p} = \gamma_\mu p^\mu$$

A.1.1 Minkowski representation

The metric of Minkowski space is given by the signature $(+, -, -, -)$, written as a matrix:

$$g_{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad (\text{A.3})$$

One representation of the gamma matrices that satisfies the Dirac algebra with this metric is

$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma^n = \begin{pmatrix} 0 & \sigma_n \\ -\sigma_n & 0 \end{pmatrix}, \quad (\text{A.4})$$

where every 0 represents an empty 2×2 matrix and every 1 represents the 2×2 identity matrix. The σ_n serve as placeholders for the three Pauli matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (\text{A.5})$$

The fifth gamma matrix can now be calculated using the equation:

$$\gamma^5 = \gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (\text{A.6})$$

Furthermore we make use of the charge conjugation operator C , which possesses properties independent of the representation in which it is given, namely:

$$C^{-1}\gamma_\mu C = -\gamma_\mu^T, \quad C\gamma_5 C^{-1} = \gamma_5^T \quad (\text{A.7})$$

In Minkowski space, it is represented by the matrix

$$C = -C^T = -C^\dagger = \begin{pmatrix} 0 & i\sigma_2 \\ i\sigma_2 & 0 \end{pmatrix}. \quad (\text{A.8})$$

A.1.2 Euclidean representation

When working in Euclidean space, it is customary to switch the indices 0 and 4. This also serves to differentiate clearly between four vectors in both representations. The space-like components keep the indices 1 through 3. Since the defining metric in Euclidean space is just the identity, the new gamma matrices need to satisfy

$$\{\gamma^\mu, \gamma^\nu\} = 2\delta^{\mu\nu} \mathbf{1}, \quad (\text{A.9})$$

$$\{\gamma^5, \gamma^\mu\} = 0. \quad (\text{A.10})$$

They can be computed with

$$\gamma_{n,E} = -i\gamma_{n,M}, \quad (\text{A.11})$$

$$\gamma_{4,E} = \gamma_{0,M}, \quad (\text{A.12})$$

where $n \in \{1, 2, 3\}$. These new representations now satisfy the anticommutation relations for this Euclidean signature.

Regarding γ_5 and C , both are unchanged by this new metric:

$$\gamma_{5,E} = -\gamma_{1,E}\gamma_{2,E}\gamma_{3,E}\gamma_{4,E} = \gamma_{5,M} \quad (\text{A.13})$$

$$C_E = C_M = \gamma_{2,E}\gamma_{4,E} \quad (\text{A.14})$$

In this work the Euclidean metric is used primarily. Due to this, the index E will be dropped and it is assumed that every gamma matrix is used in its Euclidean representation unless stated otherwise.

One very important note is that when working with the Euclidean metric as defined above, the invariant square of a four vector is opposite in sign to its Minkowski counterpart. To counteract this change, all four vector products in Minkowski space need to change their sign when transferred to Euclidean space:

$$p_E^\mu q_{\mu,E} = -p_M^\mu q_{\mu,M} \quad (\text{A.15})$$

A.2 Hyperspherical Coordinates

Over the course of this work it often becomes necessary to integrate over four dimensions. To facilitate this we utilize hyperspherical coordinates, with their general position vector being given by

$$p = |p| \begin{pmatrix} \cos \phi \sin \theta \sin \psi \\ \sin \phi \sin \theta \sin \psi \\ \cos \theta \sin \psi \\ \cos \psi \end{pmatrix}, \quad (\text{A.16})$$

where $|p|$ represents the magnitude of the vector, and θ , ϕ , and ψ are the angles parameterizing its direction.

To integrate over this coordinate system, we calculate the Jacobian and get:

$$\int d^4p = \int_0^\infty dp p^3 \int_0^{2\pi} d\phi \int_0^\pi d\theta \sin \theta \int_0^\pi d\psi \sin^2 \psi \quad (\text{A.17})$$

Again we keep in mind the translation from Minkowski to Euclidean space, which necessitates us to bring in another factor i .

$$\int d^4p_M \rightarrow i \int d^4p_E \quad (\text{A.18})$$

A.3 Dirac Spinors in Euclidean Space

The solutions of the free Dirac equation are called the Dirac spinors. In Euclidean space, their representations were extracted from reference

$$u_1(P) = \sqrt{\frac{E+M}{2M}} \begin{pmatrix} 1 \\ 0 \\ \frac{P_z}{E+M} \\ \frac{P_x+iP_y}{E+M} \end{pmatrix}, \quad (\text{A.19})$$

$$u_2(P) = \sqrt{\frac{E+M}{2M}} \begin{pmatrix} 0 \\ 1 \\ \frac{P_x-iP_y}{E+M} \\ \frac{-P_z}{E+M} \end{pmatrix}, \quad (\text{A.20})$$

$$v_1(P) = \sqrt{\frac{|E|+M}{2M}} \begin{pmatrix} \frac{-P_z}{|E|+M} \\ \frac{-(P_x+iP_y)}{|E|+M} \\ 1 \\ 0 \end{pmatrix}, \quad (\text{A.21})$$

$$v_2(P) = \sqrt{\frac{|E|+M}{2M}} \begin{pmatrix} \frac{-(P_x-iP_y)}{|E|+M} \\ \frac{P_z}{|E|+M} \\ 0 \\ 1 \end{pmatrix}, \quad (\text{A.22})$$

for a given four momentum of the form $P = (P_x, P_y, P_z, iE)$. They are normalized with the conditions:

$$u_r^\dagger(P) u_{r'}(P) = \frac{|E|}{M} \delta_{r,r'} \quad (\text{A.23})$$

$$u_r^\dagger(P) = u_r^*(P) \gamma_4 \quad (\text{A.24})$$

Appendix B

Kinematics

This chapter seeks to visualize the kinematics of the reactions taking place. It serves to explain conventions and chosen reference frames, while also deriving the momenta used in the work above.

B.1 Breit System

The Breit system is a frame of reference often used to describe scattering processes. In it, we demand that the current coupling with a particle transfers only spatial momentum, so the transferred energy Q_4 is zero. Also known as the "brick wall frame" or "infinite momentum frame", the collision can therefore be understood as an ideal reflection on an outside particle, at least if the scattered particle does not change its mass. In the context of this work we use the Breit system to describe the coupling of a meson to a baryon, so for the calculation of the form factors

B.1.1 Pion-Nucleon-Nucleon Coupling

To calculate the kinematics of a pion-nucleon coupling in the Breit frame, we choose our coordinates so that all x and y momenta are 0:

$$Q = Q_3 \hat{e}_3, \quad P_{i,3} = -P_{f,3} \quad (\text{B.1})$$

Considering momentum conservation results in the following four momenta:

$$Q = (0, 0, Q_3, 0) \quad (\text{B.2})$$

$$P_i = \left(0, 0, -\frac{1}{2}Q_3, i\sqrt{M_p^2 + \frac{1}{4}Q_3^2}\right) \quad (\text{B.3})$$

$$P_f = \left(0, 0, +\frac{1}{2}Q_3, i\sqrt{M_p^2 + \frac{1}{4}Q_3^2}\right) \quad (\text{B.4})$$

B.1.2 Kaon-Nucleon-Lambda Coupling

The result from above cannot simply be copied, unfortunately. In the case of nucleon-kaon absorption, the resulting lambda baryon is heavier than the incoming proton. For this to be possible without energy transfer by the kaon, the spatial momentum of the outgoing particle

needs to be different in magnitude than its incoming counterpart. Again working only in z -direction and taking both energy and momentum conservation into account, we formulate the following four momenta, parameterized by the momentum transfer Q .

$$Q = (0, 0, Q_3, 0) \quad (\text{B.5})$$

$$P_i = \left(0, 0, \frac{M_p^2 - M_\Lambda^2 - Q_3^2}{2Q_3}, i \sqrt{M_p^2 + \left(\frac{M_p^2 - M_\Lambda^2 - Q_3^2}{2Q_3} \right)^2} \right) \quad (\text{B.6})$$

$$P_f = \left(0, 0, \frac{M_p^2 - M_\Lambda^2 + Q_3^2}{2Q_3}, i \sqrt{M_\Lambda^2 + \left(\frac{M_p^2 - M_\Lambda^2 + Q_3^2}{2Q_3} \right)^2} \right) \quad (\text{B.7})$$

Although the validity of this approach seems to be sufficiently correct at first, the kinematics break down for small momentum transfers. If the supplied momentum by the exchange meson does not suffice to create a lambda baryon, the calculation becomes problematic. Therefore we solve for the condition

$$M_\Lambda^2 \leq E^2 = M_\Lambda^2 + \left(\frac{M_p^2 - M_\Lambda^2 + Q_3^2}{2Q_3} \right)^2, \quad (\text{B.8})$$

which leaves us with a lower bound for the calculation of

$$Q^2 \geq M_\Lambda^2 - M_p^2 =: \Lambda_{\text{BF}}^2 \approx 0.4 \text{ GeV}^2 \quad (\text{B.9})$$

Inserting a Q^2 below this threshold Λ_{BF}^2 will yield nonphysical results, a problem discussed further in.

B.2 in the Center of Mass System

When analyzing moving objects, it is convenient to work in a resting frame of reference, the center of mass system (CM system). Its convenience stems from the property that at any given time the sum of all space-like momenta is 0:

$$\sum_i \mathbf{p}_i = 0 \quad (\text{B.10})$$

In order to describe the kinematics of the reaction we will call the four momenta of all participating particles P , $\bar{P}$, Λ , $\bar{\Lambda}$, and K . Now we can choose our coordinate system so that the space-like momentum of Λ and $\bar{\Lambda}$ lays in the x -axis. The total energy E_{cm} of each baryon as well as the magnitude of the space-like momentum of the protons can easily be calculated, since all baryons involved in this reaction are on-shell:

$$E_{cm} = \sqrt{\Lambda_x^2 + M_\Lambda^2} \quad (\text{B.11})$$

$$|\mathbf{p}_{cm}| = \sqrt{E_{cm}^2 - M_p^2} \quad (\text{B.12})$$

With this, we can outline the kinematic situation of the reaction, where the angle θ is a new degree of freedom granted by the change in direction before/after the interaction. Without losing generality, we can restrict the reaction to the x - y -plane.




Figure B.1: space-like momenta of the reaction

Using this figure, we write out the four momenta of each hadron involved. Due to conservation of energy and momentum, the kaon is bound to the condition $K = \Lambda - P = \bar{P} - \bar{\Lambda}$.

$$P = (-|\mathbf{p}_{cm}| \cos \theta, |\mathbf{p}_{cm}| \sin \theta, 0, iE_{cm}) \quad (\text{B.13})$$

$$\bar{P} = (|\mathbf{p}_{cm}| \cos \theta, -|\mathbf{p}_{cm}| \sin \theta, 0, iE_{cm}) \quad (\text{B.14})$$

$$\Lambda = (-\Lambda_x, 0, 0, iE_{cm}) \quad (\text{B.15})$$

$$\bar{\Lambda} = (\Lambda_x, 0, 0, iE_{cm}) \quad (\text{B.16})$$

$$K = (|\mathbf{p}_{cm}| \cos \theta - \Lambda_x, -|\mathbf{p}_{cm}| \sin \theta, 0, 0) \quad (\text{B.17})$$

The center of mass energy E_{cm} is shared by every hadron except the kaon, which carries no energy and is thereby off-shell and a virtual exchange particle.