

Math 120

PSet 2

Sep 12 2024

# Contents

Chapter 1

Page 2







1.1 PSet 2

2

# Chapter 1

## 1.1 PSet 2

### Question 1



- (a)  $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$
- (b)  $\vec{r}(t) = t \langle \cos t, \sin t, t \rangle$
- (c)  $\vec{r}(t) = \langle \cos t, \sin t, t^3 \rangle$
- (d)  $\vec{r}(t) = \langle \cos(t^3), \sin(t^3), t^3 \rangle$
- (e)  $\vec{r}(u) = \langle \cos u, \sin u, 1 + \sin(4u) \rangle$
- (f)  $\vec{r}(u) = \langle \cos u, \sin u, 1 + 4 \sin(u) \rangle$
- (g)  $\vec{r}(t) = \langle 2 \cos t, 1 + 4 \cos t, 3 \cos t \rangle$

**Solution:**

### Question 2

2. Find a vector function that represents the curve of intersection of the plane  $z = -2$  and the sphere  $x^2 + (y - 1)^2 + (z + 1)^2 = 9$ .

**Solution:**

Question 3

Consider the vector-valued function  $\vec{r}_1(t) = \langle 2 \sin t, -3 \cos t, 0 \rangle$ ,  $0 \leq t \leq 2\pi$ .

- (a) Sketch the plane curve given by  $\vec{r}_1(t)$ .
- (b) Compute and draw on your sketch from part (a) the position vector  $\vec{r}_1\left(\frac{2\pi}{3}\right)$  and the tangent vector  $\vec{r}'_1\left(\frac{2\pi}{3}\right)$ .
- (c) The vector-valued function  $\vec{r}_2(t) = \langle 2 \cos(3t), -3 \sin(3t) \rangle$  parameterizes the same curve. Find the smallest  $t^* > 0$  such that  $\vec{r}_2(t^*) = \vec{r}_1\left(\frac{2\pi}{3}\right)$ , and compute  $\vec{r}'_2(t^*)$ . Explain how and why  $\vec{r}'_2(t^*)$  differs from the tangent vector  $\vec{r}'_1\left(\frac{2\pi}{3}\right)$  you computed in part (b).

**Solution:**

Question 4

Find parametric equations for the tangent line to the curve parameterized by

$$x = 2t + 1, \quad y = e^{t^2-4}, \quad z = \ln(1 + t^2)$$

at the point  $(5, 1, \ln 5)$ .

**Solution:**

### Question 5

- (a) Evaluate the integral  $\int (\tan t \hat{i} + \sin^2 t \hat{j} + \sec^2 t \tan t \hat{k}) dt$ .
- (b) Suppose a particle is at the point  $(-2, 1, 4)$  at time  $t = 0$ , and moves according to the velocity function  $\vec{v}(t) = \tan t \hat{i} + \sin^2 t \hat{j} + \sec^2 t \tan t \hat{k}$ . Find the particle's position at time  $t = \frac{\pi}{4}$ .

**Solution:**

### Question 6

Consider the curve parameterized by  $\vec{r}(t) = \langle e^{2t}, e^{-2t}, \sqrt{8t} \rangle$ ,  $0 \leq t \leq 1$ .

- (a) Sketch the projections of  $\vec{r}(t)$  in the  $xy$ -,  $zx$ -, and  $yz$ -planes.
- (b) Find the length of the curve. *Hint:* To integrate, you will need to write  $\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2$  as a perfect square.

**Solution:** a)

$$x(t) = e^{2t} \quad y(t) = e^{-2t}$$

$$x \cdot y = e^{2t} \cdot e^{-2t} = 1$$

$$x(t) = e^{2t} \quad z(t) = \sqrt{8t}$$

b)

$$L = \int_{t_1}^{t_2} \|\vec{r}'(t)\| dt$$

$$\vec{r}(t) = \langle e^{2t}, e^{-2t}, \sqrt{8t} \rangle$$

$$\vec{r}'(t) = \left\langle \frac{d}{dt}(e^{2t}), \frac{d}{dt}(e^{-2t}), \frac{d}{dt}(\sqrt{8t}) \right\rangle$$

$$\vec{r}'(t) = \left\langle 2e^{2t}, -2e^{-2t}, \frac{4}{\sqrt{8t}} \right\rangle$$

$$\|\vec{r}'(t)\| = \sqrt{(2e^{2t})^2 + (-2e^{-2t})^2 + \left(\frac{4}{\sqrt{8t}}\right)^2}$$

### Question 7

Let  $C$  be the curve of intersection of the cylinder  $x^2 + y^2 = 4$  and the plane  $2x + y + z = 4$ .

- (a) Find a parameterization of  $C$ .
- (b) Write down an integral for the length of  $C$ .
- (c) Find the length accurate to five decimal places by using Desmos: <https://www.desmos.com/calculator>. (Click on the keyboard icon, then “functions”, then “Misc”, to find the integral symbol.)

**Solution:**

### Question 8

Find the velocity and position vectors of a particle that has acceleration given by

$$\vec{a}(t) = 2\hat{i} + 6t\hat{j} + 12t^2\hat{k},$$

and initial velocity and position given by

$$\vec{v}(0) = \hat{i} \quad \text{and} \quad \vec{r}(0) = \hat{j} - \hat{k}.$$

***Solution:***

#### Question 9

Consider the function  $f(x, y) = \frac{\sqrt{y}-3x}{\ln(4-x^2-y^2)}$ .

- (a) Find and sketch the domain of  $f$ .
- (b) On your sketch from part (a), mark where  $f(x, y) = 0$ , and indicate the region(s) where  $f(x, y)$  is positive and negative.

***Solution:***

### Question 10

Here are several surfaces.

Match each function with its graph. Justify your answers.

(a)  $f(x, y) = x^2$

(b)  $f(x, y) = \sqrt{x^2 + y^2}$

(c)  $f(x, y) = e^{x^2+y^2} - 1$

(d)  $f(x, y) = y \sin x$

(e)  $f(x, y) = \sin(x + y)$

(f)  $f(x, y) = \sin\left(\sqrt{x^2 + y^2}\right)$

**Solution:**

### Question 11

Draw a contour map of the function  $f(x, y) = x^2 e^{-y}$  showing several level curves.

### Question 12

Match the function with its graph (labeled A-F below) and with its contour map (labeled I-VI). Give reasons for your choices.

(a)  $z = e^x \cos y$

(b)  $z = \sin x - \sin y$

(c)  $z = \frac{x-y}{1+x^2+y^2}$