

Math 120

PSet 2

Sep 12 2024

# Contents

Chapter 1

Page 2

1.1 PSet 2

2

# Chapter 1

## 1.1 PSet 2

### Question 1

Consider the line  $L_1$  given by  $x + 2y = 7$  and the line  $L_2$  given by  $5x - y = 2$ .

1. There are two unit vectors that are parallel to  $L_1$ . What are they?
2. There are two unit vectors that are perpendicular to  $L_1$ . What are they?
3. Find the acute angle between the lines  $L_1$  and  $L_2$ . First find an exact expression and then approximate to the nearest degree.

**Solution:**

a)

$$L_1 = x + 2y = 7$$

$$y = -\frac{1}{2}x + \frac{7}{2}$$

$$v_1 = (1, m_1) = (1, -\frac{1}{2})$$

$$|v_1| = \sqrt{1^2 + \left(-\frac{1}{2}\right)^2} = \sqrt{1 + \frac{1}{4}} = \sqrt{\frac{5}{4}} = \frac{\sqrt{5}}{2}$$

$$u_1 = \frac{1}{\frac{\sqrt{5}}{2}} \left(1, -\frac{1}{2}\right) = \frac{2}{\sqrt{5}} \left(1, -\frac{1}{2}\right) = \left(\frac{\sqrt{2}}{5}, -\frac{1}{\sqrt{5}}\right)$$

$$-u_1 = \left(-\frac{\sqrt{2}}{5}, \frac{1}{\sqrt{5}}\right)$$

b)

$$\langle A, B \rangle = \langle 1, 2 \rangle$$

$$|n| = \sqrt{1^2 + 2^2} = \sqrt{5}$$

$$u_1 = \frac{1}{\sqrt{5}}, \frac{2}{\sqrt{5}}$$

$$u_2 = -u_1 = -\frac{1}{\sqrt{5}}, -\frac{2}{\sqrt{5}}$$

c)

$$\vec{v}_1 = \langle 1, -2 \rangle$$

$$\vec{v}_2 = \langle 5, 1 \rangle$$

$$\cos(\theta) = \frac{v_1 \cdot v_2}{|v_1||v_2|}$$

$$\cos(\theta) = \frac{-2+5}{\sqrt{526}} = \frac{3}{\sqrt{130}}$$

$$\theta = \arccos\left(\frac{3}{\sqrt{130}}\right)$$

### Question 2

Find all values of  $x$  such that the angle between the vectors  $\langle 1, -1, 0 \rangle$  and  $\langle 2, x, 1 \rangle$  is  $\frac{\pi}{3}$ .

**Solution:**

$$\begin{aligned} v_1 &= \langle 1, -1, 0 \rangle \\ v_2 &= \langle 2, x, 1 \rangle \\ \cos\left(\frac{\pi}{3}\right) &= \frac{1}{2} \\ \frac{1}{2} &= \frac{v_1 \cdot v_2}{|v_1||v_2|} = \frac{2-x}{(\sqrt{2})\sqrt{5+x^2}} \\ 4-2x &= \sqrt{10+2x^2} \\ 10+2x^2 &= 16-16x+4x^2 \\ -2x^2+16x-6 &= 0 \\ x^2-8x+3 &= 0 \\ x &= \frac{8 \pm \sqrt{(-8)^2 - 4(1)(3)}}{2 \cdot 1} \\ x &= \frac{8 \pm \sqrt{52}}{2} \\ x &= 4 - \sqrt{13} \end{aligned}$$

### Question 3

Find the scalar and vector projections of  $\vec{b} = \hat{i} + \hat{j}$  onto  $\vec{a} = -\hat{i} + 3\hat{j}$ , and illustrate your answers with a sketch.

**Solution:**

Scalar Projection:

$$\begin{aligned} \vec{b} &= \hat{i} + \hat{j} \\ \vec{a} &= -\hat{i} + 3\hat{j} \\ \text{comp}_a \mathbf{b} &= \frac{a \cdot b}{|a|} \\ \frac{a \cdot b}{|a|} &= \frac{2}{\sqrt{10}} = \frac{2\sqrt{10}}{10} = \frac{\sqrt{10}}{5} \end{aligned}$$

Vector Projection:

$$\begin{aligned} \text{proj}_a \mathbf{b} &= \left( \frac{a \cdot b}{|a|^2} \right) a \\ \left( \frac{a \cdot b}{|a|^2} \right) a &= \frac{2}{\sqrt{10}^2} a = \frac{2}{10} a = \frac{1}{5} a \\ \frac{1}{5} a &= \frac{1}{5} (-\hat{i}, 3\hat{j}) = \left\langle -\frac{1}{5} \hat{i}, \frac{3}{5} \hat{j} \right\rangle \end{aligned}$$

#### Question 4

Find two vectors of length 2 that are orthogonal to both  $\vec{v} = \langle 2, 4, 4 \rangle$  and  $\vec{w} = \langle 1, -1, -3 \rangle$ .

**Solution:**

$$\vec{v} \times \vec{w} = \langle 4(-3) - 4(-1), 4(1) - 2(-3), 2(-1) - 4(1) \rangle = \langle -8, 10, -6 \rangle$$

$$|u| = \sqrt{(-8)^2 + 10^2 + (-6)^2} = \sqrt{200}$$

$$2 = x \cdot \sqrt{200}$$

$$x = \frac{2}{\sqrt{200}} = \frac{1}{5\sqrt{2}}$$

$$u_1 = \frac{1}{5\sqrt{2}} \langle -8, 10, -6 \rangle$$

$$u_2 = -\frac{1}{5\sqrt{2}} \langle -8, 10, -6 \rangle$$

#### Question 5

Let  $\vec{a} = \langle 3, 1, 0 \rangle$ . Find all vectors  $\vec{b} = \langle b_1, b_2, b_3 \rangle$  such that  $\vec{a} \times \vec{b}$  is parallel to the z-axis and pointing in the positive z direction. Illustrate with a sketch, in which all vectors are drawn as position vectors, i.e., with the tail at the origin.

**Solution:**

$$\vec{a} \times \vec{b} = \langle 0, 0, c \rangle$$

$$\langle 3, 1, 0 \rangle \times \langle b_1, b_2, b_3 \rangle = \langle 0, 0, c \rangle$$

$$\langle 3, 1, 0 \rangle \times \langle b_1, b_2, b_3 \rangle = \langle 1(b_3) - 0(b_2), 0(b_1) - 3(b_3), 3(b_2) - 1(b_1) \rangle$$

$$\langle 3, 1, 0 \rangle \times \langle b_1, b_2, b_3 \rangle = \langle b_3, -3b_3, 3b_2 - b_1 \rangle$$

$$3b_2 - 3b_1 > 0$$

$$b_3 = 0$$

$$-3b_3 = 0$$

#### Question 6

Consider the four points in  $\mathbb{R}^3$ ,  $K(1, 2, 3)$ ,  $L(1, 3, 6)$ ,  $M(3, 8, 6)$ , and  $N(3, 7, 3)$ .

1. Show that the vectors  $\vec{KL}$ ,  $\vec{KM}$ , and  $\vec{KN}$  are coplanar. Explain why this means that  $K$ ,  $L$ ,  $M$ , and  $N$  all lie in the same plane.
2. From part (a), we know that  $K$ ,  $L$ ,  $M$ , and  $N$  are the vertices of a quadrilateral. Explain how you can tell that this quadrilateral is actually a parallelogram.
3. Find the area of the parallelogram with vertices  $K$ ,  $L$ ,  $M$ , and  $N$ .

**Solution:** a)

$$\vec{KL} = \langle 1 - 1, 3 - 2, 6 - 3 \rangle = \langle 0, 1, 3 \rangle$$

$$\vec{KM} = \langle 3 - 1, 8 - 2, 6 - 3 \rangle = \langle 2, 6, 3 \rangle$$

$$\vec{KN} = \langle 3 - 1, 7 - 2, 3 - 3 \rangle = \langle 2, 5, 0 \rangle$$

$$\vec{KL} \cdot (\vec{KM} \times \vec{KN}) = \vec{KL} \cdot \langle 6(0) - 3(5), 3(2) - 2(0), 2(5) - 6(2) \rangle$$

$$\vec{KL} \cdot (\vec{KM} \times \vec{KN}) = \vec{KL} \cdot \langle -15, 6, -2 \rangle$$

$$\begin{aligned}\vec{KL} \cdot (\vec{KM} \times \vec{KN}) &= \vec{KL} \cdot \langle -15, 6, -2 \rangle = 0(15) + 6(1) + 3(-2) = 0 \\ |\vec{KL} \cdot (\vec{KM} \times \vec{KN})| &= 0\end{aligned}$$

They are coplanar because the volume determined by the vectors is 0, therefore they must lie on the same plane.  
b).

#### Question 7

Find the vector equation and parametric equations for the line through the point  $(1, 2, -2)$  parallel to the line  $x = t - 2$ ,  $y = -2t + 1$ ,  $z = 3$ .

**Solution:**

$$\begin{aligned}x &= t - 2 & y &= -2t + 1 & z &= 3 \\ \vec{d} &= \langle 1, -2, 0 \rangle \\ \vec{r}(t) &= \vec{r}_0 + t\vec{d} \\ \vec{r}(t) &= \langle 1, 2, -2 \rangle + t\langle 1, -2, 0 \rangle \\ x(t) &= 1 + t & y(t) &= 2 - 2t & z(t) &= -2\end{aligned}$$

Vector Equation:  $\vec{r}(t) = \langle 1, 2, -2 \rangle + t\langle 1, -2, 0 \rangle$

Parametric Equation:  $x(t) = 1 + t$ ,  $y(t) = 2 - 2t$ ,  $z(t) = -2$

#### Question 8

Consider the lines  $L_1 : x = t + 3$ ,  $y = 2t - 1$ ,  $z = -t$ , and  $L_2 : x = t - 1$ ,  $y = t - 4$ ,  $z = -t + 4$ . Determine whether the  $L_1$  and  $L_2$  are parallel, skew, or intersecting. If they intersect, find the point of intersection.

**Solution:**

$$\begin{aligned}d_1 &= \langle 1, 2, -1 \rangle & d_2 &= \langle 1, 1, -1 \rangle \\ \frac{1}{1} &\neq \frac{2}{1} \neq \frac{-1}{-1} \\ t_1 + 3 &= t_2 - 1 & 2t_1 - 1 &= t_2 - 4 & -t_1 &= -t_2 + 4 \\ t_1 + 3 &= t_2 - 1 \Rightarrow t_1 - t_2 &= -4 \\ -t_1 &= -t_2 + 4 \Rightarrow t_1 &= t_2 - 4 \\ (t_2 - 4) - t_2 &= -4 \Rightarrow -4 &= -4 \\ 2t_1 - 1 &= t_2 - 4 \\ 2(t_2 - 4) - 1 &= t_2 - 4 \\ 2t_2 - 9 &= t_2 - 4 \\ t_2 &= 5 \\ t_1 &= t_2 - 4 = 5 - 4 = 1 \\ x_1 &= 1 + 3 = 4 & y_1 &= 2(1) - 1 = 1 & z_1 &= -1\end{aligned}$$

Point of Intersection:  $(4, 1, -1)$

#### Question 9

Consider the planes  $x + y + 2z = 4$  and  $2x - y - 2z = 1$ .

1. Find a vector equation for the line of intersection of the planes.
2. Find the angle between the planes. First find an exact expression and then approximate to the nearest degree.

**Solution:**

a)

$$\begin{aligned}
 n_1 &= \langle 1, 1, 2 \rangle \quad n_2 = \langle 2, -1, -2 \rangle \\
 d &= n_1 \times n_2 = \langle 1(-2) - 2(-1), 2(2) - 1(-2), 1(-1) - 1(2) \rangle = \langle -4, 6, -3 \rangle \\
 P_1 : x + y + 2z &= 4 \quad P_2 : 2x - y - 2z = 1 \\
 x + y + 2(0) &= 4 \Rightarrow x + y = 4 \Rightarrow x = -y + 4 \\
 2x - y - 2(0) &= 1 \Rightarrow 2x - y = 1 \\
 2(-y + 4) - y &= 1 \Rightarrow -2y - 8 - y = 1 \Rightarrow -3y = 9 \Rightarrow y = -3 \\
 x &= -(-3) + 4 = 7 \\
 (7, 9, 0) \\
 r &= (7, 9, 0) + t\langle -4, 6, -3 \rangle
 \end{aligned}$$

b)

$$\begin{aligned}
 n_1 &= \langle 1, 1, 2 \rangle \quad n_2 = \langle 2, -1, -2 \rangle \\
 \cos(\theta) &= \frac{n_1 \cdot n_2}{|n_1||n_2|} = \frac{1(2) + 1(-1) + 2(-2)}{\sqrt{1^2 + 1^2 + 2^2}\sqrt{2^2 + (-1)^2 + (-2)^2}} = \frac{-1}{\sqrt{6}\sqrt{9}} = -\frac{1}{3\sqrt{6}} \\
 \theta &= \arccos\left(-\frac{1}{3\sqrt{6}}\right)
 \end{aligned}$$

#### Question 10

Let  $P$  be the plane  $x + y + 2z = 1$  and let  $A$  be the point  $(1, 1, 1)$ .

- Find an equation of the plane through point  $A$  parallel to plane  $P$ .
- Find a vector equation for the line through the point  $A$  which is perpendicular to the plane  $P$ . Call this line  $L$ .
- Find the point of intersection of the line  $L$  (from part (b)) and the plane  $P$ .
- Find the point on the plane  $P$  closest to the point  $A$ , and then find the shortest distance from the point  $A$  to the plane  $P$ .

**Solution:**

a)

$$\begin{aligned}
 P : x + y + 2z &= 1 \quad n_1 = \langle 1, 1, 2 \rangle \\
 n_2 &= t\langle 1, 1, 2 \rangle \\
 P_2 : tx + ty + 2tz &= d \\
 t1 + t1 + 2t(1) &= d \Rightarrow t + t + 2t = d \Rightarrow d = 4t \\
 tx + ty + 2tz &= 4t \Rightarrow x + y + 2z = 4
 \end{aligned}$$

b)

$$\begin{aligned}
 n &= (1, 1, 2) \\
 r(t) &= (1, 1, 1) + t(1, 1, 2)
 \end{aligned}$$

c)

$$\begin{aligned}
 P : x + y + 2z &= 1 \\
 L : x &= 1 + t \quad y = 1 + t \quad z = 1 + 2t \\
 (1 + t) + (1 + t) + 2(1 + 2t) &= 1 \\
 2 + 2t + 2 + 4t &= 1 \Rightarrow 6t + 4 = 1
 \end{aligned}$$

$$6t = -3 \Rightarrow t = -\frac{1}{2}$$

$$x : -\frac{1}{2} + 1 \quad y : -\frac{1}{2} + 1 \quad z : 1 + 2\left(-\frac{1}{2}\right)$$

Point:  $(\frac{1}{2}, \frac{1}{2}, 0)$

d) Point:  $(\frac{1}{2}, \frac{1}{2}, 0)$  because it is when the line through point A is perpendicular to the plane P

$$d = \sqrt{\left(1 - \frac{1}{2}\right) + \left(1 - \frac{1}{2}\right)^2 + (1 - 0)^2} = \sqrt{\frac{1}{4} + \frac{1}{4} + 1} = \sqrt{\frac{3}{2}}$$