

Math 120

PSet 7

Oct 24 2024

Contents

Chapter 1

Page 2

1.1 PSet 7

2

Chapter 1

1.1 PSet 7

Question 1

Evaluate the scalar line integral

$$\int_C (3x + y) ds,$$

where C is the line segment from $(-1, 3)$ to $(4, 2)$.

Solution:

$$\int_C (3x + y) ds$$

$$(-1, 3) \quad (4, 2)$$

$$f(t) = (-1, 3) + t((4, 2) - (-1, 3))$$

$$f(t) = (-1, 3) + t(5, -1) = \langle -1 + 5t, 3 - t \rangle$$

$$x = -1 + 5t \quad y = 3 - t \quad t \in [0, 1]$$

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$\frac{dx}{dt} = 5 \quad \frac{dy}{dt} = -1$$

$$ds = \sqrt{5^2 + (-1)^2} dt = \sqrt{26} dt$$

$$3x + y \Rightarrow 3(-1 + 5t) + (3 - t) \Rightarrow -3 + 15t + 3 - t = 14t$$

$$\int_0^1 14t \sqrt{26} dt \Rightarrow \sqrt{26} \int_0^1 14t dt$$

$$7\sqrt{26}t^2 \Big|_0^1 = 7\sqrt{26}(1)^2 - 7\sqrt{26}(0)^2 = 7\sqrt{26}$$

Question 2

In this problem we will sketch part of the argument that a scalar line integral $\int_C f ds$ is independent of the parameterization of C that we choose to compute the integral. Suppose $\vec{r}_1(t)$, $a \leq t \leq b$, and $\vec{r}_2(t)$, $c \leq t \leq d$, are two smooth parameterizations of the same smooth curve C . Assuming that both parameterizations are in the same direction it can be shown that $\vec{r}_2(t) = \vec{r}_1(w(t))$, for some increasing function $w(t)$ satisfying $w(c) = a$ and $w(d) = b$. If this is the case, show that

$$\int_a^b f(\vec{r}_1(t)) |\vec{r}_1'(t)| dt = \int_c^d f(\vec{r}_2(t)) |\vec{r}_2'(t)| dt$$

for any continuous function f .

Solution:


$$\begin{aligned} & \int_c^d f ds \\ & \begin{matrix} \vec{r}_1(t) & a \leq t \leq b \\ \vec{r}_2(t) & c \leq t \leq d \end{matrix} \\ & \vec{r}_2(r) = \vec{r}_1(w(t)) \quad w(c) = a \quad w(d) = b \\ & \int_a^b f(\vec{r}_1(t)) |\vec{r}_1'(t)| dt = \int_c^d f(\vec{r}_2(t)) |\vec{r}_2'(t)| dt \\ & \vec{r}_2'(t) = \frac{d}{dt} \vec{r}_2(t) = \frac{d}{dt} \vec{r}_1(w(t)) = \vec{r}_1'(w(t)) w'(t) \\ & |\vec{r}_2'(t)| = |\vec{r}_1'(w(t))| \cdot |w'(t)| \\ & \int_c^d f(\vec{r}_2(t)) |\vec{r}_2'(t)| dt = \int_a^b f(\vec{r}_1(t)) |\vec{r}_1'(w(t))| \cdot |w'(t)| dt \\ & w \text{ maps } [c, d] \text{ to } [a, b], \text{ when } t = c, s = a, \text{ and when } t = d, s = b \\ & \int_a^b f(\vec{r}_1(w(t)) |\vec{r}_1'(w(t))| \cdot |w'(t)| dt = \int_a^b f(\vec{r}_1(s)) |\vec{r}_1'(s)| ds \\ & \int_a^b f(\vec{r}_1'(t)) |\vec{r}_1'(t)| dt = \int_a^b f(r_1(s)) |\vec{r}_1'(s)| ds = \int_c^d f(\vec{r}_2(t)) |\vec{r}_2'(t)| dt \end{aligned}$$

\therefore the scalar line integral is independent of the parameterization and the equality holds true for any continuous function f

Question 3

Sketch the vector field $\vec{F}(x, y) = xy \hat{i} + \frac{1}{2} \hat{j}$.

Solution:



Question 4

Given the contour diagram for a function f shown below, in which dark colors correspond to low values of f and light colors correspond to high values of f , sketch the gradient vector field $\vec{F} = \nabla f$.

Question 5

A thin wire has the shape of the curve C parameterized by $x = \cos t$, $y = \sin t$, $z = t$, $0 \leq t \leq 4\pi$, where x , y , and z are measured in centimeters. The linear density of the wire is given by $\rho(x, y, z) = x^2 z$ grams per centimeter. Find the mass of the wire.

Solution:

$$x = \cos t \quad y = \sin t \quad z = t$$

$$0 \leq t \leq 4\pi$$

$$\rho(x, y, z) = x^2 z \frac{\text{grams}}{\text{cm}}$$

$$\text{Mass:} = \int_C \rho(x, y, z) ds$$

$$\frac{dx}{dt} = -\sin t \quad \frac{dy}{dt} = \cos t \quad \frac{dz}{dt} = 1$$

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt$$

$$ds = \sqrt{(-\sin(t))^2 + (\cos(t))^2 + (1)^2} dt$$

$$ds = \sqrt{1 + 1} dt = \sqrt{2} dt$$

$$\rho(x, y, z) = x^2 z \Rightarrow \cos^2 t \cdot t \Rightarrow t \cos^2 t$$

$$\text{Mass:} \int_0^{4\pi} \rho(t) ds = \int_0^{4\pi} t \cos^2 t \sqrt{2} dt$$

$$\sqrt{2} \int_0^{4\pi} t \cos^2 t dt \quad \cos^2 t = \frac{1 + \cos 2t}{2}$$

$$\sqrt{2} \int_0^{4\pi} t \left(\frac{1 + \cos 2t}{2} \right) dt \Rightarrow \frac{\sqrt{2}}{2} \int_0^{4\pi} t(1 + \cos 2t) dt$$

$$\frac{\sqrt{2}}{2} \int_0^{4\pi} t dt + \frac{\sqrt{2}}{2} \int_0^{4\pi} t \cos 2t dt$$

$$\int_0^{4\pi} t dt = \frac{t^2}{2} \Big|_0^{4\pi} \Rightarrow \frac{\sqrt{2}}{2} \frac{16\pi^2}{2} - \frac{\sqrt{2}}{2} \frac{0}{2} = 4\sqrt{2}\pi^2$$

$$u = t \quad du = dt$$

$$v = \frac{1}{2} \sin 2t \quad dv = \cos 2t$$

$$\int t \cos 2t dt = t \cdot \frac{1}{2} \sin 2t - \int \frac{1}{2} \sin 2t dt = \frac{1}{2} t \sin 2t + \frac{1}{4} \cos 2t + k$$

$$\left[\frac{1}{2} t \sin 2t + \frac{1}{4} \cos 2t \right]_0^{4\pi} = \left(\frac{1}{2} \cdot 4\pi \cdot 0 + \frac{1}{4} \cdot 1 \right) - \left(0 + \frac{1}{4} \cdot 1 \right) = 0$$

$$4\sqrt{2}\pi^2 + 0 = 4\sqrt{2}\pi^2$$

Question 6

Let \vec{F} be the vector field shown below, and let C be the unit circle, oriented clockwise. Is the vector line integral

$$\int_C \vec{F} \cdot d\vec{r}$$

positive, negative, or zero? Explain your reasoning.

Question 7

Evaluate the line integral

$$\int_C \sin x \, dx + \cos y \, dy$$

where C consists of the top half of the circle $x^2 + y^2 = 1$ from $(1, 0)$ to $(-1, 0)$ and the line segment from $(-1, 0)$ to $(-2, 3)$. (Remember that when you see an integral that looks like

$$\int_C P(x, y) \, dx + \int_C Q(x, y) \, dy$$

it is a shorthand notation for

$$\int_C \vec{F}(\vec{r}(t)) \cdot d\vec{r}$$

where $\vec{F}(x, y) = \langle P(x, y), Q(x, y) \rangle$. The analogous thing is true in three dimensions.)

Solution:

$$x^2 + y^2 = 1 \quad x = \cos t \quad y = \sin t \quad t \in [0, \pi]$$

$$x(t) = (1 - t)(-1) + t(-2) \quad y(t) = (1 - t)(0) + t(3) \quad t \in [0, 1]$$

$$\vec{F}(x, y) = \langle \sin x, \cos y \rangle$$

$$\int_C \sin x \, dx + \cos y \, dy$$

$$x(t) = \cos t, \quad y(t) = \sin t, \quad t \in [0, \pi]$$

$$\frac{dx}{dt} = -\sin t, \quad \frac{dy}{dt} = \cos t$$

$$\int_{C_1} \sin x \, dx + \cos y \, dy = \int_0^\pi [\sin(\cos t)(-\sin t) + \cos(\sin t) \cos t] \, dt$$

$$\int_0^\pi \sin(\cos t)(-\sin t) \, dt + \int_0^\pi \cos(\sin t) \cos t \, dt$$

$$f(x, y) = -\cos x + \sin y$$

At $(1, 0)$:

$$f(1, 0) = -\cos(1) + \sin(0) = -\cos(1)$$

At $(-1, 0)$:

$$f(-1, 0) = -\cos(1)$$

$$\int_{C_1} \sin x \, dx + \cos y \, dy = f(-1, 0) - f(1, 0) = 0$$

$$x(t) = -1 - t, \quad y(t) = 3t, \quad t \in [0, 1]$$

$$dx = -1 \, dt, \quad dy = 3 \, dt$$

$$\int_{C_2} \sin x \, dx + \cos y \, dy = \int_0^1 [\sin(-1-t)(-1) + \cos(3t)(3)] \, dt$$

Using $\sin(-1-t) = -\sin(1+t)$, the integral becomes:

$$\int_0^1 \sin(1+t) \, dt + 3 \int_0^1 \cos(3t) \, dt$$

$$\int_0^1 \sin(1+t) \, dt = -\cos(1+t) \Big|_0^1 = -\cos(2) + \cos(1)$$

$$3 \int_0^1 \cos(3t) \, dt = 3 \left(\frac{1}{3} \sin(3t) \Big|_0^1 \right) = \sin(3)$$

$$\int_{C_2} \sin x \, dx + \cos y \, dy = \cos(1) - \cos(2) + \sin(3)$$

$$\int_C \sin x \, dx + \cos y \, dy = \int_{C_1} \sin x \, dx + \cos y \, dy + \int_{C_2} \sin x \, dx + \cos y \, dy$$

Since $\int_{C_1} \sin x \, dx + \cos y \, dy = 0$, the total integral is:

$$\int_C \sin x \, dx + \cos y \, dy = \cos(1) - \cos(2) + \sin(3)$$

Question 8

Compute the line integral of the vector field

$$\vec{F}(x, y) = \frac{x}{\sqrt{x^2 + y^2}} \hat{i} + \frac{y}{\sqrt{x^2 + y^2}} \hat{j}$$

along the parabola $x = 1 + y^2$ from $(2, -1)$ to $(2, 1)$.

Question 9

Evaluate the line integral of the vector field

$$\vec{F}(x, y, z) = (x + y)\hat{i} + (y - z)\hat{j} + z^2\hat{k}$$

along the path parameterized by

$$\vec{r}(t) = t^2\hat{i} + t^3\hat{j} + t^2\hat{k}, \quad 0 \leq t \leq 1.$$

Question 10

For each of the following vector fields \vec{F} and curves C , find a function f such that $\vec{F} = \nabla f$ and use this function to evaluate

$$\int_C \vec{F} \cdot d\vec{r}$$

along the given directed curve C .

1. $\vec{F}(x, y) = \langle x^2, y^2 \rangle$, C is the arc of the parabola $y = 2x^2$ from $(-1, 2)$ to $(2, 8)$.
2. $\vec{F}(x, y, z) = \langle e^y, xe^y, (z+1)e^z \rangle$, $C : \vec{r}(t) = \langle t, t^2, t^3 \rangle$, $0 \leq t \leq 1$.

Question 11

Clairaut's Theorem implies that if the vector field $\vec{F} = P\hat{i} + Q\hat{j} + R\hat{k}$ is conservative and P, Q , and R have continuous first-order partial derivatives, then

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}, \quad \frac{\partial P}{\partial z} = \frac{\partial R}{\partial x}, \quad \frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y}.$$

1. Use the statement above to show that the vector line integral

$$\int_C x \, dx + 2x \, dy + xz \, dz$$

is not independent of path.

2. Find two directed curves C_1 and C_2 that start at the same point and end at the same point, such that

$$\int_{C_1} x \, dx + 2x \, dy + xz \, dz \neq \int_{C_2} x \, dx + 2x \, dy + xz \, dz.$$

Question 12

The force exerted by an electric charge at the origin on a charged particle at a point (x, y, z) with position vector $\vec{r} = \langle x, y, z \rangle$ is

$$\vec{F}(\vec{r}) = K \frac{\vec{r}}{|\vec{r}|^3},$$

where K is a constant. Find the work done on the particle as it moves along the straight line from $(0, 3, 0)$ to $(1, 3, 2)$ in two ways:

1. Parameterize the line segment, and compute

$$\int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) \, dt$$

directly.

2. Although \vec{F} is not defined at the origin, it turns out that \vec{F} is conservative on its domain. Find a potential function f , and use the Fundamental Theorem of Line Integrals to compute the work done on the particle.