

# Math 120 QR

Alex Hernandez Juarez

Fall 2024

# Contents

## Chapter 1

## Page 2

1.1	12.1 notes	2
1.2	12.2 Notes	2
1.3	12.3 Notes	3
1.4	12.4 Notes (Cross Product)	5

# Chapter 1

## 1.1 12.1 notes

### Definition 1.1.1: Distance Formula

Defintion:

$$|P_1P_2| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$



### Definition 1.1.2: Equation of a sphere

Defintion: An equation of a sphere with center  $C(h, k, l)$ , and radius  $r$  is

$$(x - h)^2 + (y - k)^2 + (z - l)^2$$

In particular, if the center is the origin  $O$ , than an equation of the sphere is

$$x^2 + y^2 + z^2$$



## 1.2 12.2 Notes

### Definition 1.2.1: Vector Addition

If  $\mathbf{u}$  and  $\mathbf{v}$  are vectors positioned so the initial point of  $\mathbf{v}$  is at the terminal point of  $\mathbf{u}$ , then the **sum**  $\mathbf{u} + \mathbf{v}$  is the vector from the initial point of  $\mathbf{u}$  to the terminal point of  $\mathbf{v}$ .



### Definition 1.2.2: Scalar Multiplication

If  $c$  is a scalar and  $\mathbf{v}$  is a vector, then the **scalar multiple**  $c\mathbf{v}$  is the vector whose length is  $|c|$  times the length of  $\mathbf{v}$  and whose direction is the same as  $\mathbf{v}$  if  $c > 0$  and is opposite to  $\mathbf{v}$  if  $c < 0$  or  $\mathbf{v} = \mathbf{0}$ , then  $c\mathbf{v} = \mathbf{0}$ .



### Example 1.2.1:

Given the points  $A(x_1, y_1, z_1)$  and  $B(x_2, y_2, z_2)$ , the vector  $\mathbf{a}$  with representation  $\overrightarrow{AB}$  is:

$$\mathbf{a} = \langle x_2 - x_1, y_2 - y_1, z_2 - z_1 \rangle$$



### Example 1.2.2:

If  $\mathbf{a} = \langle a_1, a_2 \rangle$  and  $\mathbf{b} = \langle b_1, b_2 \rangle$ , then:

$$\mathbf{a} + \mathbf{b} = \langle a_1 + b_1, a_2 + b_2 \rangle$$

$$\mathbf{a} - \mathbf{b} = \langle a_1 - b_1, a_2 - b_2 \rangle$$

$$c\mathbf{a} = \langle ca_1, ca_2 \rangle$$

Similarly, for three dimensional vectors,

$$\langle a_1, a_2, a_3 \rangle + \langle b_1, b_2, b_3 \rangle = \langle a_1 + b_1, a_2 + b_2, a_3 + b_3 \rangle$$

$$\langle a_1, a_2, a_3 \rangle - \langle b_1, b_2, b_3 \rangle = \langle a_1 - b_1, a_2 - b_2, a_3 - b_3 \rangle$$

$$c\langle a_1, a_2, a_3 \rangle = \langle ca_1, ca_2, ca_3 \rangle$$



### Note:-

Properties of vectors: If  $\mathbf{a}$ ,  $\mathbf{b}$ , and  $\mathbf{c}$  are vectors in  $V_n$  and  $c$  and  $d$  are scalars then

- $\mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a}$
- $\mathbf{a} + (\mathbf{b} + \mathbf{c}) = (\mathbf{a} + \mathbf{b}) + \mathbf{c}$
- $\mathbf{a} + \mathbf{0} = \mathbf{a}$
- $\mathbf{a} + \mathbf{a} + -\mathbf{a} = \mathbf{0}$
- $c(\mathbf{a} + \mathbf{b}) = c\mathbf{a} + c\mathbf{b}$
- $(c + d)\mathbf{a} = c\mathbf{a} + d\mathbf{a}$
- $(cd)\mathbf{a} = c(d\mathbf{a})$
- $l\mathbf{a} = \mathbf{a}$



## 1.3 12.3 Notes

### Definition 1.3.1: Dot Product

If  $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$  and  $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$ , then the **dot product** of  $\mathbf{a}$  and  $\mathbf{b}$  is the number  $\mathbf{a} \cdot \mathbf{b}$  given by

$$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$$

Properties of the Dot Product: If  $\mathbf{a}$ ,  $\mathbf{b}$ , and  $\mathbf{c}$  are vectors in  $V_3$  and  $c$  is a scalar, then

1.  $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$
2.  $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$
3.  $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$
4.  $(c\mathbf{a}) \cdot \mathbf{b} = c(\mathbf{a} \cdot \mathbf{b}) = \mathbf{a} \cdot (c\mathbf{b})$
5.  $\mathbf{0} \cdot \mathbf{a} = 0$




### Definition 1.3.2: Geometric Definition of the Dot Product

If  $\theta$  is the angle between vectors  $\mathbf{a}$  and  $\mathbf{b}$ , then

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos(\theta)$$

Proof:



$$|AB|^2 = |OA|^2 + |OB|^2 - 2|OA||OB| \cos \theta$$

Corollary: If  $\theta$  is the angle between nonzero vectors  $\mathbf{a}$  and  $\mathbf{b}$ , then

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}||\mathbf{b}|}$$



#### Note:-

Two vectors  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal if and only if  $\mathbf{a} \cdot \mathbf{b} = 0$



#### Example 1.3.1 (Direction Angles and Cosines)

The **direction angles** of a nonzero vector  $\mathbf{a}$  are the angles  $\alpha$ ,  $\beta$ , and  $\gamma$  (in the interval  $[0, \pi]$ ) that  $\mathbf{a}$  makes with the positive  $x$ -,  $y$ -, and  $z$ -axes, respectively.

The cosines of these direction angles,  $\cos \alpha$ ,  $\cos \beta$ , and  $\cos \gamma$ , are called the **direction cosines** of the vector  $\mathbf{a}$ . Using Corollary 6 with  $\mathbf{b}$  replaced by  $\mathbf{i}$ , we obtain:

$$\cos \alpha = \frac{\mathbf{a} \cdot \mathbf{i}}{|\mathbf{a}||\mathbf{i}|} = \frac{a_1}{|\mathbf{a}|} \quad (1)$$

Similarly, we also have:

$$\cos \beta = \frac{a_2}{|\mathbf{a}|} \quad \text{and} \quad \cos \gamma = \frac{a_3}{|\mathbf{a}|} \quad (2)$$

By squaring the expressions in Equations 8 and 9 and adding, we see that:

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1 \quad (3)$$

We can also use Equations 8 and 9 to write:

$$\mathbf{a} = \langle a_1, a_2, a_3 \rangle = \langle |\mathbf{a}| \cos \alpha, |\mathbf{a}| \cos \beta, |\mathbf{a}| \cos \gamma \rangle = |\mathbf{a}| \langle \cos \alpha, \cos \beta, \cos \gamma \rangle$$

Therefore,

$$\frac{1}{|\mathbf{a}|} \mathbf{a} = \langle \cos \alpha, \cos \beta, \cos \gamma \rangle \quad (4)$$

which says that the direction cosines of  $\mathbf{a}$  are the components of the unit vector in the direction of  $\mathbf{a}$ .

### Definition 1.3.3: Projections

The **scalar projection** of  $\mathbf{b}$  onto  $\mathbf{a}$  (also called the **component of  $\mathbf{b}$  along  $\mathbf{a}$** ) is defined to be the signed magnitude of the vector projection, which is the number  $|\mathbf{b}| \cos \theta$ , where  $\theta$  is the angle between  $\mathbf{a}$  and  $\mathbf{b}$ . This is denoted by  $\text{comp}_{\mathbf{a}} \mathbf{b}$ . Observe that it is negative if  $\pi/2 < \theta \leq \pi$ . The equation

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta = |\mathbf{a}|(|\mathbf{b}| \cos \theta)$$

shows that the dot product of  $\mathbf{a}$  and  $\mathbf{b}$  can be interpreted as the length of  $\mathbf{a}$  times the scalar projection of  $\mathbf{b}$  onto  $\mathbf{a}$ . Since

$$|\mathbf{b}| \cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|} = \frac{\mathbf{a}}{|\mathbf{a}|} \cdot \mathbf{b}$$

the component of  $\mathbf{b}$  along  $\mathbf{a}$  can be computed by taking the dot product of  $\mathbf{b}$  with the unit vector in the direction of  $\mathbf{a}$ . We summarize these ideas as follows.

**Scalar projection of  $\mathbf{b}$  onto  $\mathbf{a}$ :**  $\text{comp}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|}$

**Vector projection of  $\mathbf{b}$  onto  $\mathbf{a}$ :**  $\text{proj}_{\mathbf{a}} \mathbf{b} = \left( \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|^2} \right) \mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|^2} \mathbf{a}$



## 1.4 12.4 Notes (Cross Product)

### Definition 1.4.1: Cross Product

Given two nonzero vectors  $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$  and  $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$ , suppose that a nonzero vector  $\mathbf{c} = \langle c_1, c_2, c_3 \rangle$  is perpendicular to both  $\mathbf{a}$  and  $\mathbf{b}$ . Then  $\mathbf{a} \cdot \mathbf{c} = 0$  and  $\mathbf{b} \cdot \mathbf{c} = 0$ , and so:

$$a_1 c_1 + a_2 c_2 + a_3 c_3 = 0 \quad (1)$$

$$b_1 c_1 + b_2 c_2 + b_3 c_3 = 0 \quad (2)$$

To eliminate  $c_3$ , we multiply (1) by  $b_3$  and (2) by  $a_3$  and subtract:

$$(a_1 b_3 - a_3 b_1) c_1 + (a_2 b_3 - a_3 b_2) c_2 = 0 \quad (3)$$

Equation (3) has the form  $p c_1 + q c_2 = 0$ , for which an obvious solution is  $c_1 = q$  and  $c_2 = -p$ . So, a solution of (3) is:

$$c_1 = a_2 b_3 - a_3 b_2$$

$$c_2 = a_3 b_1 - a_1 b_3$$

Substituting these values into (1) and (2), we then get:

$$c_3 = a_1 b_2 - a_2 b_1$$

This means that a vector perpendicular to both  $\mathbf{a}$  and  $\mathbf{b}$  is:

$$\langle c_1, c_2, c_3 \rangle = \langle a_2 b_3 - a_3 b_2, a_3 b_1 - a_1 b_3, a_1 b_2 - a_2 b_1 \rangle$$

The resulting vector is called the **cross product** of  $\mathbf{a}$  and  $\mathbf{b}$  and is denoted by  $\mathbf{a} \times \mathbf{b}$ .



#### Definition 1.4.2: Cross Product of two vectors

If  $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$  and  $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$  then the **cross product** of  $\mathbf{a}$  and  $\mathbf{b}$  is:

$$\mathbf{a} \times \mathbf{b} = \langle a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1 \rangle$$



#### Note:-

Determinant of order 2:

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$



#### Note:-

Determinant of order 3:

$$\begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = a_1 \begin{vmatrix} b_2 & b_3 \\ c_2 & c_3 \end{vmatrix} - a_2 \begin{vmatrix} b_1 & b_3 \\ c_1 & c_3 \end{vmatrix} + a_3 \begin{vmatrix} b_1 & b_2 \\ c_1 & c_2 \end{vmatrix}$$



#### Definition 1.4.3: Second definition of cross product

Arithmetic Definition:

$$\begin{aligned} \mathbf{a} \times \mathbf{b} &= \begin{bmatrix} i & j & k \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{bmatrix} = |\mathbf{a}| |\mathbf{b}| \sin(\theta) \\ &= \begin{bmatrix} a_2 & a_3 \\ b_2 & b_3 \end{bmatrix} i - \begin{bmatrix} a_1 & a_3 \\ b_1 & b_3 \end{bmatrix} j + \begin{bmatrix} a_1 & a_2 \\ b_1 & b_2 \end{bmatrix} k \\ &= (a_2b_3 - a_3b_2)i - (a_1b_3 - a_3b_1)j + (a_1b_2 - a_2b_1)k \end{aligned}$$

The vector  $\mathbf{a} \times \mathbf{b}$  is orthogonal to both  $\mathbf{a}$  and  $\mathbf{b}$



#### Example 1.4.1: Proof that $\mathbf{a} \times \mathbf{b}$ is orthogonal to both $\mathbf{a}$

$$\begin{aligned} (\mathbf{a} \times \mathbf{b}) \cdot \mathbf{a} &= \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} a_1 - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} a_2 + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} a_3 \\ &= a_1(a_2b_3 - a_3b_2) - a_2(a_1b_3 - a_3b_1) + a_3(a_1b_2 - a_2b_1) \\ &= a_1a_2b_3 - a_1a_3b_2 - a_2a_1b_3 + a_2a_3b_1 + a_3a_1b_2 - a_3a_2b_1 \\ &= 0 \end{aligned}$$



**Definition 1.4.4: sin definition of cross product**

If  $\theta$  is the angle between  $\mathbf{a}$  and  $\mathbf{b}$  (so  $0 \leq \theta \leq \pi$ ), then the length of the cross product  $\mathbf{a} \times \mathbf{b}$  is given by:

$$|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}||\mathbf{b}| \sin(\theta)$$

Proof:

$$\begin{aligned} |\mathbf{a} \times \mathbf{b}|^2 &= (a_2b_3 - a_3b_2)^2 + (a_3b_1 - a_1b_3)^2 + (a_1b_2 - a_2b_1)^2 \\ &= a_2^2b_3^2 - 2a_2a_3b_2b_3 + a_3^2b_2^2 + a_3^2b_1^2 - 2a_1a_3b_1b_3 + a_1^2b_3^2 + a_1^2b_2^2 - 2a_1a_2b_1b_2 + a_2^2b_1^2 \\ &= (a_1^2 + a_2^2 + a_3^2)(b_1^2 + b_2^2 + b_3^2) - (a_1b_1 + a_2b_2 + a_3b_3)^2 \\ &= |\mathbf{a}|^2|\mathbf{b}|^2 - (\mathbf{a} \cdot \mathbf{b})^2 \\ &= |\mathbf{a}|^2|\mathbf{b}|^2 - |\mathbf{a}|^2|\mathbf{b}|^2 \cos^2 \theta \quad (\text{by Theorem 12.3.3}) \\ &= |\mathbf{a}|^2|\mathbf{b}|^2(1 - \cos^2 \theta) \\ &= |\mathbf{a}|^2|\mathbf{b}|^2 \sin^2 \theta \end{aligned}$$

Taking square roots and observing that  $\sqrt{\sin^2 \theta} = \sin \theta$  because  $\sin \theta \geq 0$  when  $0 \leq \theta \leq \pi$ , we have

$$|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}||\mathbf{b}| \sin \theta$$

**Note:-**

Two nonzero vectors  $\mathbf{a}$  and  $\mathbf{b}$  are parallel if and only if

$$\mathbf{a} \times \mathbf{b} = \mathbf{0}$$

**Example 1.4.2: Geometric interpretation of  $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}||\mathbf{b}| \sin \theta$** 

If  $\mathbf{a}$  and  $\mathbf{b}$  are represented by directed line segments with the same initial point, then they determine a parallelogram with base  $|\mathbf{a}|$ , altitude  $|\mathbf{b}| \sin(\theta)$  and area

$$A = |\mathbf{a}|(|\mathbf{b}| \sin \theta) = |\mathbf{a} \times \mathbf{b}|$$

Thus we have the following way of interpreting the magnitude of a cross product:

The length of the cross product of  $\mathbf{a} \times \mathbf{b}$  is equal to the area of the parallelogram determined by  $\mathbf{a}$  and  $\mathbf{b}$ .

